

Exercises in orthodox geometry

Edited by A. Petrunin
`orthodox.geometry@gmail.com`

Contents

1	Curves and surfaces	5
2	Comparison geometry	9
3	Curvature free differential geometry	13
4	Metric geometry	17
5	Actions and coverings	21
6	Topology	23
7	Piecewise linear geometry	25
8	Discrete geometry	27
A	Semisolutions	29
B	Dictionary	81

For problem solvers. The meaning of signs next to number of the problem:

- — easy problem;
- * — the solution requires at least two ideas;
- + — the solution requires knowledge of a theorem;
- ‡ — there are interesting solutions based on different ideas.

To get a hint, send an e-mail to the above address with the number and the name of the problem.

For problem makers. This collection is under permanent development. If you have suitable problems or corrections please e-mail it to me.

Many thanks. I want to thank everyone sharing the problems. Also I want to thank J. Luukkainen, R. Matveyev, P. Petersen, S. Tabachnikov and number of students in my classes for their interest in this list and for correcting number of mistakes. I'm also thankful to everyone who took part in the discussion of this list on *mathoverflow*.

Chapter 1

Curves and surfaces

For the most of problems in this section it is enough to know definitions of curvature of curves, second fundamental form, Gauss curvature of surfaces, Gauss–Bonnet theorem. To solve Problem 23, it is better know Hairy Ball theorem.

1. Geodesic for birds. Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a ℓ -Lipschitz function. Consider the epigraph $W \subset \mathbb{R}^3$ of f ; i.e.,

$$W = \{ (x, y, z) \in \mathbb{R}^3 \mid z \geq f(x, y) \}$$

and equip W with the induced intrinsic metric. Show that any *geodesic* in W has *variation of turn* at most $2 \cdot \ell$.


2. Spiral. Let γ be a plane curve with strictly monotonic curvature function. Prove that γ has no self-intersections.

In other words, if you drive on the plane and turn the steering wheel to the right all the time then you can not come back to the same place.

3. Divergence. Let F be a convex figure in the \mathbb{R}^2 with smooth boundary and area $F = \pi$. Show that there is a vector field \mathbf{v} on F such that $|\mathbf{v}| \leq 1$ and $\operatorname{div} \mathbf{v} \geq 2$.

4. The moon in the puddle. A smooth closed simple plane curve with curvature less than 1 bounds a figure F . Prove that F contains a disc or radius 1.

5. Closed surface. Construct a smooth embedding $\mathbb{S}^2 \hookrightarrow \mathbb{R}^3$ with principle curvatures in the interval $(-1, 1)$ such that it does not bound a ball of radius 1.



6[‡] *A curve in a sphere.* Prove that any closed curve on unit sphere which intersects every equator has length at least 2π .

7[‡] *A spring in a tin.* Let α be a closed smooth immersed curve inside a unit disc. Prove that the average absolute curvature of α is at least 1, with equality if and only if α is the unit circle possibly traversed more than once.

8[°] *Convex hat.* Let Σ be a smooth closed convex surface in the 3-dimensional Euclidean space and Π be a plane which cuts from Σ a disc Δ . Assume that the reflection of Δ in Π lies inside Σ . Show that Δ is convex in the intrinsic metric of Σ ; that is, if the ends of a minimizing geodesic in Σ lie in Δ then whole geodesic lies in Δ .

9. *Unbended geodesic.* Let Σ be a smooth closed convex surface in the 3-dimensional Euclidean space and $\gamma: [0, \ell] \rightarrow \Sigma$ be a unit speed minimizing geodesic in Σ . Set $p = \gamma(0)$, $q = \gamma(\ell)$ and

$$p_t = \gamma(t) - t \cdot \gamma'(t),$$

for $t \in (0, \ell)$ where $\gamma'(t)$ denotes the velocity vector of γ at t .

Show that one can not see q from the point p ; i.e., the line segment $[pq]$ intersects Σ at a point distinct from q .

10. *A minimal surface.* Let Σ be a *minimal surface* in \mathbb{R}^3 which has boundary on a unit sphere. Assume Σ passes through the center of the sphere. Show that area of Σ is at least π .

11.* *Half-torus.* Consider torus T ; i.e., a surface of revolution generated by revolving a circle in \mathbb{R}^3 about an axis coplanar with the circle. Let $\gamma \subset T$ be a circle at which T is tangent to a fixed plane and Ω be an open set in T containing γ .

Assume Ω' is an other smooth open surface which is path isometric to Ω and sufficiently close to Ω . Show that Ω' contains a congruent copy of γ .

12. *Asymptotic line.* Consider a smooth surface Σ in \mathbb{R}^3 given as a graph $z = f(x, y)$. Let γ be a closed smooth asymptotic line on Σ . Assume Σ is *strictly saddle* in a neighborhood of γ . Prove that projection of γ on xy -plane is not star shaped.

13. *Closed magnetic geodesics.* Give an example of a closed Riemannian 2-manifold which has no closed smooth curve with constant geodesic curvature = 1.

14.° Non-contractable geodesics. Give an example of a non-flat metric on 2-torus such that it has non-contractible geodesics.

15.° Convex figures. Consider the set of all convex figures \mathfrak{C} in the plane equipped with Hausdorff distance. Show that the set of smooth figures forms a G-delta dense subset in \mathfrak{C} .

A convex figure in the plane is said to be smooth if it has unique supporting line at every boundary point.

16. Fat curve. Construct a *simple plane curve* with non-zero Lebesgue's measure.

17. Rectifiable curve. Assume X is a compact connected set in \mathbb{R}^2 with finite 1-dimensional Hausdorff measure. Show that X can be presented as the image of a rectifiable curve $\gamma: [0, 1] \rightarrow \mathbb{R}^2$.


18. Capture a sphere in a knot. Let B be the closed unit ball in \mathbb{R}^3 and $f: \mathbb{S}^1 \rightarrow \mathbb{R}^3 \setminus B$ be a knot. Show that there is an ambient isotopy

$$H_t: \mathbb{R}^3 \setminus B \rightarrow \mathbb{R}^3 \setminus B, \quad t \in [0, 1],$$

such that $H_0 = \text{id}$, the length of $H_t \circ f$ does not increase in t and $H_1(f(\mathbb{S}^1))$ can be disjointed from B by a plane.

19.° Linked circles. Suppose that α and β are disjoint linked Jordan curves in \mathbb{R}^3 which lie at a distance 1 from each other. Show that the length of α is at least $2 \cdot \pi$.

20. Oval in oval. Consider two closed smooth strictly convex planar curves, one inside another. Show that there is a chord of the outer curve, which is tangent to the inner curve at its midpoint.




21. Periodic asymptote. Let Σ be a smooth surface with nonpositive curvature and γ be a geodesic in Σ . Assume that γ is not periodic and the curvature of Σ vanish at every point of γ . Show that γ does not have a periodic asymptote; i.e., there is no periodic geodesic δ such that the distance from $\gamma(t)$ to δ converges to 0 as $t \rightarrow \infty$.

22. Immersed surface. Let Σ be a smooth connected immersed surface in \mathbb{R}^3 with strictly positive Gauss curvature and nonempty boundary $\partial\Sigma$. Assume $\partial\Sigma$ lies in a plane Π and whole Σ lies on one side from Π . Prove that Σ is an embedded disc.

23. Two discs. Let Σ_1 and Σ_2 be two smoothly embedded open discs in \mathbb{R}^3 which have common closed smooth curve γ . Show that there is a pair of points $p_1 \in \Sigma_1$ and $p_2 \in \Sigma_2$ with parallel tangent planes.

24. Simple geodesic. Let Σ be a closed unbounded convex surface in the 3-dimensional Euclidean. Show that there is a two-sided infinite geodesic in Σ with no self-intersections.

25. Long geodesic. Assume that the surface of convex body B in \mathbb{R}^3 admits an arbitrary long closed geodesic. Show that B is a tetrahedron with equal opposite sides.



26.° Surrounded area. Let $\gamma_1, \gamma_2: \mathbb{S}^1 \rightarrow \mathbb{R}^2$ be two simple closed plane curves. Assume

$$|\gamma_1(v) - \gamma_1(w)| \leq |\gamma_2(v) - \gamma_2(w)|$$

for any $v, w \in \mathbb{S}^1$. Show that the area surrounded by γ_1 does not exceed the area surrounded by γ_2 .

27. Corkscrew geodesic. Given line ℓ in the Euclidean space construct a closed convex body K with a minimizing geodesic σ on the surface of K which rotates 1000 times around ℓ .

Chapter 2

Comparison geometry

For doing most of problems in this section it is enough to know second variation formula. Knowledge of some basic results such as O'Neil formula, Gauss formula, Gauss–Bonnet theorem, Toponogov's comparison theorem, Soul theorem, Toponogov splitting theorems and Synge's lemma also might help. We suggest to solve problem 22 before solving problem 30, To solve problem 36, it is better know that factor positively curved Riemannian manifold by an isometry group is a positively curved Alexandrov space. Problem 37 requires Liouville's theorem for geodesic flow. Problem 45 requires a Bochner type formula.

28. *Totally geodesic hypersurface.* Prove that if a compact positively curved m -dimensional manifold M admits a totally geodesic embedded hypersurface then M or its double cover is homeomorphic to the sphere.

29. *Immersed convex hypersurface I.* Let M be a complete simply connected Riemannian manifold with nonpositive curvature with dimension at least 3. Prove that any immersed locally convex compact hypersurface in M is embedded.

30* *Immersed convex hypersurface II.* Prove that any immersed locally convex hypersurface $\iota: \Sigma \looparrowright M$ in a compact positively curved manifold M of dimension $m \geq 3$, is the boundary of an immersed ball. I.e., there is an immersion of a closed ball $\bar{B}^m \looparrowright M$ such that the induced immersion of its boundary $\partial \bar{B}^m \looparrowright M$ gives ι .

31. *Almgren's inequalities.* Let Σ be a closed m -dimensional *minimal surface* in \mathbb{S}^n . Prove that $\text{vol}_m \Sigma \geq \text{vol}_m \mathbb{S}^m$.

32. *Hypercurve.* Let $M^m \hookrightarrow \mathbb{R}^{m+2}$ be a closed smooth m -dimensional submanifold and let g be the induced Riemannian metric on M^m . Assume that sectional curvature of g is positive. Prove that curvature operator of g is positively defined.

33. *Horosphere.* Let M be a complete simply connected manifold with negatively pinched sectional curvature (i.e. $-a^2 \leq K \leq -b^2 < 0$). And let $\Sigma \subset M$ be an horosphere in M (i.e., Σ is a level set of a *Busemann function* in M). Prove that Σ with the induced intrinsic metric has *polynomial volume growth*.

34. *Minimal spheres.* Show that a compact positively curved 4-manifold can not contain infinite number of mutually *equidistant minimal 2-spheres*.

35*. *Totally geodesic immersion.* Let (M, g) be a simply connected positively curved m -manifold and $\iota: N \looparrowright M$ be a totally geodesic immersion. Prove that if $\dim N > \frac{m}{2}$ then N is embedded.

36⁺. *Positive curvature and symmetry.* Prove that effective isometric S^1 -action on a 4-dimensional positively curved closed Riemannian manifold has at most 3 isolated fixed points.

37⁺. *Scalar curvature vs. injectivity radius.* Let (M, g) be a closed Riemannian m -manifold with scalar curvature $\text{Sc}_g \geq m \cdot (m-1)$ (i.e., bigger then scalar curvature of \mathbb{S}^m). Prove that the injectivity radius of (M, g) is at most π .

38. *Almost flat manifold.* Show that for any $\varepsilon > 0$ there is $n = n(\varepsilon)$ such that there is a compact n -dimensional manifold M which is not a finite quotient of a *nil-manifold* and which admits a Riemannian metric with diameter ≤ 1 and sectional curvature $|K| < \varepsilon$.

39. *Lie group.* Let g_0 and g_1 be left invariant Riemannian metrics on a Lie group G . Assume that both g_0 and g_1 have nonnegative sectional curvature; show that so is $g_0 + g_1$.

40[‡]. *Polar points.* Let (M, g) be a compact Riemannian manifold with sectional curvature ≥ 1 . Prove that for any point $p \in M$ there is a point $p^* \in M$ such that

$$|p - x|_g + |x - p^*|_g \leq \pi$$

for any $x \in M$.

41. *Deformation to a product.* Let (M, g) be a compact Riemannian manifold with non-negative sectional curvature. Show that there is a continuous one parameter family of non-negatively curved metrics g_t on M , $t \in [0, 1]$, such that $g_0 = g$ and a finite Riemannian cover of (M, g_1) is isometric to a product of a flat torus and a simply connected manifold.

42.* *Isometric section.* Let M and W be compact Riemannian manifolds and $s: W \rightarrow M$ be a Riemannian submersion. Assume that W has positive sectional curvature. Show that s does not admit an isometric section; i.e., there is no isometric embedding $\iota: M \hookrightarrow W$ such that $s \circ \iota(p) = p$ for any $p \in M$.

43.† *Minkowski space.* Let us denote by \mathbb{M}^m the set \mathbb{R}^m equipped with the metric induced by the ℓ^p -norm. Prove that if $p \neq 2$ then \mathbb{M}^m can not be a Gromov–Hausdorff limit of Riemannian m -manifolds (M_n, g_n) such that $\text{Ric}_{g_n} \geq C$ for some constant $C \in \mathbb{R}$.

44. *Curvature hollow.* Construct a Riemannian metric g on \mathbb{R}^3 which is Euclidean outside of an open bounded set Ω and scalar curvature of g is negative in Ω .

45.+ *If hemisphere then sphere.* Let M is an m -dimensional Riemannian manifold with Ricci curvature at least $m - 1$; moreover there is a point $p \in M$ such that sectional curvature is exactly 1 at all points on distance $\leq \frac{\pi}{2}$ from p . Show that M has constant sectional curvature.

46. *Flat coordinate planes.* Assume g be a Riemannian metric on \mathbb{R}^3 , such that the coordinate planes $x = 0$, $y = 0$ and $z = 0$ are flat and totally geodesic. Assume g has sectional curvature ≥ 0 (or ≤ 0). Show that origin admits a flat neighborhood.

47.† *Two-convexity.* Assume K be a closed set in \mathbb{R}^4 which contains two coordinate planes

$$\{(x, y, 0, 0) \in \mathbb{R}^4\} \quad \text{and} \quad \{(0, 0, z, t) \in \mathbb{R}^4\}$$

in its interior and lies in the closed 1-neighborhood of these two planes. Assume that K is bounded by hypersurface W and the complement $\mathbb{R}^4 \setminus K$ is two-convex; i.e., if three faces of given tetrahedron lie in $\mathbb{R}^4 \setminus K$ then so is the forth face. Prove that W can not be C^∞ -smooth.

Chapter 3

Curvature free differential geometry

A solution of problem 48 relies on Gromov's pseudo-holomorphic curves. The problem 50 uses Liouville's theorem for geodesic flow. A solution of the problem 58 use the Birkhoff's (or an other) curve shortening process.

48.⁺ *Minimal foliation.* Consider $\mathbb{S}^2 \times \mathbb{S}^2$ equipped with a Riemannian metric g which is C^∞ -close to the product metric. Prove that there is a conformally equivalent metric $\lambda \cdot g$ and re-parametrization of $\mathbb{S}^2 \times \mathbb{S}^2$ such that each sphere $\mathbb{S}^2 \times x$ and $y \times \mathbb{S}^2$ forms a *minimal surface* in $(\mathbb{S}^2 \times \mathbb{S}^2, \lambda \cdot g)$.

49. *Loewner's theorem.* Given \mathbb{RP}^n equipped with a Riemannian metric g conformally equivalent to the canonical metric g_{can} let ℓ denote the minimal length of curves in (\mathbb{RP}^n, g) which are not null-homotopic. Prove that

$$\pi^n \cdot \text{vol}(\mathbb{RP}^n, g) \geq \ell^n \cdot \text{vol}(\mathbb{RP}^n, g_{\text{can}})$$

and in case of equality $g = c \cdot g_{\text{can}}$ for some positive constant c .

50.⁺ *Convex function vs. finite volume.* Let M be a complete Riemannian manifold which admits a non-constant convex function. Prove that M has infinite volume.

51. *Besikovitch inequality.* Let g be a Riemannian metric on a n -dimensional cube $Q = [0, 1]^n$ such that any curve connecting opposite faces has length ≥ 1 . Prove that $\text{vol}(Q, g) \geq 1$ and equality holds if and only if (Q, g) is isometric to the interior of the unit cube.

52. *Distant involution.* Construct a Riemannian metric g on \mathbb{S}^3 and involution $\iota: \mathbb{S}^3 \rightarrow \mathbb{S}^3$ such that $\text{vol}(\mathbb{S}^3, g)$ is arbitrary small and $|x - \iota(x)|_g > 1$ for any $x \in \mathbb{S}^3$.

53.^o *Normal exponential map.* Let M, N be complete connected Riemannian manifolds. Assume N is immersed into M . Show that the image of the *normal exponential map* of N is dense in M .

54. *Symplectic squeezing in the torus.* Let $\omega = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$ be the standard symplectic form on \mathbb{R}^4 and \mathbb{Z}^2 the integer lattice in (x_1, y_1) coordinate plane.

Show that arbitrary bounded domain $\Omega \subset (\mathbb{R}^4, \omega)$ admits a symplectic embedding into $(\mathbb{R}^4, \omega)/\mathbb{Z}^2$.

55.^o *Diffeomorphism test.* Let M and N be complete n -dimensional simply connected Riemannian manifolds. Assume $f: M \rightarrow N$ is a smooth map such that

$$|df(v)| \geq |v|$$

for any vector v in the tangent bundle of M . Show that f is a diffeomorphism.

56. *Volume of tubular neighborhoods.* Assume M and M' be isometric closed smooth submanifolds in \mathbb{R}^n . Show that for all small r we have

$$\text{vol } B_r(M) = \text{vol } B_r(M'),$$

where $B_r(M)$ denotes r -neighborhood of M .

57.* *Disc.* Let L be a positive real. Construct a Riemannian metric g on the disc \mathbb{D} with

$$\text{diam}(\mathbb{D}, g) \leq 1 \quad \text{and} \quad \text{length } \partial\mathbb{D} \leq 1$$

such that any homotopy of the boundary to a point in (\mathbb{D}, g) must pass through a curve of length at least L .

58.⁺ *Shortening homotopy.* Let M be a compact Riemannian manifold with diameter D . Assume that for some $L > D$, there are no geodesic loops in M with length in the interval $(L - D, L + D]$. Show that for any path γ_0 in (M, g) there is a homotopy γ_t rel. to the ends such that

- a) $\text{length } \gamma_1 < L$;
- b) $\text{length } \gamma_t \leq \text{length } \gamma_0 + 2 \cdot D$ for any $t \in [0, 1]$.

59° *Geodesic hypersurface.* Let M be a hypersurface in a closed Riemannian m -dimensional manifold W . Assume M is geodesic and convex; denote by i_M the injectivity radius of M . Show that there is a point in W which lies on the distance at least $\frac{i_M}{2 \cdot (m+1)}$ from M .

Chapter 4

Metric geometry

The necessary definitions can be found in [24]. It is very hard to do 61 without using Kuratowski embedding. To do problem 67 you should be familiar with the proof of Nash–Kuiper theorem. For the problem 68 you have to know Rademacher’s theorem on differentiability of Lipschitz maps. To solve the problem 73 one has to know basic theorem on deformation of homeomorphisms.

60° *Noncontracting map.* Let K be a compact metric space and $f: K \rightarrow K$ be a noncontracting map. Prove that f is an isometry.

61⁺ *Embedding of a compact.* Prove that any compact metric space is isometric to a subset of a compact length spaces.

62. *2-sphere is far from the disc.* Show that there is no sequence of Riemannian metrics on S^2 which converge in Gromov–Hausdorff topology to the standard ball $\bar{B}^2 \subset \mathbb{R}^2$.

63. *3-sphere is close to a ball.* Construct a sequence of Riemannian metrics on S^3 which converge in Gromov–Hausdorff topology to the standard ball $\bar{B}^3 \subset \mathbb{R}^3$.

64° *Macrodimension.* Let M be a simply connected Riemannian manifold with the following property: any closed curve can be shrunk to a point in an ϵ -neighborhood of itself. Prove that the macrodimension of M on the scale ϵ is at most 1.

65. *Anti-collapse.* Construct a sequence of Riemannian metric g_i on a 2-sphere such that $\text{vol } g_i < 1$ such that induced distance functions

$d_i: \mathbb{S}^2 \times \mathbb{S}^2 \rightarrow \mathbb{R}_+$ converge to a metric $d: \mathbb{S}^2 \times \mathbb{S}^2 \rightarrow \mathbb{R}_+$ with arbitrary large Hausdorff dimension.

66.* *No short embedding.* Construct a length-metric on \mathbb{R}^3 which admits no local *short* embeddings into \mathbb{R}^3 .

67.⁺ *Sub-Riemannian sphere.* Prove that any *sub-Riemannian metric* on the n -sphere is isometric to the intrinsic metric of a hypersurface in \mathbb{R}^{n+1} .

68.⁺ *Length preserving map.* Show that there is no *length preserving map* $\mathbb{R}^2 \rightarrow \mathbb{R}$.

69. *Hyperbolic space.* Show that the hyperbolic 3-space is *quasi-isometric* to a subset of product of two hyperbolic planes.

70. *Midpoints.* Let $d(x, y) = |x - y|$ be a metric on \mathbb{R}^n induced by a norm $|\cdot|$.

Assume that $f: (\mathbb{R}^n, d) \rightarrow (\mathbb{R}^n, d)$ is an isometry such that

$$f(a) = a \quad \text{and} \quad f(b) = b$$

for some a and b in \mathbb{R}^n . Show that

$$f\left(\frac{a+b}{2}\right) = \frac{a+b}{2}.$$

71. *Hilbert problem.* Let ρ be a length-metric on the plane \mathbb{R}^2 such that any minimizing geodesics in (\mathbb{R}^2, ρ) runs along a line segment. Show that there is uniquely defined measure μ on the set of all lines in \mathbb{R}^2 such that $\rho(x, y)$ coincides with μ -measure of the set of lines which separate x from y .

72. *Straight geodesics.* Let ρ be a length-metric on \mathbb{R}^n , which is bi-Lipschitz equivalent to the canonical metric. Assume that every *geodesic* γ in (\mathbb{R}^n, ρ) is linear (i.e., $\gamma(t) = v + w \cdot t$ for some $v, w \in \mathbb{R}^n$). Show that ρ is induced by a norm on \mathbb{R}^n .

73.⁺ *A homeomorphism near quasi-isometry.* Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a *quasi-isometry*. Show that there is a (bi-Lipschitz) homeomorphism $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ on a bounded distance from f ; i.e., there is a real constant C such that

$$|f(x) - h(x)| \leq C$$

for any $x \in \mathbb{R}^n$.

74. *A family of sets with no section.* Construct a one parameter family of closed sets C_t in \mathbb{S}^1 , $t \in [0, 1]$ which is continuous in Hausdorff topology, but which does not admit a *section*; i.e., there is no continuous map $c: [0, 1] \rightarrow \mathbb{S}^1$ such that $c(t) \in C_t$ for any t .

75. *Sasaki metric.* Let (\mathbb{TS}^2, \hat{g}) be the tangent bundle over \mathbb{S}^2 equipped with Sasaki metric induced by a Riemannian metric g on \mathbb{S}^2 . Show that (\mathbb{TS}^2, \hat{g}) lies on bounded Gromov–Hausdorff distance to the ray.

Chapter 5

Actions and coverings

To do problem 78 you have to know in addition a construction of compact manifolds of constant negative curvature of given dimension m and it is better to do the problem 77 first.

76. *Bounded orbit.* Let X be a proper metric space and $\iota: X \rightarrow X$ is an isometry. Assume that for some $x \in X$, the sequence $x_n = \iota^n(x)$, $n \in \mathbb{Z}$ has a converging subsequence. Prove that x_n is bounded.

77. *Covers of figure eight.* Let (Φ, d) be a “figure eight”; i.e., a metric space which is obtained by gluing together all four ends of two unit segments.

Prove that any compact length spaces K is a Gromov–Hausdorff limit of a sequence of metric covers $(\tilde{\Phi}_n, \tilde{d}/n) \rightarrow (\Phi, d/n)$.


78.⁺ *Constant curvature is everything.* Given integer $m \geq 2$, prove that any compact length spaces K is a Gromov–Hausdorff limit of a sequence of m -dimensional manifolds M_n with curvature $-n^2$.

79.^{*} *Diameter of m -fold cover.* Let X be a length space and \tilde{X} be a connected m -fold cover of X equipped with induced intrinsic metric. Prove that

$$\text{diam } \tilde{X} \leq m \cdot \text{diam } X.$$

80.^o *Symmetric square.* Let X be a connected topological space. Note that $X \times X$ admits natural \mathbb{Z}_2 -action by $(x, y) \mapsto (y, x)$. Show that fundamental group of $X \times X / \mathbb{Z}_2$ is commutative.

81.° *Sierpinski triangle.* Find the homeomorphism group of Sierpinski triangle.



82. *Lattices in a Lie group.* Let L and M be two discrete subgroups of a connected Lie group G , h be a left invariant metric on G . Equip the groups L and M with the induced left invariant metric from G . Assume $L \backslash G$ and $M \backslash G$ are compact and moreover

$$\text{vol}(L \backslash (G, h)) = \text{vol}(M \backslash (G, h)).$$

Prove that there is bi-Lipschitz one-to-one mapping (not necessarily a homomorphism) $f: L \rightarrow M$.

83.° *Piecewise Euclidean quotient.* Let Γ be a finite subgroup of $\text{SO}(n)$; denote by P the quotient \mathbb{R}^n / Γ equipped with induced *polyhedral metric*. Assume P is *PL-homeomorphic* to \mathbb{R}^n . Show that Γ is generated by rotations around subspaces of codimension 2.

84.° *Subgroups of free group.* Show that every finitely generated subgroup of the free group is an intersection of subgroups of finite index.

85.° *Lengths of generators of the fundamental group.* Let M be a compact Riemannian manifold and $p \in M$. Show that the fundamental group $\pi_1(M, p)$ is generated by the homotopy classes of loops with length at most $2 \cdot \text{diam } M$.

86. *Short basis.* Let M be a compact nonnegatively curved n -dimensional Riemannian manifold. Show that the minimal number of generators of $\pi_1 M$ can be bounded above only in terms of n .

Chapter 6

Topology

87. *Milnor's disks.* Construct two *topologically different* smooth immersions of the disk into the plane which coincide near the boundary. (Two immersions $f_1, f_2: D \looparrowright \mathbb{R}^2$ are topologically different if there is no diffeomorphism $h: D \rightarrow D$ such that $f_1 = f_2 \circ h$.)

88. *Positive Dehn twist.* Let Σ be an oriented surface with non empty boundary. Prove that any composition of *positive Dehn twists* of Σ is not homotopic to identity *rel* boundary.

89. *Function with no critical points.* Given $n \geq 2$, construct a smooth function f defined on a neighborhood of closed unit ball B^n in \mathbb{R}^n which has no critical points and which can not be presented in the form $\ell \circ \varphi$, where $\ell: \mathbb{R}^n \rightarrow \mathbb{R}$ is a linear function and $\varphi: B^n \rightarrow \mathbb{R}^n$ is a smooth embedding.

90. *Conic neighborhood.* Let $p \in X$ be a point in a topological space X . We say that an open neighborhood U_p of $p \in X$ is conic if there is a homeomorphism from a cone to U_p which sends its vertex to p . Show that any two conic neighborhoods of p are homeomorphic to each other. [66]

Note that for two cones $\text{Cone}(\Sigma_1)$ and $\text{Cone}(\Sigma_2)$ might be homeomorphic while Σ_1 and Σ_2 are not.

91. *No C^0 -knots.* Prove that the set of smooth embeddings $f: \mathbb{S}^1 \rightarrow \mathbb{R}^3$ equipped with C^0 -topology forms a connected space.

92. *Stabilization.* Construct two compact subsets $K_1, K_2 \subset \mathbb{R}^2$ such that K_1 is not homeomorphic to K_2 , but $K_1 \times [0, 1]$ is homeomorphic to $K_2 \times [0, 1]$.

93. Isotropy. Let K_1 and K_2 be compact subsets of the coordinate subspace \mathbb{R}^n of $\mathbb{R}^{2 \cdot n}$. Show that there is a homeomorphism $h: \mathbb{R}^{2 \cdot n} \rightarrow \mathbb{R}^{2 \cdot n}$ such that $K_2 = h(K_1)$.

Moreover, h can be chosen to be isotopic to the identity map.

94. Cantor set. Construct a subset C of \mathbb{R}^3 such that C is homeomorphic to the Cantor set and the complement $\mathbb{R}^3 \setminus C$ is not simply connected.

95. Knaster's circle. Construct a bounded open set in \mathbb{R}^2 whose boundary does not contain a *simple curve*.

96° Boundary in \mathbb{R} . Construct three disjointed non-empty open sets in \mathbb{R} which have the same boundary.

97. Deformation of homeomorphism. Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a homeomorphism such that $|x - f(x)| < \varepsilon$ for any $x \in \mathbb{R}^n$ and some fixed $\varepsilon > 0$. Construct a homomorphism $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that (1) $|x - h(x)| < \varepsilon$ for any $x \in \mathbb{R}^n$, (2) $h(x) = f(x)$ if $|x| \leq 1$ and (3) $h(x) = x$ if $|x| \geq 2$.

98° Finite topological space. Prove that every finite simplicial complex is weakly homotopy equivalent to a finite topological space, and the other way around.

Chapter 7

Piecewise linear geometry

To do Problem 99, google “cyclic polytopes”. Before solving problem 105, it is better to learn what is Delaunay triangulation.

99. *Triangulation of 3-sphere.* Construct a triangulation of \mathbb{S}^3 such with 100 vertices such that any two vertices are connected by an edge.

100. *Spherical arm lemma.* Let $A = a_1 a_2 \dots a_n$ and $B = b_1 b_2 \dots b_n$ be two simple spherical polygons with equal corresponding sides. Assume A lies in a hemisphere and $\angle a_i \geq \angle b_i$ for each i . Show that A is congruent to B .

101. *Piecewise linear isometry I.* Let P be a compact m -dimensional polyhedral space. Construct a piecewise linear isometry $f: P \rightarrow \mathbb{R}^m$.

102.⁺ *Piecewise linear isometry II.* Prove that any short map to \mathbb{R}^2 which is defined on a finite subset of \mathbb{R}^2 can be extended to a piecewise linear isometry $\mathbb{R}^2 \rightarrow \mathbb{R}^2$.

103. *Minimal polyhedron.* By polyhedral disc in \mathbb{R}^3 we understand a triangulation of a plane polygon with a map in \mathbb{R}^3 which is affine on each triangle. The area of the polyhedral disc is defined as the sum of areas of the images of the triangles in the triangulation.

Consider the class of polyhedral discs glued from n triangles in \mathbb{R}^3 with fixed broken line as the boundary. Let Σ_n be a surface of minimal area in this class. Show that Σ_n is a *saddle surface*.

Note that it is not longer true if Σ minimizes area only in the class of polyhedral surfaces with fixed triangulation.

104. *Coherent triangulation.* A triangulation of a convex polygon is called coherent if there is a convex function which is linear on each triangle and changes the gradient if you come through any edge of the triangulation. Find a non-convex triangulation of triangle.

105.⁺ *Characterization of polytope.* Let P be a compact subset of Euclidean space. Assume for every point $x \in P$ there is a cone K_x with tip at x and $\varepsilon > 0$ such that

$$B_\varepsilon(x) \cap P = B_\varepsilon(x) \cap K_x.$$

Show that P is a polytope; i.e., P is a union of finite collection of simplices.

106.^{*} *A sphere with one edge.* Given a spherical polyhedral space P , denote by P_s the subset of its singular points.

Construct spherical polyhedral space P which is homeomorphic to \mathbb{S}^3 and such that P_s is formed by a knotted circle. Show that in such an example the total length of P_s can be arbitrary large and the angle around P_s can be made strictly less than 2π .

107. *Triangulation of a torus.* Show that torus does not admit a triangulation such that one vertex has 5 edges, one has 7 edges and all other vertexes have 6 edges.

108.[°] *Unique geodesics imply CAT(0).* Let P be a polyhedral space. Assume that any two points in P are connected by unique geodesic. Show that P is a CAT(0) space.

109.[°] *No simple geodesic.* Construct a convex polyhedron P which surface does not admit closed simple geodesic.

Chapter 8

Discrete geometry

One of the solutions of 110 uses mixed volumes. In order to solve problem 112, it is better to know what is the genus of complex curve of degree d . To solve problem 113 one has to use axiom of choice. In order to solve problem 118, it is better to know Dehn–Sommerville equations, see an outline on page 82.

110. *Box in a box.* Assume that a parallelepiped with sizes a, b, c lies inside another with parallelepiped sizes a', b', c' . Show that $a' + b' + c' \geq a + b + c$.

111.^o *Round circles in \mathbb{S}^3 .* Suppose that you have a finite collection of round circles in the unit 3-sphere, not necessarily all of the same radius, such that each pair is linked. Prove that there is an isotopy in the space of such collections of circles so that afterwards, they are all great circles.

112.⁺ *Harnack's circles.* Prove that a smooth algebraic curve of degree d in \mathbb{RP}^2 consists of at most $n = \frac{1}{2} \cdot (d^2 - 3 \cdot d + 4)$ connected components.

113.⁺ *Two points on each line.* Construct a set in the Euclidean plane, which intersect each line at exactly 2 points.

114.^o *Bodies with the same of shadows.* Two convex bodies K_1 and K_2 in Euclidean 3-space are said to have the same shadows if any shape which can appear as an orthogonal projection of K_1 can also appear as an orthogonal projection of K_2 and the other way around.

Construct two noncongruent convex bodies K_1 and K_2 which have the same shadows.

115.^o *Kissing number.* Show that for any convex body W in \mathbb{R}^n

$$\text{kiss } W \geq \text{kiss } B,$$

where $\text{kiss } W$ denotes the *kissing number* of W and B denotes the unit ball in \mathbb{R}^n .

116. *Monotonic homotopy.* Let F be a finite set and $h_0, h_1: F \rightarrow \mathbb{R}^m$ be two maps. Consider \mathbb{R}^m as a subspace of $\mathbb{R}^{2 \cdot m}$. Show that there is a homotopy $h_t: F \rightarrow \mathbb{R}^{2 \cdot m}$ from h_0 to h_1 such that for any $x, y \in F$ the function

$$t \mapsto |h_t(x) - h_t(y)|$$

is monotonic.

117. *Cube.* Assume the 2^n vertices of n -dimensional cube are divided into two sets A and B with the same number of vertices in each. Show that there are at least 2^{n-1} edges with the ends in the different sets.

118.⁺ *Right-angled polyhedron.* Show that in all sufficiently large dimensions, there is no compact convex hyperbolic polyhedron with right dihedral angles.

Appendix A

Semisolutions

Curves and surfaces

1. *Geodesic for birds.* Consider a geodesic

$$t \mapsto (x(t), y(t), z(t))$$

in W ; assume it is defined in the interval $\mathbb{I} \subset \mathbb{R}$. Let us denote by φ the variation of turn; it is a measure on \mathbb{I} . We need to estimate $\varphi(\mathbb{I})$.

Denote by $s = s(t)$ the natural parameter of the plane curve

$$t \mapsto (x(t), y(t)).$$

Prove that the function $f: s \mapsto z$ is concave.

Given a semiopen interval $\mathbb{J} = (a, b] \subset \mathbb{I}$, set $\mu(\mathbb{J}) = f^+(a) - f^+(b)$, where f^+ denotes right derivatives. The function μ extends to a measure which could be also written as

$$\mu = \frac{dz^2}{d^2s} \cdot ds.$$

if $\frac{dz^2}{d^2s}$ understood in the sense of distribution.

Note that $|\frac{dz}{ds}| \leq \ell$. In particular $\mu(\mathbb{I}) \leq 2 \cdot \ell$.

Further note that $\varphi \leq \sqrt{1 + \ell^2} \cdot \mu$. In particular,

$$\varphi(\mathbb{I}) \leq 2 \cdot \ell \cdot \sqrt{1 + \ell^2}.$$

A straightforward improvement of these estimates gives the optimal bound


$$\varphi(\mathbb{I}) \leq 2 \cdot \ell.$$

Comments. The problem appears in the paper of Berg, [15]. The main observation in this proof (concavity of $s \mapsto z$) was used earlier by Liberman in [70] in order to bound on the variation of turn of a geodesic on a convex surface.

2. Spiral. Without loss of generality we may assume that the curvature of γ decreases in t .

Let $z(t)$ be the center of osculating circle at $\gamma(t)$ and $r(t)$ is its radius. Prove that

$$|z'(t)| \leq r'(t).$$



Conclude that the osculating discs are nested; i.e., $D_{t_1} \supset D_{t_0}$ for $t_1 > t_0$. Hence the result follows.

Comments. The problem can be considered as a continuous analog of the Leibniz's test for alternating series.

It seems that the problem first discovered by Tait in [107] and later reproved by Kneser in [64]; see also [83].

3. Divergence. Fix a Cartesian coordinate system (x, y) on the plane. Denote by D the unit disc centered at the origin; so $\text{area } F = \text{area } D$.

Prove that there is a smooth area preserving map $F \rightarrow D$, of the form $(x, y) \mapsto (v(x), w(x, y))$.

Show that the vector field $\mathbf{v}(x, y) = (v(x), w(x, y))$ is a solution for the problem.

Comments. This is the key ingredient in the famous proof of isoperimetric inequality — to finish the proof apply the divergence formula for \mathbf{v} .

The same argument works in all dimensions. See the Gromov's appendix in [77] for more details.

4. The moon in the puddle. Consider the *cut locus* of F with respect to ∂F ; it is defined as the closure of the set of points $x \in F$ such that there are two or more points in ∂F which minimize distance to x . We will denote this cut locus by W .

Note that after a small perturbation of ∂F we may assume that W is a graph embedded in F with finite number of edges.


Note that W is a deformation retract of F . The retraction can be obtained by moving each point $y \in F \setminus W$ to W along the geodesic from the closest point to y on ∂F which pass through y .

In particular, W is a tree. Therefore W has at least two end vertices; Denote one of them by z .

Prove that the disc of radius 1 centered at z lies completely in F .

Comments. A spherical version of this problem was used by Panov and me in [89].

5. Closed surface. The solution should be guessed from the picture.



Comments. This solution is based on so called *Bing's House*, see [18].

6. A curve in a sphere. Let α be a closed curve in \mathbb{S}^2 of length $2 \cdot \ell$ which intersects each equator.

A solution with Crofton formula. Note that we can assume that α is a broken line.

Given a unit vector u denote by e_u the equator with pole at u . Let $k(u)$ the number of intersections of the α and e_u .

Note that for almost all $u \in \mathbb{S}^2$, the value $k(u)$ is even. Since each equator intersects α , we get $k(u) \geq 2$ for almost all u .

Applying Crofton formula, we get

$$\begin{aligned} 2 \cdot \ell &= \frac{1}{4} \cdot \int_{\mathbb{S}^2} k(u) \cdot d_u \text{ area} \geq \\ &\geq \frac{1}{2} \cdot \text{area } \mathbb{S}^2 = \\ &= 2 \cdot \pi. \end{aligned}$$

The *Crofton formula* (i.e., the first identity above) it is easy to obtain it for each geodesic segment of α and then sum it up.

Solution with symmetry. Let $\tilde{\alpha}$ be a subarc of α of length ℓ , with endpoints p and q . Let z be the midpoint of a minimizing geodesic $[pq]$ in \mathbb{S}^2 .

Let r be a point of intersection of α with the equator with pole at z . Without loss of generality we may assume that $r \in \tilde{\alpha}$.

The arc $\tilde{\alpha}$ together with its reflection in z form a closed curve of length $2 \cdot \ell$ that passes through r and its antipodal point r' . Therefore

$$\ell = \text{length } \tilde{\alpha} \geq |r - r'|_{\mathbb{S}^2} = \pi.$$

Comments. This problem is a the first step in the proof of Reshetnyak's motorization theorem for CAT[1] spaces, see [9].

7. A spring in a tin. Let α be a closed curve in the unit disc; denote by ℓ its length.

Let us equip the plane with complex coordinates so that 0 is the center of the unit disc. We can assume that α equipped with ℓ -periodic parametrization by length.

Consider the curve $\beta(t) = t - \frac{\alpha(t)}{\alpha'(t)}$. Note that

$$\beta(t + \ell) = \beta(t) + \ell$$

for any t . In particular

$$\text{length}(\beta|_{[0, \ell]}) \geq |\beta(\ell) - \beta(0)| = \ell.$$

Note that

$$\begin{aligned} |\beta'(t)| &= \left| \frac{\alpha(t) \cdot \alpha''(t)}{\alpha'(t)^2} \right| \leq \\ &\leq |\alpha''(t)|. \end{aligned}$$

Since $|\alpha''(t)|$ is the curvature of α at t , we get the result.

Comment. This proof admits straightforward generalization to the higher dimensions.

If instead of a disc we have a region bounded by closed convex curve γ then it is still true that the average absolute curvature of α is at least as big as average absolute curvature of γ . The proof is not that simple, see [104] and the reference there in.

8. Convex hat. Let γ be a minimizing geodesic with the ends in Δ .

Assume $\gamma \setminus \Delta \neq \emptyset$. Denote by γ' the curve formed by $\gamma \cap \Delta$ and the reflection on $\gamma \setminus \Delta$ in Π . Note

$$\text{length } \gamma' = \text{length } \gamma$$

and γ' runs partly in and partly outside of the surface, but does not get inside of Σ .

Denote by γ'' the closest point projection of γ' on Σ . The curve γ'' lies in Σ has the same ends as γ . It remains to note that

$$\text{length } \gamma'' < \text{length } \gamma;$$

the later leads to a contradiction.

9. Unbended geodesic. Let W be the closed unbounded set formed by Σ and its exterior points.

Prove that for any $x \in \Sigma$ the distance $|x - p_t|$ is nondecreasing in t .

Use the later statement, to prove the same for $|x - p_t|_W$, where $|x - p_t|_W$ stays for the intrinsic distance from x to p_t in W .

Prove that $|q - p|_W = |q - p|_\Sigma$, where $|q - p|_\Sigma$ denotes the intrinsic distance from p to q in Σ .

Conclude that the distance $|q - p_t|_W = |q - p|_\Sigma$ for any t . It follows that the curve

$$\gamma_t(\tau) = \begin{cases} (\tau - t) \cdot \gamma'(\tau) & \text{if } \tau < t; \\ \gamma(\tau) & \text{if } \tau > t. \end{cases}$$

is a minimizing geodesic from p_t to q in the intrinsic metric of W .

If q is visible from p_t for some t then the line segment $[qp_t]$ intersects Σ only at q . From above, γ_t coincides with the line segment $[qp_t]$ which is impossible.

Comment. The problem appears in Milka's paper [76].

10. A minimal surface. Without loss of generality we may assume that the sphere is centered at $0 \in \mathbb{R}^3$.

Consider the restriction h of the function $x \mapsto |x|^2$ to the surface Σ . Prove that $\Delta_\Sigma h \leq 2$ and apply the divergence theorem for $\nabla_\Sigma h$. It follows that the function

$$f: r \mapsto \frac{\text{area}(\Sigma \cap B_r(0))}{r^2}$$

is non-decreasing in the interval $(0, 1)$. Hence the result follows.

Comments. We described a partial case of so called *monotonicity formula*.

Note that if we assume in addition that the surface is a disc, then the statement holds for any saddle surface. Indeed, denote by S_r the sphere of radius r concentrated with the unit sphere. Then according to Problem 6, $\text{length } \Sigma \cap S_r \geq 2 \cdot \pi \cdot r$. Then coarea formula leads to the solution.

On the other hand there are saddle surfaces homeomorphic to the cylinder may have arbitrary small area in the ball.

If Σ does not pass through the center and we only know the distance r from center to Σ then optimal bound is expected to be $\pi \cdot (1 - r^2)$. This is known if Σ is topological disc, see [4]. An analogous result for area-minimizing submanifolds holds for dimensions and codimensions, see [5].

11. Half-torus. Let K be the convex hull of Ω' . Consider the boundary curve γ' of $\partial K \cap \Omega'$ in Ω' .

First note that the Gauss curvature of Ω' has to vanish at the points of γ' ; in other words, γ' is the image of γ under the length-preserving map. Indeed since γ' lies on convex part, the Gauss curvature at the points of γ' has to be nonnegative. On the other hand γ' bounds a flat disc in ∂K ; therefore its integral intrinsic curvature has to be $2\cdot\pi$. If the Gauss curvature is positive at some point of γ' then total intrinsic curvature of γ' has to be $< 2\cdot\pi$, a contradiction.

Assume that the asymptotic direction goes transversally to $\gamma'(t)$ at some point $\gamma'(t)$. Show that $\gamma(t) \notin \partial K$. It follows that γ is an asymptotic curve of Ω' .

Without loss of generality, we can assume that the length of γ is $2\cdot\pi$ and its intrinsic curvature is $\equiv 1$. Therefore, as the space curve, γ' has to be a curve with constant curvature 1 and it should be closed. Any such curve is congruent to a flat circle.

Comments. It is not known if Ω' is congruent to Ω .

The solution presented above is based on my answer to the question of O'Rourke, see [84]. Here are some related statements.

- ◊ Half-torus is second order rigid; this was proved in [97] and [34].
- ◊ Any second order rigid surface does not admit analytic deformation and for the surfaces of revolution, the assumption of analyticity can be removed.

12. Asymptotic line. Arguing by contradiction, assume that the projection $\bar{\gamma}$ of γ on xy -plane is star shaped with respect to the origin.

Consider the function

$$h(t) = (d_{\bar{\gamma}(t)}f)(\gamma(t)).$$

Prove that $h'(t) \neq 0$. In particular $h(t)$ is a strictly monotonic function of \mathbb{S}^1 , a contradiction.

Comments. The problem discussed by Panov in [87].

13. Closed magnetic geodesics. Take Σ to be a compact surface with constant curvature -1 ; i.e., $\Sigma = \mathbb{H}^2/\Gamma$, where \mathbb{H}^2 is the hyperbolic plane and Γ is a discrete subgroup of isometries of \mathbb{H}^2 which acts freely.

Prove that Γ contains only hyperbolic elements; i.e., any transformation $\gamma \in \Gamma$ fix two distinct points on the absolute.

Assume Σ contains a closed curve with curvature 1. Note that this curve lifts to a horocycle in \mathbb{H}^2 . Therefore Γ has to contain an element which preserves this horocycle. Such an element has to be parabolic; i.e., it fixes a single point on the absolute of \mathbb{H}^2 , a contradiction.

Comments. The problem appears in the Ginzburg's paper [42].

14. Non contractable geodesics. Take a torus of revolution T ; the rotations of the circle produce a family closed geodesics which we will call *meridians*.


Note that a geodesic on T is either a meridian or it is transversal to all the meridians. Note that closed curve of these types can not be contractable.

Comments. This problem appears from Gromov's book [48], where it is attributed to Y. Colin de Verdière.

15. Convex figures. Consider the set Ω_n of all convex figures $F \subset \mathbb{R}^2$ such that for any $x \in \partial F$ there are $y, z \in F$ such that $\angle yxz > \pi - \frac{1}{n}$.

Prove that Ω_n is open and dense in \mathfrak{C} . Finally note that the intersection $\bigcap_n \Omega_n$ forms the subset of all smooth figures in \mathfrak{C} .

16. Fat curve. Modify your favorite space filling to keep area nearly the same and removing self-intersections.



Say, you can modify the Hilbert curve which can be constructed as a limit of recursively defined sequence of curve; see the 5-th iteration on the diagram.

17. Rectifiable curve. The 1-dimensional Hausdorff measure will be denoted as \mathcal{H}_1 .

Set $L = \mathcal{H}_1(K)$. Without loss of generality, we may assume that K has diameter 1.

Assume that $0 < \varepsilon < \frac{1}{2}$. Prove that

$$(*) \quad \mathcal{H}_1(B_\varepsilon(x) \cap K) \geq \varepsilon$$

for any $x \in K$.

Let x_1, \dots, x_n be a maximal set of points in K such that

$$|x_i - x_j| \geq \varepsilon$$

for all $i \neq j$. From $(*)$ we have $n \leq 2 \cdot L / \varepsilon$.

Construct is a curve γ_ε such that (1) γ_ε is passing through all x_i , (2) length $\gamma_\varepsilon \leq 10 \cdot L$ and (3) γ_ε lies in ε -neighborhood of K . We can assume that γ_ε is parametrized by length.

The needed curve can be obtained by passing to a partial limit of γ_ε as $\varepsilon \rightarrow 0$.

Comments. This is Exercise 3.5 in the book [36] by Falconer.

18. Capture a sphere in a knot. We can assume that the knot is given by a diagram on the the sphere.

Fix a Möbius transformation $\mathbb{S}^2 \rightarrow \mathbb{S}^2$. Denote by u its conformal factor. Since the Möbius transformation preserves total area, we get

$$\frac{1}{\text{area } \mathbb{S}^2} \cdot \int_{\mathbb{S}^2} u^2 = 1.$$

Therefore,

$$\frac{1}{\text{area } \mathbb{S}^2} \cdot \int_{\mathbb{S}^2} u < 1.$$

It follows that after a suitable rotation of \mathbb{S}^2 , the length of the knot decreases.

Similar argument gives a continuous one parameter family of Möbius transformations which moves the knot in a hemisphere and allows the ball to escape.

Comments. This is a question of Brady, see [29].

19. Linked circles. Fix a point $x \in \alpha$. Note that one can find another point $x' \in \alpha$ such that the interval $[xx']$ intersects β , say at the point z . Otherwise we can move each point of α along the line segment to x . This deformation of α will not cross β ; the later contradicts that α and β are linked.

Consider the curve α' which is the central projection of α from z onto the unit sphere around z ; clearly


$$\text{length } \alpha \geq \text{length } \alpha'.$$

Note that α' passes through two antipodal points of the sphere; therefore

$$\text{length } \alpha' \geq 2 \cdot \pi.$$

Hence the result follows.

Comments. This is the simplest case of so called *Gehring's problem*. The solution above was given by Edelstein and Schwatz in [33]; later the same solution was rediscovered few times.



20. Oval in oval. Given a unit vector u , denote by x_u the point on the inner curve with outer normal vector u . Draw a chord of outer curve which is tangent to the inner curve at x_u ; denote by $r = r(u)$ and $l = l(u)$ the lengths of this chord at the right and left from x_u .

Arguing by contradiction, assume $r(u) \neq l(u)$ for any $u \in \mathbb{S}^1$. Since the functions r and l are continuous, we can assume that

$$(*) \quad r(u) > l(u) \quad \text{for any } u \in \mathbb{S}^1.$$

Prove that each of the following two integrals

$$\frac{1}{2} \cdot \int_{\mathbb{S}^1} r^2(u) \cdot du \quad \text{and} \quad \frac{1}{2} \cdot \int_{\mathbb{S}^1} l^2(u) \cdot du$$

give the area between the curves. In particular the integrals are equal to each other. The later contradicts (*).

Comments. This is a problem of Tabachnikov, see [105]. A closely related, so called *equal tangents problem* is discussed by the same author in [106].

21. Periodic asymptote. Assume contrary. Passing to a finite cover, we can assume that the asymptote has no self intersections. In this case the restriction $\gamma|_{[a, \infty)}$ has no self-intersections if a is large large enough.

Cut Σ along $\gamma([a, \infty))$ and then cut from the obtained surface an infinite triangle \triangle with two sides formed by both sides of cuts along γ ; let us denote these sides of \triangle by γ_- and γ_+ . Note that

$$\text{area } \triangle < \text{area } \Sigma < \infty$$

and both sides γ_{\pm} form infinite minimizing geodesics in \triangle .

Consider the Buseman function f for γ_+ ; denote by $\ell(t)$ the length of the level curve $f^{-1}(t)$. Let $-\kappa(t)$ be the total curvature of the suplevel set $f^{-1}([t, \infty))$. Note that for all large t we have

$$\ell'(t) = \kappa(t) \quad \text{and} \quad \kappa'(t) \leq C \cdot \ell(t)^2$$

where C is a fixed constant. The later implies that there is $\varepsilon > 0$ such that

$$\ell(t) \geq \frac{\varepsilon}{t - a}$$

for any large t . In particular,

$$\int_a^{\infty} \ell(t) = \infty.$$

By coarea formula we get

$$\text{area } \triangle = \infty,$$

a contradiction.

Comment. I've learned the problem from Dmitri Burago and Sergei Ivanov, it is originated from a discussions between Keith Burns, Michael Brin and Yakov Pesin.

22. Immersed surface. Let ℓ be a linear function which vanish on Π and positive in Σ .

Let z be a point of maximum of ℓ on Σ ; set $s_0 = \ell(z)$. Given $s < s_0$, denote by Σ_s the connected component of z in $\Sigma \cap \ell^{-1}([s, s_0])$. Note that for all s sufficiently close to s_0 we have

- ◊ Σ_s is an embeded disc;
- ◊ $\partial\Sigma_s$ is convex plane curve.

Applying open-close argument, we get that the same holds for all $s \in [0, s_0]$.

Since Σ is connected, $\Sigma_0 = \Sigma$. Hence the result follows.

Comments. This problem is a discussed in Gromov's lectures [47].

23. Two discs. Choose a continuous map $h: \Sigma_1 \rightarrow \Sigma_2$ which is identical on γ . Let us prove that for some $p_1 \in \Sigma_1$ and $p_2 = h(p_1) \in \Sigma_2$ the tangent plane $T_{p_1}\Sigma_1$ is parallel to the tangent plane $T_{p_2}\Sigma_2$; this is stronger than required.

Arguing by contradiction, assume that such point does not exist. Then for each $p \in \Sigma_1$ there is unique line $\ell_p \ni p$ which is parallel to each of the the tangent planes $T_p\Sigma_1$ and $T_{h(p)}\Sigma_2$.

Note that the lines ℓ_p form a tangent line distribution over Σ_1 and ℓ_p is tangent to γ at any $p \in \gamma$.

Let D be the disc in Σ_1 bounded by γ . Consider the doubling of D in γ ; it is diffeomorphic to \mathbb{S}^2 . The line distribution ℓ lifts to a line distribution on the doubling; the later contradicts the hairy ball theorem.

Comments. This proof was suggested nearly simultaneously by Steven Sivek and user damiano, see [91].

Note that the same proof works in case if Σ_i are oriented open surfaces such that γ cuts a compact domain in each Σ_i .

There are examples of three disks Σ_1 , Σ_2 and Σ_3 with common closed curve γ such that there no triple of points $p_i \in \Sigma_i$ with parallel tangent plane. Such examples can be found among ruled surfaces, see [96].

24. Simple geodesic. Let γ be a two-sided infinite geodesic in Σ . The following is the key statement in the proof.

Claim. *The geodesic γ contains at most one simple loop.*

To prove the claim use the following observations.

- ◊ The total curvature ω of Σ can not exceed $2\cdot\pi$.

- ◇ If φ is the angle at the base of a simple geodesic loop then the total curvature surrounded by the loop equals to $\pi + \varphi$.

Once the claim is proved, note that if a geodesic γ has a self-intersection then it contains a simple loop. From above there is only one such loop; it cuts a disc from Σ and can go around it either clockwise or counterclockwise. This way we divide all the self-intersecting geodesics into two sets which we will call *clockwise* and *counterclockwise*.

Note that the geodesic $t \mapsto \gamma(t)$ is clockwise if and only if $t \mapsto \gamma(-t)$ is counterclockwise. The sets of clockwise and counterclockwise are open and the space of geodesics is connected. It follows that there are geodesics which neither clockwise nor counterclockwise; by the definition, these geodesics have no self-intersections.

Comment. The claim in the proof appears in the Bangert's paper [14, Corollary 2].

25. Long geodesic. Denote by a the area of the surface.


Cut the surface along a long closed simple geodesic γ . We get two discs with nonnegative curvature and large perimeter, say ℓ . Note that the area of each disc is bounded above by a .

Choose one of the discs D and equip it with intrinsic metric. Note that D is non-negatively curved in the sense of Alexandrov. Denote by p and q be the points in D which lie on the maximal distance from each other.

Fix $\varepsilon > 0$. Fix a geodesic $[pq]$ in D . Show that if ℓ is large enough in terms of ε the distance from any point in D to $[pq]$ is at most ε and the curvature of ε -neighborhood of p in D is at least $\pi - \varepsilon$.

By Gauss–Bonnet formula the total curvature of Σ is $4 \cdot \pi$. Since $\varepsilon > 0$ is arbitrary, we get that there are 4 point in Σ , each with curvature π and remaining part of Σ is flat.

It remains to show that any surface with this property is isometric to the surface of tetrahedron with equal opposite edges. To do this cut Σ along three geodesics which connect one singular point to the remaining three, develop the obtained flat surface on the plane and think.



Comments. The problem was suggested by Arseniy Akopyan.

26. Surrounded area. Denote by C_1 and C_2 the compact regions bounded by γ_1 and γ_2 correspondingly.

By Kirszbraun theorem, any short map $X \rightarrow \mathbb{R}^2$ defined on $X \subset \mathbb{R}^2$ can be extended to a short map on whole \mathbb{R}^2 . In particular there is a short map $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $f(\gamma_2(v)) = f(\gamma_1(v))$ for any $v \in \mathbb{S}^1$.

Note that $f(C_2) \supset C_1$. Whence the statement follows.

27. Corkscrew geodesic. An example can be found among surfaces of the convex hulls K of three broken lines, say γ_1 , γ_2 and γ_3 with 10000 edges each and common ends, say a and b . Think that the broken lines γ_i are very close to the line segment $[ab]$ and each edge of γ_i appears as an edge of the convex hull.

For the right choice of the broken lines the minimizing geodesic σ from a to b on the surface of K has to cross γ_i in the cyclic order in such a way that each edge of γ_1 is crossed. It follows that σ rotates around the line ab more than 1000 times.

Comments. The problem appears in the article [59] by Imre, Kuiperberg and Zamfirescu.

Comparison geometry

28. Totally geodesic hypersurface. Assume Σ is a totally geodesic embedded hypersurface in M . Without loss of generality, we can assume that Σ is connected.

The complement $M \setminus \Sigma$ has one or two connected components. Assume that the number of connected components is two, let us show that M is homeomorphic to sphere.

Cut M along Σ , you get two manifolds with geodesic boundaries. Prove that the distance function to the boundary is convex in the interior of each manifold. The distance function is not smooth but it can be smoothed while keeping it convex; this can be done by applying Greene–Wu Theorem ([43, Theorem 2]). In particular it has single critical point which is its maximum.

Applying Morse lemma, we get that each manifold is homeomorphic to a ball; hence M is homeomorphic to the sphere.

If $M \setminus \Sigma$ is connected, passing to a double cover of M we reduce the problem to the case when the number of connected components is two.

Comments. The problem was suggested by Peter Petersen.

29. Immersed convex hypersurface I. Observe first that any closed embedded locally convex hypersurface in a non-positively curved simply connected complete manifold bounds a convex region.

Let Σ be immersed locally convex hypersurface in M . Set

$$m = \dim \Sigma = \dim M - 1$$

Given a point in p on Σ denote by p_r the point on distance r from p which lies on the geodesic starting from p in the outer normal direction to Σ . For fixed $r \geq 0$, the points p_r sweep an immersed locally convex hypersurface which we denote by Σ_r .

Fix $z \in \Sigma$. Denote by S_r the sphere of radius r centered at z . Note that S_r is diffeomorphic to m -dimensional sphere.

Denote by d the diameter of Σ . Note that for all $r > 0$ any point on Σ_r lies on the distance at most d from S_r . Conclude that for large r the closest point projection $\varphi_r: \Sigma_r \rightarrow S_r$ is an immersion.

Since Σ is connected and $m \geq 2$, it follows that φ_r is a diffeomorphism for all large r .

By the observation above, Σ_r bounds a convex region for all large r . By open-close argument, the same holds for all $r \geq 0$. Hence the result follows.

Comments. The proof above appears in [7].

30. Immersed convex hypersurface II. Equip Σ with the induced intrinsic metric. Denote by κ the lower bound for principle curvatures of Σ . Note that we can assume that $\kappa > 0$.

Fix sufficiently small $\varepsilon = \varepsilon(M, \kappa) > 0$. Given $p \in \Sigma$ consider the lift $\tilde{h}_p: B_\varepsilon(p) \rightarrow T_{h(p)}$ along the exponential map $\exp_{h(p)}: T_{h(p)} \rightarrow M$. More precisely

1. Connect each point $q \in B_\varepsilon(p) \subset \Sigma$ to p by a the minimizing geodesic path $\gamma_q: [0, 1] \rightarrow \Sigma$
2. Consider the lifting $\tilde{\gamma}_q$ in $T_{h(p)}$; this is the curve such that $\tilde{\gamma}_q(0) = 0$ and $\exp_{h(p)} \circ \tilde{\gamma}_q(t) = \gamma_q(t)$ for any $t \in [0, 1]$.
3. Set $\tilde{h}(q) = \tilde{\gamma}_q(1)$.

Show any \tilde{h}_p has principle curvatures at least $\frac{\kappa}{2}$.

Use the same idea as in problem 22 to show that one can fix $\delta = \delta(M, \kappa) > 0$ such that the restriction of $\tilde{h}_p|_{B_\delta(p)}$ is injective. Conclude that the restriction $h|_{B_\delta(p)}$ is injective for any $p \in \Sigma$.


Now consider locally equidistant surfaces Σ_t in the inward direction for small t . The principle curvatures of Σ_t remain at least κ in the barrier sense. By the same argument as above, any δ -ball in Σ_t is embedded.

Applying open-close argument we get a one parameter family of locally convex locally equidistant surfaces Σ_t for defined in a maximal interval $[0, a]$; the surface Σ_a degenerates to a point, say p .

The point p can be taken as the image of the center of the ball for the needed embedding $B \looparrowright M$ and the surfaces Σ_t can be taken as the restrictions of the embedding to the spheres.

Comments. As you see on the picture the statement does not hold in the two-dimensional case.

The proof presented above was indicated in Gromov's lectures [47]; it was written rigorously by Eschenburg in [35]. A variation of this proof was ob-



tained independently by Andrews in [12]; instead of equidistant deformation it use so called *inverse mean curvature flow*.

31. Almgren's inequalities. Fix a geodesic m -dimensional sphere \mathbb{S}^m in \mathbb{S}^n .

Given $r \in (0, \frac{\pi}{2}]$, denote by U_r and \tilde{U}_r the tubular r -neighbourhood of Σ and \mathbb{S}^m in \mathbb{S}^n correspondingly.

Note that

$$(*) \quad U_{\frac{\pi}{2}} = \tilde{U}_{\frac{\pi}{2}} = \mathbb{S}^n.$$

Prove that for any $x \in \partial U_r$ we have

$$H_r(x) \geq \tilde{H}_r,$$

where $H_r(x)$ denotes the mean curvature of ∂U_r at point x and \tilde{H}_r its mean curvature of $\partial \tilde{U}_r$.

Set

$$\begin{aligned} a(r) &= \text{vol}_{n-1} \partial U_r, & \tilde{a}(r) &= \text{vol}_{n-1} \partial \tilde{U}_r, \\ v(r) &= \text{vol}_n U_r, & \tilde{v}(r) &= \text{vol}_n \tilde{U}_r. \end{aligned}$$

by coarea formula,

$$\frac{d}{dr} v(r) = a(r), \quad \frac{d}{dr} \tilde{v}(r) = \tilde{a}(r).$$

for almost all r . Note that

$$\begin{aligned} \frac{d}{dr} a(r) &\leq \int_{\partial U_r} H_r(x) \cdot d_x \text{vol}_{n-1} \leq \\ &\leq a(r) \cdot \tilde{H}_r \end{aligned}$$

and

$$\frac{d}{dr} \tilde{a}(r) = \tilde{a}(r) \cdot \tilde{H}_r.$$

It follows that

$$\frac{v''(r)}{v(r)} \leq \frac{\tilde{v}''(r)}{\tilde{v}(r)}$$

for almost all r . Therefore

$$v(r) \leq \frac{\text{area } \Sigma}{\text{area } \mathbb{S}^m} \cdot \tilde{v}(r)$$

for any $r > 0$.

According to (*),

$$v(\frac{\pi}{2}) = \tilde{v}(\frac{\pi}{2}) = \text{vol } \mathbb{S}^n.$$

Hence the result follows.

Comments. This problem is the most geometric part in the proof of Almgren's isoperimetric inequality [11]. The argument presented here is very similar to the proof of Gromov–Levy isometric inequality given in the Gromov's appendix to [77].

32. Hypercurve. Fix $p \in M$. Denote by s the second fundamental form of M at p ; it is a symmetric bi-linear form on the tangent space $T_p M$ of M with values in the normal space $N_p M$ to M , see page 85. Note that the normal space $N_p M$ is two-dimensional.

Prove that if sectional curvature of M is positive, then

$$(*) \quad \langle s(X, X), s(Y, Y) \rangle > 0$$

for any pair of nonzero vectors $X, Y \in T_p M$.

Show that (*) implies that there is an orthonormal basis e_1, e_2 in $N_p M$ such that the real-valued quadratic forms

$$s_1(X, X) = \langle s(X, X), e_1 \rangle, \quad s_2(X, X) = \langle s(X, X), e_2 \rangle$$

are positive definite.

Note that the curvature operators R_1 and R_2 defined by the following identity

$$R_i(X \wedge Y), V \wedge W = s_i(X, W) \cdot s_i(Y, V) - s_i(X, V) \cdot s_i(Y, W)$$

are positive. Finally, note that $R_1 + R_2$ is the curvature operator of M at p .

Comments. The problem appears in Weinstein's paper [113].

Note that follows from [73]/[20] it follows that that the universal cover of M is homeomorphic/diffeomorphic to a standard sphere.

33. Horosphere. Set $m = \dim \Sigma = \dim M - 1$.

Let $b: M \rightarrow \mathbb{R}$ be the Busemann function such that $\Sigma = b^{-1}(\{0\})$. Set $\Sigma_r = b^{-1}(\{r\})$, so $\Sigma_0 = \Sigma$.

Let us equip each Σ_r with induced Riemannian metric. Note that all Σ_r have bounded curvature. In particular, unit ball in Σ_r has volume bounded above by universal constant, say v_0 .

Given $x \in \Sigma$ denote by γ_x the (necessary unique) unit-speed geodesic such that $\gamma_x(0) = x$ and $b(\gamma_x(t)) = t$ for any t . Consider the map $\varphi_r: \Sigma \rightarrow \Sigma_r$ defined as $\varphi_r: x \mapsto \gamma_x(r)$.

Notice that φ_r is a bi-Lipschitz map with the Lipschitz constants $e^{a \cdot r}$ and $e^{b \cdot r}$. In particular, the ball of radius R in Σ is mapped by φ_r to a ball of radius $e^{a \cdot r} \cdot R$ in Σ_r . Therefore

$$\text{vol}_m B(x, R)_\Sigma \leq e^{m \cdot b \cdot r} \cdot \text{vol}_m B(x, e^{a \cdot r} \cdot R)_{\Sigma_r}$$

for any $R, r > 0$. Applying this formula in case $e^{a \cdot r} \cdot R = 1$ implies that

$$\text{vol}_m B(x, R)_\Sigma \leq v_0 \cdot R^{m \cdot \frac{b}{a}}.$$

Comment. The problem was suggested by Vitali Kapovitch.

There are examples of horospheres as above with degree of polynomial growth higher than m . For example, consider the horosphere Σ in the complex hyperbolic space of real dimension 4. Clearly $m = \dim \Sigma = 3$ but the degree of its volume growth is 4. The later follows since Σ admits has left-invariant metric on the Heizenber's group.

34. Minimal spheres. Choose a pair of sufficiently close minimal spheres Σ and Σ' , say assume that the distance a between Σ and Σ' is strictly smaller than the injectivity radius of the manifold. Note that in this case there is a bijection $\Sigma \rightarrow \Sigma'$, which will be denoted by $p \mapsto p'$ such that the distance $|p - p'| = a$ for any $p \in \Sigma$.

Let $\iota_p: T_p \rightarrow T_{p'}$ be the parallel translation along the (necessary unique) minimizing geodesic from p to p' . Use hairy ball theorem to show that there is a pair (p, p') such that $\iota_p(T_p \Sigma) = T_{p'} \Sigma'$.

Consider pairs of unit-speed geodesics α and α' in Σ and Σ' which start at p and p' correspondingly and go in the parallel directions. Set $\ell(t) = |\alpha(t) - \alpha'(t)|$.

Use the second variation formula to show that $\ell''(0)$ has negative average for all directions from p . In particular $\ell''(0) < 0$ for a pair α and α' as above. It follows that there are points $v \in \Sigma$ near p and $v' \in \Sigma'$ near p' such that

$$|v - v'| < |p - p'|;$$

the later leads to a contradiction.

Comment. The problem was suggested by D. Burago, it is related to Frankel's theorem on minimal surfaces; see [39].

It seems pleasurable that a compact positively curved 4-dimensional manifold can not contain a pair of equidistant spheres; if such pair exists, then the distance between them is at least the injectivity radius of the manifold.

Here is a short list of classical problems with similar solutions:

- ◇ (Synge's problem [103]) *Any compact even-dimensional orientable manifold with strictly positive sectional curvature is simply connected.*
- ◇ (Frankel's problem [39]) *Show that any two compact minimal hypersurfaces in a Riemannian manifold with positive Ricci curvature must intersect.*
- ◇ (Bochner's problem [19].) *Let (M, g) be a closed Riemannian manifold with negative Ricci curvature. Prove that (M, g) does not admit an isometric \mathbb{S}^1 -action.*

The problem 35 can be considered as further development of this idea.

35. Totally geodesic immersion. Set $n = \dim N$ and $m = \dim M$.

Fix a smooth increasing concave function φ . Consider the function $f = \varphi \circ \text{dist}_N$. Note that if f is smooth at x the the Hessian $\text{Hess}_x f$ has at least $n + 1$ negative eigenvalues.

Moreover, at any point $x \notin \iota(N)$ the same holds in the barrier sense; i.e., there is a smooth function $h \geq f$ defined on M of x such that $h(x) = f(x)$ and $\text{Hess}_x f$ has at least $n + 1$ negative eigenvalues.

Use that $m < 2 \cdot n$ and the property to prove the following analog of Morse lemma for f .

Claim. *Given $x \notin \iota(N)$ there is a neighborhood $U \ni x$ such that the set*

$$U_- = \{ z \in U \mid f(z) < x \}$$

is simply connected.

Since M is simply connected, we can shrink by a disc $f_0: \mathbb{D} \rightarrow M$ any closed curve in $\iota(N)$. According to the claim, there is a homotopy $f_t: \mathbb{D} \rightarrow M$, $t \in [0, 1]$ such that $f_t(\partial\mathbb{D}) \subset \iota(N)$ for any t and $f_1(\mathbb{D}) \subset \iota(N)$. It follows that $\iota(N)$ is simply connected.

Finally, note that if $\iota(N)$ is simply connected then ι has to be an embedding.

Comments. The statement was proved by Fang, Mendonça and Rong in [37]. The main idea in this proof was discovered by Wilking, see [115].

36. Positive curvature and symmetry. Let M be a 4-dimensional Riemannian manifold with isometric \mathbb{S}^1 -action. Consider the quotient space $X = M/\mathbb{S}^1$.

Note that X is a positively curved 3-dimensional Alexandrov space; see [9] if in doubt. In particular the angle $\angle[x \frac{y}{z}]$ between geodesics any two $[xy]$ and $[xz]$ is defined and

$$(*) \quad \angle[x \frac{y}{z}] + \angle[y \frac{z}{x}] + \angle[z \frac{x}{y}] > \pi.$$

for any triangle $[xyz]$ formed by the minimizing geodesics $[xy]$, $[yz]$ and $[zx]$ in X .

Assume $p \in X$ corresponds to a fixed point of \mathbb{S}^1 -action. Show that for any three geodesics $[px]$, $[py]$ and $[pz]$ in X we have

$$(**) \quad \angle[x_y^p] + \angle[y_z^p] + \angle[z_x^p] \leq \pi.$$

Arguing by contradiction, assume that there are 4 fixed points q_1 , q_2 , q_3 and q_4 . Connect each pair $q_i \neq q_j$ by a minimizing geodesic $[q_i q_j]$.

Denote by ω the sum of all 12 angles of the type $\angle[q_i q_j q_k]$. By (*), we have

$$\omega > 4 \cdot \pi.$$

By (**) at each vertex q_i , we have

$$\omega \leq 4 \cdot \pi,$$

a contradiction.

Comment. The problem appears in the paper of Hsiang and Kleiner [58]. The connection of this proof to Alexandrov geometry was noticed by Grove in [49]. An interesting development of this idea is given by Grove and Wilking in [50].

37. Scalar curvature vs. injectivity radius. We will prove that if the injectivity radius of the manifold (M, g) is at least π then the average of sectional curvatures on (M, g) is at most 1. This is stronger than required.

Fix a point $p \in M$ and two orthonormal vectors $U, V \in T_p M$. Consider the geodesic γ in M such that $\dot{\gamma}(0) = U$.

Set $U_t = \dot{\gamma}(t) \in T_{\gamma(t)}$ and let $V_t \in T_{\gamma(t)}$ be the parallel translation of $V = V_0$ along γ .

Consider the field $W_t = \sin t \cdot V_t$ on γ . Set

$$\gamma_\tau(t) = \exp(\tau \cdot W_t), \quad \ell(\tau) = \text{length}(\gamma_\tau|_{[0, \pi]}), \quad q(U, V) = \ell''(0).$$

Note that

$$(*) \quad q(U, V) = \int_0^\pi [(\cos t)^2 - K(U_t, V_t) \cdot (\sin t)^2] \cdot dt,$$

where $K(U, V)$ denotes the curvature in the sectional direction spanned by U and V .

Denote by s and k the average values of $q(U, V)$ and $K(U, V)$.

Since any geodesics of length π is minimizing, we get $q(U, V) \geq 0$ for any pair of orthonormal vectors U and V . In particular, $s \geq 0$.

Let us take average value of right hand side in (*). By Liouville's theorem we can switch the order of integrals; therefore $s = \frac{\pi}{2} \cdot (1 - k)$. Hence the result follows.

38. Almost flat manifold. First prove that for given $\varepsilon > 0$, there is big enough n and $n \times n$ integer matrix A such that all its eigenvalues are ε -close to 1.

Consider $(n + 1)$ -dimensional manifold S obtained from $\mathbb{T}^n \times [0, 1]$ by gluing $\mathbb{T}^n \times 0$ to $\mathbb{T}^n \times 1$ along the map given by A .

Assuming that ε is small, show that S admits a metric with curvature and diameter sufficiently small.

Comment. This example was constructed by Guzhvina, see [52]. The celebrated theorem of Gromov states that there are no such examples of fixed dimension.

It is expected that for small enough $\varepsilon > 0$, a manifold of any dimension with diameter ≤ 1 and sectional curvature at most ε has nontrivial fundamental group.

✕ 39. Lie group. Consider the product $(G, g_0) \times (G, g_1)$; it is the Lie group $G \times G$ with a left invariant metric.

Consider the diagonal subgroup

$$\Delta = \{(g, g) \in G \times G\}.$$

The quotient space $(G, g_0) \times (G, g_1) / \Delta$ is isometric to $(G, ???g_0 + g_1)$. Hence the result follows.

Comment. The construction described in this solution is a partial case of so called “Cheeger’s trick”.

40. Polar points. Fix a unit-speed geodesic γ which starts at p ; i.e., $\gamma(0) = p$. Set $p^* = \gamma(\pi)$. It remains to prove that p^* is a solution.

Alternative proof. Assume contrary; i.e., for any $x \in M$ there is a point x' such that

$$|x - x'|_g + |p - x'|_g > \pi.$$

Show that there is a continuous map $x \mapsto x'$ such that the above inequality holds for any x .

Fix sufficiently small $\varepsilon > 0$. Prove that the set $W_\varepsilon = M \setminus B_\varepsilon(p)$ is homeomorphic to a ball and the map $x \mapsto x'$ sends W_ε into itself.

It remains to apply Brouwer’s fixed-point theorem.

Comments. The problem with the first proof was discussed by Milka’s in [75].

✕ 41. Deformation to a product. Denote by Γ the fundamental group of M ; if Γ is finite then the universal cover will do the trick.

Let (\tilde{M}, \tilde{g}) be universal cover of (M, g) with induced Riemannian metric. The space (\tilde{M}, \tilde{g}) is isometric to a product $\mathbb{R}^k \times K$, where K is a compact Riemannian manifold.

Denote by G the isometry group of K . Given a continuous one parameter family of homomorphisms $\varphi_t: \mathbb{R}^k \rightarrow G$, consider the diffeomorphism

$$\Phi_t: (x, k) \mapsto (x, \varphi_t(x) \cdot k)$$

of $\mathbb{R}^k \times K$ to itself and denote by \tilde{g}_t pullback of \tilde{g} via Φ_t , so

$$\Phi_t: (\mathbb{R}^k \times K, \tilde{g}_t) \rightarrow (\mathbb{R}^k \times K, \tilde{g})$$

is an isometry.

If \tilde{g}_t is Γ -invariant, the quotient $(\mathbb{R}^k \times K, \tilde{g}_t)/\Gamma$ is

It remains to find the one parameter family of homomorphisms $\varphi_t: \mathbb{R}^k \rightarrow G$ so that φ_0 is the trivial homomorphism, \tilde{g}_t is Γ -invariant for all t and $(M, g_1) = (\mathbb{R}^k \times K, \tilde{g}_1)/\Gamma$ admits a finite cover isometric to product of flat k -dimensional torus with K .

Comment. The problem appears in Wilking's paper [114].

42. Isometric section. Arguing by contradiction, assume $\iota: M \rightarrow W$ is an isometric section, it makes possible to treat M as a submanifold in W .

Denote by $\nu_p M$ the sphere of unit normal vectors to M at $p \in M$. Given $v \in \nu_p$ and real value k , set

$$p^{k \cdot v} = s \circ \exp_{\iota(p)}(k \cdot v).$$

Note that

$$(*) \quad p^0 = p \text{ for any } p \in M.$$

Fix sufficiently small $\delta > 0$. By Rauch comparison, if $w \in \nu_q$ is the parallel translation of $v \in \nu_p$ along the minimizing geodesic from p to q in M then

$$(**) \quad |p^{k \cdot v} - q^{k \cdot w}|_M < |p - q|_M$$

assuming $|k| \leq \delta$. More generally, the same comparison implies that

$$(***) \quad |p^{k \cdot v} - q^{k' \cdot w}|_M^2 < |p - q|_M^2 + (k - k')^2$$

assuming $|k|, |k'| \leq \delta$.

Choose p and $v \in \nu_p$ so that $r = |p - p^{\delta \cdot v}|$ takes the maximal possible value. From $(**)$ it follows that $r > 0$.

Let γ be the extension of unit-speed minimizing geodesic from p_v to p ; denote by v_t the parallel translation of v to $\gamma(t)$ along γ .

We can choose the parameter of γ so that $p = \gamma(0)$, $p^v = \gamma(-r)$. Set $p_n = \gamma(n \cdot r)$, so $p = p_0$ and $p^v = p_{-1}$, Fix large integer N and set $w_n = (1 - \frac{n}{N}) \cdot v_{n \cdot r}$ and $q_n = p_n^{w_n}$.

From (**), there is a constant C independent of N such that

$$|q_k - q_{k+1}| < r + \frac{C}{N^2} \cdot \delta^2.$$

Therefore

$$|q_{k+1} - p_{k+1}| > |q_k - p_k| - \frac{C}{N^2} \cdot \delta^2.$$

By induction, we get

$$|q_N - p_N| > r - \frac{C}{N} \cdot \delta^2.$$

Since N is large we get

$$|q_N - p_N| > 0.$$

By (*) we get $q_N = p_N^0 = p_N$, a contradiction.

Comment. This proof is the core of Perelman's proof of Soul conjecture, see [90].

43. Minkowski space. Fix increasing function $\varphi: (0, r) \rightarrow \mathbb{R}$ such that $\varphi'' + (n-1) \cdot (\varphi')^2 + C = 0$. Note that if $\text{Ric}_M \geq C$ then the function $x \mapsto \varphi(|p - x|_M)$ is subharmonic. In particular, for arbitrary array of points p_i and positive reals λ_i the function

$$f(x) = \sum_i \lambda_i \cdot \varphi(|p_i - x|_M).$$

It remains to show that if $p \neq 2$ then there is an array points p_i and positive reals λ_i such that the function

$$f(x) = \sum_i \lambda_i \cdot \varphi(|p_i - x|_M)$$

has strict local minimum.

Comment. The argument given here is very close to the proof of Abresch–Gromoll inequality, see [3]. An alternative solution of this problem can be build on almost splitting theorem of Cheeger and Colding, see [31].

✱ **44. Curvature hollow.** Construct a metric that the connected sum $M = \mathbb{R}^3 \# \mathbb{S}^2 \times \mathbb{S}^1$ admits a metric which is flat outside a compact set and has non positive scalar curvature. Further, note that such metric can be constructed in such a way that it has a closed geodesic γ with trivial holonomy and with constant negative curvature in its a tubular neighborhood.

Cut the tubular neighbourhood $D^2 \times \mathbb{S}^1$ of γ , prepare a metric g on $\mathbb{S}^1 \times D^2$ with negative scalar curvature which is identical to the original metric near the boundary. The needed patch $(\mathbb{S}^1 \times D^2, g)$ can be found among wrap products $\mathbb{S}^1 \times_f D^2$.

Note that after the surgery we get a manifold diffeomorphic to \mathbb{R}^3 , it flat outside of compact set and has negative scalar curvature somewhere.

Comments. This construction was given by Lohkamp in [1]. Hollows with negative Ricci curvature also exist, see ????. On the other hand there are no hollows with positive scalar curvature; the later follows from Positive Mass Conjecture.

✂ **45.** *If hemisphere then sphere.*

Comments. The problem appears in paper of Hang and Wang [53]. The problem is still interesting if instead of the first condition one has that sectional curvature ≥ 1 . If instead of first condition one only has that scalar curvature $\geq m \cdot (m-1)$, then the conclusion does not hold; it was conjectured by Min-Oo (1995) and disproved in [23]

46. *Flat coordinate planes.* Fir $\varepsilon > 0$ such that there is unique geodesic between any two points on distance $< \varepsilon$ from the origin of \mathbb{R}^3 .

Consider three points a, b and c on the coordinate lines which are ε -close to the origin.

Prove that the angles of the triangle $\triangle abc$ coincide with its model angles. It follows that there is a flat geodesic triangle in (\mathbb{R}^3, g) with vertex at a, b and c .

Use the family of constructed flat triangles to show that at any x point in the $\frac{\varepsilon}{10}$ -neighborhood of the origin the sectional curvature vanish in an open set of sectional directions. The later implies that the curvature is identically zero in this neighborhood.

Moving the origin and apply the same argument we get that the curvature is identically zero everywhere. Hence the result follows.

Comment. This problem appears in the paper of Panov and me [88]; it is based on an observation discovered by Buyalo in [28].

47. *Two-convexity.* Assume W is a hypersurface of needed type.

Morse-style solution. Let us equip \mathbb{R}^4 with (x, y, z, t) -coordinates.

Consider a generic linear function $\ell: \mathbb{R}^4 \rightarrow \mathbb{R}$ which is close to the sum of coordinates $x+y+z+t$. Note that ℓ has non-degenerate critical points on W and all its critical values are different.

Consider the sets

$$W_s = \{ w \in \mathbb{R}^4 \setminus K \mid \ell(w) < s \}.$$

Note that W_{-1000} contains a closed curve, say α , which is contactable in $\mathbb{R}^4 \setminus K$, but not constructable in the set W_{-1000} .

Set s_0 to be the infimum of the values s such that the α is contactable in W_s .

Note that s_0 is a critical value of ℓ on W ; denote by p_0 the corresponding critical point. By 2-convexity of $\mathbb{R}^4 \setminus K$, the index of p_0 has to be at most 1. On the other hand, since the disc hangs at this point, its index has to be at least 2, a contradiction.

Alexandrov-style proof. Fix a constant Riemannian metric g on \mathbb{R}^4 . According to the main result of Alexander Bishop and Berg in [8], $X_g = (\mathbb{R}^4 \setminus (\text{Int } K), g)$ has nonpositive curvature in the sense of Alexandrov. In particular the universal cover of \tilde{X}_g of X_g is a CAT[0] space.

By rescaling g and passing to the limit we obtain that universal Riemannian cover Z_g of (\mathbb{R}^4, g) branching in the coordinate planes is a CAT[0] space. Show that Z_g is CAT[0] space if and only if the two planes are orthogonal with respect to g ; the later leads to a contradiction.

Comments. Note that the closed 1-neighborhood of these two planes has two-convex complement, but the boundary of this neighborhood is not smooth.

The Morse-style is closely related to Gromov's lectures [47, §1/2].

Curvature free differential geometry

48. Minimal foliation. First show that there is a self-dual harmonic 2-form on $(\mathbb{S}^2 \times \mathbb{S}^2, g)$; i.e., a 2-form ω such that $d\omega = 0$ and $\star\omega = \omega$, where \star denotes the Hodge star operator.

Fix $p \in \mathbb{S}^2 \times \mathbb{S}^2$. Use the identity $\star\omega_p = \omega_p$ to show that there is a real number λ_p and the isometry $J_p: T_p \rightarrow T_p$ such that $J_p \circ J_p = -\text{id}$ and $\omega(X, Y) = \lambda_p \cdot g(X, J_p Y)$ for any $X, Y \in T_p$.

Consider canonical symplectic form ω_0 on $\mathbb{S}^2 \times \mathbb{S}^2$; that is sum of pullbacks of volume forms on \mathbb{S}^2 for the two projections $\mathbb{S}^2 \times \mathbb{S}^2 \rightarrow \mathbb{S}^2$. Note that for the canonical metric on $\mathbb{S}^2 \times \mathbb{S}^2$, the form ω_0 is harmonic and self-dual. Since g is close to the standard metric, we can assume that ω is close to ω_0 . In particular $\lambda_p \neq 0$ for any $p \in \mathbb{S}^2 \times \mathbb{S}^2$.

It follows that ω defines symplectic structure on $\mathbb{S}^2 \times \mathbb{S}^2$ and J is its pseudocomplex structure. It remains to take the pseudoholomorphic surfaces in the homology classes of $\mathbb{S}^2 \times x$ and $x \times \mathbb{S}^2$ for some fixed $x \in \mathbb{S}^2$.

49. Loewner's theorem. Denote by λ the conformal factor of g ; i.e., $g = \lambda^2 \cdot g_{\text{can}}$.

Denote by s the average of g -lengths of the lines in \mathbb{RP}^n . Prove that

$$\ell \leq s = \pi \cdot \oint_{\mathbb{RP}^n} \lambda \cdot d \operatorname{vol}_{\operatorname{can}},$$

where $\operatorname{vol}_{\operatorname{can}}$ denotes the volume for g_{can} and \oint denoted the average value.

Note that

$$\operatorname{vol}(\mathbb{RP}^n, g) = \operatorname{vol}(\mathbb{RP}^n, g_{\operatorname{can}}) \cdot \oint_{\mathbb{RP}^n} \lambda^n \cdot d \operatorname{vol}_{\operatorname{can}}.$$

By Hölder's inequality, we have

$$\left(\oint_{\mathbb{RP}^n} \lambda \cdot d \operatorname{vol}_{\operatorname{can}} \right)^n \leq \oint_{\mathbb{RP}^n} \lambda^n \cdot d \operatorname{vol}_{\operatorname{can}}.$$

Hence the result follows.

50. Convex function vs. finite volume. Assume contrary; i.e., there is a complete Riemannian manifold M with finite volume which admits a convex function f .

Denote by $\tau: T^1M \rightarrow M$ the unit tangent bundle over M . Clearly $\operatorname{vol} T^1M$ is finite.

Note that there is a nonempty bounded open set $U \subset T^1M$ such that $df(u) > \varepsilon$ for any $u \in U$ and some fixed $\varepsilon > 0$.

Denote by φ^t the geodesic flow on T^1M . Given $u \in U$, consider the function $h: t \mapsto f \circ \tau \circ \varphi^t(u)$. Note that $h'(t) > \varepsilon$ for any $t \geq 0$.

Prove that there is an infinite sequence of positive reals t_1, t_2, \dots such that

$$\varphi^{t_i}(U) \cap \varphi^{t_j}(U) = \emptyset$$

if $i \neq j$. The later implies that $\operatorname{vol} T^1M = \infty$, a contradiction.

Comment. The problem appears in Yau's paper [119].

51. Besikovitch inequality. Set

$$A_i = \{ (x_1, x_2, \dots, x_n) \in [0, 1]^n \mid x_i = 0 \}.$$

Consider functions $f_i: [0, 1]^n \rightarrow \mathbb{R}$ defined by $f_i(x) = \operatorname{dist}_{A_i} x$. Prove that Jacobian of the map $\mathbf{f}: [0, 1]^n \rightarrow \mathbb{R}^n$ defined as

$$\mathbf{f}: x \mapsto (f_1(x), f_2(x), \dots, f_n(x))$$

is at most 1 and $\mathbf{f}([0, 1]^n) \supset [0, 1]^n$. Hence the result follows.

Comments. This inequality appears in Besicovitch's paper [16].

52. Distant involution. Given $\varepsilon > 0$, construct a disc D in the plane with

$$\text{length } \partial D < 10 \quad \text{and} \quad \text{area } D < \varepsilon$$

which admits an continuous involution ι such that

$$|\iota(x) - x| \geq 1$$

for any $x \in \partial D$. An example of D can be guessed from the picture.

Take the product $D \times D \subset \mathbb{R}^4$; it is homeomorphic to a 4-dimensional ball. Note that

$$\text{vol}_3[\partial(D \times D)] = 2 \cdot \text{area } D \cdot \text{length } \partial D < 20 \cdot \varepsilon.$$

The boundary $\partial(D \times D)$ homeomorphic to \mathbb{S}^3 and the restriction of the involution $(x, y) \mapsto (\iota(x), \iota(y))$ has the needed property.

It remains to smooth $\partial(D \times D)$.

Comments. This example was discovered by Croke in [32].

It is instructive to show that for \mathbb{S}^2 such thing is not possible.

53. Normal exponential map. Assume contrary; i.e., there is a point $p \in M$ such that the image of normal exponential map to N does not touch ε -neighborhood of p .

Show that given $R > 0$ there is $\delta > 0$ such that if $x \in N$ and $|p - x|_M < R$ then there is a unit speed curve in N which moves to p with velocity at least δ . (In fact, the value δ depends on ε , R and the curvature bounds in the R -ball centered at p .)

Following this curve for sufficient time brings us to p ; i.e., $p \in N$, a contradiction.

Comments. The problem was suggested by Alexander Lytchak.



From the picture, you should guess an example of immersion $\iota: \mathbb{R} \looparrowright \mathbb{R}^2$ such that one point does not lie in the image of the corresponding normal exponential.

54. Symplectic squeezing in the torus. Equip \mathbb{R}^4 with (x_1, y_1, x_2, y_2) -coordinates so that

$$\omega = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$$

is the symplectic form.

The embedding will be given as a composition of a linear symplectomorphism λ with the quotient map $\varphi: \mathbb{R}^4 \rightarrow \mathbb{T}^2 \times \mathbb{R}^2$ by the integer (x_1, y_1) -lattice. Clearly $\varphi \circ \lambda$ preserves the symplectic structure, it remains to find λ such that the restriction $\varphi \circ \lambda|_\Omega$ is injective.



Without loss of generality, we can assume that Ω is a ball centered at the origin. Choose an oriented 2-dimensional subspace V subspace of \mathbb{R}^4 such that the integral of ω over $\Omega \cap V$ is small positive number, say $\frac{\pi}{4}$.

Note that there is a linear symplectomorphism λ which maps planes parallel to V to planes parallel to the (x_1, y_1) -plane, and that maps the disk $V \cap \Omega$ to a disk. It follows that the intersection of $\lambda(\Omega)$ with any plane parallel to the (x_1, y_1) -plane is equal to a disk of radius at most $\frac{1}{2}$. In particular $\varphi \circ \lambda|_{\Omega}$ is injective.

Comments. This construction is given by Guth in [51] and attributed to Leonid Polterovich.

Note that according to Gromov's theorem, analogous statement with $\mathbb{C} \times \mathbb{D}$ as the target does not hold, here $\mathbb{D} \subset \mathbb{C}$ is the open disc with induced symplectic structure. In particular, the projection of $\lambda(\Omega)$ as above to (x_1, y_1) -plane can not be made arbitrary small.

55. Diffeomorphism test. Since N is simply connected, it is sufficient to show that $f: M \rightarrow N$ is a covering map.

Note that f is an open immersion. Let h be the pullback metric on M for $f: M \rightarrow N$. Clearly $h \geq g$. In particular (M, h) is complete and the map $f: (M, h) \rightarrow N$ is a local isometry.

It remains to prove that any local isometry between complete connected Riemannian manifolds of the same dimension is a covering map.

56. Volume of tubular neighborhoods. Let us denote by NM and TM the normal and tangent bundle of M in \mathbb{R}^n .

Consider the normal exponential map $\exp_M: NM \rightarrow \mathbb{R}^n$ and denote by J_V its Jacobian at $V \in N_p M$. Note that for all small $\varepsilon > 0$, we have

$$(*) \quad \text{vol } B_\varepsilon(M) = \int_M d_p \text{vol}_m \cdot \int_{B_r \subset N_p M} J_V \cdot d_V \text{vol}_{n-m}.$$

Set $m = \dim M$. Given $p \in M$, denote by $s_p: T_p \times T_p \rightarrow N_p$ the second fundamental form of M . Recall that the curvature tensor of M at p can be expressed the following way

$$R_p(X \wedge Y, V \wedge W) = \langle s_p(X, W), s_p(Y, V) \rangle - \langle s_p(X, V), s_p(Y, W) \rangle.$$

Given $V \in N_p M$, express J_V in terms of $\langle s_p(X, Y), V \rangle$. Show that for small r the integral


$$v(r) = \int_{B_r \subset N_p M} J_V \cdot d_V \text{vol}_{n-m}$$

is a polynomial of r and its coefficients can be expressed in terms of the curvature tensor R_p .

It follows that the right hand side in (*) can be expressed in terms of curvature tensor of M . The problem follows since the curvature tensor can be expressed in terms of metric tensor of M .

Comments. The formula for volume of tubular neighborhood of submanifolds used in the proof was described by Weyl in [112].

57. Disc. Show that given a positive integer n one can construct a tree T embedded into the disc such that any homotopy of the boundary of the disc to a point pass through a curve which intersects n different edges. (For the tree on the diagram $n = 3$.)



Fix small $\varepsilon > 0$, say $\varepsilon = \frac{1}{10}$. Consider the disc with embedded tree T as above. We will construct a metric on the disc with diameter and length of its boundary below 1 such that the distance between any two edges of T of without common vertex is at least ε .

To construct such a metric, first fix a metric on the cylinder $\mathbb{S}^1 \times [0, 1]$ such that

- ◊ The ε -neighborhoods of the boundary components are product metrics.
- ◊ Any vertical segment $x \times [0, 1]$ has length $\frac{1}{2}$.
- ◊ One of the boundary component is ε -long.
- ◊ The other boundary component has length $2 \cdot m \cdot \varepsilon$, where m is the number of edges in T .

Equip T with a metric so that each edge has length ε and glue the long boundary component of the cylinder to T by piecewise isometry so that the resulting space is homeomorphic to disc and the tree corresponds to it-self.

According to the first construction, for any null-homotopy of the boundary the least length is at least $n \cdot \frac{\varepsilon}{10}$. The obtained metric is not Riemannian, but is easy to smooth. Since n is arbitrary the result follows.

Comments. The problem appears in the paper [38] by Frankel and Katz.

58. Shortening homotopy. Set $p = \gamma_0(0)$ and $\ell_0 = \text{length } \gamma_0$.

By compactness argument, there exists $\delta > 0$ such that no geodesic loops based at p with has length in the interval $(L - \delta, L + \delta + \delta]$.

Assume $\ell_0 \geq L + \delta$. Choose $t_0 \in [0, 1]$ such that

$$\text{length } \gamma_0|_{[0, t_0]} = L + \delta$$

Let σ be a the minimizing geodesic from $\gamma(t_0)$ to p . Note that γ_0 is homotopic to the joint

$$\gamma'_0 = \gamma_0|_{[0,t_0]} * \sigma * \bar{\sigma} * \gamma|_{[t_0,1]},$$

where $\bar{\sigma}$ denotes the backward parametrization of σ .

Consider the loop λ_0 at p formed by joint of $\gamma|_{[0,t_0]}$ and σ . Applying a curve shortening process to λ_0 , we get a curve shortening homotopy λ_t rel. its ends from the loop λ_0 to a geodesic loop λ_1 at p . From above,

$$\text{length } \lambda_1 \leq L - D.$$

The joint $\gamma_t = \lambda_t * \bar{\sigma} * \gamma|_{[t_0,1]}$ is a homotopy from γ'_0 to an other curve γ_1 . From the construction it is clear that

$$\begin{aligned} \text{length } \gamma_t &\leq \text{length } \gamma_0 + 2 \cdot \text{length } \sigma \leq \\ &\leq \text{length } \gamma_0 + 2 \cdot D \end{aligned}$$

for any $t \in [0, 1]$ and

$$\begin{aligned} \text{length } \gamma_1 &= \text{length } \lambda_1 + \text{length } \sigma + \text{length } \gamma|_{[t_0,1]} \leq \\ &\leq L - D + D + \text{length } \gamma - (L + \delta) = \\ &= \ell_0 - \delta. \end{aligned}$$

Repeating the procedure few times we get we get curves $\gamma_2, \gamma_3, \dots, \gamma_n$ joint by the needed homotopies so that $\ell_{i+1} \leq \ell_i - \delta$ and $\ell_n < L + \delta$, where $\ell_i = \text{length } \gamma_i$.

If $\ell_n \leq L$, we are done. Otherwise repeat the argument once more for $\delta' = \ell_n - L$.

Comments. The problem discussed by Nabutovsky and Rotman in [81]. The reader might wonder whether there are manifolds which satisfy the above condition for some L . Such examples can be found among the Zoll spheres; the later is shown by Balachev, Croke and Katz in [13].

59. *Geodesic hypersurface.* Let r be the maximal distance from points in W to M .

Fix a fine triangulation of W so that M becomes a subcomplex. Say, let us assume that the diameter of each simplex in τ is less than ε . We can assume that τ is a barycentric subdivision of an other triangulation, so all the vertices of τ can be colored into colors $(0, \dots, m+1)$ such that the vertices of each simplex get different colors. Denote by τ_i the i -dimensional subcomplex of τ formed by all vertices with colors $(0, \dots, i)$.

For each vertex v in τ choose a point $v' \in M$ on the distance $\leq r$. Note that if v and w are the vertices of one simplex then

$$|v' - w'|_M < 2 \cdot r + \varepsilon.$$

If $\frac{i_M}{2 \cdot (m+1)} > r$, we can assume that $\varepsilon < \frac{i_M}{2 \cdot (m+1)} - r$. It makes possible to extend the map $v \mapsto v'$ to a continuous map $W \rightarrow M$. The map is already defined on τ_0 , using the cone construction we can extend it to τ_1 ; we can do this since the distance between vertices in one simplex are below injectivity radius of M . Repeat the cone construction recursively, to extend the map $\tau_{m+1} = \tau$ (some distance estimate are needed here).

It follows that fundamental class of M vanish in the homology ring of M , a contradiction.

Comment. This problem is a simplified version of Gromov's bound on filling radius given in [45].

Metric geometry

60. Noncontracting map. Given any pair of point $x_0, y_0 \in K$, consider two sequences x_0, x_1, \dots and y_0, y_1, \dots such that $x_{n+1} = f(x_n)$ and $y_{n+1} = f(y_n)$ for each n .

Since K is compact, we can choose an increasing sequence of integers n_k such that both sequences $(x_{n_i})_{i=1}^\infty$ and $(y_{n_i})_{i=1}^\infty$ converge. In particular, both of these sequences are Cauchy; i.e.,

$$|x_{n_i} - x_{n_j}|_K, |y_{n_i} - y_{n_j}|_K \rightarrow 0 \quad \text{as} \quad \min\{i, j\} \rightarrow \infty.$$

Since f is noncontracting, we get

$$|x_0 - x_{|n_i - n_j|}| \leq |x_{n_i} - x_{n_j}|.$$

It follows that there is a sequence $m_i \rightarrow \infty$ such that

$$(*) \quad x_{m_i} \rightarrow x \quad \text{and} \quad y_{m_i} \rightarrow y \quad \text{as} \quad i \rightarrow \infty.$$

Set

$$\ell_n = |x_n - y_n|_K,$$

where $|\ast - \ast|_K$ denotes the distance between points in K . Since f is noncontracting, (ℓ_n) is a nondecreasing sequence.

By $(*)$, it follows that $\ell_{m_i} \rightarrow \ell_0$ as $m_i \rightarrow \infty$. It follows that (ℓ_n) is a constant sequence.

In particular

$$|x_0 - y_0|_K = \ell_0 = \ell_1 = |f(x_0) - f(y_0)|_K$$

for any pair of points (x_0, y_0) in K . I.e., f is distance preserving, in particular injective.

From (*), we also get that $f(K)$ is everywhere dense. Since K is compact $f: K \rightarrow K$ is surjective. Hence the result follows.

Comment. This is a basic lemma in the introduction to Gromov–Hausdorff distance; see for example [24, 7.3.30]. The proof presented here is not quite standard; it was given by Travis Morrison, one of my students at MASS program, Fall 2011 at Penn State.

61. *Embedding of a compact.* Let K be a compact metric space. Denote by $B(K)$ the space of bounded functions on K equipped with sup norm; i.e.,

$$|f| = \sup_{x \in K} |f(x)|.$$

Note that the map $\varphi: K \rightarrow B(K)$, defied by $x \mapsto \text{dist}_x$ is a distance preserving embedding.

Denote by W the linear convex hull of the image $\varphi(K) \subset B(K)$ with the metric induced from $B(K)$. It remains to show that W forms a compact length space.

Comment. The map φ is called *Kuratowski embedding*, although it was essentially discovered by Fréchet in the same paper he introduced metric spaces.

62. *2-sphere is far from a ball.* Assume contrary, let (\mathbb{S}^2, g) is sufficiently close to B^2 .

Choose a closed simple curve γ in \mathbb{S}^2 which is close to the boundary of B^2 . Choose two points p_1 and p_2 in \mathbb{S}^2 on the opposite sides of γ which are sufficiently close to the center of B^2 .

On one had p_1 and p_2 have to be close in \mathbb{S}^2 . On the other hang, to get from p_1 to p_2 in \mathbb{S}^2 , one has to cross γ . Hence the distance from p_1 to p_2 in \mathbb{S}^2 has to be close to 2, a contradiction.

63. *3-sphere is close to a ball.* Make fine burrows in the standard 3-ball which do not change its topology, but at the same time a come sufficiently close to any point in the ball.

Consider the doubling of obtained ball in its boundary. Clearly the obtained space is homeomorphic to \mathbb{S}^3 . prove that the burrows can be made so that it is sufficiently close to the original ball in the Gromov–Hausdorff metric.

It remains to smooth the obtained space slightly to get a genuine Riemannian metric with needed property.

Comment. I learned this construction from Sergei Ivanov.

64. Macrodimension. Choose a point $p \in M$, denote by f the distance function from p .

Let us cover M by the connected components of the preimages $f^{-1}((n-1, n+1))$. Clearly any point in M is covered by at most two such components. It remains to show that each of these components has diameter less than 100.

Assume contrary; let x and y be two points in such connected component and $|x - y|_M \geq 100$. Connect x to y by a curve τ in the component. Consider the closed curve σ formed by two geodesics $[px]$, $[py]$ and τ .

Prove that σ can be divided into 4 arcs α , β , γ and δ in such a way that the minimal distance from α to γ as well as the minimal distance from β to δ is at least 10.

Use the last statement to show that σ can not be shrank by a disc in its 1-neighborhood; the later contradicts the assumption.

Comment. The problem was discussed at a talk by Nikita Zinoviev around 2004.

✂ **65. Anti-collapse.** Fix a sequence $\varepsilon_n \rightarrow 0$.

Assume S be a surface with a flat disc $D_0 \subset S$ of radius r_0 . Let us cut from D_0 two discs D_1 and D'_1 of radii $r_1 = r_0/10$ and glue instead a cylinder with high ε_0 with discs on the top. Now repeat the operation for each D_1 and D'_1 cutting from each two discs of radius $r_2 = r_1/10$ and glue instead cylinder with high ε_0 with discs on the top. Continue the process, we get an increasing sequence of Riemannian metrics on S .

Comments. The problem appears in the paper [26] by Burago, Ivanov and Shoenthal.

✂ **66. No short embedding.** Let c_0, c_1, \dots, c_n be a chain of disjoint circles in \mathbb{R}^3 ; i.e., c_i and c_{i-1} are linked for each i . Assume that \mathbb{R}^3 is equipped with a length-metric d , such that the total length of the circles is ℓ .

Note that for any short homeomorphism $f: (\mathbb{R}^3, d) \rightarrow \mathbb{R}^3$ the distance from $f(c_0)$ to $f(c_n)$ is at most ℓ .

Fix a line segment $[ab]$ in \mathbb{R}^3 . Modify the metric on \mathbb{R}^3 in arbitrary small neighborhood of $[ab]$ so that there is a chain going along \mathbb{I} as above with arbitrary small total length ℓ , but at the same time the obtained metric is arbitrary close to the canonical; i.e.,

$$|d(x, y) - |x - y|| < \varepsilon$$

for any two points $x, y \in \mathbb{R}^3$ and fixed in advanced small $\varepsilon > 0$.

Now choose a converging sequence of line segments $[a_n b_n] \rightarrow [a_\infty b_\infty]$ and apply the above construction in nonoverlapping neighborhood of $[a_n b_n]$ and for a sequence ε_n which converges to zero very fast.

Note that for the obtained length metric d , the space (\mathbb{R}^3, d) does not admit a short homeomorphism to \mathbb{R}^3 . Indeed, if such homeomorphism h exists then from the above construction, we get

$$|h(a_\infty) - f(b_\infty)| \leq \ell_n + \varepsilon_n + \varepsilon_{n+1} + \dots$$

The right hand side converges to 0 as $n \rightarrow \infty$. Therefore $h(a_\infty) = f(b_\infty)$, a contradiction.

Comments. The problem appears in the paper [26] by Burago, Ivanov and Shoenthal.

67. Sub-Riemannian sphere. Prove that there is a nondecreasing sequence of Riemannian metric tensors $g_0 \leq g_1 \leq \dots$ such that the induced metrics converge to the given sub-Riemannian metrics. The metric g_0 can be assumed to be a metric on round sphere.

Applying the construction as in Nash–Kuiper theorem, one can produce a sequence of smooth embeddings $h_n: \mathbb{S}^n \rightarrow \mathbb{R}^{n+1}$ with the induced metrics g'_n such that $|g'_n - g_n| \rightarrow 0$.

Moreover, assume we assign a positive real number $\varepsilon(h)$ for any smooth embedding $h: \mathbb{S}^n \rightarrow \mathbb{R}^{n+1}$. Then we can assume that

$$|h_{n+1}(x) - h_n(x)| < \varepsilon(h_n)$$

for any $x \in \mathbb{S}^n$ and n . Choosing $\varepsilon(h_n)$ sufficiently small guarantees that h_n converges to say h_∞ and the metric induced by h_∞ coincides with the given sub-Riemannian metric.

Comments. The original papers of Nash [82] and Kuiper [65] are very readable.

It seems that the problem appeared on this list first it was re-discovered later by Le Donne in [69]. Yet my paper [94] is closely relevant.

68. Length preserving map. Assume there is a length-preserving map $f: \mathbb{R}^2 \rightarrow \mathbb{R}$. Note that f is Lipschitz. Therefore by Rademacher's theorem, f is differentiable almost everywhere.

Fix a unit vector u . Prove that, for almost all x , the length of curve $\alpha: t \mapsto x + t \cdot u$, $t \in [0, 1]$ can be expressed as the integral

$$\int_0^1 (d_{\alpha(t)} f)(u) \cdot dt.$$

It follows that $|d_x f(v)| = |v|$ for almost all $x, v \in \mathbb{R}^2$; in particular $d_x f$ is defined and has rank 2 at some point x , a contradiction.

Comment. Here is an exercise which can be solved the same way.

Let \mathbb{M}^2 be a Minkowski plane which is not isometric to the Euclidean plane. Show that \mathbb{M}^2 does not admit a length preserving map to \mathbb{R}^3 .

69. Hyperbolic space. Note that 2-dimensional hyperbolic space viewed as (\mathbb{R}^2, g) , where

$$g(x, y) = \begin{pmatrix} 1 & 0 \\ 0 & e^x \end{pmatrix}.$$

The same way 3-dimensional hyperbolic space viewed as (\mathbb{R}^3, h) , where where

$$h(x, y, z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^x & 0 \\ 0 & 0 & e^x \end{pmatrix}.$$

Prove that the map $\mathbb{R}^3 \rightarrow \mathbb{R}^4$ defined as

$$(x, y, z) \mapsto (x, y, x, z)$$

is a quasi-isometry from (\mathbb{R}^3, h) to its image in $(\mathbb{R}^2, g) \times (\mathbb{R}^2, g)$.

Comments. In the proof we used that horosphere in the hyperbolic space is isometric to the Euclidean plane. This observation appears already in the book of Lobachevsky, see [71, 34].

70. Midpoints. Without loss of generality, we can assume that $b + a = 0$.

Consider the sequence of isometries f_0, f_1, \dots defined recursively

$$f_{n+1}(x) = -f_n^{-1}(-f_n(x))$$

with $f_0 = f$. Note that $f_n(a) = a$ and $f_n(b) = b$ for any n and

$$|f_{n+1}(0)| = 2 \cdot |f_n(0)|.$$


The later condition implies that if $f(0) \neq 0$ then $|f_n(0)| \rightarrow \infty$ as $n \rightarrow \infty$. On the other hand, since f_n is isometry and $f(a) = a$, we get $|f_n(0)| \leq 2 \cdot |a|$, a contradiction.

Comment. Note that the problem is not trivial since in general metric midpoint of a and b in (\mathbb{R}^n, d) are not defined uniquely.

The solution above is the main step in the Väisälä's proof of Mazur–Ulam theorem; it states that any isometry $(\mathbb{R}^n, d) \rightarrow (\mathbb{R}^n, d)$ has to be an affine. See [109] and [80].

✂ 71. *Hilbert problem.*

Hilbert's construction. Let p and q be points in the interior of F . Consider the line ℓ passing through P and Q , denote by a and b the points in the intersection $\ell \cap \partial F$; we can assume that the points a, p, q and b appear on ℓ in the same order.



Set

$$\rho(p, q) = \ln \frac{|a - q| \cdot |p - b|}{|a - p| \cdot |q - b|}.$$

It remains to prove that ρ is a Finsler metric and any line segment in the interior F forms a geodesic for the metric ρ .

Pogorelov's construction. Fix a measure μ on the space of chords of F which is positive on any open set.

Given two points $x, y \in F$, denote by $\rho(x, y)$ the μ -measure of all chords which separate x from y in F . Show that ρ is a metric and the geodesics for this metric are formed by arcs of great circles in \mathbb{S}^2 .

It remains to choose the measure μ so that ρ is complete; such t measure can be found among regular measures with big density on short chords.

Comments. This metric was introduced by Hilbert in [56]; it was the motivating example of his 4-th problem [57]. The following construction given by Pogorelov is closely related.

Let μ be a measure on the space of equators in \mathbb{S}^2 which is positive on any open set. Given two points $x, y \in \mathbb{S}^2$, denote by $\rho(x, y)$ the measure of all equators which separate x from y . Then ρ is a metric and the geodesics for this metric are formed by arcs of great circles in \mathbb{S}^2 .

72. Straight geodesics. From uniqueness of straight segment between given points in \mathbb{R}^m , it follows that any straight line in \mathbb{R}^m forms a geodesic in (\mathbb{R}^m, ρ) .

Set

$$\|\mathbf{v}\|_{\mathbf{x}} = \rho(\mathbf{x}, (\mathbf{x} + \mathbf{v})).$$

Note that

$$\|\lambda \cdot \mathbf{v}\|_{\mathbf{x}} = |\lambda| \cdot \|\mathbf{v}\|_{\mathbf{x}}$$

for any $\mathbf{x}, \mathbf{v} \in \mathbb{R}^m$ and $\lambda \in \mathbb{R}$.

Prove that

$$\|\lambda \cdot \mathbf{v}\|_{\mathbf{x}} - \|\lambda \cdot \mathbf{v}\|_{\mathbf{x}'} \leq \text{Const} \cdot |\mathbf{x} - \mathbf{x}'|$$

for any $\mathbf{x}, \mathbf{x}', \mathbf{v} \in \mathbb{R}^m$, $\lambda \in \mathbb{R}$ and some fixed $\text{Const} \in \mathbb{R}$. Passing to the limit as $\lambda \rightarrow \infty$, we get $\|\mathbf{v}\|_{\mathbf{x}}$ does not depend on \mathbf{x} ; hence the result follows.

73. *A homeomorphism near quasi-isometry.* Let $M \geq 1$ and $A \geq 0$. Define (M, A) -quasi-isometry as a map $f: X \rightarrow Y$ between metric spaces X and Y such that

$$\frac{1}{M} \cdot |x - y| - A \leq |f(x) - f(y)| \leq M \cdot |x - y| + A$$

and any point in Y lies on the distance at most A from a point in the image $f(X)$.

Note that $(M, 0)$ -quasi-isometry is a $[\frac{1}{M}, M]$ -bi-Lipschitz map. Moreover, if $f_n: \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a $(M, \frac{1}{n})$ -quasi-isometry for each n then any partial limit of f_n as $n \rightarrow \infty$ is a $[\frac{1}{M}, M]$ -bi-Lipschitz map.

It follows that given $M \geq 1$ and $\varepsilon > 0$ there is $\delta > 0$ such that for any (M, δ) -quasi-isometry $f: \mathbb{R}^m \rightarrow \mathbb{R}^m$ and any $p \in \mathbb{R}^m$ there is an $[\frac{1}{M}, M]$ -bi-Lipschitz map $h: B_1(p) \rightarrow \mathbb{R}^m$ such that

$$|f(x) - h(x)| < \varepsilon$$

for any $x \in B_1(p)$.

Applying recaling, we can get the following equivalent formulation. Given $M \geq 1$, $A \geq 0$ and $\varepsilon > 0$ there is big enough $R > 0$ such that for any (M, A) -quasi-isometry $f: \mathbb{R}^m \rightarrow \mathbb{R}^m$ and any $p \in \mathbb{R}^m$ there is a $[\frac{1}{M}, M]$ -bi-Lipschitz map $h: B_R(p) \rightarrow \mathbb{R}^m$ such that

$$|f(x) - h(x)| < \varepsilon \cdot R$$

for any $x \in B_R(p)$.

Now cover \mathbb{R}^m by balls $B_R(p_n)$, construct a $[\frac{1}{M}, M]$ -bi-Lipschitz map $h_n: B_R(p_n) \rightarrow \mathbb{R}^m$ for each n . The maps h_n are close to f on $B_R(p_n)$.

It remains to deform slightly each h_n so that they agree on the overlaps. This can be done by Siebenmann's Theorem, see [99]. If instead you apply Sullivan's theorem, you get a bi-Lipschitz homeomorphism $h: \mathbb{R}^m \rightarrow \mathbb{R}^m$, see [102] or [108].

Comments. The problem was suggested by Dmitri Burago.

74. *A family of sets with no section.* Let us identify \mathbb{S}^1 with $[0, 1]/0 \sim 1$. Consider one parameter family of Cantor sets K_t formed by all possible sums $\sum_{n=1}^{\infty} a_n \cdot t^n$, where a_i is 0 or 1 and $t \in [0, \frac{1}{2}]$.

Note that $K_{\frac{1}{2}} = \mathbb{S}^1$.

Denote by $\rho_\alpha: \mathbb{S}^1 \rightarrow \mathbb{S}^1$ the rotation by angle α . Set $Z_t = \rho_{\frac{1}{1-2 \cdot t}}(K_t)$ for $t \in [0, \frac{1}{2})$ and $Z_{\frac{1}{2}} = \mathbb{S}^1$.

Prove that the family of sets Z_t is a continuous in Hausdorff topology and it does not have a section.

Comments. The problem is suggested by Stephan Stadler.

It is instructive to check that any Hausdorff continuous family of closed sets in \mathbb{R} admits a continuous section.

75. Sasaki metric. Show that there is a constant ℓ such that for any two unit tangent vectors $v \in T_p\mathbb{S}^2$ and $w \in T_q\mathbb{S}^2$ there is a path $\gamma: [0, 1] \rightarrow \mathbb{S}^2$ from p to q such that

$$\text{length } \gamma \leq \ell$$

and w is the parallel transformation of v along γ .

Note that once it is proved, it follows that diameter of the set of all vectors of fixed length in $T\mathbb{S}^2$ has diameter at most ℓ ; in particular the map $T\mathbb{S}^2 \rightarrow \mathbb{R}_{\geq 0}$ defined as $v \mapsto |v|$ preserves the distance with the maximal error ℓ . Hence the result follows.

Actions and coverings

76. Bounded orbit. Note that we can assume that the orbit $x_n = \iota^n(x)$ is dense in X ; otherwise pass to the closure of this orbit. In particular, we can choose a finite number of positive integers values n_1, n_2, \dots, n_k such that the points $x_{n_1}, x_{n_2}, \dots, x_{n_k}$ form a $\frac{1}{10}$ -net in $B_{10}(x_0)$.

Prove that that if $x_m \in B_1(x_0)$ then $x_{m+n_i} \in B_1(x_0)$ for some $i \in \{1, \dots, k\}$.

Set $N = \max_i \{n_i\}$. It follows that among any N elements in a row x_{i+1}, \dots, x_{i+N} there is at least one in $B_1(x_0)$. In particular, N isometric copies of $B_1(x_0)$ cover whole X . Hence the result follows.

Comments. The problem appears in the Calka's paper [30].

✂ **77. Covers of figure eight.** First show that any compact metric space can be presented as a limit of a sequence of finite metric graphs Γ_n ; moreover one can assume the degree of each vertex in G_n is 3 and each edge length of Γ_n is multiple of ε_n for some sequence $\varepsilon_n \rightarrow 0+$.

It remains to approximate any finite metric graph by finite coverings of figure eight. Guess this part of the proof from the following picture.

Comments The problem appears in thesis of Sahovic [98].

✂ **78. Constant curvature is everything.** Assume a fundamental group of a compact manifold M admits an epimorphism to the free group with two generators. In other words, there are two disjoint closed hypersurfaces S_1 and S_2 in M which do not separate points in M but each S_i separates points of any of its small neighborhoods.

Prepare many copies of M , cut one copy along S_1 and S_2 , n_1 copies along S_1 and n_2 copies along S_2 then glue them as it shown on the

picture. It produce a finite cover of M which is after rescaling is close to figure eight.

Finally applying Problem 77.

Comments The problem appears in thesis of Sahovic [98].

79. *Diameter of m -fold cover.* Let $\tilde{p}, \tilde{q} \in \tilde{M}$ and $\tilde{\gamma}: [0, 1] \rightarrow \tilde{M}$ be a minimizing geodesic from \tilde{p} to \tilde{q} . Denote by p, q, γ their projections to M .

We need to show that

$$\text{length } \tilde{\gamma} \leq m \cdot \text{diam}(M).$$

Suppose the contrary.

Represent γ as joint of m paths of equal length,

$$\gamma = \gamma_1 * \dots * \gamma_m,$$

so

$$\text{length}(\gamma_i) = \text{length}(\gamma)/m > \text{diam}(M).$$

Let σ_i be a minimizing geodesic in M connecting the endpoints of γ_i . Note that

$$\text{length } \sigma_i \leq \text{diam } M < \text{length } \gamma_i.$$

Consider $m+1$ paths $\alpha_0, \dots, \alpha_m$ defined as

$$\alpha_i = \sigma_1 * \dots * \sigma_i * \gamma_{i+1} * \dots * \gamma_m.$$

Consider their liftings $\tilde{\alpha}_0, \dots, \tilde{\alpha}_m$ with \tilde{q} as the endpoint. Note that two curves, say α_i and α_j for $i < j$, have the same starting point in \tilde{M} . Consider the path

$$\beta = \gamma_1 * \dots * \gamma_i * \sigma_{i+1} * \dots * \sigma_j * \gamma_{j+1} * \dots * \gamma_m.$$

Prove that there is lift $\tilde{\beta}$ of β which connects \tilde{p} to \tilde{q} in \tilde{M} . Clearly $\text{length } \beta < \text{length } \gamma$, a contradiction.

Comments. The question was originally asked by Alex Nabutovsky; it was answered by Sergei Ivanov, see [93].

80. *Symmetric square.* Let $\Gamma = \pi_1 X$ and $\Delta = \pi_1((X \times X)/\mathbb{Z}_2)$. Consider the homomorphism $\varphi: \Gamma \times \Gamma \rightarrow \Delta$ induced by the projection $X \times X \rightarrow (X \times X)/\mathbb{Z}_2$.

Prove that the restrictions $\varphi|_{\Gamma \times \{1\}}$ and $\varphi|_{\{1\} \times \Gamma}$ are onto.

It remains to note that

$$\varphi(\alpha, 1)\varphi(1, \beta) = \varphi(1, \beta)\varphi(\alpha, 1)$$

for any α and β in Γ .

Comments. The problem was suggested by Rostislav Matveyev.

81. Sierpinski triangle. Denote the Sierpinski triangle by Δ .

Let us show that any homeomorphism of Δ is also its isometry. Therefore the group homeomorphisms is the symmetric group S_3 .

Let $\{x, y, z\}$ be a 3-point set in Δ such that $\Delta \setminus \{x, y, z\}$ has 3 connected components. Prove that there is unique choice for the set $\{x, y, z\}$ and it is formed by the midpoints of its big sides.

It follows that any homeomorphism of Δ permutes the set $\{x, y, z\}$.

A similar argument shows that this permutation uniquely describes the homeomorphism.

Comments. The problem was suggested by Bruce Kliener.

82. Lattices in a Lie group. Denote by V_ℓ and W_m the Voronoi domain of for each $\ell \in L$ and $m \in M$ correspondingly; i.e.,

$$V_\ell = \{ g \in G \mid |g - \ell|_G \leq |g - \ell'| \text{ for any } \ell' \in L \}$$

$$W_m = \{ g \in G \mid |g - m|_G \leq |g - m'| \text{ for any } m' \in M \}$$

Note that for any $\ell \in L$ and $m \in M$ we have

$$\begin{aligned} \text{vol } V_\ell &= \text{vol}(L \setminus (G, h)) = \\ (*) \quad &= \text{vol}(M \setminus (G, h)) = \\ &= \text{vol } W_m. \end{aligned}$$

Consider the bipartite graph Γ with vertices formed by the elements of L and M such that $\ell \in L$ is adjacent to $m \in M$ if and only if $V_\ell \cap W_m \neq \emptyset$.

By (*) the graph Γ satisfies the condition in the Hall's marriage theorem. Therefore there is a bijection $f: L \rightarrow M$ such that

$$V_\ell \cap W_{f(\ell)} \neq \emptyset$$

for any $\ell \in L$.

It remains to notice that f is bi-Lipschitz..

Comments. The problem is discussed in the paper [27] by Burago and Kleiner. For a finitely generated group G it is not known if G and $G \times \mathbb{Z}_2$ can fail to be bi-Lipschitz. (The groups are assumed to be equipped with word metric.)

83. Piecewise Euclidean quotient. Note that the group Γ serves as holonomy group of the quotient space $P = \mathbb{R}^n / \Gamma$ with the induced polyhedral metric. More precisely, one can identify \mathbb{R}^n with the tangent space of a regular point x_0 of P in such a way that for any $\gamma \in \Gamma$ there is a loop ℓ in P which pass only through regular points and has the holonomy γ .

Fix $\gamma \in \Gamma$. Let ℓ be the corresponding loop. Since P is simply connected, we can shrink ℓ by a disc. By general position argument we can assume that the disc only pass through simplices of codimension 0, 1 and 2 and intersect the simplices of codimension 2 transversely.

In other words, ℓ can be presented as a product of loops such that each loop goes around a single simplex of codimension 2 and comes back. The holonomy for each of these loops is a rotation around a hyperplane. Hence the result follows.

Comments. The converse to the problem also holds; it was proved by Lange in [67], his proof based ealier results of Mikhailova, see [74].

Note that the cone over spherical suspension over Poincaré sphere is homeomorphic to \mathbb{R}^5 and it is quotient of \mathbb{R}^5 by a finite subgroup of $\mathrm{SO}(5)$. I.e., if in this problem one exchanges “PL-homeomorphism” to “homeomorphism” then the answer is different; a complete classification of such actions was also obtained in [67].

84. Subgroups of free group. Let G be a finitely generated subgroup of free group with m generators, further denoted by F_m .

Let W be the wedge sum of n circles, so $\pi_1(W, p) = F_m$. Equip W with length metric so that that each circle has unit length.

Pass to the metric cover \tilde{W} of W such that $\pi_1(\tilde{W}, \tilde{p}) = G$ for a lift \tilde{p} of p .

Fix sufficiently large integer n and consider doubling of the closed ball $\tilde{B}_{n+\frac{1}{2}}(\tilde{p})$ in its boundary. Let us denote the obtained doubling by Z_n and set $G_n = \pi(Z_n, \tilde{p})$.

Prove that Z_n is a metric covering of W ; it makes possible to consider G_n as a subgroup of F_m . By construction, Z_n is compact; therefore G_n has finite order in F_m .

It remains to show that that

$$G = \bigcap_{n>k} G_n,$$

where k is the maximal length of word in the generating set of G .

Comments. Originally the problem was solved by Hall in [78]. The proof presented here is close to the solution of Stalings in [101]; see also [116].

Many results about free groups which can be proven similar way. For example:

- ◊ Subgroups of free groups are free.
- ◊ Two elements of the free groups u and v commute if and only if they are powers of some third element.

85. Lengths of generators of the fundamental group. Choose a length-minimizing geodesic loop γ which represents a given element $a \in \pi_1 M$.

Fix $\varepsilon > 0$. Represent γ as a joint

$$\gamma = \gamma_1 * \dots * \gamma_n$$

of paths with length $\gamma_i < \varepsilon$ for each i .

Denote by $p = p_0, p_1, \dots, p_n = p$ the endpoints of these arcs. Connect p to p_i by a length minimizing geodesic σ_i . Note that γ is homotopic to a product of loops

$$\alpha_i = \sigma_{i-1} * \gamma_i * \sigma_{i-1}$$

and length $\alpha_i < 2 \cdot \text{diam } M + \varepsilon$ for each i .

It remains to show that for sufficiently small $\varepsilon > 0$ any loop with length less than $2 \cdot \text{diam } M + \varepsilon$ is homotopic to a loop with length at most $2 \cdot \text{diam } M$.

Comments. The statement was observed by Gromov, see [48, Proposition 3.22].

✂ **86.** *Short basis.* Fix a point $p \in M$; set $\Gamma = \pi_1(M, p)$.

Given $a \in \Gamma$ denote by γ_a a length minimizing geodesic loop in the class a . Set $|a| = \text{length } \gamma_a$.

Let us construct so called *short basis* of $a_1, a_2, \dots \in \Gamma$.

1. Choose $a_1 \in \Gamma$ so that $|a_1|$ takes the minimal value.
2. Choose $a_2 \in \Gamma \setminus \langle a_1 \rangle$ so that $|a_2|$ takes the minimal value.
3. Choose $a_2 \in \Gamma \setminus \langle a_1, a_2 \rangle$ so that $|a_2|$ takes the minimal value.
4. and so on

Note that the sequence terminates after n steps if and only if a_1, \dots, a_n forms a generating set of Γ .

Given $a \in \Gamma$, set $w_a = \gamma'_a(0) \in T_p$. Assume

$$\angle(w_{a_i}, w_{a_j}) < \frac{\pi}{3}$$


for some $i < j$. By Toponogov comparison theorem we get $|a_j a_i^{-1}| < |a_j|$; the later leads to a contradiction.

It follows that $\angle(w_{a_i}, w_{a_j}) \geq \frac{\pi}{3}$ for all $i \neq j$. It gives a bound on n in terms of $\dim M$.

Comments. This construction introduced by Gromov in his paper on almost flat manifolds [44].

Topology

✂ **87.** *Milnor's disks.* The immersed circle below is called Bennequins curve. It is a good exercise to count different discs bounded by the given immersed circle.



[46]

✂ 88. *Positive Dehn twist.* Consider the universal covering $\tilde{\Sigma} \rightarrow \Sigma$. The surface $\tilde{\Sigma}$ comes with the orientation induced from Σ .

Note that we may assume that $\tilde{\Sigma}$ has infinite number of boundary components.

Fix a point x_0 on the boundary of $\tilde{\Sigma}$. Given other points y and z we will write $y \prec z$ if z lies on the left side from one (and therefore any) simple curve in $\tilde{\Sigma}$ which goes from x to y . Note that \prec defines a linear order on $\partial\tilde{\Sigma} \setminus \{x\}$. We will write $y \preceq z$ if $y \prec z$ or $y = z$.

Note that any homeomorphism $h: \Sigma \rightarrow \Sigma$ which is identity on the boundary lifts to unique homeomorphism $\tilde{h}: \tilde{\Sigma} \rightarrow \tilde{\Sigma}$ is such a way that $\tilde{h}(x) = x$.

Prove that if h is positive Dehn twist then $y \preceq \tilde{h}(y)$ for any $y \neq x$ in $\partial\tilde{\Sigma} \setminus \{x\}$ and there is a point y_0 such that $y_0 \prec \tilde{h}(y_0)$.

Finally note that the later property is a homotopy invariant and it survives under compositions of maps. Hence the statement follows.

Comments. The problem was suggested by Rostislav Matveyev.

✂ 89. *Function with no critical points.* Construct an immersion $\psi: B^n \rightarrow \mathbb{R}^n$ such that

$$\ell \circ \varphi \neq \ell \circ \psi$$

for any embedding $\varphi: B^n \rightarrow \mathbb{R}^n$.

It remains to note that the composition $f = \ell \circ \psi$ has no critical points.

✂ 90. *Conic neighborhood.* Let V and W be two cone neighborhoods of point p . Without loss of generality, we may assume that $V \subset W$.


Let us construct a sequence of embeddings $f_n: V \rightarrow W$ such that (1) for any compact set K there is a positive ineteger n such that $f_n(k) = f_m(k)$ for any $k \in K$ and $m \geq n$. (2)??? $\bigcup_n f_n(V) = W$.

Once such sequence is constructed, set $f(x) = f_n(x)$ for all large values of n .

✂ **91.** *No C^0 -knots.*

Comment. This problem was suggested by Greg Kuperberg, see [92].

92. *Simple stabilization.* The example can be guessed from the diagram.



Caomments. It was one of the special problems in my analysis class taught by Galuzina. Likely this is not the original source.

93. *Isotropy.* Fix a hoemeomrphism $\varphi: K_1 \rightarrow K_2$.

By Tietze extension theorem, the hoemeomrphisms $\varphi: K_1 \rightarrow K_2$ and $\varphi^{-1}: K_2 \rightarrow K_1$ can be extended to a continuous maps, say $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^n \rightarrow \mathbb{R}^n$ correspondingly.

Consider the homeomorphisms $h_1, h_2, h_3: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ defined the following way

$$\begin{aligned} h_1(x, y) &= (x, y + f(x)), \\ h_2(x, y) &= (x - g(y), y), \\ h_3(x, y) &= (y, -x). \end{aligned}$$

It remains to prove that each homeomorphism h_i is isotopic to the indentity map and

$$h = h_3 \circ h_2 \circ h_1.$$

Comments. The problem had been discussed by Klee in [63].

✂ **94.** *Cantor set.*

✂ **95.** *Knaster's circle.* A map $f: \mathbb{S}^1 \rightarrow \mathbb{S}^1$ will be called ε -crooked if for any arc $\mathbb{I} \subset \mathbb{S}^1$ with end points a and b there are points $x, y \in \mathbb{I}$ such that the points a, y, x, b appear on \mathbb{I} in the same order and

$$|f(x) - f(a)|_{\mathbb{S}^1}, |f(y) - f(b)|_{\mathbb{S}^1} < \varepsilon.$$

Show that for any $\varepsilon > 0$ there is an ε -crooked map $f: \mathbb{S}^1 \rightarrow \mathbb{S}^1$ of degree 1.

Take a sequence of ε_n -crooked maps for a sequence ε_n which converge fast to 0. and use this map to construct a nested sequence of embedding of annuli in the plane. Each annulus bounds a disc and the intersection of all these annuli bound a disc which is the union of all these discs.

It remains to show that the boundary of the obtained disc does not contain a simple curve.

Comments. [79].

An other related problem is to construct a set in \mathbb{R}^3 homeomorphic to the Cantor's set with non simply connected complement.

96. *Boundary in \mathbb{R} .* Prove that the Cantor's set forms a boundary of three disjoint open set in \mathbb{R} .

Comments. In the plane one can assume in addition that each set is connected. This examples are called *lakes of Wada*; these are three disjoint open discs in the plane which share the same boundary. This example described by Yoneyama in [120].

✖ **97.** *Deformation of homeomorphism.* Fix a smooth increasing concave function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ such that $\varphi(r) = r$ for any $r \leq 1$ and $\sup_r \varphi(r) = 2$.

Consider \mathbb{R}^n with polar coordinates (u, r) , where $u \in \mathbb{S}^{n-1}$ and $r \geq 0$. Let $\Phi: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is defined by $\Phi(u, r) = (u, \varphi(r))$.

Set $h(x) = \Phi \circ f \circ \Phi^{-1}(x)$ is $|x| < 2$ and $h(x) = x$ otherwise.

Prove that $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a solution.

Comments. ???

✖ **98.** *Finite topological space.* Let F be a finite topological space. Given two points $p, q \in F$ we will write $p \preceq q$ if p lies in any closed set containing q .

Prepare a cell for each point in F

Consider a finite CW complex W .

Denote by S the set of all cells of W and equip S with the topology such that ???

Piecewise linear geometry

99. *Triangulation of 3-sphere.* Choose 100 distinct points $x_1, x_2, \dots, \dots, x_{100}$ on the curve

$$\gamma: t \mapsto (t, t^2, t^3, t^4)$$

in \mathbb{R}^4 . Let P be the convex hull of $\{x_1, x_2, \dots, x_{100}\}$.

Prove that for any two points x_i and x_j there is a hyperplane H in \mathbb{R}^4 which pass through x_i and x_j and leaves γ on one side. The later statement implies that any two vertices x_i and x_j of P are connected by an edge.

The statement follows since the surface of P is homeomorphic to \mathbb{S}^2 .

Comments. The polyhedron P above is an example of so called *cyclic polytopes*.

100. Spherical arm lemma. Let us cut the polygon A from the sphere and glue instead the polygon B . Denote by Σ the obtained spherical polyhedral space. Note that

- ◇ Σ is homeomorphic \mathbb{S}^2 .
- ◇ Σ has curvature ≥ 1 in the sense of Alexandrov; i.e., the total angle around each singular point is less than 2π .
- ◇ All the singular points of Σ lie outside of an isometric copy of a hemisphere $\mathbb{S}_+^2 \subset \Sigma$

It is sufficient to show that Σ is isometric to the standard sphere. Assume contrary. If n denotes be the number of singular points in Σ , it means that $n > 0$.

We will arrive to a contradiction applying induction on n . The base case $n = 1$ is trivial; i.e., Σ can not have single singular point.

Now assume Σ has $n > 1$ singular points. Choose two singular points p, q , cut Σ along a geodesic $[pq]$ and patch the hole so that the obtained new polyhedron Σ' has curvature ≥ 1 . The patch is obtained by doubling a spherical triangle in two sides. For the right choice of the triangle, the points p and q become regular in Σ' and exactly one new singular point appears in the patch.

This way, constructed a spherical polyhedral space Σ' with $n - 1$ singular points which satisfy the same conditions as Σ

By induction hypothesis Σ' does not exist. Hence the result follow.

Alternative end of proof. By Alexandrov embedding theorem, Σ is isometric to the surface of convex polyhedron P in the unit 3-dimensional sphere \mathbb{S}^3 . The center of hemisphere has to lie in a facet, say F of P . It remains to note that F contains the equator and therefore P has to be hemisphere in \mathbb{S}^3 or intersection of two hemispheres. In both cases its surface is isometric to \mathbb{S}^2 .

Comments. The problem appear in Zalgaller's paper [117].

The patch construction above was introduced by Alexandrov in his proof of convex embeddability of polyhedrons; the earliest reference we have found is [10, VI, §7].

An alternative proof can be build of Alexandrov's embedding theorem, see [88].

✂ **101. Piecewise linear isometry I.** Given a triangulation of P consider the Voronoi domain for each vertex. Prove that the triangulation can be subdivided if necessary so that Voronoi domain of each vertex is isometric to a convex subset in the cone with vertex corresponding to the tip.

Note that the boundaries of all the Voronoi domains form a graph with straight edges. One can triangulate P so that each triangle has such edge as the base and the remaining vertex is the center of an adjusted Voronoi domain.

Let us map all the vertices of the original triangulation in the origin of \mathbb{R}^2 . ???

✂ **102. Piecewise linear isometry II.** Let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be two collections of points in \mathbb{R}^2 such that $|a_i - a_j| \geq |b_i - b_j|$ for all pairs i, j . We need to construct a piecewise linear length-preserving map $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $f(a_i) = b_i$ for each i .

Assume that the problem is solved if $n < m$. Then there is a piecewise linear length-preserving map $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $f(a_i) = b_i$ for each $i > 1$. Consider the set

$$\Omega = \{ x \in \mathbb{R}^2 \mid |f(x) - b_1| > |x - a_1| \}.$$

If $\Omega = \emptyset$ then $f(a_1) = b_1$; i.e., the problem is solved.

Prove that Ω is the interior of a polygon which is star-shaped with respect to a_1 . Redefine the map f inside Ω so that it remains piecewise linear length-preserving and $f(a_1) = b_1$.

Comments. This problem appears in paper of Brehm, see [22]. The same proof works in all dimensions. This problem is closely related to Kirszbraun's theorem stated below. It was proved by Kirszbraun in his thesis [62], later it was reproved by Valentine in [110]; the proof of Brehm is very close to the one given by Valentine.

103. Minimal polyhedron. Arguing by contradiction, assume T is a minimal polyhedral surface which is not saddle.

Prove that one can move one of the vertices of T in such a way that the lengths of all edges starting at this vertex decrease.


Prove that if, by this deformation, the area does not decrease then there are two adjusted triangles in the triangulation, say $\triangle pxy$ and $\triangle qxy$ such that

$$\angle xpy + \angle xqy > \pi.$$


Finally show that in this case exchanging triangles $\triangle pxy$ and $\triangle qxy$ to the triangles $\triangle pxq$ and $\triangle pyq$ leads to a polyhedral surface with smaller area. I.e., T was not minimal, a contradiction.

Comments. This problem is discussed in my paper [95].

For general polyhedral surface, the deformation which decrease the legths of all edges may not decrease the area. Moreover, the surface which minimize the area among all surfaces with fixed triagulation may be not saddle. An example of such surface can be seen on the picture.



104. Coherent triangulation. Look at the diagram and think.



Comments. The problem discussed in section 7C of the book [41] by Gelfand, Kapranov and Zelevinsky.

105. Characterization of polytope. Arguing by contradiction, let us assume that $P \subset \mathbb{R}^m$ is a counterexample and m takes minimal possible value.

Choose a finite cover B_1, B_2, \dots, B_n of K , where $B_i = B_{\varepsilon_i}(z_i)$ and $B_i \cap P = B_i \cap K_i$, where K_i is a cone with the tip at z_i .

For each i , consider function $f_i(x) = |z_i - x|^2 - \varepsilon_i^2$. Note that

$$W_{i,j} = \{ x \in \mathbb{R}^n \mid f_i(x) = f_j(x) \}$$

is a hyperplane for any pair $i \neq j$.

The subset $P_{i,j} = P \cap W_{i,j}$ satisfies the same asumtions as P , but lies in a hyperplane. Since m is minimal, we get that $P_{i,j}$ is a polytope for any pair i, j .

Consider Voronoi domains

$$V_i = \{ x \in \mathbb{R}^n \mid f_i(x) \geq f_j(x) \text{ for any } j \}.$$

Note that $P \cap V_i$ is formed by the points which lie on the segments from z_i to a point in $P \cap \partial V_i$.

The statement follows since ∂V_i is covered by the hyperplanes $W_{i,j}$.

Comments. The problem is mentioned in the paper [68] by Lebedeva and Petrunin.

106. *A sphere with one edge.* Such example P can be found among the spherical polyhedral spaces which admit an isometric \mathbb{S}^1 -action with geodesic orbits.

Fix large relatively prime integers $p > q$. Consider the triangle Δ with angles $\frac{\pi}{p}$, $\frac{\pi}{q}$ and say $\pi \cdot (1 - \frac{1}{p})$ in the sphere of radius $\frac{1}{2}$. Denote by $\hat{\Delta}$ the doubling of Δ in its boundary. Note that $\hat{\Delta}$ is homeomorphic to \mathbb{S}^2 , it has 3 singular points with total angles $2 \cdot \frac{\pi}{p}$, $2 \cdot \frac{\pi}{q}$ and $2 \cdot \pi \cdot (1 - \frac{1}{p})$.

Consider \mathbb{S}^1 -action on $\mathbb{S}^3 \subset \mathbb{C}^2$ by the diagonal matrices $\begin{pmatrix} z^p & 0 \\ 0 & z^q \end{pmatrix}$, $z \in \mathbb{S}^1 \subset \mathbb{C}$. Construct a spherical polyhedral metric d on \mathbb{S}^3 such that the \mathbb{S}^1 -orbits become geodesics and the quotient $(\mathbb{S}^3, d)/\mathbb{S}^1$ is isometric to $\hat{\Delta}$.

In the constructed example the singular points with total angles $2 \cdot \frac{\pi}{p}$ and $2 \cdot \frac{\pi}{q}$ should correspond to the points with isotropy groups \mathbb{Z}/p and \mathbb{Z}/q of the action. The points in $P = (\mathbb{S}^3, d)$ on the orbits over these points will be regular points of P . The singular locus P^* of P will be formed by the orbit corresponding to the remaining singular point of $\hat{\Delta}$. By construction,

- ◇ P^* is a closed geodesic with angle $2 \cdot \pi \cdot (1 - \frac{1}{p})$ around it.
- ◇ P^* forms a (p, q) -torus knot in the ambient \mathbb{S}^3 .

Comments. Likely only the torus knots can appear this way.

The construction appears in the paper [85] by Panov. The cone K over P is a polyhedral space with natural complex structure; i.e., one can cut simplices from \mathbb{C}^2 and glue the cone from them in such a way that complex structures will agree along the gluings. Moreover the cone K can be holomorphically parametrized by \mathbb{C}^2 in such a way that the cone over P^* becomes an algebraic curve $z^p = w^q$ in (z, w) -coordinates of \mathbb{C}^2 .

107. *Triangulation of a torus.* Let us equip the torus with the flat metric such that each triangle is equilateral. The metric will have two singular cone points, the first corresponds to the vertex v_5 with 5 triangles, the total angle around this point is $\frac{5}{3} \cdot \pi$ and the second corresponds to the vertex v_7 with 7 triangles, the total angle around this point is $\frac{7}{3} \cdot \pi$.

Prove the following.

Observation *The holonomy group of this metric is generated by rotation by $\frac{\pi}{3}$.*

Consider a closed geodesic γ_1 which minimize the length of all circles which are not null-homotopic. Let γ_2 be an other closed geodesic which minimize the length and is not homotopic to any power of γ_1 .

Show that γ_1 and γ_2 intersect at a single point.

Show that γ_i can not pass v_5 .

Apply the observation above to show that if γ_i pass through v_7 then the measure of one of two angles which γ_i cuts at v_7 equals to π . Use the later statement to show that one can push γ_i aside so it does not longer pass through v_7 , but remains a closed geodesic.

Cut \mathbb{T}^2 along γ_1 and γ_2 . In the obtained quadrilateral, connect v_5 to v_7 by a minimizing geodesic and cut along it. This way we obtain an annulus with flat metric. Look at the neighborhood of the boundary components and show that the anulus can and can not be isometrically immersed into the plane; this is a contradiction.

Comments. There are flat metrics on the torus with two singular points which have the total angles $\frac{5}{3}\cdot\pi$ and $\frac{7}{3}\cdot\pi$. Such example can be obtained by identifying the the hexagon on the picture according to the arrows. But the holonomy group of the obtained torus is generated by the rotation by angle $\frac{\pi}{6}$. In particular, the observation is necessary in the proof.



The same argument shows that holonomy group of flat torus with exactly two singular points with total angle $2\cdot(1\pm\frac{1}{n})\cdot\pi$ has more than n elements. In the solution we did the case $n = 6$.

The problem was originally discovered and solved in [61], their proof is combinatorial. The solution described above was given by Rostislav Matveyev in his lectures [72]. A complex-analytic proof was later found in the paper [60] by Izmistiev, Rote, Springborn and Kusner.

108. *Unique geodesics imply CAT(0).* Uniqueness of geodesics implies that P is contractable. In particular, P is simply connected.

It remains to prove that P is locally CAT(0); equivalently, the space of directions Σ_p at any point $p \in P$ is a CAT(1) space.

We can assume that the statement holds in all dimensions less than $\dim P$. In particular, Σ_p is locally CAT(1). If Σ_p is not CAT(1) then it contains a periodic geodesic γ of length $\ell < 2\cdot\pi$, such that any arc of γ of length $\frac{\ell}{2}$ is length minimizing.

Consider two points x and y in the tangent cone of p in directions $\gamma(0)$ and $\gamma(\frac{\ell}{2})$. Show that there are two distinct minimizing geodesics between x and y . The later leads to a contradiction.

Comments. The existence of geodesic γ seem to be proved first by Bowditch in [21]; a simpler proof can be found in [9].

109. *No simple geodesic.* The curvature of a vertex on the surface of a convex polyhedron is defined as the $2\cdot\pi - \theta$, where θ is the total angle around the vertex.

Notice that a simple closed geodesic cuts the surface into two discs with total curvature $2\cdot\pi$ each. Therefore it is sufficient to construct a convex polyhedron with curvatures of the vertices $\omega_1, \omega_2, \dots, \omega_n$ such that $2\cdot\pi$ can not be obtained as sum of some of some of ω_i . An example of that type can be found among 3-simplexes.

Comments. The problem discussed in the Galperin's paper [40].

Discrete geometry

110. *Box in a box.* Let Π be a parallelepiped with dimensions a, b and c . Denote by $v(r)$ the volume of r -neighborhoods of Π ,

Note that for all positive r we have

$$(*) \quad v(r) = w_3 + w_2 \cdot r + w_1 \cdot r^2 + w_0 \cdot r^3,$$

where

- ◇ $w_0 = \frac{4}{3} \cdot \pi$ is the volume of unit ball,
- ◇ $w_1 = \pi \cdot (a + b + c)$,
- ◇ $w_2 = 2 \cdot (a \cdot b + b \cdot c + c \cdot a)$ is the surface area of Π ,
- ◇ $w_3 = a \cdot b \cdot c$ is the volume of Π ,

Assume Π' be an other parallelepiped with dimensions a', b' and c' . For the volume $v'(r)$ the volume of r -neighborhoods of Π' we have a formula similar (*).

Note that if $\Pi \subset \Pi'$ then $v(r) \leq v'(r)$ for any r . Checking this inequality for $r \rightarrow \infty$, we get

$$a + b + c \leq a' + b' + c'.$$

Comments. The problem was discussed by Shen in [100].

A formula analogous to (*) holds for arbitrary convex body B in arbitrary dimension n . The coefficient $w_i(B)$ in the polynomial with different normalization constants appear under different names most commonly *intrinsic volume* and *quermassintegral*. The later can be also defined as the average of area of projections of B to the i -dimensional planes. In particular if B' and B are convex bodies such that $B' \subset B$ then $w_i(B') \leq w_i(B)$ for any i . This generalize our problem quite a bit. Further generalizations lead to so called *mixed volumes*, see [25] for more on the subject.

111. *Round circles in \mathbb{S}^3 .* For each circle consider the containing it plane in \mathbb{R}^4 . Note that the circles are linked if and only if the corresponding planes intersect at a single point inside \mathbb{S}^3 .

Take the intersection of the planes with the sphere of radius $R \geq 1$, rescale and pass to the limit as $R \rightarrow \infty$. This way we get needed isotopy.

Comments. The problem was discussed in the thesis of Walsh, see [111].

112. Harnack's circles. Let $\sigma \subset \mathbb{RP}^2$ be a smooth algebraic curve of degree d . Consider the complexification $\Sigma \subset \mathbb{CP}^2$ of σ . Without loss of generality, we may assume that Σ is regular.

Prove that all regular complex algebraic curves of degree d in \mathbb{RP}^2 are homeomorphic to each other. Straightforward calculation show that Σ has genus $n = \frac{1}{2} \cdot (d^2 - 3 \cdot d + 4)$.

The real curve σ forms the fixed point set of Σ by complex conjugation. Prove that each connected component of σ adds 1 to the genus of Σ . Hence the result follows.

Comment. This problem was suggested by Greg Kuperberg, see [92].

113. Two points on each line. Take any complete ordering of the set of all lines so that each beginning interval has cardinality less than continuum.

Assume we have a set of points X such that each line intersects X at at most 2 points and cardinality of X is less than continuum.

Choose the least line ℓ in the ordering which intersect X by 0 or 1 point. Note that the set of all lines intersecting X at two points has cardinality less than continuum. Therefore we can choose a point on ℓ and add it to X so that the remaining lines are not overloaded.

It remains to apply well ordering principle.

114. Bodies with the same of shadows. Let B be the unit ball in \mathbb{R}^3 centered at the origin.

Fix small $\varepsilon > 0$. Consider two bodies

$$\begin{aligned} B'' &= \{ (x, y, z) \in B \mid x \leq 1 - \varepsilon, y \leq 1 - \varepsilon \}, \\ B''' &= \{ (x, y, z) \in B \mid x \leq 1 - \varepsilon, y \leq 1 - \varepsilon, z \leq 1 - \varepsilon \}. \end{aligned}$$

Prove that B'' and B''' have the same shadows.

Comments. The problem based on the question of Hamkins [55] answered by Ivanov.

115. Kissing number. Let $m = \text{kiss } B$ and B_1, B_2, \dots, B_m the the copies of B which touch B and have no common interior points. For each B_i consider the vector v_i from the center of B to the center of B_i . Note that $\angle(v_i, v_j) \geq \frac{\pi}{3}$ if $i \neq j$.

For each i , consider supporting hyperplane Π_i to W with outer normal vector v_i . Denote by W_i the reflection of W in Π_i .

Prove that W_i and W_j have no common interior points if $i \neq j$; the later gives the needed inequality.

Comments. The proof is given by Halberg, Levin and Straus in [54].

It expected that the same inequality holds for the orientation-preserving version of kissing number.

116. Monotonic homotopy. Note that we can assume that $h_0(F)$ and $h_1(F)$ both lie in the coordinate n -spaces of $\mathbb{R}^{2 \cdot n} = \mathbb{R}^n \times \mathbb{R}^n$; i.e., $h_0(F) \subset \mathbb{R}^n \times \{0\}$ and $h_1(F) \subset \{0\} \times \mathbb{R}^n$.

Show that the following homotopy is monotonic

$$h_t(x) = (h_0(x) \cdot \cos \frac{\pi \cdot t}{2}, h_1(x) \cdot \sin \frac{\pi \cdot t}{2}).$$

Comment. This homotopy was discovered by Ralph Alexander in [6]. It has number of applications, one of the most beautiful is the given by Bezdek and Connolly [17] in their proof of Kneser–Poulsen and Klee–Wagon conjectures in dimension 2.

117. Cube. Consider the cube $[-1, 1]^n \subset \mathbb{R}^n$. Any vertex this cube has the form $\mathbf{q} = (q_1, q_2, \dots, q_n)$, where $q_i = \pm 1$.

For each vertex \mathbf{q} , consider the intersection of the corresponding octant with the unit sphere; i.e., the set

$$V_{\mathbf{q}} = \{ (x_1, x_2, \dots, x_n) \in \mathbb{S}^{n-1} \mid q_i \cdot x_i \geq 0 \text{ for each } i \}.$$

Consider the set $\mathcal{A} \subset \mathbb{S}^{n-1}$ formed by the union of all the sets $V_{\mathbf{q}}$ for $\mathbf{q} \in A$. Note that

$$\text{vol}_{n-1} \mathcal{A} = \frac{1}{2} \cdot \text{vol}_{n-1} \mathbb{S}^{n-1}$$

and

$$\text{vol}_{n-2} \partial \mathcal{A} = \frac{k}{2^{n-1}} \cdot \text{vol}_{n-2} \mathbb{S}^{n-2},$$

where k is the number of edges of the cube with one end in A and the other in B .

It remains to show that

$$\text{vol}_{n-2} \partial \mathcal{A} \geq \text{vol}_{n-2} \mathbb{S}^{n-2}.$$

The later follows from the isoperimetric inequality for \mathbb{S}^n .

Comment. The problem was suggested by Greg Kuperberg, see [92].

118. Right-angled polyhedron. Before coming into proof read about *Dehn–Sommerville equations* on page 82.

Let P be a right-angled hyperbolic polyhedron of dimension m . Note that P is simple; i.e., exactly m facets meet at each vertex of P .

From the projective model of hyperbolic plane, one can see that for any simple compact hyperbolic polyhedron there is a simple Euclidean polyhedron with the same combinatorics. In particular Dehn–Sommerville equations hold for P .

Denote by (f_0, f_1, \dots, f_m) and (h_0, h_1, \dots, h_m) the f - and h -vectors of P . Recall that $h_i \geq 0$ for any i and $h_0 = h_m = 1$. By Dehn–Sommerville equations, we get

$$(*) \quad f_2 > \frac{m-2}{4} \cdot f_1.$$

Since P is hyperbolic, each 2-dimensional face of P has at least 5 sides. It follows that

$$f_2 \leq \frac{m-1}{5} \cdot f_1.$$

The later contradicts $(*)$ for $m \geq 6$.

Comments. The proof above is the core of Vinberg’s proof of nonexistence of compact hyperbolic Coxeter’s polyhedra of large dimensions.

Playing a bit more with the same inequalities, one gets nonexistence of right-angled hyperbolic polyhedra, in all dimensions starting from 5. In 4-dimensional case, an example of a bonded right-angled hyperbolic polyhedron can be found among regular *120-cells*.

Appendix B

Dictionary

Busemann function. Let X be a metric spaces and γ is a *geodesic* ray in X ; i.e., $\gamma: [0, \infty) \rightarrow X$ is a curve such that for any $t_0, t_1 \in [0, \infty)$, we have

$$|\gamma(t_0) - \gamma(t_1)|_X = |t_1 - t_0|$$

where $|x - y|_X$ denotes the distance from x to y in X . The Busemann function $b_\gamma: X \rightarrow \mathbb{R}$ is defined by

$$b_\gamma(p) = \lim_{t \rightarrow \infty} (|p - \gamma(t)|_X - t).$$

From the triangle inequality, the expression under the limit is nonincreasing in t ; therefore the limit above is well defined.

Curvature operator. The Riemannian curvature tensor R can be viewed as an operator \mathbf{R} on bi-vectors defined by

$$\langle \mathbf{R}(X \wedge Y), Z \wedge T \rangle = \langle R(X, Y)Z, T \rangle,$$

The operator $\mathbf{R}: \bigwedge^2 T \rightarrow \bigwedge^2 T$ is called *curvature operator* and it is said to be *positively defined* if $\langle \mathbf{R}(\varphi), \varphi \rangle > 0$ for all non zero bi-vector $\varphi \in \bigwedge^2 T$.

Dehn twist. Let Σ be a surface and $\gamma: \mathbb{R}/\mathbb{Z} \rightarrow \Sigma$ be noncontractible closed *simple curve*. Let U_γ be a neighborhood of γ which admits a homeomorphism $h: U_\gamma \rightarrow \mathbb{R}/\mathbb{Z} \times (0, 1)$. Dehn twist along γ is a homeomorphism $f: \Sigma \rightarrow \Sigma$ which is identity outside of U_γ and such that

$$h \circ f \circ h^{-1}: (x, y) \mapsto (x + y, y).$$

If Σ is orientable, then the Dehn twist described above is called *positive* if h is orientation preserving.

Dehn–Sommerville equations. Assume P is a *simple* Euclidean m -dimensional polyhedron; i.e., every vertex of P exactly m facets are meeting. Denote by f_k the number of k -dimensional faces of P ; the array (f_0, f_1, \dots, f_m) is called f -vector of P .

Fix an order of the vertices v_1, v_2, \dots, v_{f_0} of P so that for some linear function ℓ , we have $\ell(v_i) > \ell(v_j) \Leftrightarrow i < j$. The *index* of the vertex v_i is defined as the number of edges $[v_i v_j]$ such that $j < i$. The number of vertices of given index k will be denoted as h_k . the array (h_0, h_1, \dots, h_m) is called h -vector of P . Clearly $h_0 = h_m = 1$ and $k_k \geq 0$ for all k .

Each k -face contains unique vertex which maximize ℓ ; if the vertex has index i then $i \geq k$ and then it is the maximal vertex for exactly $\frac{i!}{k! \cdot (i-k)!}$ faces of dimension k . This observation can be packed in the following polynomial identity

$$\sum_k h_k \cdot (t+1)^k = \sum_k f_k \cdot t^k.$$

Note that the identity above implies that h -vector does not depend on the choice of order of the vertices. The Dehn–Sommerville equation can be written as

$$h_k = h_{m-k};$$

this identity obtained by reversing the order of vertices.

Doubling of a manifold M with boundary ∂M is is two copies of M_1, M_2 identified along corresponding points on their boundaries $\partial M_1, \partial M_2$.

Euclidean cone. Let Σ be a metric space with diameter $\leq \pi$. A metric space K is called Euclidean cone over Σ if its underling set coincides with the quotient $\Sigma \times [0, \infty)/\sim$ by the minimal equivalence relation \sim such that $(x, 0) \sim (y, 0)$ for any $x, y \in \Sigma$ and the metric is defined by cosine rule; i.e.,

$$|(x, a) - (y, b)|_K^2 = a^2 + b^2 - 2 \cdot a \cdot b \cdot \cos |x - y|_\Sigma.$$

Equidistant subsets. Two subsets A and B in a metric space are called equidistant if dist_A is constant on B and dist_B is constant on A .

Exponential map. Let (M, g) be a Riemannian manifold; denote by TM the tangent bundle over M and by $T_p = T_p M$ the tangent space at point $p \in M$.

Given a vector $v \in T_p M$ denote by γ_v the geodesic in (M, g) such that $\gamma(0) = p$ and $\gamma'(0) = v$. The map $\exp: TM \rightarrow M$ defined by $v \mapsto \gamma_v(1)$ is called exponential map.

The restriction of \exp to the T_p is called exponential map at p and denoted as \exp_p .

Given a smooth submanifold $S \subset M$; denote by NS the normal bundle over S . The restriction of \exp to NS is called normal exponential map of S and denoted as \exp_S .

Geodesic. Let X be a metric space and \mathbb{I} be a real interval. A locally isometric immersion $\gamma: \mathbb{I} \rightarrow X$ is called geodesic. In other words, γ is a geodesic if for any $t_0 \in \mathbb{I}$ we have

$$|\gamma(t) - \gamma(t')|_X = |t - t'|$$

for all $t, t' \in \mathbb{I}$ sufficiently close to t_0 . Note that in our definition geodesic has unit speed (that is not quite standard).

Kissing number. Let W_0 be a convex body in \mathbb{R}^n . The kissing number of W_0 is the maximal integer k such that there are k bodies W_1, W_2, \dots, W_k such that each W_i is congruent to W_0 , for each i we have $W_i \cap W_0 \neq \emptyset$ and have no pair W_i, W_j has common interior points.

Length space. A complete metric space X is called *length space* if the distance between any pair of points in X is equal to the infimum of lengths of curves connecting these points.

Macrodimension. Let X be a metric space $a > 0$ and n is an integer. We say that the macrodimension of X at the scale a is at most n if there is a map f from X to an n -dimensional simplicial complex K such that for any $k \in K$ the preimage $f^{-1}\{k\}$ has diameter less than a .

If macrodimension of X at the scale a is at most n , but not at most $n - 1$, we say that n is the macrodimension of X at the scale a .

Equivalently, the macrodimension of X on scale a can be defined as the least integer n such that X admits an open covering with diameter of each set less than a and such that each point in X is covered by at most $n + 1$ sets in the cover.

Length preserving map. A map $f: X \rightarrow Y$ of *length spaces* X and Y is a length preserving map if for any path $\alpha: [0, 1] \rightarrow X$, we have

$$\text{length}(\alpha) = \text{length}(f \circ \alpha).$$

Minimal surface. Let Σ be a k -dimensional smooth surface in a Riemannian manifold M and $T = T\Sigma$ and $N = N\Sigma$ correspondingly tangent and normal bundle. Let $s: T \otimes T \rightarrow N$ denotes the second fundamental form of Σ . Let e_i is an orthonormal basis for T_x , set $H_x = \sum_i s(e_i, e_i) \in N_x$; it is the mean curvature vector at $x \in \Sigma$. We say that Σ is *minimal* if $H \equiv 0$.

Minkowski space — \mathbb{R}^m with a metric induced by a norm.

Nil-manifolds form the minimal class of manifolds which includes a point, and has the following property: the total space of any principle \mathbb{S}^1 -bundle over a nil-manifold is a nil-manifold.

Any nil-manifold is diffeomorphic to the quotient of a connected nilpotent Lie group by a lattice.

The celebrated Gromov's theorem states that almost flat manifolds admit a finite folded cover by a nil-manifold

Polyhedral space — a simplicial complex with a metric such that each simplex is isometric to a simplex in a Euclidean space.

It admits the following generalizations: *spherical (hyperbolic)* polyhedral space — a simplicial complex with a metric such that each simplex is isometric to a simplex in a unit sphere (correspondingly hyperbolic space of constant curvature -1).

A point in a polyhedral space is called *regular* if it has a neighborhood isometric to an open set in a Euclidean space; otherwise it is called *singular*. To define regular/singular points in spherical or hyperbolic space, one has to exchange in the above definition Euclidean space to unit sphere or hyperbolic space with curvature -1 .

Polynomial volume growth. A Riemannian manifold M has polynomial volume growth if for some (and therefore any) $p \in M$, we have

$$\text{vol } B_r(p) \leq C \cdot (r^k + 1),$$

where $B_r(p)$ is the ball in M and C, k are real constants.

Proper metric space. A metric space X is called *proper* if any closed bounded set in X is compact.

Piecewise linear isometry — a piecewise linear map from a polyhedral space which is isometric on each simplex.

More precisely: Let P and Q be polyhedral spaces, a map $f: P \rightarrow Q$ is called piecewise linear isometry if there is a triangulation \mathcal{T} of P such that at any simplex $\Delta \in \mathcal{T}$ the restriction $f|_{\Delta}$ is globally isometric.

PL-homeomorphism or piecewise linear homeomorphism. A map $h: P \rightarrow Q$ between polyhedral spaces P and Q is called PL-homeomorphism if it is a homeomorphism and both spaces P and Q admit triangulations such that each simplex of P is mapped to a simplex of Q by an affine map.

Quasi-isometry. A map $f: X \rightarrow Y$ is called a quasi-isometry if there is a positive real constant C such that $f(X)$ is a C -net in Y and

$$\frac{1}{C} \cdot |x - y|_X - C \leq |f(x) - f(y)|_Y \leq C \cdot |x - y|_X + C.$$

Note that a quasi-isometry is not assumed to be continuous, for example any map between compact metric spaces is a quasi-isometry.

Saddle surface. A smooth surface Σ in \mathbb{R}^3 is saddle (correspondingly strictly saddle) if the product of the principle curvatures at each point is ≤ 0 (correspondingly < 0).

It admits the following generalization to non-smooth case and arbitrary dimension of the ambient space: A surface Σ in \mathbb{R}^n is saddle if the restriction $\ell|_{\Sigma}$ of any linear function $\ell: \mathbb{R}^3 \rightarrow \mathbb{R}$ has no strict local minima at interior points of Σ .

One can generalize it further to an arbitrary ambient space, using convex functions instead of linear functions in the above definition.

Sasaki metric. Let (M, g) be a Riemannian manifold. The Sasaki metric is the most natural choice of metric on the tangent space TM . It is uniquely defined by the following properties:

- (i) The natural projection $\tau: TM \rightarrow M$ is a Riemannian submersion.
- (ii) The metric on each tangent space $T_p \subset TM$ is the Euclidean metric induced by g .
- (iii) Assume $\gamma(t)$ is a curve in M and $v(t) \in T_{\gamma(t)}$ is a parallel vector field along γ . Note that $v(t)$ forms a curve in TM and $T_{\gamma(t)}M$ forms a submanifold in TM . For the Sasaki metric, we have $\dot{v}(t) \perp T_{\gamma(t)}M$ for any t .

A more constructive way to describe Sasaki metric is given by identifying $T_u TM$ for any $u \in T_p M$ with the direct sum of so called vertical and horizontal vectors $T_p M \oplus T_p M$. The projection of this splitting defined by the differential of τ and the Levi-Civita connection. Then $T_u TM$ is equipped with the metric defined as

$$\hat{g}(X, Y) = g(X^V, Y^V) + g(X^H, Y^H),$$

where $X^V, X^H \in T_p$ denotes the vertical and horizontal components of $X \in T_u TM$.

Second fundamental form. Assume $f: M \hookrightarrow \mathbb{R}^n$ be an immersion of smooth manifold M . Given a point $p \in M$ denote by T_p and $N_p = T_p^\perp$ the tangent and normal spaces of L at p . The second fundamental form for f at $p \in M$ is defined as

$$s(v, w) = (\nabla_v w)^\perp, \quad (*)$$

where $(\nabla_v w)^\perp$ denotes the orthogonal projection of covariant derivative $\nabla_v w$ onto the normal bundle.

Assume $\gamma_v: \mathbb{R} \rightarrow M$ is a geodesic with tangent vector $v \in T_p$; i.e., such that $\gamma_v(0) = p$ and $\gamma'_v(0) = v$. Then

$$s(v, v) = (f \circ \gamma_v)''(0).$$

This property can be also used to define second fundamental form via the identity

$$s(v, w) = \frac{1}{2} \cdot [s(v + w, v + w) - s(v, v) - s(w, w)].$$

The formula (*) can be used to define the second fundamental form for smooth immersions from into Riemannian manifold.

Short map — the same as 1-Lipschitz of distance nonexpanding map.

Simple curve — an image of a continuous injective map of a real segment or a circle in a topological space.

Sub-Riemannian metric. Let (M, g) is a Riemannian manifold.

Assume that in the tangent bundle TM a choice of sub-bundle H is given; the sub-bundle H which will be called *horizontal distribution*. The tangent vectors which lie in H will be called *horizontal*. A piecewise smooth curve will be called *horizontal* if all its tangent vectors are horizontal.

The sub-Riemannian distance between points x and y is defined as infimum of lengths of horizontal curves connecting x to y .

Alternatively, the distance can be defined as a limit of Riemannian distances for the metrics $g_\lambda(X, Y) = g(X^h, Y^h) + \lambda \cdot g(X^v, Y^v)$ as $\lambda \rightarrow \infty$, where X^h denotes the horizontal part of X ; i.e., the orthogonal projection of X to H and X^v denotes the vertical part of X ; i.e., $X^v = X - X^h$.

One usually adds a condition which ensure that any curve in M can be arbitrary well approximated by a horizontal curve with the same endpoints. (In particular this ensures that the distance will not take infinite values.) The most common condition is so called *complete non-integrability*; it means that for any $x \in M$, one can choose a basis in $T_x M$ from the vectors of the following type: $A(x)$, $[A, B](x)$, $[A, [B, C]](x)$, $[A, [B, [C, D]]](x)$, ... where all vector fields A, B, C, D, \dots are horizontal.

Variation of turn. Let $\gamma: [a, b] \rightarrow \mathbb{R}^n$ be a curve. The variation of turn of γ is defined as supremum of sum of external angles for broken lines inscribed in γ . Namely,

$$\sup \left\{ \sum_{i=1}^{n-1} \alpha_i \mid a = t_0 < t_1 < \dots < t_n = b \right\},$$

where $\alpha_i = \pi - \angle \gamma(t_{i-1}) \gamma(t_i) \gamma(t_{i+1})$.

Bibliography

- [1] J. Lohkamp, ???.
- [2] A. Lytchak, ???.
- [3] Uwe Abresch and Detlef Gromoll, *On complete manifolds with nonnegative Ricci curvature*, J. Amer. Math. Soc. **3** (1990), no. 2, 355–374.
- [4] H. Alexander and R. Osserman, *Area bounds for various classes of surfaces*, Amer. J. Math. **97** (1975), no. 3, 753–769.
- [5] H. Alexander, D. Hoffman, and R. Osserman, *Area estimates for submanifolds of Euclidean space*, Symposia Mathematica, Vol. XIV (Convegno di Teoria Geometrica dell'Integrazione e Varietà Minimali, INDAM, Rome, 1973), Academic Press, London, 1974, pp. 445–455.
- [6] Ralph Alexander, *Lipschitzian mappings and total mean curvature of polyhedral surfaces. I*, Trans. Amer. Math. Soc. **288** (1985), no. 2, 661–678.
- [7] S. Alexander, *Locally convex hypersurfaces of negatively curved spaces*, Proc. Amer. Math. Soc. **64** (1977), no. 2, 321–325.
- [8] Stephanie B. Alexander, I. David Berg, and Richard L. Bishop, *Geometric curvature bounds in Riemannian manifolds with boundary*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 703–716.
- [9] S. Alexander, V. Kapovitch, and Petrunin A., *Alexandrov geometry*, in preparation.
- [10] A. D. Aleksandrov, *Vnutrennyaya Geometriya Vypuklykh Poverhnosteĭ*, OGIZ, Moscow-Leningrad, 1948.
- [11] F. Almgren, *Optimal isoperimetric inequalities*, Indiana Univ. Math. J. **35** (1986), no. 3, 451–547.
- [12] Ben Andrews, *Contraction of convex hypersurfaces in Riemannian spaces*, J. Differential Geom. **39** (1994), no. 2, 407–431.
- [13] Florent Balacheff, Christopher Croke, and Mikhail G. Katz, *A Zoll counterexample to a geodesic length conjecture*, Geom. Funct. Anal. **19** (2009), no. 1, 1–10.
- [14] Victor Bangert, *Geodesics and totally convex sets on surfaces*, Invent. Math. **63** (1981), no. 3, 507–517.
- [15] I. D. Berg, *An estimate on the total curvature of a geodesic in Euclidean 3-space-with-boundary*, Geom. Dedicata **13** (1982), no. 1, 1–6.
- [16] A. S. Besicovitch, *On two problems of Loewner*, J. London Math. Soc. **27** (1952), 141–144.

- [17] Károly Bezdek and Robert Connelly, *Pushing disks apart—the Kneser-Poulsen conjecture in the plane*, J. Reine Angew. Math. **553** (2002), 221–236.
- [18] R. H. Bing, *Some aspects of the topology of 3-manifolds related to the Poincaré conjecture*, Lectures on modern mathematics, Vol. II, Wiley, New York, 1964, pp. 93–128.
- [19] S. Bochner, *Vector fields and Ricci curvature*, Bull. Amer. Math. Soc. **52** (1946), 776–797. MR0018022 (8,230a)
- [20] Christoph Böhm and Burkhard Wilking, *Manifolds with positive curvature operators are space forms*, Ann. of Math. (2) **167** (2008), no. 3, 1079–1097.
- [21] B. H. Bowditch, *Notes on locally CAT(1) spaces*, Geometric group theory (Columbus, OH, 1992), Ohio State Univ. Math. Res. Inst. Publ., vol. 3, de Gruyter, Berlin, 1995, pp. 1–48. MR1355101 (97e:53070)
- [22] Ulrich Brehm, *Extensions of distance reducing mappings to piecewise congruent mappings on \mathbf{R}^m* , J. Geom. **16** (1981), no. 2, 187–193.
- [23] Simon Brendle, Fernando C. Marques, and Andre Neves, *Deformations of the hemisphere that increase scalar curvature*, Invent. Math. **185** (2011), no. 1, 175–197.
- [24] Dmitri Burago, Yuri Burago, and Sergei Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001.
- [25] Yu. D. Burago and V. A. Zalgaller, *Geometric inequalities*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 285, Springer-Verlag, Berlin, 1988. Translated from the Russian by A. B. Sosinskiĭ; Springer Series in Soviet Mathematics. MR936419 (89b:52020)
- [26] D. Burago, S. Ivanov, and D. Shoenthal, *Two counterexamples in low-dimensional length geometry*, Algebra i Analiz **19** (2007), no. 1, 46–59.
- [27] D. Burago and B. Kleiner, *Rectifying separated nets*, Geom. Funct. Anal. **12** (2002), no. 1, 80–92.
- [28] S. V. Buyalo, *Volume and fundamental group of a manifold of nonpositive curvature*, Mat. Sb. (N.S.) **122(164)** (1983), no. 2, 142–156 (Russian).
- [29] Zarathustra Elessar Brady, *Is it possible to capture a sphere in a knot?*. URL: <http://mathoverflow.net/q/8091>.
- [30] Aleksander Całka, *On conditions under which isometries have bounded orbits*, Colloq. Math. **48** (1984), no. 2, 219–227.
- [31] Jeff Cheeger and Tobias H. Colding, *Lower bounds on Ricci curvature and the almost rigidity of warped products*, Ann. of Math. (2) **144** (1996), no. 1, 189–237, DOI 10.2307/2118589. MR1405949 (97h:53038)
- [32] Christopher B. Croke, *Small volume on big n -spheres*, Proc. Amer. Math. Soc. **136** (2008), no. 2, 715–717.
- [33] Michael Edelstein and Binyamin Schwarz, *On the length of linked curves*, Israel J. Math. **23** (1976), no. 1, 94–95.
- [34] N. V. Efimov, *Qualitative problems of the theory of deformation of surfaces*, Uspehi Matem. Nauk (N.S.) **3** (1948), no. 2(24), 47–158.
- [35] J.-H. Eschenburg, *Local convexity and nonnegative curvature—Gromov’s proof of the sphere theorem*, Invent. Math. **84** (1986), no. 3, 507–522.
- [36] K. J. Falconer, *The geometry of fractal sets*, Cambridge Tracts in Mathematics, vol. 85, Cambridge University Press, Cambridge, 1986.

- [37] Fuquan Fang, Sérgio Mendonça, and Xiaochun Rong, *A connectedness principle in the geometry of positive curvature*, Comm. Anal. Geom. **13** (2005), no. 4, 671–695.
- [38] S. Frankel and M. Katz, *The Morse landscape of a Riemannian disk*, Ann. Inst. Fourier (Grenoble) **43** (1993), no. 2, 503–507.
- [39] T. Frankel, *On the fundamental group of a compact minimal submanifold*, Ann. of Math. (2) **83** (1966), 68–73.
- [40] G. Galperin, *Convex polyhedra without simple closed geodesics*, Regul. Chaotic Dyn. **8** (2003), no. 1, 45–58.
- [41] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, resultants and multidimensional determinants*, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2008. Reprint of the 1994 edition. MR2394437 (2009a:14065)
- [42] Viktor L. Ginzburg, *On the existence and non-existence of closed trajectories for some Hamiltonian flows*, Math. Z. **223** (1996), no. 3, 397–409.
- [43] R. E. Greene and H. Wu, *On the subharmonicity and plurisubharmonicity of geodesically convex functions*, Indiana Univ. Math. J. **22** (1972/73), 641–653.
- [44] M. Gromov, *Almost flat manifolds*, J. Differential Geom. **13** (1978), no. 2, 231–241.
- [45] Mikhael Gromov, *Filling Riemannian manifolds*, J. Differential Geom. **18** (1983), no. 1, 1–147. MR697984 (85h:53029)
- [46] ———, *Partial differential relations*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 9, Springer-Verlag, Berlin, 1986. MR864505 (90a:58201)
- [47] M. Gromov, *Sign and geometric meaning of curvature*, Rend. Sem. Mat. Fis. Milano **61** (1991), 9–123 (English, with English and Italian summaries).
- [48] Misha Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2007.
- [49] Karsten Grove, *Geometry of, and via, symmetries*, Conformal, Riemannian and Lagrangian geometry (Knoxville, TN, 2000), Univ. Lecture Ser., vol. 27, Amer. Math. Soc., Providence, RI, 2002, pp. 31–53.
- [50] Karsten Grove and Burkhard Wilking, *A knot characterization and 1-connected nonnegatively curved 4-manifolds with circle symmetry*, Geom. Topol. **18** (2014), no. 5, 3091–3110.
- [51] Larry Guth, *Symplectic embeddings of polydisks*, Invent. Math. **172** (2008), no. 3, 477–489.
- [52] G. Guzhvina, *Gromov’s pinching constant*, arXiv:0804.0201 [math.DG].
- [53] Fengbo Hang and Xiaodong Wang, *Rigidity theorems for compact manifolds with boundary and positive Ricci curvature*, J. Geom. Anal. **19** (2009), no. 3, 628–642.
- [54] Charles J. A. Halberg Jr., Eugene Levin, and E. G. Straus, *On contiguous congruent sets in Euclidean space*, Proc. Amer. Math. Soc. **10** (1959), 335–344.
- [55] Joel David Hamkins, *Is the sphere the only surface all of whose projections are circles? Or: Can we deduce a spherical Earth by observing that its shadows on the Moon are always circular?*, MathOverflow. URL: <http://mathoverflow.net/q/39127>.

- [56] David Hilbert, *Ueber die gerade Linie als kürzeste Verbindung zweier Punkte*, Math. Ann. **46** (1895), no. 1, 91–96.
- [57] ———, *Mathematical problems*, Bull. Amer. Math. Soc. **8** (1902), no. 10, 437–479.
- [58] Wu-Yi Hsiang and Bruce Kleiner, *On the topology of positively curved 4-manifolds with symmetry*, J. Differential Geom. **29** (1989), no. 3, 615–621.
- [59] Imre Bárány, Krystyna Kuperberg, and Tudor Zamfirescu, *Total curvature and spiralling shortest paths*, Discrete Comput. Geom. **30** (2003), no. 2, 167–176. U.S.-Hungarian Workshops on Discrete Geometry and Convexity (Budapest, 1999/Auburn, AL, 2000).
- [60] Ivan Izvestiev, Robert B. Kusner, Günter Rote, Boris Springborn, and John M. Sullivan, *There is no triangulation of the torus with vertex degrees 5, 6, ..., 6, 7 and related results: geometric proofs for combinatorial theorems*, Geom. Dedicata **166** (2013), 15–29.
- [61] S. Jendrol' and E. Jucovič, *On the toroidal analogue of Eberhard's theorem*, Proc. London Math. Soc. (3) **25** (1972), 385–398.
- [62] M.D. Kirszbraum, *Über die zusammenziehenden und Lipschitzschen Transformationen.*, Fundam. Math. **22** (1934), 77–108.
- [63] V. L. Klee Jr., *Some topological properties of convex sets*, Trans. Amer. Math. Soc. **78** (1955), 30–45.
- [64] A. Kneser, *Bemerkungen über die Anzahl der Extreme der Krümmung auf geschlossenen Kurven und über verwandte Fragen in einer nichteuklidischen Geometrie*, Festschrift H. Weber (1912), 170–180.
- [65] Nicolaas H. Kuiper, *On C^1 -isometric imbeddings. I, II*, Nederl. Akad. Wetensch. Proc. Ser. A. **58** = Indag. Math. **17** (1955), 545–556, 683–689.
- [66] Kyung Whan Kwun, *Uniqueness of the open cone neighborhood*, Proc. Amer. Math. Soc. **15** (1964), 476–479.
- [67] Christian Lange, *When is the underlying space of an orbifold a topological manifold*, [arXiv:1307.4875 \[math.GN\]](#).
- [68] N. Lebedeva and A. Petrunin, *Local characterization of polyhedral spaces*, Geometriae Dedicata (to appear) [arXiv:1402.6670 \[math.DG\]](#).
- [69] Enrico Le Donne, *Lipschitz and path isometric embeddings of metric spaces*, Geom. Dedicata **166** (2013), 47–66.
- [70] J. Liberman, *Geodesic lines on convex surfaces*, C. R. (Doklady) Acad. Sci. URSS (N.S.) **32** (1941), 310–313.
- [71] N. I. Lobachevsky, *Geometrische Untersuchungen zur Theorie der Parallelinien*, Berlin: F. Fincke, 1840.
- [72] Rostislav Matveev, *Surfaces with polyhedral metrics*, International Mathematical Summer School for Students 2011 (Jacobs University, Bremen).
- [73] Mario J. Micallef and John Douglas Moore, *Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes*, Ann. of Math. (2) **127** (1988), no. 1, 199–227.
- [74] M. A. Mikhaïlova, *A factor space with respect to the action of a finite group generated by pseudoreflections*, Izv. Akad. Nauk SSSR Ser. Mat. **48** (1984), no. 1, 104–126.
- [75] A. D. Milka, *Multidimensional spaces with polyhedral metric of nonnegative curvature. I*, Ukrain. Geometr. Sb. Vyp. **5–6** (1968), 103–114.

- [76] ———, *Shortest lines on convex surfaces*, Dokl. Akad. Nauk SSSR **248** (1979), no. 1, 34–36.
- [77] Vitali D. Milman and Gideon Schechtman, *Asymptotic theory of finite-dimensional normed spaces*, Lecture Notes in Mathematics, vol. 1200, Springer-Verlag, Berlin, 1986. With an appendix by M. Gromov.
- [78] Marshall Hall Jr., *Subgroups of finite index in free groups*, Canadian J. Math. **1** (1949), 187–190.
- [79] Wayne Lewis, *The pseudo-arc*, Bol. Soc. Mat. Mexicana (3) **5** (1999), no. 1, 25–77.
- [80] S. Mazur and S. Ulam, *Sur les transformations isométriques des espaces vectoriels normés*, C. R. Acad. Sci., Paris **194** (1932), no. 1, 946–948.
- [81] Alexander Nabutovsky and Regina Rotman, *Length of geodesics and quantitative Morse theory on loop spaces*, Geom. Funct. Anal. **23** (2013), no. 1, 367–414.
- [82] John Nash, *C^1 isometric imbeddings*, Ann. of Math. (2) **60** (1954), 383–396.
- [83] V. Ovsienko and S. Tabachnikov, *Projective differential geometry old and new*, Cambridge Tracts in Mathematics, vol. 165, Cambridge University Press, Cambridge, 2005.
- [84] Joseph O’Rourke, *Why is the half-torus rigid?*.
URL: <http://mathoverflow.net/q/77760>.
- [85] Dmitri Panov, *Polyhedral Kähler manifolds*, Geom. Topol. **13** (2009), no. 4, 2205–2252.
- [86] ———, *Foliations with unbounded deviation on \mathbb{T}^2* , J. Mod. Dyn. **3** (2009), no. 4, 589–594.
- [87] ———, *Parabolic curves and gradient mappings*, Proc. Steklov Inst. Math. **2** (221) (1998), 261–278.
- [88] Dmitri Panov and Anton Petrunin, *Sweeping out sectional curvature*, Geom. Topol. **18** (2014), no. 2, 617–631.
- [89] ———, *Ramification conjecture*, arXiv:1312.6856 [math.GT].
- [90] G. Perelman, *Proof of the soul conjecture of Cheeger and Gromoll*, J. Differential Geom. **40** (1994), no. 1, 209–212.
- [91] Anton Petrunin, *Two discs with no parallel tangent planes*.
URL: <http://mathoverflow.net/q/17486>.
- [92] ———, *One-step problems in geometry*.
URL: <http://mathoverflow.net/q/8247>.
- [93] ———, *Diameter of m -fold cover*.
URL: <http://mathoverflow.net/q/7732>.
- [94] A. Petrunin, *Intrinsic isometries in Euclidean space*, St. Petersburg Math. J. **22** (2011), no. 5, 803–812.
- [95] ———, *Area minimizing polyhedral surfaces are saddle*, American Mathematical Monthly, to appear.
- [96] Petya Pushkar, *A generalization of Cauchy’s mean value theorem*.
URL: <http://mathoverflow.net/q/16335>.
- [97] Eduard Rembs, *Verbiegungen höherer Ordnung und ebene Flächenrinnen.*, Math. Z. **36** (1932), 110–121.

- [98] V. Sahovic, *Approximations of Riemannian Manifolds with Linear Curvature Constraints*, Dissertation, Westfälische Wilhelms-Universität Münster, 2009.
- [99] L. C. Siebenmann, *Deformation of homeomorphisms on stratified sets. I, II*, Comment. Math. Helv. **47** (1972), 123–136; *ibid.* **47** (1972), 137–163. MR0319207 (47 #7752)
- [100] A. Shen, *Unexpected proofs: Boxes in a Train*, Math. Intelligencer **21** (1999), no. 3, 48–50.
- [101] John R. Stallings, *Topology of finite graphs*, Invent. Math. **71** (1983), no. 3, 551–565.
- [102] Dennis Sullivan, *Hyperbolic geometry and homeomorphisms*, Geometric topology (Proc. Georgia Topology Conf., Athens, Ga., 1977), Academic Press, New York-London, 1979, pp. 543–555.
- [103] J. L. Synge, *On the connectivity of spaces of positive curvature*, C. R. Congr. internat. Math., Oslo 1936, **2** (1937), 138–139.
- [104] Serge Tabachnikov, *The tale of a geometric inequality*, MASS selecta, Amer. Math. Soc., Providence, RI, 2003, pp. 257–262.
- [105] ———, end of 80's.
- [106] ———, *The (Un)equal Tangents Problem*, The American Mathematical Monthly **119** (May 2012), no. 5, 398–405.
- [107] P. Tait, *Note on the circles of curvature of a plane curve*, Proc. Edinburgh Math. Soc. **14** (February 1895), 26.
- [108] Pekka Tukia and Jussi Väisälä, *Bi-Lipschitz extensions of maps having quasiconformal extensions*, Math. Ann. **269** (1984), no. 4, 561–572, DOI 10.1007/BF01450765. MR766014 (86c:30041)
- [109] Jussi Väisälä, *A proof of the Mazur-Ulam theorem*, Amer. Math. Monthly **110** (2003), no. 7, 633–635.
- [110] F. A. Valentine, *On the extension of a vector function so as to preserve a Lipschitz condition*, Bull. Amer. Math. Soc. **49** (1943).
- [111] Genevieve Walsh, *Great circle links in the three-sphere*, Ph.D. thesis, 2003.
- [112] H. Weyl, *On the volume of tubes*, Amer. J. Math. **61** (1939), 461472.
- [113] Alan Weinstein, *Positively curved n -manifolds in \mathbb{R}^{n+2}* , J. Differential Geometry **4** (1970), 1–4.
- [114] Burkhard Wilking, *On fundamental groups of manifolds of nonnegative curvature*, Differential Geom. Appl. **13** (2000), no. 2, 129–165.
- [115] ———, *Torus actions on manifolds of positive sectional curvature*, Acta Math. **191** (2003), no. 2, 259–297.
- [116] Henry Wilton, *In Memoriam J. R. Stallings — Topology of Finite Graphs* (2008), <https://1dtopology.wordpress.com/2008/12/01/>.
- [117] V. A. Zalgaller, *On deformations of a polygon on a sphere*, Uspehi Mat. Nauk (N.S.) **11** (1956), no. 5(71), 177–178 (Russian).
- [118] ———, *Isometric imbedding of polyhedra*, Dokl. Akad. Nauk SSSR **123** (1958), 599–601 (Russian).
- [119] Shing Tung Yau, *Non-existence of continuous convex functions on certain Riemannian manifolds*, Math. Ann. **207** (1974), 269–270.
- [120] Kunizô Yoneyama, *Theory of Continuous Set of Points*, Tohoku Mathematical Journal, First Series **12** (1917), 43–158.