

# ON THE TOTAL CURVATURE OF MINIMIZING GEODESICS ON CONVEX SURFACES

NINA LEBEDEVA AND ANTON PETRUNIN

ABSTRACT. We give a universal upper bound for the total curvature of minimizing geodesic on a convex surface in the Euclidean space.

## 1. INTRODUCTION

**1.1. Main theorem.** *The total curvature of a minimizing geodesic on a convex surface in  $\mathbb{R}^3$  can not exceed  $10^{10}$ .*

The main theorem was stated as an open question in [1], [2] and [3], but we have learned it from Dmitry Burago only few years ago.

Let us briefly discuss the related results.

- ◊ In [4], Liberman gives a bound on the total curvature of short geodesic in terms of the ratio diameter and inradius of  $K$ . In the proof he use now so called Liberman's lemma 3.1 discussed below. This statement was rediscovered in [3].
- ◊ In [5], Usov gives the optimal bound for total curvature of geodesic on the graph of  $\ell$ -Lipschitz convex function. Namely, he proves that if  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  is  $\ell$ -Lipschitz and convex then any geodesic in its graph

$$\Gamma_f = \{ (x, y, z) \in \mathbb{R}^3 \mid z = f(x, y) \}$$

has total curvature at most  $2 \cdot \ell$ . This statement was also rediscovered in [3]. Yet an amusing generalization of Usov's result is given by Berg in [6].

- ◊ In [7], Pogorelov conjectured that any the spherical image of geodesic on convex surface has to be contractible. It is easy to see that the length of spherical image of geodesic can not be smaller than its total curvature, so this conjecture (if it would be true) would be stronger than Liberman's theorem. Counterexamples were found indepenently by Milka in [8], Usov in [9] and yet much later rediscovered by Pach in [2].
- ◊ In [3], Bárány, Kuperberg, and Zamfirescu have constructed a corkscrew minimizing geodesic on a closed hypersurface; that is a minimizing geodesic which twists around given line arbitrary many times. In the same paper they also constructed a minimizing geodesic on a convex surface in  $\mathbb{R}^3$  with total curvature bigger than  $2 \cdot \pi$ . (Note that  $2 \cdot \pi$  is the optimal bound for the analogous problem in the plane.)

## 2. PRELIMINARIES

**Semicontinuity of total curvature.** Recall that the *total curvature* of a curve  $\gamma: [0, \ell] \rightarrow \mathbb{R}^3$  (briefly  $\text{TotCurv } \gamma$ ) is defined as supremum of sum of exterior angles for the polygonal lines inscribed in  $\gamma$ .

---

1991 *Mathematics Subject Classification.* 53A05, 53C45, 53C22.

N. Lebedeva was partially supported by RFBR grant 14-01-00062.

A. Petrunin was partially supported by NSF grant DMS 1309340.

Note that for a polygonal line  $\sigma$ , its total curvature coincide with the sum of its exterior angles.

If  $\gamma$  is a smooth curve with unit-speed parametrization, then

$$\text{TotCurv } \gamma = \int_0^\ell \kappa(t) \cdot dt,$$

where  $\kappa(t) = |\ddot{\gamma}(t)|$  is the curvature of  $\gamma$  at  $t$ .

**2.1. Proposition.** *Assume  $\gamma_n: \mathbb{I} \rightarrow \mathbb{R}^3$  is a sequence of curves converging pointwise to a curve  $\gamma_\infty: \mathbb{I} \rightarrow \mathbb{R}^3$ . Then*

$$\liminf_{n \rightarrow \infty} \text{TotCurv } \gamma_n \geq \text{TotCurv } \gamma_\infty.$$

*Proof.* Choose a polygonal line  $\sigma_\infty$  inscribed in  $\gamma_\infty$  with total curvature sufficiently close to  $\text{TotCurv } \gamma_\infty$ . Let  $\gamma_\infty(t_0), \dots, \gamma_\infty(t_k)$  be the vertices of  $\sigma_\infty$ . Consider the polygonal lines  $\sigma_n$  with the vertices  $\gamma_n(t_0), \dots, \gamma_n(t_k)$ . Note that

$$\text{TotCurv } \sigma_n \rightarrow \text{TotCurv } \sigma_\infty.$$

Hence the statement follows.  $\square$

**Convergence.** Given a closed set  $\Sigma \subset \mathbb{R}^3$ , denote by  $\text{dist}_\Sigma$  the distance function from  $\Sigma$ ; that is

$$\text{dist}_\Sigma(x) = \inf \{ |x - y| \mid y \in \Sigma \}.$$

We say that a sequence of closed sets  $\Sigma_n \subset \mathbb{R}^3$  converges to the closed set  $\Sigma_\infty \subset \mathbb{R}^3$  if for any  $x \in \mathbb{R}^3$ , we have  $\text{dist}_{\Sigma_n}(x) \rightarrow \text{dist}_{\Sigma_\infty}(x)$  as  $n \rightarrow \infty$ .

**Convex surfaces.** By convex surface in the Euclidean 3-space  $\mathbb{R}^3$  we understand the boundary of closed convex set with nonempty interior.

**2.2. Proposition.** *Assume  $\Sigma_n$  be a sequence of convex surfaces which converge to a convex surface  $\Sigma_\infty$ . Then for any minimizing geodesic  $\gamma_\infty$  in  $\Sigma_\infty$  there is a sequence of minimizing geodesics  $\gamma_n$  in  $\Sigma_n$  which pointwise converge to  $\gamma_\infty$  as  $n \rightarrow \infty$ .*

*Proof.* Assume  $\gamma_\infty$  parametrized by its length  $[0, \ell]$ .

Fix a subinterval  $[a, b] \subset [0, \ell]$  such that  $0 < a$  and  $b < \ell$ . Set  $p_\infty = \gamma_\infty(a)$  and  $q_\infty = \gamma_\infty(b)$ .

Let  $p_n, q_n \in \Sigma_n$  be a two sequences of points which converge to  $p_\infty$  and  $q_\infty$  correspondingly.

Denote by  $\gamma_n$  a minimizing geodesic from  $p_n$  to  $q_n$  in  $\Sigma_n$ . Note that  $\gamma_n$  converges to  $\gamma_\infty|_{[a, b]}$  as  $n \rightarrow \infty$ .

Taking the subinterval  $[a, b]$  closer and closer to  $[0, \ell]$  and applying diagonal procedure, we get the result.  $\square$

### 3. LIBERMAN'S LEMMA.

In this section we give a slight generalization of the construction of Liberman in [4].

**Development.** Let  $\gamma: [0, \ell] \rightarrow \mathbb{R}^3$  be a curve parametrized by length and a point  $p$  does not lie on  $\gamma$ .

Assume  $\tilde{\gamma}_p: [0, \ell] \rightarrow \mathbb{R}^2$  is a plane curve parametrized by length and  $\tilde{p}$  is a point in the plane such that

$$|\tilde{p} - \tilde{\gamma}(t)| = |\tilde{p} - \tilde{\gamma}(t)|$$

for any  $t \in [0, \ell]$  and the direction from  $\tilde{p}$  to  $\tilde{\gamma}(t)$  changes counterclockwise. Then  $\tilde{\gamma}_p$  is called *development* of  $\gamma$  with respect to  $p$ .

We say that the development  $\tilde{\gamma}_p$  is convex in the interval  $[a, b]$  if the curvilinear triangle bounded by arc  $\tilde{\gamma}|_{[a, b]}$  and the segments  $[\tilde{p}\tilde{\gamma}(a)]$ ,  $[\tilde{p}\tilde{\gamma}(b)]$  is convex. Correspondingly we say that  $\tilde{\gamma}$  is convex in the interval  $[a, b]$  if the complement of this triangle in the angle bounded by half-lines  $[\tilde{p}\tilde{\gamma}(a))$  and  $[\tilde{p}\tilde{\gamma}(b))$  is convex.

We say that  $\tilde{\gamma}_p$  is locally convex (concave) in the interval  $[a, b]$  if any point  $x \in [a, b]$  admits a closed neighborhood  $[a', b']$  in  $[a, b]$  such that  $\tilde{\gamma}_p$  is convex (correspondingly concave) in the interval  $[a', b']$ .

If we pass to the limit of this construction as  $p$  moves to infinity along a half-line in the direction of a unit vector  $u$  then the limit curve is called development of  $\gamma$  in direction  $u$  and denoted as  $\tilde{\gamma}_u$ .

The development  $\tilde{\gamma}_u$  could be also defined directly. Namely assume a curve  $\tilde{\gamma}_u: [0, \ell] \rightarrow \mathbb{R}^2$  is parametrized by length and for a fixed unit vector  $\tilde{u} \in \mathbb{R}^2$ , we have

$$\langle \tilde{u}, \tilde{\gamma}_u(t) \rangle = \langle u, \gamma_u(t) \rangle$$

for any  $t \in [0, \ell]$  and the projection of  $\tilde{\gamma}_u(t)$  to the line normal to  $\tilde{u}$  is monotonic in  $t$ . Then the curve  $\tilde{\gamma}_u$  is called development of  $\gamma$  in the direction  $u$ .

We can assume that  $\tilde{u}$  is the vertical vector in the coordinate plane. In this case  $\tilde{\gamma}_u$  is a graph of a function  $y = f(x)$  defined in some real interval. We say that  $\tilde{\gamma}_u$  is concave (convex) in the interval  $[a, b]$  if the function  $f$  is concave (correspondingly convex) in the corresponding real interval.

**Dark and light sides.** Let  $\Sigma \subset \mathbb{R}^3$  be a smooth convex surface and  $z \notin \Sigma$  and  $p \in \Sigma$ .

We say that  $p$  lies on the dark (light) side of  $\Sigma$  with from  $z$  if all the points  $p + t \cdot (p - z)$  lie outside of  $\Sigma$  for  $t > 0$  (correspondingly for  $t < 0$ ). The intersection of dark and light side is called horizon of  $z$ ; it is denoted by  $\omega_z$ .

Note that if  $z$  lies inside  $\Sigma$  then all the points on  $\Sigma$  lies on dark side from  $z$ .

If  $\Sigma$  is smooth we can define the outer normal vector  $\nu_p$  to  $\Sigma$  at  $p$ . In this case  $p$  lies on dark (light) side of  $\Sigma$  with from  $z$  if and only if  $\langle z - p, \nu_p \rangle \geq 0$  (correspondingly for  $\langle z - p, \nu_p \rangle \leq 0$ ). If in addition  $\Sigma$  is strongly convex then the horizon is formed by a collection of disjoint smooth curves.

We could also define light/dark sides and horizon in the limit case, if  $p$  escapes to infinity along a half-line in direction  $u$ . Let us also define it directly, we say that a point  $p \in \Sigma$  lies on dark (light) side for the unit vector  $u$  if all the points  $p + u \cdot t$  lie outside of  $\Sigma$  for all  $t > 0$ , (correspondingly  $t < 0$ ). As before the intersection of light and dark side is called horizon and denoted by  $\omega_u$ .

In the smooth case the later means that  $\langle \nu_p, u \rangle \geq 0$  (correspondingly  $\langle \nu_p, u \rangle \leq 0$ ). If  $\Sigma$  is strongly convex then  $\omega_u$  is locally a smooth curve.

**3.1. Liberman's Lemma.** Assume  $\gamma$  be a geodesic on convex surface  $\Sigma \subset \mathbb{R}^3$ . Then for any point  $z \notin \Sigma$  the development  $\tilde{\gamma}_z$  is locally convex (locally concave) if  $\gamma$  lies on dark (correspondingly light) side of  $\Sigma$  from  $z$ .

Similarly for any unit vector  $u$ , the development  $\tilde{\gamma}_u$  is locally convex (locally concave) if  $\gamma$  lies on dark (correspondingly light) side of  $\Sigma$  for  $u$ .

Note that for any space curve  $\gamma$  and any unit vector  $u$  we have

$$\text{TotCurv } \tilde{\gamma}_u \leq \text{TotCurv } \gamma.$$

On the other hand total curvature of few developments gives an estimate for the total curvature of the original curve. For example if  $i, j, k$  is the orthonormal basis then

$$\text{TotCurv } \gamma \leq \text{TotCurv } \tilde{\gamma}_i + \text{TotCurv } \tilde{\gamma}_j + \text{TotCurv } \tilde{\gamma}_k.$$

If  $\gamma$  lies on one dark or light side for direction  $\mathbf{u}$  then by Liberman's lemma we get

$$\text{TotCurv } \tilde{\gamma}_{\mathbf{u}} \leq \pi.$$

It follows that  $\gamma$  cross the horizons  $\omega_i$ ,  $\omega_j$  and  $\omega_k$  at most  $N$  times then

$$\begin{aligned} \text{TotCurv } \gamma &\leq \text{TotCurv } \tilde{\gamma}_i + \text{TotCurv } \tilde{\gamma}_j + \text{TotCurv } \tilde{\gamma}_k \leq \\ &\leq (N+1) \cdot \pi. \end{aligned}$$

Therefore if  $\gamma$  violates Main Theorem then it has to cross the horizons  $\omega_i$ ,  $\omega_j$  and  $\omega_k$  more that  $10^9$  times.

**3.2. Corollary.** *Let  $\Sigma \subset \mathbb{R}^3$  be a strongly convex smooth surface and  $\gamma: [0, \ell] \rightarrow \Sigma$  be a unit speed geodesic. Assume that for some unit vector  $u$ , the  $\gamma$  cross the horizon  $\omega_u$  transversally at  $t_1 < \dots < t_k$ . Set  $\alpha_i = \angle(\dot{\gamma}(t_i), u)$  for each  $i$ . Then*

$$\text{TotCurv}_u \gamma \leq 3 \cdot \pi + 2 \cdot |\alpha_1 - \alpha_2 + \dots - (-1)^k \cdot \alpha_k|$$

Most of the remaining part of paper devoted to finding an estimate for

$$|\alpha_1 - \alpha_2 + \dots - (-1)^k \cdot \alpha_k|.$$

*Proof.* By Liberman's lemma,

$$\text{TotCurv}_u(\gamma|_{[t_i, t_{i+1}]}) = \pm(\alpha_i - \alpha_{i+1})$$

where we have  $+$  sign if  $[t_i, t_{i+1}]$  lies on the dark side from  $u$  and  $-$  if it lies on the light side. Summing all this up we get

$$\text{TotCurv}_u(\gamma|_{[t_1, t_k]}) = \alpha_1 - 2 \cdot \alpha_2 - \dots + (-1)^k \cdot 2 \cdot \alpha_{k-1} - (-1)^k \cdot \alpha_k.$$

By Liberman's lemma we also have

$$\text{TotCurv}_u(\gamma|_{[0, t_1]}), \text{TotCurv}_u(\gamma|_{[t_k, \ell]}) \leq \pi$$

Since  $|\alpha_i| \leq \frac{\pi}{2}$ , the statement follows.  $\square$

#### 4. KEY LEMMA

Let  $\Sigma$  be a smooth strongly convex surface in  $\mathbb{R}^3$ . Given a unit vector  $\mathbf{u}$ , denote by  $\omega_{\mathbf{u}}$  the  $\mathbf{u}$ -horizon on  $\Sigma$ .

Let  $\gamma: [a, b] \rightarrow \Sigma$  be a geodesic which lies completely on the dark side and both ends  $\gamma(a)$  and  $\gamma(b)$  lie on  $\omega_{\mathbf{u}}$ .

Set  $\alpha = \angle(\dot{\gamma}(a), \mathbf{u}) - \frac{\pi}{2}$  and  $\beta = \angle(\dot{\gamma}(b), \mathbf{u}) - \frac{\pi}{2}$ .

Assume  $\Omega$  is one of the discs which  $\gamma$  cuts from the dark side of  $\Sigma$ .

Then


$$\int_{\Omega} K \cdot d\text{area}$$

takes one of the following four values  $\alpha - \beta$ ,  $2 \cdot \pi - (\alpha - \beta)$ ,  $\pi \pm (\alpha + \beta)$ .

Note that  $\text{TotCurv}_{\mathbf{u}} \gamma = \alpha - \beta$  therefore in all 4 cases we have

$$\text{TotCurv}_{\mathbf{u}} \gamma \leq \int_{\Omega} K \cdot d\text{area}.$$

**Snake case.** This observation makes possible to estimate  $\text{TotCurv}_{\mathbf{u}} \gamma$  if  $\gamma$  cross  $\omega_{\mathbf{u}}$  as shown on the picture. That is, points  $\gamma(t_1), \dots, \gamma(t_k)$  lie in the same order on cross  $\omega_{\mathbf{u}}$ .



In this case we can choose nonoverlapping discs  $\Omega_1, \dots, \Omega_k$  which  $\gamma$  cuts from light and dark side alternatively. Then estimates above show that


$$\begin{aligned} \text{TotCurv}_{\mathbf{u}} \gamma &= \text{TotCurv}_{\mathbf{u}}(\gamma|_{[0, t_0]}) + \text{TotCurv}_{\mathbf{u}}(\gamma|_{[t_0, t_1]}) + \dots \\ &\quad \dots + \text{TotCurv}_{\mathbf{u}}(\gamma|_{[t_{k-1}, t_k]}) + \text{TotCurv}_{\mathbf{u}}(\gamma|_{[t_k, \ell]}) \leq \\ &\leq \pi + K(\Omega_1) + \dots + K(\Omega_k) + \pi \leq \\ &\leq 6 \cdot \pi. \end{aligned}$$

The last inequality holds since the discs  $\Omega_i$  do not overlap and by Gauss–Bonnet we have  $K(\Sigma) \leq 4 \cdot \pi$ .

This gives the first idea that the Key Lemma might be useful. It does not help that easily if the  $\gamma$  cross  $\omega$  in a different order, say as the spiral shown on the picture.

Assume that one of the arcs of  $\omega_{\mathbf{u}}$  from  $\gamma(a)$  and  $\gamma(b)$ , say  $\omega_{\mathbf{u}}^+$ , together with  $\gamma$  bound a disc in  $\Sigma$ . In particular,

$$\text{TotCurv}_{\mathbf{u}} \gamma \leq \int_{\Omega} K \cdot d \text{ area}$$



**4.1. Key Lemma.** *Let  $\gamma: [0, \ell] \rightarrow \Sigma$  be a geodesic on the convex surface in the Euclidean space and  $u \in \mathbb{S}^2$ . Assume that  $0 = t_0 < t_1 < \dots < t_n = \ell$  be the values such that each arcs  $\gamma|_{[t_{i-1}, t_i]}$  alternating light and dark side of  $\Sigma$  with respect to  $u$ . Set  $\alpha_i = \angle(\dot{\gamma}(t_i), u)$ . Then*

$$\text{TotCurv}_{\mathbf{u}} \gamma = \left| \sum_i (-1)^i \alpha_i \right|.$$

*Moreover, if  $1 < i < n$  and  $\Omega_i$  denotes the domain of  $\Sigma$  bounded by the arc  $\gamma|_{[t_{i-1}, t_i]}$  and the  $u$ -horizon then*

$$|\alpha_i - \alpha_{i-1}| \leq \text{curv } \Omega_i,$$

*where  $\text{curv } \Omega_i$  denotes the total curvature of  $\Omega_i$ . In particular,*

$$\text{TotCurv}_{\mathbf{u}} \gamma \leq 4 \cdot \pi + \sum_i \text{curv } \Omega_i.$$

**Remarks.** Clearly  $\text{TotCurv}_z \gamma \leq \text{TotCurv } \gamma$  for any curve  $\gamma$  in  $\Sigma$ .

On the other hand given few points  $z_i$  which do not lie in one plane one can estimate  $\text{TotCurv } \gamma$  in terms of  $\text{TotCurv}_{z_i} \gamma$  the distances between  $z_i$  and the maximal distance to  $\gamma$ .

Let  $N = N(\Sigma, \gamma, u)$  be the maximal integer such that at most  $N$  of the domains  $\Omega_i$  intersect at one point. Note that from [3], it follows that the value  $N$  can take arbitrary large value. The number  $N$  can be estimated through the maximal rotation number of subarcs of  $\gamma$  with respect to the lines. In particular the total curvature of geodesic  $\gamma$  can be bounded in terms of maximal rotation number of subarcs of  $\gamma$  around the lines. The later was claimed in [3] without a proof.

Then

$$\sum_{i=2}^{n-1} \text{curv } \Omega_i \leq N \cdot \text{curv } \Sigma \leq 4 \cdot N \cdot \pi.$$

Therefore, we get an estimate

$$\text{TotCurv}_u \gamma \leq 4 \cdot N \cdot \pi + |\alpha_0 - \alpha_1| + |\alpha_{n-1} - \alpha_n| \leq (4 \cdot N + 2) \cdot \pi.$$

Since the same holds for any vector  $u$ , we can taking average we get

$$\text{TotCurv} \gamma \leq 3 \cdot (4 \cdot N + 2) \cdot \pi.$$

## 5. LENGTH AND DIAMETER

Let  $\varepsilon > 0$ . A curve  $\gamma: [a, b] \rightarrow \mathbb{R}^3$  will be called  $\varepsilon$ -straight if

$$\text{length} \gamma \leq e^\varepsilon \cdot |\gamma(b) - \gamma(a)|$$

**5.1. Lemma.** *Given  $\varepsilon > 0$  there is  $\delta > 0$  (any  $\delta < (1 - e^{-\varepsilon})/2$  will do) such that in any minimizing geodesic of length  $\ell$  on a convex surface  $\Sigma$  in  $\mathbb{R}^3$  there an  $\varepsilon$ -straight arc of length at least  $\delta \cdot \ell$ ;*

*Proof.* Set  $\alpha = \arccos e^{-\varepsilon}$ . Let  $N$  be the maximal number of points in  $\mathbb{S}^2$  which lie on distance at least  $2 \cdot \alpha$  from each other.

Let  $\gamma: [0, \ell] \rightarrow \Sigma$  be a minimizing geodesic parametrized by its length.

Given a value  $t \in [0, \ell]$ , set  $t'$  to be the maximal value in  $[0, \ell]$  such that the interval  $[t, t']$  is  $\varepsilon$ -straight.

Consider the maximal sequence  $0 = t_0 < t_1 < \dots < t_n < \ell$  such that  $t_{i+1} = t'_i$ .

Denote by  $\nu_i$  the outer unit normal vector to  $\Sigma$  at  $\gamma(t_i)$ . Note that  $\angle(\nu_i, \nu_j) > 2 \cdot \alpha$  for all  $i$  and  $j$ . It follows that the sequence  $(t_i)$  terminates after at most  $N$  steps. Therefore any  $\delta < \frac{1}{N+1}$  does the job.  $\square$

**5.2. Lemma.** *Assume  $\gamma$  is a minimizing geodesic on a convex surface in  $\mathbb{R}^3$ . Then*

$$\text{length} \gamma < 4 \cdot \text{diam} \gamma.$$

*Proof.* Assume contrary; that is, there is convex surface  $\Sigma \subset \mathbb{R}^3$  and a geodesic  $\gamma: [0, 4] \rightarrow \Sigma$  is parametrized by its length with  $\text{diam} \gamma \leq 1$ .

Denote by  $\nu_0, \nu_2$  and  $\nu_4$  the outer unit normal vectors to  $\Sigma$  at  $\gamma(0), \gamma(2)$  and  $\gamma(4)$  correspondingly.

Note that  $\angle(\nu_0, \nu_2), \angle(\nu_2, \nu_4) \geq \frac{2}{3} \cdot \pi$  and  $\angle(\nu_0, \nu_4) > \frac{2}{3} \cdot \pi$ , a contradiction.  $\square$

## 6. REDUCTION TO A MONOTONIC CASE

In this section we show that to prove the Main theorem, it is sufficient to consider only the geodesics which go almost in one direction. The following proposition will be applied to  $\varepsilon = \frac{\pi}{4}$ ; in this case one can take  $\delta = 10^{-10}$ .

**6.1. Proposition.** *Given  $\varepsilon > 0$  there is  $\delta > 0$  such that the following statement holds.*

*If  $\gamma: [0, \ell] \rightarrow \Sigma$  is a minimizing geodesic on a smooth strongly convex surface  $\Sigma$  in  $\mathbb{R}^3$  then there is an interval  $[a, b] \subset [0, \ell]$  such that*

$$\text{TotCurv}(\gamma|_{[a,b]}) > \delta \cdot \text{TotCurv} \gamma.$$

*and*

$$\angle(\dot{\gamma}(t), \mathbf{k}) < \varepsilon$$

*for any  $t \in [a, b]$  and a fixed unit vector  $\mathbf{k}$ .*

*Proof.* Applying rescaling, we can assume that  $\text{diam} \gamma = 3$ . By Lemma 5.2 length  $\gamma_n < 12$ . Therefore we can subdivide  $\gamma$  into 12 arcs  $\gamma_1, \dots, \gamma_{12}$  such that for each  $n$  there is a point  $p_n \in K$  which lies on the distance at least 1 from  $\gamma_n$  and

length  $\gamma_n \leq 1$ . Choose an arc  $\gamma' = \gamma_n$  with the maximal total curvature and set  $p' = p_n$ . Clearly

$$\text{TotCurv } \gamma' \geq \frac{1}{12} \cdot \text{TotCurv } \gamma.$$

Applying Liberman's Lemma to  $\gamma'$  with the reference point  $p'$  we get that

$$\text{TotCurv}_{p'} \gamma' < \pi + 1 < 5.$$

Choose an integer  $N > \frac{2}{\varepsilon}$ . Note that we can divide  $\gamma'$  into  $N$  arcs  $\gamma'_1, \dots, \gamma'_N$  so that

$$\text{TotCurv}_{p'} \gamma'_n \leq \frac{5}{N}$$

for each  $n$ . Choose among these arcs the one with maximal total curvature, denote it further by  $\gamma''$ . Clearly

$$\text{TotCurv } \gamma'' > \frac{\varepsilon}{10^3} \cdot \text{TotCurv } \gamma'.$$

Fix a parameter  $t$  of  $\gamma''$  and denote by  $\alpha$  the angle between  $\dot{\gamma}''(t)$  and  $p - \gamma''(t)$ .

If  $\alpha < \frac{\varepsilon}{2}$  or  $\alpha > \pi - \frac{\varepsilon}{2}$ , then the problem is solved.

Otherwise applying Lemma 5.1 we get a nondegenerate (say equilateral) triangle  $\triangle a_1 a_2 a_3$  in  $K_n$  of the size comparable to  $\text{diam } \gamma'$  and on the distance comparable to  $\text{diam } \gamma'$  from any point of  $\gamma''$ , say side of triangle can be taken to be  $\frac{\varepsilon^2}{1000} \cdot \text{diam } \bar{\gamma}$  and the distance to any point can be assumed to be between  $\text{diam } \bar{\gamma}$  and  $2 \cdot \text{diam } \bar{\gamma}$ .

Apply the construction to each vertex of the triangle. We pass to an arc of  $\hat{\gamma}$  such that the angle between  $\dot{\gamma}(t)$  and  $a_i - \gamma(t)$  and the distance  $|\gamma(t) - a_i|$  are nearly constant for each  $i$ . The later imply that  $\dot{\gamma}$  is nearly constant.  $\square$

## 7. ELEVATING GEODESICS

In this section we fix notations which will be used further without additional explanation.

Fix a  $(x, y, z)$ -coordinates on the Euclidean space; denote by  $(\mathbf{i}, \mathbf{j}, \mathbf{k})$  the standard basis.

The lines parallel to the  $z$ -axis will be called *vertical*; the lines and planes parallel to  $(x, y)$ -plane will be called *horizontal*.

**7.1. Definition.** A smooth curve  $\gamma: [0, \ell] \rightarrow \mathbb{R}^3$  is called *elevating* if both ends  $\gamma(0)$  and  $\gamma(\ell)$  lie on the  $z$ -axis and  $\langle \dot{\gamma}(t), \mathbf{k} \rangle > 0$  for all  $t$ .

**$(\lambda, \mu, \nu)$ -frame.** Let  $\Sigma$  be a convex surface and  $\gamma: [0, \ell] \rightarrow \Sigma$  is an elevating minimizing geodesic with unit-speed parametrization.

Given  $t \in [0, \ell]$ , consider the oriented orthonormal frame  $\lambda(t), \mu(t), \nu(t)$  such that  $\nu(t)$  is the outer normal to  $\Sigma$  at  $\gamma(t)$ , the vector  $\mu(t)$  is horizontal and therefore the vector  $\lambda(t)$  lies in the plane spanned by  $\nu(t)$  and the  $z$ -axis. We assume in addition that  $\langle \lambda, \mathbf{k} \rangle \geq 0$ .

Since  $\langle \dot{\gamma}(t), \mathbf{k} \rangle > 0$ ,  $\nu(t)$  can not be vertical and therefore the frame  $(\lambda, \mu, \nu)$  is uniquely defined for any  $t \in [0, \ell]$ .

**Angle functions.** Set

$$\varphi(t) = \angle(\mathbf{k}, \dot{\gamma}(t)), \quad \psi(t) = \frac{\pi}{2} - \angle(\mathbf{k}, \nu(t)), \quad \alpha(t) = \frac{\pi}{2} - \angle(\mu(t), \dot{\gamma}(t)),$$

From the above definitions it follows that  $|\alpha(t)|, |\psi(t)| \leq \frac{\pi}{2}$  and for each  $t$  there is a right spherical triangle with legs  $|\alpha(t)|, |\psi(t)|$  and hypotenuse  $\varphi(t)$ . In particular  $\cos \alpha \cdot \cos \psi = \cos \varphi$ . Whence we get the following.

**7.2. Claim.** For any  $t$  we have

$$\varphi(t) \geq |\psi(t)| \quad \text{and} \quad \varphi(t) \geq |\alpha(t)|$$

Applying Liberman's Lemma in the direction  $\mathbf{k}$  we also get the following.

**7.3. Claim.** *If an arc  $\gamma|_{[a,b]}$  lies in the dark (light) side for  $\mathbf{k}$  then the function  $\varphi$  is nondecreasing (correspondingly nonincreasing) in  $[a, b]$ .*

## 8. PLANE SECTIONS

Assume  $\gamma$  is a curve on a smooth strictly convex surface  $\Sigma$  in  $\mathbb{R}^3$ . Consider a plane  $L$  passing through two points of  $\gamma$ , say  $p = \gamma(a)$  and  $q = \gamma(b)$  with  $a < b$ . Let  $L_{\pm}$  be a half-planes in  $L$  bounded by the line through  $p$  and  $q$ . Set  $\sigma_{\pm} = \Sigma \cap L_{\pm}$ ; note that  $\sigma_{\pm}$  are a smooth convex plane curve connecting  $p$  to  $q$  in  $\Sigma$ .

**8.1. Observation.** *If  $\gamma$  is a minimizing geodesic in the convex surface  $\Sigma \subset \mathbb{R}^3$  and  $a, b$  and  $\sigma_{\pm}$  as above then*

$$\text{length } \sigma_{\pm} \geq \text{length}(\gamma|_{[a,b]}).$$

Based on this observation we give couple of estimates on elevating minimizing geodesics.

**8.2. Propostion.** *Assume  $\gamma: [0, \ell] \rightarrow \Sigma$  is an elevating minimizing geodesic in the convex surface  $\Sigma \subset \mathbb{R}^3$ . Assume that for a subsegment  $[a, b] \subset [0, \ell]$  the following conditions hold*

- (i) *The points  $\gamma(a)$  and  $\gamma(b)$  lie in a half-plane with boundary line formed by the  $z$ -axis and the arc  $\gamma|_{[a,b]}$  goes around the  $z$ -axis at least once.*
  - (ii)  *$\gamma(a)$  lies above the horizontal plane through  $\frac{1}{2} \cdot (\gamma(0) + \gamma(\ell))$ .*
- Then  $\gamma(b)$  lies on the dark side of  $\Sigma$  with respect to  $\mathbf{k}$ .*

*Proof.* Let us apply Observation 8.1 to the plane containing  $z$ -axis,  $\gamma(a)$  and  $\gamma(b)$ .

We can assume that  $\gamma(0)$  is the origin of the  $(x, y, z)$ -coordinate system and both points  $p = \gamma(a)$  and  $q = \gamma(b)$  lie in the  $(x, z)$ -coordinate half-plane with  $x \geq 0$ , denoted by  $\Pi$ . We can assume that  $\sigma_{+} \subset \Pi$ . Let  $(x_p, 0, z_p)$  and  $(x_q, 0, z_q)$  be the coordinates of  $p$  and  $q$ .

From the assumptions we get  $z_p < z_q < 2 \cdot z_p$ .

By convexity of the curve  $\Pi \cap \Sigma$  we get

$$\text{length } \sigma_{+} \leq \sqrt{(z_q - z_p)^2 + x_p^2}$$

On the other hand, since  $\gamma|_{[a,b]}$  goes around  $z$ -axis at least once, we get

$$\text{length } \gamma|_{[a,b]} \geq \sqrt{(z_q - z_p)^2 + (x_p + x_q)^2}.$$

These two estimates contradict Observation 8.1. □

**8.3. Corollary.** *If  $\Sigma, \gamma, \ell, a$  and  $b$  as in the Proposition and the arc  $\gamma|_{[a,b]}$  goes around the  $z$ -axis at least twice then the arc  $\gamma|_{[b,\ell]}$  lies on the dark side with respect to  $\mathbf{k}$ .*

*Proof.* Fix  $b' \in [b, \ell]$ . Note that one can find  $a' \in [a, b]$  such that the the assumptions of Proposition 8.2 hold for the interval  $[a', b']$ . Applying Proposition we get the result. □

**8.4. Propostion.** *Assume  $\gamma: [0, \ell] \rightarrow \Sigma$  is elevating minimizing geodesic in the convex surface  $\Sigma \subset \mathbb{R}^3$ . Assume that the arc  $\gamma|_{[b,\ell]}$  lies in the dark side of  $\Sigma$  with respect to  $\mathbf{k}$ . Set  $\varphi(t) = \angle(\mathbf{k}, \dot{\gamma}(t))$  and  $\psi(t) = \frac{\pi}{2} - \angle(\mathbf{k}, \nu(t))$ . If  $b \leq s < t \leq \ell$  and the point  $\gamma(s)$  lies in the plane  $\Pi$  through  $\gamma(t)$  spanned by  $\nu(t)$  and  $\lambda(t)$  then*

$$\varphi(s) \leq \psi(t).$$

*Proof.* Let us apply Observation 8.1 to the plane  $\Pi$  and  $p = \gamma(s)$  and  $q = \gamma(t)$ .



Let  $z_p$  and  $z_q$  be the  $z$ -coordinates of  $p$  and  $q$ .

Since  $\gamma|_{[s,t]}$  lies in the dark side, its Liberman's development  $\tilde{\gamma}|_{[s,t]}$  with respect to  $\mathbf{k}$  is concave. In particular

$$\text{length}(\gamma|_{[s,t]}) = \text{length}(\tilde{\gamma}|_{[s,t]}) \geq \frac{z_q - z_p}{\cos \varphi(s)}.$$

On the other hand, convexity of  $\sigma_+$  imply that

$$\text{length } \sigma_+ \leq \frac{z_q - z_p}{\cos \psi(t)}.$$

It remains to apply Observation 8.1.  $\square$

## 9. $\mathbf{s}$ -PAIRS

Let  $\Sigma \subset \mathbb{R}^3$  be a strongly convex surface and  $\gamma: [0, \ell] \rightarrow \Sigma$  be an elevating minimizing geodesic.

After rotating  $(x, y)$ -plane if necessary, we can assume that the border of shadow in the directions of  $x$ -axis, say  $\omega_x$ , is a smooth curve and  $\gamma$  intersects them transversely.

Let  $t_1 < t_2 < \dots < t_k$  be the time moments in  $[0, \ell]$  at which  $\gamma$  crossing  $\omega_x$ . Note that

$$\mu(t_n) = s_n \cdot e_x \quad \text{for some } s_n = \pm 1.$$

Set

$$\varphi_n = \varphi(t_n) \quad \psi_n = \psi(t_n) \quad \alpha_n = \alpha(t_n)$$

We say that a pair of indexes  $i < j$  forms an  $\mathbf{s}$ -pair if

$$\sum_{n=i}^j s_n = 0 \quad \text{and} \quad \sum_{n=i}^{j'} s_n > 0$$

if  $i < j' < j$ .

Note that for any index  $i$  appears in at most one  $\mathbf{s}$ -pair and for any  $\mathbf{s}$ -pair  $(i, j)$  we have

- ◇  $s_i = 1$ ; that is,  $i$ -th bracket has to be opening.
- ◇  $s_j = -1$ ; that is,  $j$ -th bracket has to be closing.

In particular,

$$s_i \cdot \alpha_i + s_j \cdot \alpha_j = \alpha_i - \alpha_j.$$

**Bracket interpretation.** If you exchange “+1” and “−1” in  $\mathbf{s}$  by “(” and “)” correspondingly then  $(i, j)$  is an  $\mathbf{s}$ -pair if and only if the  $i$ -th bracket forms a pair with  $j$ -bracket.

**Embedded disc interpretation.** Assume  $(i, j)$  is an  $\mathbf{s}$ -pair. Note that in this case there is an arc of  $\omega_i$  from  $\gamma(t_i)$  to  $\gamma(t_j)$  with monotonic  $z$ -coordinate. Moreover this arc, say  $\sigma$  together with  $\gamma|_{[t_i, t_j]}$  bounds an immersed disc in  $\Sigma$ . That is there is an immersion  $\iota: \mathbb{D} \rightarrow \Sigma$  such that the closed curve  $\iota|_{\partial \mathbb{D}}$  is formed by joint of  $\sigma$  and  $\gamma|_{[t_i, t_j]}$ .

The proof can be guessed from the diagram. It shows a lift of  $\gamma$  in the universal cover of strip of  $\Sigma$  between horizontal planes through  $\gamma(t_i)$  and  $\gamma(t_j)$ ; the solid vertical lines correspond are lifts of  $\sigma$  and the dashed lines corresponds to the lifts of the other component of  $\omega_i$  between the planes.

We say that  $q$  is the depth of an  $\mathbf{s}$ -pair  $(i, j)$  (briefly  $q = \text{depth}_{\mathbf{s}}(i, j)$ ) if  $q$  is the maximal number such that there is  $q$ -long nested sequence of  $\mathbf{s}$ -pairs starting with  $(i, j)$ ; that is a sequence of  $\mathbf{s}$ -pairs  $(i, j) = (i_1, j_1), (i_2, j_2), \dots, (i_q, j_q)$  such that

$$i = i_1 < \dots < i_q < j_q < \dots < j_1 = j.$$

We say that  $q$  is the depth of an  $\mathbf{s}$ -pair  $(i, j)$  (briefly  $q = \text{depth}_{\mathbf{s}}(i, j)$ ) if  $q$  is the maximal number such that there is a  $q$ -long nested sequence of  $\mathbf{s}$ -pairs starting with  $(i, j)$ ; that is a sequence of  $\mathbf{s}$ -pairs  $(i, j) = (i_1, j_1), (i_2, j_2), \dots, (i_q, j_q)$  such that

$$i = i_1 < \dots < i_q < j_q < \dots < j_1 = j.$$

Note that the  $\mathbf{s}$ -pair of the same depth do not overlap; that is if for two distinct  $\mathbf{s}$ -pairs  $(i, j)$  and  $(i', j')$ , we have  $\text{depth}(i, j) = \text{depth}(i', j')$  then either  $i < j < i' < j'$  or  $i' < j' < i < j$ .

The following proposition follow directly from the definitions above.

**9.1. Proposition.** *Let  $(i, j)$  be an  $\mathbf{s}$ -pair. Then the arcs  $\gamma|_{[t_i, t_j]}$  and an arc of  $\omega_i$  bound an immesed disc in  $\Sigma$  which lies between horizontal planes through  $\gamma(t_i)$  and  $\gamma(t_j)$ . Moreover the maximal multiplicity of the disc is at most  $\text{depth}_{\mathbf{s}}(i, j)$ .*

**9.2. Corollary.** *Let us denote by  $S_q$  the subset of indexes  $\{1, \dots, k\}$  which are the parts of  $\mathbf{s}$ -pairs with depth  $q$ . Then*

$$\sum_{n \in S_q} s_n \cdot \alpha_n \leq 4 \cdot \pi \cdot q.$$

*Proof.* For each  $n$  denote by  $K_n$  the integral of Gauss curvature of the part of surface  $\Sigma$  which lies below horizontal plane through  $\gamma(t_n)$ . Note that

$$0 \leq K_1 \leq \dots \leq K_k \leq 4 \cdot \pi.$$

By Proposition 9.1 and the Key Lemma, we get

$$s_i \cdot \alpha_i + s_j \cdot \alpha_j = \alpha_i - \alpha_j \leq q \cdot (K_j - K_i)$$

The statement follows since the  $\mathbf{s}$ -pairs with the same depth do not overlap.  $\square$

**9.3. Corollary.** *Assume*

$$q = \max_{1 \leq i < j \leq k} \left\{ \left| \sum_{n=i}^j s_n \right| \right\}$$

*Then*

$$\left| \sum_{n=1}^k s_n \cdot \alpha_n \right| \leq 2 \cdot q \cdot \left(q + \frac{3}{2}\right) \cdot \pi.$$

*Proof.* Denote by  $S$  the set of all indexes which appear in some  $\mathbf{s}$ -pair.

Note that depth of any  $\mathbf{s}$  pair is at most  $q$ . That is,

$$S = S_1 \cup \dots \cup S_q.$$

By Corollary 9.2,

$$\sum_{n \in S} s_n \cdot \alpha_n \leq 2 \cdot q \cdot (q + 1) \cdot \pi. \quad \blacksquare$$

Set  $R = \{1, \dots, k\} \setminus S$ ; this is the set of indexes which do not appear in an  $\mathbf{s}$ -pair.

Given  $r$ , set  $i \in Q_r$  if

$$\sum_{n=1}^i s_n = r.$$

Note that  $Q_r \neq \emptyset$  for at most  $q$  values of  $r$  and in each set  $Q_r$  there are at most 2 indexes which do not appear in an  $\mathbf{s}$ -pair; that is  $Q_r \cap R$  has at most two indexes for each  $r$ .

Sine  $|a_n| < \frac{\pi}{2}$ , we get

$$\left| \sum_{n \in R} s_n \cdot \alpha_n \right| \leq q \cdot \pi.$$

The later inequality together with ❶ implies the statement in the corollary.  $\square$

## 10. GEOMETRIC GROWTH

**10.1. Claim.** *Assume  $\psi(t) > \varepsilon$  for ant  $t \in [t_i, t_j]$  and  $s_i + \dots + s_j = 2$  Then*

$$|\alpha_j - \alpha_i| > \pi \cdot \sin \varepsilon.$$

**10.2. Claim.** *Let  $\gamma$  be elevating minimizing geodesic on a graph  $z = f(x, y)$  of a concave function. Then for any pair of indexes  $j > i$ , such that*

$$\left| \sum_{n=i}^j s_n \right| \geq 5$$

*we have*

$$\varphi_j > \frac{3}{2} \cdot \varphi_i.$$

*Proof.* Without loss of generality, we may assume that

$$\sum_{n=i}^j s_n = 6$$

Let  $j'$  be the least index such that

$$\left| \sum_{n=i}^{j'} s_n \right| = 5.$$

Note that for any  $b > t_j$  there is  $a \in [t_i, t_j]$  such that intrval  $[a, b]$  satisfies the assumptions of Proposition 8.4. In particular  $\psi(b) > \varphi_i$  for any  $b > t_j$ . Applying Claim 10.1, we get that  $|\alpha_j| > \frac{\pi}{2} \cdot \varphi_i$  or  $|\alpha_{j'}| > \frac{\pi}{2} \cdot \varphi_i$ . Since  $\varphi_n$  is nondecreasing, and  $\varphi_n \geq |\alpha_n|$  for any  $n$ , in both cases we get

$$\varphi_j > \frac{\pi}{2} \cdot \varphi_i.$$

$\square$

## 11. AN ESTIMATE FOR GRAPHS

**11.1. Proposition.** *There is c constant  $\omega'$  ( $\omega' = 10 \cdot \pi$  will do) such that if  $\gamma$  is an elevating minimizing geodesic on a graph  $z = f(x, y)$  of a concave function  $f$  then*

$$\text{TotCur}_i \gamma \leq \omega'.$$

*Proof.* We can assume that  $\gamma$  cross the  $i$  horizon  $\omega_i$  tranfersally. Let  $t_1 < \dots < t_k$  be the values of parameter at which  $\gamma$  cross  $\omega_i$  and  $s_1, \dots, s_k$  the signs as in ...

Recall that  $S_q$  denotes the subset of indexes  $\{1, \dots, k\}$  which appear in  $s$ -pair with depph  $q$ . According to Corollary 9.2,

$$\left| \sum_{n \in S_q} s_n \cdot \alpha_n \right| \leq 4 \cdot q \cdot \pi.$$

In particular,

$$\left| \sum_{n \in S_1 \cup \dots \cup S_5} s_n \cdot \alpha_n \right| \leq 40 \cdot \pi.$$

Set  $R = \{1, \dots, k\} \setminus (S_1 \cup \dots \cup S_5)$ ; this is the set of indexes which appear in  $s$ -pairs with depth at least 6 as well as those which do not appear in any  $s$ -pair.

According to ???

$$\left| \sum_{n \in R} s_n \cdot \alpha_n \right| \leq \sum_{n \in R} \varphi_n.$$

To estimate the last sum will use the results in Section 10. First let us subdivide  $R$  into 5 subsets  $R_1, \dots, R_5$ , by setting  $n \in R_m$  if  $m \equiv n \pmod{5}$ .

Given  $n \in R_m$ , denote by  $n'$  the least index in  $R_m$  which is larger  $n$ ;  $n'$  is defined for any  $n \in R_m$  except the largest one. According to ???  $\varphi_{n'} > 2 \cdot \varphi_n$ . Since  $\varphi_n$  is nondecreasing in  $n$ , we get

$$\sum_{n \in R_m} \varphi_n \leq 2 \cdot \varphi_k.$$

It follows that

$$\sum_{n \in R} \varphi_n \leq 10 \cdot \varphi_k < 5 \cdot \pi.$$

According to Liberman's lemma

$$\begin{aligned} \text{TotCurv}_i \gamma &\leq 4 \cdot \pi + 2 \cdot [s_1 \cdot \alpha_1 + \dots + s_k \cdot \alpha_k] \leq \\ &\leq 100 \cdot \pi. \end{aligned}$$

□

## 12. FINAL ASSEMBLING

Assume there is a minimizing geodesic  $\gamma: [0, \ell] \rightarrow \Sigma$  in a convex surface  $\Sigma \subset \mathbb{R}^3$  such that

$$\text{TotCurv} \gamma = \omega.$$

According to ??? we can assume that  $\Sigma$  is strongly convex.

According to ???, we can pass to an elevating arc, of  $\gamma$  for some  $(x, y, z)$ -coordinate system with total curvature  $> \frac{\omega}{10^6}$ . Rename this arc by  $\gamma$  and let us use the notations in Section 7.

Rotating  $(x, y)$ -coordinate plane we can ensure that

$$\text{TotCurv} \gamma \leq 10 \cdot \text{TotCurv}_i \gamma$$

and that  $\gamma$  cross the horizon  $\omega_i$  transversally.

Let us subdivide  $\gamma$  into three arcs lower middle and upper arcs  $\gamma_-$ ,  $\gamma_0$  and  $\gamma_+$  the the following way.

Note that according to ???  $\gamma_+$  lies on a graph of concave function. By Proposition ??, we get

$$\text{TotCurv}_i \gamma_+ \leq 100 \cdot \pi. \quad \textcircled{2}$$

Similarly  $\gamma_-$  lies on a graph of convex function and the same proposition implies

$$\text{TotCurv}_i \gamma_- \leq 100 \cdot \pi. \quad \textcircled{3}$$

By Corollary 9.3,

$$\text{TotCurv}_i \gamma_0 \leq 100 \cdot \pi. \quad \textcircled{4}$$

$$\text{TotCurv} \gamma_0 \leq ???$$

Together with  $\textcircled{2}$  and  $\textcircled{3}$ , the later implies that

$$\text{TotCurv}_i \gamma \leq 300 \cdot \pi.$$

□

## REFERENCES

- [1] P. K. Agarwal, S. Har-Peled, M. Sharir, and K. R. Varadarajan. “Approximating shortest paths on a convex polytope in three dimensions”. *J. ACM* 44.4 (1997), pp. 567–584.
- [2] J. Pach. “Folding and turning along geodesics in a convex surface”. *Geombinatorics* 7.2 (1997), pp. 61–65.
- [3] I. Bárány, K. Kuperberg, and T. Zamfirescu. “Total curvature and spiralling shortest paths”. *Discrete Comput. Geom.* 30.2 (2003). U.S.-Hungarian Workshops on Discrete Geometry and Convexity (Budapest, 1999/Auburn, AL, 2000), pp. 167–176.
- [4] J. Liberman. “Geodesic lines on convex surfaces”. *C. R. (Doklady) Acad. Sci. URSS (N.S.)* 32 (1941), pp. 310–313.
- [5] V. V. Usov. “The length of the spherical image of a geodesic on a convex surface”. *Sibirsk. Mat. Ž.* 17.1 (1976), pp. 233–236.
- [6] I. D. Berg. “An estimate on the total curvature of a geodesic in Euclidean 3-space-with-boundary”. *Geom. Dedicata* 13.1 (1982), pp. 1–6.
- [7] A. V. Pogorelov. *Extrinsic geometry of convex surfaces*. Translated from the Russian by Israel Program for Scientific Translations, Translations of Mathematical Monographs, Vol. 35. American Mathematical Society, Providence, R.I., 1973, pp. vi+669.
- [8] A. D. Milka. “A shortest path with nonrectifiable spherical representation. I”. *Ukrain. Geometr. Sb.* 16 (1974), pp. 35–52, ii.
- [9] V. V. Usov. “The three-dimensional swerve of curves on convex surfaces”. *Sibirsk. Mat. Ž.* 17.6 (1976), pp. 1427–1430, 1440.

N. LEBEDEVA  
 MATH. DEPT. ST. PETERSBURG STATE UNIVERSITY, UNIVERSITETSKY PR., 28, STARY PETERHOF, 198504, RUSSIA.  
 STEKLOV INSTITUTE, 27 FONTANKA, ST. PETERSBURG, 191023, RUSSIA.  
*E-mail address:* lebed@pdmi.ras.ru

A. PETRUNIN  
 MATH. DEPT. PSU, UNIVERSITY PARK, PA 16802, USA  
*E-mail address:* petrunin@math.psu.edu