

SOME VC-DENSITY COMPUTATIONS IN SHELAH-SPENCER GRAPHS

ANTON BOBKOV

ABSTRACT. We compute vc-densities of minimal extension formulas in Shelah-Spencer random graphs.

We fix the density of the graph α .


Lemma 0.1. *For any $\mathcal{A} \in \mathbf{K}_\alpha$ and $\epsilon > 0$ there exists an \mathcal{B} such that $(\mathcal{A}, \mathcal{B})$ is minimal and $\delta(\mathcal{B}/\mathcal{A}) < \epsilon$.*

Proof. Let m be an integer such that $m\alpha < 1 < (m+1)\alpha$. Suppose \mathcal{A} has less than $m+1$ vertices. Make a construction $\mathcal{A}_0 = \mathcal{A}$ and \mathcal{A}_{i+1} is \mathcal{A}_i with one extra vertex connected to every single vertex of \mathcal{A}_i . Stop when the total number of vertices is $m+1$. Proceed as in [1] 4.1. Resulting construction is still minimal. \square

Lemma 0.2. *Let $\mathcal{A}_1 \subset \mathcal{B}_1$ and $\mathcal{A}_2 \subset \mathcal{B}_2$ be \mathbf{K}_α structures with $(\mathcal{A}_2, \mathcal{B}_2)$ a minimal pair with $\epsilon = \delta(\mathcal{B}_2/\mathcal{A}_2)$. Let M be some ambient structure. Fix embeddings of $\mathcal{A}_1, \mathcal{B}_1, \mathcal{A}_2$ into M . Assume that it is not the case that $\mathcal{A}_2 \subset \mathcal{B}_2$ and \mathcal{A}_1 is disjoint from \mathcal{A}_2 (No!). Now consider all possible embeddings $f: \mathcal{B}_2 \rightarrow M$ over \mathcal{A}_1 . Let $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ and $\mathcal{B}_f = \mathcal{B}_1 \cup f(\mathcal{B}_2)$ with $\delta_f = \delta(\mathcal{B}_f/\mathcal{A})$. Then δ_f is at most $\delta(\mathcal{B}_1 \cup \mathcal{A}/\mathcal{A}) + \epsilon$*

Fix an embedding f . It induces the following substructure diagram in M . Denote

$$\begin{aligned}\mathcal{A} &= \mathcal{A}_1 \cup \mathcal{A}_2 \\ \mathcal{B}_f^* &= \mathcal{B}_1 \cup f(\mathcal{B}_2) \\ \mathcal{B}_1^* &= \mathcal{B}_1 \cup \mathcal{A} \\ \mathcal{B}_2^* &= f(\mathcal{B}_2) \cup \mathcal{A} \\ \mathcal{B}^* &= \mathcal{B}_1^* \cap \mathcal{B}_2^*\end{aligned}$$



From the diagram we see that

$$\delta(\mathcal{B}_f/\mathcal{A}) \leq \delta(\mathcal{B}_1^*/\mathcal{A}) + \delta(\mathcal{B}_2^*/\mathcal{B}^*)$$

Thus all we need to do is to verify that

$$\delta(\mathcal{B}_2^*/\mathcal{B}^*) \leq \epsilon$$

Let \mathcal{B}' denote graph induced on all the vertices in $(f(B_2)/B_1) \cup A_2$. Then \mathcal{B}' is a substructure of \mathcal{B}_2 over \mathcal{A}_2 . By minimality we get that $\delta(\mathcal{B}'/\mathcal{A}_2) \leq \epsilon$. We need to show $\delta(\mathcal{B}_2^*/\mathcal{B}^*) \leq \delta(\mathcal{B}'/\mathcal{A}_2)$. Do the vertex computation

$$\begin{aligned}
 B_2^* - B^* &= \\
 f(B_2) - (B_1 \cap f(B_2)) - A &= \\
 f(B_2) - B_1 - A &= \\
 f(B_2) - B_1 - A_2
 \end{aligned}$$

and

$$B' - A_2 = f(B_2) - B_1 - A_2$$

So the sets of the extra vertices in the extension are the same. The base $\mathcal{B}_2^*/\mathcal{B}^*$ is larger so we can introduce some extra edges but no new vertices. This means that $\delta(\mathcal{B}_2^*/\mathcal{B}^*) \leq \delta(\mathcal{B}'/\mathcal{A}_2)$ giving us the original statement.

Let $\phi(x, y)$ be a formula in a random graph with $|x| = |y| = 1$ saying that there exists \mathcal{D} over $\mathcal{C} = \{x, y\}$ such that $(\mathcal{D}, \mathcal{C})$ is minimal with relative dimension ϵ . Let n be such that $n\epsilon < 1 < (n+1)\epsilon$. Then we argue that $vc(\phi) = n$.

Fix a m -strong (for any $m > |M|$) set of non-connected vertices A . Fix some a . We investigate the trace of $\phi(x, a)$ on B . Suppose we have b_1, \dots, b_k satisfying $\phi(b_i, a)$ as witnessed by M_j . Relative dimension of $M_1 \cup M_2 \cup \dots \cup M_j \cup a$ is minimized when all M_j are disjoint (by minimality). Thus for that dimension to be positive we can have at most n extensions.

Consider sets $\mathcal{B}_1 \dots \mathcal{B}_n$ with

- (1) $a_i \in \mathcal{B}_i$
- (2) $a_i \in A$

- (3) $a_i \neq a_j$
- (4) $a^* \in \bigcap \mathcal{B}_i$

and s.t. $\mathcal{B}_i/\{a^*, a_i\}$ is isomorphic to \mathcal{B}/A . We look at all the possible embeddings with those properties. We argue that a disjoint configuration minimizes total dimension of the whole construction.

We argue by induction on n . Fix an embedding $\mathcal{B}_1, \dots, \mathcal{B}_n$ and consider possible choices for $\mathcal{B}_{n+1}, a_{n+1}$. We can pick a_n to be an element of A not used so far and embed \mathcal{B}_{n+1} over $\{a^*, a_i\}$ disjoint from the entire construction. On the other hand suppose it is embedded such that there is an intersection. We set up to apply the previous lemma. Let

$$\begin{aligned}\mathcal{B}_1 &= \bigcup_{1..n} \mathcal{B}_i \\ \mathcal{A}_1 &= \{a_1, \dots, a_n\} \\ \mathcal{B}_2 &= \mathcal{B}_{n+1} \\ \mathcal{A}_2 &= \{a^*, a_{n+1}\}\end{aligned}$$

Applying the lemma say that the extra dimension cannot be larger then ϵ .

REFERENCES

- [1] Michael C. Laskowski, *A simpler axiomatization of the Shelah-Spencer almost sure theories*, Israel J. Math. **161** (2007), 157-186. MR MR2350161
E-mail address: bobkov@math.ucla.edu