

a
K fits for EG and PEG 35k at 25 wt.%
PEG 35k 25.0 wt.% fit: K = (7.2e + 09 ± 1.8e + 09) M⁻¹
EG 25.0 wt.% fit: K = (1.1e + 08 ± 6.2e + 06) M⁻¹
PEG 35k 25.0 wt.% data
EG 25.0 wt.% data
donor count rate [cps]
C_DNA (acceptor) [nM]
b
K vs crowder concentration
K [M⁻¹]
crowder concentration [wt.%]
buffer
EG
PEG 200
PEG 400
PEG 600
PEG 1k
PEG 1.5k
PEG 3k
PEG 6k
PEG 12k
PEG 20k
PEG 35k
c
ΔΔG vs crowder concentration
ΔΔG [kJ/mol]
crowder concentration [wt.%]
d
Molecular crowding fits for EG and PEG 35k
K [M⁻¹]
EG fit
lnK = 20.9 + 0.1x [M⁻¹] + -0.2x² [M⁻²]
EG uncertainty band
PEG 35k fit
lnK = 21.7 + 521.0x [M⁻¹] + -45190.7x² [M⁻²]
PEG 35k uncertainty band
crowder concentration [wt.%]
e
a1 and a2 coefficients vs polymer length
×10⁴
a1 [M⁻¹]
a2 [M⁻²]
a1
a2
degree of polymerization
f
Position of ΔΔG_minimum vs R_g
concentration at ΔΔG_minimum [c/c*]
R_g [nm]