

Simplex Method Using the Dictionaries Formulation Approach

Antonio Flores-Tlacuahuac

ITESM, México

August 19, 2019

Solve by Simplex Method:

$$\begin{array}{ll}\max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\ \text{s.t.} & 2x_1 + 3x_2 + x_3 \leq 5 \\ & 4x_1 + x_2 + 2x_3 \leq 11 \\ & 3x_1 + 4x_2 + 2x_3 \leq 8 \\ & x_1, x_2, x_3 \geq 0\end{array}$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\max_{x_1, x_2, x_3} \Omega = 5x_1 + 4x_2 + 3x_3$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}\max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\ \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5\end{array}$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}\max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\ \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\ & 4x_1 + x_2 + 2x_3 + x_5 = 11\end{array}$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}
 \max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\
 \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\
 & 4x_1 + x_2 + 2x_3 + x_5 = 11 \\
 & 3x_1 + 4x_2 + 2x_3 + x_6 = 8
 \end{array}$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}
 \max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\
 \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\
 & 4x_1 + x_2 + 2x_3 + x_5 = 11 \\
 & 3x_1 + 4x_2 + 2x_3 + x_6 = 8 \\
 & x_1, x_2, x_3, x_4, x_5, x_6 \geq 0
 \end{array}$$

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}
 \max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\
 \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\
 & 4x_1 + x_2 + 2x_3 + x_5 = 11 \\
 & 3x_1 + 4x_2 + 2x_3 + x_6 = 8 \\
 & x_1, x_2, x_3, x_4, x_5, x_6 \geq 0
 \end{array}$$

- ▶ where x_4, x_5 and x_6 are the so-called **slack variables**

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}
 \max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\
 \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\
 & 4x_1 + x_2 + 2x_3 + x_5 = 11 \\
 & 3x_1 + 4x_2 + 2x_3 + x_6 = 8 \\
 & x_1, x_2, x_3, x_4, x_5, x_6 \geq 0
 \end{array}$$

- ▶ where x_4, x_5 and x_6 are the so-called **slack variables**
- ▶ Note that the slacks variables also become decision variables and are bounded by a non-negativity bound

- ▶ Transform the set of inequalities constraints into a set of Equality constraints introducing **slack variables**
- ▶ Use a slack variable for each inequality constraint

$$\begin{array}{ll}
 \max_{x_1, x_2, x_3} & \Omega = 5x_1 + 4x_2 + 3x_3 \\
 \text{s.t.} & 2x_1 + 3x_2 + x_3 + x_4 = 5 \\
 & 4x_1 + x_2 + 2x_3 + x_5 = 11 \\
 & 3x_1 + 4x_2 + 2x_3 + x_6 = 8 \\
 & x_1, x_2, x_3, x_4, x_5, x_6 \geq 0
 \end{array}$$

- ▶ where x_4, x_5 and x_6 are the so-called **slack variables**
- ▶ Note that the slacks variables also become decision variables and are bounded by a non-negativity bound

- Write the optimization problem using the *Dictionary* Approach

- ▶ Write the optimization problem using the *Dictionary* Approach
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

- ▶ Write the optimization problem using the *Dictionary* Approach
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

- ▶ Write the optimization problem using the *Dictionary* Approach
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

$$\begin{array}{rcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 \end{array}$$

- ▶ Write the optimization problem using the *Dictionary* Approach
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

$$\begin{array}{rcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 \\ x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 \end{array}$$

- ▶ Write the optimization problem using the *Dictionary* Approach
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

x_4	$=$	$5 - 2x_1 - 3x_2 - x_3$
x_5	$=$	$11 - 4x_1 - x_2 - 2x_3$
x_6	$=$	$8 - 3x_1 - 4x_2 - 2x_3$
Ω	$=$	$5x_1 + 4x_2 + 3x_3$

- ▶ Write the optimization problem using the *Dictionary Approach*
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

x_4	$=$	$5 - 2x_1 - 3x_2 - x_3$
x_5	$=$	$11 - 4x_1 - x_2 - 2x_3$
x_6	$=$	$8 - 3x_1 - 4x_2 - 2x_3$
Ω	$=$	$5x_1 + 4x_2 + 3x_3$

- ▶ Variables in the LHS (x_4, x_5, x_6) are called **Basic variables**

- ▶ Write the optimization problem using the *Dictionary Approach*
- ▶ Move the slacks variables to LHS and keep the original decision variables in the RHS

x_4	$=$	$5 - 2x_1 - 3x_2 - x_3$
x_5	$=$	$11 - 4x_1 - x_2 - 2x_3$
x_6	$=$	$8 - 3x_1 - 4x_2 - 2x_3$
Ω	$=$	$5x_1 + 4x_2 + 3x_3$

- ▶ Variables in the LHS (x_4, x_5, x_6) are called **Basic variables**
- ▶ Variables in the RHS (x_1, x_2, x_3) are called **Non-Basic variables**

- Write the optimization problem using the *Dictionary* Approach
- Move the slacks variables to LHS and keep the original decision variables in the RHS

x_4	$=$	$5 - 2x_1 - 3x_2 - x_3$
x_5	$=$	$11 - 4x_1 - x_2 - 2x_3$
x_6	$=$	$8 - 3x_1 - 4x_2 - 2x_3$
Ω	$=$	$5x_1 + 4x_2 + 3x_3$

- Variables in the LHS (x_4, x_5, x_6) are called **Basic variables**
- Variables in the RHS (x_1, x_2, x_3) are called **Non-Basic variables**

x_4

x_5

x_6

Basic Variables

x_1

x_2

x_3

Non-Basic Variables

- Write the optimization problem using the *Dictionary* Approach
- Move the slacks variables to LHS and keep the original decision variables in the RHS

x_4	$=$	$5 - 2x_1 - 3x_2 - x_3$
x_5	$=$	$11 - 4x_1 - x_2 - 2x_3$
x_6	$=$	$8 - 3x_1 - 4x_2 - 2x_3$
Ω	$=$	$5x_1 + 4x_2 + 3x_3$

- Variables in the LHS (x_4, x_5, x_6) are called **Basic variables**
- Variables in the RHS (x_1, x_2, x_3) are called **Non-Basic variables**

x_4

x_5

x_6

Basic Variables

x_1

x_2

x_3

Non-Basic Variables

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- Fix to zero the values of the remaining variables: $x_2=0, x_3=0$

First Iteration

- ▶ Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- ▶ Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- ▶ Amount by which x_1 can be increased keeping feasibility?

First Iteration

- ▶ Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- ▶ Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- ▶ Amount by which x_1 can be increased keeping feasibility?

$$2x_1 + 3x_2 + x_3 \leq 5 \quad \begin{array}{l} \text{Basic Variable} \\ \rightarrow x_4 \end{array}$$

First Iteration

- ▶ Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- ▶ Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- ▶ Amount by which x_1 can be increased keeping feasibility?

Basic Variable

$$2x_1 + 3x_2 + x_3 \leq 5 \quad \rightarrow x_4$$

$$4x_1 + x_2 + 2x_3 \leq 11 \quad \rightarrow x_5$$

First Iteration

- ▶ Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- ▶ Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- ▶ Amount by which x_1 can be increased keeping feasibility?

Basic Variable

$2x_1 + 3x_2 + x_3$	\leq	5	$\rightarrow x_4$
$4x_1 + x_2 + 2x_3$	\leq	11	$\rightarrow x_5$
$3x_1 + 4x_2 + 2x_3$	\leq	8	$\rightarrow x_6$

First Iteration

- ▶ Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- ▶ Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- ▶ Amount by which x_1 can be increased keeping feasibility?

Basic Variable

$$2x_1 + 3x_2 + x_3 \leq 5 \quad \rightarrow x_4$$

$$4x_1 + x_2 + 2x_3 \leq 11 \quad \rightarrow x_5$$

$$3x_1 + 4x_2 + 2x_3 \leq 8 \quad \rightarrow x_6$$

after setting $x_2 = 0$ and $x_3 = 0$:

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- Amount by which x_1 can be increased keeping feasibility?

Basic Variable

$$2x_1 + 3x_2 + x_3 \leq 5 \quad \rightarrow x_4$$

$$4x_1 + x_2 + 2x_3 \leq 11 \quad \rightarrow x_5$$

$$3x_1 + 4x_2 + 2x_3 \leq 8 \quad \rightarrow x_6$$

after setting $x_2 = 0$ and $x_3 = 0$:

$$2x_1 \leq 5 \rightarrow x_1 = 2.5$$

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- Amount by which x_1 can be increased keeping feasibility?

		Basic Variable
$2x_1 + 3x_2 + x_3$	≤ 5	$\rightarrow x_4$
$4x_1 + x_2 + 2x_3$	≤ 11	$\rightarrow x_5$
$3x_1 + 4x_2 + 2x_3$	≤ 8	$\rightarrow x_6$

after setting $x_2 = 0$ and $x_3 = 0$:

$$\begin{array}{rclcl} 2x_1 & \leq & 5 & \rightarrow x_1 & = & 2.5 \\ 4x_1 & \leq & 11 & \rightarrow x_1 & = & 2.75 \end{array}$$

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- Amount by which x_1 can be increased keeping feasibility?

		Basic Variable
$2x_1 + 3x_2 + x_3$	≤ 5	$\rightarrow x_4$
$4x_1 + x_2 + 2x_3$	≤ 11	$\rightarrow x_5$
$3x_1 + 4x_2 + 2x_3$	≤ 8	$\rightarrow x_6$

after setting $x_2 = 0$ and $x_3 = 0$:

$2x_1$	≤ 5	$\rightarrow x_1$	$= 2.5$
$4x_1$	≤ 11	$\rightarrow x_1$	$= 2.75$
$3x_1$	≤ 8	$\rightarrow x_1$	$= 2.66$

First Iteration

- Want to increase the value of the objective function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

select the largest coefficient variable in Ω : x_1

- Fix to zero the values of the remaining variables: $x_2=0, x_3=0$
- Amount by which x_1 can be increased keeping feasibility?

		Basic Variable
$2x_1 + 3x_2 + x_3$	≤ 5	$\rightarrow x_4$
$4x_1 + x_2 + 2x_3$	≤ 11	$\rightarrow x_5$
$3x_1 + 4x_2 + 2x_3$	≤ 8	$\rightarrow x_6$

after setting $x_2 = 0$ and $x_3 = 0$:

$2x_1$	≤ 5	$\rightarrow x_1$	$= 2.5$
$4x_1$	≤ 11	$\rightarrow x_1$	$= 2.75$
$3x_1$	≤ 8	$\rightarrow x_1$	$= 2.66$

hence,

► for $x_1 = 2.5$

hence,

► for $x_1 = 2.5$

$$2(2.5) = 5 \leq 5$$

hence,

► for $x_1 = 2.5$

$$\begin{array}{rcl} 2(2.5)=5 & \leq & 5 \\ 4(2.5)=10 & \leq & 11 \end{array}$$

hence,

► for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

hence,

► for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

► for $x_1 = 2.75$

hence,

► for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

► for $x_1 = 2.75$

$$2(2.75)=5.5 \not\leq 5$$

hence,

► for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

► for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11

hence,

► for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

► for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

hence,

- ▶ for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

- ▶ for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

- ▶ for $x_1 = 2.66$

hence,

- ▶ for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

- ▶ for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

- ▶ for $x_1 = 2.66$

$2(2.66)=5.32$	$\not\leq$	5
----------------	------------	---

hence,

- ▶ for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

- ▶ for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

- ▶ for $x_1 = 2.66$

$2(2.66)=5.32$	$\not\leq$	5
$4(2.66)=10.64$	\leq	11

hence,

- ▶ for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

- ▶ for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

- ▶ for $x_1 = 2.66$

$2(2.66)=5.32$	$\not\leq$	5
$4(2.66)=10.64$	\leq	11
$3(2.66)=7.98$	\leq	8

hence,

- ▶ for $x_1 = 2.5$

$2(2.5)=5$	\leq	5
$4(2.5)=10$	\leq	11
$3(2.5)=7.5$	\leq	8

- ▶ for $x_1 = 2.75$

$2(2.75)=5.5$	$\not\leq$	5
$4(2.75)=11$	\leq	11
$3(2.75)=8.25$	$\not\leq$	8

- ▶ for $x_1 = 2.66$

$2(2.66)=5.32$	$\not\leq$	5
$4(2.66)=10.64$	\leq	11
$3(2.66)=7.98$	\leq	8

- ▶ Then we pick $x_1=2.5$ (to meet constraints)

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3 = 5 - 2(2.5) - 3(0) - (0) = 0$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \end{array}$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\ x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5 \end{array}$$

- ▶ In summary, from the first iteration:

$$\begin{array}{l} x_1 = 2.5 \\ x_2 = 0 \end{array}$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

$$\Omega(x) = 12.5$$

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

$$\Omega(x) = 12.5$$

Things relevant to recall:

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\ x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

$$\Omega(x) = 12.5$$

Things relevant to recall:

- ▶ x_1 was the increasing variable

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclclcl}
 x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\
 x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\
 x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5
 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

$$\Omega(x) = 12.5$$

Things relevant to recall:

- ▶ x_1 was the increasing variable
- ▶ x_4 was the basic variable related to the equation from which maximum increase in x_1 , without constraints violation, was achieved

- ▶ Then we pick $x_1=2.5$ (to meet constraints)
- ▶ and because $x_2=0$ and $x_3=0$ from:

$$\begin{array}{rclclclcl} x_4 & = & 5 - 2x_1 - 3x_2 - x_3 & = & 5 - 2(2.5) - 3(0) - (0) & = & 0 \\ x_5 & = & 11 - 4x_1 - x_2 - 2x_3 & = & 11 - 4(2.5) - (0) - 2(0) & = & 1 \\ x_6 & = & 8 - 3x_1 - 4x_2 - 2x_3 & = & 8 - 3(2.5) - 4(0) - 2(0) & = & 0.5 \end{array}$$

- ▶ In summary, from the first iteration:

$$x_1 = 2.5$$

$$x_2 = 0$$

$$x_3 = 0$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0.5$$

and,

$$\Omega(x) = 12.5$$

Things relevant to recall:

- ▶ x_1 was the increasing variable
- ▶ x_4 was the basic variable related to the equation from which maximum increase in x_1 , without constraints violation, was achieved

Second Iteration

x_4

x_5

x_6

Basic Variables


x_1

x_2


x_3

Non-Basic Variables


Second Iteration






Basic Variables



Non-Basic Variables




Basic Variables




Non-Basic Variables


Second Iteration





Basic Variables




Non-Basic Variables




Basic Variables



Non-Basic Variables




Basic Variables




Non-Basic Variables


Second Iteration






Basic Variables




Non-Basic Variables




Basic Variables



Non-Basic Variables



Basic Variables



Non-Basic Variables

- ▶ How to turn a non-basic into a basic variable?

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$x_5 = 11 - 4x_1 - x_2 - 2x_3$$

$$x_6 = 8 - 3x_1 - 4x_2 - 2x_3$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$x_5 = 11 - 4x_1 - x_2 - 2x_3$$

$$x_6 = 8 - 3x_1 - 4x_2 - 2x_3$$

For x_5 :

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$x_5 = 11 - 4x_1 - x_2 - 2x_3$$

$$x_6 = 8 - 3x_1 - 4x_2 - 2x_3$$

For x_5 :

$$x_5 = 11 - 4x_1 - x_2 - 2x_3$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \end{aligned}$$

For x_5 :

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ &= 11 - 4 \left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \right) - x_2 - 2x_3 \end{aligned}$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \end{aligned}$$

For x_5 :

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ &= 11 - 4\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - x_2 - 2x_3 \\ &= 1 + 5x_2 + 2x_4 \end{aligned}$$

- ▶ How to turn a non-basic into a basic variable?
- ▶ Pick from the **Dictionary formulation** that equation in which x_4 is the basic variable:

$$x_4 = 5 - 2x_1 - 3x_2 - x_3$$

and solve for x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

and replace this equation into the two remaining dictionary equations:

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \end{aligned}$$

For x_5 :

$$\begin{aligned} x_5 &= 11 - 4x_1 - x_2 - 2x_3 \\ &= 11 - 4\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - x_2 - 2x_3 \\ &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_6 :

For x_6 :

$$x_6 = 8 - 3x_1 - 4x_2 - 2x_3$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3\end{aligned}$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\Omega = 5x_1 + 4x_2 + 3x_3$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3\end{aligned}$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

$$\boxed{x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4}$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

$$\left| \begin{array}{lcl} x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 & = & 1 + 5x_2 + 2x_4 \end{array} \right|$$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

x_1	$=$	$\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$
x_5	$=$	$1 + 5x_2 + 2x_4$
x_6	$=$	$\frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

x_1	$=$	$\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$
x_5	$=$	$1 + 5x_2 + 2x_4$
x_6	$=$	$\frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$
Ω	$=$	$\frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4$

For x_6 :

$$\begin{aligned}x_6 &= 8 - 3x_1 - 4x_2 - 2x_3 \\&= 8 - 3\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) - 4x_2 - 2x_3 \\&= \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= 5x_1 + 4x_2 + 3x_3 \\&= 5\left(\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4\right) + 4x_2 + 3x_3 \\&= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4\end{aligned}$$

In summary, the new dictionary reads:

x_1	$=$	$\frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$
x_5	$=$	$1 + 5x_2 + 2x_4$
x_6	$=$	$\frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$
Ω	$=$	$\frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4$

- Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 = \frac{5}{2} - \frac{1}{2}x_3 \rightarrow x_3=5$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 = \frac{5}{2} - \frac{1}{2}x_3 \rightarrow x_3 = 5^1$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{rclclcl} x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\ x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=5 & ^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow & x_5=1 & \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=1 &
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=5 & ^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow & x_5=1 & \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=1 & ^2
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=5 & ^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow & x_5=1 & \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow & x_3=1 & ^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{rclclcl}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 = \frac{5}{2} - \frac{3}{2}(0) - \frac{1}{2}(1) - \frac{1}{2}(0) = 2$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{rclclcl}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

$$\begin{array}{rclclcl}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{3}{2}(0) - \frac{1}{2}(1) - \frac{1}{2}(0) & = & 2 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 + 5(0) + 2(0) & = & 1
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{rclclcl}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

$$\begin{array}{rclclcl}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{3}{2}(0) - \frac{1}{2}(1) - \frac{1}{2}(0) & = & 2 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 + 5(0) + 2(0) & = & 1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} + \frac{1}{2}(0) - \frac{1}{2}(1) + \frac{3}{2}(0) & = & 0
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

$$\begin{array}{llllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{3}{2}(0) - \frac{1}{2}(1) - \frac{1}{2}(0) & = & 2 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 + 5(0) + 2(0) & = & 1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} + \frac{1}{2}(0) - \frac{1}{2}(1) + \frac{3}{2}(0) & = & 0 \\
 \Omega & = & \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 & = & \frac{25}{2} - \frac{7}{2}(0) + \frac{1}{2}(1) - \frac{5}{2}(0) & = & 13
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- ▶ Now, pick the non-basic variable featuring the largest positive coefficient in Ω : x_3
- ▶ Check the maximum amount by which x_3 can be increased without constraints violation and set to zero the remaining non-basic variables ($x_2 = 0, x_4 = 0$)
- ▶ From the present dictionary:

$$\begin{array}{llllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{1}{2}x_3 & \rightarrow x_3=5^1 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 & \rightarrow x_5=1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} - \frac{1}{2}x_3 & \rightarrow x_3=1^2
 \end{array}$$

- ▶ Hence, setting $x_3 = 1$:

$$\begin{array}{llllll}
 x_1 & = & \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 & = & \frac{5}{2} - \frac{3}{2}(0) - \frac{1}{2}(1) - \frac{1}{2}(0) & = & 2 \\
 x_5 & = & 1 + 5x_2 + 2x_4 & = & 1 + 5(0) + 2(0) & = & 1 \\
 x_6 & = & \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4 & = & \frac{1}{2} + \frac{1}{2}(0) - \frac{1}{2}(1) + \frac{3}{2}(0) & = & 0 \\
 \Omega & = & \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 & = & \frac{25}{2} - \frac{7}{2}(0) + \frac{1}{2}(1) - \frac{5}{2}(0) & = & 13
 \end{array}$$

¹when solving this equation set the basic variable x_1 to zero

²when solving the basic variable this equation set x_6 to zero

- In summary, from the second iteration:

- In summary, from the second iteration:

$$x_1 = 2$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

$$\Omega(x) = 13$$

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

$$\Omega(x) = 13$$

Things relevant to recall:

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

$$\Omega(x) = 13$$

Things relevant to recall:

- x_3 was the increasing variable

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

$$\Omega(x) = 13$$

Things relevant to recall:

- x_3 was the increasing variable
- x_6 was the basic variable related to the equation from which maximum increase in x_3 , without constraints violation, was achieved

- In summary, from the second iteration:

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

and,

$$\Omega(x) = 13$$

Things relevant to recall:

- x_3 was the increasing variable
- x_6 was the basic variable related to the equation from which maximum increase in x_3 , without constraints violation, was achieved

Third Iteration

Third Iteration


 x_1 x_5 x_6

Basic Variables


 x_4 x_2 x_3

Non-Basic Variables


Third Iteration






Basic Variables



Non-Basic Variables




Basic Variables




Non-Basic Variables


Third Iteration





Basic Variables




Non-Basic Variables




Basic Variables



Non-Basic Variables




Basic Variables




Non-Basic Variables


Third Iteration





Basic Variables




Non-Basic Variables




Basic Variables



Non-Basic Variables



Basic Variables



Non-Basic Variables

- Once again, turn a non-basic into a basic variable (x_3)

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_1 :

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_1 :

$$x_1 = \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_1 :

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{1}{2}x_4 \end{aligned}$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_1 :

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{1}{2}x_4 \\ &= 2 - 2x_2 - 2x_4 + x_6 \end{aligned}$$

- ▶ Once again, turn a non-basic into a basic variable (x_3)
- ▶ Similarly to the previous iteration, pick from the past **Dictionary formulation** that equation in which x_6 is the basic variable:

$$x_6 = \frac{1}{2} + \frac{1}{2}x_2 - \frac{1}{2}x_3 + \frac{3}{2}x_4$$

and solve for x_3 :

$$x_3 = 1 + x_2 + 3x_4 - 2x_6$$

and replace in the remaining dictionary equations:

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ x_5 &= 1 + 5x_2 + 2x_4 \end{aligned}$$

For x_1 :

$$\begin{aligned} x_1 &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}x_3 - \frac{1}{2}x_4 \\ &= \frac{5}{2} - \frac{3}{2}x_2 - \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{1}{2}x_4 \\ &= 2 - 2x_2 - 2x_4 + x_6 \end{aligned}$$

Also replace into the Objective Function:

Also replace into the Objective Function:

$$\Omega = \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

The new dictionary reads:

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

The new dictionary reads:

x_1	=	$2 - 2x_2 - 2x_4 + x_6$
x_3	=	$1 + x_2 + 3x_4 - 2x_6$
x_5	=	$1 + 5x_2 + 2x_4$
Ω	=	$13 - 3x_2 - x_4 - x_6$

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

The new dictionary reads:

x_1	=	$2 - 2x_2 - 2x_4 + x_6$
x_3	=	$1 + x_2 + 3x_4 - 2x_6$
x_5	=	$1 + 5x_2 + 2x_4$
Ω	=	$13 - 3x_2 - x_4 - x_6$

since all coefficients in the past objective function are negative, no further improvement of the Objective Function can be achieved.

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

The new dictionary reads:

x_1	$=$	$2 - 2x_2 - 2x_4 + x_6$
x_3	$=$	$1 + x_2 + 3x_4 - 2x_6$
x_5	$=$	$1 + 5x_2 + 2x_4$
Ω	$=$	$13 - 3x_2 - x_4 - x_6$

since all coefficients in the past objective function are negative, no further improvement of the Objective Function can be achieved.

This means that the optimal solution has been found.

Also replace into the Objective Function:

$$\begin{aligned}\Omega &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}x_3 - \frac{5}{2}x_4 \\ &= \frac{25}{2} - \frac{7}{2}x_2 + \frac{1}{2}(1 + x_2 + 3x_4 - 2x_6) - \frac{5}{2}x_4 \\ &= 13 - 3x_2 - x_4 - x_6\end{aligned}$$

The new dictionary reads:

x_1	$=$	$2 - 2x_2 - 2x_4 + x_6$
x_3	$=$	$1 + x_2 + 3x_4 - 2x_6$
x_5	$=$	$1 + 5x_2 + 2x_4$
Ω	$=$	$13 - 3x_2 - x_4 - x_6$

since all coefficients in the past objective function are negative, no further improvement of the Objective Function can be achieved.

This means that the optimal solution has been found.

$$x_1 = 2$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

$$x_5 = 1$$

$$x_6 = 0$$

$$\Omega(x) = 13$$