RESEARCH

A sample article title

Jane E. Doe* and John R.S. Smith

*Correspondence:
jane.e.doe@cambridge.co.uk
ETSI Informática, Universidad de Málaga, Málaga, España
Full list of author information is available at the end of the article

Abstract

Keywords: sample; article; author

Content

Text and results for this section, as per the individual journal's instructions for authors. Here, we reference the figure 1 and figure 2 but also the table 1.

Section title

Text for this section...

In this section we examine the growth rate of the mean of Z_0, Z_1 and Z_2. In addition, we examine a common modeling assumption and note the importance of considering the tails of the extinction time T_x in studies of escape dynamics. We will first consider the expected resistant population at vT_x for some $v > 0$, (and temporarily assume $\alpha = 0$)

$$E\big[Z_1(vT_x)\big] = \int_0^{v\wedge 1} Z_0(uT_x)\exp(\lambda_1)\,du.$$

If we assume that sensitive cells follow a deterministic decay $Z_0(t) = xe^{\lambda_0 t}$ and approximate their extinction time as $T_x \approx -\frac{1}{\lambda_0}\log x$, then we can heuristically estimate the expected value as

$$\begin{aligned} &E\big[Z_1(vT_x)\big] \\ &\quad = \frac{\mu}{r}\log x \int_0^{v\wedge 1} x^{1-u}x^{(\lambda_1/r)(v-u)}\,du. \end{aligned} \tag{1}$$

Thus we observe that this expected value is finite for all $v > 0$ (also see [**?**, **?**, **?**, **?**, **?**, **?**]).

Table 1 Sample table title. This is where the description of the table should go

	B1	B2	B3
A1	0.1	0.2	0.3
A2	...	..	.
A3	..	.	.

Sub-heading for section

Text for this sub-heading...




Figure 1 Sample figure title




Figure 2 Sample figure title

Sub-sub heading for section

Text for this sub-sub-heading...

Sub-sub-sub heading for section Text for this sub-sub-sub-heading...

1 Introducción

2 Materiales y métodos

3 Resultados

El HP con nombre *Tall stature* y código HP:0000098 tiene 182 genes que al mapearlos en la red de STRING [1] con un umbral de corte de 900, quedan 166 genes y 300 conexiones. La red resultante tiene una transitivity de 0,5016.

El HP con nombre *Aortic dissection* y código HP:0002647 tiene 41 genes que al mapearlos en la red de STRING [1] con un umbral de corte de 700, quedan 39 genes y 79 conexiones. La red resultante tiene una transitivity de 0,5758.

4 Discusión

5 Conclusiones

Abreviaciones
Indicar lista de abreviaciones mostrando cada acrónimo a que corresponde

Disponibilidad de datos y materiales
Debéis indicar aquí un enlace a vuestro repositorio de github.

Contribución de los autores
Usando las iniciales que habéis definido al comienzo del documento, debeis indicar la contribución al proyecto en el estilo: J.E : Encargado del análisis de coexpresión con R, escritura de resultados; J.R.S : modelado de red con python y automatizado del código, escritura de métodos; ... OJO: que sea realista con los registros que hay en vuestros repositorios de github.

Author details
ETSI Informática, Universidad de Málaga, Málaga, España.

References

1. Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A.L., Fang, T., Doncheva, N.T., Pyysalo, S., Bork, P., Jensen, L.J., Mering, C.V.: The string database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Research **51** (2023). doi:10.1093/nar/gkac1000