

# Pima Indians | Exploratory Analysis

Antonio Pano Flores

9/11/2022

## Variables Used:

Pregnancy: Number of times pregnant.

Glucose: Oral Glucose Tolerance Test.

Blood Pressure: Diastolic blood pressure (mm Hg)

Skin Thickness: Triceps skin fold thickness (mm)

Insulin: 2-Hour serum insulin (mu U/ml)

BMI: Body mass index (weight in kg/(height in m)<sup>2</sup>)

DiabetesPedigreeFunction: Diabetes pedigree function (function which scores likelihood of diabetes based on family history).

Age: Years of Age.

Outcome: 0 = Does Not Have Diabetes, 1 = Does Have Diabetes

## Preparing the Data for Use:

1. Adjusting Outcome Integer into a Factor.
2. Replacing 0's with NA values within the `Insulin`, `BMI`, and `SkinThickness` fields.
3. Creating a 'GlucoseGroup' variable based on labels set by the American Diabetes Association (<https://diabetes.org/diabetes/a1c/diagnosis>)
4. Creating a new variable 'WeightGroup' based on BMI value. Using bins set by the American Cancer Society (<https://www.cancer.org/healthy/cancer-causes/diet-physical-activity/body-weight-and-cancer-risk/adult-bmi.html>).

```
glimpse(PI)
```

```
## Rows: 768
## Columns: 9
## $ Pregnancies      <int> 6, 1, 8, 1, 0, 5, 3, 10, 2, 8, 4, 10, 10, 1, ~
## $ Glucose          <int> 148, 85, 183, 89, 137, 116, 78, 115, 197, 125~
## $ BloodPressure    <int> 72, 66, 64, 66, 40, 74, 50, 0, 70, 96, 92, 74~
## $ SkinThickness    <int> 35, 29, 0, 23, 35, 0, 32, 0, 45, 0, 0, 0, 0, ~
## $ Insulin          <int> 0, 0, 0, 94, 168, 0, 88, 0, 543, 0, 0, 0, 0, ~
## $ BMI              <dbl> 33.6, 26.6, 23.3, 28.1, 43.1, 25.6, 31.0, 35.~
## $ DiabetesPedigreeFunction <dbl> 0.627, 0.351, 0.672, 0.167, 2.288, 0.201, 0.2~
## $ Age              <int> 50, 31, 32, 21, 33, 30, 26, 29, 53, 54, 30, 3~
## $ Outcome          <int> 1, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 1, ~
```

```

PI$Outcome = as.factor(PI$Outcome)
PI$Glucose = na_if(PI$Glucose, 0)
PI$Insulin = na_if(PI$Insulin, 0)
PI$BMI = na_if(PI$BMI, 0)
PI$SkinThickness = na_if(PI$SkinThickness, 0)
PI$BloodPressure = na_if(PI$BloodPressure, 0)
#PI$GlucosePerInsulin = PI$Glucose / PI$Insulin

PI <- PI %>%
  mutate(WeightGroup = cut(PI$BMI, breaks = c(0, 18.4, 24.9, 29.9, Inf),
    labels = c("Underweight", "NormalWeight", "Overweight", "Obese")),
    GlucoseGroup = cut(PI$Glucose, breaks = c(0, 99.9, 124.9, Inf),
    labels = c("Normal", "Prediabetes", "Diabetes"))) %>%
  select(Pregnancies, Glucose, GlucoseGroup, BMI, WeightGroup, everything())

glimpse(PI)

```

```

## Rows: 768
## Columns: 11
## $ Pregnancies      <int> 6, 1, 8, 1, 0, 5, 3, 10, 2, 8, 4, 10, 10, 1, ~
## $ Glucose          <int> 148, 85, 183, 89, 137, 116, 78, 115, 197, 125~
## $ GlucoseGroup     <fct> Diabetes, Normal, Diabetes, Normal, Diabetes,~
## $ BMI              <dbl> 33.6, 26.6, 23.3, 28.1, 43.1, 25.6, 31.0, 35.~
## $ WeightGroup      <fct> Obese, Overweight, NormalWeight, Overweight, ~
## $ BloodPressure    <int> 72, 66, 64, 66, 40, 74, 50, NA, 70, 96, 92, 7~
## $ SkinThickness    <int> 35, 29, NA, 23, 35, NA, 32, NA, 45, NA, NA, N~
## $ Insulin          <int> NA, NA, NA, 94, 168, NA, 88, NA, 543, NA, NA,~
## $ DiabetesPedigreeFunction <dbl> 0.627, 0.351, 0.672, 0.167, 2.288, 0.201, 0.2~
## $ Age              <int> 50, 31, 32, 21, 33, 30, 26, 29, 53, 54, 30, 3~
## $ Outcome          <fct> 1, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 0, 1, 1, ~

```

Critique 1: This dataset only contains the diastolic blood pressure levels. In order to get a better picture of artery health, systolic blood pressure levels should have been included.

Critique 2: No data was recorded on the births themselves and whether there were miscarriages or failed births.

Critique 3: No data on whether the group was measured after a fast so I don't feel confident enough to use a calculation for Insulin Resistance.

All 0 values were changed to NA values as they aren't close to typical measurements in BMI, BloodPressure, SkinThickness, or Insulin levels in any population. I assume they are missing values, instead

## Exploratory Data Analysis

- First, looking at statistic summaries with `skim()`— keeping the data *with* NA values. The reason for `skim()` over boxplots is because after glancing at the data, I can only assume that standard deviations may be small for some variables. This prompts me to look the spread of the data numerically, rather than visually, in order to be precise.

Noticing that the oldest recorded woman is 81 years old— an outlier and 11 years older than the second oldest woman in these samples.

If we look at the `skim()`'s BMI histogram, it seems right-skewed which can be expected if we have more samples in which the person was negative for diabetes (assuming younger people naturally have diabetes less often). Using `skim()`'s second output, I learn that there are 500 rows for those that did not have diabetes which definitely is more than 268 for those that did. Still,

since BMI is a strong predictor of diabetes, I plot the distribution individually– assigning color red for the subset that have diabetes/ blue for those without diabetes for presentation purposes.

After doing so, I find that BMI statistics are actually normally distributed despite the right-skewed histogram found in `skimr`. I find that the mean is ~33 for both those with and without diabetes– indicating that the general population *is* very prone to the health condition. This can be confirmed if you perform research on the Arizona Pima Indians & their extraordinarily high rate of kidney disease and failure as a leading cause of death.

Insulin, on the other hand, *is* right-skewed. This means the samples tend to have lower Insulin values more often than higher Insulin values.

```
skimr::skim(PI)
```









#### Data summary

Name	PI
Number of rows	768
Number of columns	11
Column type frequency:	
factor	3
numeric	8
Group variables	
	None

#### Variable type: factor


skim_variable	n_missing	complete_rate	ordered	n_unique	top_counts
GlucoseGroup	5	0.99	FALSE	3	Dia: 311, Pre: 260, Nor: 192
WeightGroup	11	0.99	FALSE	4	Obe: 472, Ove: 179, Nor: 102, Und: 4
Outcome	0	1.00	FALSE	2	0: 500, 1: 268

#### Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
Pregnancies	0	1.00	3.85	3.37	0.00	1.00	3.00	6.00	17.00	
Glucose	5	0.99	121.69	30.54	44.00	99.00	117.00	141.00	199.00	
BMI	11	0.99	32.46	6.92	18.20	27.50	32.30	36.60	67.10	
BloodPressure	35	0.95	72.41	12.38	24.00	64.00	72.00	80.00	122.00	
SkinThickness	227	0.70	29.15	10.48	7.00	22.00	29.00	36.00	99.00	
Insulin	374	0.51	155.55	118.78	14.00	76.25	125.00	190.00	846.00	
DiabetesPedigreeFunction	0	1.00	0.47	0.33	0.08	0.24	0.37	0.63	2.42	
Age	0	1.00	33.24	11.76	21.00	24.00	29.00	41.00	81.00	


```
PI_BMI <- PI[!is.na(PI$BMI),] %>% select(BMI, Outcome)

ggplot(PI_BMI, aes(x=BMI, fill=Outcome)) + theme_minimal() +
  geom_density(color=NA) +
  facet_wrap(vars(Outcome), scale="free_x") +
  geom_vline(aes(xintercept = median(BMI)), size = .8, color = "black") +
  geom_vline(aes(xintercept = mean(BMI)), size=.5, linetype="dashed") +
  scale_fill_manual(values=c("#4479E4", "red3"),
                    name = "Diabetes", labels = c("Absent", "Present"))
```



```
PI_Insulin <- PI[!is.na(PI$Insulin),] %>% select(Insulin, Outcome)

ggplot(PI_Insulin, aes(x=Insulin, fill=Outcome)) + theme_minimal() +
  geom_density(color=NA) + facet_wrap(vars(Outcome), scale="free_x") +
  geom_vline(aes(xintercept = median(Insulin)), size = .8, color = "black") +
  geom_vline(aes(xintercept = mean(Insulin)), size=.5, linetype="dashed") +
  scale_fill_manual(values=c("#4479E4", "red3"),
                    name = "Diabetes", labels = c("Absent", "Present"))
```




Creating two vectors in which I eliminate NA values for each field, independently. This way, I avoid eliminating data from the neighboring columns and portray an accurate distribution.

- Keeping only necessary fields from the vectors in order to save space in the Global Environment.


- Second, looking at distributions for the samples that had Diabetes.

- Then, I look at distributions for the samples that did *not* have Diabetes.
- After creating distributions using `facet_wrap()` and adjusting for a free x-scale, I find that there are either normal or right-skewed distributions only:
  - Right-Skewed: This is true for variables Age, Insulin, Pregnancies, and Pedigree Function.

```
PI %>% filter(Outcome == 0) %>%
  purrr::keep(is.numeric) %>%
  gather() %>%
  ggplot(aes(value)) +
    facet_wrap(~ key, scales = "free") +
    geom_density()
```



```
PI %>% filter(Outcome == 1) %>%
  purrr::keep(is.numeric) %>%
  gather() %>%
  ggplot(aes(value)) +
    facet_wrap(~ key, scales = "free") +
    geom_density()
```



## Exploring same statistics for all samples that had every attribute present.

The number of women with all attributes recorded are only at 392 rows rather than 768. This is around half of the dataframe: ~ 51%.

- Noticing that every field's average dropped slightly.
  - BMI still large.
  - Average Glucose levels are normal.
  - Average Blood Pressure is normal but I still see a high max at p100. This is expected, however, as the number of those diagnosed with diabetes now make up ~33% of the total samples.
  - Average Insulin levels for those with all attributes recorded are higher than what is considered normal by 16 mg/dL.
- For those with all attributes recorded, using the quartile ranges, I can determine that at least 75% of the women in the samples will:
  - Have had at least one pregnancy.
  - Be 23 years of age or older.
  - Have glucose levels of 99 or higher. (The average is higher than what's considered normal by 16 miligrams / decilitre).
  - Have a BMI of 28.4 or higher.
  - Have blood pressure of 62 or more (Nothing out of the ordinary).

```
skimr::skim(PI %>% drop_na())
```






#### Data summary

Name	PI %>% drop_na()
Number of rows	392
Number of columns	11
Column type frequency:	
factor	3
numeric	8
Group variables	
None	

#### Variable type: factor

skim_variable	n_missing	complete_rate	ordered	n_unique	top_counts
GlucoseGroup	0	1	FALSE	3	Dia: 168, Pre: 122, Nor: 102
WeightGroup	0	1	FALSE	4	Obe: 262, Ove: 85, Nor: 44, Und: 1
Outcome	0	1	FALSE	2	0: 262, 1: 130

#### Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
Pregnancies	0	1	3.30	3.21	0.00	1.00	2.00	5.00	17.00	
Glucose	0	1	122.63	30.86	56.00	99.00	119.00	143.00	198.00	
BMI	0	1	33.09	7.03	18.20	28.40	33.20	37.10	67.10	
BloodPressure	0	1	70.66	12.50	24.00	62.00	70.00	78.00	110.00	
SkinThickness	0	1	29.15	10.52	7.00	21.00	29.00	37.00	63.00	

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
Insulin	0	1	156.06	118.84	14.00	76.75	125.50	190.00	846.00	
DiabetesPedigreeFunction	0	1	0.52	0.35	0.09	0.27	0.45	0.69	2.42	
Age	0	1	30.86	10.20	21.00	23.00	27.00	36.00	81.00	

## Comparing the mean values for women who have diabetes against those who did not.

- Both subsets of the data contain an obesity-level average BMI. Those who weren't diagnosed with diabetes fell, on average, into Class 1 Obesity (BMI of 30 to < 35) while those who were, fell into Class 2 Obesity (BMI of 35 to < 40).
- Women with diabetes tend to have had more pregnancies. As previously stated, no data is recorded on the births themselves and whether there were miscarriages or failed births.

```
num_0 <- PI %>% filter(Outcome==0) %>% drop_na() %>% select(Pregnancies, Glucose, BMI, BloodPressure, SkinThick
kness, Insulin, DiabetesPedigreeFunction, Age) %>%
  colMeans() %>% as.data.frame() %>% tibble::rownames_to_column() %>%
  rename("0" = ".")

num_1 <- PI %>% filter(Outcome==1) %>% drop_na() %>% select(Pregnancies, Glucose, BMI, BloodPressure, SkinThick
kness, Insulin, DiabetesPedigreeFunction, Age) %>%
  colMeans() %>% as.data.frame() %>% tibble::rownames_to_column() %>%
  rename("1" = ".")

stat_means <- inner_join(num_0,num_1, by="rowname") %>% rename("Variable" = "rowname")
stat_means
```

```
##           Variable           0           1
## 1      Pregnancies    2.7213740    4.4692308
## 2           Glucose   111.4312977   145.1923077
## 3              BMI    31.7507634    35.7776923
## 4    BloodPressure    68.9694656    74.0769231
## 5    SkinThickness    27.2519084    32.9615385
## 6           Insulin   130.8549618   206.8461538
## 7 DiabetesPedigreeFunction    0.4721679    0.6255846
## 8              Age    28.3473282    35.9384615
```

## Checking correlations between numeric variables in order to avoid multi-collinearity for future predictions.

Noticing that the variables that have the highest correlation are: 1) Pregnancy and Age 2) Skin Thickness and BMI 3) Insulin and Glucose

