

Fundamentals of Computer Graphics

Spectral geometry

Emanuele Rodolà
rodola@di.uniroma1.it



SAPIENZA
UNIVERSITÀ DI ROMA

Course quality questionnaire

The questionnaire is completely **anonymous** – your privacy is respected

Instructions:

- Disable pop-up block in your browser
- Go to <https://www.uniroma1.it>
- Click on **Studenti** and access **Infostud 2.0**
- Click on **Corsi di laurea**
- On the left menu, click on **Opinioni studenti**
- Enter the OPIS code and click on **Vai al questionario**

OPIS codes are:

- **LNM5IDE3**
- **CUP342HA** for the curriculum “Networks and security”

Self-adjointness of Δ

$$\langle F, \nabla g \rangle_{T\mathcal{X}} = -\langle \operatorname{div} F, g \rangle_{\mathcal{X}}$$

Self-adjointness of Δ

$$\langle \nabla f, \nabla g \rangle_{T\mathcal{X}} = -\langle \operatorname{div} \nabla f, g \rangle_{\mathcal{X}}$$

Self-adjointness of Δ

$$\langle \nabla f, \nabla g \rangle_{T\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

Self-adjointness of Δ

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \nabla f, \nabla g \rangle_{T\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

Self-adjointness of Δ

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\lambda \langle \phi, \phi \rangle_{\mathcal{X}} = \langle \lambda \phi, \phi \rangle_{\mathcal{X}}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}}\end{aligned}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \phi, \Delta \phi \rangle_{\mathcal{X}}\end{aligned}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \phi, \Delta \phi \rangle_{\mathcal{X}} \\ &= \overline{\langle \Delta \phi, \phi \rangle}_{\mathcal{X}}\end{aligned}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \phi, \Delta \phi \rangle_{\mathcal{X}} \\ &= \overline{\langle \Delta \phi, \phi \rangle_{\mathcal{X}}} \\ &= \overline{\langle \lambda \phi, \phi \rangle_{\mathcal{X}}}\end{aligned}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \phi, \Delta \phi \rangle_{\mathcal{X}} \\ &= \overline{\langle \Delta \phi, \phi \rangle_{\mathcal{X}}} \\ &= \overline{\langle \lambda \phi, \phi \rangle_{\mathcal{X}}} \\ &= \overline{\lambda} \langle \phi, \phi \rangle_{\mathcal{X}}\end{aligned}$$

Laplacian eigenvalues

$$\langle f, \Delta g \rangle_{\mathcal{X}} = \langle \Delta f, g \rangle_{\mathcal{X}}$$

This means that the Laplace-Beltrami operator Δ is **self-adjoint**

Now consider the (in general, **complex**) eigenvalue equation:

$$\Delta \phi = \lambda \phi$$

For the eigenfunction/eigenvalue pair (ϕ, λ) , we have:

$$\begin{aligned}\lambda \langle \phi, \phi \rangle_{\mathcal{X}} &= \langle \lambda \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \Delta \phi, \phi \rangle_{\mathcal{X}} \\ &= \langle \phi, \Delta \phi \rangle_{\mathcal{X}} \\ &= \overline{\langle \Delta \phi, \phi \rangle_{\mathcal{X}}} \\ &= \overline{\langle \lambda \phi, \phi \rangle_{\mathcal{X}}} \\ &= \bar{\lambda} \langle \phi, \phi \rangle_{\mathcal{X}}\end{aligned}$$

Hence, the eigenvalues of Δ are **real**

Laplacian eigenfunctions

$$\Delta\phi = \lambda\phi$$

Consider distinct eigenfunctions ϕ_i, ϕ_j with $\lambda_i \neq \lambda_j$. We have:

$$\langle \phi_i, \Delta\phi_j \rangle_{\mathcal{X}} = \langle \Delta\phi_i, \phi_j \rangle_{\mathcal{X}}$$

Laplacian eigenfunctions

$$\Delta\phi = \lambda\phi$$

Consider distinct eigenfunctions ϕ_i, ϕ_j with $\lambda_i \neq \lambda_j$. We have:

$$\lambda_j \langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \lambda_i \langle \phi_i, \phi_j \rangle_{\mathcal{X}}$$

Laplacian eigenfunctions

$$\Delta\phi = \lambda\phi$$

Consider distinct eigenfunctions ϕ_i, ϕ_j with $\lambda_i \neq \lambda_j$. We have:

$$\lambda_j \langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \lambda_i \langle \phi_i, \phi_j \rangle_{\mathcal{X}}$$

This can only be true if $\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = 0$

Laplacian eigenfunctions

$$\Delta\phi = \lambda\phi$$

Consider distinct eigenfunctions ϕ_i, ϕ_j with $\lambda_i \neq \lambda_j$. We have:

$$\lambda_j \langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \lambda_i \langle \phi_i, \phi_j \rangle_{\mathcal{X}}$$

This can only be true if $\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = 0$

Therefore, the eigenfunctions ϕ of Δ are **orthogonal**, and can be rescaled to be **orthonormal**:

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij}$$

where $\delta_{ij} = 1$ if $i = j$, and 0 otherwise

Spectral theorem

$$\Delta\phi = \lambda\phi$$

Now recall that the Laplacian $\Delta : \mathcal{F}(\mathcal{X}) \rightarrow \mathcal{F}(\mathcal{X})$ operates on a **vector space** of functions $\mathcal{F}(\mathcal{X})$

Spectral theorem

$$\Delta\phi = \lambda\phi$$

Now recall that the Laplacian $\Delta : \mathcal{F}(\mathcal{X}) \rightarrow \mathcal{F}(\mathcal{X})$ operates on a **vector space** of functions $\mathcal{F}(\mathcal{X})$


Theorem: The eigenfunctions $\{\phi_i\}$ of Δ form an orthonormal basis of $\mathcal{F}(\mathcal{X})$

Spectral theorem

$$\Delta\phi = \lambda\phi$$

Now recall that the Laplacian $\Delta : \mathcal{F}(\mathcal{X}) \rightarrow \mathcal{F}(\mathcal{X})$ operates on a **vector space** of functions $\mathcal{F}(\mathcal{X})$

Theorem: The eigenfunctions $\{\phi_i\}$ of Δ form an orthonormal basis of $\mathcal{F}(\mathcal{X})$



Discretization

In the **discrete setting** we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies$$

Discretization

In the **discrete setting** we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

Discretization

In the **discrete setting** we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies$$

Discretization

In the **discrete setting** we have:

$$\begin{aligned}\langle f, g \rangle_{\mathcal{X}} &\implies \mathbf{f}^\top \mathbf{A} \mathbf{g} \\ \langle f, \Delta g \rangle_{\mathcal{X}} &\implies \mathbf{f}^\top \mathbf{S} \mathbf{g}\end{aligned}$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \mathbf{\Phi}^\top \mathbf{A} \mathbf{\Phi} = \mathbf{I}$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \boldsymbol{\Phi}^\top \mathbf{A} \boldsymbol{\Phi} = \mathbf{I}$$

$$c_i = \langle \phi_i, f \rangle_{\mathcal{X}} \implies$$

Discretization

In the **discrete setting** we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \boldsymbol{\Phi}^\top \mathbf{A} \boldsymbol{\Phi} = \mathbf{I}$$

$$c_i = \langle \phi_i, f \rangle_{\mathcal{X}} \implies \mathbf{c} = \boldsymbol{\Phi}^\top \mathbf{A} \mathbf{f}$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \boldsymbol{\Phi}^\top \mathbf{A} \boldsymbol{\Phi} = \mathbf{I}$$

$$c_i = \langle \phi_i, f \rangle_{\mathcal{X}} \implies \mathbf{c} = \boldsymbol{\Phi}^\top \mathbf{A} \mathbf{f}$$

And we consider the generalized eigenvalue problem:

$$\mathbf{A}^{-1} \mathbf{S} \boldsymbol{\Phi} = \boldsymbol{\Phi} \boldsymbol{\Lambda}$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$

$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \boldsymbol{\Phi}^\top \mathbf{A} \boldsymbol{\Phi} = \mathbf{I}$$

$$c_i = \langle \phi_i, f \rangle_{\mathcal{X}} \implies \mathbf{c} = \boldsymbol{\Phi}^\top \mathbf{A} \mathbf{f}$$

And we consider the generalized eigenvalue problem:

$$\mathbf{S} \boldsymbol{\Phi} = \mathbf{A} \boldsymbol{\Phi} \boldsymbol{\Lambda}$$

Discretization

In the discrete setting we have:

$$\langle f, g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{A} \mathbf{g}$$


$$\langle f, \Delta g \rangle_{\mathcal{X}} \implies \mathbf{f}^\top \mathbf{S} \mathbf{g}$$

$$\langle \phi_i, \phi_j \rangle_{\mathcal{X}} = \delta_{ij} \implies \Phi^\top \mathbf{A} \Phi = \mathbf{I}$$

$$c_i = \langle \phi_i, f \rangle_{\mathcal{X}} \implies \mathbf{c} = \Phi^\top \mathbf{A} \mathbf{f}$$

And we consider the generalized eigenvalue problem:

$$\mathbf{S} \Phi = \mathbf{A} \Phi \Lambda$$



Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f =$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f =$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f = - \int_{\mathcal{X}} \langle \nabla 1, \nabla f \rangle =$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f = - \int_{\mathcal{X}} \langle \nabla 1, \nabla f \rangle = 0$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f = - \int_{\mathcal{X}} \langle \nabla 1, \nabla f \rangle = 0$$

In particular, let $\Delta \phi = \lambda \phi$. Then we have:

$$\int_{\mathcal{X}} \phi =$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f = - \int_{\mathcal{X}} \langle \nabla 1, \nabla f \rangle = 0$$

In particular, let $\Delta\phi = \lambda\phi$. Then we have:

$$\int_{\mathcal{X}} \phi = \frac{1}{\lambda} \int_{\mathcal{X}} \Delta\phi =$$

Integral of eigenfunctions

It is easy to see that the following holds for any $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$\int_{\mathcal{X}} \Delta f = \int_{\mathcal{X}} 1 \Delta f = - \int_{\mathcal{X}} \langle \nabla 1, \nabla f \rangle = 0$$

In particular, let $\Delta\phi = \lambda\phi$. Then we have:

$$\int_{\mathcal{X}} \phi = \frac{1}{\lambda} \int_{\mathcal{X}} \Delta\phi = 0$$

In other words, eigenfunctions have **zero average**

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 =$$

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle =$$

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle = \int_{\mathcal{X}} \phi \Delta \phi =$$

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle = \int_{\mathcal{X}} \phi \Delta \phi = \lambda \int_{\mathcal{X}} \phi \phi =$$

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle = \int_{\mathcal{X}} \phi \Delta \phi = \lambda \int_{\mathcal{X}} \phi \phi = \lambda$$

So **eigenvalues are the Dirichlet energies of eigenfunctions**

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle = \int_{\mathcal{X}} \phi \Delta \phi = \lambda \int_{\mathcal{X}} \phi \phi = \lambda$$

So **eigenvalues are the Dirichlet energies of eigenfunctions**

Further, $\lambda = 0$ only when $\nabla \phi = 0$, i.e., when ϕ is **constant**

Dirichlet energy

The Dirichlet energy of a function $f : \mathcal{X} \rightarrow \mathbb{R}$ measures its **smoothness**:

$$\mathcal{E}(f) = \int_{\mathcal{X}} \|\nabla f(x)\|^2 dx$$

For a Laplacian eigenfunction ϕ with eigenvalue λ :

$$\int_{\mathcal{X}} \|\nabla \phi\|^2 = \int_{\mathcal{X}} \langle \nabla \phi, \nabla \phi \rangle = \int_{\mathcal{X}} \phi \Delta \phi = \lambda \int_{\mathcal{X}} \phi \phi = \lambda$$

So **eigenvalues are the Dirichlet energies of eigenfunctions**

Further, $\lambda = 0$ only when $\nabla \phi = 0$, i.e., when ϕ is **constant**; in particular, $\lambda = 0$ is **always** an eigenvalue of Δ , since $\Delta f = 0$ for any constant f

Discrete spectrum


The eigenvalues of the Laplacian are **countable** and are canonically ordered non-decreasingly:

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \rightarrow \infty$$

Discrete spectrum

The eigenvalues of the Laplacian are **countable** and are canonically ordered non-decreasingly:

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \rightarrow \infty$$



Discrete spectrum

The eigenvalues of the Laplacian are **countable** and are canonically ordered non-decreasingly:

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \cdots \rightarrow \infty$$

The Laplacian eigenvalues follow **Weyl's asymptotic law**:

$$\lambda_i \approx \frac{\pi}{\int_{\mathcal{X}} da} i \quad \text{for } i \rightarrow \infty$$

That is, they grow **linearly** with growth rate that is inversely proportional to **surface area**

Laplacian eigenvalues: Properties

The Laplacian spectrum is particularly useful in shape analysis:

Laplacian eigenvalues: Properties

The Laplacian spectrum is particularly useful in shape analysis:

- It is **isometry invariant**

Laplacian eigenvalues: Properties

The Laplacian spectrum is particularly useful in shape analysis:

- It is **isometry invariant**
- It depends **continuously** on the Riemannian metric

Laplacian eigenvalues: Properties

The Laplacian spectrum is particularly useful in shape analysis:

- It is **isometry invariant**
- It depends **continuously** on the Riemannian metric
- It is easy and **efficient** to compute

Laplacian eigenvalues: Properties

The Laplacian spectrum is particularly useful in shape analysis:


- It is **isometry invariant**
- It depends **continuously** on the Riemannian metric
- It is easy and **efficient** to compute
- It is **informative** as it encodes metric and topological information of the surface (e.g. surface area, presence of symmetries, etc.)

Laplacian eigenvalues: Properties


The Laplacian spectrum is particularly useful in shape analysis:

- It is **isometry invariant**
- It depends **continuously** on the Riemannian metric
- It is easy and **efficient** to compute
- It is **informative** as it encodes metric and topological information of the surface (e.g. surface area, presence of symmetries, etc.)

However, one “cannot hear the shape of the drum”




Laplacian eigenfunctions: Euclidean




First eigenfunctions of 1D Euclidean Laplacian = standard Fourier basis

Laplacian eigenfunctions: manifold



First eigenfunctions of a manifold Laplacian

Laplacian eigenfunctions: graph




First eigenfunctions of a graph Laplacian

Fourier analysis: Euclidean space

A function $f : [-\pi, \pi] \rightarrow \mathbb{R}$ can be written as Fourier series


$$f(x) = \sum_{k \geq 0} \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x') e^{-ikx'} dx' e^{ikx}$$



Fourier analysis: Euclidean space

A function $f : [-\pi, \pi] \rightarrow \mathbb{R}$ can be written as Fourier series


$$f(x) = \sum_{k \geq 0} \underbrace{\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x') e^{-ikx'} dx'}_{\hat{f}_k = \langle f, e^{ikx} \rangle_{L^2([-\pi, \pi])}} e^{ikx}$$



Fourier analysis: Euclidean space

A function $f : [-\pi, \pi] \rightarrow \mathbb{R}$ can be written as Fourier series

$$f(x) = \sum_{k \geq 0} \underbrace{\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x') e^{-ikx'} dx'}_{\hat{f}_k = \langle f, e^{ikx} \rangle_{L^2([-\pi, \pi])}} e^{ikx}$$




Fourier basis = Laplacian eigenfunctions: $\frac{d^2}{dx^2} e^{ikx} = k^2 e^{ikx}$

Fourier analysis: non-Euclidean space

A function $f : \mathcal{X} \rightarrow \mathbb{R}$ can be written as Fourier series

$$f(x) = \sum_{k \geq 1} \underbrace{\int_{\mathcal{X}} f(x') \phi_k(x') dx'}_{\hat{f}_k = \langle f, \phi_k \rangle_{L^2(\mathcal{X})}} \phi_k(x)$$



Fourier basis = Laplacian eigenfunctions: $\Delta \phi_k(x) = \lambda_k \phi_k(x)$


Teaser exercise: Maps in the Laplacian eigenbasis

Consider two **isometric** shapes \mathcal{X}, \mathcal{Y} with Laplacian eigenfunctions $\{\phi_i\}, \{\psi_j\}$ respectively spanning the functional spaces $\mathcal{F}(\mathcal{X}), \mathcal{F}(\mathcal{Y})$.

Teaser exercise: Maps in the Laplacian eigenbasis

Consider two **isometric** shapes \mathcal{X}, \mathcal{Y} with Laplacian eigenfunctions $\{\phi_i\}, \{\psi_j\}$ respectively spanning the functional spaces $\mathcal{F}(\mathcal{X}), \mathcal{F}(\mathcal{Y})$.


Now let $T : \mathcal{F}(\mathcal{X}) \rightarrow \mathcal{F}(\mathcal{Y})$ be the ground-truth map.



Teaser exercise: Maps in the Laplacian eigenbasis

Consider two **isometric** shapes \mathcal{X}, \mathcal{Y} with Laplacian eigenfunctions $\{\phi_i\}, \{\psi_j\}$ respectively spanning the functional spaces $\mathcal{F}(\mathcal{X}), \mathcal{F}(\mathcal{Y})$.

Now let $T : \mathcal{F}(\mathcal{X}) \rightarrow \mathcal{F}(\mathcal{Y})$ be the ground-truth map.



How does the matrix representation of T look like, in the **Laplacian eigenbases**?