

# Fundamentals of Computer Graphics

The Laplace-Beltrami operator

Emanuele Rodolà  
[rodola@di.uniroma1.it](mailto:rodola@di.uniroma1.it)



SAPIENZA  
UNIVERSITÀ DI ROMA

# Course quality questionnaire

The questionnaire is completely **anonymous** – your privacy is respected

Instructions:

- Disable pop-up block in your browser
- Go to <https://www.uniroma1.it>
- Click on **Studenti** and access **Infostud 2.0**
- Click on **Corsi di laurea**
- On the left menu, click on **Opinioni studenti**
- Enter the OPIS code and click on **Vai al questionario**

OPIS codes are:

- **LNM5IDE3**
- **CUP342HA** for the curriculum “Networks and security”

## Motivation: Heat diffusion

For a subset  $U \subset \mathbb{R}^2$  the diffusion of heat is described by the [heat equation](#):

$$\begin{aligned}\frac{\partial}{\partial t} u(x, t) &= \Delta u(x, t) \\ u(x, 0) &= u_0(x)\end{aligned}$$

## Motivation: Heat diffusion

For a subset  $U \subset \mathbb{R}^2$  the diffusion of heat is described by the [heat equation](#):

$$\begin{aligned}\frac{\partial}{\partial t} u(x, t) &= \Delta u(x, t) \\ u(x, 0) &= u_0(x)\end{aligned}$$

where the [Laplacian](#) is defined as:

$$\Delta u(x, t) = -\operatorname{div}(\nabla u) = \frac{\partial^2 u(x, t)}{\partial x_1^2} + \frac{\partial^2 u(x, t)}{\partial x_2^2}$$


## Motivation: Heat diffusion

For a subset  $U \subset \mathbb{R}^2$  the diffusion of heat is described by the [heat equation](#):

$$\begin{aligned}\frac{\partial}{\partial t} u(x, t) &= \Delta u(x, t) \\ u(x, 0) &= u_0(x)\end{aligned}$$

where the [Laplacian](#) is defined as:

$$\Delta u(x, t) = -\operatorname{div}(\nabla u) = \frac{\partial^2 u(x, t)}{\partial x_1^2} + \frac{\partial^2 u(x, t)}{\partial x_2^2}$$



## Motivation: Heat diffusion on a surface

For a **surface**  $\mathcal{X} \in \mathbb{R}^3$  the diffusion of heat is described by the **heat equation**:

$$\begin{aligned}\frac{\partial}{\partial t} u(x, t) &= \Delta u(x, t) \\ u(x, 0) &= u_0(x)\end{aligned}$$


where the **Laplacian** is defined as:

$$\Delta u(x, t) = -\text{div}(\nabla u) = ?$$


## Inner product on a manifold

Given two **scalar functions**  $f, g : \mathcal{X} \rightarrow \mathbb{R}$ , we define their inner product as:

$$\langle f, g \rangle_{\mathcal{X}} = \int_{\mathcal{X}} f(x)g(x)dx$$



$f : \mathcal{X} \rightarrow \mathbb{R}$



$g : \mathcal{X} \rightarrow \mathbb{R}$

## Inner product on a manifold


Given two **scalar functions**  $f, g : \mathcal{X} \rightarrow \mathbb{R}$ , we define their inner product as:

$$\langle f, g \rangle_{\mathcal{X}} = \int_{\mathcal{X}} f(x)g(x)dx$$


When discretized on a triangle mesh of  $n$  vertices, this boils down to:

$$\mathbf{f}^\top \text{diag}(a_i) \mathbf{g}$$

where  $a_i$  with  $i = 1, \dots, n$  are the local **area elements** at each vertex



$f : \mathcal{X} \rightarrow \mathbb{R}$



$g : \mathcal{X} \rightarrow \mathbb{R}$

## Inner product on a manifold

Given two **tangent vector fields**  $F, G : \mathcal{X} \rightarrow T\mathcal{X}$ , we define their inner product as:

$$\langle F, G \rangle_{T\mathcal{X}} = \int_{\mathcal{X}} \langle F(x), G(x) \rangle_{T_x \mathcal{X}} dx$$




$F : \mathcal{X} \rightarrow T\mathcal{X}$



$G : \mathcal{X} \rightarrow T\mathcal{X}$


# Geometric intuition



Smooth **scalar field**  $f$

# Geometric intuition


- Gradient  $\nabla f(x)$   
‘direction of the steepest increase  
of  $f$  at  $x$ ’



Smooth vector field  $F$

# Geometric intuition


- Gradient  $\nabla f(x)$   
‘direction of the steepest increase  
of  $f$  at  $x$ ’
- Divergence  $\operatorname{div}(F(x))$   
‘scalar density of an outward flux of  
 $F$  from an infinitesimal volume  
around  $x$ ’




Smooth vector field  $F$

# Geometric intuition


- Gradient  $\nabla f(x)$   
‘direction of the steepest increase of  $f$  at  $x$ ’
- Divergence  $\operatorname{div}(F(x))$   
‘scalar density of an outward flux of  $F$  from an infinitesimal volume around  $x$ ’
- Laplacian  $\Delta f(x) = -\operatorname{div}(\nabla f(x))$   
‘scalar difference between  $f(x)$  and the average of  $f$  on an infinitesimal sphere around  $x$ ’




# Adjointness



# Adjointness




# Adjointness



Gradient and divergence are (negative) **adjoint** operators:

$$\langle F, \nabla f \rangle_{T\mathcal{X}} = -\langle \operatorname{div} F, f \rangle_{\mathcal{X}}$$

# Adjointness



Gradient and divergence are (negative) **adjoint** operators:

$$\langle F, \nabla f \rangle_{T\mathcal{X}} = -\langle \operatorname{div} F, f \rangle_{\mathcal{X}}$$

Notice that the first inner product is among **vector fields**, while the second is among **scalar functions**

# Finite elements method (FEM)

We will **discretize** the Laplace-Beltrami operator using a new approach

Overall idea:

# Finite elements method (FEM)

We will **discretize** the Laplace-Beltrami operator using a new approach

Overall idea:

- Consider the equation

$$\Delta f(x) = g(x)$$

We are interested in computing  $g$ , and then directly “solve for”  $\Delta$

# Finite elements method (FEM)

We will **discretize** the Laplace-Beltrami operator using a new approach

Overall idea:

- Consider the equation

$$\Delta f(x) = g(x)$$

We are interested in computing  $g$ , and then directly “solve for”  $\Delta$

- To do so, we instead look at the **weak formulation**

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

where  $h_j : \mathcal{X} \rightarrow \mathbb{R}$  are some **test functions**

# Finite elements method (FEM)

We will **discretize** the Laplace-Beltrami operator using a new approach

Overall idea:

- Consider the equation

$$\Delta f(x) = g(x)$$

We are interested in computing  $g$ , and then directly “solve for”  $\Delta$

- To do so, we instead look at the **weak formulation**

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

where  $h_j : \mathcal{X} \rightarrow \mathbb{R}$  are some **test functions**

- We obtain expressions for the left- and right-hand sides, from which we explicitly compute a **matrix representation** for  $\Delta$

## Hat basis

Recall that on triangle meshes we approximate scalar functions as

$$f(x) \approx \sum_{i=1}^n f(v_i) h_i(x)$$


where  $h_i$  are **hat basis functions** defined at each vertex:

## Hat basis

Recall that on triangle meshes we approximate scalar functions as

$$f(x) \approx \sum_{i=1}^n f(v_i) h_i(x)$$

where  $h_i$  are **hat basis functions** defined at each vertex:



## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}}$$

## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\begin{aligned}\langle \Delta f, h_j \rangle_{\mathcal{X}} &= -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}} \\ &= \langle \nabla f, \nabla h_j \rangle_{T\mathcal{X}}\end{aligned}$$

## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\begin{aligned}\langle \Delta f, h_j \rangle_{\mathcal{X}} &= -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}} \\ &= \langle \nabla f, \nabla h_j \rangle_{T\mathcal{X}} \\ &= \langle \nabla \sum_i f(v_i) h_i(x), \nabla h_j \rangle_{T\mathcal{X}}\end{aligned}$$

## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\begin{aligned}\langle \Delta f, h_j \rangle_{\mathcal{X}} &= -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}} \\ &= \langle \nabla f, \nabla h_j \rangle_{T\mathcal{X}} \\ &= \langle \nabla \sum_i f(v_i) h_i(x), \nabla h_j \rangle_{T\mathcal{X}} \\ &= \sum_i f(v_i) \langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}}\end{aligned}$$

## FEM discretization

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\begin{aligned}\langle \Delta f, h_j \rangle_{\mathcal{X}} &= -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}} \\ &= \langle \nabla f, \nabla h_j \rangle_{T\mathcal{X}} \\ &= \langle \nabla \sum_i f(v_i) h_i(x), \nabla h_j \rangle_{T\mathcal{X}} \\ &= \sum_i f(v_i) \underbrace{\langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}}}_{s_{ij}=s_{ji}}\end{aligned}$$

## FEM discretization: Stiffness

We use the hat basis as test functions in the equation:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

We can rewrite the left-hand side as:

$$\begin{aligned}\langle \Delta f, h_j \rangle_{\mathcal{X}} &= -\langle \operatorname{div} \nabla f, h_j \rangle_{\mathcal{X}} \\ &= \langle \nabla f, \nabla h_j \rangle_{T\mathcal{X}} \\ &= \langle \nabla \sum_i f(v_i) h_i(x), \nabla h_j \rangle_{T\mathcal{X}} \\ &= \sum_i f(v_i) \underbrace{\langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}}}_{s_{ij}=s_{ji}}\end{aligned}$$

where  $\mathbf{S} = (s_{ij}) \in \mathbb{R}^{n \times n}$  is the symmetric **stiffness matrix**; so we have:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = (\mathbf{S}\mathbf{f})_j$$

## FEM discretization

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

## FEM discretization

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

For the right-hand side we have:

$$\langle g, h_j \rangle_{\mathcal{X}} = \left\langle \sum_i g(v_i) h_i(x), h_j \right\rangle_{\mathcal{X}}$$

## FEM discretization

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

For the right-hand side we have:

$$\begin{aligned}\langle g, h_j \rangle_{\mathcal{X}} &= \left\langle \sum_i g(v_i) h_i(x), h_j \right\rangle_{\mathcal{X}} \\ &= \sum_i g(v_i) \langle h_i, h_j \rangle_{\mathcal{X}}\end{aligned}$$

## FEM discretization: Mass

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

For the right-hand side we have:

$$\begin{aligned}\langle g, h_j \rangle_{\mathcal{X}} &= \left\langle \sum_i g(v_i) h_i(x), h_j \right\rangle_{\mathcal{X}} \\ &= \sum_i g(v_i) \underbrace{\langle h_i, h_j \rangle_{\mathcal{X}}}_{a_{ij}=a_{ji}}\end{aligned}$$

where  $\mathbf{A} = (a_{ij}) \in \mathbb{R}^{n \times n}$  is the symmetric **mass matrix**; so we have:

$$\langle g, h_j \rangle_{\mathcal{X}} = (\mathbf{A}\mathbf{g})_j$$

## Discrete Laplace-Beltrami operator

We get to:

$$\langle \Delta f, h_j \rangle_{\mathcal{X}} = \langle g, h_j \rangle_{\mathcal{X}}$$

## Discrete Laplace-Beltrami operator

We get to:

$$(\mathbf{S}\mathbf{f})_j = (\mathbf{A}\mathbf{g})_j \quad \text{for } j = 1, \dots, n$$

## Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{S}\mathbf{f} = \mathbf{A}\mathbf{g}$$

## Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

## Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

Our **discrete Laplacian** is thus given by:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

# Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

Our **discrete Laplacian** is thus given by:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

Some important **properties** that we will see of this discretization:

- The matrices involved are **sparse** and **symmetric**

# Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

Our **discrete Laplacian** is thus given by:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

Some important **properties** that we will see of this discretization:

- The matrices involved are **sparse** and **symmetric**
- They have the same structure as the vertex **adjacency** matrix

# Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

Our **discrete Laplacian** is thus given by:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

Some important **properties** that we will see of this discretization:

- The matrices involved are **sparse** and **symmetric**
- They have the same structure as the vertex **adjacency** matrix
- The stiffness **S** is **positive semi-definite**

# Discrete Laplace-Beltrami operator

We get to:

$$\mathbf{A}^{-1}\mathbf{S}\mathbf{f} = \mathbf{g}$$

Our **discrete Laplacian** is thus given by:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

Some important **properties** that we will see of this discretization:

- The matrices involved are **sparse** and **symmetric**
- They have the same structure as the vertex **adjacency** matrix
- The stiffness **S** is **positive semi-definite**
- They can be computed easily and **efficiently** for any given mesh

## Mass integral


Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):

$$a_{ij} = \langle \mathbf{h}_i, \mathbf{h}_j \rangle_{\mathcal{X}}$$

# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):


$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \end{aligned}$$



# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):


$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \\ &= \int_{T_{jii'}} h_i(x) h_j(x) dx + \int_{T_{ji''i}} h_i(x) h_j(x) dx \end{aligned}$$



# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \\ &= \int_{T_{jii'}} h_i(x) h_j(x) dx + \int_{T_{ji''i}} h_i(x) h_j(x) dx \end{aligned}$$




Functions  $h_i$  and  $h_j$  decrease linearly from 1 to 0

# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \\ &= \int_{T_{jii'}} h_i(x) h_j(x) dx + \int_{T_{ji''i}} h_i(x) h_j(x) dx \end{aligned}$$



Functions  $h_i$  and  $h_j$  decrease linearly from 1 to 0


So, in parameter space, each of the two integrals can be written as:

$$2A(T_{\ell}) \int_0^1 \int_0^{1-u} \mathbf{v} \mathbf{u} dv du$$

# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \\ &= \int_{T_{jii'}} h_i(x) h_j(x) dx + \int_{T_{ji''i}} h_i(x) h_j(x) dx \end{aligned}$$



Functions  $h_i$  and  $h_j$  decrease linearly from 1 to 0


So, in parameter space, each of the two integrals can be written as:

$$2A(T_{\ell}) \int_0^1 \int_0^{1-u} \mathbf{v} \mathbf{u} dv du = 2A(T_{\ell}) \frac{1}{24}$$

# Mass integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{ij} &= \langle h_i, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_i(x) h_j(x) dx \\ &= \int_{T_{jii'}} h_i(x) h_j(x) dx + \int_{T_{ji''i}} h_i(x) h_j(x) dx \end{aligned}$$



Functions  $h_i$  and  $h_j$  decrease linearly from 1 to 0

So, in parameter space, each of the two integrals can be written as:

$$2A(T_{\ell}) \int_0^1 \int_0^{1-u} \mathbf{v} \mathbf{u} dv du = 2A(T_{\ell}) \frac{1}{24} = \frac{1}{12} A(T_{\ell})$$

## Mass integral


Now assume  $i = j$  (diagonal of  $\mathbf{A}$ ):

$$a_{jj} = \langle \mathbf{h}_j, \mathbf{h}_j \rangle_{\mathcal{X}}$$

# Mass integral

Now assume  $i = j$  (diagonal of  $\mathbf{A}$ ):


$$\begin{aligned} a_{jj} &= \langle \mathbf{h}_j, \mathbf{h}_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} \mathbf{h}_j(x) \mathbf{h}_j(x) dx \end{aligned}$$



# Mass integral

Now assume  $i = j$  (diagonal of  $\mathbf{A}$ ):


$$\begin{aligned} a_{jj} &= \langle h_j, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_j(x) h_j(x) dx \\ &= \sum_{i \in \mathcal{N}(j)} \int_{T_{jii'}} h_j(x) h_j(x) dx \end{aligned}$$



# Mass integral

Now assume  $i = j$  (diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{jj} &= \langle \mathbf{h}_j, \mathbf{h}_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} \mathbf{h}_j(x) \mathbf{h}_j(x) dx \\ &= \sum_{i \in \mathcal{N}(j)} \int_{T_{jii'}} \mathbf{h}_j(x) \mathbf{h}_j(x) dx \end{aligned}$$




Like before, each integral in parameter space looks like:

$$2A(T_{\ell}) \int_0^1 \int_0^{1-u} \mathbf{u}^2 dv du$$

# Mass integral

Now assume  $i = j$  (diagonal of  $\mathbf{A}$ ):

$$\begin{aligned} a_{jj} &= \langle h_j, h_j \rangle_{\mathcal{X}} \\ &= \sum_{\ell} \int_{T_{\ell}} h_j(x) h_j(x) dx \\ &= \sum_{i \in \mathcal{N}(j)} \int_{T_{jii'}} h_j(x) h_j(x) dx \end{aligned}$$



Like before, each integral in parameter space looks like:

$$2A(T_{\ell}) \int_0^1 \int_0^{1-u} u^2 dv du = \frac{1}{6} A(T_{\ell})$$

## Stiffness integral


Assume  $i \neq j$  (off-diagonal of  $\mathbf{S}$ ):

$$s_{ij} = \langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}}$$

# Stiffness integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{S}$ ):


$$\begin{aligned}s_{ij} &= \langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}} \\ &= \sum_{\ell} \int_{T_\ell} \langle \nabla h_i(x) \nabla h_j(x) \rangle dx\end{aligned}$$



# Stiffness integral

Assume  $i \neq j$  (off-diagonal of  $\mathbf{S}$ ):

$$\begin{aligned}s_{ij} &= \langle \nabla h_i, \nabla h_j \rangle_{T\mathcal{X}} \\&= \sum_{\ell} \int_{T_\ell} \langle \nabla h_i(x) \nabla h_j(x) \rangle dx \\&= -\frac{1}{2}(\cot \alpha_{ij} + \cot \beta_{ij})\end{aligned}$$



The integrals are non-zero in the same positions as the mass matrix

The formulas involve internal angles (also known as “cotangent Laplacian”)

# Stiffness and mass matrices

The discrete (FEM) Laplace-Beltrami operator is the  $n \times n$  matrix:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

where

$$s_{ij} = \begin{cases} -\frac{1}{2}(\cot\alpha_{ij} + \cot\beta_{ij}) & \text{if } e_{ij} \in E \\ -\sum_{k \neq i} s_{ik} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

$$a_{ij} = \begin{cases} \frac{1}{12}(A(T_{jii'}) + A(T_{ji''i})) & \text{if } e_{ij} \in E \\ \frac{1}{6} \sum_{k \in \mathcal{N}(i)} A(T_k) & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

# Stiffness and mass matrices

The discrete (FEM) Laplace-Beltrami operator is the  $n \times n$  matrix:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

where

$$s_{ij} = \begin{cases} -\frac{1}{2}(\cot\alpha_{ij} + \cot\beta_{ij}) & \text{if } e_{ij} \in E \\ -\sum_{k \neq i} s_{ik} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$
$$a_{ij} = \begin{cases} \frac{1}{12}(A(T_{jii'}) + A(T_{ji''i})) & \text{if } e_{ij} \in E \\ \frac{1}{6} \sum_{k \in \mathcal{N}(i)} A(T_k) & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

A **lumped mass** matrix (easier to invert) is obtained as:

$$\tilde{a}_{ij} = \begin{cases} \sum_{k=1}^n a_{ik} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

# Stiffness and mass matrices

The discrete (FEM) Laplace-Beltrami operator is the  $n \times n$  matrix:

$$\mathbf{L} = \mathbf{A}^{-1}\mathbf{S}$$

where

$$s_{ij} = \begin{cases} -\frac{1}{2}(\cot\alpha_{ij} + \cot\beta_{ij}) & \text{if } e_{ij} \in E \\ -\sum_{k \neq i} s_{ik} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$
$$a_{ij} = \begin{cases} \frac{1}{12}(A(T_{jii'}) + A(T_{ji''i})) & \text{if } e_{ij} \in E \\ \frac{1}{6} \sum_{k \in \mathcal{N}(i)} A(T_k) & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

A **lumped mass** matrix (easier to invert) is obtained as:

$$\tilde{a}_{ij} = \begin{cases} \frac{1}{3} \sum_{k \in \mathcal{N}(i)} A(T_k) & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

# Exercise: Discrete Laplacian

Implement the FEM Laplacian

Test it similarly to what we did with the graph Laplacian to:

- Denoise a scalar function on a surface
- Skeletonize a mesh
- Implement least-squares meshes

## Suggested reading

- S. Axler, “Linear algebra done right – 3rd edition”. Springer, 2015  
*Section 7.A*
- Reuter et al., “Discrete Laplace-Beltrami operators for shape analysis and segmentation”. CAG 33, 2009.  
*Sections 1 to 2.1.2*