

Notes for *Foundations of Modern Analysis* by
Avner Friedman

Anton Ottosson – antonott@kth.se

August 27, 2024

Chapter 1 – Measure Theory

Section 1.1 – Rings and algebras

Problems

1.1.1

$$\left(\varinjlim_n E_n\right)^c = \varprojlim_n E_n^c, \quad \left(\varprojlim_n E_n\right)^c = \varinjlim_n E_n^c.$$

Solution. For the first identity, note that

$$\begin{aligned} x \in \left(\varinjlim_n E_n\right)^c &\iff x \notin \varinjlim_n E_n \\ &\iff x \notin E_n \text{ for infinitely many } n \\ &\iff x \in E_n^c \text{ for infinitely many } n \\ &\iff x \in \varprojlim_n E_n^c. \end{aligned}$$

For the second,

$$\begin{aligned} x \in \left(\varprojlim_n E_n\right)^c &\iff x \notin \varprojlim_n E_n \\ &\iff x \in E_n \text{ for finitely many } n \\ &\iff x \in E_n^c \text{ for all but finitely many } n \\ &\iff x \in \varinjlim_n E_n^c. \end{aligned}$$

1.1.2

$$\varprojlim_n E_n = \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_n, \quad \varinjlim_n E_n = \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_n.$$

Solution. Suppose $x \in \varprojlim_n E_n$. Then $x \in E_n$ for infinitely many n . It follows that $x \in \bigcup_{n=k}^{\infty} E_n$ for all $k \in \mathbb{N}$, and hence that $x \in \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_n$.

Conversely, assume that $x \in \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_n$. Then $x \in \bigcup_{n=k}^{\infty} E_n$ for all $k \in \mathbb{N}$. It follows that $x \in E_n$ for infinitely many n , and thus that $x \in \varlimsup_n E_n$. This proves the first identity.

Next, suppose that $x \in \varliminf_n E_n$. Then $x \in E_n$ for all but finitely many n , so there is some $k' \in \mathbb{N}$ such that $x \in E_n$ for all $n \geq k'$. It follows that $x \in \bigcap_{n=k'}^{\infty} E_n$, and hence that $x \in \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_n$.

Conversely, assume that $x \in \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_n$. Then $x \in \bigcap_{n=k'}^{\infty} E_n$ for some $k' \in \mathbb{N}$, which means that $x \in E_n$ for all $n \geq k'$. It follows that $x \in E_n$ for all but finitely many n ; that is, $x \in \varliminf_n E_n$.

1.1.3

If \mathcal{R} is a σ -ring and $E_n \in \mathcal{R}$, then

$$\bigcap_{n=1}^{\infty} E_n \in \mathcal{R}, \quad \overline{\lim}_n E_n \in \mathcal{R}, \quad \varliminf_n E_n \in \mathcal{R}.$$

Solution. Let $Y = \bigcup_{n=1}^{\infty} E_n$. Then $E_n \subset Y$ for all n , and it follows that

$$\bigcap_{n=1}^{\infty} E_n = Y \cap \left(\bigcap_{n=1}^{\infty} E_n \right) = Y - \left(Y - \bigcap_{n=1}^{\infty} E_n \right).$$

Notice that

$$Y - \bigcap_{n=1}^{\infty} E_n = \bigcup_{n=1}^{\infty} (Y - E_n) \in \mathcal{R},$$

by properties (b) and (e). (The equality is analogous to the identity (1.1.2), but with Y in place of X .) Hence, applying (b) again, we find that

$$\bigcap_{n=1}^{\infty} E_n = Y - \left(Y - \bigcap_{n=1}^{\infty} E_n \right) \in \mathcal{R}.$$

For later reference, let us call this result (x).

Given $k \in \mathbb{N}$, let $A_n = \emptyset$ for $n < k$, and let $A_n = E_n$ for $n \geq k$. Then $A_n \in \mathcal{R}$ for all n by (a), hence

$$\bigcup_{n=k}^{\infty} E_n = \bigcup_{n=1}^{\infty} A_n \in \mathcal{R}$$

by (e). It then follows by (x) that

$$\overline{\lim}_n E_n = \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_n \in \mathcal{R}.$$

By a similar argument we find that (x) implies

$$\bigcap_{n=k}^{\infty} E_n \in \mathcal{R}$$

for all $k \in \mathbb{N}$. Hence

$$\varliminf_n E_n = \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_n \in \mathcal{R}$$

by (e).

1.1.4

The intersection of any collection of rings (algebras, σ -rings, or σ -algebras) is also a ring (an algebra, σ -ring, or σ -algebra).

Solution. Let \mathcal{C} be a collection of classes. Let $\bigcap \mathcal{C}$ denote the intersection of all classes in \mathcal{C} . We will show that if one of the properties (a)-(e) is satisfied by all classes in \mathcal{C} , then $\bigcap \mathcal{C}$ satisfies that property as well. The result requested in the problem then follows as an immediate corollary.

It is clear that if every $\mathcal{R} \in \mathcal{C}$ satisfies (a), then so does $\bigcap \mathcal{C}$. Suppose every $\mathcal{R} \in \mathcal{C}$ satisfies (b). If $A, B \in \bigcap \mathcal{C}$ then $A, B \in \mathcal{R}$ for every $\mathcal{R} \in \mathcal{C}$. Hence $A - B \in \mathcal{R}$ for all $\mathcal{R} \in \mathcal{C}$, and it follows that $A - B \in \bigcap \mathcal{C}$. The argument for (c) is similar (with $A \cup B$ in place of $A - B$), and (d) is obvious.

Finally, suppose that every $\mathcal{R} \in \mathcal{C}$ satisfies (e). If $A_1, A_2, \dots \in \bigcap \mathcal{C}$ then $A_1, A_2, \dots \in \mathcal{R}$ for every $\mathcal{R} \in \mathcal{C}$. Hence $\bigcup_{n=1}^{\infty} A_n \in \mathcal{R}$ for all $\mathcal{R} \in \mathcal{C}$, and it follows that $\bigcup_{n=1}^{\infty} A_n \in \bigcap \mathcal{C}$.

1.1.5

If \mathcal{D} is any class of sets, then there exists a unique ring \mathcal{R}_0 such that (i) $\mathcal{R}_0 \supset \mathcal{D}$, and (ii) any ring \mathcal{R} containing \mathcal{D} contains also \mathcal{R}_0 . \mathcal{R}_0 is called the *ring generated* by \mathcal{D} , and is denoted by $\mathcal{R}(\mathcal{D})$.

Solution. Let \mathcal{R}_0 be the intersection of all rings containing \mathcal{D} . This is a ring by the previous exercise, and it satisfies the properties (i) and (ii). To see that it is unique, let \mathcal{R}'_0 also be a ring satisfying (i) and (ii). Then $\mathcal{R}_0 \subset \mathcal{R}'_0$ and $\mathcal{R}'_0 \subset \mathcal{R}_0$ by property (ii), so $\mathcal{R}_0 = \mathcal{R}'_0$.

1.1.6

If \mathcal{D} is any class of sets, then there exists a unique σ -ring \mathcal{S}_0 such that (i) $\mathcal{S}_0 \supset \mathcal{D}$, and (ii) any σ -ring containing \mathcal{D} contains also \mathcal{S}_0 . We call \mathcal{S}_0 the *σ -ring generated* by \mathcal{D} , and denote it by $\mathcal{S}(\mathcal{D})$. A similar result holds for σ -algebras, and we speak of the *σ -algebra generated* by \mathcal{D} .

Solution. By the same argument as in the previous exercise, \mathcal{S}_0 is the intersection of all σ -rings containing \mathcal{D} . Similarly the σ -algebra generated by \mathcal{D} is the intersection of all σ -algebras containing \mathcal{D} .