

Faster Algebraic Shifting

Antony Della Vecchia, Michael Joswig and Fabian Lenzen



TU Berlin

ISSAC 2025-08-01

Algebraic Shifting



Definition A k -uniform hypergraph is a collection $S \subseteq \binom{[n]}{k}$.

S is shifted if for any $\sigma \in S$, $\sigma|_i^j \in S$ for $i \in \sigma$ and $j < i$.



Algebraic Shifting

Definition A k -uniform hypergraph is a collection $S \subseteq \binom{[n]}{k}$.
 S is shifted if for any $\sigma \in S$, $\sigma|_i^j \in S$ for $i \in \sigma$ and $j < i$.



Theorem Kalai '84, Björner-Kalai '88 Given Σ a simplicial complex and a field $\mathbb{F}$, there exists a shifted simplicial complex $\Delta^{\mathbb{F}}(\Sigma)$ with the same f -vector and Betti numbers as Σ .

Theorem DV, Joswig, Lenzen '25 There exists a fast in practice Las Vegas algorithm for determining $\Delta^{\mathbb{F}}(S)$.

```
julia> using Oscar
```

```
julia> S = uniform_hypergraph([[1, 2], [2, 3], [3, 4], [4, 1]])  
UniformHypergraph{4, 2, [[1, 2], [1, 4], [2, 3], [3, 4]]}
```

```
julia> exterior_shift(GF(2), S)  
UniformHypergraph{4, 2, [[1, 2], [1, 3], [1, 4], [2, 3]]}
```

Theorem DV, Joswig, Lenzen '25 There exists a fast in practice Las Vegas algorithm for determining $\Delta^{\mathbb{F}}(S)$.

```
julia> using Oscar
```

```
julia> S = uniform_hypergraph([[1, 2], [2, 3], [3, 4], [4, 1]])  
UniformHypergraph{4, 2, [[1, 2], [1, 4], [2, 3], [3, 4]]}
```

```
julia> exterior_shift(GF(2), S)  
UniformHypergraph{4, 2, [[1, 2], [1, 3], [1, 4], [2, 3]]}
```

Problem Kalai '02 Does there exist a deterministic or Las Vegas polynomial time algorithm for determining $\Delta(S)$?

Theorem Keehn, Nevo '24+ There exists a deterministic polynomial time algorithm for computing $\Delta^{\mathbb{Q}}(\Sigma)$ when Σ is a triangulation of either the torus, the klein bottle or the real projective plane.

Exterior Algebraic Shift

Definition Let $\mathbb{E} \supset \mathbb{F}$ be a field extension and let $g \in \mathrm{GL}(n, \mathbb{E})$.

- The k th compound matrix $g^{\wedge k}$ is the $\binom{n}{k} \times \binom{n}{k}$ -matrix of k -minors of g .
- $g^{\wedge S}$ is the $|S| \times \binom{n}{k}$ -submatrix of $g^{\wedge k}$ with rows indexed by S .

Let $\sigma_i \in S$, $|S| = m$ and $\tau_i \in \binom{[n]}{k}$ such that

$$g^{\wedge S} = \begin{pmatrix} \det g_{\sigma_1 \tau_1} & \det g_{\sigma_1 \tau_2} & \cdots & \det g_{\sigma_m \tau_{\binom{n}{k}}} \\ \vdots & \vdots & \vdots & \vdots \\ \det g_{\sigma_m \tau_1} & \det g_{\sigma_m \tau_2} & \cdots & \det g_{\sigma_m \tau_{\binom{n}{k}}} \end{pmatrix}.$$

Exterior Algebraic Shift

Definition Let $\mathbb{E} \supset \mathbb{F}$ be a field extension and let $g \in \mathrm{GL}(n, \mathbb{E})$.

- The k th compound matrix $g^{\wedge k}$ is the $\binom{n}{k} \times \binom{n}{k}$ -matrix of k -minors of g .
- $g^{\wedge S}$ is the $|S| \times \binom{n}{k}$ -submatrix of $g^{\wedge k}$ with rows indexed by S .

Let $\sigma_i \in S$, $|S| = m$ and $\tau_i \in \binom{[n]}{k}$ such that

$$g^{\wedge S} = \begin{pmatrix} \det g_{\sigma_1 \tau_1} & \det g_{\sigma_1 \tau_2} & \cdots & \det g_{\sigma_m \tau_1} \\ \vdots & \vdots & \vdots & \vdots \\ \det g_{\sigma_m \tau_1} & \det g_{\sigma_m \tau_2} & \cdots & \det g_{\sigma_m \tau_k} \end{pmatrix}.$$

The (exterior) algebraic shift is,

$$\Delta_g^{\mathbb{F}}(S) := \left\{ \sigma \in \binom{[n]}{k} : \sigma \text{ indexes a column with a pivot of } g^{\wedge S} \right\}.$$


Small Example

For $g = (x_{ij})_{1 \leq i, j \leq 4}$ $S = \{\{1, 2\}, \{2, 3\}, \}$ $\subset \binom{[4]}{2}$, $\mathbb{F} = \text{GF}(2)$.

$$g^{\wedge S} = \begin{pmatrix} x_{11}x_{22}+x_{12}x_{21} & x_{11}x_{23}+x_{13}x_{21} & x_{11}x_{24}+x_{14}x_{21} & x_{12}x_{23}+x_{22}x_{13} & x_{12}x_{24}+x_{14}x_{22} & x_{13}x_{24}+x_{14}x_{23} \\ x_{21}x_{32}+x_{22}x_{31} & x_{21}x_{33}+x_{23}x_{31} & x_{21}x_{34}+x_{24}x_{31} & x_{22}x_{33}+x_{23}x_{32} & x_{22}x_{34}+x_{24}x_{32} & x_{23}x_{34}+x_{24}x_{33} \end{pmatrix}.$$

Which row reduces to,

$$\begin{pmatrix} x_{11}x_{22}+x_{12}x_{21} & x_{11}x_{23}+x_{13}x_{21} & x_{11}x_{24}+x_{14}x_{21} & x_{12}x_{23}+x_{13}x_{22} & x_{12}x_{24}+x_{14}x_{22} & x_{13}x_{24}+x_{14}x_{23} \\ 0 & x_{11}x_{21}x_{22}x_{33} + x_{11}x_{21}x_{23}x_{32} + x_{12}x_{21}^2x_{33} + x_{13}x_{21}^2x_{32} + x_{12}x_{21}x_{22}x_{31} + x_{13}x_{21}x_{22}x_{31} & x_{11}x_{21}x_{22}x_{34} + x_{11}x_{21}x_{24}x_{32} + x_{12}x_{21}^2x_{34} + x_{14}x_{21}^2x_{32} + x_{12}x_{21}x_{22}x_{31} + x_{12}x_{21}x_{24}x_{31} & x_{11}x_{22}^2x_{33} + x_{11}x_{22}x_{23}x_{32} + x_{12}x_{21}x_{22}x_{33} + x_{13}x_{21}x_{22}x_{32} + x_{12}x_{22}^2x_{23}x_{31} + x_{13}x_{22}^2x_{23}x_{31} & x_{11}x_{22}^2x_{34} + x_{11}x_{22}x_{24}x_{32} + x_{12}x_{21}x_{22}x_{34} + x_{14}x_{21}x_{22}x_{32} + x_{12}x_{22}^2x_{24}x_{31} + x_{14}x_{22}^2x_{24}x_{31} & x_{11}x_{22}x_{23}x_{34} + x_{11}x_{22}x_{24}x_{33} + x_{12}x_{21}x_{23}x_{34} + x_{12}x_{21}x_{24}x_{33} + x_{13}x_{21}x_{24}x_{32} + x_{14}x_{21}x_{23}x_{32} + x_{13}x_{22}x_{24}x_{31} + x_{14}x_{22}x_{23}x_{31} \end{pmatrix}.$$



LU decomposition

Lemma $\Delta_{gb}(S) = \Delta_g(S)$ for any upper triangular b and any simplicial complex S .

LU decomposition For $g \in \text{GL}(n, \mathbb{E})$ with non zero leading principal minors, there exists ℓ lower triangular, u upper triangular such that

$$g = \ell u .$$

LU decomposition

Lemma $\Delta_{gb}(S) = \Delta_g(S)$ for any upper triangular b and any simplicial complex S .

LU decomposition For $g \in GL(n, \mathbb{E})$ with non zero leading principal minors, there exists ℓ lower triangular, u upper triangular such that

$$g = \ell u .$$

Example

$K = \{\{1, 2\}, \{2, 3\}, \{4\}\}$, and $\ell^{\wedge S}$ reduces to

$$\begin{pmatrix} x_{13}x_{24} + x_{14}x_{23} & x_{12}x_{24} + x_{14} & x_{24} & x_{12}x_{23} + x_{13} & x_{23} & 1 \\ 0 & x_{12}x_{23}x_{24}x_{34} + x_{12}x_{24}^2 & x_{23}x_{24}x_{34} & x_{12}x_{23}^2x_{34} + x_{12}x_{23}x_{24} & x_{23}^2x_{34} & x_{23}x_{34} \\ & + x_{13}x_{24}x_{34} + x_{14}x_{24} & + x_{24}^2 & + x_{13}x_{23}x_{34} + x_{14}x_{23} & + x_{23}x_{24} & + x_{24} \end{pmatrix},$$

Monte Carlo Algorithm

- Randomized algorithm.
- Current Standard for computing $\Delta^{\mathbb{F}}(S)$.
- Pick u by sampling from $\mathbb{F}$.
- Computing $\Delta_u^{\mathbb{F}}(S)$ is fast.
- Successful with high probability in characteristic 0.
- Not the case for small fields.

Verification Algorithm (Las Vegas)

Algorithm Verify u computes the algebraic shift of S

Input: $u \in \mathbb{E}^{n \times n}$, S a k -uniform hypergraph.

Output: true if $\Delta^{\mathbb{F}}(S) = \Delta_u^{\mathbb{F}}(S)$; otherwise false

- 1: $S' \leftarrow \Delta_u(S)$
 - 2: $\sigma_{\max} \leftarrow \max_{\text{lex}} S'$
 - 3: $T \leftarrow \{\tau \in \binom{[n]}{k} : \tau <_{\text{lex}} \sigma_{\max}\}$
 - 4: **if** $|S'| = |T| + 1$ **then**
 - 5: **return** true
 - 6: **end if**
 - 7: $m \leftarrow$ row echelon form of $\ell_{*T}^{\wedge S}$
 - 8: **if** columns of pivots of $m = S' \setminus \{\sigma_{\max}\}$ **then**
 - 9: **return** true
 - 10: **end if**
 - 11: **return** false
-

$$\ell^{\wedge S} = \begin{pmatrix} | & | & \dots & | & | & \dots \\ c_1 & c_2 & \dots & c_{|T|} & \sigma_{\max} & \dots \\ | & | & \dots & | & | & \dots \end{pmatrix}$$

Simplicial Complexes


Definition An abstract simplicial complex Σ , is a collection of non empty sets of $[n] := \{1, 2, \dots, n\}$, such that for any $\tau, \sigma \subset [n]$, if $\tau \subset \sigma \in K$ then $\tau \in \Sigma$.
change lower case k

- $\sigma \in \Sigma$ a k-simplex or a face, $\dim(\sigma) = k + 1$.
- f-vector number of k -faces for each k .

Simplicial Complexes

Definition An abstract simplicial complex Σ , is a collection of non empty sets of $[n] := \{1, 2, \dots, n\}$, such that for any $\tau, \sigma \subset [n]$, if $\tau \subset \sigma \in K$ then $\tau \in \Sigma$.
change lower case k

- $\sigma \in \Sigma$ a k -simplex or a face, $\dim(\sigma) = k + 1$.
- f -vector number of k -faces for each k .



The f -vector is $(4, 5, 1)$

Simplicial Homology

Definition Given a field $\mathbb{F}$, and a simplicial complex Σ we define the k th simplicial homology to be the vector space

$$H_k^{\mathbb{F}}(\Sigma) := k\text{-cycles} / k+1\text{-boundaries}.$$

The k th Betti number of Σ is the dimension of $H_k^{\mathbb{F}}(\Sigma)$.

Simplicial Homology

Definition Given a field $\mathbb{F}$, and a simplicial complex Σ we define the k th simplicial homology to be the vector space

$$H_k^{\mathbb{F}}(\Sigma) := k\text{-cycles} / k+1\text{-boundaries.}$$

The k th Betti number of Σ is the dimension of $H_k^{\mathbb{F}}(\Sigma)$.

The Betti numbers are $(1, 1, 0)$




Table of timings

						$\mathbb{Q}$		GF(2)		GF(4)		
n	o	g	i	#2-faces		time	#trials	time	#trials	time	#trials	...
7	x	1	2	12		0.1	1 *	6.5	14	6.9	6	...
7	✓	0	5	10		0.3	1	1.9	3	2.5	1	...
7	✓	1	1	14		2.7	1	578.7	2	554.0	1	...
8	x	1	4	14		2.4	1	-	>100 *	11.9	5	...
8	x	1	5	14		0.1	1 *	8.7	26	9.6	2	...
8	x	1	6	14		0.1	1 *	10.6	75	11.7	2	...
8	x	1	7	14		0.1	1 *	3.3	26	4.3	2	...
8	x	2	1	16		0.1	1 *	1338.8	25	1350.0	2	...
8	x	2	2	16		0.1	1 *	oot	oot	oot	oot	...
8	✓	0	1	12		1.1	1	4.6	84	5.5	2	...
8	✓	0	2	12		1.8	1	-	>100	8.1	2	...
8	✓	1	1	16		3.6	1	578.1	26	571.1	2	...
8	✓	1	3	16		27.0	1	oot	oot	oot	oot	...
8	✓	1	7	16		oom	oom	oot	oot	oot	oot	...
:	:	:	:	:		:	:	:	:	:	:	.

Table of timings

n	o	g	i	#2-faces	$\mathbb{Q}$			GF(2)		GF(4)		...
					time	#trials		time	#trials	time	#trials	
7	x	1	2	12	0.1	1	*	6.5	14	6.9	6	...
7	✓	0	5	10	0.3	1		1.9	3	2.5	1	...
7	✓	1	1	14	2.7	1		578.7	2	554.0	1	...
8	x	1	4	14	2.4	1		—	>100 *	11.9	5	...
8	x	1	5	14	0.1	1	*	8.7	26	9.6	2	...
8	x	1	6	14	0.1	1	*	10.6	75	11.7	2	...
8	x	1	7	14	0.1	1	*	3.3	26	4.3	2	...
8	x	2	1	16	0.1	1	*	1338.8	25	1350.0	2	...
8	x	2	2	16	0.1	1	*	oot	oot	oot	oot	...
8	✓	0	1	12	1.1	1		4.6	84	5.5	2	...
8	✓	0	2	12	1.8	1		—	>100	8.1	2	...
8	✓	1	1	16	3.6	1		578.1	26	571.1	2	...
8	✓	1	3	16	27.0	1		oot	oot	oot	oot	...
8	✓	1	7	16	oom	oom		oot	oot	oot	oot	...
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮

Thank you!

Table of sizes

[illegible]