
SIMULTANEOUSLY ESTIMATING FOOD WEB **COMPLEXITY** CONNECTANCE AND STRUCTURE WITH UNCERTAINTY

A PREPRINT

Anubhav Gupta *

Department of Evolutionary Biology and Environmental Studies

University of Zurich

8057 Zurich, Switzerland

`anubhav.gupta@ieu.uzh.ch`

Reinhard Furrer

Department of Mathematics and Department of Computational Science

University of Zurich

8057 Zurich, Switzerland

`reinhard.furrer@math.uzh.ch`

Owen L. Petchey

Department of Evolutionary Biology and Environmental Studies

University of Zurich

8057 Zurich, Switzerland

`owen.petchey@ieu.uzh.ch`

November 27, 2021

Abstract

- 1 3) We fit the ADBM using approximate Bayesian computation to 12 observed food webs
2 from a wide variety of ecosystems. Connectance was consistently overestimated in
3 the new parameterisation method. In some of the food webs, considerable variation
4 in estimated parameter distributions occurred, and resulted in considerable variation
5 (i.e. uncertainty) in predicted food web structure.

*Corresponding author

- 4) These results lend weight to the possibility that the observed food web data is missing some trophic links that do actually occur. It also seems likely that the ADBM likely predicts some links that do not exist. The latter could be addressed by accounting in the ADBM for additional traits other than body size. Further work could also address the significance of uncertainty in parameter estimates for predicted food web responses to environmental change.
- 5) Food web models explain and predict the trophic interactions in a food web, and they can infer missing interactions among the organisms. The allometric diet breadth model (ADBM) is a food web model based on the foraging theory. In the ADBM the foraging parameters are allometrically scaled to body sizes of predators and prey. In Petchey et al. (2008), the parameterisation of the ADBM had two limitations: (a) the model parameters were point estimates, and (b) food web connectance was not estimated.
- 6) The novelty of our current approach is: (a) we consider multiple predictions from the ADBM by parameterising it with approximate Bayesian computation, to estimate parameter distributions and not point estimates. (b) Connectance emerges from the parameterisation, by measuring model fit using the true skill statistic, which takes into account prediction of both the presences and absences of links.
- ~~7) We fit the ADBM using approximate Bayesian computation to 16 observed food webs from a wide variety of ecosystems. Connectance was consistently overestimated in the new parameterisation method. In some of the food webs, considerable variation in estimated parameter distributions occurred, and resulted in considerable variation (i.e. uncertainty) in predicted food web structure.~~
- ~~8) We conclude that the observed food web data is likely missing some trophic links that do actually occur, and that the ADBM likely predicts some links that do not exist. The latter could be addressed by accounting in the ADBM for additional traits other than body size. Further work could also address the significance of uncertainty in parameter estimates for predicted food web responses to environmental change.~~

Keywords connectance · ABC · ADBM · food web · true skill statistic · uncertainty

1 Introduction

Knowledge about the trophic interactions among the organisms in a community is crucial for understanding the structure and dynamics of ecological communities and for predicting their response to environmental change (Dunne, Williams, and Martinez 2002; Tylianakis and Binzer 2014; O'Connor et al. 2009; Bergamino, Lercari, and Defeo 2011; Krause et al. 2003; Lurgi, L'opez, and Montoya 2012; Morris, Sinclair, and Burwell

2015). The network of trophic interactions is often referred to as a food web. The food web structure can provide answers to key ecological questions: which species are more vulnerable to environmental changes such as temperature ([Owen L. Petchey et al. 1999](#)); how robust a food web is to extinctions (Dunne, Williams, and Martinez 2002); and how a food web reacts if the ~~top~~-predators are removed (Knight et al. 2005)?

Trophic interactions information from multiple sources can be used to infer a food web, e.g. gut contents (Peralta-Maraver, ~~L'opez-Rodríguez~~[L'opez-Rodríguez](#), and de Figueroa 2017), ~~stable isotope composition of tissues (Layman et al. 2007), and experimentation (Warren 1989). Sometimes the methods used to infer the interactions may lead to uncertainty in the constructed food web. E.g. In gut content analysis of some fish predators, there can be tissues which are not identifiable and [2016](#)) and cannot be assigned with certainty to a specific prey item (Baker, Buckland, and Sheaves 2014). With stable isotope ratios of tissues, uncertainty may be due to factors such as variability in the isotopic fractionation values across multiple combinations of diets and tissues/species, unquantified temporal or spatial variation in prey isotopic values, and variation caused by routing of particular dietary nutrients into particular tissues (Crawford, McDonald, and Bearhop 2008). Furthermore, complete recording of all interactions usually requires a large sampling effort even at small spatial and temporal scales (Hobson, Piatt, and Pitocchelli 1994). Food web structure is very difficult to record at larger spatial and temporal scales without losing resolution (spatial, temporal, and taxonomic) (Gravel et al. 2013; Martinez 1991; Jord'an and Osv'ath 2009). Less than complete sampling of interactions can result in no interaction being observed between a pair of individuals that in fact do interact, which results in missing links in a food web. Due to under-sampling, food webs can be poorly understood, which may hinder further advances in the field (Martinez et al. 1999).~~

When interactions are difficult to observe, and hence well-documented food webs are not available, models which predict species interactions may provide a solution (Tamaddoni-Nezhad et al. 2013; Gravel et al. 2013; [Owen L. Petchey et al. 2008](#); Allesina, Alonso, and Pascual ~~2008~~[2008a](#); Cohen, Newman, and Steele 1985). A food web model can be used to predict missing information about species interactions. For example, [Owen L. Petchey et al. \(2008\)](#) showed how a model of species interactions (and therefore food web structure) could be parameterised from data on the known presence and absence of trophic interactions. The model and its parameter values encode the rules for occurrence or absence of species interactions to predict food web structure. Observed data may be used to select and parameterise the model. Tamaddoni-Nezhad et al. (2013) used large agricultural datasets, logic-based machine learning and text mining to assign interactions between nodes to automatically construct food webs. Gravel et al. (2013), inspired by the niche model of food web structure developed a method that used the statistical relationship between predator and prey body size to infer the food web.

Food web models are also useful for ecological forecasting. Lindegren et al. (2010) used a stochastic food web model driven by regional climate scenarios to produce quantitative forecasts of cod dynamics in the twenty-first century. Hattab et al. (2016) forecasted the potential impacts of climate change on the local food

web structure of the highly threatened Gulf of Gabes ecosystem, located in the south of the Mediterranean Sea. Hence, food web models have an important role in filling gaps in knowledge about species interactions, including predicting future changes in food web structure.

The allometric diet breadth model (ADBM) was the first model able to predict food web ~~complexity~~ connectance (i.e. the number of trophic links) and structure (i.e. the arrangement of those links) (Beckerman, Petchey, and Warren 2006; Owen L. Petchey et al. 2008). ~~The ADBM Models such as the Cascade model require connectance as an input parameter whereas the ADBM does not (“A Stochastic Theory of Community Food Webs I. Models and Aggregated Data” 1985). It uses foraging theory, specifically the contingency model (MacArthur and Pianka 1966), to predict which set of the available prey species would be consumed by a predator. This set is the prey species that maximises the energy intake rate of the predator. The model requires the foraging related traits of species, such as energy content of a potential prey item, the rate of space clearance (also known as attack rate), the density of prey items, and handling time (the amount of time required to handle food items). The model is termed “allometric” because each of these quantities is derived from the body size of the prey and predator using several allometric relationships. The ADBM has also been used to investigate the effect of temperature on an observed food web structure (O’Gorman et al. 2019).~~

diet of each potential consumer and thereby the food web structure (further details are in the Material and Methods section). Connectance is the number of realised trophic links divided by the total number of possible trophic links. The ADBM had variable success in explaining the structure of 15 different food webs, with the proportion of links correctly predicted ranging from 5 % to 65 % (Table 1). The ADBM correctly predicted 65% of the presence of links in the Coachella valley food web. The poorest prediction of presence of links was for the Grasslands food web with only 7% of the presence of links correctly predicted. When trophic interactions were more strongly dependent on size, the model correctly predicted a greater proportion of links. Indeed, constructing a food web based only on body size (i.e. ignoring taxonomy) resulted in almost twice the number of correctly predicted links, i.e. 83%, in contrast to taxonomy (Woodward et al. 2010).

Although Owen L. Petchey et al. (2008) demonstrated that foraging theory could predict food web structure, their implementation of the ADBM included at least two limitations. The parameterisation method provided estimates of the parameters with no uncertainty: a single set of parameter values that maximised the explanatory power was selected. In other words, the parameterisation method led to point estimates of the parameters that predicted a single food web structure (because the ADBM is purely deterministic). Moreover, the best predicted food web was not exactly the same as the observed one. In a sense then, the parameterisation method used in Owen L. Petchey et al. (2008) was akin to estimating the intercept and slope of a regression line, but not any uncertainty in those parameters. Given that uncertainty is an essential dimension in ecological models, and in predictions about the future state of ecological communities (Owen L. Petchey et al. 2015; Carpenter 2016), this is an important limitation.

The second limitation was ~~in the estimation of the~~ that the connectance of the ~~food web, which is the~~
~~number of realised trophic links divided by the total number of possible trophic links~~ predicted food web was
~~not estimated~~. Although the ADBM can in principle ~~predict connectance, estimate connectance,~~ Owen L.
Petchey et al. (2008) prevented the model from doing so. They set the value of relevant parameters in the
model to instead ensure the predicted connectance was equal to the observed connectance. ~~The ADBM~~
~~was not therefore used to simultaneously predict complexity and structure of food webs.~~ Moreover, fixing
predicted connectance to be equal to observed connectance does not account for the possibility that the
observed connectance was imperfectly measured. Indeed, if low effort was used to observe the trophic links
in a community, the observed connectance are likely to be lower than if all trophic links were observed.
Connectance is an important driver for the stability and dynamics of a food web (May 1972) and most of the
structural properties of food webs co-vary with connectance (Dunne, Williams, and Martinez 2002; Poisot
and Gravel 2014), thus this limitation must be addressed.

In this article we report on how we address these limitations. We removed the first limitation by
applying approximate Bayesian computation (ABC). The approach originated in population genetics and
has been used in a wide range of research fields: systems biology (Toni et al. 2009), ecology (Jabot and
Chave 2009), epidemiology (Shriner et al. 2006) and ecological networks (Ibanez 2012; Poisot and Stouffer
2016). One of the advantages of ABC is that it does not require a likelihood function. As ADBM is a
complex deterministic model where the likelihood can not be explicitly evaluated, ABC is a good choice of
parameterisation.

We addressed the second limitation by allowing estimation of number of links as well as arrangement of
links. To accomplish this, we measured model fit by using the true skill statistic, which takes into account
both the number of presences and absences of links correctly predicted. High values of the true skill statistic
occurs when both the predicted arrangement of links and the predicted number of links are close to the
observed arrangement and number of links, respectively.

Table 1: Information about the food webs predicted using the ADBM.

Common food web name (Original Publication)	Predation matrix source	Body size source	General ecosystem	Number of species	Connectance	Body size range (approximate)	Proportion of presence of links correct	Type of interactions
Benguela Pelagic (Yodzis 1998)	Brose et al. (2008)	Brose et al. (2008)	Marine	30	0.21	10^{-8} to 10^6	0.54	Predation
Broadstone Stream (taxonomic aggregation) (Woodward and Hildrew 2001; Woodward et al. 2005)	Brose et al. (2008)	Brose et al. (2008)	Freshwater	29	0.19	10^{-6} to 10^{-2}	0.40	Predation
Broom (Mommott et al. 2000)	Brose et al. (2008)	Brose et al. (2008)	Terrestrial	60	0.03	10^{-6} to 10^0	0.09	Herbivory, Parasitism, Predation, Pathogenic
Capinteria (Lafferty et al. 2006)	Hechinger et al. (2011)	Hechinger et al. (2011)	Marine (Salt Marsh)	88	0.08	10^{-6} to 10^4	0.33	Predator-parasite, Parasite-parasite
Caricaie Lakes (Cattin et al. 2004)	Brose et al. (2008)	Brose et al. (2008)	Freshwater	158	0.05	10^{-5} to 10^5	0.13	Predation, Parasitism
Grasslands (Dawah et al. 1995)	Brose et al. (2008)	Brose et al. (2008)	Terrestrial	65	0.03	10^{-3} to 10^{-2}	0.07	Herbivory, Parasitism
Mill Stream (Ledger, Edwards, Woodward unpublished)	Brose et al. (2008)	Brose et al. (2008)	Freshwater	80	0.06	10^{-6} to 10^{-2}	0.36	Herbivory, Predation
Skipwith Pond (Warren 1989)	Brose et al. (2008)	Brose et al. (2008)	Freshwater	71	0.07	10^{-4} to 10^{-1}	0.14	Predation
Small Reef (Opitz 1996 Table 8.6.2)	Alyssa R. Cirtwill and Anna Eklöf (2018)	Alyssa R. Cirtwill and Anna Eklöf (2018)	Marine (Reef)	239	0.06	10^{-11} to 10^5	0.30	Predation, Herbivory
Tuesday Lake (Jonsson et al. 2005)	Brose et al. (2008)	Brose et al. (2008)	Freshwater	73	0.08	10^{-11} to 10^3	0.46	Predation
Ythan (Emmerson and Raffaelli 2004)	Alyssa R. Cirtwill and Anna Eklöf (2018)	Alyssa R. Cirtwill and Anna Eklöf (2018)	Marine (Estuarine)	85	0.04	10^{-6} to 10^4	0.17	Predation
Broadstone Stream (size aggregation) (Woodward et al. 2010)	Guy Woodward. (2021)	Guy Woodward. (2021)	Freshwater	29	0.24	10^{-7} to 10^2	0.83	Predation

2 Materials and Methods

In the upcoming sections, we present a detailed account of the application of ABC to parameterise the ADBM, the description of the ADBM and of the food web data we used. We explain the rejection Monte Carlo ABC method in the main text, and Markov chain Monte Carlo ABC and sequential Monte Carlo ABC methods in the Supplementary information (hereafter SI) Section S1 (hereafter SI-S1). We computed an accuracy measure known as true skill statistic to assess the ADBM's predictions and calculated different food web properties to compare these predictions across food webs.

2.1 Allometric Diet Breadth Model (ADBM)

The allometric diet breadth model (ADBM) is based on optimal foraging theory, specifically the contingency foraging model (MacArthur and Pianka 1966). The ADBM predicts the set of prey species a consumer should feed upon to maximise its rate of energy intake (Owen L. Petchey et al. 2008) (hereafter referred as PBRW study). The species in this set are assumed to have the trophic link with the predator. To make these predictions, the model assumes that a foraging predator is in one of two exclusive states: searching for prey or handling a prey item. The notation used below corresponds to that of (Owen L. Petchey et al. 2008). The model requires four variables for each potential predator-prey interaction:

- The energy content of the resources E_i (only prey i specific) (energy).
- The handling times H_{ij} , which is the time not spent searching caused by consuming a prey item (prey i and predator j specific) (time).
- The space clearance rates A_{ij} (also known as the attack rate; prey i and predator j specific) (area or volume per time).
- The prey densities N_i (only prey i specific) (individuals per area or volume).

The term “Allometric” in the ADBM refers to the use of four allometric relationships, one for each of these four variables, including the body size of the predator M_j and prey M_i (Table 2). With these four allometric relationships, and the body size of each of the species in a community,

Table 2: Traits with their allometric function and corresponding parameters in ADBM.

Traits (Unit)	Allometric function	Parameters	Comments
Energy (<i>Joules</i>)	$E_i = eM_i$	e	Arbitrary. No effect on structure*.
Abundance (individual/ m^2 or individual/ m^3)	$N_i = nM_i^{a_i}$	n	Connectance affected by the product nah^* .
		n_i	Assumed value of $\frac{-3}{4}$ based on empirical data
Space Clearance Rate (m^2/s or m^3/s)	$A_{ij} = aM_i^{a_i}M_j^{a_j}$	a	Connectance affected by the product nah^* .
			Estimated using ABC
		a_i	Estimated using ABC
		a_j	Estimated using ABC
Handling time (s)	$H_{ij} = \frac{h}{b - \frac{M_i}{M_j}}$ if $\frac{M_i}{M_j} < b$	h	Connectance affected by the product nah^* .
	$H_{ij} = \infty$ if $\frac{M_i}{M_j} \geq b$	b	Estimated using ABC

* See SI S4 for further explanation for why only four parameters were estimated.

we can predict the four variables (energy, handling time, space clearance rate, and prey density), and then use the contingency foraging model to predict diets.

Each of the four allometric equations has parameters: a constant and/or at least one exponent (Table 2). It is the value of some of these parameters that can be estimated to have the predicted food web structure match (as closely as possible) the structure of an observed food web. This is akin to choosing values of slope and intercept of a linear regression that maximises the fit of the regression line to the observed data.

Please note that in the notation of the allometric exponents (a_i, a_j), the i and j refer to the exponent for the prey mass and predator mass respectively, i.e. the i and j refers to the role and not species specific.

Because some of the allometric constants and exponents are known, and because others are redundant with respect to each other (see Table 2 for details), we estimate only the following parameters: a, a_i, a_j and b in the model (Table 2).

In the ADBM, some species can be predicted to eat others, but to not be eaten by others, i.e. be predicted to be a top predator. This can occur for relatively large species when the exponent b is less than 1, which can cause the handling time of this large species to be infinite for all potential consumer species.

2.2 Observed food web data

The observed food webs that we fit the ADBM to belong to marine, freshwater and terrestrial ecosystems (Table 1). The observed connectance of these food webs is from 0.02 to 0.34 and there are

179 19 to 158 species. The food webs contain primary producers, herbivores, carnivores, parasites, and
 180 parasitoids. They also contain various types of feeding interactions, including predation, herbivory,
 181 bacterivory, parasitism, pathogenic, and parasitoid.

182 The goodness of fit of the ADBM’s predictions depends on the types of interactions in the
 183 food webs in the PBRW study. Because some of the interactions are more size structured than other
 184 interactions. Predacious and aquatic herbivore interactions were predicted better than parasitoid
 185 and herbivory ones (PBRW study). ~~Moreover, the ADBM’s predictions improve when the species~~
 186 ~~(taxonomic) are replaced by size classes (ignoring taxonomy) (Woodward et al. 2010).~~

187 All food webs with one exception (Broadstone Stream) was available only at the species level,
 188 i.e. with information about interactions between species and the body size of species. We use the
 189 term “species” in this study to indicate a “node” in a food web in which nodes are connected by
 190 trophic interactions, and nodes are a collection of individuals that share links. These species/nodes
 191 are not always taxonomic species, but can be broader taxonomic ranks.

192 In contrast, the Broadstone Stream food web data contained interactions between individuals
 193 and the individual body sizes. Thus, the Broadstone Stream food web can be constructed by
 194 aggregating by either taxonomy or size (Woodward et al. 2010). ~~The ADBM can predict the~~
 195 ~~food web irrespective of the aggregation method, and aggregation by size, ignoring taxonomy led~~
 196 ~~to higher explanatory power i.e. the match between observed and predicted food web structure was~~
 197 ~~higher. With aggregation by size, 83% of the links were correctly predicted than with taxonomic~~
 198 ~~aggregation where 40% of links were correctly predicted (Woodward et al. 2010).~~

199 2.3 Parameter estimation: Approximate Bayesian Computation

200 We used approximate Bayesian computation (ABC) to identify sets of parameter values that resulted
 201 in predicted food webs that were close in structure to the observed food web. ABC is an approach
 202 that does not require a likelihood function. Instead, there is a distance function that measures the
 203 distance between a model’s prediction and the observed data. The approximation of the likelihood
 204 depends on the ABC method used, which is further discussed below and SI. The model parameter
 205 values are sampled from a prior distribution. The accepted parameter values form an approximate
 206 posterior distribution for the model parameter. We implemented three ABC methods to parameterise

the ADBM: namely rejection Monte Carlo (Fig. 1), Markov chain Monte Carlo, and sequential Monte Carlo. The three methods produced very similar results (SI Figs S33-S34) and we therefore only include the simplest (rejection) in this main text.

2.3.1 Prior distribution

The prior distributions for a_i and a_j were chosen to be uniform distributions. The range of distribution was from -1.5 to 1.5 and 0 to 3 for a_i and a_j respectively, informed by the estimates in Rall et al. (2012). However, we chose a prior range specific to food webs for the parameter b because body size varies greatly among the species in the observed food webs. For example: in the Benguela Pelagic food web, the body sizes of species range from the order of 10^{-8} gm to 10^5 gm. Hence, the range of prey-predator ratio was from the order of 10^{-14} to 10^{14} . To take this into account, we took the prior of $\log_{10}(b)$ from a uniform distribution ranging from -15 to 15. In the case of parameter a , we chose the prior of $\log_{10}(a)$ to be a uniform distribution. However, the prior range varied between food webs. For example, the prior range for Benguela Pelagic was chosen to be -12 to 10. ~~The method for choosing the specific range of the prior distribution of the~~

~~parameter is detailed in SI-S4.~~ For the prior of $\log_{10}(a)$ to be a uniform distribution. Since, the ADBM estimated connectance to be higher than the real connectance, lower values of a were favoured in the parameterisation. Hence, the upper bound of the prior was set to 10. To set the lower bound, we investigated how the true skill statistic varied with $\log_{10}(a)$ (e.g. SI Fig. S35). We found that the TSS increased with decreasing $\log_{10}(a)$ and then remained constant, for a constant value of $\log_{10}(b)$. We therefore decided to set the lower bound of $\log_{10}(a)$ such that the maximum variation of TSS was taken into account, while attempting to keep the range of prior as small as possible. In the case of Benguela Pelagic as shown in SI Fig. S35, the lower bound of $\log_{10}(a)$ was taken to be -12.

2.3.2 Comparison of observed and predicted

The difference between the model's prediction and the observed data (e.g. the sum of squared residuals is such a distance in linear regression) is quantified by a distance measure. The distance is lower when there is a closer match between the model's prediction and the observation. A perfect match would result in zero distance.




Figure 1: Flowchart of rejection approximate Bayesian computation method implemented to parameterise the ADBM.

235 The magnitude of the distance is used for the acceptance or rejection of a set of parameter
 236 values. An accepted set of parameter values contributes to the posterior distribution, rejected ones
 237 do not. This makes the distance measure one of the important features of ABC. A threshold distance
 238 is chosen, and if the distance for a particular set of parameter values is less than the threshold, then
 239 that set of parameter values contributes to the posterior distribution. When the distance is greater
 240 than the threshold, the parameter values do not contribute to the posterior. Hence, the magnitude
 241 of the distance threshold determines the proportion of a model's parameters that are accepted. A
 242 higher threshold causes a high proportion of acceptances but less accuracy with the acceptance
 243 of some parameter sets that result in predictions quite unlike the observed data. Below, we first
 244 describe and justify our choice of distance measure, and then our choice of threshold.

245 **Choice of distance measure** In the PBRW study the measure of distance was equivalent to
 246 ~~$1 - a/(a + c)$~~ , where ~~a~~ $1 - TP/(TP + FN)$, where TP is the number of observed links that were
 247 predicted (the number of true positives) and ~~c~~ FN is the number of observed links that were not
 248 predicted (the number of false negatives). A distance of 0 indicates that all observed links were
 249 correctly predicted. One way for the ADBM to achieve this is to predict that every species has a
 250 trophic link with every other species including itself – a fully connected food web with connectance
 251 of 1. The PBRW study prevented this by constraining the number of predicted links to be equal to
 252 the number of observed links, i.e. the model connectance was fixed to be the same as the observed
 253 connectance. In this study, we relaxed this constraint, with the number of links as well as the
 254 arrangement of links being estimated. The first step was to choose an appropriate distance measure.

255 The distance measure used in this study is 1 minus the true skill statistic: distance = $1 - \text{TSS}$.
 256 This distance ranges from 0 to 2.

257 TSS is defined as:

$$\text{TSS} = \frac{ad - bc}{(a + c)(b + d)} \frac{TP \cdot TN - FP \cdot FN}{(TP + FN)(FP + TN)}$$

258 where ~~a~~ TP is the number of observed links that are predicted by the model (true positives), ~~d~~ TN
 259 is the number of observed absences of links that are correctly predicted (true negatives), ~~b~~ FP is
 260 the number of false positives, and ~~c~~ FN is the number of false negatives.

261 The TSS ranges from -1 to 1 , where $+1$ indicates a perfect prediction. A TSS value of zero
 262 or less indicates a performance no better than random.

263 The inclusion of true and false negatives in the distance measure means that the best theoreti-
 264 cally possible prediction (smallest distance) is a unique prediction, and specifically the one in which
 265 the predicted presence and absence of links matches exactly with the observed presence and absence
 266 of links.

267 **Choice of threshold value of distance** Food web dynamics and stability are strongly dependent
 268 on connectance (May 1972), we therefore set the distance threshold (for acceptance) such that
 269 the model had a reasonable chance of predicting the observed value of connectance. Note that in
 270 the following section (*The Rejection ABC method*) we use the term tol to denote the value of the
 271 distance threshold.

272 To do this, we examined how the predicted connectance varied with the distance threshold.
 273 An example of this relationship is given in Fig. 2 for the Benguela Pelagic food web ~~and in SI~~
 274 Fig. S1 for other food webs. We chose the minimum threshold value that gave a range of predicted
 275 connectance containing the observed connectance.

276 Furthermore, it is useful to note that in Fig. 2 there are no connectance values below a distance
 277 threshold value of less than 0.5 because for this particular food web there were no sets of parameter
 278 values that achieved a better model fit than is indicated by $1 - TSS = 0.5$. I.e. it is impossible for
 279 the ADBM to make better predictions than this. One reason for this is that the ADBM, when body
 280 size is the only trait, can only predict contiguous diets in trait space, whereas the observed data
 281 contains gaps in the diet.

282 2.3.3 The Rejection ABC method

283 In the rejection ABC method, a set of parameter values are sampled from the prior distributions.
 284 This set of parameter values is either accepted, and thereby added to the posterior distribution of the
 285 parameter values, or it is rejected (based on if the distance $1 - TSS$ is less than or greater than the
 286 threshold distance, as mentioned above). This process is repeated until there are enough acceptances
 287 to give stable (approximate) posterior distributions. ~~In addition, we used a kernel function that~~
 288 ~~assigns weight to each~~ Acceptance or rejection of a set of parameter values is probabilistic and




Figure 2: The prediction interval of the predicted connectance increases with increasing distance threshold for the Benguela Pelagic food web. The green line and black line represent the observed connectance and mean of predicted connectance respectively.

depends on a weight assigned to that set of parameter values. This weight is given by a kernel function, where the weight is inversely proportional to the distance (1 - TSS). This weight is then used in a way that makes it the probability of the set of parameter values being accepted.

In the upcoming section, we further detail the rejection ABC method.

Properties:

- A prior distribution $\pi(\theta)$: π is the uniform distribution for parameters $\theta = (a, a_i, a_j, b)$
- A model prediction $model(\theta)$: $ADBM(\theta)$. This is a predicted food web, x_i , given by a particular set of parameter values θ_i . Hence, $x_i = ADBM(\theta_i)$
- A summary statistic $s(x)$: x is the predation matrix predicted by the ADBM.

A kernel function $K(u)$: epanechnikov $K(u) = \frac{3}{4}(1 - \frac{u}{tol})$ if $u \leq tol$ A kernel function $K(u)$: epanechnikov $K(u) = 0$ otherwise

where tol is the distance threshold

300 • A distance function $d(x_i, y)$: $d(x_i, y) = 1 - TSS(x_i, y)$

301 • An observed food web y , in the form of a predation matrix containing zeros and ones.

302 *Sampling:*

303 for $i = 1 \dots n = 1000$

304 • Draw a set of parameter values θ_i from the prior distribution $\pi(\theta)$.

305 • Compute the model result $x_i = model(\theta_i)$

306 • Compute $s(x_i)$ and $d(s(x_i), s(y))$

307 • Accept or reject the parameter set probabilistically:

308 – Assign a ~~probability-weight~~ p_i to θ_i as per the kernel K ; ~~$p = K(d)$~~ $p_i = \frac{K(d)}{K(a)}$, where d

309 is the distance evaluated in the previous step.

310 – Compute $\alpha \sim U(0, 1)$

311 – If ~~$p_i \leq \alpha$~~ $p_i \leq \alpha$, then accept θ_i and $i = i + 1$

312 *Output:*

313 An approximate joint posterior distribution using the accepted $\theta_1, \dots, \theta_n$.

314 2.4 Assessment of model fit

315 Accuracy is how close the model prediction is to the observation. The ADBM's prediction is a

316 predation matrix that consists of the presence and absence of links thus comparing how close the

317 prediction is to the observation is not straightforward as comparing two numerical values. We

318 defined the accuracy of the ADBM using true skill statistics to take into account the true and false

319 predictions of both the presence and absence of links, which is defined above.

320 We examined how closely structural properties of the predicted food web matched those of the

321 observed food webs ~~using the R *cheddar* package (Hudson et al. 2013).~~ We evaluated properties

322 such as proportion of basal species, proportion of intermediate species, proportion of top species,

323 proportion of herbivores, mean omnivory, clustering coefficient, standard deviation of generality,

324 standard deviation of vulnerability, diet similarity, mean path length and nestedness. We ~~did~~ could

325 not compute mean trophic level and maximum trophic level because ~~their computation did not~~

326 ~~converge in the R~~ the networks contained too many paths for the *cheddar* package ~~(Hudson et al.~~
327 ~~2013) for all the food webs.~~ algorithm to compute.

328 We investigated the performance of the ADBM parameterised with the ABC by computing
329 standardised error of the food web properties, where the standardised error is the absolute raw error
330 (the difference between observed and predicted value) divided by the maximum absolute raw error
331 for that property. We did not calculate the standardised error for mean omnivory and mean path
332 length because it had some NA values and infinite values for all the food webs respectively.

333 3 Results

334 As an example of the model outcomes, we first present the results for the Benguela food web
335 (e.g. predicted food web structure, variation in predicted food web structure, and posterior parameter
336 distributions). We chose this food web as it was well explained using the method of Petchey et
337 al (2008) (hereafter referred as PBRW study). The results of the other food webs are included in
338 the SI Figs S1-S32. We then compare model outcomes across all empirical food webs between the
339 PBRW study and our current work. We compare the true skill statistic of the two approaches and
340 compare some food web properties, such as proportions of basal, intermediate, and top species.

341 The true skill statistic (TSS) of the predicted Benguela Pelagic food web varied between 0.4
342 and 0.52. This variation in the TSS is represented in terms of predation matrices displayed in Fig.
343 3(a), which overlays 1000 independent predation matrices ~~created from the posterior parameter~~
344 ~~distributions~~ accepted from the ABC method. In all the 1000 independent predation matrices, the
345 predicted links are mostly present in the upper triangular portion of the matrix where most of the
346 observed links are also present. Links in the upper right triangle of the predation matrix are for
347 predators feeding on prey smaller than themselves.

348 In the 1000 predicted predation matrices, there predators are sometimes smaller than their
349 predicted prey, the links in the lower left triangle of the predation matrix. This is also portrayed
350 in the marginal distribution of $\log_{10}(b)$ in Fig. 4(d), as it includes values greater than $b = 2$
351 ($\log_{10}(b) = 0.3$). This is relevant as values of $b = 2$ make the most profitable prey item equal in size
352 to the predator size. Lower values of b make the most profitable prey item smaller than the size of
353 the predator.

354 There were around 250 potential links in the lower left triangle of the predation matrix that
 355 were never predicted in any of the 1000 predicted predation matrix (Fig. 3(b)). This strongly
 356 suggests that the predator-prey size ratio of these links is so small (i.e. very large prey, very small
 357 predator) that the links cannot occur, given that the preponderance of observed links are predators
 358 consuming prey smaller than themselves.

359 The marginal posterior of parameter b in the Benguela Pelagic food web was more constrained
 360 than the marginal posterior distribution of the other three allometric parameters (Fig. 4) as the
 361 posterior range was the narrowest.

362 The mean true skill statistic using the ABC approach was higher than the point estimates
 363 from the PBRW study (Fig. 5(a)) across all food webs except one. Our present approach led to
 364 estimates of connectance greater than the values of connectance of the PBRW study, which were
 365 fixed to equal the observed values of connectance.

366 We did not find a consistent relationship between the parameters estimated using the current
 367 approach and those estimated in the PBRW study (Fig. 5(c-f)), except for in the case of parameter
 368 b . The mean using the ABC approach was always higher than the estimates from the PBRW study
 369 (Fig. 5(f)) and the 95% credible interval of the posterior of b includes the estimate from the PBRW
 370 study.

371 The marginal posterior of parameter b was more constrained than the other three allometric
 372 parameters, i.e. the posterior range was the narrowest (SI Figs S17-S32). In most of the food webs,
 373 the parameter b had a unimodal distribution (SI Figs S17-S32). EcoWEB60 and Grasslands had a
 374 bimodal distribution and Sierra Lakes had three modes.

375 The structural food web properties proportion of intermediate species, mean omnivory, cluster-
 376 ing coefficient, sd of generality, sd of vulnerability, diet similarity and nestedness estimated from
 377 the current ABC approach were generally higher than the PBRW study (~~SI Fig. S36~~ Fig. 6(b, e-j)).
 378 The properties proportion of basal species, proportion of top species, and proportion of herbivores
 379 were generally lower (~~SI Fig. S36~~ Fig. 6(a, c, d)).

380 The real values of the proportion of intermediate species, mean omnivory, clustering coefficient,
 381 sd of generality, sd of vulnerability and nestedness, were mostly within the lower range of the




Figure 3: (a) Observed and predicted predation matrices for Benguela Pelagic food web. Body size increases from left to right and top to bottom along the predation matrix. Black circles show where there is an observed trophic link. The intensity of the pink circles shows the proportion of 1000 predicted food webs that had a trophic link between the corresponding species. This type of overlay is shown for two examples predicted in panel (c). (b) The histogram of the number of times a link was predicted across 1000 independently predicted food webs. There were 841 species pairs in this food web. About 150 of these were predicted to have a trophic link in all 1000 predicted predation matrices. The red bar shows the number of pairs of species for which a trophic link was never predicted. (c) Two predicted predation matrices for Benguela Pelagic food web corresponding to the minimum and the maximum value of estimated b , and their sum.




Figure 4: Marginal prior and marginal posterior distribution of the ADBM parameters for the Benguela Pelagic food web estimated using rejection ABC. The black vertical line in (d) corresponds to the value of $b (=2)$ above which the most profitable prey item is larger in respect to the predator size.

382 predicted 95% interval. The proportion of basal species, proportion of top species, proportion of
 383 herbivores were underestimated in comparison to the real values for most of the food webs.

384 The ADBM, when parameterised with the ABC, generally better predicted the structural food
 385 web properties, such as proportion of basal species when the true skill statistics was higher (Fig.
 386 7(a)) across the ~~16~~ 12 food webs. However, the ABC parameterised ADBM less accurately predicted
 387 food web properties on average than in the PBRW study (Fig. 7(b)).

388 Within each food web, we found various relationships between the standardised error and true
 389 skill statistic (SI Figs S37 and S38). E.g. For Skipwith Pond food web (SI Fig. S37(1)), high values
 390 of TSS were associated with high error, whereas the opposite was true for other food webs, such as




Figure 5: TSS (a), connectance (b) and ADBM parameters (c, d, e, f) computed using the ABC method compared with the corresponding point estimates from Petchev et al (2008). The red lines are the 95% credible/prediction intervals and the black filled circles represent the corresponding means. The grey region represents the intervals of the prior distributions for a_i and a_j . The grey lines represent the prior range of the parameters a and b in the \log_{10} scale. The prior range for the parameter b extends above and below the y-axis limits for some food webs and so the values of the limits are shown on the plot. The dashed black lines are the 1:1 relationships for reference.




Figure 6: Structural properties of predicted food webs with 95% prediction interval parameterised using the ABC method plotted against the point estimates from Petchey et al. (2008). The black filled circles correspond to the mean, and green filled circles correspond to the properties of the observed food webs. The dashed black lines are the 1:1 relationships for reference.




Figure 7: (a) The mean standardised error of the food web properties predicted from the ADBM parameterised using rejection ABC plotted against the mean true skill statistic for each food webs. The vertical and horizontal bars correspond to 95% prediction intervals of the standardised error and true skill statistic respectively. Solid blue line is linear regression through the means ($t = -2.44$, 2.335 , $df = 14$, $P = 0.028$, 0.041). (b) The mean standardised error computed from the ABC method plotted against the mean standardised error from Petchey et al. (2008). The dashed line is the 1:1 relationship for reference.

Broadstone Stream (SI Fig. S37(b, p)). The other food webs showed more complex relationships. The shapes of the point scatters in SI Figs S37-38 are caused by the structural constraints of the ADBM (e.g. it can only predict contiguous diets) interacting with the TSS and the connectance of the observed food web. These features prevent, for example, a predicted food web with very high connectance having high TSS. Similarly, a predicted food web with connectance equal to that of the highest TSS prediction tends to have low TSS.

4 Discussion

The ABC parameterisation method employed here improves on the basic parameterisation methods applied in Owen L. Petchey et al. (2008) (PBRW). The ABC method provides uncertainty in parameter estimates, and thereby a range of predicted food webs (Fig. 5(c-f)). It also allowed us to

401 estimate parameters that were fixed by the PBRW study, and thereby also predicts connectance
 402 (Fig. 5(b)). Including uncertainty and predicting connectance are significant advances in ADBM.
 403 They allow predictions in changes of food web structure caused by environmental changes that
 404 include uncertainty in the predicted food web structure and including uncertainty in such predictions
 405 is critical ([Owen L. Petchey et al. 2015](#); [Cressie et al. 2009](#); [Lindegren et al. 2010](#)). A future
 406 development will be to partition the contribution of different sources of uncertainty such as incomplete
 407 sampling and model deficiencies to make improvements in the model with the aim of reducing
 408 uncertainty. Future research should investigate the functional and dynamical significance of the
 409 uncertainty in the predicted food web structure. Below we discuss some of the results of our study,
 410 and expand on these opportunities and priorities for future research.

411 In all cases, the predicted connectance was greater than the observed connectance (Fig. 5(b)).
 412 Why did this occur? Firstly, it is important to recognise that the ADBM (when using only body
 413 size as a trait) can only predict diets that are contiguous with respect to the size of prey. I.e. it
 414 cannot predict that a predator will consume prey of size 1 and 3, and not prey of size 2. Such
 415 patterns can however be predicted if a trait ~~other than~~ in addition to size and which is not perfectly
 416 correlated with size, influences foraging parameters ([Owen L. Petchey et al. 2008](#); [Allesina, Alonso,](#)
 417 [and Pascual 2008b](#); [Stouffer, Camacho, and Amaral 2006](#)). Secondly, it is important to note that
 418 the observed diets were not contiguous when prey are ordered by their size. The estimation process
 419 will result in a greater number of predicted links than observed given these features, and the model
 420 attempts to maximise the coincidence of predicted and observed link presence and absence (i.e. the
 421 true skill statistic).

422 These findings raise the question as to whether the model or the observed data is incorrect. We
 423 expect that ~~the observed data does not contain some links that would occur in reality~~ some of the
 424 links that do in reality occur are not present in the empirical datasets. This could be caused by low
 425 empirical sampling effort or rare prey-predator interactions even when sampling is extensive. This
 426 ~~can be possible due to low sampling effort causing some links that do occur to be not observed.~~ In
 427 this case, ~~the model may correctly predict a link that was not yet observed as the data was incorrect~~
 428 a false positive may actually be a correctly predicted link. More intensive and more complete sampling
 429 of links in food webs has been recognised as important, due to the potential that a low sampling
 430 effort will influence the perceived food web structure ([Martinez et al. 1999](#)).

431 We expect there are ~~eases~~ also cases of real false positives, where the model ~~incorrectly~~ predicts
 432 a feeding link despite no possibility that such a link ~~would~~ could occur in reality. This may be the
 433 case when a trait other than, or in addition to, prey size is influential. For example, a particular
 434 prey species may have a defensive trait that means it takes longer to consume it than an undefended
 435 prey of the same size. Incorporating traits other than body size in the ADBM would allow for
 436 discontinuous diets along the size axis. It is also possible that better estimates of parameters could
 437 cause lower estimated values of connectance. Furthermore, the ADBM's current form is a biology-
 438 only model; it does not include an observation process, although this could be included. The model
 439 would then be able to predict the absence of a link due to incomplete observations.

440 It would be interesting to take a very well sampled food web ~~;~~ ~~and~~ (real or simulated) and
 441 remove links at random to create a less well sampled version, and to test if the ~~ABC parameter~~
 442 ~~estimation applied to a subset of the observed links in a simulated poorly sampled food web predicts~~
 443 ~~the connectance of a well sampled food web. Such an outcome would indicate the~~ very well sampled
 444 version can be predicted from the less well sampled version (with ABC parameter estimation). If
 445 it could, then there is potential to compensate for under-sampling with an appropriate food web
 446 model and estimation procedure.

447 The ABC parameterisation resulted in a lower prediction accuracy of structural features of the
 448 food webs (Fig. 7 (b)) due to the overestimation of connectance. This was confirmed by principal
 449 component analysis of variation in the food web structural properties which revealed a first PC axis
 450 representing on average 62% of the overall variance, and this first axis was highly correlated with
 451 connectance, with an average Spearman correlation of 0.87 ~~;~~ (see SI S7 for details). Furthermore,
 452 there was a strong positive relationship between the mean standardised error in structural properties
 453 of the food webs and mean standardised error in connectance of the food webs (SI Fig. S33).

454 Our parameterisation approach was to maximise the true skill statistic (the coincidence of
 455 predicted and observed link presences, and the coincidence of predicted and observed link absences).
 456 The TSS assigns equal importance to the collection of presence and absence of observed links with
 457 the weight of an observed single presence or absence link being dependent on the connectance of the
 458 food web. If the connectance is less than 0.5, the TSS assigns more weight to a presence of link
 459 than to an absence of a link and vice versa.

Because the connectance of the observed food webs is less than 0.5 (Table 4), the TSS implicitly assigned more weight to a single-presence of link than to a single-absence of link. This result is as expected, as the chance that a recorded link is a correct is likely to be greater than the chance that a recorded absence is correct. This is because upweighting of link presences seems appropriate since observing a feeding interaction is unambiguous, whereas not observing one may be caused by various processes. That is to say, the observation of a single feeding interaction is sufficient to record the presence of a link. However, whereas this is not true for the absence of links: one observation of a predator not consuming a prey does not mean that it will never do so. Nevertheless, if we observe no interaction between two species during the sampling period, we conclude that there is an absence of link.

To improve our estimation procedure we could quantify the uncertainty in the recorded absence of links and include this uncertainty in the parameterisation method. Weight/importance could be assigned to true positives, true negatives, false positives and false negatives calculated from empirical studies which may be specific to that food web. Alternatively, an observation process could be added to the model, such that the biological part of the model can predict that a feeding link is possible, but then the observation process in the model leads to that link not being predicted.

In the PBRW study, the parameter b played a major role in maintaining the maximum predictive power of the ADBM. Indeed, they found that estimating b only, and not estimating either a_i or a_j slightly decreased model performance, and that estimating only b and a_j did not decrease model performance relative to when all three parameters were estimated.

We found that the posterior distribution of the parameter b was the most constrained of all the parameters (Fig. 4). Parameter b defines the range of prey body size which has a finite handling time, and the prey size with the highest energetic profitability. As the parameter b relates to the prey-predator body size ratio, the constrained posterior of b (Fig. 4(d)) indicates the importance of the ratio of body size of prey and predator in determining the food web structure with the ADBM.

The marginal posterior of parameter a was right-skewed (Fig. 4(c)). This may be because the ABC parameterisation overestimates the connectance, which means that lower values of a are preferred over higher values of a (a lower value of a leads to a lower space clearance/attack rate, and a lower space clearance rate results in a higher connectance).

Information about who eats ~~who~~ whom can be collected from multiple sources, such as gut contents of organisms, stable isotope composition of tissues, and experimentation (Peralta-Maraver, L'opez-Rodríguez, and de Figueroa 2017; Layman et al. 2007; Warren 1989). Moreover, experimentation provides independent estimates of allometric foraging parameters, such as b , a_i , and a_j (Rall et al. 2012). Diverse data could be used to parameterise the ADBM's predictions to test how uncertainty in the different datasets influences the ADBM's predictions using ABC. Appropriate summary statistics in the ABC method could be used to address such challenges. We could use, as an example, the approximate trophic position inferred from stable isotope ratio data from an individual tissue and gut content data of a predator simultaneously to parameterise the ADBM. The trophic position and the gut content information would be the summary statistics in this example. A further question that could be addressed in future studies is how the quantity of data affects the ADBM's predictions. The outcome of such a study could help food web researchers decide on how much data from a specific source is needed to predict the food web structure, and help further optimise the deployment of limited sampling resources.

When only partial food web data is available (Patonai and Jord'an 2017), the summary statistics in ABC can be used to infer these food web structures from the ADBM. It would be possible to use gut content data of only some of the species in a food web to parameterise the ADBM and predict the food web structure. Summary statistics opens up a broad spectrum of possibilities in parameterising food web models. There are multiple empirical and theoretical studies on a range of different food web properties of food webs across different ecosystems (Williams and Martinez 2000; Goldwasser and Roughgarden 1993; Martinez 1991). These can conceivably be used in parameterising food web models using ABC to constrain the model predictions.

5 Acknowledgements

This work was supported by the University Research Priority Program Global Change and Biodiversity (Grant number: U-704-04-11) of the University of Zurich. We thank the Petchey group members for their valuable suggestions in the manuscript. We thank Debra Zuppingner-Dingley for proofreading the manuscript.

6 Author contributions

Anubhav Gupta: Conceptualization (equal), [data curation \(lead\)](#), formal analysis (lead), [investigation \(lead\)](#), methodology (lead), [project administration \(lead\)](#), software (lead), [validation \(lead\)](#), [visualisation \(lead\)](#), writing – original draft preparation (lead), writing - review and editing (equal). Reinhard Furrer: Methodology (supporting). Owen L. Petchey: Conceptualization (equal), funding acquisition (lead), methodology (supporting), resources (lead), [supervision \(lead\)](#), writing – original draft preparation (supporting), writing - review and editing (equal).

7 Data Accessibility Statement

All the data used in this study was collected in other studies [and is openly available](#). We list those studies [and the open access source](#) in Table 1. ~~Some of those studies provide open access to their data and others do not. We have permission to use the data but do not have permission to distribute it.~~ The code used for data curation is available in the repository https://github.com/anubhav3/C1_data. The complete code used in the analysis is available in the repository https://github.com/anubhav3/C1_method_v2.

References

- [“A Stochastic Theory of Community Food Webs I. Models and Aggregated Data.” 1985. *Proceedings of the Royal Society of London. Series B. Biological Sciences* 224 \(1237\): 421–48. <https://doi.org/10.1098/rspb.1985.0042>.](#)
- Allesina, Stefano, David Alonso, and Mercedes Pascual. 2008. ~~“A General Model for~~ 2008a. [“A General Model for Food Web Structure.”](#) *Science* 320 (5876): 658–61. <https://doi.org/10.1126/science.1156269>. ~~American Association for the Advancement of Science~~
- . 2008b. [“A General Model for Food Web Structure.”](#) *Science* 320 (5876): 658–61. <https://doi.org/10.1126/science.1156269>.
- Baker, Ronald, Amanda Buckland, and Marcus Sheaves. 2014. “Fish Gut Content Analysis: Robust Measures of Diet Composition.” *Fish and Fisheries* 15 (1): 170–77. <https://doi.org/10.1111/faf.12026>.

- Beckerman, A. P., O. L. Petchey, and P. H. Warren. 2006. “Foraging Biology Predicts Food Web Complexity.” *Proceedings of the National Academy of Sciences of the United States of America* 103: ~~13745–9.~~13745–49. [://000240648300036.](https://doi.org/10.1016/j.ecss.2010.12.007)
- Bergamino, Leandro, Diego Lercari, and Omar Defeo. 2011. “Food Web Structure of Sandy Beaches: ~~Temporal~~Temporal and Spatial Variation Using Stable Isotope Analysis.” *Estuarine, Coastal and Shelf Science* 91 (4): 536–43. <https://doi.org/10.1016/j.ecss.2010.12.007>.
- Brose, Ulrich, Lara Cushing, Eric L. Berlow, Tomas Jonsson, Carolin Banasek-Richter, Louis-Felix Bersier, Julia L. Blanchard, et al. 2005. ~~“Body Sizes of Consumers and Their Resources.”~~“Body Sizes of Consumers and Their Resources.” *Ecology* 86 (9): ~~2545–52~~2545–45. <https://doi.org/10.1890/05-0379>.
- Carpenter, Stephen R. 2016. “Ecological Futures: Building an Ecology of the Long Now.” *Ecology*, October, 2069–83. [https://doi.org/10.1890/0012-9658\(2002\)083%5B2069:EFBAE0%5D2.0.CO;2@10.1002/\(ISSN\)1939-9170.MacArthurAward](https://doi.org/10.1890/0012-9658(2002)083%5B2069:EFBAE0%5D2.0.CO;2@10.1002/(ISSN)1939-9170.MacArthurAward).
- Cattin, Marie-France, Louis-F’elix Bersier, Carolin Banašek-Richter, Richard Baltensperger, and Jean-Pierre Gabriel. 2004. “Phylogenetic Constraints and Adaptation Explain Food-Web Structure.” *Nature* 427 (6977, 6977)-~~Nature Publishing Group~~: 835–39. <https://doi.org/10.1038/nature02327>.
- Cohen, Joel E. 1989. “Just Proportions in Food Webs.” *Nature* 341 (6238, 6238)-~~Nature Publishing Group~~: 104–5. <https://doi.org/10.1038/341104b0>.
- Cohen, Joel E., C. M. Newman, and John Hyslop Steele. 1985. “A Stochastic Theory of Community Food Webs ~~I-Models-I.~~Models and Aggregated Data.” *Proceedings of the Royal Society of London. Series B. Biological Sciences* 224 (1237)-~~Royal Society~~: 421–48. <https://doi.org/10.1098/rspb.1985.0042>.
- Crawford, Kerry, Robbie A. McDonald, and Stuart Bearhop. 2008. “Applications of Stable Isotope Techniques to the Ecology of Mammals.” *Mammal Review* 38 (1): 87–107. <https://doi.org/10.1111/j.1365-2907.2008.00120.x>.
- Cressie, Noel, Catherine A. Calder, James S. Clark, Jay M. Ver Hoef, and Christopher K. Wikle. 2009. “Accounting for Uncertainty in Ecological Analysis: The Strengths and Limitations of

- 570 Hierarchical Statistical Modeling.” *Ecological Applications* 19 (3): 553–70. [https://doi.org/](https://doi.org/10.1890/07-0744.1)
571 10.1890/07-0744.1.
- 572 Dawah, Hassan Ali, Bradford A. Hawkins, and Michael F. Claridge. 1995. “Structure of
573 the ~~Parasitoid Communities of Grass-Feeding Chalcid Wasps.~~” [Parasitoid Communities of](#)
574 [Grass-Feeding Chalcid Wasps.](#)” *The Journal of Animal Ecology* 64 (6): 708. [https://doi.](https://doi.org/10.2307/5850)
575 [org/10.2307/5850](https://doi.org/10.2307/5850).
- 576 Dunne, Jennifer A., Richard J. Williams, and Neo D. Martinez. 2002. “Network Structure and
577 Biodiversity Loss in Food Webs: Robustness Increases with Connectance.” *Ecology Letters* 5 (4):
578 558–67.
- 579 Emmerson, Mark C., and Dave Raffaelli. 2004. “Predator–Prey Body Size, Interaction Strength
580 and the Stability of a Real Food Web.” *Journal of Animal Ecology* 73 (3): 399–409. [https:](https://doi.org/10.1111/j.0021-8790.2004.00818.x)
581 [//doi.org/10.1111/j.0021-8790.2004.00818.x](https://doi.org/10.1111/j.0021-8790.2004.00818.x).
- 582 Goldwasser, Lloyd, and Jonathan Roughgarden. 1993. “Construction and ~~Analysis of a~~ [Analysis of](#)
583 [a](#) Large Caribbean Food Web: ~~Ecological Archives E074-001.~~” [Ecological Archives E074-001.](#)”
584 *Ecology* 74 (4): 1216–33. <https://doi.org/10.2307/1940492>.
- 585 Gravel, Dominique, Timoth’ee Poisot, Camille Albouy, Laure Velez, and David Mouillot. 2013.
586 “Inferring Food Web Structure from Predator-Prey Body Size Relationships.” Edited by Robert
587 Freckleton. *Methods in Ecology and Evolution* 4 (11): 1083–90. [https://doi.org/10.1111/](https://doi.org/10.1111/2041-210X.12103)
588 [2041-210X.12103](https://doi.org/10.1111/2041-210X.12103).
- 589 Harper-Smith, Sarah, Eric L. Berlow, Roland A. Knapp, Richard J. Williams, and Neo D. Mar-
590 tinez. 2005. ~~“COMMUNICATING ECOLOGY THROUGH FOOD WEBS: VISUAL-~~
591 ~~IZING AND QUANTIFYING THE EFFECTS OF STOCKING ALPINE LAKES WITH~~
592 ~~TROUT.”~~ ~~In~~” [In](#) *Dynamic Food Webs*, 407–23. ~~Elsevier~~ [Elsevier](#). [https://doi.org/10.1016/](https://doi.org/10.1016/B978-012088458-2/50038-2)
593 [B978-012088458-2/50038-2](https://doi.org/10.1016/B978-012088458-2/50038-2).
- 594 Hattab, Tarek, Fabien Leprieur, Frida Ben Rais Lasram, Dominique Gravel, François Le Loc’h,
595 and Camille Albouy. 2016. “Forecasting Fine-Scale Changes in the Food-Web Structure of
596 Coastal Marine Communities Under Climate Change.” *Ecography* 39 (12): 1227–37. [https:](https://doi.org/10.1111/ecog.01937)
597 [//doi.org/10.1111/ecog.01937](https://doi.org/10.1111/ecog.01937).

- 598 Hobson, Keith A., John F. Piatt, and Jay Pitocchelli. 1994. ~~“Using Stable Isotopes to~~ “Using
 599 Stable Isotopes to Determine Seabird Trophic Relationships.” *Journal of Animal Ecology* 63
 600 (4). ~~Wiley, British Ecological Society~~: 786–98. <https://doi.org/10.2307/5256>.
- 601 Hudson, Lawrence N., Rob Emerson, Gareth B. Jenkins, Katrin Layer, Mark E. Ledger, Doris E.
 602 Pichler, Murray S. A. Thompson, Eoin J. O’Gorman, Guy Woodward, and Daniel C. Reuman.
 603 2013. “Cheddar: Analysis and Visualisation of Ecological Communities in ~~R~~.” R” *Methods in*
 604 *Ecology and Evolution* 4 (1): 99–104. <https://doi.org/10.1111/2041-210X.12005>.
- 605 Ibanez, S’ebastien. 2012. “Optimizing Size Thresholds in a Plant–Pollinator Interaction Web:
 606 Towards a Mechanistic Understanding of Ecological Networks.” *Oecologia* 170 (1): 233–42.
 607 <https://doi.org/10.1007/s00442-012-2290-3>.
- 608 Jabot, Franck, and J’erôme Chave. 2009. “Inferring the Parameters of the Neutral Theory of
 609 Biodiversity Using Phylogenetic Information and Implications for Tropical Forests.” *Ecology*
 610 *Letters* 12 (3): 239–48. <https://doi.org/10.1111/j.1461-0248.2008.01280.x>.
- 611 Jonsson, Tomas. 1998. “Food Webs and the Distribution of Body Sizes.” ~~PhD~~.
- 612 Jonsson, Tomas, Joel E. Cohen, and Stephen R. Carpenter. 2005. ~~“Food Webs, Body Size, and~~
 613 ~~Species Abundance in~~ “Food Webs, Body Size, and Species Abundance in Ecological Community
 614 Description.” ~~In~~ In *Advances in Ecological Research*, 36:1–84. ~~Elsevier~~ Elsevier. [https:](https://doi.org/10.1016/S0065-2504(05)36001-6)
 615 [//doi.org/10.1016/S0065-2504\(05\)36001-6](https://doi.org/10.1016/S0065-2504(05)36001-6).
- 616 Jord’an, Ferenc, and Györgyi Osv’ath. 2009. “The Sensitivity of Food Web Topology to Temporal
 617 Data Aggregation.” *Ecological Modelling* 220 (22): ~~3141–63~~ 141–46. [https://doi.org/10.1016/](https://doi.org/10.1016/j.ecolmodel.2009.05.002)
 618 [j.ecolmodel.2009.05.002](https://doi.org/10.1016/j.ecolmodel.2009.05.002).
- 619 Knight, Tiffany M., Michael W. McCoy, Jonathan M. Chase, Krista A. McCoy, and Robert D.
 620 Holt. 2005. “Trophic Cascades Across Ecosystems.” *Nature* 437 (7060): 880–83. [https:](https://doi.org/10.1038/nature03962)
 621 [//doi.org/10.1038/nature03962](https://doi.org/10.1038/nature03962).
- 622 Krause, Ann E., Kenneth A. Frank, Doran M. Mason, Robert E. Ulanowicz, and William W.
 623 Taylor. 2003. “Compartments Revealed in Food-Web Structure.” *Nature* 426 (6964): 282–85.
 624 <https://doi.org/10.1038/nature02115>.

- 625 Lafferty, K. D., A. P. Dobson, and A. M. Kuris. 2006. “Parasites Dominate Food Web Links.”
 626 *Proceedings of the National Academy of Sciences* 103 (30): ~~11211–6~~11211–16. [https://doi.](https://doi.org/10.1073/pnas.0604755103)
 627 [org/10.1073/pnas.0604755103](https://doi.org/10.1073/pnas.0604755103).
- 628 Layman, Craig A., D. Albrey Arrington, Carmen G. Montaña, and David M. Post. 2007. ~~“Can~~
 629 ~~“Can~~ Stable Isotope Ratios Provide ~~for Community-Wide Measures of Trophic Structure?”~~ ~~for~~
 630 Community-Wide Measures of Trophic Structure?” *Ecology* 88 (1): 42–48. [https://doi.org/](https://doi.org/10.1890/0012-9658(2007)88%5B42:CSIRPF%5D2.0.CO;2)
 631 [10.1890/0012-9658\(2007\)88%5B42:CSIRPF%5D2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88%5B42:CSIRPF%5D2.0.CO;2).
- 632 Lindegren, Martin, Christian Möllmann, Anders Nielsen, Keith Brander, Brian R. MacKenzie,
 633 and Nils Chr. Stenseth. 2010. “Ecological Forecasting Under Climate Change: The Case of
 634 ~~Baltic~~-Baltic Cod.” *Proceedings of the Royal Society B: Biological Sciences* 277 (1691): 2121–30.
 635 <https://doi.org/10.1098/rspb.2010.0353>.
- 636 Lurgi, Miguel, Bernat C. L’opez, and Jos’e M. Montoya. 2012. “Climate Change Impacts on Body
 637 Size and Food Web Structure on Mountain Ecosystems.” *Philosophical Transactions of the Royal*
 638 *Society B: Biological Sciences* 367 (1605): ~~3050–7~~3050–57. [https://doi.org/10.1098/rstb.](https://doi.org/10.1098/rstb.2012.0239)
 639 [2012.0239](https://doi.org/10.1098/rstb.2012.0239).
- 640 MacArthur, Robert H., and Eric R. Pianka. 1966. ~~“On Optimal Use of a Patchy Environment.”~~
 641 “On Optimal Use of a Patchy Environment.” *The American Naturalist* 100 (916): 603–9.
- 642 Martinez, Neo D. 1991. “Artifacts or ~~Attributes? Effects of Resolution on the~~ ~~Attributes? Effects~~
 643 of Resolution on the Little Rock Lake Food Web.” *Ecological Monographs* 61 (4): 367–92.
 644 <https://doi.org/10.2307/2937047>.
- 645 Martinez, Neo D., Bradford A. Hawkins, Hassan Ali Dawah, and Brian P. Feifarek. 1999. “Ef-
 646 fects of ~~Sampling Effort on Characterization of Food-Web Structure.~~ Sampling Effort on
 647 Characterization of Food-Web Structure.” *Ecology* 80 (3): 1044–55. [https://doi.org/10.](https://doi.org/10.1890/0012-9658(1999)080%5B1044:E0SE0C%5D2.0.CO;2)
 648 [1890/0012-9658\(1999\)080%5B1044:E0SE0C%5D2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080%5B1044:E0SE0C%5D2.0.CO;2).
- 649 May, Robert M. 1972. “Will a Large Complex System ~~Be Stable?”~~ Be Stable?” *Nature* 238 (5364):
 650 413. <https://doi.org/10.1038/238413a0>.

- Memmott, J., N. D. Martinez, and J. E. Cohen. 2000. "Predators, Parasitoids and Pathogens: Species Richness, Trophic Generality and Body Sizes in a Natural Food Web." *Journal of Animal Ecology* 69 (1): 1–15. <https://doi.org/10.1046/j.1365-2656.2000.00367.x>.
- Morris, Rebecca J., Frazer H. Sinclair, and Chris J. Burwell. 2015. "Food Web Structure Changes with Elevation but Not Rainforest Stratum." *Ecography* 38 (8): 792–802. <https://doi.org/10.1111/ecog.01078>.
- O'Connor, Mary I., Michael F. Piehler, Dina M. Leech, Andrea Anton, and John F. Bruno. 2009. "Warming and Resource Availability Shift Food Web Structure ~~and Metabolism.~~" [and Metabolism.](#) Edited by Michel Loreau. *PLoS Biology* 7 (8): e1000178. <https://doi.org/10.1371/journal.pbio.1000178>.
- ~~O'Gorman, Eoin J., Owen L. Petchey, Katy J. Faulkner, Bruno Gallo, Timothy A. C. Gordon, Joana Neto-Cerejeira, Jon S. 'Olafsson, Doris E. Pichler, Murray S. A. Thompson, and Guy Woodward. 2019. "A Simple Model Predicts How Warming Simplifies Wild Food Webs." *Nature Climate Change* 9 (8): 611–16.~~
- Opitz, Silvia. 1996. "Quantitative Models of Trophic Interactions in Caribbean Coral Reefs." ~~Journal of~~
- Patonai, Katalin, and Ferenc Jord'an. 2017. "Aggregation of Incomplete Food Web Data May Help to Suggest Sampling Strategies." *Ecological Modelling* 352 (May): 77–89. <https://doi.org/10.1016/j.ecolmodel.2017.02.024>.
- Peralta-Maraver, I., M. J. L'opez-Rodríguez'opez-Rodríguez, and J. M. Tierno de Figueroa. ~~2017.~~ [2016.](#) "Structure, Dynamics and Stability of a ~~Mediterranean~~ [Mediterranean](#) River Food Web." *Marine and Freshwater Research* 68 (3). ~~CSIRO PUBLISHING~~: 484–95. <https://doi.org/10.1071/MF15154>.
- ~~Petchey, Owen L., Andrew P. Beckerman, Jens O. Riede, and Philip H. Warren. 2008. "Size, Foraging, and Food Web Structure." [Peralta-Maraver, I., M. J. L'opez-Rodríguez, and J. M. Tierno de Figueroa. 2017. "Structure, Dynamics and Stability of a Mediterranean River Food Web." *Proceedings of the National Academy of Sciences* \[Marine and Freshwater Research\]\(#\) 405 \(11\). National Acad Sciences: 4191–6.](#)~~ [68 \(3\): 484–95.](#) <https://doi.org/10.1071/MF15154>.

- Petchey, Owen L., P. Timon McPhearson, Timothy M. Casey, and Peter J. Morin. 1999. "Environmental Warming Alters Food-Web Structure and Ecosystem Function." *Nature* 402 (6757): 69–72. <https://doi.org/10.1038/47023>.
- Petchey, Owen L., Mikael Pontarp, Thomas M. Massie, Sonia K'efi, Arpat Ozgul, Maja Weilenmann, Gian Marco Palamara, et al. 2015. "The Ecological Forecast Horizon, and Examples of Its Uses and Determinants." *Ecology Letters* 18 (7): 597–611. <https://doi.org/10.1111/ele.12443>.
- Petchey, Owen L, Andrew P Beckerman, Jens O Riede, and Philip H Warren. 2008. "Size, Foraging, and Food Web Structure." *Proceedings of the National Academy of Sciences* 105 (11): 4191–96.
- Poisot, Timoth'ee, and Dominique Gravel. 2014. "When Is an Ecological Network Complex? ~~Connectance~~ Connectance Drives Degree Distribution and Emerging Network Properties." *PeerJ* 2 (February): ~~PeerJ Inc.~~ e251. <https://doi.org/10.7717/peerj.251>.
- Poisot, Timoth'ee, and Daniel B. Stouffer. 2016. "How Ecological Networks Evolve." ~~Preprint~~ Preprint. Ecology. Eeology. <https://doi.org/10.1101/071993>.
- Polis, Gary A. 1991. ~~"Complex Trophic Interactions in Deserts: of Food-Web Theory."~~ "Complex Trophic Interactions in Deserts: An Empirical Critique of Food-Web Theory." *The American Naturalist* 138 (1): 123–55. <http://www.jstor.org/stable/2462536>.
- Rall, B. C., U. Brose, M. Hartvig, G. Kalinkat, F. Schwarzmuller, O. Vucic-Pestic, and O. L. Petchey. 2012. "Universal Temperature and Body-Mass Scaling of Feeding Rates." *Philosophical Transactions of the Royal Society B: Biological Sciences* 367 (1605): 2923–34. <https://doi.org/10.1098/rstb.2012.0242>.
- Shriner, Daniel, Yi Liu, David C. Nickle, and James I. Mullins. 2006. "Evolution of ~~Intrahost Hiv~~ Intrahost Hiv - 1 Genetic Diversity During Chronic Infection." *Evolution* 60 (6): 1165–76. <https://doi.org/10.1111/j.0014-3820.2006.tb01195.x>.
- Stouffer, Daniel B., Juan Camacho, and Lu'is A. Nunes Amaral. 2006. "A Robust Measure of Food Web Intervality." *Proceedings of the National Academy of Sciences* 103 (50): 19015–20. <https://doi.org/10.1073/pnas.0603844103>.

- 706 Tamaddoni-Nezhad, Alireza, Ghazal Afroozi Milani, Alan Raybould, Stephen Muggleton, and David
 707 A. Bohan. 2013. “Construction and ~~Validation of~~ Validation of Food Webs Using ~~Logic-Based~~
 708 ~~Machine Learning and Text Mining.~~” In Logic-Based Machine Learning and Text Mining. In
 709 *Advances in Ecological Research*, 49:225–89. ~~Elsevier~~Elsevier. [https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-420002-9.00004-4)
 710 B978-0-12-420002-9.00004-4.
- 711 Toni, Tina, David Welch, Natalja Strelkowa, Andreas Ipsen, and Michael P. H. Stumpf. 2009.
 712 ~~“Approximate Bayesian.”~~ “Approximate Bayesian Computation Scheme for Parameter Inference
 713 and Model Selection in Dynamical Systems.” *Journal of the Royal Society Interface* 6 (31):
 714 187–202. <https://doi.org/10.1098/rsif.2008.0172>.
- 715 Tylianakis, Jason M., and Amrei Binzer. 2014. “Effects of Global Environmental Changes on
 716 ~~Parasitoid Host~~ Parasitoid host Food Webs and Biological Control.” *Biological Control* 75 (Au-
 717 gust): 77–86. <https://doi.org/10.1016/j.biocontrol.2013.10.003>.
- 718 Warren, Philip H. 1989. “Spatial and ~~Temporal Variation in the Structure of a~~ Temporal Variation
 719 in the Structure of a Freshwater Food Web.” *Oikos* 55 (3): 299. [https://doi.org/10.2307/](https://doi.org/10.2307/3565588)
 720 3565588.
- 721 Williams, Richard J., and Neo D. Martinez. 2000. “Simple Rules Yield Complex Food Webs.” *Nature*
 722 404 (6774, 6774): ~~Nature Publishing Group~~: 180–83. <https://doi.org/10.1038/35004572>.
- 723 Woodward, Guy, Julia Blanchard, Rasmus B. Lauridsen, Francois K. Edwards, J. Iwan Jones,
 724 David Figueroa, Philip H. Warren, and Owen L. Petchey. 2010. “Chapter 6 - ~~Individual-Based~~
 725 ~~Food Webs: Species Identity, Body Size and Sampling Effects.~~” In Individual-Based Food Webs:
 726 Species Identity, Body Size and Sampling Effects. In *Advances in Ecological Research*, edited
 727 by Guy Woodward, 43:211–66. Integrative ~~Ecology: From Molecules to Ecosystems~~Ecology:
 728 From Molecules to Ecosystems. Academic Press. ~~Academic Press.~~[https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-385005-8.00006-X)
 729 B978-0-12-385005-8.00006-X.
- 730 Woodward, Guy, and Alan G. Hildrew. 2001. “Invasion of a Stream Food Web by a New Top
 731 Predator.” *Journal of Animal Ecology* 70 (2): 273–88. [https://doi.org/10.1111/j.1365-2656.](https://doi.org/10.1111/j.1365-2656.2001.00497.x)
 732 2001.00497.x.

- 733 Woodward, Guy, Dougie C. Speirs, and Alan G. Hildrew. 2005. "Quantification and ~~Resolution of a~~
734 ~~Complex, Size-Structured Food Web.~~" ~~In~~ Resolution of a Complex, Size-Structured Food Web.
735 In *Advances in Ecological Research*, 36:85–135. ~~Elsevier~~Elsevier. [https://doi.org/10.1016/](https://doi.org/10.1016/S0065-2504(05)36002-8)
736 [S0065-2504\(05\)36002-8](https://doi.org/10.1016/S0065-2504(05)36002-8).
- 737 Yodzis, Peter. 1998. "Local Trophodynamics and the Interaction of Marine Mammals and Fisheries
738 in the ~~Benguela~~-Benguela Ecosystem." *Journal of Animal Ecology* 67 (4): 635–58. [https:](https://doi.org/10.1046/j.1365-2656.1998.00224.x)
739 [//doi.org/10.1046/j.1365-2656.1998.00224.x](https://doi.org/10.1046/j.1365-2656.1998.00224.x).