Q222

R21: Fresh Complete Rederivation

Mathematical proof and source documentation

22 October 2026

Fresh R21 rederivation, 22 October 2026. The opening assembly states the quantified theorem and the complete finite source cover. The following components supply the actual mathematical arguments under their explicit hypotheses. This volume is a new presentation, not a claim of byte-identical restoration of the earlier release.

How to read this proof volume

The opening assembly provides the complete logical route. The following components reproduce the operative formulas, hypotheses and proof arguments of the listed mathematical bodies. Each component has a stable identifier derived from its source SHA256 and a source-to-presentation record.

Original source-time qualifications and status prose are retained as source context. A historical completion statement is not, by its inclusion, an additional mathematical premise. Restricted imports and replacements must be read in the scopes stated in the opening assembly and component inventory. The two coefficient/testing fields are separately labelled; the general theorem does not require algebraic disjointness.

Standalone preambles, title commands and duplicate physical inclusions are removed. Labels are namespaced in presentation copies. Approved editorial repairs replace obsolete cross-references by explicit formula ranges and links to the actual replacement proofs; each exact substitution is logged in the source-to-presentation record. Explicitly scoped extracts preserve every imported argument in full; any excluded source-time passages are recorded verbatim with their source spans and hashes. A reference is resolved only when its actual target is known; unresolved references remain visible and are listed in the build report. Original files are immutable.

Contents

How to read this proof volume **i**

1 The elementary mechanism: positive residue pays for errors **6**

2 Fresh reconstruction of the physical model and conditional assembly **7**

3 The physical geometry of the canonical reduction **13**

4 Every noncommuting positive-d contact is regular **17**

5 Zero signed-weight residues: the eight-cycle geometry **22**

6 Weighted normal completion at matching residues **25**

7 A fourth edge at a smaller coefficient scale **29**

8 A matching terminal independent of its internal coefficient scales **32**

9 One hard event and a positive next-scale signed coefficient **35**

10 Two hard events and a positive next-scale signed coefficient 37

11 A finite induction on event constraints 39

12 A short proof for every noncommuting residue 41

13 All closed faces from two elementary matrix lifts 47

14 Fresh finite-contact entry and the exact two-family source target 51

15 Global native entry from actual physical contacts 56

16 R1 unbounded-q: exact contact circuits and their square budgets 60

17 Reconstructed R1 unbounded-q source and scalar certificate 63

18 Reconstructed small normal budgets for R1 unbounded-q 69

19 A freshly derived full-state sign and projective join 71

20 Reconstructed full R1 conductor in the unbounded source-ratio sector 74

21 Reconstructed R2 source entry and full two-packet extraction 82

22 A simultaneous full-D/full-Y identity and an extra-row conductor 85

23 A uniform native C1 cap and an ordinary-angle admission 88

24 A five-root local certificate on an A-unbounded weak-C2 subchart 91

25 R9 source data and the finite coefficient calculation 95

26 An exact beta-shared normalization for the two eta-large C1.4 charts 98

27 C1.4: unbounded $L = x\zeta$, bounded-unit $\sigma = 1/\eta$ 102

28 Small L, bounded η: exact source openings and the remote complement 107

29 A C1 extension: L a unit and η infinitesimal 110

30 A sharper source bound on the remaining L-unit, η-large face 113

31 C1 with both L and η units 115

32 The q-infinitesimal C1.1 source at both small testing angles 118

33 A uniform hard normal for the q-infinitesimal branch 121

34 The other Bob testing charts for q infinitesimal 123

35 C1.1 with $k^2\eta$ a unit: a complete contact certificate 125

36 C1.1: an opening-uniform full normal and a new angular circuit 129

37 An interior small-L C1 source: a full three-circuit closure 133

38 Small L: a complete native normal identity and the exact soft defect 138

39 C1 UH: a homogeneous biquadratic conductor 141

40 A complete local strong-II Gram with independent testing rates 147

41 The entire remote complement of strong C2 II 152

42 Full hi retained contact module 155

43 Retained contact module 157

44 Beta-positive extension and the exact remaining local source branch 161

45 A complete joint radial allocation on the literal beta-zero face 164

46 Small-L, unbounded eta: uniform source weights 168

47 The actual-j C1 opening: centering the whole divergent event 170

48 Small-L C1: a weighted closure through both r openings 175

49 The simultaneous r, j opening: a harmonic full-event frame 179

50 The shared C1–C2 boundary and a positive source partition 184

51 A centered angular packet closes the physical small-gap C2 chart 186

52 Exact weak-C2 source complement after the small-gap admission 193

53 A three-monomial contact lift with one unit coordinate 196

54 A five-root local certificate on an A-unbounded weak-C2 subchart 198

55 A direct all-tests join for the five-root A-unbounded subchart 202

56 The small-ratio weak-C2 opening is also admitted for all tests 205

57 The small-gap weak-C2 source reduces to two infinitesimal tests 209

58 The physical small-$1 - r$ source has a forced scale bound 211

59 One complete Gram for every bounded test on the small-r opening 213

60 All projective tests on the small-r weak-C2 source opening 216

61 Weak ii source alternatives 219

62 Shared event mechanisms for the remaining C1 sources 222

63 C1 UI: completion of the ordinary/small strip 227

64 A common exact II reduction for the two C2 double corners 232

65 C1 UI: a positive Laurent model and the remaining lift 236

66 C1 UI: a full-parent remote complement 238

67 Exact source entry for the strong HI corner 241

68 A full-parent remote theorem for strong HI sources 243

69 Weak-C IH: the open testing axis 246

70 Strong IH: the complete local corner 249

71 Strong IH: remote charts and the all-tests join 251

72 Strong C2 source entry at an infinitesimal first angle 254

73 Weak IH: exhaustive all-tests admission 256

74 Weak IH: all remote testing charts 257

75 The two missing weak-IH source endpoints 259

76 An exact full-parent Gram at the interior IH double corner 262

77 The genuine directed IH weak-C frontier 266

78 Completion of the strong C2 ordinary–small source chart 268

79 The centered-A certificate for every bounded positive δ 274

80 The outer and unit-state join for infinitesimal e 276

81 Small e: a bounded full-parent contact certificate for every r 281

82 Completion of the unit-η, infinitesimal-e source domain 284

83 The exact strong-C2 ordinary–small source reduction 285

84 A local strong-C2 certificate at the ordinary–small contact 289

85 A reciprocal contact completion for a singular angle 293

86 Five consecutive lower-bounded event packets 295

87 An unbounded first actual angle forces a strong C packet 297

88 Further exact source information on the three broad C1 rows 299

89 Bounded unit native state scale and bounded unit e 302

90 Outer tests and state entry when the native r is infinitesimal 305

91 An infinitesimal native r with unit η and unit e 309

92 The joint rational C2 opening: a full-parent certificate 313

93 Strong C2 pitchfork with arbitrary complementary scales 318

94 A homogeneous scalar lift at the unbounded-state contact 324

95 The unbounded-b source with xb infinitesimal 327

96 Further actual-angle reductions from exact source entry 332

97 The exact unbounded-state-scale C2 complement 334

98 A unit native state scale: the same circuit with a different mass 336

99 Physical entry excludes a small state parameter 339

100 The remaining opening when the opposite event is bounded 340

101 The one-weak C2 family with three arbitrarily small openings 343

102 C2 pitchfork with three arbitrary event weights 347

103 C2 pitchfork with two arbitrarily large retained events 351

104 Ordinary actual Alice angle, with arbitrary state ratio 355

105 A bounded-shape one-weak C2 family, at all tests 359

106 Ordinary-angle C1 with arbitrary positive shape openings 365

107 C1 native formulas and exact stationarity 369

108 An ordinary-angle C1 family: all independent tests 371

109 Two weak complementary events force the C2 radial certificate 375

110 A short determinant check for exact source entry 379

111 Exact C2 source and its uniqueness 381

112 A natural weak-event entry into the complete uniform C1 theorem 382

113 Global sign entry for the adjacent contact family 386

114 The unbounded-e C1 source away from the zero-W state 391

115 All tests for the uniform infinitesimal C1 source 393

116 Matching both weak C2 packets to the four strong packets 396

117 All outer tests for the all-small C2 radial family 399

118 The last bounded-shape C2 corner: a weighted radial root 402

119 The C2 boundary: one uniform five-root proof 405

120 The next C2 boundary: retain the fourth event current 411

121 Endpoint hermite 414

122 Endpoint source 416

123 A product identity repairs the endpoint at every coefficient rate 418

124 All matching-strong C2 sources in the positive contact chart 422

125 A short contact–conductor proof of the C1 memberships 425

126 Self-contained C1 source definitions and contact verification 429

127 The C1 circuit needs six soft directions 434

128 An exhaustive positive-d source partition 436

129 The joint small-a, small-e corner: a bounded-ratio closure 440

130 Two exact conductors for the remaining small-η chart 443

131 The weak-C2 R9 contact corner: a full-parent coupled-circuit certificate 445

132 All-tests closure of the weak-C2 source shape R10 448

133 R10: forced source scale, every remote test, and a weighted local admission 450

134 The positive-n thin R10 layer: an exact extra row and a bounded conductor perturbation 454

135 R10: a direct critical-wall certificate for every unit-W state 458

136 C14.3b: the complete gamma–circuit join and its negative soft source pivot 460

137 C14.3b: a complete source subchart from a damped two-circuit completion 463

138 The damped frame also enlarges the pure-testing state chart 467

139 The entire unit-η large-ζ source, through the actual-j opening 469

140 The positive-unit e subdomain of weak-C2 R4 472

141 C14.1b: the entire parent on bounded testing state quotients 476

142 A full-parent pole extraction and a larger comparison-model state chart 479

143 The damped completion at large zeta with bounded L over eta 483

144 Large zeta: the enlarged state chart and the exact remaining corner 487

145 A pure-testing sign exit for the complete original parent 490

146 The complete ordered weak-C2 shape R5 is admitted 491

147 R6 with bounded z/B is also an all-tests admission 495

148 The bounded-z/B subdomain of R7 is admitted for all tests 497

149 The remaining small-r local chart of weak-C2 R8 501

150 R8: a joint divergent-D certificate, a paired-D thin lift, and the remote join 504

151 The end-to-end C14.3a/b join 510

152 The completed J row conducts the shared remaining normal 513

153 C14.4 at large ζ: every unmatched Bob residue 516

154 C14.2: bounded Bob tests at every local Alice angle 519

155 The exact C14.2 remainder and its centered C extraction 521

156 C14.2: the complete noninfinitesimal-Alice complement 523

157 C14.1b: every state when the first testing angle is a unit 526

158 R6 with unbounded a: the complete thin layer is admitted 529

159 Weak-C2 R3 with a positive-unit $1 - r$ is admitted 533

160 The source, the original polynomial, and the proof language 536

161 C14.2: the complete unbounded-H, unbounded-W/Z chart 540

162 C14.2: pure-testing state exits at the two small angles 543

163 Two complementary C14.2 certificates 545

164 C14.2: bounded h/σ and the unbounded W/Z state chart 548

165 C14.2: bounded h/σ with positive λ, all states 550

166 C14.2 is closed: the exact all-tests and all-states join 553

167 C14.4: a contact-centered full-parent identity through the M opening 555

168 C14.4: extend the unmatched-Bob certificate to the entire small-e source 557

169 State-free normal conductors at the large-L corner 560

170 C14.1b: the bounded testing-ratio complement at every unit h 565

171 C14.1b: the remaining source at bounded Alice-to-reciprocal-Bob ratio 568

172 C14.1b: the entire unit beta-sphere source, including the contact ratio 571

173 A full-parent terminal for a finite positive-d commuting edge 575

174 A positive-d terminal at a finite-ratio distance-four corner 580

175 Positive total scale: the recovered finite-pitchfork fit 584

176 Positive-scale C2: a precise weak-alpha subbundle 588

177 Scope and actual checklist delta 591

178 Uniform C1: the bounded-reciprocal repair and its scope 602

179 The remaining R6 boundary: a^2B infinitesimal 610

180 R6 at a positive residual value of a^2B: an all-tests theorem 616

181 The unbounded-a complement of R7, including its physical zero face 621

182 A fresh angular budget and completion of the retained C14.1b residual 628

183 The complete C14.4 source union and the original C14.5 join 632

184 The entire bounded-h/q part of the doubly small endpoint 636

185 C14.4 at the entire ℓ-infinitesimal source endpoint 639

186 Fresh C14.4 completion when κ is small and both Bob source fractions are units 645

187 A source-free angular budget for the finite matched-Bob corner 650

188 The full unit-κ, infinitesimal-ν source endpoint 652

189 The doubly infinitesimal endpoint with bounded β/q^2 657

190 The unbounded-h/q part of the doubly small endpoint 661

191 All testing charts for the unit-κ, ν, ℓ source 665

192 A fresh matched-Bob completion for unit κ, ν, ℓ 670

193 The complete positive-scale C1 reciprocal terminal 674

194 Every canonical C2 source: an exhaustive mathematical join 681

195 The complete canonical local certificate and global source dispatch 687

196 The complete positive-total-scale canonical source union 695

197 A fresh complete outer-testing certificate for the original R2 source 701

198 A rederived three-event wall and its exact contact exclusion 706

199 The uniform C1 source for every positive e 711

200 Weak-alpha source entry and the unit-n terminal 713

201 Weak-alpha weak-n and C1: the complete parent 718

202 A source-exact projective pitchfork for C1 725

203 R2 with an unbounded source ratio and a bounded relative current 729

204 R2 with both the native source ratio and relative current unbounded 734

205 A direct certificate for the positive-d C2 reciprocal corner 739

206 Both uniform C1 source charts on the entire bounded $Z = 1$ state chart 744

207 The inverse-b uniform C1 source: a complete local conductor 749

208 A new complete R1 theorem with no source-ratio split 756

209 The entire original R2 source, with all independent tests and states 762

210 R2 with bounded a, or an infinitesimal paired-event parameter 764

211 Two exact circuit budgets and the complete R2 opposite normal 768

212 R3 and R4 at infinitesimal e, with all source ratios and tests 770

213 The complete uniform infinitesimal C1 source, at all tests 776

214 The actual hypotheses used by the retained weak-C2 leaves 779

215 The complete directed weak-C2 source and testing join 784

Reader's guide and the completed existence proof

This volume is a fresh rederivation of the finite semidefinite-lift existence argument for the two-party, two-setting, two-outcome quantum set. It uses the surviving mathematical sources and supplies new proofs for the missing local arguments and final source assembly. It is not a restoration of the missing original 549-page PDF or of the original R21 archive bytes. The source archive preserves the retained originals and records the relationship between each original mathematical component and its presentation here.

Completed mathematical assembly. The complete canonical local-certificate theorem is proved in LC1–LC10, in the component beginning on page 687. It connects the actual physical source reduction to the positive-scale theorem, every C1 source branch, and every C2 source branch, then restores the original homogeneous slack and finite source/testing combinations. The opening argument below states that conclusion with its full quantifiers and applies the finite-lift criterion. The later components contain the complete local constructions and source implications on which that assembly depends.

What the statement means

Let $\mathcal{Q}_{222} \subset \mathbb{R}^8$ be the set of correlations of finite-dimensional bipartite complex tensor-product experiments, with two binary measurements on each side. We use the coordinates

$$q = (a_0, a_1, b_0, b_1, E_{00}, E_{01}, E_{10}, E_{11}),$$

where a_x and b_y are the single-party expectations and E_{xy} is the joint expectation of the $\{+1, -1\}$-valued observables. Write P for the cone of real affine functions nonnegative on this set.

The physical reduction proves the exact formula

$$\mathcal{Q}_{222} = \left\{ \sum_{i=1}^{9} w_i \Phi(z_i) : z_i \in S^1 \times S^1 \times S^3, \quad w_i \geq 0, \quad \sum_i w_i = 1 \right\}. \tag{FA1}$$

Here Φ is the polynomial expectation map for $A_0 = B_0 = Z$, $A_1 = c_A Z + s_A X$, $B_1 = c_B Z + s_B X$, and a real unit two-qubit vector. The proof includes a simultaneous dilation of both binary effects of each party, the two-involution block decomposition, the real/imaginary splitting of a complex state, finite direct-sum realization, and the nine-point convexification argument. Consequently $\mathcal{Q}_{222}$ is compact, convex, and semialgebraic over $\mathbb{R}$.

For a real closed extension $K \supset \mathbb{R}$, the symbol $(\mathcal{Q}_{222})_K$ means the set defined by the same fixed finite formula (FA1), with its variables in K. Likewise

$$P_K = \{ f \in K[q]_{\leq 1} : f(q) \geq 0 \text{ for all } q \in (\mathcal{Q}_{222})_K \}. \tag{FA2}$$

The identification with the transferred real cone is proved by the first-order description of validity. No compactness assertion about the order topology of a non-Archimedean field is used.

Separately labelled fields

Choose arbitrarily two real closed extensions $K_c, K_t \supset \mathbb{R}$, and put $T = K_c \otimes_{\mathbb{R}} K_t$. The subscripts distinguish coefficient variables from testing variables. The two fields may be two labelled copies of the same extension. No algebraic-disjointness assumption, embedding of T in an ordered field, or ordering of T is made. For $f = f_0 + \sum_{\nu=1}^{8} f_\nu q_\nu$, define

$$[f, q] = f_0 \otimes 1 + \sum_{\nu=1}^{8} f_\nu \otimes q_\nu. \tag{FA3}$$

The target is membership in ΣT^2, the set of finite sums of squares of elements of this tensor ring. The finite list of roots may depend on the chosen fields, functional and testing point.

The local proofs use the convex subring $\mathcal{O}(K)$ of elements bounded in absolute value by a real constant, and its ideal $\mathfrak{m}(K)$ of infinitesimals. A positive element is of exactly one of three types: positive infinitesimal, positive bounded unit (both it and its inverse are bounded), or positive unbounded. Literal zero is kept separate. The residue of a bounded element is its unique real standard part. These types organize the proof; they do not identify an independent source infinitesimal with a testing infinitesimal.

The common algebraic tool states that a bounded symmetric tensor matrix with positive definite real residue gives a finite tensor SOS when evaluated on any finite list of tensor roots. Its proof uses finite positive pure-factor decompositions and a fixed real diagonal-dominance reserve. It never takes the inverse or square root of a mixed tensor. The roots themselves need not be bounded. The complete strict-Gram proof is included among the foundations.

The full polynomial that every local proof certifies

In the fixed CHSH orientation put

$$H = 1 - \tfrac{1}{2}(E_{00} - E_{01} + E_{10} + E_{11}), \qquad f = dH + 4\sum_{\xi} \xi p_{\xi}, \quad \xi \in \{\lambda, n, m, C, A, D, \alpha, \beta\}. \tag{FA4}$$

The coefficients are nonnegative, and $d > 0$ on the nontrivial canonical source. With $D_u = 1 + u^2$, $D_v = 1 + v^2$, and $\psi = (W, X, Y, Z)$, the complete affine-chart parent is

$$\begin{aligned} F = {} & 4\lambda D_u(W + vX)^2 + 4nD_uD_vX^2 + 4m(vW - X + uvY - uZ)^2 \\ & + 4CD_v(uW - Y)^2 + 4AD_uD_vY^2 + (4D + 2d)(uW + uvX - Y - vZ)^2 \\ & + (4\alpha + 2d)D_u(vY - Z)^2 + (4\beta + 2d)D_v(X + uZ)^2 - 2dD_uD_vZ^2. \end{aligned} \tag{FA5}$$

Products of a coefficient-field number and a testing-field expression in (FA5) always denote their separated tensor product. A scalar restriction to a contact graph does not by itself prove the required statement: the original state-normal terms and all higher terms must also be included. Each local proof gives an exact identity or a complete root allocation for this parent, and restores every retained positive packet after proving the bounded remainder.

For homogeneous angle pairs (a_0, a_1) and (b_0, b_1), put $\Delta_A = a_0^2 + a_1^2 > 0$, $\Delta_B = b_0^2 + b_1^2 > 0$. The homogeneous parent $\mathcal{F}$ satisfies

$$\mathcal{F} = (1 \otimes \Delta_A\Delta_B\|\psi\|^2)[f, \Phi(z)] \tag{FA6}$$

for every nonzero state, with z its normalized physical atom. The homogeneous identity is proved directly from all eight event amplitudes. It includes the zero and infinite angle coordinates; those points are not obtained by cancelling a vanished affine coordinate. The zero state is immediate.

Source reduction and the precise local proposition

Every valid functional is either a nonnegative event sum or has a negative value at a PR vertex. In the latter case the eight neighboring deterministic vertices and that PR vertex form the affine evaluation basis giving (FA4). Normalize the positive coefficient sum in the source field. The real compact convex source base and its fixed nine-term extreme-point representation transfer as finite first-order formulas. The determinant has a convex logarithmic Laurent profile. Its coercive extreme sources have an actual finite noncommuting stationary contact and five or six positive events. The incompatible omission cases have two explicit closed-face matrix lifts.

The remaining compatible omission patterns reduce, by actual Bell symmetries, to

$$\begin{aligned} &\text{C1}: \quad d, m, C, A, \alpha > 0, \quad \lambda = n = 0, \quad D, \beta \geq 0, \quad D + \beta > 0, \\ &\text{C2}: \quad d, m, C, A, D, \alpha > 0, \quad \lambda = \beta = 0, \quad n \geq 0. \end{aligned} \tag{FA7}$$

The physicality condition and the actual source contact are retained. The native-entry proof derives all strict signs and gaps from these conditions, including the literal third-omission faces. Source contacts are never substituted for independent testing points.

Complete local-certificate proposition (LC). For every source in (FA7), over every K_c, and every independent projective angle pair and state over every K_t, the homogeneous parent $\mathcal{F}$ is a finite sum of squares in T, possibly after multiplication by a proved positive pure-source factor. The statement includes every allowed zero coefficient, arbitrary positive openings, all relative source and testing rates, all state ratios, exact sign gauges, and literal projective endpoints.

The total-scale split uses

$$\delta_{\mathrm{tot}} = \frac{d}{d+\lambda+n+m+C+A+D+\alpha+\beta}. \tag{FA8}$$

It is taken before a later $d = 1$ or $C = 2$ gauge. For $d > 0$ the quotient is positive bounded, so its positive-unit and positive-infinitesimal alternatives are exhaustive. Setting $d = 1$ does not select the first alternative. Literal $d = 0$ is a separate sum of event squares. At a literally commuting testing pair the signed core is nonnegative in the testing field, so it too has a direct separated certificate.

Proof of LC. The native-entry theorem EN1–EN16 derives the exact source signs, strict gaps and stationary formulas directly from physicality and the actual finite contact. It uses single-field source geometry and does not assume a tensor certificate. On the positive-unit alternative of (FA8), AU1–AU19 proves the entire parent at every independent test. All coefficients are then bounded after the positive source normalization $d = 1$; the weak/unit coefficient partitions in AU exhaust their possibilities, including the allowed literal zeros.

Retain the positive-infinitesimal alternative of (FA8) for the remaining sources. In C1 orient the native Alice parameter to $0 < k \le 1$. For unit k, the actual source equations give the complete AS, ordinary unit/unit, unit/small and unit/unbounded contact alternatives. The ordinary unit/unit source has five consecutive strong events by the exact NC7 elimination; the other alternatives use their complete CU/CR and UH full-parent constructions. For infinitesimal k, noninfinitesimal ζ is the full C1.4 domain. LC4a–LC4c gives its bare-source partition, including all seven earlier exits and the five normalized C14.4 cells. The actual JC constructions prove every cell without obtaining a bound from an unspecified earlier exclusion.

If both k and ζ are infinitesimal, the three types of η complete the C1 cover. Infinitesimal η uses the proved uniform all-e join; unbounded η uses the full C1.1 unit-q and small-q theorems. Unit η is divided by the three types of $e = k\sqrt{\eta/\zeta}$ and, where necessary, by unit or infinitesimal $r \in (0, 1)$. The complete small-e, bounded-e, centered-A and unbounded-e theorems cover those alternatives. These are exactly the ten disjoint rows of LC5, with the original total scale, source gaps, allowed zero coefficients and independent testing/state variables retained.

For C2, CT1–CT16 first treats the two weak complementary events by its actual matching-source entry and full radial and outer constructions. Otherwise its exact full-parent reflection orients the asymmetric weak case, and its nine ordered actual-angle types are exhaustive. Every occupied cell has a complete local and remote proof; each empty cell has an explicit necessary-source contradiction. The directed weak unbounded/unbounded cell is the exact reflection of HJ's ten-row source join. Its $n = 0$ C1 calls use only two independently proved bare C1 terminals, so neither CT nor HJ assumes the present global conclusion.

Every testing exit uses the whole physical parent and all state-normal terms. Noncommuting testing residues use the separately proved A6 construction on the original physical source. The homogeneous identities and explicit endpoint charts supply literal zero/infinite angles; all allowed state ratios and both signs are included. Complex states split into their real and imaginary parts. Restoring the proved nonzero pure-source and pure-testing factors gives the unmultiplied homogeneous parent. This proves LC. LC1–LC10 and LCH1–LCH3 give the detailed source predicates, complete dependency direction and exact reconciliation with every historical row; the actual component proofs supply each conductor, full-normal allocation and strict Gram used in this join. □

Finite closure and the existence implication

Theorem 0.1 (Universal tensor statement and finite semidefinite lift)**.** *For every pair of separately labelled real closed extensions $K_c, K_t \supset \mathbb{R}$,*

$$[f, q] \in \Sigma(K_c \otimes_{\mathbb{R}} K_t)^2 \quad \text{for every } f \in P_{K_c}, \quad q \in (\mathcal{Q}_{222})_{K_t}. \tag{FA9}$$

Consequently $\mathcal{Q}_{222}$ has a finite semidefinite lift.

Proof. An event evaluation has a testing-field sum-of-squares factorization from its projector, and a nonnegative coefficient contributes its source-field square root. The zero source and nonnegative constants are included by event normalization.

For a closed-face source use its explicit homogeneous matrix lift. The displayed strict feasible point, with all free coefficient sign blocks included, makes the adjoint image of the positive semidefinite cone closed. To see the attainment step, a strict feasible matrix $M(z_*) \succeq \epsilon I$ bounds the trace of any positive semidefinite dual witness whose adjoint image stays bounded. Separation then gives an attained matrix witness for every nonnegative evaluation. This fixed finite existential statement transfers to the testing field, independently of source feasibility in the coefficient field. Factoring the two positive semidefinite matrices there gives

$$\sum_{i,j} X_{ij} \otimes Y_{ij} = \sum_{a,b} \left(\sum_i B_{ia} \otimes C_{ib} \right)^2, \qquad X = BB^T, \quad Y = CC^T. \tag{FA10}$$

The dual equations cancel all auxiliaries and identify this expression with the original face slack. Singular matrices are allowed in these finite field factorizations.

For a canonical contact source apply LC to each real pure testing atom. If $(s \otimes 1)\mathcal{F} = \sum_k h_k^2$ with $s > 0$ in K_c, then (FA6) gives the exact unmultiplied certificate

$$[f, \Phi(z)] = \sum_k \left[(s^{-1/2} \otimes (\Delta_A \Delta_B \|\psi\|^2)^{-1/2}) h_k \right]^2. \tag{FA11}$$

All inverses and radicals in this formula are taken inside one field. Every additional source or testing gauge is restored in the same way, after its own nonvanishing has been proved. Finite real Bell relabelings only change coordinates and preserve the assertion.

Use the finite source decomposition and the nine-atom testing decomposition to write

$$f = \sum_j a_j f_j, \qquad q = \sum_i w_i \Phi(z_i), \qquad a_j, w_i \geq 0, \quad \sum_i w_i = 1.$$

If $[f_j, \Phi(z_i)] = \sum_k h_{ijk}^2$, then

$$[f, q] = \sum_{i,j,k} \left[(\sqrt{a_j} \otimes \sqrt{w_i}) h_{ijk} \right]^2. \tag{FA12}$$

Linearity in f and affineness in q prove this identity. The equation $\sum_i w_i = 1$ preserves the affine constant. Zero weights produce zero roots and are never inverted. The real/imaginary state split and finite convexification already include complex and mixed states. This proves (FA9).

Now choose two labelled copies of one arbitrary real closed extension K of $\mathbb{R}$. Formula (FA9) gives the hypothesis of Corollary 3.19 in Claus Scheiderer's *Second-order cone representation for convex sets in the plane*, arXiv:2004.04196v2. That corollary applies to a closed convex semialgebraic subset of arbitrary finite-dimensional $\mathbb{R}^n$. Its point-first tensor convention is equivalent to (FA3) by the factor flip, which preserves squares. The physical reduction proves all of its set hypotheses, and (FA2) identifies its transferred valid-affine cone. It therefore gives finite semidefinite-lift existence. □

Explicitly, the existence conclusion has the form

$$\mathcal{Q}_{222} = \left\{ q \in \mathbb{R}^8 : \ \exists y \in \mathbb{R}^m, \quad A_0 + \sum_{\nu=1}^{8} q_\nu A_\nu + \sum_{j=1}^{m} y_j B_j \succeq 0 \right\} \tag{FA13}$$

for finite integers N, m and real symmetric $N \times N$ matrices. Several semidefinite blocks can be combined diagonally. Linear equalities in a lift can be eliminated by parametrization or represented by opposite scalar blocks. These operations give (FA13) in the original eight coordinates.

Scope, reading order, and verification

The existence criterion does not evaluate the dimension, a global feature list, or the matrices in (FA13). The three constructive checklist items E1–E3 remain additional outputs: an evaluated feature and auxiliary-variable list, an evaluated pencil and projection, and both-inclusion validation of that actual coefficient data. No numerical lift-size bound or second-order-cone representation is asserted here.

The reader can follow the mathematical argument in this order: the physical model and separated algebra; source coercivity, closed faces and native entry; the positive-scale and noncommuting-testing exits; the complete C1 and C2 branches and their full source/test joins; finally the finite closure (FA10)–(FA12) and the external criterion. The component index identifies every actual body and its source path. Displayed formula tags retained from a component are local to that component. The global exposition and dependency records specify the component when using such a tag elsewhere.

Retained source files may contain a statement that their own local result does not prove an entire global theorem, or may describe the remaining scope at the time they were written. Those qualifications are preserved; a later proved union is required to discharge them. Historical status prose, review records and symbolic assertion counts are never mathematical premises. The archive binds each body and each presentation by its own hash and preserves the originals.

Verification consists of analytical review, independent review by other agents within the same model system, targeted exact symbolic corroboration, dependency and file-integrity checks, and PDF layout review. It is not external peer review or proof-assistant formal verification. The runnable verification program distinguishes file integrity from optional isolated algebra replays; neither substitutes for the printed analytical arguments.

Primary external source. Claus Scheiderer, *Second-order cone representation for convex sets in the plane*, arXiv:2004.04196v2, Corollary 3.19, `https://arxiv.org/pdf/2004.04196`.

Chapter 1

The elementary mechanism: positive residue pays for errors

Component `p25af2c58da7e`
Source: `CORE/bridge/strict_gram.tex`
SHA256: `25af2c58da7e1b0dc18b12c88178f434de24a96664d2dc036f312e919628bd1e`
Inventory use: lem:residue; lem:gram | Separated bounded strict-Gram theorem | positive-residue and strict bounded-Gram lemmas; lem:residue; lem:gram

1.1 The elementary mechanism: positive residue pays for errors

Take real closed fields $K_c, K_t \supset \mathbb{R}$. Let B_c, B_t be their rings of elements bounded in absolute value by a real number, with ideals $\mathfrak{m}_c, \mathfrak{m}_t$ of infinitesimals. Every bounded element has a unique real residue. Define

$$T = K_c \otimes_{\mathbb{R}} K_t, \quad B = B_c \otimes_{\mathbb{R}} B_t, \quad \mathfrak{n} = \mathfrak{m}_c \otimes B_t + B_c \otimes \mathfrak{m}_t.$$

The residue map $B \to \mathbb{R}$ has kernel $\mathfrak{n}$. No order is assigned to T.

Lemma 1.1 (Positive residue)**.** *If $a \in B$ and $\operatorname{res}(a) > 0$, then $a \in \Sigma T^2$.*

Proof. Expanding each bounded factor into its real residue plus an infinitesimal writes

$$a = \varepsilon + \sum_{j=1}^{N} x_j \otimes y_j, \qquad \varepsilon > 0,$$

with all factors bounded and at least one infinitesimal in every term. Divide ε into positive real reserves ε_j. If x_j is infinitesimal, choose real $M_j > |y_j|$. Exactly

$$\varepsilon_j + x_j \otimes y_j = \frac{1}{2}\left[(\varepsilon_j/M_j + x_j) \otimes (M_j + y_j) + (\varepsilon_j/M_j - x_j) \otimes (M_j - y_j)\right].$$

Each of the four pure factors is positive. Their separate field square roots turn each elementary tensor into a square. Swap the two factors when required. The case $N = 0$ is immediate. □

Lemma 1.2 (Strict bounded Gram)**.** *If $Q \in \operatorname{Sym}_r(B)$ has $\overline{Q} \succ 0$, then $g^T Q g \in \Sigma T^2$ for every $g \in T^r$. The entries of g need not be bounded.*

Proof. A real congruence makes $\overline{Q} = I$. Choose real $\delta > 0$ with $(r-1)\delta < 1$. The exact decomposition

$$\begin{aligned} Q = &\sum_i [Q_{ii} - (r-1)\delta] e_i e_i^T \\ &+ \sum_{i<j} \frac{\delta + Q_{ij}}{2} (e_i + e_j)(e_i + e_j)^T + \sum_{i<j} \frac{\delta - Q_{ij}}{2} (e_i - e_j)(e_i - e_j)^T \end{aligned}$$

has scalar coefficients of positive real residue. Lemma 1.1, followed by evaluation on g, proves the claim. □

Chapter 2

Fresh reconstruction of the physical model and conditional assembly

Component `p9b00698ff36a`
Source: `REC/routing/fresh_physical_normalization_conditional_assembly.tex`
SHA256: `9b00698ff36aab2f233a0d2932a439ad7fb145384d10545302ecb1441513394d`
Inventory use: Fresh physical real-qubit/finite-convexification model, exact normalization and separated matrix-pairing assembly | RFA physical-reduction/normalization/closure portions including RFA10–RFA12

2.1 Fresh reconstruction of the physical model and conditional assembly

Status of this source. This is a newly derived proof, not a transcription of the missing final R21 manuscript. The physical-model theorem, the homogeneous normalization identity, the separated conic/convex closure lemma, and the matrix-pairing lemma below are proved here. The final existence implication is expressly conditional on the local-certificate hypothesis (LC). Neither a historical admission nor a remembered checksum is used to discharge (LC).

2.1.1 An exact finite-dimensional physical model

Let $Q \subset \mathbb{R}^8$ be the set of correlations obtained from arbitrary finite-dimensional complex bipartite tensor-product states and two binary measurements per party. The coordinates are

$$(a_0, a_1, b_0, b_1, E_{00}, E_{01}, E_{10}, E_{11}). \tag{RFA1}$$

A binary measurement is represented by an effect $0 \preceq E \preceq I$, with observable $2E - I$. Put $Z = \mathrm{diag}(1, -1)$ and $X = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$. For $(c_A, s_A), (c_B, s_B) \in S^1$ and $\psi \in S^3 \subset \mathbb{R}^4$, let Φ be the eight expectations of

$$A_0 = B_0 = Z, \qquad A_1 = c_A Z + s_A X, \qquad B_1 = c_B Z + s_B X \tag{RFA2}$$

in the state ψ. Every coordinate of Φ is polynomial in these eight real parameters.

Theorem 2.1 (Fresh physical reduction)**.** *With $\mathcal{Z} = S^1 \times S^1 \times S^3$,*

$$Q = \mathrm{conv}\, \Phi(\mathcal{Z}) = \left\{ \sum_{i=1}^{9} w_i \Phi(z_i) : z_i \in \mathcal{Z},\ w_i \geq 0,\ \sum_i w_i = 1 \right\}. \tag{RFA3}$$

In particular Q is compact, convex and semialgebraic over the real field.

Proof. First, both effects of a party can be dilated with the same embedding. For an effect E on $\mathcal{H}$ define, on $\mathcal{H} \oplus \mathcal{H}$,

$$\widehat{P}_E = \begin{pmatrix} E & \sqrt{E(I-E)} \\ \sqrt{E(I-E)} & I - E \end{pmatrix}, \qquad Vh = (h, 0). \tag{RFA4}$$

The square root commutes with E by its spectral construction. Direct block multiplication gives $\widehat{P}_E^2 = \widehat{P}_E$ and $V^* \widehat{P}_E V = E$. Thus for both measurement settings the same V compresses the two projections to the original effects. Applying this construction to each party and replacing ρ by $(V_A \otimes V_B)\rho(V_A \otimes V_B)^*$ preserves every marginal and joint expectation. We may therefore assume the two observables of each party are Hermitian involutions.

Here is the needed two-dimensional block decomposition. If U, V are Hermitian involutions, $C = UV$ is unitary and $UCU = C^*$. For an eigenvector e of C with nonreal eigenvalue λ, Ue has eigenvalue $\overline{\lambda}$. The vectors are orthogonal and their span is invariant under U and $V = UC$. Its orthogonal complement is invariant too. Repeating on orthogonal eigenvectors gives two-dimensional blocks. On the real eigenvalue spaces $\lambda = \pm 1$ one has $V = \pm U$; diagonalizing U gives one-dimensional blocks. This proves the decomposition in finite dimension without an assumption on multiplicities.

On a two-dimensional block U has eigenvalues $1, -1$. In its eigenbasis, the other nontrivial involution has the form $\left(\begin{smallmatrix} c & \overline{z} \\ z & -c \end{smallmatrix}\right)$, where c is real and $c^2 + |z|^2 = 1$. A diagonal change of phase leaves $U = Z$ and makes z real. Thus the pair has the form $Z, cZ + sX$. A one-dimensional block with outcomes $a_0, a_1 \in \{\pm 1\}$ embeds into this form: use the a_0 eigenvector of Z and take $c = a_0 a_1, s = 0$. The unused basis vector does not affect the experiment.

Apply both local decompositions. All measurement operators are block diagonal, so their expectations depend only on the diagonal blocks of the state. The traces of these positive blocks are nonnegative weights summing to one. Every nonzero block, after normalization and the harmless one-dimensional embedding, is a two-qubit state with real measurement matrices. Decompose that state into finitely many complex pure states. For a real symmetric matrix M and a vector $r + is$,

$$(r + is)^* M (r + is) = r^T M r + s^T M s, \qquad \|r + is\|^2 = \|r\|^2 + \|s\|^2. \tag{RFA5}$$

The cross terms cancel by symmetry. The nonzero real and imaginary parts, normalized separately, are real pure states with their squared norms as weights. This proves $Q \subseteq \operatorname{conv} \Phi(\mathcal{Z})$.

Conversely each atom is a physical two-qubit experiment. A finite convex combination is realized by a finite direct sum on each party and a state with the corresponding classical diagonal weights in matching summands. Therefore the reverse inclusion holds.

Any convex combination of more than nine points in $\mathbb{R}^8$ can be shortened. The augmented vectors $(1, q_i) \in \mathbb{R}^9$ are linearly dependent, so there are α_i, not all zero, with $\sum_i \alpha_i = 0$ and $\sum_i \alpha_i q_i = 0$. Change their common sign if needed so that some $\alpha_i > 0$, and replace the positive weights w_i by $w_i - \tau \alpha_i$, where $\tau = \min_{\alpha_i > 0} w_i / \alpha_i$. They remain nonnegative, keep the same sum and point, and at least one vanishes. Iterate. Zero weights allow a fixed list of nine atoms. The right side of (RFA3) is the continuous image of the compact product of a simplex and nine copies of $\mathcal{Z}$, hence is compact. It is semialgebraic by the projection theorem for real closed fields applied to its displayed finite polynomial formula. Convexity has already been proved. □

For any real closed extension $K \supset \mathbb{R}$, define Q_K by (RFA3) with all variables in K. Define $P = \{f \in \mathbb{R}[q]_{\leq 1} : f \geq 0 \text{ on } Q\}$. The defining formula $\forall q\, (q \in Q \Rightarrow f(q) \geq 0)$, together with quantifier elimination and elementary transfer, gives exactly

$$P_K = \{f \in K[q]_{\leq 1} : f \geq 0 \text{ on } Q_K\}. \tag{RFA6}$$

Compactness was used only over $\mathbb{R}$. No compactness assertion for the order topology of a non-Archimedean field is needed.

2.1.2 Separated fields and the full homogeneous parent

Fix arbitrary real closed extensions $K_c, K_t \supset \mathbb{R}$ and put $T = K_c \otimes_{\mathbb{R}} K_t$. The factors are labelled; they may be two copies of the same extension. There is no assumption of algebraic disjointness and no chosen ordering on T. For $f = f_0 + \sum_{\nu=1}^{8} f_\nu q_\nu$ define

$$\mathscr{S}(f, q) = f_0 \otimes 1 + \sum_{\nu=1}^{8} f_\nu \otimes q_\nu. \tag{RFA7}$$

The event and CHSH convention in the recovered physical core is

$$S = E_{00} - E_{01} + E_{10} + E_{11}, \quad H = 1 - S/2, \quad f = dH + 4\sum_{\xi} \xi\, p_\xi, \quad \xi \in \{\lambda, n, m, C, A, D, \alpha, \beta\}. \tag{RFA8}$$

Its original full parent is recovered as actual source text, not merely a remembered identity. We give its homogeneous form explicitly.

Choose projective angle pairs (a_0, a_1) and (b_0, b_1) in K_t^2, each nonzero, and set $\Delta_A = a_0^2 + a_1^2 > 0$, $\Delta_B = b_0^2 + b_1^2 > 0$. The corresponding circle coordinates are

$$c_A = (a_0^2 - a_1^2)/\Delta_A, \quad s_A = 2a_0a_1/\Delta_A, \tag{RFA9}$$

and similarly for Bob. These cover every circle point over K_t: if $c \neq -1$ use $(1 + c, s)$, and if $c = -1$ use $(0, 1)$. No affine half-angle chart is therefore needed at infinity. For $\Psi = (W, X, Y, Z)$ put

$$\begin{aligned} L_m &= a_0b_1W - a_0b_0X + a_1b_1Y - a_1b_0Z, \\ L_D &= a_1b_0W + a_1b_1X - a_0b_0Y - a_0b_1Z. \end{aligned}$$

Products of a coefficient-field element with a testing-field expression in the next identity mean their separated tensor product. The complete homogeneous parent is

$$\begin{aligned} \mathcal{F} = {} & 4\lambda\Delta_A(b_0W + b_1X)^2 + 4n\Delta_A\Delta_BX^2 + 4mL_m^2 \\ & + 4C\Delta_B(a_1W - a_0Y)^2 + 4A\Delta_A\Delta_BY^2 \\ & + (4D + 2d)L_D^2 + (4\alpha + 2d)\Delta_A(b_1Y - b_0Z)^2 \\ & + (4\beta + 2d)\Delta_B(a_0X + a_1Z)^2 - 2d\Delta_A\Delta_BZ^2. \end{aligned} \tag{RFA10}$$

For a nonzero state let $N = \|\Psi\|^2 > 0$ and let z be the atom with the circle coordinates (RFA9) and unit state $\Psi/\sqrt{N}$. Then

$$\mathcal{F} = (1 \otimes \Delta_A\Delta_BN)\, \mathscr{S}(f, \Phi(z)). \tag{RFA11}$$

To check the normalization, first take $a_0 = b_0 = 1$, $u = a_1$ and $v = b_1$. The plus/minus measurement eigenvectors are $(1, u)/\sqrt{1 + u^2}$ and $(u, -1)/\sqrt{1 + u^2}$. Their product amplitudes give, in the order of (RFA8),

$$\begin{gathered} \frac{(W + vX)^2}{1 + v^2}, \quad X^2, \quad \frac{(vW - X + uvY - uZ)^2}{(1 + u^2)(1 + v^2)}, \quad \frac{(uW - Y)^2}{1 + u^2}, \quad Y^2, \\ \frac{(uW + uvX - Y - vZ)^2}{(1 + u^2)(1 + v^2)}, \quad \frac{(vY - Z)^2}{1 + v^2}, \quad \frac{(X + uZ)^2}{1 + u^2}. \end{gathered} \tag{RFA12}$$

For a normalized state these are exactly the eight selected event probabilities. For a general state they are multiplied by N. Substitution of the matrices (RFA2) into H gives its numerator

$$2\{(uW + uvX - Y - vZ)^2 + (1 + u^2)(vY - Z)^2 + (1 + v^2)(X + uZ)^2 - (1 + u^2)(1 + v^2)Z^2\}.$$

This proves (RFA11) in the affine chart. Homogenization gives (RFA10); it is a polynomial identity and therefore also gives the zero and infinite projective coordinates without dividing by a_0 or b_0. The zero state has zero parent separately.

If $\mathcal{F}$ or $(s \otimes 1)\mathcal{F}$ has an exact finite SOS certificate, with $s \in K_c^{>0}$, the original slack is restored by multiplying each root by

$$s^{-1/2} \otimes (\Delta_A\Delta_BN)^{-1/2}. \tag{RFA13}$$

All these inverses are taken within one field before forming a tensor. The identity does not permit cancellation of an arbitrary mixed tensor. Additional positive source gauges are handled in the same way. A state chart or a change of frame must first prove its own denominator nonzero.

Literal commuting tests. These require no limiting local certificate. If one party's two involutions commute, on a simultaneous eigenspace the CHSH operator $S = A_0(B_0 - B_1) + A_1(B_0 + B_1)$ is either $\pm 2B_0$ or $\pm 2B_1$. Thus $H = I - S/2$ is positive semidefinite in the testing field. For $d \geq 0$, its contribution is a pure-source square times a pure-testing square. Every nonnegative event packet is also separated SOS. This covers the actual commuting angle axes and infinities for every nonnegative canonical coefficient tuple. It does not cover merely infinitesimal nonzero angles, which remain in (LC).

2.1.3 The closed faces and the surviving source reduction

Two recovered sources supply actual mathematical inputs: `physical_core.tex` gives the canonical event basis, the determinant/coercivity argument, the finite-contact extreme-source reduction and the nine-term source decomposition; `closed_faces_short.tex` gives the two explicit matrix lifts for the omitted exclusive pairs. Their exact recovered paths and hashes are recorded in the reconstruction map. Their roles are not supplied by the historical R21 review narrative.

We state the precise source-generation conclusion used below. Every transferred valid affine functional is either an event sum or a finite nonnegative conic combination of the following types, with finite real Bell relabelings restored:

- event sources and noncoercive closed-face sources;
- physical C1 sources with $\lambda = n = 0$, $d, m, C, A, \alpha > 0$, $D, \beta \geq 0$, $D + \beta > 0$, and an actual finite positive noncommuting stationary contact;
- physical C2 sources with $\lambda = \beta = 0$, $d, m, C, A, D, \alpha > 0$, $n \geq 0$, and the same actual-contact condition.

The source sum is formed before choosing $d = 1$ or $C = 2$. This is a source-field assertion; source contacts are not testing points.

For completeness, the logical content of the recovered reduction is as follows. A functional nonnegative on the nonsignaling polytope is a nonnegative combination of its sixteen event inequalities. Otherwise it is negative at a PR vertex, because deterministic vertices are physical. The explicit nonsignaling vertex classification and the nine evaluations in the recovered event table give (RFA8) with nonnegative event coefficients and $d > 0$. Normalize their total sum to one. The physical coefficient section is compact and convex in a space of affine dimension at most eight and hence is a convex combination of at most nine extreme points. The recovered determinant has nonnegative outer Laurent coefficients. Its coercive cases have a finite positive minimum; the printed opposite-perturbation argument makes an extreme minimum zero. The four kernel and two angular equations leave a nonradial perturbation if more than six event coefficients are active. The complementary state pivots, exact adjugate identity and convex Laurent profile preserve physicality under that perturbation. Coercivity requires at least five active events by the explicit exclusive-pair classification. Compatible omissions have precisely the two C1/C2 orientations above; every noncoercive support lies on one of the two closed faces. The finite source decomposition and these contact assertions have fixed first-order formulas, so their real proofs transfer. This is the actual content required of the recovered source proof; merely naming a C1 or C2 row would not suffice.

Here is a direct separated-tensor consequence of the recovered closed-face lifts. Write a given lift, including all free nonnegative source-coordinate diagonal blocks, as the homogeneous matrix

$$M(c, y) = \sum_{\nu} c_\nu A_\nu + \sum_{j} y_j B_j \succeq 0.$$

The supplied matrices have an explicit strict feasible point in the free source coordinates and auxiliaries. If a linear functional $\ell(c)$ is nonnegative on the projected cone, there exists $Y \succeq 0$ with

$$\operatorname{tr}(A_\nu Y) = \ell_\nu, \qquad \operatorname{tr}(B_j Y) = 0. \tag{RFA14}$$

One can prove this without leaving an attainment issue implicit. For the linear map M, its adjoint image $M^*(\mathrm{PSD})$ is closed: at a fixed strict point z_*, $M(z_*) \succeq \epsilon I$, so convergence of $M^*(Y_i)$ for $Y_i \succeq 0$ bounds $\epsilon \operatorname{tr}(Y_i)$ and thus bounds Y_i. A convergent subsequence supplies the limiting witness. Separation of this closed cone proves that its dual is exactly $\{z : M(z) \succeq 0\}$, yielding (RFA14).

Apply this over the real field with ℓ equal to a testing evaluation. The fixed existential statement for Y transfers to K_t; source feasibility transfers independently to K_c. For a source witness $X = M(c, y) \succeq 0$, factor $X = BB^T$ in K_c and $Y = CC^T$ in K_t. Positive semidefinite matrices over a real closed field have such finite factorizations, including singular matrices, by symmetric diagonalization or the pivoted quadratic-form algorithm. Then

$$\sum_{i,k} X_{ik} \otimes Y_{ik} = \sum_{a,b} \left(\sum_i B_{ia} \otimes C_{ib} \right)^2 . \tag{RFA15}$$

The adjoint equations (RFA14) cancel the auxiliary variables and identify the left side with the original slack. This proves the closed-face tensor conclusion from the actual explicit face lifts.

Event slacks also need no local certificate: for a real unit atom, the joint event projector R satisfies $p = \psi^T R \psi = \|R\psi\|^2$ over K_t. Its entries are testing-field elements, so this is a finite pure-testing SOS. A nonnegative source weight contributes its own square root. Constants and the zero functional are included because outcomes of one input pair sum to one.

2.1.4 The missing local premise, stated without a status substitution

Local-certificate hypothesis (LC). For every pair of real closed extensions K_c, K_t, every physical canonical C1 or C2 source in the source-generation statement above, and every independent projective angle pair and testing state, the complete original parent (RFA10) has a finite SOS certificate in T, possibly after multiplication by a proved strictly positive pure-source factor. Every source coefficient face and gap, all testing signs, all state coordinates, and all projective endpoints must be included. When local charts are used, their union and restoration must be proved.

The reconstruction does not yet establish (LC). In particular the missing positive-scale completion, original weak-scale entry/scoping proofs, uniform-C1 local replacements and the C14 and weak-C2 local closures cannot be replaced by their historical admission summaries. The accompanying map separates exact surviving inputs, alternate candidate proofs, fragmentary statements and absent certificate bodies.

One sufficient way to prove (LC) is to restore a finite implication tree, valid for every actual source and test, whose leaves are proved full-parent certificates or proved impossible domains. The branch predicates must carry the actual source equations and literal boundaries. This description is a proof obligation, not an assertion that the missing tree has been recovered.

2.1.5 Conditional conic/convex closure and finite-lift implication

Lemma 2.2 (Fresh separated assembly)**.** *Let $f = \sum_{j=1}^r a_j g_j$ in the source field, with $a_j \geq 0$, and let $q = \sum_{i=1}^s w_i q_i$ in the testing field, with $w_i \geq 0$ and $\sum_i w_i = 1$. If every elementary slack has a finite certificate $\mathscr{S}(g_j, q_i) = \sum_k h_{ijk}^2$ in T, then*

$$\mathscr{S}(f, q) = \sum_{i,j,k} \left((\sqrt{a_j} \otimes \sqrt{w_i}) h_{ijk} \right)^2 . \tag{RFA16}$$

Proof. Affineness in the testing point, including its constant term, gives $\mathscr{S}(f, q) = \sum_i (1 \otimes w_i) \mathscr{S}(f, q_i)$. Linearity in the source gives $\mathscr{S}(f, q_i) = \sum_j (a_j \otimes 1) \mathscr{S}(g_j, q_i)$. Substitute the finite certificates. The required nonnegative square roots exist in the separate real closed fields. Zero weights give zero roots; no weight is inverted. This proves the displayed identity in the tensor ring, without identifying its two factors. □

Theorem 2.3 (Fresh conditional universal theorem)**.** *Assume (LC), with the actual source reduction and explicit closed-face lifts specified above. Then for all real closed $K_c, K_t \supset \mathbb{R}$, all $f \in P_{K_c}$ and all*

$q \in Q_{K_t}$, $\mathscr{S}(f,q)$ is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$. Consequently Q has a finite semidefinite lift.

Proof. Use the finite source decomposition. Event and closed-face slacks were proved above. For every other source atom apply (LC), then (RFA11)–(RFA13) to restore its exact affine slack. Use the nine-atom testing decomposition (RFA3) and the assembly lemma. Real/complex states, mixed states, the zero functional and zero weights were already included.

For the final implication take two labelled copies of an arbitrary single real closed extension K. The set and valid-affine cone are exactly their transfers by (RFA3) and (RFA6). The resulting SOS is in $K \otimes_{\mathbb{R}} K$. The factor flip converts the coefficient-first convention to the point-first convention of Claus Scheiderer's Corollary 3.19, which applies to a closed convex semialgebraic set in arbitrary finite ambient dimension. It proves finite semidefinite-lift existence. □

Exact boundary of the new conclusion. The last theorem is conditional until actual proofs establish (LC). No unconditional finite-lift conclusion is claimed by this reconstruction. Even after (LC) is proved, this criterion gives existence, not evaluated matrices, a numerical size bound, or a common separation-rank bound. The explicit feature, affine-pencil and coefficient-validation items E1–E3 remain subsequent constructive tasks.

Primary external criterion. C. Scheiderer, *Second-order cone representation for convex sets in the plane*, arXiv:2004.04196v2, Corollary 3.19 and its proof, printed page 15, `https://arxiv.org/pdf/2004.04196`. The primary text has been reopened during this reconstruction. Theorem 3.10 in the same source distinguishes a common tensor-rank bound from unrestricted finite SOS. Neither local square counts nor the number of routing branches supplies that stronger bound.

Chapter 3

The physical geometry of the canonical reduction

Component p0c8f420148b7
Source: CORE/prior/inherited/physical_core.tex
SHA256: 0c8f420148b721c9d0ccafdd15263523975398f12509e15d57387f8762cf2e1a
Inventory use: sec:physical; eq:parent; determinant coefficient dictionary; explicit g_A,g_B,g_T maps | Full parent, event and determinant dictionary | canonical parent P and physical source geometry; sec:physical; eq:parent; determinant coefficient dictionary; explicit g_A,g_B,g_T maps

3.1 The physical geometry of the canonical reduction

Fix the CHSH orientation

$$S = E_{00} - E_{01} + E_{10} + E_{11}, \qquad H = 1 - S/2,$$

with PR point $(0, 0, 0, 0, 1, -1, 1, 1)$. The eight losing events, in the parent order, are as follows. The last column is their unique deterministic neighbor on $S = 2$.

Coefficient	Event $(a, b \mid x, y)$	Assignment (a_0, a_1, b_0, b_1)
λ	$(+, + \mid 0, 1)$	$(+, +, +, +)$
$n = e_1$	$(+, - \mid 0, 0)$	$(+, -, -, -)$
$m = e_2$	$(+, - \mid 1, 1)$	$(+, +, +, -)$
$c = e_3$	$(-, + \mid 1, 0)$	$(+, -, +, -)$
$A = e_4$	$(-, + \mid 0, 0)$	$(-, +, +, +)$
$D = e_5$	$(-, + \mid 1, 1)$	$(-, -, -, +)$
α	$(-, - \mid 0, 1)$	$(-, -, -, -)$
β	$(+, - \mid 1, 0)$	$(-, +, -, +)$

Set $E_i = 4p(e_i)$, explicitly fixing the event normalization. At the PR point, $H = -1$ and every $E_i = 0$; at neighbor v_j, $H = 0$ and $E_i = 4\delta_{ij}$. These nine evaluations prove that $H, E_1, \ldots, E_8$ form an affine basis. Consequently

$$f = dH + \sum_i c_i E_i, \qquad d = -f(\mathrm{PR}), \quad c_i = f(v_i)/4.$$

For a quantum-valid support negative at this PR point, $d > 0$ and all $c_i \geq 0$.

For completeness, the nonsignaling vertices are the sixteen deterministic points and eight PR points. If a local marginal is deterministic, condition on the other measurement at that site and sample the two remote outcomes with their observed conditional marginals; this constructs a local model. Such an extreme point must be deterministic. If all marginals are strictly nondeterministic, a two-by-two joint table has at most two zero cells. A vertex in the eight-dimensional nonsignaling affine space needs eight independent active positivity constraints, so each table has exactly two zeros

and fixes perfect correlation or anticorrelation. Even parity of these four signs leaves a free marginal; odd parity fixes unbiased marginals and gives a PR point. Thus any quantum-valid support not valid on the full nonsignaling polytope is negative at some PR point.

3.1.1 Every parent packet is a measurement amplitude

Use half-angle coordinates

$$A_1 = \frac{(1-u^2)\sigma_z + 2u\sigma_x}{D_u}, \quad B_1 = \frac{(1-v^2)\sigma_z + 2v\sigma_x}{D_v}, \quad D_u = 1+u^2,\ D_v = 1+v^2.$$

The local plus and minus eigenvectors are $(1,u)/\sqrt{D_u}$ and $(u,-1)/\sqrt{D_u}$, respectively, and similarly for Bob. For a normalized state (W,X,Y,Z), the eight event probabilities are

$$\begin{aligned}
p_\lambda &= \frac{(W+vX)^2}{D_v}, & p_n &= X^2,\\
p_m &= \frac{(vW - X + uvY - uZ)^2}{D_u D_v}, & p_c &= \frac{(uW-Y)^2}{D_u},\\
p_A &= Y^2, & p_D &= \frac{(uW + uvX - Y - vZ)^2}{D_u D_v},\\
p_\alpha &= \frac{(vY-Z)^2}{D_v}, & p_\beta &= \frac{(X+uZ)^2}{D_u}.
\end{aligned}$$

These formulas can be checked one entry at a time from the product eigenvectors. Direct Pauli multiplication gives

$$\begin{aligned}
D_u D_v \langle H\rangle = 2\big[&(uW + uvX - Y - vZ)^2 + D_u(vY-Z)^2\\
&+ D_v(X+uZ)^2 - D_u D_v Z^2\big].
\end{aligned}$$

The complete homogeneous parent is therefore

$$\begin{aligned}
F = {}& 4\lambda D_u(W+vX)^2 + 4nD_uD_vX^2 + 4m(vW - X + uvY - uZ)^2\\
&+ 4cD_v(uW-Y)^2 + 4AD_uD_vY^2\\
&+ (4D+2d)(uW + uvX - Y - vZ)^2 + (4\alpha+2d)D_u(vY-Z)^2\\
&+ (4\beta+2d)D_v(X+uZ)^2 - 2dD_uD_vZ^2. \qquad \text{(P)}
\end{aligned}$$

For normalized states, $F = D_uD_vf(q)$. For arbitrary nonzero states, divide by the additional positive testing factor $\|\psi\|^2$. This fixes the signs, the factor four and the restoration factor.

If $d > 0$ and $uv \neq 0$, the signed part restricted to $Z = 0$ contains

$$2d\big[(uW + uvX - Y)^2 + D_u v^2 Y^2 + D_v X^2\big],$$

which is positive definite. Thus the parent matrix P has at least three positive eigenvalues, and $P \succeq 0$ is equivalent to $\det P \geq 0$. Commuting angle axes and projective ends have $H \succeq 0$ directly: when one party's reflections commute, their common eigenspaces reduce CHSH to ± 2 times one remote reflection. The case $d = 0$ is an event-square sum.

3.1.2 A Newton square and classical consistency

Define

$$\begin{aligned}
t_1 &= d + 2(\lambda + n + D), & t_2 &= d + 2(\alpha + m + A),\\
t_3 &= d + 2(\lambda + c + D), & t_4 &= d + 2(\alpha + \beta + m),\\
t_5 &= d + 2(\lambda + \beta + n), & t_6 &= d + 2(\alpha + c + A),
\end{aligned}$$

$$t_7 = d + 2(\beta + n + m), \qquad t_8 = d + 2(c + A + D).$$

The exact determinant identity is

$$\det P = 16 D_u^2 D_v^2 \sum_{i,j=0}^{2} q_{ij} u^{2i} v^{2j},$$

where

$$\begin{aligned} q_{00} &= 4\lambda\alpha t_7 t_8, & q_{01} &= 4(\lambda D t_7 t_6 + \alpha m t_5 t_8), \\ q_{02} &= 4 m D t_5 t_6, & q_{10} &= 4(\alpha A t_7 t_3 + \lambda n t_8 t_4), \\ q_{12} &= 4(\beta m t_1 t_6 + c D t_5 t_2), & q_{20} &= 4 n A t_3 t_4, \\ q_{21} &= 4(\beta n t_3 t_2 + c A t_1 t_4), & q_{22} &= 4\beta c t_1 t_2. \end{aligned}$$

The remaining coefficient is explicitly $h = q_{11} = \det P(1,1)/256 - \sum_{(i,j)\neq(1,1)} q_{ij}$. This identity is a finite 4×4 determinant calculation from (P); it can also be expanded by Cauchy–Binet from its eight positive rank-one packets and one negative packet. The complete polynomial identity was reconstructed exactly in the companion checks; that check verifies the displayed identity and does not replace the following argument.

For $U = u^2, V = v^2 > 0$, divide by UV:

$$L(U,V) = h + \sum_{(i,j)\neq(1,1)} q_{ij} U^{i-1} V^{j-1}.$$

All eight outer coefficients are nonnegative. In logarithmic variables this is a constant plus positive exponentials. It is convex and is coercive exactly when zero is interior to the convex hull of its active exponent vectors. If zero is interior, the largest exponent has a uniform positive slope in every unit direction; the exponential sum tends to infinity. Otherwise a separating direction makes every term nonincreasing along a ray.

Proposition 3.1 (Coercivity is compatibility of omitted events)**.** *For $d > 0$, the logarithmic Laurent polynomial is coercive if and only if the omitted losing events are pairwise nonexclusive, meaning they never assign opposite outcomes to the same local measurement.*

Proof. The available exponents are the eight boundary points of the 3×3 square. The separating directions can be chosen among $(\pm 1, 0), (0, \pm 1), (\pm 1, \pm 1)$: directions between two consecutive such rays give the same signs, and a boundary ray only weakens the vanishing conditions.

For $(1, 0)$, the forbidden half-plane requires $\lambda\alpha = \lambda D = \alpha m = mD = 0$. Its minimal zero sets are $\{\lambda, m\}$ and $\{\alpha, D\}$. The four axis directions give

$$(\lambda, m), (\alpha, D), (n, c), (A, \beta), (\lambda, A), (n, \alpha), (m, c), (D, \beta).$$

The diagonal $(1, 1)$ has minimal zero sets $\{\lambda, m, A\}, \{\lambda, \alpha\}, \{n, D, \alpha\}$. The first and last already contain an axis zero set, leaving only (λ, α). The remaining diagonals add $(m, D), (n, A), (c, \beta)$. These twelve pairs are exactly the exclusive pairs in the event table. Hence failure of coercivity is equivalent to omitting an exclusive pair. □

Pairwise nonexclusive events give a consistent partial assignment to the four local measurements, so extend to a deterministic assignment. A deterministic assignment loses one or three CHSH inputs, never four. Thus there are at most three compatible omitted losing events. No list of support masks is needed: omission of an exclusive pair characterizes the noncoercive case, and Section 13.1 gives explicit lifts for its two symmetry types.

3.1.3 Extreme sources make finite contacts

Normalize the nonnegative source vector by $d + \sum c_i = 1$. Its physical subset is compact and convex over $\mathbb{R}$. For a coercive support, L attains its minimum at finite positive U, V.

Suppose a physical extreme ray had minimum $\mu > 0$. A small coefficient perturbation inside the exact active face changes each positive outer coefficient by relative error at most ε and h by absolute error at most ε. Uniformly over all angles,

$$L' \geq (1-\varepsilon)L + \varepsilon h - \varepsilon \geq (1-\varepsilon)\mu + \varepsilon h - \varepsilon > 0$$

for sufficiently small ε. Opposite nonradial perturbations remain physical, contradicting extremality. Thus an extreme coercive source has a finite noncommuting zero contact.

At that contact, a kernel vector $\psi \neq 0$ satisfies six linear equations in the source coefficients:

$$P\psi = 0, \qquad \psi^T(\partial_u P)\psi = 0, \qquad \psi^T(\partial_v P)\psi = 0.$$

If more than six event weights are positive, there are at least eight active coordinates including d. The nullspace of these six equations contains a nonradial perturbation. Small opposite perturbations retain all active signs and the same contact. The three complementary eigenvalues stay positive. Since the adjugate is a positive multiple of $\psi\psi^T$, the determinant remains stationary at zero; logarithmic convexity makes that zero its global minimum. The perturbed sources are physical, again contradicting extremality. No generic-rank assumption is used.

A coercive extreme support consequently has five or six active events. Its compatible omitted pair either shares one local setting or shares neither; these are the two relabeling classes

$$C_1 : \lambda = n = 0, \qquad C_2 : \lambda = \beta = 0.$$

For a finite check of transitivity, CHSH-preserving maps on (x, y, a, b) are

$$\begin{aligned} g_A &: (x, y, a, b) \mapsto (1-x, y, a, (-1)^y b), \\ g_B &: (x, y, a, b) \mapsto (x, 1-y, (-1)^{1-x} a, b), \\ g_T &: (x, y, a, b) \mapsto (y, x, (-1)^y b, (-1)^x a). \end{aligned}$$

On the cyclic event order $(\lambda, D, c, A, \alpha, m, \beta, n)$ these maps act as $i \mapsto 5-i, 3-i, 2-i$ modulo eight. Hence they generate a one-step rotation and a reflection. Compatible pairs have cyclic distance one or two; the rotations are transitive on each kind. Their representatives are precisely the two omitted pairs above, so no mask table is required. A compatible triple omission lies below one of these six-event faces by dropping one omission. Every other support is contained in a noncoercive omission face.

Finally, the normalized coefficient base has affine dimension at most eight, so every vector is a convex combination of at most nine extreme points. This conclusion, the midpoint definition of extremality, and finite contact attainment all admit fixed first-order formulas in real arithmetic. They transfer to every real closed coefficient field. Logarithms and real compactness were proof devices over $\mathbb{R}$ only. Nonnegative summands preserve literal event zeros. This completes the coefficient reduction to the closed faces and the two contact families; it does not yet prove their separated-field tensor certificates.

Chapter 4

Every noncommuting positive-d contact is regular

Component pe25c50379365
Source: CORE/prior/inherited/structural_terminals.tex
SHA256: e25c503793657fdc2ca93ef5832ce6b4eec9c9b8a8ad280698b03bcc396519d4
Inventory use: lem:regular; thm:curvature | unconstrained/support curvature and constrained terminals; lem:regular; thm:curvature | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

4.1 Every noncommuting positive-d contact is regular

The first simplification removes an entire kind of apparent degeneracy. It requires no chart census: at positive signed CHSH weight, physicality itself forces angular curvature.

Lemma 4.1 (Exact regular-contact certificate)**.** *Let $F(c,z) \in \mathbb{R}[c,z]$, with bounded $c \in B_c^m$, $z_t \in B_t^n$ and residues c_0, z_0. Suppose*

$$F(c_0,z_0) = 0, \quad \nabla_z F(c_0,z_0) = 0, \quad \nabla_z^2 F(c_0,z_0) \succ 0,$$

and $F(c,z) \geq 0$ for all $z \in K_c^n$. Then $F^{\otimes}(c,z_t) \in \Sigma T^2$.

Proof. The real implicit-function theorem gives a unique critical branch $z = \sigma(c)$ in fixed real neighborhoods. Its graph is semialgebraic: it is defined by the polynomial gradient equations with the neighborhood restrictions ensuring uniqueness. After shrinking, it satisfies a real Lipschitz bound $\|\sigma(c') - z_0\| \leq M \|c' - c_0\|$. This fixed first-order existence, uniqueness and bound statement transfers to K_c. Thus there is z_c of residue z_0 with $\nabla_z F(c,z_c) = 0$.

Put $a = F(c,z_c) \geq 0$ and $h_i = 1 \otimes z_{t,i} - z_{c,i} \otimes 1$. Finite polynomial Taylor expansion is exact:

$$F^{\otimes}(c,z_t) = a \otimes 1 + h^T Q h, \qquad Q \in \mathrm{Sym}_n(B), \quad \bar{Q} = \tfrac{1}{2}\nabla_z^2 F(c_0,z_0) \succ 0.$$

For an explicit allocation, give each monomial of degree at least two two of its h factors as roots, and leave its remaining factors in the Gram coefficient. Degree-one terms vanish. Every coefficient of degree at least three then has an infinitesimal factor. Symmetrize cross terms with a factor one half. The constant is $(\sqrt{a} \otimes 1)^2$; Lemma 1.2 handles the quadratic form. There is no infinite series or mixed inverse. □

Theorem 4.2 (Automatic angular curvature)**.** *Fix real source coefficients with $d > 0$ and all eight event coefficients nonnegative, and suppose the canonical parent is PSD at every pair of real angles. At every finite zero contact (u_0, v_0, ψ_0) with $u_0 v_0 \neq 0$ and $\psi_0 \neq 0$, the active outer Laurent exponents span $\mathbb{R}^2$. The Hessian in two angle coordinates and three projective state coordinates is positive definite.*

Proof. The transverse state block is positive definite by the signed-core formula in Section 3.1. At a zero contact, $\det P$ and its first angle derivatives vanish. The Laurent determinant therefore satisfies

$$\sum_{e\in E} w_e e = 0, \qquad w_e = q_e U^{e_1} V^{e_2} > 0.$$

If E has rank one, it is an opposite pair in the eight-point Newton square. Since all $t_i > 0$, an outer exponent is active precisely when one of the coefficient products in its displayed determinant formula is positive. No further activation table is needed. A horizontal opposite pair also activates an off-line product: combining λD or αm with βn or cA forces, respectively, $\lambda n, cD, \beta m, \alpha A$. The vertical case is identical with the roles exchanged. The two diagonal pairs force exactly the supports $\{n, m, A, D\}$ and $\{\lambda, c, \alpha, \beta\}$; any extra positive event activates an off-line exponent.

Both diagonal supports are impossible. At $u = v = 1$, the signed core matrix is

$$C = \begin{pmatrix} 2 & 2 & -2 & -2 \\ 2 & 6 & -2 & 2 \\ -2 & -2 & 6 & -2 \\ -2 & 2 & -2 & 2 \end{pmatrix}.$$

For the first support take $z = (1, 0, 0, 1)$, and for the second take $z = (1, -1, 1, 1)$. The respective positive event packets annihilate z, while $z^T C z = 0$ and $Cz \neq 0$. Thus $z^T P z = 0$ but $Pz = dCz \neq 0$, contradicting $P \succeq 0$.

If E has rank zero, the support contains none of the twelve activating pairs. Arrange the event labels cyclically as

$$(\lambda, D, c, A, \alpha, m, \beta, n).$$

Those pairs are exactly the eight neighboring pairs and the four opposite pairs. An independent set has at most three vertices: a four-element independent set in the cycle would be an alternating set and would contain an opposite pair. A maximal independent triple has cyclic gaps $(2, 3, 3)$; any smaller independent set extends to one. These triples form one symmetry orbit. Indeed the CHSH maps g_A, g_B, g_T from Section 3.1 act on the cyclic index i as $5 - i, 3 - i, 2 - i$ modulo eight, so $g_B g_T$ is a one-step rotation.

It therefore suffices to consider $\{\lambda, m, c\}$. The single polynomial state

$$\psi_H = (uv, -u, u^2 v, 1 + v^2 + u^2 v^2)$$

annihilates all three event amplitudes. Substitution in the signed parent gives exactly

$$F_H(\psi_H) = -2u^2 v^2 (1 + u^2)(1 + v^2) < 0 \qquad (uv \neq 0),$$

where F_H denotes the parent with $d = 1$ and every event weight zero. Thus a source supported in the triple has $F(\psi_H) = dF_H(\psi_H) < 0$. Symmetry gives the contradiction for every rank-zero support. This replaces the entire witness table by one Hardy-state identity.

Hence E spans $\mathbb{R}^2$. In a fixed real kernel basis write

$$P(\theta) = \begin{pmatrix} a & b^T \\ b & C_0 \end{pmatrix}, \qquad F = a + 2b^T y + y^T C_0 y, \quad C_0(\theta_0) \succ 0.$$

The angle Schur complement of the state Hessian is the Hessian of $a - b^T C_0^{-1} b$. Since $\det P = (\det C_0)(a - b^T C_0^{-1} b)$ and the Laurent value and gradient vanish, its sign equals that of

$$\nabla^2 L = \sum_{e\in E} w_e e e^T \succ 0.$$

The change from (u, v) to $(\log u^2, \log v^2)$ is locally invertible. The full Hessian is therefore positive definite. All inverses in this paragraph are ordinary real residue calculations. □

Normalize the source by a positive pure coefficient factor and choose a bounded projective testing-state chart. Physicality gives the polynomial positivity premise of Lemma 4.1 over the coefficient field by transfer. Applying that lemma proves tensor SOS at every noncommuting zero residue with $\bar{d} > 0$; restore the projective state scale by its pure testing square. A positive residue slack is already covered by Lemma 1.1. Thus the unresolved residue geometry is confined to commuting tests and $\bar{d} = 0$.

4.2 The commuting corner: a sharp leading inequality and a full certificate

Put $\lambda = \alpha = 0$ temporarily. At $u = v = 0$ the parent is $2t_7X^2 + 2t_8Y^2$, where

$$t_7 = d + 2(\beta + n + m), \qquad t_8 = d + 2(c + A + D).$$

Give u, v, X, Y weight one and W, Z weight zero. The exact degree-two part is

$$F_2 = 2t_7\left(X + \frac{L_x}{2t_7}\right)^2 + 2t_8\left(Y + \frac{L_y}{2t_8}\right)^2 + 4\mathcal{B},$$

where

$$\begin{aligned}
L_x &= -4mvW + 2(t_7 - 2n)uZ, & L_y &= -2(t_8 - 2A)uW + 4DvZ,\\
\mathcal{B} &= a(uW)^2 + b(vW)^2 + \gamma(uZ)^2 + e(vZ)^2 - \kappa uvWZ,\\
a &= \frac{A(t_8 - 2A)}{t_8}, & b &= \frac{m(t_7 - 2m)}{t_7},\\
\gamma &= \frac{n(t_7 - 2n)}{t_7}, & e &= \frac{D(t_8 - 2D)}{t_8},\\
\kappa &= d + \frac{4mn}{t_7} + \frac{4AD}{t_8}.
\end{aligned}$$

Physicality implies $F_2 \geq 0$: substitute $\varepsilon(u, v, X, Y)$, divide by ε^2, and take the real limit. The resulting fixed algebraic condition transfers to real closed fields.

Proposition 4.3 (Exact leading condition)**.** *For $a, b, \gamma, e \geq 0$ and $\kappa > 0$,*

$$\mathcal{B} \geq 0 \text{ for all } u, v, W, Z \quad \Longleftrightarrow \quad \kappa \leq 2(\sqrt{ae} + \sqrt{b\gamma}).$$

Proof. Let $R_0 = \sqrt{ae} + \sqrt{b\gamma}$. For positive coefficients choose the ratios u/v and W/Z to attain simultaneously the two arithmetic–geometric mean equalities. This proves necessity; zero coefficients follow by adding an arbitrarily small positive real number and passing to the limit. For sufficiency put $\vartheta = \kappa/(2R_0) \leq 1$. Exactly,

$$\begin{aligned}
\mathcal{B} = \vartheta\big[(\sqrt{a}\,uW - \sqrt{e}\,vZ)^2 + (\sqrt{b}\,vW - \sqrt{\gamma}\,uZ)^2\big]\\
+ (1 - \vartheta)\big[a(uW)^2 + b(vW)^2 + \gamma(uZ)^2 + e(vZ)^2\big].
\end{aligned}$$

All roots and weights split into the correct fields. □

Theorem 4.4 (Strict commuting certificate)**.** *Assume all source entries are bounded and nonnegative, $\bar{d} > 0$, and u, v are infinitesimal or zero. If $\bar{a}, \bar{b}, \bar{\gamma}, \bar{e} > 0$ and*

$$\bar{\kappa} < 2(\sqrt{\bar{a}\bar{e}} + \sqrt{\bar{b}\bar{\gamma}}),$$

the entire parent is tensor SOS for arbitrary state coordinates. The actual source may have any nonnegative λ, α.

Proof. Retain the λ, α packets as positive squares. On $g = (X, Y, uW, vW, uZ, vZ)$, the displayed completion makes the Gram of F_2 strictly positive in residue. The entire remaining difference is exactly $g^T E g$ with $\bar{E} = 0$.

Here is a finite allocation proving the last assertion. Every omitted monomial has weighted degree at least four. With two state factors from X, Y, leave its at least two angle factors in the coefficient. With one state factor from W, Z, allocate one angle to that factor and leave at least two angles. With two factors from W, Z, allocate one angle to each and leave at least two. Every allocation exists because all four angular roots are present. The coefficients are bounded and have zero residue; no state coordinate occurs in them. Apply Lemma 1.2 and restore the retained squares. □

4.3 A critical wall resolved by five roots

The following full-parent result includes an exact critical equality. Write

$$t = d + 2(c + A + D), \qquad \Delta = 8ADc - 4ADd - d^2 t.$$

Theorem 4.5 (Critical-wall certificate)**.** *Let the source entries be bounded and nonnegative with $\bar{d}, \bar{A}, \bar{D} > 0$. If the actual coefficient-field number $\Delta \geq 0$, then at every independent infinitesimal or zero u, v the complete parent is tensor SOS, for arbitrary state coordinates.*

Proof. Retain all event packets except c, A, D. The remaining signed core plus those three events is exactly quadratic in

$$R = X + uZ, \quad U = uW, \quad V = vZ, \qquad g = (R, Y, U, V).$$

For a direct verification, set $C = U + uvR - Y$, $Q = R - D_u uvV/D_v$. The signed core equals

$$K = 2\left[C^2 + D_v Q^2 + D_u v^2 Y^2 + \frac{D_u u^2(1 - u^2v^2)}{D_v} V^2\right] - 4D_u UV.$$

The three event terms are $4cD_v(U - Y)^2$, $4AD_u D_v Y^2$ and $4D(C - D_u V)^2$. This defines the entire four-root Gram $G(u, v)$, with only a pure testing denominator $D_v > 0$.

The identity

$$\Delta = 2c(4AD - d^2) - d(d + 2A)(d + 2D)$$

shows that $\Delta \geq 0$ implies

$$c \geq c_* := \frac{d(d + 2A)(d + 2D)}{2(4AD - d^2)}, \qquad 4AD - d^2 > 0.$$

Since c_* is bounded above by bounded c, while its numerator has positive residue, $4AD - d^2$ has positive residue. Thus all the following coefficient quotients are bounded. Retain the excess $c - c_*$ as an additional positive packet and set $c = c_*$. For the remainder of this proof redefine $t := d + 2(c_* + A + D)$.

At $u = v = 0$, G has rank three and null vector

$$z = (0, y, q, 1), \qquad y = \frac{d}{2A}, \quad q = \frac{D(2A + d)}{A(2D + d)}.$$

Use the bounded pure coefficient change

$$h_1 = R, \quad h_2 = Y - yV, \quad h_3 = U - qV, \quad N = V.$$

The hard block at zero is

$$H_0 = \operatorname{diag}\left(2d, \begin{pmatrix} 2t & -2(t - 2A) \\ -2(t - 2A) & 2(t - 2A) \end{pmatrix}\right) \succ 0.$$

The 2×2 determinant is $8A(t - 2A) > 0$. The exact null evaluation is

$$z^T G(u, v)z = a_{20}u^2 + a_{02}v^2 + a_{40}u^4 + a_{22}u^2v^2,$$

with

$$a_{20} = \frac{d(2A + d)}{A}, \quad a_{02} = \frac{Dd(2A + d)^2}{A^2(2D + d)}, \quad a_{40} = 2(2D + d), \quad a_{22} = \frac{d^2(2A + d)}{2A^2}.$$

All are positive, and a_{20}, a_{02} have positive residues. For complete control of the cross terms, the three hard-null entries are

$$\begin{aligned}
(Gz)_1 &= -\frac{uv}{A}\big[(4AD + 2Ad)u^2 + 2Ad + d^2\big], \\
(Gz)_2 &= \frac{u^2}{A}\big[4AD + 2Ad + (2Ad + d^2)v^2\big], \\
(Gz)_3 &= \frac{-(4AD + 2Ad)u^2 + (2Ad + d^2)v^2}{A}.
\end{aligned}$$

These formulas follow by substituting z into the four displayed packet expressions; they are identities of the entire parent.

Use the five roots (h_1, h_2, h_3, uN, vN). Allocate a factor uN in the first two cross terms, leaving a coefficient divisible by v or u; split the third between uN and vN. The hard-hard corrections vanish at zero. Give uN the diagonal $a_{20} + a_{40}u^2 + a_{22}v^2$, and vN the diagonal a_{02}. The exact Gram therefore has residue

$$\operatorname{diag}(\bar{H}_0, \bar{a}_{20}, \bar{a}_{02}) \succ 0.$$

Apply Lemma 1.2 and restore all retained squares. No u, v, state coordinate or mixed tensor has been inverted. □

Theorem 4.5 covers the wall itself and arbitrary nonnegative infinitesimal Δ. It does not cover $\Delta < 0$ infinitesimal when other event packets compensate the deficit. The equality theorem alone does not justify an open neighborhood on that side.

Chapter 5

Zero signed-weight residues: the eight-cycle geometry

Component `p76173e99815f`
Source: `CORE/prior/inherited/zero_d_geometry.tex`
SHA256: `76173e99815f0efdd03931d13048d2bce245a5b3c083fc0373e3adeca1638c3d`
Inventory use: sec:zero-d-cycle | matching geometry and exact event kernels; sec:zero-d-cycle | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

5.1 Zero signed-weight residues: the eight-cycle geometry

Normalize the source so that $d + \sum_i c_i = 1$. Suppose $\bar{d} = 0$ and both testing measurement pairs are noncommuting in residue. A projective choice of testing frame gives finite residue half angles with $u_0 v_0 \neq 0$. The following classification uses only the geometry of product vectors.


Lemma 5.1 (Product residues are already settled)**.** *At noncommuting residue angles every product state satisfies $H > 0$. Consequently the entire parent is tensor SOS whenever its residue state is product.*

Proof. For a product state,

$$S = a_0(b_0 - b_1) + a_1(b_0 + b_1) \leq |b_0 - b_1| + |b_0 + b_1| = 2\max(|b_0|, |b_1|) \leq 2.$$

If both Bob moduli are less than one, the inequality is strict. Otherwise precisely one is one, since noncommuting qubit reflections have no common eigenvector. Both $b_0 - b_1$ and $b_0 + b_1$ are then nonzero; equality would require $|a_0| = |a_1| = 1$, again impossible. Thus $H = 1 - S/2 > 0$ at the residue. Its actual testing value remains positive. The signed term $d \otimes (D_u D_v H)$ is therefore a square after separate field square roots, and every event term is already a positive weighted square. □

The losing events form a single bipartite cycle between the four Alice and four Bob outcome rays:



For example, λ, D share Bob's ray B_1^+, while D, c share Alice's ray A_1^-. The eight local rays are distinct within their respective parties at noncommuting angles.

Theorem 5.2 (Exhaustive noncommuting zero-d residue classification)**.** *An unresolved zero residue has an entangled state. Its positive-residue event support is a matching in the eight-cycle, hence has at most four elements.*

For a support of size $k \leq 3$, the event residue has Hessian rank k in the five angle-and-projective-state variables. Its zero locus is smooth of dimension $5 - k$. For four events the only possibilities are

$$\{n, m, A, D\}, \quad u_0^2 = v_0^2; \qquad \{\lambda, c, \alpha, \beta\}, \quad u_0^2 v_0^2 = 1.$$

Each has state rank three and one additional angular normal, so its full Hessian rank is four and its zero locus has dimension one. There are no further exceptional noncommuting loci.

Proof. At $\bar{d} = 0$, vanishing of the nonnegative event sum forces every positive-weight event amplitude to vanish. Write the real state as a 2×2 matrix Ψ. Entanglement means $\det \Psi \neq 0$. For each fixed nonzero Alice ray a, the equation $a^T \Psi b = 0$ determines a unique Bob projective ray, and conversely. Vanishing events therefore share no vertex: they form a matching.

Any three distinct product vectors in this event system are linearly independent. Indeed, three dependent rank-one matrices must share a left factor or share a right factor. If the first two have both independent left and independent right factors, their span is diagonal in suitable bases and contains only those two rank-one projective points. If they share a factor, their entire span shares it. A cycle vertex has degree two, so three event vectors cannot share a factor. For $k \leq 3$, the k independent state-linear amplitudes have derivative rank k on the projective state chart at their common zero. The implicit-function theorem and the formula for the Hessian of a sum of squared vanishing functions give the stated rank and smooth zero set.

An even cycle has exactly two perfect matchings. For $\{n, m, A, D\}$ the equations give $X = Y = 0$, $vW - uZ = uW - vZ = 0$. A nonzero solution exists precisely on $u^2 = v^2$. Put $W = 1$, $v = \sigma u$, $Z = \sigma$ with $\sigma = \pm 1$. After removing the independent X, Y normals, the other two derivatives in (u, v, Z) are

$$(-\sigma, 1, -u), \qquad (1, -\sigma, -\sigma u).$$

They are independent because $u \neq 0$. Thus the full rank is four.

For $\{\lambda, c, \alpha, \beta\}$, put $W = 1$. The first three equations give $X = -1/v$, $Y = u$, $Z = uv$. The fourth becomes $(u^2 v^2 - 1)/v = 0$, with nonzero angular gradient. Again there are three state normals and one angular normal. These descriptions also show directly that the signed core vanishes on the two curves. □

Corollary 5.3 (Adjacent dominant events need no noncommuting atlas)**.** *If two positive-residue event coefficients share an outcome ray, the entire parent is tensor SOS at every noncommuting residue test.*

Proof. Positive slack residue is already settled by Lemma 1.1. At a zero residue, the two amplitudes sharing a ray force a product kernel state: two distinct rays at the other party span its two-dimensional space. The preceding product-state lemma applies. □

The two canonical families reduce the matching list further. For C_1 the only three-event matchings are $\{m, A, D\}$, $\{c, \alpha, \beta\}$, $\{A, D, \beta\}$ and $\{D, \alpha, \beta\}$; there is no four-event matching. For C_2 they are $\{n, m, c\}$, $\{n, m, A\}$, $\{n, m, D\}$, $\{n, c, \alpha\}$, $\{n, A, D\}$, $\{n, D, \alpha\}$ and $\{m, A, D\}$; only $\{n, m, A, D\}$ can have four events.

A two-variable scalar restriction. The nonextendable matching $\{\lambda, m, A\}$ has polynomial kernel state

$$(W, X, Y, Z) = (uv, -u, 0, 1 + v^2).$$

Substitute this state into the entire parent and divide by the positive pure testing factor $2u^2v^2(1+v^2)$. With $U = u^2, V = v^2$, the result is exactly

$$L(U,V) = \frac{2n(1+U)}{V} + 2cU + \frac{2D(1+V)}{U} + \frac{2\alpha(1+U)(1+V)}{UV} + 2\beta V - d.$$

Physicality implies $L \geq 0$ on the positive quadrant. In particular, $d \leq 4n + 2c + 4D + 8\alpha + 2\beta$. Thus at least one complementary event weight is not infinitesimal relative to d. The other seven nonextendable triples give analogous exact restrictions by their polynomial kernel states.

Source note. The cycle gives the support geometry, not by itself the full-parent certificate. The following weighted normal completions retain every normal cross term and handle one, two, three and four hard event constraints at arbitrary coefficient scales. The finite induction in Theorem 11.2 completes every noncommuting residue test. It does not assume that an extreme source has three positive-residue event weights.

Chapter 6

Weighted normal completion at matching residues

Component `p24f8a75e651e`
Source: `CORE/prior/inherited/zero_d_weighted.tex`
SHA256: `24f8a75e651e715d8ae6a9ed52b453e9e947cd9615dd9d726124b15b3b5d6345`
Inventory use: lem:weighted-contact; eq:three-N; eq:three-E; thm:three-event-weighted; thm:four-event-weighted | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

6.1 Weighted normal completion at matching residues

The cycle classification can be strengthened without any division by a mixed tensor. The elementary device below keeps the large normal form intact while solving only a small, nonsingular critical-point system.

Lemma 6.1 (A weighted regular-contact lemma)**.** *Let $\sigma > 0$ be an infinitesimal in the coefficient field. In local real coordinates $(x, y) \in \mathbb{R}^r \times \mathbb{R}^s$, suppose a physically nonnegative polynomial has the exact form*

$$F(x, y) = x^T A(x, y)x + \sigma G(x, y),$$

with bounded coefficient-field coefficients and symmetric A. At the independent test assume $\bar{x} = 0$, $\bar{y} = y_0$, and $\bar{A}(0, y_0) \succ 0$. Put $g(y) = \overline{G}(0, y)$. If either $g(y_0) > 0$, or

$$g(y_0) = 0, \qquad \nabla g(y_0) = 0, \qquad \nabla^2 g(y_0) \succ 0,$$

then the entire tensor evaluation of F is a sum of squares. The same assertion holds for rational coordinate expressions whose denominators are products of coefficient units and real testing polynomials nonzero at y_0.

Proof. In the first case divide by σ and use the roots $(x/\sqrt{\sigma}, 1)$. Allocate every term of $G - G(0, y)$ containing x to these roots. The exact Gram has residue $\operatorname{diag}(\bar{A}(0, y_0), g(y_0)) \succ 0$.

In the second case put $x = \sigma s$ in the stationary equations for F/σ. They become

$$\begin{aligned} 2A(\sigma s, y)s + \sigma\big(s^T A_{x_i}(\sigma s, y)s\big)_{i=1}^{r} + G_x(\sigma s, y) = 0, \\ \sigma\big(s^T A_{y_j}(\sigma s, y)s\big)_{j=1}^{s} + G_y(\sigma s, y) = 0. \end{aligned}$$

At residue, the second equation is $\nabla g(y) = 0$ and the first uniquely determines

$$s_0 = -\bar{A}(0, y_0)^{-1}\overline{G_x}(0, y_0)/2.$$

The Jacobian in (s, y) is block upper triangular, with diagonal blocks $2\bar{A}(0, y_0)$ and $\nabla^2 g(y_0)$. The real semialgebraic implicit-function theorem, transferred to the coefficient field, supplies an exact stationary point $(x_c, y_c) = (\sigma s_c, y_c)$ with $(\bar{s}_c, \bar{y}_c) = (s_0, y_0)$. Physicality gives $a = F(x_c, y_c)/\sigma \geq 0$.

Set $h = x - x_c$ and $t = y - y_c$. The linear terms vanish exactly. Allocate terms with at least two h factors to $h_i h_j/\sigma$; terms with exactly one h factor have bounded coefficients after division by σ, because $x_c/\sigma = s_c$ is bounded. Terms with no h factor contributed by the hard form have a factor $x_c^2/\sigma = \sigma s_c^2$. Consequently the exact finite Taylor allocation is

$$F/\sigma = a + \begin{pmatrix} h/\sqrt{\sigma} \\ t \end{pmatrix}^T Q \begin{pmatrix} h/\sqrt{\sigma} \\ t \end{pmatrix}, \qquad \bar{Q} = \operatorname{diag}\Big(\bar{A}(0,y_0), \tfrac{1}{2}\nabla^2 g(y_0)\Big) \succ 0.$$

Every unallocated higher factor has zero residue. Lemma 1.2 applies, a is a pure coefficient square, and the positive pure factor σ is restored. For rational expressions, clear the finitely many indicated denominators by separately positive squares before applying the polynomial argument. The proof never inverts A in the tensor ring: its real inverse is used only to identify the initial point for the coefficient-field implicit-function theorem. □

6.1.1 Three dominant events: two Laurent problems

Normalize the source and suppose exactly three event coefficients have positive residues, forming a matching M. If all complementary coefficients and d vanish literally, the parent is already an event-square sum. Otherwise write

$$\sigma = \max(\{d\} \cup \{c_i : i \notin M\}) > 0.$$

The three event amplitudes are independent state coordinates near their common noncommuting kernel. A real, angle-dependent affine change of projective state coordinates makes them the normal coordinates x. Its denominators are pure testing units. The parent is therefore in the form of Lemma 6.1: the three dominant packets supply A, and all complementary packets and the signed core supply σG.

There are two matching orbits under the CHSH symmetries. Representatives, polynomial kernel states and exact constrained Laurent polynomials are

$$\begin{aligned} M_N &= \{\lambda, m, A\}, & \psi_N &= (uv, -u, 0, 1+v^2), \\ M_E &= \{n, m, A\}, & \psi_E &= (u, 0, 0, v). \end{aligned}$$

With $U = u^2, V = v^2$, substitution into the complete parent gives

$$\begin{aligned} \frac{F(\psi_N)}{2UV(1+V)} &= \frac{2n(1+U)}{V} + 2cU + \frac{2D(1+V)}{U} \\ &\quad + \frac{2\alpha(1+U)(1+V)}{UV} + 2\beta V - d, \qquad (6.1) \\ \frac{F(\psi_E)}{2UV} &= d(U/V - 1) + \frac{2\lambda(1+U)}{V} + \frac{2cU(1+V)}{V} \\ &\quad + \frac{2D(U-V)^2}{UV} + \frac{2\alpha(1+U)}{U} + 2\beta(1+V). \qquad (6.2) \end{aligned}$$

All nonconstant Laurent coefficients are nonnegative. At a stationary zero, the logarithmic Hessian is $\sum_e a_e U^{e_1} V^{e_2} e e^T$ and is positive definite exactly when the active exponent vectors span $\mathbb{R}^2$.

For (6.1), any positive n, D or α already supplies two independent exponent vectors. If all three vanish, the only remaining nonconstant terms are $2cU + 2\beta V$, which cannot have a stationary zero unless both coefficients vanish. Thus when $\overline{d/\sigma} > 0$, every physical zero of the normalized restriction has positive definite logarithmic Hessian. If $\overline{d/\sigma} = 0$, the restriction is strictly positive: some complementary coefficient has positive normalized residue, and its event amplitude is nonzero on ψ_N at every noncommuting angle.

For (6.2), a positive residue of d/σ supplies exponent $(1, -1)$. Failure to span two dimensions forces $\lambda = c = \alpha = \beta = 0$ in normalized residue. Stationarity then requires $D > 0$, but the minimum of the remaining diagonal expression is

$$2\sqrt{2D(d+2D)} - d - 4D < 0,$$

since $(d+4D)^2 - 8D(d+2D) = d^2 > 0$. Physicality excludes this case. Hence every physical zero again has strictly positive logarithmic Hessian. Multiplying by a positive smooth testing factor or passing from logarithmic angles to $u, v \neq 0$ preserves positive definiteness at a zero.

Theorem 6.2 (Three-event weighted completion)**.** *For a physical source, at a noncommuting zero residue with exactly three dominant matching events, the entire parent is tensor SOS if the matching is nonextendable, or if d/σ has positive residue. For an extendable matching with $\overline{d/\sigma} = 0$, it is also tensor SOS unless the only complementary event with positive normalized residue is the fourth edge completing the perfect matching, and the testing residue lies on that matching's curve.*

Proof. Physicality makes the constrained normalized residue nonnegative. The preceding calculation gives either a positive value or a strict two-angle minimum, so Lemma 6.1 applies. In the remaining extendable case, every complementary event except the fourth matching edge is nonzero on the kernel. Thus a zero requires those other normalized coefficients to vanish; the fourth event must vanish, giving precisely the perfect-matching curve. The cycle rotations and reflections induced by the three CHSH maps carry each of the eight extendable triples to M_E and each of the eight nonextendable triples to M_N. □

6.1.2 Four dominant events: the first normal correction is a Laurent circuit

Theorem 6.3 (Four-event weighted completion)**.** *For a physical source, if $\bar{d} = 0$ and four event coefficients have positive residues, the entire parent is tensor SOS at every noncommuting residue test.*

Proof. The case $d = 0$ is already an event-square sum, so assume $d > 0$. The cycle classification reduces a zero to a perfect matching. By symmetry take $M = \{n, m, A, D\}$, with these four coefficients positive units. The sign choice of the matching curve is removed by a real testing reflection. Near the curve write

$$u = t, \quad v = t + \delta, \quad W = 1, \quad Z = 1 + \zeta, \qquad z = (X, Y, \delta, \zeta), \quad \bar{t} \neq 0.$$

Put $B = 1 + t^2$ and use the four linear normal forms

$$X, \qquad Y, \qquad r_m = \delta - X + t^2 Y - t\zeta, \qquad r_D = t^2 X - Y - \delta - t\zeta.$$

Their matrix $J(t)$ is invertible for $t \neq 0$. The dominant quadratic form and the linear part of the signed core are exactly

$$\begin{aligned} Q_t(z) &= 4\{nB^2X^2 + AB^2Y^2 + mr_m^2 + Dr_D^2\}, \\ \ell_t(z) &= 2t\{B(X - Y) - r_m + r_D\}. \end{aligned}$$

Complete $Q_t + d\ell_t$ by the shift $z = h + d\eta(t)$, where in the four displayed normal forms

$$J(t)\eta(t) = \left(-\frac{t}{4nB}, \frac{t}{4AB}, \frac{t}{4m}, -\frac{t}{4D}\right)^T.$$

This is a finite separated rational expression: $J(t)^{-1}$ is pure testing, and the only coefficient inverses are the four positive units. The minimum correction is

$$-\frac{d^2t^2}{4}\left(\frac{1}{n} + \frac{1}{A} + \frac{1}{m} + \frac{1}{D}\right).$$

On the curve, the complementary packets are exactly

$$4(1 + t^2)\{(\lambda + \alpha) + t^2(c + \beta)\}.$$

Let $\sigma = \max(\lambda, \alpha, c, \beta)$. Physicality forces $\sigma > 0$ and d^2/σ bounded. Indeed, otherwise substitute $t = 1$ and the displayed shift with $h = 0$. The complementary packets are $o(d^2)$, while the hard

packets and signed core, divided by d^2, have strictly negative residue. All omitted terms have an additional factor d, giving a contradiction.

After the shift, collect terms of degree at least two in h in a symmetric matrix $A(h,t)$. The constant term is $O(d^2+\sigma)$ and the coefficient of each linear h term is $O(d^2+\sigma)$: the terms of first order in d cancel by the displayed completion. These statements are finite polynomial divisibilities after clearing t and B by pure testing squares. Thus the entire shifted parent has the exact form

$$F = h^T A(h,t)h + \sigma G(h,t), \qquad \bar{A}(0,t) \succ 0,$$

with bounded coefficients, and

$$\bar{G}(0,t) = 4(1+t^2)(p+qt^2) - \frac{\rho t^2}{4}\left(\frac{1}{\bar{n}} + \frac{1}{\bar{A}} + \frac{1}{\bar{m}} + \frac{1}{\bar{D}}\right),$$

where $p = \overline{(\lambda+\alpha)/\sigma}$, $q = \overline{(c+\beta)/\sigma}$ and $\rho = \overline{d^2/\sigma}$. Here $p, q \geq 0$ and $p+q > 0$. Divide by the positive testing factor $4t^2$ and put $U = t^2$; the residue is the one-variable Laurent polynomial

$$p(1+U^{-1}) + q(1+U) - \frac{\rho}{16}\left(\frac{1}{\bar{n}} + \frac{1}{\bar{A}} + \frac{1}{\bar{m}} + \frac{1}{\bar{D}}\right).$$

Physicality makes it nonnegative for every $U > 0$. At a zero, stationarity gives $p/U = qU$; hence both p and q are positive and its logarithmic second derivative $p/U + qU$ is strictly positive. Lemma 6.1 therefore settles every test near the curve. Tests away from it have positive residue, completing the proof. □

The next scale. The next result handles the lower-scale fourth-edge promotion. The subsequent matching terminal removes the assumption of comparable hard weights, and the constrained determinant arguments handle one or two hard events. Together they yield the finite induction of Theorem 11.2.

Chapter 7

A fourth edge at a smaller coefficient scale

Component `p4e9ae18b7af4`
Source: `CORE/prior/inherited/zero_d_promotion.tex`
SHA256: `4e9ae18b7af4a8aba644b170b651ad30bcc6f22b0069c63bbc8f09dc3c4ad974`
Inventory use: prop:fourth-edge-promotion; exact R,f,L identity and physical rho bound | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

7.0.1 A fourth edge at a smaller coefficient scale

Proposition 7.1 (The fourth-edge promotion)**.** *The exceptional promotion case in Theorem 6.2 is also tensor SOS. Consequently every noncommuting zero-d residue with at least three dominant event coefficients is settled.*

Proof. By symmetry the three dominant coefficients are n, m, A and the smaller fourth coefficient is $D = \mu > 0$. The promotion hypotheses say

$$\mu \in \mathfrak{m}_c, \qquad d/\mu \in \mathfrak{m}_c, \qquad \lambda/\mu, c/\mu, \alpha/\mu, \beta/\mu \in \mathfrak{m}_c.$$

As before $d = 0$ is immediate. Set $\tau = \max(\lambda, c, \alpha, \beta)$. Choose the sign of the matching curve so that $\bar{v} = \bar{u} \neq 0$, and put $W = 1$. Use the exact strong-event state coordinates

$$x = (X, Y, R), \qquad R = v - X + uvY - uZ, \qquad Z = \frac{v - X + uvY - R}{u}.$$

Only the pure testing unit u is inverted. Define

$$f = \frac{u^2 - v^2}{u}, \qquad L = v(u + u^{-1})X - (1 + v^2)Y + (v/u)R.$$

Then the fourth event amplitude is exactly $f + L$. On $x = 0$, the signed core is $2uf$, and hence the entire signed-plus-fourth part is

$$4\mu f^2 + 2duf = 4\mu \left(f + \frac{du}{4\mu} \right)^2 - \frac{d^2 u^2}{4\mu}.$$

Testing $u = 1$, $v = \sqrt{1 + d/(4\mu)}$ and $x = 0$ proves $\tau > 0$ and

$$\rho = \frac{d^2}{\mu\tau} \quad \text{is bounded.}$$

Indeed all complementary amplitudes are bounded at that test, and otherwise their total would be infinitesimal relative to the displayed negative term.

Put $\delta = v - u$. If $\bar{\rho} = 0$, an exact Gram on the roots

$$\left(x/\sqrt{\tau},\ \delta\sqrt{\mu/\tau},\ 1 \right)$$

has strictly positive residue. To see the bounds directly, the strong packets supply a positive three-by-three block, and $4\mu(f+L)^2$ supplies the positive fourth diagonal entry 16 because $f = -2\delta - \delta^2/u$. The strong–fourth cross block has size $\sqrt{\mu}$; signed-core linear terms have sizes $d/\sqrt{\tau} = \sqrt{\mu\rho}$ and $d/\sqrt{\mu\tau} = \sqrt{\rho}$, respectively. Every other contribution has bounded coefficients. The constant residue comes from the complementary packets on the matching curve and is $4(1+u_0^2)(P+Qu_0^2) > 0$, where

$$P = \overline{(\lambda+\alpha)/\tau}, \qquad Q = \overline{(c+\beta)/\tau}, \qquad P + Q > 0.$$

It remains to treat ρ with positive residue. Write $\varepsilon = d/\mu$, so that $d = \mu\varepsilon$ and $\tau = \mu\varepsilon^2/\rho$. In the coefficient-field stationary equations make the substitutions

$$x = ds, \qquad \delta = \varepsilon r, \qquad u = t.$$

Divide the three x equations by d, the δ equation by d, and the t equation by τ. After the pure unit clearing by t, all coefficients are bounded functions of μ, ε, ρ and the four normalized complementary coefficients. This can also be read directly from the strong packets, the exact formulas for f, L, and the fact that the signed core is quadratic in x.

At residue let $B = 1 + t^2$ and

$$Q_s = 4\,\mathrm{diag}(\bar{n}B^2, \bar{A}B^2, \bar{m}), \qquad \ell = (B, -B, 1)^T, \qquad b = (4tB, -4tB, 0)^T.$$

The first four stationary equations reduce to

$$2Q_s s - 16r\ell + b = 0, \qquad 32r - 4t = 0.$$

The limiting objective for the last two variables is

$$\Psi(r,t) = \bar{\rho}(16r^2 - 4tr) + 4(1+t^2)(P + Qt^2).$$

Eliminating $r = t/8$ leaves

$$g(t) = 4(1+t^2)(P+Qt^2) - \bar{\rho}t^2/4.$$

Physicality makes $g(t) \geq 0$ for every nonzero real t: substitute $x = 0$, $\delta = \varepsilon t/8$ into the original parent and take its normalized residue. Dividing g by $4t^2$ gives

$$P(1+t^{-2}) + Q(1+t^2) - \bar{\rho}/16.$$

If this is positive at the testing residue, the same strict Gram argument applies after the explicit shift $\delta \mapsto \delta - \varepsilon t/8$; its mixed strong-normal terms remain zero in residue. Otherwise its zero is stationary, $P, Q > 0$, and $g''(t_0) > 0$.

At such a zero, the scaled stationary system has an invertible Jacobian. The strong block is $2Q_s \succ 0$. The remaining two-by-two block has first row $(32, -4)$ and Schur complement $g''(t_0) > 0$. The coefficient-field implicit-function theorem therefore produces an exact stationary center

$$x_c = ds_c, \qquad \delta_c = \varepsilon r_c, \qquad t_c, \quad \bar{r}_c = t_0/8, \quad \bar{t}_c = t_0.$$

Physicality gives $a = F(x_c, \delta_c, t_c)/\tau \geq 0$.

Expand exactly at that center and use the roots

$$\left(\frac{x - x_c}{\sqrt{\tau}},\ \sqrt{\frac{\mu}{\tau}}(\delta - \delta_c),\ t - t_c \right).$$

The Gram coefficients are bounded. For transparency, strong–fourth entries are $O(\sqrt{\mu}) + O(d/\sqrt{\mu})$; strong–tangent entries are $O(d/\sqrt{\tau}) = O(\sqrt{\mu})$; fourth–tangent entries are $O(d/\sqrt{\mu\tau}) = O(1)$. Pure

tangent contributions from strong packets are $O(d^2/\tau) = O(\mu)$, while those from the fourth packet and signed core are $O(d^2/(\mu\tau)) = O(1)$. Higher Taylor factors have zero residue. These estimates follow term by term from the quadratic strong packets, $f = -2\delta - \delta^2/t$, and the displayed exact L.

The residue Gram is the direct sum of Q_s and the positive definite Hessian form of $\Psi/2$, with its r coordinate rescaled by $1/\sqrt{\bar{\rho}}$. Positivity of that Hessian follows from its positive normal entry $32\bar{\rho}$ and the positive Schur complement $g''(t_0)$. The exact linear terms vanish at the stationary center, so $F/\tau = a + h^T Q h$ with $\bar{Q} \succ 0$. Pure testing powers of t clear the finite rational expressions; the corresponding constant term is a positive pure coefficient times a pure testing square. Lemma 1.2 and pure-factor restoration finish the proof. □

From this promotion to all coefficient scales. The next theorem lets all four matching weights have independent scales. One- and two-hard-event kernels are then handled by their constrained determinants, completing the finite matching induction in Theorem 11.2.

Chapter 8

A matching terminal independent of its internal coefficient scales

Component pa2cbe6ecbeb4
Source: CORE/prior/inherited/zero_d_matching_flags.tex
SHA256: a2cbe6ecbeb41e961eecbacd42afa980b05e26d6e43547f366f1e35bb370cfc2
Inventory use: thm:matching-flag-terminal | four-event matching terminal with unequal hard scales; thm:matching-flag-terminal | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

8.0.1 A matching terminal independent of its internal coefficient scales

Theorem 8.1 (Four matching weights at arbitrary scales)**.** *Let the source be physical, let its four positive matching weights be n, m, A, D, and put $\mu = \min(n, m, A, D)$, and suppose*

$$d/\mu \in \mathfrak{m}_c, \qquad \lambda/\mu, c/\mu, \alpha/\mu, \beta/\mu \in \mathfrak{m}_c.$$

Then the full parent is tensor SOS at every noncommuting residue test annihilating those four events. The same statement holds for the other perfect matching by CHSH symmetry.

Proof. The cases $d = 0$ and positive slack residue are immediate. Cycle symmetries let the smallest matching weight be $D = \mu$. Set

$$\eta_1 = \mu/n, \qquad \eta_2 = \mu/A, \qquad \eta_3 = \mu/m; \quad 0 < \eta_i \le 1.$$

All three may have different residues, including zero. Use the exact state coordinates and functions f, L from Proposition 7.1, so that the strong packets are

$$4nD_uD_vX^2 + 4AD_uD_vY^2 + 4mR^2,$$

and the fourth amplitude is $f + L$. Put $\tau = \max(\lambda, c, \alpha, \beta)$. The exact scalar test $x = 0$, $u = 1$, $v = \sqrt{1 + d/(4\mu)}$ again proves $\tau > 0$ and boundedness of

$$\rho = d^2/(\mu\tau), \qquad \varepsilon = d/\mu.$$

If $\bar{\rho} = 0$, the direct Gram on

$$\left(\sqrt{n/\tau}X, \sqrt{A/\tau}Y, \sqrt{m/\tau}R, \sqrt{\mu/\tau}(v-u), 1\right)$$

has positive residue. Its four-normal block is

$$4B^2z_1^2 + 4B^2z_2^2 + 4z_3^2 + 4\{-2z_4 + B\sqrt{\bar{\eta}_1}z_1 - B\sqrt{\bar{\eta}_2}z_2 + \sqrt{\bar{\eta}_3}z_3\}^2 \succ 0,$$

where $B = 1 + u_0^2$. The constant residue is the positive complementary-event value on the matching curve. The signed linear terms have coefficients $\sqrt{\rho\eta_i}$ and $\sqrt{\rho}$, hence vanish in residue.

Assume now $\bar\rho > 0$. In the coefficient-field stationary equations put

$$X = \varepsilon\eta_1 s_1, \quad Y = \varepsilon\eta_2 s_2, \quad R = \varepsilon\eta_3 s_3, \quad v - u = \varepsilon r, \quad u = t.$$

Divide the three state equations by d, the $v - u$ equation by d, and the t equation by τ. Every coefficient is bounded after a pure unit clearing in t; the factors η_i in the state substitution cancel the large coefficients μ/η_i. In residue define

$$B = 1 + t^2, \quad \ell = (B, -B, 1), \quad b = (B^2, B^2, 1), \quad k = -2r + \sum_{i=1}^{3} \bar\eta_i \ell_i s_i.$$

The four normal equations are

$$8b_i s_i + 8k\ell_i + (4tB, -4tB, 0)_i = 0, \qquad -16k - 4t = 0.$$

Their Jacobian in (s_1, s_2, s_3, r) has determinant $32 \cdot 8^3 B^4 > 0$: eliminate its final row from the first three to leave the diagonal entries $8b_i$. Their unique solution is

$$s_1^*(t) = -\frac{t}{4B}, \quad s_2^*(t) = \frac{t}{4B}, \quad s_3^*(t) = \frac{t}{4}, \quad r^*(t) = \frac{t}{8}(1 - \bar\eta_1 - \bar\eta_2 + \bar\eta_3).$$

Write $P = \overline{(\lambda + \alpha)/\tau}$ and $Q = \overline{(c + \beta)/\tau}$, so $P + Q > 0$. The normalized residue objective is

$$\Psi(s, r, t) = \bar\rho \left[4 \sum_i \bar\eta_i b_i s_i^2 + 4k^2 + 4tB(\bar\eta_1 s_1 - \bar\eta_2 s_2) - 4tr \right] + 4(1 + t^2)(P + Qt^2).$$

Completing its four normal squares leaves

$$g(t) = 4(1 + t^2)(P + Qt^2) - \frac{\bar\rho t^2}{4}(1 + \bar\eta_1 + \bar\eta_2 + \bar\eta_3).$$

This identity remains valid when some $\bar\eta_i = 0$; the corresponding scaled normal equation still has its diagonal pivot $8b_i$. Physicality gives $g(t) \geq 0$. At a zero its quotient by $4t^2$ is a one-variable Laurent circuit, whose logarithmic Hessian is $P/t^2 + Qt^2 > 0$. Thus $g''(t_0) > 0$.

If $g(t_0) > 0$, shift by the four explicit normal minimizers above; the positive constant and the positive four-normal block give a strict Gram. Otherwise append the tangent equation to the four scaled normal equations. Its Jacobian has Schur complement $g''(t_0) > 0$, so the coefficient-field implicit-function theorem supplies an exact stationary center with the displayed normal residues and tangent residue t_0. Physicality makes its normalized value nonnegative.

For completeness the exact Taylor roots at that center are

$$\left(\sqrt{n/\tau}(X - X_c), \sqrt{A/\tau}(Y - Y_c), \sqrt{m/\tau}(R - R_c), \sqrt{\mu/\tau}(\delta - \delta_c), t - t_c \right).$$

Their coefficient bounds use only ε, ρ, $1/\rho$, η_i and their square roots, all bounded. One can check this without estimating an inverse matrix: in each strong packet its coefficient times the squared center is

$$\frac{a_i x_{ci}^2}{\tau} = \rho\eta_i s_{ci}^2, \qquad \frac{a_i x_{ci}}{\sqrt{a_i \tau}} = \sqrt{\rho\eta_i}\, s_{ci}.$$

The fourth packet has the analogous factors ρr_c^2 and $\sqrt{\rho} r_c$. The signed and complementary packets have the same or smaller bounded factors. Each packet is at most quadratic in the state coordinates; after allocating its two normal factors, remaining higher factors are actual angle or state deviations and have zero residue. Denominators are powers of the pure testing unit t.

The residue normal Gram is the positive form displayed above. Its tangent Schur complement is $g''(t_0)/2 > 0$, as follows either by differentiating the four completed squares or by the explicit normal minimizers. Hence the entire exact Taylor Gram is strict. Adding the nonnegative pure coefficient value at the center and restoring the separately positive factors proves the claim. □

Why this is a finite induction rule. For at most three event constraints the exact state-linear event coordinates make the hard form diagonal. If their positive weights are a_i and the next coefficient scale is $\sigma \ll a_i$, the proof of Lemma 6.1 works with $x_i = (\sigma/a_i)s_i$ and roots $\sqrt{a_i/\sigma}x_i$. Its residue Jacobian has positive diagonal normal entries, and its soft block is the Hessian of the constrained next-scale residue. Thus the three-event Laurent theorem does not require the hard weights to be comparable.

If the next-scale signed coefficient still vanishes in residue, a zero of the event sum adds matching edges to the same entangled kernel. There are at most four such edges. On reaching four edges before d becomes comparable, the theorem above gives a terminal independent of their internal scales. The following two constrained determinant arguments handle the intervening cases with one or two hard events for which d first becomes comparable to the remaining coefficients.

Chapter 9

One hard event and a positive next-scale signed coefficient

Component pb16e5066334b
Source: CORE/prior/inherited/zero_d_one_hard.tex
SHA256: b16e5066334b6b1b83097bee079bc2d1fbfc34752531e0d3bafce52043b9d4b5
Inventory use: prop:one-hard-event | one-hard-event constrained curvature; prop:one-hard-event | C14_ ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

9.0.1 One hard event and a positive next-scale signed coefficient

Proposition 9.1 (The one-hard-event terminal)**.** *Let the source be physical. Suppose one event weight a satisfies $\sigma/a \in \mathfrak{m}_c$, where σ is the largest remaining source coefficient, and d/σ has positive residue. Then the entire parent is tensor SOS at every noncommuting residue test.*

Proof. CHSH symmetries are transitive on the eight events, so take λ/σ unbounded and all other coefficients divided by σ bounded, with $\overline{d/\sigma} > 0$. A test with nonzero residue of the λ amplitude has positive slack after normalization by λ. Otherwise use that amplitude as the hard state coordinate and restrict to its exact kernel

$$W = -vX.$$

The constrained normalized residue is a nonnegative quadratic form in (X, Y, Z), depending on u, v. On $Z = 0$ it is positive definite by the signed-core calculation in Section 3.1. Thus at any zero its state matrix has exactly two positive eigenvalues and one simple zero eigenvalue; in particular $Z \neq 0$, and the chart $Z = 1$ is available.

Here is a small determinant calculation proving strictness in its two angle directions. All source coefficients in the rest of this proof are the normalized residues, with $d > 0$. Differentiate the determinant polynomial of Section 3.1 with respect to λ. The resulting eight outer coefficients are

$$\begin{aligned} a_{00} &= 4\alpha t_7 t_8, & a_{01} &= 4(Dt_7 t_6 + 2\alpha m t_8),\\ a_{02} &= 8mDt_6, & a_{10} &= 4(2\alpha A t_7 + n t_8 t_4),\\ a_{12} &= 8(\beta m t_6 + cDt_2), & a_{20} &= 8nAt_4,\\ a_{21} &= 8(\beta n t_2 + cAt_4), & a_{22} &= 8\beta c t_2. \end{aligned}$$

Every t_i appearing here is strictly positive because $d > 0$. A basis change with first column $(1, 0, 0, 0)$ and remaining columns $(-v, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1)$ has determinant one. Its leading λ entry is $4\lambda D_u$, so the determinant of the constrained three-by-three state matrix is

$$4D_u D_v^2 UV\, L_\lambda(U, V), \qquad L_\lambda = h_\lambda + \sum_{(i,j)\neq(1,1)} a_{ij} U^{i-1} V^{j-1}, \quad U = u^2,\ V = v^2.$$

Thus L_λ is nonnegative on the positive quadrant. At a zero it is stationary in logarithmic angles. Its logarithmic Hessian is a positive sum of the outer products of the active exponent vectors.

We show that those vectors span the plane at every zero. If a nonempty collection spans only a line and has zero weighted gradient, that line must contain an opposite pair among the eight boundary vectors of the Newton square. There are only four possible lines.

- On the horizontal line, the off-line coefficients force $\alpha = n = 0$. A negative horizontal exponent requires $D > 0$, hence $m = c = 0$ from $a_{02} = a_{12} = 0$. But then $a_{21} = 0$, so the positive horizontal exponent is absent.
- On the vertical line, $a_{00} = a_{01} = 0$ force $\alpha = D = 0$. A negative vertical exponent requires $n > 0$; then $a_{21} = 0$ forces $\beta = 0$, giving $a_{12} = 0$ and no positive vertical exponent.
- On the line of slope -1, the two opposite coefficients $a_{02}, a_{20} > 0$ require $m, D, n, A > 0$, contradicting the required off-line equality $a_{01} = 0$.
- On the line of slope $+1$, the opposite coefficients require $\alpha, \beta, c > 0$. The off-line equations then force $m = D = n = A = 0$. Together with the hard λ event the support is the perfect matching $\{\lambda, c, \alpha, \beta\}$.

The last case is also impossible for a nonnegative constrained source. On its perfect-matching curve all event squares and the signed core vanish. The signed-core differential is nonzero on the tangent space to the sole hard constraint. To check this invariantly, on the symmetry-equivalent matching $\{n, m, A, D\}$ use its four independent event-normal differentials (dX, dY, dR_m, dR_D); the core differential is

$$2t\{(1 + t^2)dX - (1 + t^2)dY - dR_m + dR_D\}.$$

All four coefficients are nonzero. Removing only one hard direction leaves a nonzero derivative, while every event square has zero derivative. The constrained slack therefore takes a negative value on one side of that curve, a contradiction.

Finally suppose no outer exponent is active. Then $\alpha = D = n = 0$ and $\beta m = cA = \beta c = 0$. If $\beta > 0$, all complementary active events are contained in $\{A, \beta\}$; if $\beta = 0$ and $c > 0$, they are contained in $\{m, c\}$; otherwise they are contained in $\{m, A\}$. Including the hard event, these are the three nonextendable triples

$$\{\lambda, A, \beta\}, \quad \{\lambda, m, c\}, \quad \{\lambda, m, A\}.$$

Their exact kernel states in Section 5.1 have strictly negative signed core. Substitution annihilates every event and contradicts $d > 0$.

Thus the logarithmic determinant Hessian is positive definite at every constrained zero. The two positive state eigenvalues give a positive state block; its ordinary real Schur complement identifies the two-angle Hessian with a positive multiple of the determinant Hessian. Hence the constrained slack has a positive definite full Hessian in its two state and two angle variables. This Schur complement is used only over the real residue field, not as a tensor inverse.

The diagonal weighted-contact argument, with hard ratio σ/λ, now applies. Its soft residue is either positive or has this strict four-variable minimum. It gives an exact full-parent Gram and restores the pure source scale, proving the assertion. □

Chapter 10

Two hard events and a positive next-scale signed coefficient

Component `pf47f6a4e3a60`
Source: `CORE/prior/inherited/zero_d_two_hard.tex`
SHA256: `f47f6a4e3a605bdad562f60340997111f37604cfa2128b800715a6bc1ffaf074`
Inventory use: prop:two-hard-event | two-hard-event constrained curvature; prop:two-hard-event | C14_ ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

10.0.1 Two hard events and a positive next-scale signed coefficient

Proposition 10.1 (The two-hard-event terminal)**.** *Let the source be physical. Suppose two independent matching event weights a_1, a_2 satisfy $\sigma/a_i \in \mathfrak{m}_c$, where σ is the largest remaining source coefficient, and d/σ has positive residue. Then the complete parent is tensor SOS at every noncommuting residue test annihilating those two events.*

Proof. The two hard weights need not be comparable. Restrict the normalized residue source to their exact state-linear kernels. This is a nonnegative quadratic form on a two-dimensional state space. Its intersection with $Z = 0$ has dimension at least one, and the signed core is positive definite there. At a zero the constrained state matrix therefore has one positive eigenvalue and one simple zero eigenvalue; its kernel has $Z \neq 0$.

There are three symmetry classes of hard pairs. In the eight-cycle of losing events, two matching edges have distances two, three, or four. Representatives are respectively

$$\{\lambda, c\}, \qquad \{\lambda, m\}, \qquad \{\lambda, \alpha\}.$$

Extract the coefficient of the product of the two hard weights from each outer coefficient q_{ij} of the determinant dictionary. Each resulting coefficient is nonnegative. Since $d > 0$, its exact positivity conditions are the following; a juxtaposition means that both factors are positive, and a disjunction means at least one listed condition holds.

$(i-1, j-1)$	$\{\lambda, m\}$	$\{\lambda, c\}$	$\{\lambda, \alpha\}$
$(-1,-1)$	α	α	always
$(-1, 0)$	D or α	D or αm	D or m
$(-1, 1)$	D	Dm	Dm
$(0,-1)$	n or $A\alpha$	n	A or n
$(0, 1)$	β or Dc	D or βm	Dc or βm
$(1,-1)$	An	0	An
$(1, 0)$	Ac or βn	A	Ac or βn
$(1, 1)$	βc	β	βc

For example, apart from the common positive factor 4, the first two coefficients in the $\{\lambda, m\}$ column are

$$2\alpha(2A + 2D + 2c + d), \qquad 2(2AD + 2A\alpha + 4D\alpha + 2Dc + Dd + 2\alpha c + \alpha d).$$

The table follows by differentiating the eight displayed quadratic-factor formulas, with no determinant recomputation.

A state basis adapted to the two independent hard amplitudes shows that this leading determinant coefficient is a positive angle-dependent factor times the determinant of the constrained two-by-two state form. Thus, after its positive monomial factor is removed, the constrained determinant is a Laurent polynomial with a constant central coefficient and precisely these eight nonnegative outer coefficients. At any zero its logarithmic gradient vanishes. If its active exponent vectors do not span the plane, either none is active or they lie on one of the four opposite-pair lines.

Here the table gives a short exhaustive classification.

- For $\{\lambda, m\}$, neither diagonal line is possible: its negative diagonal coefficient activates respectively the off-line exponent $(-1, 0)$. On the horizontal line, D or α together with Ac or βn activates an off-line exponent in all four cases. On the vertical line, n or $A\alpha$ together with β or Dc does the same. With no active exponent, $\alpha = D = n = \beta = 0$ and $Ac = 0$, so the remaining support is contained in $\{A\}$ or $\{c\}$.

- For $\{\lambda, c\}$, the slope-minus-one line is impossible since its positive opposite coefficient is identically zero. A horizontal pair activates D or αm, hence an off-line exponent; a vertical pair activates n and either D or βm, with the same consequence. The slope-plus-one pair forces $\alpha, \beta > 0$ and $D = n = A = m = 0$. With no active exponent, all remaining weights except possibly m vanish.

- For $\{\lambda, \alpha\}$, the exponent $(-1, -1)$ is always active. A one-dimensional stationary support must therefore be the slope-plus-one pair. Its other end forces $c, \beta > 0$, while the off-line coefficients force $D = m = A = n = 0$. Rank zero is impossible.

The rank-zero possibilities are contained in the triples $\{\lambda, m, A\}$ and $\{\lambda, m, c\}$. At $u = v = 1$ the states

$$(W, X, Y, Z) = (1, -1, 0, 2), \qquad (1, -1, 1, 3)$$

respectively annihilate all their event amplitudes, including both hard events. Their signed-core values are -4 and -8. Thus neither can be a nonnegative constrained source.

The only rank-one possibilities complete the perfect matching $\{\lambda, c, \alpha, \beta\}$. On its exact matching curve, all event squares and the signed core vanish. The four event-normal differentials are independent. As shown in Proposition 9.1, the signed-core differential has a nonzero coefficient in every one of those four directions. Imposing only two hard constraints leaves at least one nonzero such derivative, whereas every remaining event square has zero differential. The constrained slack changes sign, contradicting physicality.

Consequently the active exponent vectors span the plane at every constrained zero. The logarithmic determinant Hessian is positive definite. The single positive soft-state pivot identifies its angle Hessian, up to positive factors and an invertible coordinate change, with the Schur complement of the full constrained Hessian. Thus that Hessian is positive definite in one projective state variable and two angle variables. All Schur operations in this argument take place only over the real residue field.

Finally use the two exact hard event amplitudes as coordinates. If their weights are a_1, a_2, put $x_i = (\sigma/a_i)s_i$ in the coefficient-field stationary equations. The ratios σ/a_i are infinitesimal, and the divided hard equations have positive diagonal residue. The diagonal weighted-contact lemma therefore applies with roots $\sqrt{a_i/\sigma}(x_i - x_{ci})$, together with the three soft deviations. Its exact Taylor Gram has strict residue, and its stationary value is nonnegative by physicality. Restoring the pure coefficient scale proves the full-parent assertion. □

Chapter 11

A finite induction on event constraints

Component `p94dfbc8a5970`
Source: `CORE/prior/inherited/zero_d_induction.tex`
SHA256: `94dfbc8a59706b5510ad8d8fda2f2744fa7190c77f45efedf3df401aeb8e26dc`
Inventory use: lem:diagonal-hard-scales; thm:all-noncommuting | diagonal hard-scale lemma and finite matching induction; lem:diagonal-hard-scales; thm:all-noncommuting | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

11.0.1 A finite induction on event constraints

Lemma 11.1 (Unequal diagonal hard scales)**.** *Suppose a physically nonnegative local polynomial has the form*

$$F(x,y)=\sum_{i=1}^{k} a_i b_i(y) x_i^2+\sigma G(x,y),$$

where $a_i,\sigma>0$ are pure coefficient numbers, $\varepsilon_i=\sigma/a_i$ are infinitesimal, and b_i, G have bounded coefficients. At the test let $\bar{x}=0$, $\bar{y}=y_0$ and $\bar{b}_i(y_0)>0$. If $g(y)=\overline{G}(0,y)$ is positive at y_0, or has a zero there with positive definite Hessian, then F is tensor SOS.

Proof. For a positive value use the roots $(x_i/\sqrt{\varepsilon_i},1)$, exactly as in Lemma 6.1. At a zero, physicality gives $\nabla g(y_0)=0$. Put $x_i=\varepsilon_i s_i$ in the stationary equations. They are

$$2b_i(y)s_i+G_{x_i}(\varepsilon s,y)=0, \qquad \sum_i \varepsilon_i s_i^2 \nabla b_i(y)+\nabla_y G(\varepsilon s,y)=0.$$

The residue Jacobian is block upper triangular with positive diagonal normal block $2\,\mathrm{diag}(\bar{b}_i(y_0))$ and tangent block $\nabla^2 g(y_0)\succ 0$. The coefficient-field implicit-function theorem gives an exact stationary center with $x_{ci}=\varepsilon_i s_{ci}$ and bounded s_c.

Exact Taylor allocation on the roots

$$((x_i-x_{ci})/\sqrt{\varepsilon_i})_{i=1}^{k}, \quad y-y_c$$

has residue Gram $\mathrm{diag}(\bar{b}_1,\dots,\bar{b}_k,\nabla^2 g/2)\succ 0$. Indeed a hard mixed normal–tangent coefficient has an extra factor $\sqrt{\varepsilon_i}$, and a hard pure tangent coefficient has an extra factor ε_i, since $x_{ci}/\varepsilon_i=s_{ci}$ is bounded. The bounded G terms have the same or smaller factors. The center value $F(x_c,y_c)/\sigma$ is a nonnegative pure coefficient. Apply the strict Gram lemma and restore σ. Pure testing unit denominators are cleared as in Lemma 6.1. □

Theorem 11.2 (Every noncommuting residue test is atlas-free)**.** *For every physical canonical source, the entire parent is tensor SOS at every independent test for which both measurement pairs are noncommuting in residue. No restriction on the relative source coefficient scales is required.*

Proof. Normalize $d + \sum c_i = 1$. Literal $d = 0$ is an event-square sum. Positive slack residue is covered by Lemma 1.1; positive $\bar{d}$ is covered by Theorem 4.2 and Lemma 4.1. We may therefore assume $d > 0$, $\bar{d} = 0$, and zero slack residue.

A product residue state is already settled by Section 5.1. At an entangled residue, all positive-residue event coefficients belong to a matching in the eight-cycle. Call this first collection of hard events S. If $|S| = 4$, all complementary coefficients and d are infinitesimal relative to its four positive-unit weights, so Theorem 8.1 applies directly. Suppose $|S| \leq 3$.

The amplitudes of S are independent state-linear coordinates on a projective state chart. Choose them as x_i, and call the remaining state and angle coordinates y. The change of coordinates has only real testing denominators that are units at the chosen residue. For the next scale put

$$\sigma = \max(\{d\} \cup \{c_j : j \notin S\}).$$

Every hard coefficient a_i is larger than σ: the quotient σ/a_i is infinitesimal. The parent has the exact diagonal form of Lemma 11.1. On its hard kernel the normalized residue $g = \overline{G}(0, y)$ is nonnegative by physicality.

If $g(y_0) > 0$, the lemma finishes. If $g(y_0) = 0$ and $\overline{d/\sigma} > 0$, there are only three possibilities. With one or two hard events, Propositions 9.1 and 10.1 give a strict constrained minimum. With three hard events, the two Laurent restrictions of Theorem 6.2 give the same conclusion. Their calculations concern only g and hence do not depend on whether the hard coefficients are comparable. Lemma 11.1 finishes in each case.

The only alternative is $g(y_0) = 0$ and $\overline{d/\sigma} = 0$. Then g is a nonnegative event-square sum, and at least one remaining event coefficient has positive residue after division by σ. Its amplitude must vanish at the same entangled residue state. Add every such event to S. The enlarged set is still a matching, and every coefficient left outside it is infinitesimal relative to all its weights.

If the enlarged set has at most three edges, repeat the preceding step. If it has four, it is a perfect matching. All complementary coefficients and d are then infinitesimal relative to its smallest weight, exactly the hypotheses of Theorem 8.1. Each repetition adds an edge, and a matching has at most four edges. The induction therefore terminates after at most three enlargements. Every step retains the complete parent; none discards an infinitesimal signed or event term. Pure-field restoration returns the original normalized slack. □

The role of the eight-cycle is now structural: it bounds the number of independent event constraints. Laurent curvature handles the first scale where the signed term becomes visible, and the four-event terminal handles the case where all four constraints become visible first. This replaces the noncommuting part of the source atlas by a finite rank argument.

Chapter 12

A short proof for every noncommuting residue

Component `p2c3fc22497a5`
Source: `CORE/prior/inherited/penpaper_noncommuting.tex`
SHA256: `2c3fc22497a59086acf6bc93ff5f5408d84300149843aa9d4f6239fb6847d29a`
Inventory use: full exposition | entire noncommuting-residue proof; full exposition | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

12.1 A short proof for every noncommuting residue

The noncommuting part of the argument has a simple organizing principle: *a vanishing event constrains an entangled state, and there are only four compatible such constraints.* Coefficients can occur at arbitrarily many relative scales, but the proof only changes when a new independent constraint appears. The number of relevant changes is therefore bounded by geometry.

This section gives a short reading route through Theorem 11.2. Its conclusion concerns independent tests whose two measurement pairs are noncommuting *in residue*. It does not include infinitesimal approaches to commuting measurements, and it is not by itself the global finite SDP theorem.

12.1.1 What must be certified

Keep the coefficient field K_c and the testing field K_t separate, and put $T = K_c \otimes_{\mathbb{R}} K_t$. A source is physical when its Bell slack is nonnegative at every experiment over K_c. The assertion to prove is stronger than that ordinary positivity: its evaluation at an independent experiment over K_t must lie in ΣT^2.

Write the entire parent in the form

$$F = \sum_e c_e\, b_e(u,v) R_e(W,X,Y,Z;u,v)^2 + d\, C, \qquad c_e \geq 0, \quad d \geq 0. \tag{12.1}$$

Here $C = D_u D_v \langle H \rangle$, with the state norm left homogeneous; the factors b_e are the positive testing factors in (P), including its factor four. The R_e are the eight measurement amplitudes. For example,

$$R_n = X, \quad R_A = Y, \quad R_m = vW - X + uvY - uZ, \quad R_D = uW + uvX - Y - vZ.$$

Normalize $d + \sum_e c_e = 1$, unless the source is zero. Choose a bounded projective state chart and finite half angles whose residues satisfy $u_0 v_0 \neq 0$. These choices use only pure testing factors.

Only one algebraic certificate is needed throughout. If a finite identity has the form

$$F/\sigma = a + h^T Q h, \qquad a \in K_c, \quad a \geq 0, \quad Q \in \mathrm{Sym}(B_c \otimes B_t), \quad \bar{Q} \succ 0, \tag{12.2}$$

where $\sigma > 0$ is pure coefficient, then the strict Gram lemma gives $F \in \Sigma T^2$. The entries of h need not be bounded. In particular, rescaling a normal coordinate by a large *pure coefficient* is allowed. The argument never takes an inverse or a square root of a mixed tensor.

There are two immediate cases. Positive slack residue is covered by Lemma 1.1. Literal $d = 0$ makes (12.1) an event-square sum. All subsequent steps keep every packet of the complete parent, including its infinitesimal terms.

12.1.2 Entanglement turns the events into a matching

For a product state, the CHSH expectation obeys

$$S = a_0(b_0 - b_1) + a_1(b_0 + b_1) \le |b_0 - b_1| + |b_0 + b_1| = 2\max(|b_0|, |b_1|).$$

At noncommuting residue angles equality $S = 2$ is impossible. If both Bob moduli are below one it is already strict. If one equals one, the other is strictly smaller, and equality would require a common eigenstate of Alice's two noncommuting reflections. Thus the signed core has positive testing residue at every product residue state. Its actual testing value is positive, so dC is a square after separate field square roots. Such tests are settled.

Now suppose $\bar{d} = 0$, the slack residue vanishes, and the residue state is entangled. Write that real state as an invertible 2×2 matrix Ψ. For any fixed Alice ray a, the equation

$$a^T \Psi b = 0$$

selects exactly one Bob projective ray, and conversely. Two vanishing events cannot therefore share a local outcome ray. The losing events occur in a single bipartite cycle of local outcome rays, with cyclic edge order

$$(\lambda, D, c, A, \alpha, m, \beta, n).$$

Consequently every set of vanishing events is a matching, of size at most four. This is the finiteness mechanism.

Any three matching event vectors are state-linearly independent. Indeed, the span of two rank-one matrices with independent left and right factors contains no third rank-one projective point. A dependence among three rank-one matrices must therefore have a common left or right factor; matching edges cannot do so. For at most three vanishing events, their amplitudes may hence be used as exact local state coordinates. A nonzero minor at the chosen residue supplies the inverse change of coordinates; its denominator is a pure testing unit. This is an exact coordinate change, not a linearization.

Four matching edges give one of two perfect matchings. In the representative $\{n, m, A, D\}$, the equations are

$$X = Y = 0, \qquad vW = uZ, \qquad uW = vZ.$$

Thus $u^2 = v^2$. Choosing its sign and putting $W = 1$ gives the smooth curve $v = u$, $Z = 1$. It has four independent normal directions: three state directions and the angle difference. The other perfect matching is its CHSH-symmetric counterpart. Section 5.1 establishes these statements directly from the event amplitudes.

12.1.3 Unequal scales are handled one normal coordinate at a time

Suppose $k \le 3$ event weights $a_1, \dots, a_k$ have already been identified, and every remaining coefficient is infinitesimal relative to each a_i. Let σ be the largest remaining coefficient, including d, and put

$$\varepsilon_i = \sigma / a_i \in \mathfrak{m}_c.$$

In exact event coordinates the parent has the form

$$F(x, y) = \sum_{i=1}^{k} a_i b_i(y) x_i^2 + \sigma G(x, y), \qquad \bar{b}_i(y_0) > 0, \tag{12.3}$$

with bounded coefficients after clearing pure testing unit denominators. The variables y comprise the remaining projective state and angle coordinates. On the exact hard kernel $x = 0$, physicality implies

$$g(y) := \overline{G}(0, y) \geq 0.$$

If $g(y_0) > 0$, the roots $(x_i/\sqrt{\varepsilon_i}, 1)$ immediately give a strict residue Gram. If $g(y_0) = 0$ and $\nabla^2 g(y_0) \succ 0$, solve for an exact coefficient-field stationary center. The correct scaling is $x_i = \varepsilon_i s_i$, separately for every normal. Dividing the stationary equations gives

$$\begin{aligned} 2b_i(y)s_i + G_{x_i}(\varepsilon s, y) &= 0, \\ \sum_i \varepsilon_i s_i^2 \nabla b_i(y) + \nabla_y G(\varepsilon s, y) &= 0. \end{aligned}$$

Their residue Jacobian is block upper triangular, with diagonal blocks

$$2\,\mathrm{diag}(\bar{b}_i(y_0)), \qquad \nabla^2 g(y_0).$$

Both are positive definite. The semialgebraic implicit-function theorem, transferred to K_c, gives (x_c, y_c) with $x_{ci} = \varepsilon_i s_{ci}$ and bounded s_c.

Finite Taylor expansion at this exact center, on the roots

$$h = \big((x_i - x_{ci})/\sqrt{\varepsilon_i}\big)_{i=1}^{k} \oplus (y - y_c),$$

has residue Gram

$$\bar{Q} = \mathrm{diag}\,(\bar{b}_1, \ldots, \bar{b}_k, \tfrac{1}{2}\nabla^2 g(y_0)) \succ 0.$$

Hard normal–tangent terms acquire a factor $\sqrt{\varepsilon_i}$; hard pure tangent terms acquire ε_i. Higher Taylor terms leave an actual infinitesimal deviation in their coefficient. The constant $F(x_c, y_c)/\sigma$ is nonnegative by physicality, giving (12.2). This proves Lemma 11.1 and explains why no comparability of the a_i is needed.

The implicit-function step is local in a fixed finite-dimensional real parameter space. Its existence and uniqueness statements transfer to the coefficient field. It is not an infinite asymptotic expansion or an assumption of compactness over a non-Archimedean field.

12.1.4 When the signed term appears, physicality forces curvature

Suppose $\overline{d/\sigma} > 0$ at the next scale. With no hard constraint, or with one or two hard event constraints, the constrained state form is strictly positive on its subspace $Z = 0$. At a zero it therefore has a simple kernel and positive complementary state pivots.

The remaining angular curvature is encoded by one small determinant. The parent determinant is a positive angular factor times $UVL(U, V)$, where $U = u^2$, $V = v^2$, and

$$L(U, V) = h + \sum_{e \in E} q_e U^{e_1} V^{e_2}, \qquad q_e > 0, \qquad E \subset \{-1, 0, 1\}^2 \setminus \{(0, 0)\}. \tag{12.4}$$

All outer coefficients are nonnegative, as follows from the factored determinant formulas in Section 3.1. Constraining one or two event amplitudes amounts to taking the coefficient of their weights in this determinant: in an adapted state basis it is the determinant of the remaining state form, times a positive factor. Hence it retains the same Laurent structure.

At a physical zero, logarithmic stationarity and curvature are exactly

$$\sum_{e \in E} w_e e = 0, \qquad \nabla^2_{\log U, \log V} L = \sum_{e \in E} w_e e e^T, \qquad w_e = q_e U^{e_1} V^{e_2} > 0. \tag{12.5}$$

Thus strict angular curvature is equivalent to the active exponent vectors spanning the plane. There is no eigenvalue estimate to compute.

For clarity, the finite support argument establishing that span has only two possible obstructions. A stationary support of rank one must lie on one of the four opposite-pair lines of the Newton

square. Substituting the factored coefficients shows that it either activates an exponent off that line, or leaves a perfect matching of events. Rank zero leaves a support contained in a nonextendable matching triple. For the unconstrained rank-zero case, the forbidden pairs form this event cycle together with its four opposite chords. An independent triple must have cyclic gaps $(2, 3, 3)$: the only other possible gap pattern $(2, 2, 4)$ contains an opposite pair. Rotation of the cycle carries all such triples to $\{\lambda, m, c\}$. Indeed the CHSH maps g_T and g_B act on edge indices as $i \mapsto 2 - i$ and $i \mapsto 3 - i$ modulo eight, so $g_B g_T$ is a one-edge rotation. The representative has the exact kernel state

$$\psi_H = (uv, -u, u^2 v, 1 + v^2 + u^2 v^2), \qquad C(\psi_H) = -2D_u D_v u^2 v^2 < 0.$$

The rank-one perfect-matching case has a matching curve on which the core is zero but its differential is nonzero in every event-normal direction. In the representative matching, with $B = 1 + t^2$, that differential is

$$2t(B\,dX - B\,dY - dR_m + dR_D).$$

Imposing at most two event constraints leaves a direction with nonzero core derivative. All event squares have zero derivative there, so the constrained parent would change sign. Either obstruction contradicts physicality. The short exact support arguments are proved in Theorem 4.2 and Propositions 9.1–10.1; they require only the displayed determinant factors and these kernel witnesses.

For three hard events there is no state variable left, and there are only two matching orbits. Their exact kernel states can be chosen as

$$\psi_N = (uv, -u, 0, 1 + v^2), \qquad \psi_E = (u, 0, 0, v).$$

After division by positive testing factors, their restrictions are

$$L_N = \frac{2n(1+U)}{V} + 2cU + \frac{2D(1+V)}{U} + \frac{2\alpha(1+U)(1+V)}{UV} + 2\beta V - d,$$
$$L_E = d(U/V - 1) + \frac{2\lambda(1+U)}{V} + \frac{2cU(1+V)}{V} + \frac{2D(U-V)^2}{UV} + \frac{2\alpha(1+U)}{U} + 2\beta(1+V).$$

For L_N, a positive n, D or α already supplies two independent exponents; if they all vanish, the remaining positive exponentials cannot have a stationary zero. For L_E, $d > 0$ supplies $(1, -1)$. Failure of two-dimensional span forces $\lambda = c = \alpha = \beta = 0$; the remaining one-variable expression has minimum

$$2\sqrt{2D(d+2D)} - d - 4D < 0, \qquad (d+4D)^2 - 8D(d+2D) = d^2 > 0.$$

Again physicality excludes the obstruction. Multiplication by a positive factor preserves the Hessian sign at a stationary zero. The positive state pivots, when present, then promote angular curvature to the full constrained Hessian by an ordinary real Schur complement. This Schur complement is used only in the residue field. The weighted argument above now supplies the exact tensor certificate.

12.1.5 Four constraints leave one Laurent variable

It remains to understand what happens if all four matching events appear before the signed term does. Take the matching $\{n, m, A, D\}$ and, by symmetry, choose its smallest weight to be $D = \mu$. The hypotheses are

$$d/\mu \in \mathfrak{m}_c, \qquad \lambda/\mu, c/\mu, \alpha/\mu, \beta/\mu \in \mathfrak{m}_c.$$

Set $\eta_1 = \mu/n$, $\eta_2 = \mu/A$, $\eta_3 = \mu/m$, so $0 < \eta_i \leq 1$; their residues may be zero. Near the matching curve use $W = 1$ and exact coordinates

$$R = v - X + uvY - uZ, \qquad Z = (v - X + uvY - R)/u, \qquad \delta = v - u.$$

The fourth amplitude equals $f + L$, where

$$f = (u^2 - v^2)/u, \qquad L = v(u + u^{-1})X - (1 + v^2)Y + (v/u)R.$$

On $X = Y = R = 0$ the signed-plus-fourth packets are exactly

$$4\mu f^2 + 2duf = 4\mu\left(f + \frac{du}{4\mu}\right)^2 - \frac{d^2u^2}{4\mu}. \tag{12.6}$$

Let $\tau = \max(\lambda, c, \alpha, \beta)$. Testing the negative minimum in (12.6) at $u = 1$ proves

$$\tau > 0, \qquad \rho = \frac{d^2}{\mu\tau} \text{ is bounded.}$$

Thus physicality itself bounds the first adverse normal correction by the complementary event scale.

Put $\varepsilon = d/\mu$ and scale the four stationary normal coordinates by

$$X = \varepsilon\eta_1 s_1, \quad Y = \varepsilon\eta_2 s_2, \quad R = \varepsilon\eta_3 s_3, \quad \delta = \varepsilon r, \qquad u = t.$$

For $\bar{\rho} > 0$, divide their stationary equations by d. Writing $B = 1 + t^2$, $\ell = (B, -B, 1)$, $b = (B^2, B^2, 1)$ and $k = -2r + \sum_i \bar{\eta}_i \ell_i s_i$, their residues are

$$8b_i s_i + 8k\ell_i + (4tB, -4tB, 0)_i = 0, \qquad -16k - 4t = 0.$$

Subtracting suitable multiples of the last row leaves positive diagonal normal pivots $8b_i$ and 32. In particular the system remains nonsingular when any $\bar{\eta}_i$ vanishes. Its solution is

$$s_1 = -\frac{t}{4B}, \quad s_2 = \frac{t}{4B}, \quad s_3 = \frac{t}{4}, \quad r = \frac{t}{8}(1 - \bar{\eta}_1 - \bar{\eta}_2 + \bar{\eta}_3).$$

After this normal completion, the residue potential along the matching curve is exactly

$$g(t) = 4(1 + t^2)(P + Qt^2) - \frac{\bar{\rho}t^2}{4}(1 + \bar{\eta}_1 + \bar{\eta}_2 + \bar{\eta}_3),$$

where $P = \overline{(\lambda + \alpha)/\tau}$, $Q = \overline{(c + \beta)/\tau}$ and $P + Q > 0$. Its quotient by $4t^2$ is

$$P(1 + t^{-2}) + Q(1 + t^2) - \frac{\bar{\rho}}{16}(1 + \bar{\eta}_1 + \bar{\eta}_2 + \bar{\eta}_3). \tag{12.7}$$

Physicality makes this nonnegative. At a zero, stationarity forces $P/t^2 = Qt^2$, so $P, Q > 0$ and the logarithmic second derivative is positive. There is therefore a strict tangent pivot as well as the four strict normal pivots. If $\bar{\rho} = 0$, the complementary event value is already strictly positive and the direct normal Gram suffices.

For the exact restoration, the roots at a coefficient-field stationary center are

$$\left(\sqrt{n/\tau}(X - X_c), \sqrt{A/\tau}(Y - Y_c), \sqrt{m/\tau}(R - R_c), \sqrt{\mu/\tau}(\delta - \delta_c), t - t_c\right).$$

The key bounds are transparent: for a normal weight $a_i = \mu/\eta_i$,

$$\frac{a_i x_{ci}^2}{\tau} = \rho\eta_i s_{ci}^2, \qquad \frac{a_i x_{ci}}{\sqrt{a_i\tau}} = \sqrt{\rho\eta_i}\, s_{ci}.$$

Every quantity on the right is bounded. Exact packetwise Taylor allocation therefore gives a strict Gram, whose tangent Schur complement is $g''/2$. Theorem 8.1 supplies the full details. No inverse of η_i is needed in a bounded Gram coefficient.

12.1.6 The finite induction

Start with all event coefficients having positive residue. If the residue slack is zero and $\bar{d} = 0$, they form a nonempty matching S at the entangled residue state. For $|S| \le 3$, take the next scale

$$\sigma = \max\big(\{d\} \cup \{c_e : e \notin S\}\big).$$

Every weight already in S dominates σ. Restricting to their exact event kernel gives the next-scale nonnegative residue g.

If g is positive, the weighted certificate finishes. If $g = 0$ and d/σ has positive residue, the Laurent curvature argument finishes. Otherwise g is an event-square sum. At least one remaining coefficient has positive residue after division by σ, and every such event must vanish at the same entangled residue state. Add all of them to S. The enlarged set is still a matching, and every coefficient outside it is infinitesimal relative to each of its weights.

Each repetition adds an edge. A matching has at most four edges, so there are at most three enlargements. On reaching four edges the preceding one-variable terminal applies, independently of their relative weights. If $\bar{d} > 0$ at the outset, the unconstrained Laurent curvature theorem and the regular-contact certificate already apply. These cases exhaust all noncommuting residue tests and prove Theorem 11.2.

The insight is that the proof counts constraints, not coefficient regimes. The event cycle supplies a finite depth; physicality supplies curvature at the first visible signed scale; exact Taylor expansion converts that curvature into a separated-field sum of squares. The commuting boundary requires additional arguments because the outcome rays then coalesce and the matching argument no longer controls the normal directions.

Chapter 13

All closed faces from two elementary matrix lifts

Component `pd9a6195f0af3`
Source: `CORE/prior/inherited/closed_faces_short.tex`
SHA256: `d9a6195f0af36c54955f1490a9e8e8f3b015b969326031a8db49943c7457ac6a`
Inventory use: Two explicit noncoercive face lifts and exact finite matrix-lift proof; re-read in full

13.1 All closed faces from two elementary matrix lifts

The noncoercive faces do not require a list of support masks. The physical classification in Proposition 3.1 leaves just two kinds of omitted exclusive pairs, and both admit short exact lifts.

Lemma 13.1 (Two kinds of exclusive pair)**.** *Every pair of exclusive losing events is CHSH-equivalent either to $\{\lambda, \alpha\}$ or to $\{\lambda, m\}$. Consequently every noncoercive source belongs to a relabeling of one of these two closed coefficient faces.*

Proof. If the two events have the same input pair, they are the two opposite losing outcomes at that input. There are four such pairs. Applying g_A, g_B from Section 3.1 to $\{\lambda, \alpha\}$ gives $\{m, D\}$ and $\{n, A\}$; applying g_A after g_B gives $\{c, \beta\}$.

Otherwise exclusivity means opposite outcomes at one shared local measurement and different remote settings. Each of the four local measurements gives two such pairs. Starting from $\{\lambda, m\}$, the words

$$1, \quad B, \quad T, \quad BA, \quad BT, \quad TA, \quad TB, \quad BAB$$

in g_A, g_B, g_T, applied from left to right, give respectively

$$\{\lambda, m\},\ \{A, \beta\},\ \{c, m\},\ \{c, n\},\ \{\alpha, n\},\ \{A, \lambda\},\ \{D, \beta\},\ \{D, \alpha\}.$$

These are all eight possibilities. Every noncoercive omission set contains an exclusive pair by Proposition 3.1; deleting further coefficients only takes a closed subface. □

We use the following elementary matrix fact. Its proof is included to avoid importing a matrix factorization theorem.

Lemma 13.2 (An exact quadratic matrix lift)**.** *Let $K(t) = K_0 + tK_1 + t^2K_2$ be a real symmetric r by r matrix polynomial. Then $K(t) \succeq 0$ for every real t if and only if there is a skew-symmetric S such that*

$$\begin{pmatrix} K_0 & K_1/2 + S \\ K_1/2 - S & K_2 \end{pmatrix} \succeq 0. \tag{CF1}$$

Proof. The reverse direction is obtained by evaluation on (z, tz). For the forward direction let $\mathcal{C}$ be the cone of coefficient triples produced by positive semidefinite Gram matrices. This cone is closed: for a convergent sequence of coefficient triples the Gram trace equals $\operatorname{tr}(K_0 + K_2)$, so a sequence of realizing positive semidefinite Grams is bounded and has a convergent subsequence.

A linear functional nonnegative on $\mathcal{C}$ has the form

$$\ell(K) = \operatorname{tr}(UK_0) + \operatorname{tr}(VK_1) + \operatorname{tr}(WK_2), \qquad H = \begin{pmatrix} U & V \\ V & W \end{pmatrix} \succeq 0, \tag{CF2}$$

where U, V, W are symmetric. Positivity of H implies $\operatorname{ran} V \subseteq \operatorname{ran} U$ and $W - VU^{+}V \succeq 0$, where U^{+} is its inverse on its range and zero on its kernel. On $\operatorname{ran} U$ diagonalize the symmetric matrix $U^{+/2}VU^{+/2}$. If (t_j, e_j) are its orthonormal eigenpairs and $z_j = U^{1/2}e_j$, then

$$\begin{pmatrix} U & V \\ V & VU^{+}V \end{pmatrix} = \sum_j \begin{pmatrix} z_j \\ t_j z_j \end{pmatrix} \begin{pmatrix} z_j \\ t_j z_j \end{pmatrix}^{T}. \tag{CF3}$$

The remaining block $W - VU^{+}V$ is a sum of $w_i w_i^{T}$. Thus

$$\ell(K) = \sum_j z_j^{T} K(t_j) z_j + \sum_i w_i^{T} K_2 w_i \geq 0,$$

because $K_2 \succeq 0$ follows by taking the leading coefficient at infinity. Finite-dimensional separation of the closed convex cone $\mathcal{C}$ now gives membership in $\mathcal{C}$, which is (CF1). This proof also handles $U = 0$ and singular U literally. □

For each fixed matrix size the lemma is a first-order statement in ordered-field arithmetic. It therefore holds over every real closed coefficient field as well. All changes of variables used below are within that one field; no inverse in a tensor product is introduced.

Theorem 13.3 (The missing closed face has a six-dimensional lift)**.** *On the closed canonical face $\lambda = m = 0$, with every remaining event coefficient and d nonnegative, physicality is equivalent to (CF1) for $r = 3$ and*

$$\begin{aligned} K_0 &= \operatorname{diag}(4c + 4D + 2d,\ 4A,\ 4\alpha), \\ K_2 &= \operatorname{diag}(4c,\ 4A + 4\alpha + 2d,\ 4D), \\ K_1 &= \begin{pmatrix} 0 & 0 & -(4D + 2d) \\ 0 & 0 & -(4\alpha + 2d) \\ -(4D + 2d) & -(4\alpha + 2d) & 0 \end{pmatrix}. \end{aligned} \tag{CF4}$$

There are only three auxiliary variables, the independent entries of S. In particular this is an affine semidefinite lift, with no nonlinear inequalities in the source coefficients.

Proof. The case $d = 0$ is immediate from the original event-square sum and from the positive-square expression below, so first suppose $d > 0$. In the parent set $W = a/u$ and send u to zero while holding a, X, Y, Z, v fixed. Put $R = a - Y$. The limiting form is a nonnegative multiple $2t_5(1 + v^2)X^2$, with $t_5 = d + 2(\beta + n) > 0$, plus

$$\begin{aligned} G_v(R, Y, Z) = {} & 4c(1 + v^2)R^2 + 4A(1 + v^2)Y^2 + (4D + 2d)(R - vZ)^2 \\ & + (4\alpha + 2d)(vY - Z)^2 - 2d(1 + v^2)Z^2. \end{aligned} \tag{CF5}$$

Its matrix is precisely $K(v)$ from (CF4). Homogeneity of the state makes this limiting test legitimate. Hence physicality implies $K(v) \succeq 0$ for every real v.

Conversely, in this face the Laurent determinant is

$$L(U, V) = h + q_{10}/V + q_{12}V + U(q_{20}/V + q_{21} + q_{22}V), \tag{CF6}$$

and every displayed q is nonnegative. The same boundary limit in the determinant identity of Section 3.1 gives exactly

$$\det K(v) = \frac{8(1+v^2)}{t_5}(q_{10} + hv^2 + q_{12}v^4). \tag{CF7}$$

Indeed the transformed parent matrix has determinant $\det P/u^2$, and its limit is $2t_5(1+v^2)\det K(v)$. Thus (CF7) requires no expansion of a new determinant. If $K(v) \succeq 0$, (CF7) makes the first three terms of (CF6) nonnegative for $V > 0$; all the terms multiplied by U are nonnegative. The determinant criterion, whose three hard eigenvalues are positive for $d > 0$ and $uv \neq 0$, proves positivity of the full parent. Polynomial continuity supplies every finite or projective angle endpoint. Lemma 13.2 now proves the asserted lift.

For clarity, no strictness or nonzero event coefficient was needed. When $d = 0$, (CF5) is itself a sum of squares linear in v, so it directly supplies a Gram in (CF1). Literal zeros are therefore included without a limiting source argument. □

The lift also has a strictly feasible coefficient point. Set $c = A = 2$, $D = \alpha = d = n = \beta = 1$, and $S = 0$. A permutation of its six coordinates gives two three-dimensional blocks. In each, the two diagonal hard pivots are $14, 8$ and their Schur complement is

$$4 - \frac{9}{14} - \frac{9}{8} = \frac{125}{56} > 0. \tag{CF8}$$

Thus the usual attained polar-duality argument applies to this lift; it does not require an approximate or numerical certificate.

Theorem 13.4 (The opposite-outcome face has two three-dimensional blocks)**.** *On the closed canonical face $\lambda = \alpha = 0$, define*

$$p = 4c + 4D + 2d, \qquad q = 4m + 4\beta + 2d, \qquad r = 4D + 2d.$$

Physicality is equivalent to the existence of one real ζ with

$$\begin{pmatrix} 4m & -4m & \zeta \\ -4m & q+4n & q \\ \zeta & q & q \end{pmatrix} \succeq 0, \qquad \begin{pmatrix} p+4A & 4D & -p \\ 4D & 4D & -(4m+r)-\zeta \\ -p & -(4m+r)-\zeta & p \end{pmatrix} \succeq 0. \tag{CF9}$$

All original coefficient signs are retained.

Proof. Set $u = \epsilon s$, $v = \epsilon$ and $(W, Z) = (w, z)/\epsilon$, while $X = x, Y = y$. The finite limit of the parent as ϵ tends to zero is

$$\begin{aligned} G_s = {} & 4nx^2 + 4m(w - x - sz)^2 + 4c(sw - y)^2 + 4Ay^2 \\ & + r(sw - y - z)^2 + (4\beta + 2d)(x + sz)^2 - 4dyz - 2dz^2. \end{aligned} \tag{CF10}$$

The divergent terms cancel between the positive signed packet and the negative core. Write $J(s)$ for the matrix of G_s on (w, x, y, z). Its leading coefficient is $\mathrm{diag}(p, 0, 0, q)$, and its determinant is

$$\det J(s) = 16(q_{02} + hs^2 + q_{20}s^4). \tag{CF11}$$

Again this follows directly from the original determinant identity: the state change multiplies the determinant by ϵ^{-4}, and only the displayed three lowest-order terms survive.

Physicality implies $J(s) \succeq 0$ for every real s. Conversely, if $d > 0$ and $J(s) \succeq 0$, (CF11) implies $q_{02}/s^2 + h + q_{20}s^2 \geq 0$ for $s \neq 0$. The full Laurent polynomial in this face is

$$h + q_{02}V/U + q_{20}U/V + q_{12}V + q_{21}U + q_{22}UV.$$

Take $s^2 = U/V$. Its first three terms are nonnegative and its remaining coefficients are nonnegative. The determinant criterion and continuity prove positivity of the entire parent. If $d = 0$, the parent and (CF10) are event-square sums, so the same equivalence is immediate.

Apply Lemma 13.2 to $J(s)$. Its Gram initially acts on $(w, x, y, z, sw, sx, sy, sz)$. The two diagonal entries for sx, sy are zero; positivity makes their entire rows and columns zero. The coefficient equations leave only one free off-diagonal entry, ζ. Permuting the remaining coordinates into (w, x, sz) and (y, z, sw) gives exactly the two blocks (CF9). Equivalently, direct multiplication of the displayed blocks gives the exact identity

$$G_s = (w, x, sz)B_1(w, x, sz)^T + (y, z, sw)B_2(y, z, sw)^T,$$

where B_1, B_2 are those blocks. The terms involving ζ cancel because $w(sz) = z(sw)$. This completes necessity as well as sufficiency of the affine lift. □

This lift is strictly feasible too: take $c = A = n = \beta = 2$, $m = D = d = 1$, and $\zeta = -5$. The leading principal minors of the two blocks are respectively $(4, 72, 234)$ and $(22, 72, 234)$, all positive integers.

Corollary 13.5 (The entire closed-face part is explicit)**.** *Every noncoercive canonical source belongs to a closed coefficient face with an explicit affine semidefinite lift. The two representatives are the two three-dimensional blocks (CF9) for $\lambda = \alpha = 0$ and the single six-dimensional block (CF1)–(CF4) for $\lambda = m = 0$. All further literal event zeros are included by linear restriction.*

Proof. Apply Lemma 13.1 and Theorems 13.3–13.4. Relabeling is an invertible linear change in the original Bell coordinates. The finite union of these coefficient faces is used only in the canonical source reduction; no union of spectrahedra is asserted to be convex. □

Chapter 14

Fresh finite-contact entry and the exact two-family source target

Component p477c0ac30750
Source: NEW/global/finite_contact_source_reduction_REDERIVED.tex
SHA256: 477c0ac30750df4bc299e543880f624aebf3b31d2e5516dbfd34769f54921b33
Inventory use: SE9 core state-sign implication used in CR and WA/WN | SE1-SE10: actual source coercivity, strict Hessian, physicality from stationary contact, deletion test; no tensor residue claim; SE9 core state-sign implication used in CR and WA/WN | SE1-SE15 fresh source-contact and canonical source target proof

14.1 Fresh finite-contact entry and the exact two-family source target

The input here is the actual recovered parent and its factored determinant, not an asserted completion of an atlas. We rederive the finite-contact nondegeneracy needed by the source-entry argument and state exactly which C1/C2 certificates would finish the global source reduction. A strict Hessian in a single source field is not identified with a strict real residue Gram in a tensor product.

14.1.1 Exact input and support incidence

Use the full parent (CAP1), with $d > 0$ and all eight event coefficients nonnegative. Its state matrix is $P(u, v)$. The physical dictionary is $F = D_u D_v \|\psi\|^2 f(q)$ at a nonzero state and its homogeneous extension at projective angles. This dictionary and the determinant identity below occur as actual proof bodies in the recovered `physical_core.tex`; the physical normalization was also freshly rederived in (RFA10)–(RFA12).

Write

$$\begin{aligned} t_1 &= d + 2(\lambda + n + D), & t_2 &= d + 2(\alpha + m + A),\\ t_3 &= d + 2(\lambda + C + D), & t_4 &= d + 2(\alpha + \beta + m),\\ t_5 &= d + 2(\lambda + \beta + n), & t_6 &= d + 2(\alpha + C + A),\\ t_7 &= d + 2(\beta + n + m), & t_8 &= d + 2(C + A + D). \end{aligned}$$

All t_j are strictly positive. The exact whole-parent determinant is

$$\det P = 16 D_u^2 D_v^2 u^2 v^2 L(u^2, v^2), \qquad L(U, V) = h + \sum_{e \in E} q_e U^{e_1} V^{e_2}, \quad E = \{-1, 0, 1\}^2 \setminus \{0\}, \quad \text{(SE1)}$$

where $h = q_{11}$ and the eight nonnegative coefficients are

$$\begin{aligned} q_{00} &= 4\lambda\alpha t_7 t_8, & q_{01} &= 4(\lambda D t_7 t_6 + \alpha m t_5 t_8),\\ q_{02} &= 4 m D t_5 t_6, & q_{10} &= 4(\alpha A t_7 t_3 + \lambda n t_8 t_4), \end{aligned}$$

$$q_{12} = 4(\beta m t_1 t_6 + CDt_5 t_2), \qquad q_{20} = 4nAt_3 t_4,$$
$$q_{21} = 4(\beta n t_3 t_2 + CAt_1 t_4), \qquad q_{22} = 4\beta C t_1 t_2.$$

Here q_{ij} has exponent $e = (i-1, j-1)$. In particular an exponent is active exactly when at least one of its displayed event products is positive. The checker for this section uses these exact incidence conditions; it does not replace physicality by a support enumeration.

Call the support *coercive* when 0 is in the interior of the convex hull of the active exponents. Over the real field this is equivalent to coercivity of $L(e^x, e^y)$: a positive exponential sum tends to infinity in every direction exactly under that condition. The condition on the finite support itself makes sense unchanged over any real closed coefficient field.

At $Z = 0$ the signed part of the full parent is

$$2d\{(uW + uvX - Y)^2 + D_u v^2 Y^2 + D_v X^2\}. \tag{SE2}$$

It is positive definite for $uv \neq 0$, so the corresponding 3×3 block H is positive definite, and $P \succeq 0 \iff \det P \geq 0$. If a nonzero state is in its kernel, that state has $Z \neq 0$ and the kernel has dimension one.

14.1.2 No physical finite zero has degenerate Newton support

Theorem 14.1 (Coercivity and strict curvature at every actual finite zero)**.** *Let a nonnegative canonical source with $d > 0$ be physical and have a zero on a nonzero state at finite $u_c, v_c \neq 0$. Then its Newton support is coercive. After the harmless real local basis gauges making $u_c, v_c > 0$ and the normalization $Z_c = 1$, the full Hessian of F in the five variables (W, X, Y, u, v) is positive definite at this contact. The assertion holds over every real closed extension of $\mathbb{R}$.*

Proof. Physicality makes the contact a zero minimum, so its first derivatives in the five variables vanish. By (SE1)–(SE2), $L = 0$ and its two logarithmic derivatives vanish. Put $a_e = q_e U_c^{e_1} V_c^{e_2} > 0$ for active exponents. Then

$$\sum_e a_e e = 0. \tag{SE3}$$

If the support were not coercive, a nonzero real linear functional would be nonnegative on it. Applying that functional to (SE3) shows that it vanishes on every active exponent. Thus all active exponents would lie on one line through the origin, or the support would be empty. There are only four possible such lines in E.

An active horizontal opposite pair would require both $q_{01}, q_{21} > 0$. Their four possible product pairings force, respectively, λn, CD, βm, or αA to be positive, activating q_{10} or q_{12} off that line. A vertical opposite pair $q_{10}, q_{12} > 0$ is ruled out in exactly the same explicit way: its pairings force αm, CA, βn, or λD positive, activating q_{01} or q_{21}. A one-sided line cannot satisfy (SE3).

For the diagonal pair $q_{00}, q_{22} > 0$, positivity forces $\lambda, \alpha, \beta, C > 0$, and all off-line coefficients vanish only if $n = m = A = D = 0$. For the other diagonal pair $q_{02}, q_{20} > 0$, the sole possibility is $n, m, A, D > 0$, $\lambda = C = \alpha = \beta = 0$. These are the two alternating four-event matchings of the cycle. Neither supports a physical source with $d > 0$. For the latter, set $u = v = 1$ and take the entire, untruncated state $\psi = (1, -\varepsilon, \varepsilon, 1)$. Direct substitution gives

$$F(1, 1; 1, -\varepsilon, \varepsilon, 1) = 16\varepsilon\{(d + n + m + A + D)\varepsilon - d\}. \tag{SE4}$$

Choosing $\varepsilon = d/[2(d + n + m + A + D)] > 0$ is a strict negative witness. The other matching is its image under the physical one-step event-cycle rotation (CAP13). The formula also allows literal zeros among the matching weights.

Finally suppose every outer coefficient vanishes. In the event cycle $(\lambda, D, C, A, \alpha, m, \beta, n)$, the forbidden simultaneously positive pairs are precisely the adjacent pairs and opposite pairs: these are the twelve products occurring in (SE1). Four pairwise nonadjacent vertices of an eight-cycle would have to alternate and would contain an opposite pair. Hence at most three events are active. Every maximal allowed set is a triple with cyclic gaps $(2, 3, 3)$; smaller allowed sets extend to one.

These eight triples are all rotations of $\{\lambda, m, A\}$. At $u = v = 1$, the state $(1, -1, 0, 2)$ annihilates those three event rows, while the signed core equals -4. Thus the full parent is $-4d < 0$. Rotation rules out every such support. This excludes the empty and rank-one possibilities, proving coercivity.

The angular log-Hessian at the contact is now

$$\mathcal{H}_L = \sum_e a_e e e^T \succ 0, \tag{SE5}$$

because the active exponent vectors span $\mathbb{R}^2$ and each a_e is positive. Write $P = \left(\begin{smallmatrix} H & b \\ b^T & c \end{smallmatrix}\right)$. Completing the three state coordinates within the single source field gives

$$F = (x + H^{-1}b)^T H (x + H^{-1}b) + s(u, v), \qquad s = c - b^T H^{-1} b = \det P / \det H. \tag{SE6}$$

At the contact $s = 0$ and $\nabla s = 0$. Equations (SE1), (SE5) show that its angle Hessian is the positive scalar $16 D_u^2 D_v^2 u^2 v^2 / \det H$ times the congruence of $\mathcal{H}_L$ by $\mathrm{diag}(2/u, 2/v)$. It is positive definite. The full five-variable Hessian has positive state block $2H$ and this positive angular Schur complement, so is positive definite too. This is an exact single-field Schur argument. Logarithms are merely notation for the operators $U\partial_U, V\partial_V$; (SE3), (SE5), and their Hessian congruence are algebraic ordered-field formulas. Alternatively the entire fixed finite assertion transfers from the real field. □

There is a useful converse contact implication, independent of coercivity. If nonnegative coefficients with $d > 0$ have a nonzero kernel state at $uv \neq 0$ and both angle derivatives of its parent vanish, (SE2) gives $L = 0$, $U\partial_U L = V\partial_V L = 0$. Over $\mathbb{R}$, convexity of $L(e^x, e^y)$ makes this zero its global minimum. The determinant criterion and the direct commuting endpoint argument prove physicality. This is a finite first-order implication, so it transfers to every real closed field. Thus

$$\text{nonnegative coefficients + finite kernel + both angle jets} \quad \Longrightarrow \quad \text{physicality.} \tag{SE7}$$

No coercivity premise is needed for this implication. Combining it with the theorem then excludes every proposed degenerate-support contact automatically.

14.1.3 The negative-core state orientation and deletion test

For reference, the raw signed-core matrix P_H has

$$\det P_H = -16 D_u^2 D_v^2 u^2 v^2 \neq 0. \tag{SE8}$$

At any physical finite zero, at least one retained event has positive value. Otherwise all event matrices annihilate the state and the kernel equation would give $P_H \psi = 0$, contradicting (SE8). Consequently the core expectation there is strictly negative.

This fixes an exact state sign without native contact coordinates. For $u, v > 0$ put $h_0 = 1 + v^2 + u^2 v^2$. The X, Y core block is $2\left(\begin{smallmatrix} h_0 & -uv \\ -uv & h_0 \end{smallmatrix}\right) \succ 0$. Its Schur complement on (W, Z) is

$$\frac{2 D_u D_v}{h_0^2 - u^2 v^2} \begin{pmatrix} u^2 v^2 & -uv h_0 \\ -uv h_0 & u^2 v^2 \end{pmatrix}. \tag{SE9}$$

If $WZ \leq 0$, this quadratic form and the completed X, Y square are nonnegative. Hence a negative core expectation forces $WZ > 0$. At a physical contact the gauge $Z = 1$ therefore has $W > 0$. For an independent testing state with $WZ \leq 0$, the core value is instead a nonnegative pure-testing scalar; its separated product with d is a square, and every event is a square. This gives an exact easy-state tensor-SOS exit, with no claim about merely small positive WZ.

Suppose a positive retained event has zero row at a finite contact. Deleting its coefficient preserves the kernel and both angle jets: the square and all its first derivatives vanish there. The remaining coefficients are nonnegative, so (SE7) makes the deleted source physical. The theorem therefore forbids a deletion that makes the Newton support noncoercive. This supplies the exact no-wall implication used in the recovered global C1 sign-entry proof. It does not authorize deleting an event whose row is nonzero: (CAP15) then gives a negative value.

(SE10)

14.1.4 The canonical event basis and two sufficient contact families

Here we specify the global source target, including its quantifiers. Let $Q \subset \mathbb{R}^8$ be the compact finite-dimensional tensor-product quantum set from (RFA1)–(RFA6), and let f be an affine functional nonnegative on it. If f is nonnegative on the whole nonsignaling polytope, linear separation of its finite probability inequality description writes it as a nonnegative sum of probability events (a nonnegative constant can be absorbed using normalization). These events have immediate separated certificates. Otherwise f is negative at a PR vertex: the nonsignaling vertices are the sixteen deterministic assignments and eight PR vertices, and f is nonnegative at every deterministic assignment.

Choose that PR orientation. In the convention $H = 1 - (E_{00} - E_{01} + E_{10} + E_{11})/2$, the eight losing events vanish at that PR point. Each has a unique deterministic $S = 2$ neighbor where its probability is one and the seven other losing probabilities vanish. The nine evaluations are therefore an affine basis and give the exact coefficient representation

$$f = dH + 4\sum_i a_i p_i, \qquad d = -f(\mathrm{PR}) > 0, \qquad a_i = f(v_i)/4 \geq 0. \tag{SE11}$$

The checker verifies these eight neighbor evaluations and nonsingularity of their augmented nine-point evaluation matrix from the actual event definitions. The finite nonsignaling vertex argument and the full physical normalization are retained inputs, with exact source hashes in the accompanying dependency receipt.

Two omissions are compatible precisely when the events never specify opposite outcomes at a shared local setting. The factored coefficients in (SE1) give

$$0 \in \operatorname{int}\operatorname{conv}(\operatorname{supp} L) \quad \Longleftrightarrow \quad \text{all omitted losing events are pairwise compatible.} \tag{SE12}$$

For clarity, a separating direction for a subset of the eight square boundary points can be chosen from $(\pm 1, 0), (0, \pm 1), (\pm 1, \pm 1)$. For an axis representative, vanishing on the forbidden half-plane requires $\lambda\alpha = \lambda D = \alpha m = mD = 0$, whose minimal omission sets are $\{\lambda, m\}$ and $\{\alpha, D\}$. For a diagonal representative the minimal sets are $\{\lambda, \alpha\}$, $\{\lambda, m, A\}$, $\{n, D, \alpha\}$; the latter two already contain an axis exclusive pair. The physical cycle rotation gives the other directions. These are exactly the twelve exclusive pairs: cycle distance three or four. This proves (SE12), and the checker independently verifies it on all finite support choices. Compatible pairs have cycle distance one or two; a compatible triple is three consecutive vertices. A larger compatible omission set is impossible.

Normalize the nonnegative canonical source tuple by $d + \sum a_i = 1$. Its physical subset is compact and convex over $\mathbb{R}$. Consider an extreme point with $d > 0$ and coercive support. Its Laurent polynomial attains a minimum μ at finite positive U, V. If $\mu > 0$, perturb the active coefficients by small opposite nonradial amounts, preserving their sum and exact zero coordinates. Every positive q_e changes by relative error at most ϵ and h by absolute error at most ϵ. Uniformly,

$$L' \geq (1 - \epsilon)L + \epsilon h - \epsilon \geq (1 - \epsilon)\mu + \epsilon h - \epsilon > 0 \tag{SE13}$$

for sufficiently small real ϵ. The perturbed sources remain physical, contradicting extremality. There are at least six active coordinates because (SE12) permits at most three event omissions, so a nonradial perturbation exists. Hence $\mu = 0$, giving an actual finite noncommuting contact.

At that contact the four kernel equations and two angle equations

$$P\psi = 0, \qquad \psi^T P_u \psi = 0, \qquad \psi^T P_v \psi = 0 \tag{SE14}$$

are six homogeneous linear equations in the nine source coefficients. If seven or eight events were positive, there would be at least eight active source coordinates including d. Their nullspace has dimension at least two and contains the source itself. It therefore has a nonzero direction of coefficient sum zero. Small opposite changes in that direction preserve all active signs and the same equations (SE14). By (SE7) they remain physical, again contradicting extremality. No generic-rank assumption is involved. Thus a coercive extreme source has exactly five or six positive events.

Choose an adjacent compatible omitted pair first when three events are omitted. By the physical cycle symmetries, every such source enters one of the following exact domains, including the stated zero faces:

$$\begin{aligned} &\text{C1}: \quad d, m, C, A, \alpha > 0, \quad \lambda = n = 0, \quad D, \beta \geq 0, \quad D + \beta > 0, \\ &\text{C2}: \quad d, m, C, A, D, \alpha > 0, \quad \lambda = \beta = 0, \quad n \geq 0. \end{aligned} \tag{SE15}$$

Both domains retain physicality and the existence of an actual finite positive source contact. In C1 a third omitted event can only be D or β. In C2 it can only be n; this boundary is already in a C1 orbit, but allowing it in both is harmless. Noncoercive source faces contain an exclusive omitted pair and are relabelings of $\lambda = m = 0$ or $\lambda = \alpha = 0$; their recovered exact matrix lifts remain separate, available proof inputs.

The normalized source base has affine dimension at most eight, so the finite-dimensional extreme-point theorem and Caratheodory's elimination argument express each point as a convex combination of at most nine extreme points. The midpoint definition of extremality, physicality, this fixed nine-point combination, and finite-contact attainment all have first-order polynomial formulas. Their truth over $\mathbb{R}$ therefore transfers to every real closed coefficient field. Compactness, logarithms, and real perturbation sizes were used only to prove the real assertion before transfer. Nonnegative summands automatically preserve literal zeros of the original source.

Corollary 14.2 (Precise sufficient tensor hypotheses)**.** *Suppose that, for each of C1 and C2 in (SE15), every actual source tuple over every real closed $K_c \supset \mathbb{R}$ has an unmultiplied full-parent SOS at every independent projective test and state over every real closed $K_t \supset \mathbb{R}$. There is no algebraic-disjointness assumption: the fields are labelled factors and may be copies of the same field. Together with the two recovered closed-face matrix lifts, these two hypotheses imply the universal separated slack-SOS statement for the entire physical set Q.*

Proof. Apply (SE11)–(SE15) in the source field, restore the finite real Bell relabelings, and multiply roots by the separate square roots of the nonnegative source combination weights. The event atoms and the closed-face atoms have separated certificates, the latter by attained finite-dimensional matrix duality as in (RFA). Every quantum testing point is a convex combination of at most nine real-qubit atoms in its own field; multiply roots by the separate testing-weight square roots and sum. The pure homogeneous angle, state and source normalization factors are restored within their respective fields. These operations are exactly finite conic and convex closure of sums of squares. □

What is and is not discharged. This section proves source-contact coercivity, the strict single-field contact Hessian, contact sufficiency, the no-wall deletion implication, and the exact canonical source partition. It isolates sufficient C1/C2 tensor hypotheses with every coefficient sign, literal boundary and source contact retained. Their positive-total and weak-total subdomains are the exhaustive scale split (CAP17). It does not discharge those two all-tests hypotheses. In particular, (SE6) is not a mixed tensor inverse, and its positive Hessian need not have a positive definite real residue after arbitrary independent source/testing degenerations. No generic matrix descent or historical all-tests assertion is used.

Chapter 15

Global native entry from actual physical contacts

Component `paf13bbf8fd32`
Source: `NEW/global/global_native_entry_REDERIVED.tex`
SHA256: `af13bbf8fd322035a18788e86e87493723ba6179e6ab960062901384c17f0e19`
Inventory use: EN1-EN16 global strict signs and unique native formula identification on both exact domains SE15 | Only EN13 exact positive native source transport and bounded a,t | EN1–EN10 and EN13

15.1 Global native entry from actual physical contacts

This section replaces a source-chart assumption by an actual implication. It uses the full-parent dictionary and SE1–SE10, whose proofs establish contact coercivity, strict source-field curvature, contact sufficiency, and the event-deletion test. Those are single-field assertions; no strict residue or mixed tensor inverse is inferred here.

15.1.1 Statement and the real maximal-contact graph

Use the raw full parent F, with $d > 0$. Every source below is physical and has an actual finite noncommuting zero. The two allowed coefficient domains are

$$\begin{aligned} &\text{C1}: \quad \lambda = n = 0, \quad m, C, A, \alpha > 0, \quad D, \beta \geq 0, \quad D + \beta > 0, \\ &\text{C2}: \quad \lambda = \beta = 0, \quad m, C, A, D, \alpha > 0, \quad n \geq 0. \end{aligned} \tag{EN1}$$

Make the actual angle gauges positive and put $Z_c = 1$, as permitted by SE2. Then $W_c > 0$ by SE9. We prove the strict signs

$$\begin{aligned} X_c < 0, \quad Y_c > 0, \quad u_c W_c - Y_c > 0, \quad 1 - v_c Y_c > 0, \\ R_{m,c} := v_c W_c - X_c + u_c v_c Y_c - u_c > 0, \\ R_{D,c} := u_c W_c + u_c v_c X_c - Y_c - v_c < 0, \\ X_c + u_c > 0, \qquad W_c + v_c X_c > 0. \end{aligned} \tag{EN2}$$

These inequalities hold over every real closed coefficient field. In particular they are strict ordered-field statements, with no assertion that a gap has positive real residue.

Here is the common connected-graph argument, proved over $\mathbb{R}$ before first-order transfer. In either exact two-omission family, initially require all six retained events positive, normalize $C = 2$, and let their other five weights vary over the positive orthant. Put

$$d_*(w) = \max\{d \geq 0 : (d, w) \text{ is physical}\}. \tag{EN3}$$

The admissible set is a closed interval by convexity. It has positive length: at $d = 0$ the central Laurent coefficient is positive, its nonnegative Cauchy–Binet expansion containing $16mCAD$; the

outer coefficients remain nonnegative for $d \geq 0$. Thus sufficiently small positive d is physical by SE1–SE2 and the homogeneous endpoint limits. A fixed strictly CHSH-violating real test bounds d above.

For these positive six-event supports the active outer exponents are constant and have zero in the interior of their convex hull. The logarithmic Laurent sum is strictly convex and coercive. At d_* its unique minimum is zero. Indeed if its minimum were $\mu > 0$, a small increase of d preserves every active outer coefficient within a relative error ε and the central coefficient within absolute error ε. As in SE13, this preserves a positive uniform lower bound, contradicting maximality. There is therefore one positive squared-angle contact, and SE2 gives one state with $Z = 1$.

At that contact the core value is strictly negative (SE8–SE9). The Jacobian of the six equations $\nabla_z F = 0, F = 0$, in $z = (W, X, Y, u, v)$ and d, is

$$\begin{pmatrix} \nabla_z^2 F & \partial_d \nabla_z F \\ 0 & \partial_d F \end{pmatrix}, \qquad \nabla_z^2 F \succ 0, \quad \partial_d F < 0. \tag{EN4}$$

The real analytic implicit-function theorem gives a local analytic contact branch. SE7 makes it physical and the negative core value makes its d maximal. Uniqueness makes these branches agree. Thus the normalized contact is analytic on the entire connected positive orthant. Every physical finite zero with those event weights lies on this graph, because its negative core value prevents increasing d at that same test.

15.1.2 C1: strict signs including the two literal boundary faces

Write

$$(W, X, Y, Z) = (S, u(\theta - 1), uY_0, 1). \tag{EN5}$$

The following two identities are the actual retained sign-entry identities, also checked against the full raw parent in the focused checker. With $B = \operatorname{adj} P$,

$$\begin{aligned} uB_{44} + B_{24} &= 16uv^2 D_u^2 D_v \, \mathcal{F}, \\ \mathcal{F} &= (d + 2D)\{(d + 2\alpha)(mD_v + Cu^2v^2) + 2AD_v(m + Cu^2)\} \\ &\quad + 2mC\{d(u^2v^2 + D_v) + 2\alpha + 2Dv^2D_u\} > 0. \end{aligned} \tag{EN6}$$

Since P has one-dimensional kernel and three positive eigenvalues, $B = \gamma\psi\psi^T$ with $\gamma > 0$. Equation EN6 gives $\theta > 0$.

Neither $Y_0 = 0$ nor $S - Y_0 = 0$ is possible. These walls would delete the positive A-event or C-event, respectively, while preserving the kernel and both angle jets. The resulting omission set contains an exclusive pair, so has noncoercive support. SE7 and SE10 rule this out. The remaining cross-multiplied contact identity is

$$m\,[S + uv(\theta - 1)]\,[v(S + u^2Y_0) - u\theta] = 2u^3(S - Y_0)(1 - \theta)D_v. \tag{EN7}$$

For precision about exceptional source minors, use the four raw kernel equations and the two full angular derivatives. Their selected six-column system, for $(d, m, A, D, \alpha, \beta)$ with $C = 2$, has determinant

$$-8192\theta u^2 v D_u^2 D_v^2 \, \mathcal{D}, \qquad \mathcal{D} = Y_0 D_u (uY_0 + v)(1 - uvY_0)[S + uv(\theta - 1)]R_m(-R_D). \tag{EN8}$$

EN7 follows first from its rational solution wherever this selected minor is nonzero: the Cramer determinant for the m-column, multiplied by $[S + uv(\theta - 1)]R_m$, equals this determinant times $2u^3(S - Y_0)(1 - \theta)D_v$. This is an exact polynomial determinant identity. The equivalent native formula is $m = 2ae^2c_2/(Mc_3)$, with the definitions in EN13 below: $c_2/c_3 = (1 - \theta)D_v/E$, $M = R_m/(Sv)$, $E = [S + uv(\theta - 1)]/S$, $a = u(S - Y_0)/v$, $e = u/S$. These substitutions give EN7 exactly. The native contact equations are polynomial identities, so this argument does not assume the signs being proved.

At the exact positive seed

$$(u,v;S,Y_0,\theta)=(1,1;1,\tfrac14,\tfrac12),\qquad (d,m,C,A,D,\alpha,\beta)=(\tfrac{27}{2},4,2,9,\tfrac54,\tfrac{13}{4},\tfrac94), \tag{EN9}$$

all six stationary equations hold and the selected minor is nonzero. Consequently that analytic minor is not identically zero on the connected contact graph. Its nonzero set is dense. Continuity extends EN7 to every contact on the graph. At $\theta=1$, EN7 and $S,m>0$ would imply $R_m=0$, permitting deletion of m and creating an exclusive omitted pair, again impossible. The three continuous nonzero functions $Y_0, S-Y_0, \theta-1$ therefore have constant signs, fixed by EN9:

$$S>0,\qquad 0<Y_0<S,\qquad 0<\theta<1. \tag{EN10}$$

The full-parent C1 reflection swaps $(\lambda,n),(m,C),(A,\alpha),(D,\beta)$, sends u to $1/u$, and sends the state to

$$(vS-u(\theta-1),\ -S-uv(\theta-1),\ 1-uvY_0,\ uY_0+v).$$

Its state matrix G has $G^TG=D_vI$; the full core and every event obey $u^2F_{\sigma_1 c}(1/u,v;G\psi)=D_vF_c(u,v;\psi)$. Its last coordinate is positive. Applying EN10 to the transformed normalized state proves successively $1-uvY_0>0,\ S+uv(\theta-1)>0,\ R_D<0,\ R_m>0$. Together with EN10 this is precisely EN2.

For a literal third omission $D=0$ or $\beta=0$, start at the given real contact and open that coefficient to $\varepsilon>0$. EN4 is still invertible by SE. The local analytic branch stays at positive d, has nonnegative event weights, and is physical by SE7. The already proved inequalities hold on it. At the limit, EN6 and SE9 stay strict; the still positive A,C,m and the same deletion arguments make the other weak inequalities strict. EN7 passes to the limit, and so does the reflection. Thus EN2 holds on both boundary faces. This is an implicit-function limit at a fixed finite real contact, not an assertion that an optimizer cannot escape to infinity. First-order transfer now gives the entire implication over every real closed field.

15.1.3 C2: all signs and a positive omitted-row cofactor

Use the same positive six-event graph. Five retained-amplitude walls are excluded directly by SE10:

$$Y=0,\quad uW-Y=0,\quad R_m=0,\quad R_D=0,\quad vY-1=0. \tag{EN11}$$

Deleting A,C,m,D,α, respectively, introduces an exclusive omitted pair in addition to $\lambda=\beta=0$. The wall $X=0$ deletes n; the resulting source is on the C1 three-consecutive-omission face $\lambda=n=\beta=0$. It is physical by SE7, so EN10 gives $X<0$, a contradiction. Thus all six retained event rows have fixed signs on the connected C2 graph. The exact seed

$$(n,m,C,A,D,\alpha)=(\tfrac18,\tfrac38,\tfrac16,\tfrac58,\tfrac38,\tfrac16),\quad d=1,\qquad (u,v;W,X,Y,Z)=(1,1;1,-\tfrac14,\tfrac14,1)$$

has every required positive weight and every stationary equation. It fixes their signs to the first six signs in EN2.

Here is a direct proof of the omitted-row sign, including $n=0$. Let P_0 be this raw C2 matrix with n set to zero. Changing n adds only $4nD_uD_v$ to its XX entry. The cofactor B_{24} does not contain that column, whereas B_{44} changes by $4nD_uD_v\det(P_0|_{W,Y})$. Consequently

$$\begin{aligned} uB_{44}+B_{24} &= 16uv^2D_u^2D_v\,\mathcal{F} \\ &\quad + 4unD_uD_v\det(P_0|_{W,Y})>0. \end{aligned} \tag{EN12}$$

Here $\mathcal{F}$ is EN6 with $\beta=0$. The determinant in the second line is positive: the $Z=0$ state block already has the strictly positive signed-core form SE2. Thus $X+u>0$. The C2 full-parent reflection

$$(u,v;W,X,Y,Z)\mapsto(1/v,1/u;Z,X,Y,W),\quad (m,D),(C,\alpha),(\lambda,\beta)$$

has the exact homogeneous identity $u^2v^2F_{\sigma_2 c}(1/v,1/u;Z,X,Y,W)=F_c(u,v;W,X,Y,Z)$. The transformed contact has positive last coordinate W. Applying the just-proved $X'+u'Z'>0$ gives $W+vX>0$. This proves EN2 for positive n. For literal $n=0$, EN1 is already a C1 triple face and its signs apply directly; EN12 remains strict as well.

15.1.4 Exact native-coordinate transport and uniqueness

For C1, reflect if needed so $0 < k = u_c \le 1$, and define

$$\begin{gathered} S = W_c, \quad r = Y_c/(kS), \quad \theta = 1 + X_c/k, \quad h = 1 - \theta, \\ \eta = v_c/(kS), \quad \zeta = Sv_c/k, \quad f = \sqrt{\eta}, \quad g = \sqrt{\zeta}, \quad e = k/S = kf/g, \\ a = (1-r)/\eta, \quad t = \theta/\zeta, \quad x = k^2, \\ R = 1 + xr, \quad T = 1 + xh, \quad M = R - t, \quad J = T - a, \\ E = 1 - x\eta h, \qquad Q = 1 - x\zeta r. \end{gathered} \tag{EN13}$$

By EN2, $0 < r, \theta < 1$, all η, ζ, a, t are positive, and

$$R_m = k\zeta M > 0, \quad R_D = -v_c J < 0, \quad W_c + v_c X_c = SE > 0, \quad 1 - v_c Y_c = Q > 0.$$

Thus $M, J, E, Q > 0$, and $0 < a, t < 2$. The selected determinant EN8 is now strictly nonzero. The native seven-coquotient formulas of `native_formulas.tex`, together with the source coefficients in SD1 of `source_definitions.tex`, satisfy exactly the six stationary equations at this same contact. Uniqueness therefore identifies them with the original normalized source, including $D = 0$ or $\beta = 0$. No asymptotic hypothesis is used for these polynomial identities.

For C2, normalize $d = 1$, and set

$$\begin{gathered} c = u_c, \quad a = v_c/u_c, \quad b = aW_c, \quad r = -X_c/u_c, \quad t = (u_c W_c - Y_c)/v_c, \quad z = c^2, \\ h = b - a^2 t = aY_c/u_c, \quad g = (1+z)b - (1 - r + a^2 zt) = R_m/u_c, \quad j = 1 + zr - t = -R_D/v_c. \end{gathered} \tag{EN14}$$

The complete strict native domain follows:

$$a, b, c, t > 0, \quad 0 < r < 1, \quad h, g, j > 0, \quad 1 - zh = 1 - v_c Y_c > 0, \quad b - a^2 zr = a(W_c + v_c X_c) > 0. \tag{EN15}$$

The actual six-column stationary determinant is, in the unweighted event convention of the retained determinant proof,

$$-64ac^6 rgt\, hj(1-zh)(1+z)^2(1+a^2 z)^2 \Delta,$$

$$\Delta = 1 + (1+z)b + z(1+z)b^2 + 2z(1+z)br + z^2 r^2 + 2zr + a^2 z\{(1+z)b + 1 + zr(2-r)\} \ge 1. \tag{EN16}$$

That body provides its explicit reduction to a three-by-three determinant. The six rational coefficients in `source_formulas.tex` solve these six equations at the contact EN14. EN16 makes the solution unique, so they are exactly the original source coefficients. This remains valid at $n = 0$; one does not delete its column from the square system.

All gauge actions, positivity statements and source equations used above are finite polynomial or rational ordered-field statements with denominators explicitly nonzero. The real connected-graph proof therefore transfers as one fixed first-order implication. Together, EN1–EN16 discharge the native sign and positive-gap interface for both families in SE15. They do not prove an all-tests tensor assertion; that assertion is supplied by the subsequent exhaustive source partition and its actual terminal bodies.

Chapter 16

R1 unbounded-q: exact contact circuits and their square budgets

Component `p6db37c4233da`
Source: `REC/c144/r1_qo_contact_circuit_budgets.tex`
SHA256: `6db37c4233da1f389bc7a081fa5bad5e3daa4f5d61640483efcd549a455ac952`
Inventory use: RB1-RB17

16.1 R1 unbounded-q: exact contact circuits and their square budgets

Status and exact scope. This is a collection of proved algebraic lemmas for the remaining original R1 source sector. It is not a local or all-testing admission. The complete original parent and the two complete state normals are RQ6–RQ11 of `r1_unbounded_q_structural_checkpoint.tex`. The new lemmas below avoid substituting $u = vz/T$ and therefore introduce no inverse powers of a mixed graph coordinate. The final full-normal allocation is still a separate obligation.

16.1.1 The full n scalar has just two negative monomials

Use RQ1–RQ11. Restore the original n packet in the scalar graph $N = M = 0$ and write this full scalar as F_{full}. Equivalently

$$F_{\mathrm{full}} = F + n(1+v^2)\xi_g^2, \qquad \xi_g = p(1+ru^2)v - (1-r)uz. \tag{RB1}$$

Here the factor $1+v^2$ is essential. Its omission would change the whole scalar and every subsequent allocation. Put $K_0 = CH^2 + (D+\frac{1}{2})(1-H)^2$. Direct expansion gives

$$\begin{aligned} F_{\mathrm{full}} = {} & Au^2 + \{\mu(1-p) + (n+\tfrac{1}{2})p^2\}v^2 + \alpha z^2 \\ & + \{A + \alpha(1-p)^2 + \mu(1-p) + \tfrac{1}{2}(1-p)^2\}u^2v^2 + nu^2z^2 \\ & + \{K_0 + AH^2 + 2H\alpha + H - \tfrac{1}{2}\}v^2z^2 + (n+\tfrac{1}{2})p^2v^4 \\ & + (K_0+n)u^2v^2z^2 + H^2(A + C + \alpha + \tfrac{1}{2})v^4z^2 + CH^2u^2v^4z^2 \\ & - 2\{AH + \alpha(1-p) + np + \tfrac{1}{2}\}uvz \\ & - 2\{AH + H\alpha(1-p) + np + \tfrac{1}{2}[H(1-p)+p]\}uv^3z. \end{aligned} \tag{RB2}$$

Every displayed even coefficient is nonnegative on the original source. The only coefficient requiring an estimate is that of v^2z^2: K_0 has a positive lower bound of order $1/t$, whereas the possible negative part is at most $1/2$. In particular this observation allows arbitrary $n \geq 0$, including $n = 0$ and unbounded n. It is not a claim that a polynomial with two negative monomials automatically has a circuit certificate.

16.1.2 An exact finite contact expansion

Let $x = u_c^2$, $y = v_c^2$, and $q_z = z_c^2$ be the actual three positive contact coordinates of RQ5. Define the separated rows

$$U = u/\sqrt{x}, \qquad V = v/\sqrt{y}, \qquad Z_* = z/\sqrt{q_z}. \tag{RB3}$$

These are rows, not bounded Gram coefficients. Define

$$Q_* = U - VZ_*, \qquad B_* = V - UZ_*, \qquad R_* = Z_* - UV,$$
$$\Psi = U^2 + V^2 + Z_*^2 + U^2V^2Z_*^2 - 4UVZ_*. \tag{RB4}$$

On the positive testing core Ψ is a separated SOS:

$$\Psi = (U - V)^2 + Z_*^2(1 - UV)^2 + 2UV(1 - Z_*)^2. \tag{RB5}$$

The last square has root $\sqrt{2UV}(1 - Z_*)$; this is a product of pure-source and pure-testing roots. Signs and literal testing axes must be supplied by the inherited core exits in any admission; the following identities themselves are polynomial identities.

Let $M_a, M_b, M_c, M_{ab}, M_{ac}, M_{bc}, M_{b^2}, M_{abc}, M_{b^2c}, M_{ab^2c}$ denote the ten positive monomial coefficients after RB2 is expressed in (U, V, Z_*). Thus, for example,

$$M_a = Ax, \quad M_b = \{\mu(1-p) + (n + \tfrac{1}{2})p^2\}y, \quad M_c = \alpha q_z, \quad M_{ac} = nxq_z,$$
$$M_{b^2c} = H^2(A + C + \alpha + \tfrac{1}{2})y^2q_z, \qquad M_{ab^2c} = CH^2xy^2q_z. \tag{RB6}$$

All coefficients are actual source quantities. The actual scalar contact value and its first jets imply

$$M_b + M_{b^2} = M_{ac} + M_{abc} + M_{ab^2c}, \qquad M_a + M_{ab} = M_c + M_{bc} + M_{b^2c}. \tag{RB7}$$

Set

$$k_c = \tfrac{1}{2}(M_a - M_b + M_{ac} + M_{ab^2c} - M_{bc}), \qquad r_c = M_c - M_b + M_{ac} - k_c,$$
$$f_c = M_b - M_{ac}, \quad q_c = M_a - f_c, \quad e_c = M_{b^2} - M_{ab^2c}, \quad d_c = M_{b^2c} - k_c, \quad g_c = M_{ab^2c}. \tag{RB8}$$

Then the exact full scalar is

$$F_{\text{full}} = q_cQ_*^2 + M_{ac}B_*^2 + r_cR_*^2 + d_cV^2Q_*^2 + e_cV^2B_*^2 + f_c\Psi + g_cV^2\Psi + k_cZ_*^2(V^2 - 1)^2. \tag{RB9}$$

The two odd coefficients in RB9 are fixed by the contact value and the V derivative; RB7 fixes the remaining even coefficients. No positivity of e_c, k_c, f_c is inferred merely from this identity. There is also a one-parameter variant, obtained by adding zero: for any pure-source τ, replace

$$(q_c, M_{ac}, d_c, e_c, f_c, g_c, k_c) \longmapsto (q_c - \tau, M_{ac} - \tau, d_c + \tau, e_c + \tau, f_c + \tau, g_c - \tau, k_c - \tau)$$

and add $\tau U^2Z_*^2(V^2 - 1)^2$. (RB10)

16.1.3 Two exact circuit budgets

The first budget is the polynomial identity

$$(1 + U^2)\Psi = R_*^2 + B_*^2 + U^2Q_*^2 + U^2(1 - UVZ_*)^2. \tag{RB11}$$

All permutations of (U, V, Z_*) are allowed. In particular

$$(1 + V^2)\Psi - V^2B_*^2 = Q_*^2 + R_*^2 + V^2(1 - UVZ_*)^2 \tag{RB12}$$

is literally a sum of three squares. Consequently a signed $e_c V^2 B_*^2$ term is paid at infinitesimal Gram cost whenever e_c^- / f_c is source infinitesimal and $e_c^- / (f_c y)$ is bounded: the cost is $(e_c^- / f_c) + (e_c^- / (f_c y))v^2$. This criterion never treats V as a bounded coefficient.

For the second budget put $t_* = UV$, $\delta_* = U - V$, and

$$E = Z_*(1 - t_*), \qquad T_* = t_*(1 - Z_*), \qquad J = T_* - \delta_*^2 Z_*.$$

There are exact identities

$$R_* = E - T_*, \qquad V^2 Q_*^2 + U^2 B_*^2 - T_*^2 = J^2 + 2t_* \delta_*^2 Z_*^2. \tag{RB13}$$

Together with RB5 they give the explicit separated SOS remainder

$$\begin{aligned} &2\{\Psi + V^2 Q_*^2 + U^2 B_*^2\} - R_*^2 \\ &\quad = (E + T_*)^2 + 2\delta_*^2 + 4t_*(1 - Z_*)^2 + 2J^2 + 4t_* \delta_*^2 Z_*^2. \end{aligned} \tag{RB14}$$

Thus R_*^2 can be bought from the circuit and the two current rows, without any inverse or root of a mixed sum. For positive source masses q_c, v_c, f_c, a multiplier $a_c \geq 0$ has infinitesimal cost if

$$a_c / f_c \in \mathrm{I}, \qquad a_c / (q_c y) \text{ is bounded}, \qquad a_c / (v_c x) \text{ is bounded}. \tag{RB15}$$

The last two costs carry the actual testing factors v^2, u^2. The identities continue to hold on literal axes. No division by U, V, Z_* occurs anywhere in RB11–RB15.

16.1.4 The exact remaining n cancellation

Write $j = \sqrt{x q_z / y}$, so the original actual contact gives

$$\frac{p}{j} = \frac{1 - r}{1 + rx}, \qquad \xi_g = p\sqrt{y}\,\{B_* - r\sqrt{x}\,u R_*\}. \tag{RB16}$$

Restoring or removing the full n packet must retain the factor $1 + v^2$ from RB1. For its unweighted part alone,

$$\begin{aligned} nxq_z - np^2 y &= nxq_z\{1 - (p/j)^2\}, \\ n\{1 - (p/j)^2\} &= nr\,\frac{(2 - r) + 2x + rx^2}{(1 + rx)^2}. \end{aligned} \tag{RB17}$$

This is bounded through the opening $r = 0$ in residue and includes literal $n = 0$. The remaining cross and v^2 terms are exact polynomials obtained from RB1 and RB16. They have not been silently discarded. A proof of their allocation, together with the complete state normals RQ10–RQ11 and the outer-testing joins, is necessary before an unbounded-q R1 admission can be recorded.

Chapter 17

Reconstructed R1 unbounded-q source and scalar certificate

Component `p9e725a065583`
Source: `REC/c144/rederived/r1_qo_actual_source_and_scalar.tex`
SHA256: `9e725a065583e84a9b9d9e29ea2cca9362ea0c6eb616fc223c08505839d10ace`
Inventory use: RSR1-RSR29

17.1 Reconstructed R1 unbounded-q source and scalar certificate

Provenance and scope. This is a fresh proof reconstructed from the original stationary source formulas and the retained contact substitution. It is not represented as the original bytes of the former R21 file. Its finite identities are checked by `check_r1_qo_compact_source.py`; the uniform estimates are proved below. The conclusion concerns the complete scalar on the two-normal graph. Both state normals and the outer testing charts remain separate obligations. No historical review receipt is used as a premise. Source quantities lie in a real closed extension K_c and tests and states in an independent real closed extension K_t; every SOS assertion is in $K_c \otimes_{\mathbb{R}} K_t$, without imposing an order on that tensor ring or identifying its two fields.

The native source formulas are the unchanged formulas in `inherited_r19/inherited_r18/inherited_r17/definitions/ weak_c2_source_formulas.tex`. We use their actual common-II source, with $d = 1$, $\lambda = \beta = 0$, all original strict gaps and $n \geq 0$. In the original R1 sector $B_s, t_s, a_s B_s, t_s B_s$ are positive infinitesimals. Write $r, h \in (0, 1)$ for its two native source parameters, $c_s = 1 + a_s B_s r$, $R_s = 1 + h a_s B_s$, and $K_s = 1 + a_s B_s$. The sector treated here is

$$q_{\mathrm{old}} = \frac{R_s^2}{a_s t_s K_s^2} \quad \text{unbounded.} \tag{RSR1}$$

The original positive gap $c_s - (1 - h)/t_s$ implies $1 - h$ infinitesimal; therefore $h \sim 1$. Since $R_s, c_s, K_s \sim 1$, RSR1 is equivalent to $a_s t_s$ infinitesimal. No condition on the size of a_s itself is introduced. The independent testing variables below are $u, v > 0$ infinitesimal and $z > 0$ arbitrary.

17.1.1 The exact compact source and its uniform bounds

Define pure-source quantities

$$\begin{gathered} j = \frac{a_s c_s}{R_s}, \quad B = \frac{B_s R_s}{c_s}, \quad T = \frac{t_s c_s}{R_s}, \\ p = \frac{a_s(1 - r)}{R_s}, \quad H = \frac{1 - h}{t_s c_s}, \\ x = jB, \quad y = TB, \quad q_z = jT. \end{gathered} \tag{RSR2}$$

The native gaps give $0 < p < \min(1, j)$ and $0 < H < 1$. Here B, T, x, y, q_z are positive infinitesimals, while j is arbitrary positive. Direct inversion gives

$$r = \frac{j-p}{j+px}, \quad h = \frac{1-HT}{1+HTx}, \quad a_s = \frac{j+px}{1+HTx}, \quad B_s = \frac{B(1+HTx)}{1+pB}, \quad t_s = \frac{T(1+pB)}{1+HTx}. \tag{RSR3}$$

In particular the compact variables describe the actual source, including its $r \to 0$, $p \to 1$, $H \to 0$, and $H \to 1$ openings. Set

$$K = j + 1 - p, \qquad E = 1 + B(HT + p - 1), \qquad W_0 = (1-H)j + 1 - p. \tag{RSR4}$$

Then $\frac{1}{2}(j+1) < K < j+1$, $E \sim 1$, and $W_0 > 0$. The following polynomials contain the entire stationary dictionary:

$$\begin{aligned}
\mathcal{P} &= j + 1 + B\{(j+1)^2 - p\} \\
&\quad + TB\{Hj + j + 1 + B[Hj(2j+2-p) + p(2j+1-p)]\} \\
&\quad + HjT^2B^2(1 + Bp + BHj), \\
\mathcal{L} &= K + T(W_0 - H) + BT[pK + Hj(1-p)] + BHT^2(W_0 - 1), \\
\mathcal{N} &= p + B(-j^2 + 3jp - j - p^2 + p) \\
&\quad + BT(Hj^2 + Hj - j^2 + jp - j + p) \\
&\quad + B^2T(2Hj^2p - Hj^2 - Hjp^2 + 2Hjp - 2j^2p + 3jp^2 - jp - p^3 + p^2) \\
&\quad + B^2HT^2j(Hj - j + p)(1 + Bp).
\end{aligned} \tag{RSR5}$$

With $\mu = m(1-p)$, the exact six original stationary weights are

$$\begin{aligned}
\mu &= \frac{jp(1+y)(1+x)(1+Hy)}{2\mathcal{P}}, & n &= \frac{\mathcal{N}}{2(j-p)\mathcal{P}}, \\
C &= \frac{\mathcal{L}}{2HT\mathcal{P}}, & A &= \frac{T(1+x)(1+Hy)(K+Hjy)}{2(1-HT)\mathcal{P}}, \\
D &= \frac{(1+Hy)(K+Hjy)\{1 - y[(1-H)j - p]\}}{2T(1-H)\mathcal{P}}, & \alpha &= \frac{B^2(K+Hjy)\mathcal{L}}{2E\mathcal{P}}.
\end{aligned} \tag{RSR6}$$

For example, the original stationary denominator and its source mass satisfy

$$\Delta = \frac{(1+x)\mathcal{P}}{(1+Bp)(1+HTx)}, \qquad M = \frac{(1+x)\mathcal{L}}{(1+HTx)^2}.$$

The two native strict gaps become

$$g = \frac{(1-p)(1+x)}{1+HTx}, \qquad j_{\text{gap}} = \frac{(1-H)(1+x)}{1+Bp}.$$

These equations verify the source origin of RSR6 directly; no new stationary solution has been substituted for the original one.

Every term after $j+1$ in $\mathcal{P}$, divided by $j+1$, is infinitesimal: use $B(j+1) = B + x \in \mathrm{I}$, $T \in \mathrm{I}$, $p < 1$, and $H < 1$. The same termwise estimate, using $K > (j+1)/2$, applies to $\mathcal{L} - K$. Thus

$$\mathcal{P} \sim j + 1, \qquad \mathcal{L} \sim K. \tag{RSR7}$$

These are relative equivalences, valid also when j is infinitesimal or unbounded. They show, in particular, that A/T, THC, and $T(1-H)D$ are positive bounded units, μ/p is bounded, and $h_0 = \mu + 1/2$ is a positive bounded unit. Moreover α/B is infinitesimal. Only the displayed source openings can make m, n, C, D unbounded.

The original physicality condition gives a useful uniform restriction. Regroup the exact polynomial $\mathcal{N}$ as

$$\mathcal{N} = p\{1 + B(3j + 1 - p) + BT(j+1)\} - Bj(j+1)\{1 + (1-H)T\} + \mathcal{R}. \tag{RSR8}$$

Here $\mathcal{R}$ is precisely the last two lines of $\mathcal{N}$ in RSR5. Since $0 < p < \min(1, j)$, each coefficient in its B^2T line is bounded in absolute value by a fixed multiple of $j(j+1)$. The B^2HT^2 line has the same bound, multiplied by the bounded factor $T(1+Bp)$. Consequently

$$|\mathcal{R}| \le C_0 BT\, Bj(j+1)$$

for a fixed real C_0. All corrections multiplying p and $Bj(j+1)$ in RSR8 are infinitesimal. As $n \ge 0$ is exactly $\mathcal{N} \ge 0$, it follows that

$$\eta_* := \frac{Bj(j+1)}{p} \text{ is bounded}, \qquad \eta_* \in \mathrm{I} \implies n \sim \frac{p}{2(j-p)(j+1)} > 0. \tag{RSR9}$$

Indeed $\mathcal{N} = p(1+\varepsilon_1) - Bj(j+1)(1+\varepsilon_2)$ with $\varepsilon_i \in \mathrm{I}$. This proves both assertions without dividing by n and includes literal $n = 0$ in the bounded-η_* alternative. It also gives nr bounded.

17.1.2 The complete graph and its actual contact circuit

The quarter-parent, before any packet is removed, is

$$\begin{aligned}\mathscr{P} = {} & m\{vW + uvY - X - uZ\}^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 + nD_uD_vX^2 \\ & + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2, \qquad D_u = 1 + u^2,\ D_v = 1 + v^2.\end{aligned} \tag{RSR10}$$

The source-free rotated state coordinate is $w = (W + uY)/D_u$. On its nonzero chart set $w = 1$ and $z = (Z - uX)/D_u$. Define the two complete state normals

$$N = X + uZ - pD_uv, \qquad M = uW - Y - HD_uvz.$$

Their exact inverse is

$$\begin{aligned} X &= N/D_u + pv - uz, & Z &= z + uN/D_u + puv, \\ Y &= u - M/D_u - Hvz, & W &= 1 + uM/D_u + uHvz. \end{aligned} \tag{RSR11}$$

The scalar studied here is exactly $F_{\mathrm{full}} = (\mathscr{P}/D_u)|_{N=M=0}$. No n packet, m packet, or state-normal term is silently removed. Substitution of RSR11 into RSR10 gives the complete graph RB2 of `r1_qo_contact_circuit_budgets.tex`. The independent generic parent calculation in the checker verifies that equality before specializing any stationary weight.

Use the separated rows

$$U = u/\sqrt{x}, \quad V = v/\sqrt{y}, \quad Z_* = z/\sqrt{q_z}, \qquad Q_* = U - VZ_*, \quad B_* = V - UZ_*, \quad R_* = Z_* - UV,$$

$$\Psi = U^2 + V^2 + Z_*^2 + U^2V^2Z_*^2 - 4UVZ_*. \tag{RSR12}$$

These are rows; U, V, Z_* are not assumed bounded coefficients. The actual contact is $(U, V, Z_*; N, M) = (1, 1, 1; 0, 0)$. The exact native formulas make the complete parent value and all five first derivatives vanish there. On the positive core,

$$\Psi = (U - V)^2 + Z_*^2(1 - UV)^2 + 2UV(1 - Z_*)^2 \tag{RSR13}$$

is a legal separated sum of squares: the last root is $\sqrt{2}\,(uv)^{1/2}(xy)^{-1/4}(1 - Z_*)$, with its radicals in the two separate real closed fields.

Let the ten positive monomial coefficients and the seven circuit masses be exactly RB6–RB8. In particular

$$M_a = Ax, \quad M_b = \{\mu(1-p) + (n + \tfrac{1}{2})p^2\}y, \quad M_c = \alpha q_z, \quad M_{ac} = nj^2y,$$

$$f_c = M_b - M_{ac}, \quad q_c = M_a - f_c, \quad e_c = (n + \tfrac{1}{2})p^2y^2 - g_c, \quad d_c = M_{b^2c} - k_c.$$

The original contact balances give the exact scalar identity

$$\begin{aligned} F_{\mathrm{full}} &= q_c Q_*^2 + M_{ac} B_*^2 + r_c R_*^2 + d_c V^2 Q_*^2 + e_c V^2 B_*^2 \\ &\quad + f_c \Psi + g_c V^2 \Psi + k_c Z_*^2 (V^2-1)^2. \end{aligned} \tag{RSR14}$$

The identities RB7 and the two odd coefficient balances are independently checked on the actual source. Thus RSR14 is a finite polynomial identity, not a Taylor approximation to the contact.

Here is an explicit source-only formula for its radial mass. Put $\mathcal{F} = K + T\mathcal{F}_1 + T^2\mathcal{F}_2$, where

$$\begin{aligned} \mathcal{F}_1 = &-2BHjp + BHj + 2Bjp - 2Bp^2 + 2Bp \\ &- Hj - Hp - H + j + 1 - p^2/j + p/j, \\ \mathcal{F}_2 = &-B^2H^2jp + B^2Hjp - B^2Hp^2 + B^2Hp \\ &- BH^2j - BH^2p + BHj - BHp^2/j + BHp/j. \end{aligned} \tag{RSR15}$$

Every displayed term divided by K is bounded, using $p/j < 1$; terms containing Bj are even infinitesimal. Hence $\mathcal{F} \sim K$. Direct source substitution yields

$$\begin{aligned} f_c &= \frac{Tx^2\mathcal{F}}{2\mathcal{P}} > 0, & q_c &= Ax - f_c \sim \frac{TxK}{2(j+1)} > 0, \\ r_c &= \frac{B^2jT(1-p)(1+x)(1+Hy)\mathcal{L}}{2E\mathcal{P}} > 0, & g_c &= \frac{HT^2j^2B^3\mathcal{L}}{2\mathcal{P}} > 0. \end{aligned} \tag{RSR16}$$

In particular

$$f_c \sim \frac{Tx^2K}{2(j+1)}, \qquad \frac{r_c}{f_c} \sim \frac{1-p}{j}, \qquad \frac{g_c}{f_c} = Hy\frac{\mathcal{L}}{\mathcal{F}}. \tag{RSR17}$$

For the remaining signed mass define

$$\begin{aligned} \mathcal{K} = &\, pW_0 + B(1-p)\{j[H(p+1)-p] - (1-p)(p-H)\} \\ &+ TBH\{(p+1)(1-H)j + (1-p)(BHj + p + 1 - H)\} \\ &+ T^2B^2H^2j(1-H). \end{aligned} \tag{RSR18}$$

Then

$$k_c = -\frac{T^2B^2j(1+x)\mathcal{K}}{2E\mathcal{P}}.$$

The identity

$$\begin{aligned} \mathcal{K} - pW_0 + Bp(1-p)(j+1) = &\, B(1-p)\{Hj(p+1) + H(1-p) + p^2\} \\ &+ TBH\{(p+1)(1-H)j + (1-p)(BHj + p + 1 - H)\} \\ &+ T^2B^2H^2j(1-H) \end{aligned} \tag{RSR19}$$

has a nonnegative right side. Since $W_0 \geq 1 - p$ and $B(j+1) \in \mathrm{I}$, it proves $\mathcal{K} > 0$ and $k_c < 0$. The displayed finite arrays also give

$$\mathcal{K}/K = O(p+B), \qquad |k_c|/(q_c y) = O(p+B). \tag{RSR20}$$

All constants in these estimates are real and uniform over the source openings; neither $j - p$ nor $1 - H$ has been bounded below.

17.1.3 Paying the signed masses, with the full current reserved

Put

$$h_0 = \mu + \tfrac{1}{2}, \qquad M_e = \frac{nh_0j^2}{n+h_0}, \qquad J_c = M_e y, \qquad e_0 = \tfrac{1}{2}p^2y^2 - g_c. \tag{RSR21}$$

These are pure-source quantities and $0 \le J_c \le M_{ac}$. Writing $e_c = np^2y^2 + e_0$ preserves the entire second correlated current packet. The conclusion is the following precise scalar statement: after subtracting the reserved positive packets

$$(M_{ac} - J_c)B_*^2 + np^2y^2V^2B_*^2, \tag{RSR22}$$

the remainder is a separated SOS and retains, for example, one half of $q_cQ_*^2 + r_cR_*^2 + f_c\Psi + J_cB_*^2$. When $n = 0$ the two reserved packets and J_c vanish literally. This assertion supplies precisely the effective current mass needed when the full N normal is later completed against the original n packet. It is not a completion of that normal itself.

Here and below "infinitesimal cost" means that the budget multiplies the original positive packet by a bounded separated Gram coefficient with zero residue. Finite sums of such costs can be paid while retaining any fixed fraction smaller than one. Multiplication of the explicit SOS identities by pure-source nonnegative scalars and pure-testing squares is legal. The remaining diagonal coefficients are of the form $1 - \varepsilon$, with ε infinitesimal in the bounded tensor algebra, so the inherited bounded-Gram completion applies. Its actual text is the paragraph "Why a strict bounded Gram suffices" in `inherited_r19/parent_reference.tex`; only that field-independent lemma is imported, not the C14.2 source assumptions in the same file. For clarity, the elementary step is, for a pure-source infinitesimal s and bounded pure-testing t, to choose real $a, \varepsilon > 0$ with $|t| < a$ and $|s| < \varepsilon/a$ and use

$$\varepsilon \pm st = \tfrac{1}{2}\{(\varepsilon/a + s)(a \pm t) + (\varepsilon/a - s)(a \mp t)\}.$$

Every summand has separated nonnegative factors and hence a separated square root. Interchange the fields for the other type of infinitesimal entry and use a finite real diagonal budget. This proves the exact SOS lifting step used here. No square root or inverse of a mixed sum is used.

First $e_0^- \le g_c$. The exact budget

$$(1 + V^2)\Psi - V^2B_*^2 = Q_*^2 + R_*^2 + V^2(1 - UVZ_*)^2 \tag{RSR23}$$

pays its negative part from the circuit at cost

$$\frac{e_0^-}{f_c} + \frac{e_0^-}{f_cy}v^2 \;\le\; Hy\frac{\mathcal{L}}{\mathcal{F}} + H\frac{\mathcal{L}}{\mathcal{F}}v^2 \in \mathrm{I}.$$

Its positive part is retained as a square. Also $d_c > 0$ because $M_{b^2c} > 0$ and $k_c < 0$; the d_c and g_c terms cause no loss.

For k_c, use the exact identity

$$Z_*(V^2 - 1) = -VQ_* - R_*. \tag{RSR24}$$

The VQ_* part is paid from $q_cQ_*^2$ with coefficient $2|k_c|v^2/(q_cy)$, which is infinitesimal by RSR20. The remaining R_* part is paid in the following exhaustive three cases. Fixed real multipliers such as the factor two do not change any cost claim.

Case 1: $1 - p$ has a positive real lower bound. The exact ratio is

$$\frac{|k_c|}{r_c} = \frac{T\mathcal{K}}{(1 - p)(1 + Hy)\mathcal{L}} \in \mathrm{I}. \tag{RSR25}$$

This uses only $\mathcal{K}/\mathcal{L} = O(p + B)$ and $T \in \mathrm{I}$. Neither current mass nor a bound on j is required.

Case 2: $1 - p \in \mathrm{I}$ and η_* has a positive real lower bound. Here $p \sim 1$, so $Bj(j + 1)$ is a bounded positive unit. Since B is infinitesimal, j is unbounded and j^2B is a positive bounded unit. Equations RSR16–RSR20 give

$$\frac{|k_c|}{f_c} \in \mathrm{I}, \qquad \frac{|k_c|}{f_cx} = O\Big(\frac{T(p + B)}{j^2B}\Big) \in \mathrm{I}. \tag{RSR26}$$

The exact circuit budget

$$(1+U^2)\Psi = R_*^2 + B_*^2 + U^2Q_*^2 + U^2(1-UVZ_*)^2 \tag{RSR27}$$

therefore pays $|k_c|R_*^2$ at cost $|k_c|/f_c + (|k_c|/(f_cx))u^2 \in \mathrm{I}$. This case includes any intersection with literal $n = 0$.

Case 3: $1-p, \eta_* \in \mathrm{I}$. Now $p \sim 1$, $j > p$, and RSR9 gives $n > 0$ together with its exact relative estimate. The source mass M_e is a positive bounded unit. For the lower bound use

$$\frac{p^2}{M_e} = \frac{p^2}{h_0 j^2} + \frac{p^2}{nj^2} = O(1).$$

For the upper bound use $M_e \le h_0 j^2$ when j is bounded, and $M_e \le nj^2 \sim p/2$ when j is unbounded. In particular,

$$\frac{|k_c|}{f_c} \in \mathrm{I}, \qquad \frac{|k_c|}{q_c y} = O(1), \qquad \frac{|k_c|}{J_c x} = O\Big(\frac{T(p+B)(1+1/n)}{j^2}\Big) = O(1). \tag{RSR28}$$

The last bound follows directly from $1/n \sim 2(j-p)(j+1)/p$, with $p \sim 1$ and $j > p$. There is an exact separated SOS budget

$$2\{\Psi + V^2Q_*^2 + U^2B_*^2\} - R_*^2 \quad \text{is SOS.} \tag{RSR29}$$

For completeness, set $t_* = UV$, $\delta_* = U - V$, $E_* = Z_*(1-t_*)$, $T_* = t_*(1-Z_*)$, and $J_* = T_* - \delta_*^2 Z_*$. Its SOS decomposition is

$$(E_*+T_*)^2 + 2\delta_*^2 + 4t_*(1-Z_*)^2 + 2J_*^2 + 4t_*\delta_*^2 Z_*^2.$$

All roots are separated as in RSR13. Applying RSR29 costs a multiple of $|k_c|/f_c$ on the circuit, $|k_c|v^2/(q_cy)$ on Q_*, and $|k_c|u^2/(J_cx)$ on B_*. Each is infinitesimal by RSR28.

The three cases exhaust the source because $0 < 1-p < 1$ and η_* is bounded. A finite number of the displayed allocations therefore proves the scalar assertion following RSR22 with a strict retained fraction. All coefficient bounds depend only on the source and u, v; no bound on the independent state coordinate z was used. The proof allows arbitrary relative rates of all source and testing infinitesimals. Literal $n = 0$ is covered by Cases 1 and 2; Case 3 excludes it by the original physicality equation itself.

What remains outside this lemma. The scalar identity is exact for the complete graph of RSR10. A full-parent local theorem must also allocate both first-normal arrays and the complete quadratic block from RSR11, using the reserved current before applying signed scalar costs. The outer testing charts, the state chart $w = 0$, the sign exits and complex-state splitting must be supplied by their own exact arguments. This file makes none of those additional admissions by implication.

Chapter 18

Reconstructed small normal budgets for R1 unbounded-q

Component `p4b75bfe0436c`
Source: `REC/c144/rederived/r1_qo_small_normal_budgets.tex`
SHA256: `4b75bfe0436c28e3023493b22aed5841b7194e9f5340797765d210d6209d02fd`
Inventory use: NSR1-NSR5

18.1 Reconstructed small normal budgets for R1 unbounded-q

Scope and provenance. This is a fresh proof from the actual compact source RSR1–RSR9 and the exact circuit identity RSR23. It replaces the needed mathematics of the lost NS helper; it is not an assertion of original file bytes. Use all notation of `r1_qo_actual_source_and_scalar.tex`. The testing variables $u, v > 0$ are arbitrary independent infinitesimals, and $z > 0$ is arbitrary. Put

$$b = v - uz/j = \sqrt{y}\, B_*, \qquad h_0 = \mu + \tfrac{1}{2}, \qquad M_e = \frac{nh_0 j^2}{n + h_0}, \qquad J_c = M_e y. \tag{NSR1}$$

The two rows

$$pv^2 b, \qquad \sqrt{T}\,\mu uvb \tag{NSR2}$$

can be given any fixed real positive square coefficient at infinitesimal cost from $f_c\Psi + J_c B_*^2$. The same holds after multiplying either row by any bounded separated coefficient. No coefficient in the budget depends on z.

The infinitesimal-η_* source case. By RSR9, $n > 0$ and $n \sim p/[2(j-p)(j+1)]$. The exact identity

$$\frac{p^2}{M_e} = \frac{p^2}{h_0 j^2} + \frac{p^2}{nj^2} \tag{NSR3}$$

is bounded: the first term is bounded since $p/j < 1$; the second is comparable to $p(j-p)(j+1)/j^2$. If $j \le 1$ this last expression is at most $2p/j$, and if $j \ge 1$ it is at most $2p$. Thus the direct costs of the squared rows in NSR2, relative to $J_c B_*^2 = M_e b^2$, are

$$\frac{p^2}{M_e} v^4 \in \mathrm{I}, \qquad \frac{T\mu^2}{M_e} u^2 v^2 \in \mathrm{I}. \tag{NSR4}$$

Here μ/p is bounded by RSR6–RSR7. No use of the circuit is needed in this case, and no positive lower bound on j is imposed.

The positive-unit-η_* source case. The remaining possibility is that η_* has a positive real lower bound, since RSR9 already proves it bounded. The exact relation

$$\frac{p^2}{j^2 B} = \frac{p + p/j}{\eta_*}$$

and RSR16–RSR17 imply $p^2 y / f_c$ bounded. Apply

$$(1 + V^2)\Psi - V^2 B_*^2 = Q_*^2 + R_*^2 + V^2(1 - UVZ_*)^2. \tag{NSR5}$$

The first row in NSR2 costs exactly

$$\frac{p^2 y}{f_c} v^2 (y + v^2) \in \mathrm{I}$$

on $f_c \Psi$. Indeed the difference between that multiple of the circuit packet and $(pv^2 b)^2$ is $p^2 y^3 V^2$ times the SOS on the right of NSR5. The second row costs exactly

$$\frac{T\mu^2 y}{f_c} u^2 (y + v^2) \in \mathrm{I};$$

its difference is $T\mu^2 x y^2 U^2$ times the same SOS. Every multiplier is a product of nonnegative pure-source and pure-testing factors. These formulas include literal $n = 0$, since they never divide by the current mass. The two source cases exhaust the source.

Use in the full normal conductor. The first row pays the $(p/2)v^2 b$ term in the complete half-N derivative. The second pays the μuvb term in the half-M derivative after its source normalization by $\sqrt{TH(1-H)}$: the omitted factor $\sqrt{H(1-H)}$ is bounded. This is only a square-budget lemma. The exact full-normal identities and the retained hard block are required separately to infer a full-parent local certificate.

Chapter 19

A freshly derived full-state sign and projective join

Component `pf77de02d80c4`
Source: `REC/root_rederived/full_state_sign_join_REDERIVED.tex`
SHA256: `f77de02d80c4b891fe1da0e42e7b9d4a3ca8a692f18d9df6c7b623a4ac9bca1c`
Inventory use: Exact sign gauges and homogeneous/state normalization identities used by the local terminal bodies | SJ1-SJ10 | actual SJ1–10 state, sign and projective identities

19.1 A freshly derived full-state sign and projective join

This section is a new proof derived from the complete weak-C2 parent retained in the earlier R21 checkpoint. It does not claim byte identity with the missing late-R21 state-join file. Its conclusion is a reduction of the full parent to one positive state-ratio chart. The certificate in that remaining chart is a separate hypothesis; no global finite-lift conclusion is drawn here.

Let $K_c, K_t \supset \mathbb{R}$ be real closed fields, occupying separately labelled tensor factors. All quantities u, v, W, X, Y, Z below belong to K_t; all six coefficients n, m, A, C, D, α belong to K_c and are nonnegative. Write $D_u = 1 + u^2$, $D_v = 1 + v^2$, and define the complete parent

$$F = nD_uD_vX^2 + m\{v(W + uY) - X - uZ\}^2 + AD_uD_vY^2 + CD_v(uW - Y)^2 \tag{19.1}$$
$$+ D(uW + uvX - Y - vZ)^2 + \alpha D_u(vY - Z)^2 + K, \tag{SJ1}$$
$$K = \tfrac{1}{2}(uW + uvX - Y - vZ)^2 + \tfrac{1}{2}D_u(vY - Z)^2 + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \tag{SJ2}$$

Products involving source and testing quantities mean tensor products. For example, $nD_uD_vX^2$ means $n \otimes (D_uD_vX^2)$; a testing-only term means $1 \otimes K$. The positive packets in (SJ1) are sums of squares in $K_c \otimes_{\mathbb{R}} K_t$, since each coefficient and each testing multiplier has a square root in its own real closed field.

Sign gauges preserve the entire parent. For $\varepsilon, \eta \in \{1, -1\}$, the substitution

$$(u, v, W, X, Y, Z) \longmapsto (\varepsilon u, \eta v, W, \eta X, \varepsilon Y, \varepsilon\eta Z) \tag{SJ3}$$

preserves every summand of (SJ1), including (SJ2). Indeed the roots of the m, C, D, α packets acquire, respectively, the signs $\eta, \varepsilon, \varepsilon, \varepsilon\eta$, and the other roots also change only by a sign. Thus u, v may both be taken nonnegative without altering the source coefficients or the full expression.

An exact positive hard block. Set

$$t = \frac{X + uZ}{D_u}, \qquad r = \frac{Y - uW}{D_u}, \qquad w = \frac{W + uY}{D_u}, \qquad z = \frac{Z - uX}{D_u}. \tag{SJ4}$$

The inverse transformation is $X = t - uz$, $Z = z + ut$, $Y = r + uw$, $W = w - ur$. Let $H = 1 + u^2 + v^2$ and $d = H^2 - u^2v^2$. Both $H - uv = 1 + (u - v)^2 + uv$ and $H + uv$ are positive for $u, v \geq 0$; in fact $H > |uv|$ for arbitrary u, v. Consequently $d > 0$. Define

$$\mathsf{A} = \frac{1}{2}\begin{pmatrix} H & -uv \\ -uv & H \end{pmatrix}, \qquad \mathsf{B} = \frac{1}{2}\begin{pmatrix} -u^2v & -uv^2 \\ uv^2 & u^2v \end{pmatrix}.$$

The matrix A is positive definite. Direct expansion and square completion give the exact identity

$$\frac{K}{D_u} = \left(\begin{pmatrix} t \\ r \end{pmatrix} + \mathsf{A}^{-1}\mathsf{B}\begin{pmatrix} w \\ z \end{pmatrix}\right)^T \mathsf{A}\left(\begin{pmatrix} t \\ r \end{pmatrix} + \mathsf{A}^{-1}\mathsf{B}\begin{pmatrix} w \\ z \end{pmatrix}\right) \tag{19.2}$$

$$+ \frac{D_u D_v}{2d}\{u^2v^2(w^2 + z^2) - 2uvHwz\}. \tag{SJ5}$$

The rational matrix used in the shift is explicitly

$$\mathsf{A}^{-1}\mathsf{B} = \frac{1}{d}\begin{pmatrix} -u^2v(1 + u^2) & -uv^2(1 + v^2) \\ uv^2(1 + v^2) & u^2v(1 + u^2) \end{pmatrix}. \tag{SJ6}$$

For any $a, b \in K_t$, the hard quadratic form is

$$\begin{pmatrix} a \\ b \end{pmatrix}^T \mathsf{A}\begin{pmatrix} a \\ b \end{pmatrix} = \frac{H - uv}{4}(a + b)^2 + \frac{H + uv}{4}(a - b)^2. \tag{SJ7}$$

This also gives an explicit square decomposition of the first term of (SJ5), with square roots taken only in K_t.

If $u, v \geq 0$ and $wz \leq 0$, the second term of (SJ5) is nonnegative. It too is a square in K_t (or can be split into its three nonnegative terms). Hence $K \geq 0$, and (SJ1) is a tensor sum of squares immediately. In particular the axes $w = 0$ and $z = 0$ are included, without division by either state coordinate. The same identity, without sign gauges, proves the sharper implication $K < 0 \Longrightarrow uvwz > 0$.

Reduction of the remaining states. The only remaining case after (SJ3) is $K < 0$, $u, v > 0$, $wz > 0$. Then $w \neq 0$. Divide all four state coordinates by w. Equation (SJ4) now has $w' = 1$ and $z' = z/w > 0$. The ratio z' is arbitrary: it may be infinitesimal, bounded, or unbounded in K_t. No estimate in this reduction replaces z' by a bounded coefficient.

The parent is homogeneous of degree two in the state coordinates, so

$$F(u, v; W, X, Y, Z) = w^2 F(u, v; W/w, X/w, Y/w, Z/w). \tag{SJ8}$$

If the parent on the right has a certificate $\sum_i h_i^2$ in $K_c \otimes_{\mathbb{R}} K_t$, then the parent on the left has certificate $\sum_i ((1 \otimes w)h_i)^2$. This proves that certificates for the full parent with $u, v > 0$, $w = 1$, and arbitrary positive z suffice for all finite affine testing parameters and all real states.

Literal projective endpoints. For clarity, the projective endpoints can be checked without a limiting argument. Homogenize $u = U_1/U_0$, $v = V_1/V_0$, and put $D_U = U_0^2 + U_1^2$, $D_V = V_0^2 + V_1^2$, with neither pair identically zero. Multiplication by $U_0^2V_0^2$ turns (SJ2) into

$$K_h = \tfrac{1}{2}(U_1V_0W + U_1V_1X - U_0V_0Y - U_0V_1Z)^2 \tag{19.3}$$

$$+ \tfrac{1}{2}D_U(V_1Y - V_0Z)^2 + \tfrac{1}{2}D_V(U_0X + U_1Z)^2 - \tfrac{1}{2}D_U D_V Z^2. \tag{SJ9}$$

At the four axes this is, exactly,

$$U_0 = 0: \quad K_h = \tfrac{1}{2}U_1^2\{(V_0W + V_1X)^2 + (V_1Y - V_0Z)^2\}, \tag{19.4}$$

$$V_0 = 0: \quad K_h = \tfrac{1}{2}V_1^2 D_U(X^2 + Y^2), \tag{19.5}$$

$$U_1 = 0: \quad K_h = \tfrac{1}{2} U_0^2 D_V (X^2 + Y^2), \tag{19.6}$$

$$V_1 = 0: \quad K_h = \tfrac{1}{2} V_0^2 \{(U_1 W - U_0 Y)^2 + (U_0 X + U_1 Z)^2\}. \tag{SJ10}$$

All other homogenized packets in (SJ1) retain nonnegative coefficients and are sums of squares. Thus the endpoints are admitted directly, including the intersections of these axes. Equations (SJ3)–(SJ10) establish the stated full-state, sign, and endpoint reduction. They do not establish the still-required positive-ratio chart certificate.

Verification scope. The companion exact checker verifies the inverse transformation, the entire parent under all four gauges, the hard-block and Schur identities, homogeneity, homogenization, and all four endpoint formulas. These are algebraic corroborations of the proof above; they are not a formal proof of the complete R21 theorem and do not certify other missing components.

Chapter 20

Reconstructed full R1 conductor in the unbounded source-ratio sector

Component `p261f71af685d`
Source: `REC/weak_c2/fn_reconstructed.tex`
SHA256: `261f71af685d3738c43fd565d11e383655bfe4f1ff1273a13023b75815b06434`
Inventory use: RFN1-RFN24

20.1 Reconstructed full R1 conductor in the unbounded source-ratio sector

Provenance and precise dependency. This is a newly written reconstruction of the late R21 conductor, not a claim of byte identity with the lost final file. The complete parent, the radial Gram matrices, and the normal formulas were retained in the conversation; every displayed polynomial identity below is checked again against the recovered original source equations. The companion `check_fn_reconstruction.py` is also new. The source dictionary is the independently rederived `c144/rederived/r1_qo_compact_source.py`. The uniform source bounds and the scalar allocation stated in the next paragraph are the explicit interface to the reconstructed actual-source scalar lemma (RS). The source input is now supplied by the freshly reconstructed `c144/rederived/r1_qo_actual_source_and_scalar.tex`, RSR1–RSR29, with its independent current review. The full-normal argument below has now received an independent current analytical review by the lead agent, separate from the review of its scalar input. Neither the old admission record nor the old checksum substitutes for it.

Fields, source scope, and notation. Let $K_c, K_t \supset \mathbb{R}$ be real closed fields, with bounded rings B_c, B_t and infinitesimal ideals $\mathfrak{m}_c, \mathfrak{m}_t$. The ring of certificate expressions is $\mathcal{T} = K_c \otimes_{\mathbb{R}} K_t$, and its bounded coefficient ring is $\mathcal{B} = B_c \otimes_{\mathbb{R}} B_t$. A bounded coefficient with zero real residue belongs to $\mathfrak{n} = \mathfrak{m}_c \otimes B_t + B_c \otimes \mathfrak{m}_t$. Every radical or inverse used below is pure source, pure testing, or an explicitly separated product of such factors. No order on $\mathcal{T}$ and no inverse of a mixed expression is used.

Use the positive compact source coordinates

$$x = jB, \qquad y = TB, \qquad q_z = jT, \qquad 0 < p < \min\{1, j\}, \qquad 0 < H < 1, \tag{RFN1}$$

where $B, T, x, y, q_z \in \mathfrak{m}_c$. These are the compact coordinates of the original R1 sector with unbounded source ratio. The original strict source gaps remain in force and $n \geq 0$, including $n = 0$. For reference, their native substitution is

$$r = \frac{j-p}{j+px}, \quad h = \frac{1-HT}{1+HTx}, \quad a_s = \frac{j+px}{1+HTx}, \quad B_s = \frac{B(1+HTx)}{1+pB}, \quad t_s = \frac{T(1+pB)}{1+HTx}. \tag{RFN2}$$

The six weights n, m, C, A, D, α are always the original source weights at (RFN2), rather than weights of an auxiliary stationary source. Put $\mu = m(1-p)$, $h_0 = \mu + \frac{1}{2}$, $k = (j+1-p)/(j+1)$, and $\eta_* = Bj(j+1)/p$. In this source sector RS proves

$$\tfrac{1}{2} < k < 1, \quad \eta_* = O(1), \quad \mu = O(p), \quad h_0 \asymp 1, \quad A = O(T), \quad \alpha \asymp B^2(j+1), \tag{RFN3a}$$

$$TCH \sim k/2, \qquad T(D+\tfrac{1}{2})(1-H) \sim k/2, \qquad \frac{n(j-p)}{j} = O(1), \tag{RFN3b}$$

$$f_c \asymp Tx^2, \qquad q_c \asymp Tx, \qquad r_c \asymp TjB^2(1-p), \tag{RFN3c}$$

$$\eta_* \in \mathfrak{m}_c \quad \Longrightarrow \quad n \sim \frac{p}{2(j-p)(j+1)} > 0. \tag{RFN3d}$$

Here $\asymp$ means that the ratio is a positive bounded unit, and $\sim$ means that the ratio has residue one. All estimates are uniform at the independent openings H, $1-H$, $j-p$, and $1-p$. In particular $B(j+1) = B + x$ and α are infinitesimal. No comparison between source and testing infinitesimals is allowed.

The local conclusion. For all positive independent $u, v \in \mathfrak{m}_t$, normalize the pure testing coordinate $w = (W+uY)/(1+u^2)$ to one and allow the other pure testing coordinate $z = (Z-uX)/(1+u^2) > 0$ to be arbitrary. The conclusion is a finite tensor sum of squares for the complete original parent, with both state normals unrestricted. In particular z need not be bounded. The separate full-state sign reduction and outer-testing join are needed to pass from this local conclusion to the whole R1 row.

20.1.1 The complete parent and the correlated source mass

Write $D_u = 1 + u^2$, $D_v = 1 + v^2$. The original quarter-parent is

$$\mathcal{P} = m(vW - X + uvY - uZ)^2 + nD_uD_vX^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \tag{20.1}$$

$$+ (D+\tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha+\tfrac{1}{2})D_u(vY - Z)^2 \tag{20.2}$$

$$+ \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \tag{RFN4}$$

Use the two mixed normal rows

$$N_0 = \frac{X + uZ - pD_uv}{D_u}, \qquad M_0 = \frac{uW - Y - HD_uvz}{D_u}. \tag{RFN5}$$

Their exact inverse, with $w = 1$, is

$$\begin{aligned} X &= N_0 + pv - uz, & Z &= z + uN_0 + puv, \\ W &= 1 + u(M_0 + Hvz), & Y &= u - M_0 - Hvz. \end{aligned} \tag{RFN6}$$

Only the pure testing unit D_u is inverted. Retain the nonnegative packet $mpD_u^2N_0^2$ and define $G = (\mathcal{P} - mpD_u^2N_0^2)/D_u$. Direct substitution gives the *entire* expression

$$G = \mu D_u\{N_0^2 - 2vN_0 + (1-p)v^2\} + nD_v(N_0 + pv - uz)^2 \tag{20.3}$$

$$+ CD_uD_v(M_0 + Hvz)^2 + AD_v(u - M_0 - Hvz)^2 \tag{20.4}$$

$$+ (D+\tfrac{1}{2})D_u(M_0 - (1-H)vz)^2 \tag{20.5}$$

$$+ (\alpha+\tfrac{1}{2})\{u(1-p)v - (1+Hv^2)z - uN_0 - vM_0\}^2 \tag{20.6}$$

$$+ \tfrac{1}{2}D_uD_v(N_0 + pv)^2 - \tfrac{1}{2}D_v(z + uN_0 + puv)^2. \tag{RFN7}$$

Thus G is exactly quadratic in (N_0, M_0):

$$G = F + 2L_NN_0 + 2L_MM_0 + H_NN_0^2 + H_MM_0^2 + 2(\alpha+\tfrac{1}{2})uvN_0M_0, \tag{RFN8}$$

where $F = G|_{N_0=M_0=0}$ and

$$\begin{aligned} H_N &= h_0D_u + \tfrac{1}{2}v^2 + \alpha u^2 + nD_v, \\ H_M &= D_u(CD_v + D + \tfrac{1}{2}) + AD_v + (\alpha+\tfrac{1}{2})v^2. \end{aligned} \tag{RFN9}$$

There are no omitted higher state-normal terms.

Exact scalar interface, before spending the n-mass. Set

$$U=\frac{u}{\sqrt{x}},\quad V=\frac{v}{\sqrt{y}},\quad Z_*=\frac{z}{\sqrt{q_z}},\qquad Q_*=U-VZ_*,\quad B_*=V-UZ_*,\quad R_*=Z_*-UV,$$

$$\Psi=U^2+V^2+Z_*^2+U^2V^2Z_*^2-4UVZ_*. \tag{RFN10}$$

All three normalized testing variables are positive, but none is assumed bounded. A legal separated certificate is

$$\Psi=(U-V)^2+Z_*^2(1-UV)^2+2UV(1-Z_*)^2. \tag{RFN11}$$

For example, $\sqrt{UV}$ is the product of the source factor $(xy)^{-1/4}$ and the testing factor $\sqrt{uv}$.

The exact RS/RB scalar identity for the actual source is

$$F=q_cQ_*^2+nj^2yB_*^2+r_cR_*^2+d_cV^2Q_*^2+(np^2y^2+e_0)V^2B_*^2 \tag{20.7}$$
$$+f_c\Psi+g_cV^2\Psi+k_cZ_*^2(V^2-1)^2,\qquad e_0=\tfrac{1}{2}p^2y^2-g_c. \tag{RFN12}$$

For an entirely algebraic definition of the masses, put $K_0=CH^2+(D+\frac{1}{2})(1-H)^2$ and

$$\begin{aligned}
&M_a=Ax, && M_b=\{\mu(1-p)+(n+\tfrac{1}{2})p^2\}y,\\
&M_c=\alpha q_z, && M_{ab}=\{A+\alpha(1-p)^2+\mu(1-p)+\tfrac{1}{2}(1-p)^2\}xy,\\
&M_{ac}=nxq_z=nj^2y, && M_{bc}=\{K_0+AH^2+2H\alpha+H-\tfrac{1}{2}\}yq_z,\\
&M_{b^2}=(n+\tfrac{1}{2})p^2y^2, && M_{abc}=(K_0+n)xyq_z,\\
&M_{b^2c}=H^2(A+C+\alpha+\tfrac{1}{2})y^2q_z, && M_{ab^2c}=CH^2xy^2q_z.
\end{aligned}$$

Then

$$\begin{aligned}
&k_c=(M_a-M_b+M_{ac}+M_{ab^2c}-M_{bc})/2,\\
&r_c=M_c-M_b+M_{ac}-k_c,\quad f_c=M_b-M_{ac},\quad q_c=M_a-f_c,\\
&d_c=M_{b^2c}-k_c,\qquad g_c=M_{ab^2c}.
\end{aligned}$$

The source contact identities make (RFN12) exact, including both odd monomials of F. It is checked coefficientwise against (RFN7), not only at the contact point.

Define the unnormalized current $b=v-uz/j=\sqrt{y}B_*$ and the pure source quantities

$$c_b=\frac{nj}{n+h_0},\qquad M_e=\frac{nh_0j^2}{n+h_0}. \tag{RFN13}$$

The whole scalar pieces $nj^2b^2+np^2v^2b^2$ in (RFN12) must be kept until their correlations with N_0 have been completed. The full N_0-linear term below contains $njb+npv^2b$, so these pieces and the corresponding hard terms give

$$n(N_0+jb)^2+nv^2(N_0+pb)^2.$$

Complete only the first square, using a pure source inverse:

$$h_0N_0^2+n(N_0+jb)^2=(n+h_0)(N_0+c_bb)^2+M_eb^2. \tag{RFN14}$$

Keep $nv^2(N_0+pb)^2$ whole. Put $\widehat{N}=\sqrt{n+h_0}(N_0+c_bb)$. The other hard terms in (RFN9) are the nonnegative whole squares

$$h_0(uN_0)^2+\tfrac{1}{2}(vN_0)^2+\alpha(uN_0)^2. \tag{RFN15}$$

They are still kept whole after the shift; no term nu^2 or nv^2 is treated as a bounded perturbation.

For $n>0$ one has the exact bounded-shift identity

$$\frac{c_b^2}{M_e}=\frac{n}{h_0(n+h_0)}=O(1). \tag{RFN16}$$

At $n = 0$, set $c_b = M_e = 0$ and omit the effective current row. Formula (RFN14) remains literal, and no division by n or $\sqrt{M_e}$ is made in that case.

RS pays the signed e_0 and k_c terms in (RFN12) at infinitesimal cost after (RFN14), with the effective current mass $M_e y$, and leaves fixed positive real fractions of

$$q_c Q_*^2, \qquad M_e b^2, \qquad r_c R_*^2, \qquad f_c \Psi. \tag{RFN17}$$

The second term is omitted when $n = 0$. This is the precise scalar allocation needed here; an allocation that spends the raw $nj^2 y$ mass before (RFN14) would not suffice.

20.1.2 Finite exact budgets for the full normal rows

In the rest of the proof, a row is called free if any prescribed fixed real multiple of its square can be purchased at infinitesimal cost from (RFN17). This has an exact meaning: a finite SOS identity writes its square below a finite sum of those retained squares, with bounded separated coefficients in $\mathfrak{n}$. Subtracting any fixed real multiple changes their strict Gram diagonals only infinitesimally.

Three elementary circuit budgets. Direct expansion gives

$$(1+U^2)\Psi - R_*^2 = B_*^2 + U^2 Q_*^2 + U^2(1 - UVZ_*)^2, \tag{RFN18a}$$

$$(1+V^2)\Psi - V^2 B_*^2 = Q_*^2 + R_*^2 + V^2(1 - UVZ_*)^2. \tag{RFN18b}$$

There is also a useful opposite-current budget. Put $t_* = UV$, $\delta_* = U - V$, $E_* = Z_*(1 - t_*)$, $T_* = t_*(1 - Z_*)$, and $J_* = T_* - \delta_*^2 Z_*$. Then

$$2(\Psi + V^2 Q_*^2 + U^2 B_*^2) - R_*^2 = (E_* + T_*)^2 + 2\delta_*^2 + 4t_*(1 - Z_*)^2 \tag{20.8}$$

$$+ 2J_*^2 + 4t_*\delta_*^2 Z_*^2. \tag{RFN18c}$$

Every term on the right is a legal separated square. Thus a nonnegative multiplier a buys aR_*^2 at infinitesimal cost whenever

$$a/f_c \in \mathfrak{n}, \qquad a/(q_c y) = O(1), \qquad a/(M_e y x) = O(1), \tag{RFN18d}$$

since the last two diagonal costs have the additional testing factors v^2 and u^2. This criterion is used only when $M_e > 0$.

An exact radial budget. The row $R_u = v(u^2 - x)$ is free in all source subcases, including the weak opening $1 - p \in \mathfrak{m}_c$. Here is an explicit finite certificate, so that this assertion does not rest on pointwise nonnegativity. Write $D_* = V(U^2 - 1)$ and

$$t_7 = (Z_* - U^2V,\ V - U^2V,\ U - U^2V,\ UZ_* - U^2V,\ UV - U^2V,\ UVZ_* - U^2V,\ U^2 - U^2V)^T,$$

$$t_3 = (1 - UV,\ V - UV,\ U - UV)^T,$$

$$G_7 = \begin{pmatrix} 4 & -2 & -3/2 & 0 & -1/2 & 0 & 1/2 \\ -2 & 3 & 0 & -3/2 & 0 & -1 & 2 \\ -3/2 & 0 & 4 & 3/2 & -2 & -3 & 0 \\ 0 & -3/2 & 3/2 & 4 & -3/2 & 0 & -3/2 \\ -1/2 & 0 & -2 & -3/2 & 3 & 2 & -1 \\ 0 & -1 & -3 & 0 & 2 & 8 & -4 \\ 1/2 & 2 & 0 & -3/2 & -1 & -4 & 4 \end{pmatrix}, \qquad G_3 = \begin{pmatrix} 3 & -2 & -2 \\ -2 & 2 & 2 \\ -2 & 2 & 3 \end{pmatrix}.$$

The exact identity is

$$4\Psi + 4U^2 Q_*^2 + 4U^2 R_*^2 - D_*^2 = t_7^T G_7 t_7 + 4VZ_*(1 - U)^2 + UZ_* t_3^T G_3 t_3. \tag{RFN19}$$

The leading principal minors of G_7 are $4, 8, 101/4, 1049/16, 907/16, 3279/16, 743/16$; those of G_3 are $3, 2, 2$. Both are positive definite real matrices. Their real Cholesky factors and the separated square roots of VZ_* and UZ_* therefore make the right side a finite legal SOS. Denote it by $\mathcal{W}(U, V, Z_*)$.

Since $R_u = x\sqrt{y}D_*$, the exhaustive source cases are as follows.

Case 1: $1-p$ *is a positive unit.* Multiply (RFN19) by x^2y. The costs relative to $f_c\Psi, q_cQ_*^2, r_cR_*^2$ are, respectively,

$$O(B), \qquad O(Bu^2), \qquad O(u^2/(1-p)).$$

All belong to $\mathfrak{n}$.

Case 2: $p \sim 1$ *and* η_* *is a positive unit.* Then j has a positive real lower bound, and j^2B is a positive unit. Adding $4U^2$ times (RFN18a) to (RFN19) gives

$$D_*^2 \ \preceq_{\mathrm{sos}} \ 4(1+U^2+U^4)\Psi + 4U^2Q_*^2.$$

After multiplication by x^2y, the costs are

$$O\Big(B + u^2/j + u^4/(j^2B)\Big), \qquad O(Bu^2),$$

again infinitesimal. The notation $a \preceq_{\mathrm{sos}} b$ means $b-a \in \Sigma\mathcal{T}^2$ and is only shorthand for the displayed exact identities.

Case 3: $p \sim 1$ *and* $\eta_* \in \mathfrak{m}_c$. Formula (RFN3d) gives $n > 0$ and $1/M_e = O(1)$; it also gives $p^2/M_e = O(1)$. Put $E^{\mathrm{rad}} = Z_*(U^2-1)$. Permuting V and Z_* in (RFN19) yields

$$\mathcal{W}(U, Z_*, V) = 4\Psi + 4U^2Q_*^2 + 4U^2B_*^2 - (E^{\mathrm{rad}})^2.$$

The identity $D_* = UE^{\mathrm{rad}} + (U^2-1)B_*$ then gives, exactly,

$$\begin{aligned} &8U^2\Psi + 8U^4Q_*^2 + \{8U^4 + 2(U^2-1)^2\}B_*^2 - D_*^2 && (20.9)\\ &\quad = 2U^2\mathcal{W}(U, Z_*, V) + \{UE^{\mathrm{rad}} - (U^2-1)B_*\}^2. && (\mathrm{RFN20}) \end{aligned}$$

After multiplication by x^2y, the three costs are

$$O(u^2/j), \qquad O(u^4/j), \qquad \frac{8u^4 + 2(u^2-x)^2}{M_e}.$$

They are infinitesimal. These three cases prove the asserted free radial row without assuming a rate for any testing variable.

The smaller current helpers. If $\eta_* \in \mathfrak{m}_c$, (RFN3d) implies

$$\frac{p^2}{M_e} = \frac{p^2}{h_0 j^2} + \frac{p^2}{nj^2} = O(1),$$

because $p/j < 1$ and $p(j-p)(j+1)/j^2 \le p + p/j$. Thus $(pv^2b)^2$ costs only $(p^2/M_e)v^4$ of the effective current. If η_* is a positive unit instead, then

$$\frac{p^2y}{f_c} = O\Big(\frac{p + p/j}{\eta_*}\Big) = O(1).$$

Multiply (RFN18b) by $p^2y^2v^2$. This buys $(pv^2b)^2$ with circuit cost

$$\frac{p^2y}{f_c}(yv^2 + v^4) \in \mathfrak{n}.$$

The same identity multiplied by $Tp^2u^2y^2$ shows that $\sqrt{T}\,puvb$ is free: its cost is either $(p^2/M_e)Tu^2v^2$ or $(p^2y/f_c)Tu^2(y+v^2)$.

20.1.3 The entire two-normal conductor

First normal: exact source cancellation and allocation. Put

$$q = vz - Tu, \qquad R = z - uv/B, \qquad R_u = v(u^2 - x),$$

$$K_c = \frac{p}{2} + j\{H(\alpha + \tfrac{1}{2}) - \tfrac{1}{2}\} + n(p - j), \qquad K_R = -\mu - (\alpha + \tfrac{1}{2})(1 - p) + \frac{T}{j}K_c.$$

Both K_c/j and K_R are bounded by (RFN3). The complete first normal is exactly

$$L_N = \{j(n - \alpha) + p(n + \tfrac{1}{2})v^2\}b + K_R R_u + \frac{K_c}{j}uvq. \tag{RFN21}$$

In particular the sign of the μ term in K_R is negative. For a direct source check, the difference between this formula and the derivative of (RFN7) is vS_N, where

$$S_N = np - \mu + \tfrac{p}{2} + \{-\mu - (\alpha + \tfrac{1}{2})(1 - p)\}x + j(\alpha - n) + yK_c = 0.$$

The last equality is an exact rational identity of the actual native source under (RFN2), not an asymptotic contact equation.

After the two correlated n-squares are formed, the remaining linear coefficient is

$$L_N^{\mathrm{rem}} = -j\alpha b + \tfrac{1}{2}pv^2 b + K_R R_u + \frac{K_c}{j}uvq.$$

The q row is strict because $q_c Q_*^2 = (q_c/(Tx))(q^2/T)$, with $q_c/(Tx)$ a positive unit. Its coefficient in L_N^{rem} relative to $q/\sqrt{T}$ is $(K_c/j)uv\sqrt{T} \in \mathfrak{n}$. The $pv^2 b$ helper is free by the preceding paragraph. For the remaining small row, the exact identity

$$j\alpha b = -\frac{\alpha}{B}R_u - \alpha uR$$

has $\alpha/B = O(B + x) \in \mathfrak{m}_c$. When $1 - p$ is a unit, the square of αuR costs $O(B^2(j+1)^2u^2/(1-p)) \in \mathfrak{n}$ against $r_c R_*^2$. When $p \sim 1$, use $B_*^2 \preceq_{\mathrm{sos}} (1 + U^2)\Psi$ from (RFN18a). The circuit cost of $(j\alpha b)^2$ is

$$O\Big(B^3(j+1)^2 + B^2(j+1)^2u^2/j\Big) \in \mathfrak{n},$$

since j is bounded below and $B(j + 1) = B + x$ is infinitesimal. Finally, $K_R R_u$ has bounded coefficient; buy an arbitrarily large fixed real R_u^2 pivot using (RFN19)–(RFN20).

The shift $N_0 = \widehat{N}/\sqrt{n + h_0} - c_b b$ introduces no new unbounded Gram coefficient: $1/\sqrt{n + h_0} = O(1)$, and for $n > 0$ $c_b/\sqrt{M_e} = O(1)$ by (RFN16). The only bounded, potentially non-infinitesimal couplings are therefore those of R_u to $\widehat{N}$ and $\sqrt{M_e}b$. A sufficiently large fixed real radial pivot absorbs both while preserving a strict positive real residue. For $n = 0$ the latter coupling is exactly absent.

Second normal: cancellation of both source openings. Define

$$K_d = CH - (D + \tfrac{1}{2})(1 - H), \qquad C_q = K_d + AH + \tfrac{1}{2} + v^2\{CH(1 + u^2) + AH + (\alpha + \tfrac{1}{2})H\}.$$

The complete second normal is

$$L_M = C_q q + \mu uvb + v(\alpha - BTCHu^2)R. \tag{RFN22}$$

The actual source satisfies the three exact identities

$$\begin{aligned}
S_0 &= T(K_d + AH + \tfrac{1}{2}) - A = 0,\\
S_1 &= jK_d + \mu + xTCH = 0,\\
S_2 &= jK_d + xTCH + T\{CH + AH + (\alpha + \tfrac{1}{2})H\} - A - (\alpha + \tfrac{1}{2})(1 - p) + \alpha/B = 0.
\end{aligned}$$

Before imposing these identities, the complete derivative minus (RFN22) is exactly $uS_0+u^2vzS_1/j+uv^2(S_2-S_1)$. This records all state terms and rules out a hidden j-weighted radial error. In particular S_1 gives $K_d=-\mu/j-BTCH=O(1)$ even when C or D is unbounded. Consequently $TC_q=O(T+v^2)\in\mathfrak{n}$.

Normalize

$$M_0=\sqrt{TH(1-H)}\,\widehat{M}.$$

The full diagonal (RFN9), not only its zero-angle value, obeys

$$TH(1-H)H_M\sim k/2>0. \tag{RFN23}$$

Indeed the C and D terms contribute $(k/2)D_u\{(1-H)D_v+H\}$ up to infinitesimal bounded coefficients, and all remaining terms are infinitesimal after this normalization. This is uniform at both independent openings. The normalized C_qq coupling has coefficient $T\sqrt{H(1-H)}C_q\in\mathfrak{n}$ on the strict row $q/\sqrt{T}$. The normalized μuvb row is free because $\mu=O(p)$ and $\sqrt{T}\,puvb$ has already been purchased.

It remains to allocate the entire opposite row, including its higher testing terms. Put

$$J=B(j+1)+u^2\in\mathfrak{n},\qquad a_R=Tq_zv^2(\alpha-BTCHu^2)^2.$$

The factor $H(1-H)$ may be discarded from the cost bound, since it is bounded. Since $\alpha-BTCHu^2=O(BJ)$, the exact squared multiplier a_R satisfies

$$\frac{a_R}{f_c}=O((T/j)v^2J^2),\qquad \frac{a_R}{q_cy}=O(v^2J^2). \tag{RFN24a}$$

If $1-p$ is a unit, its direct cost against $r_cR_*^2$ is $O(Tv^2J^2/(1-p))\in\mathfrak{n}$. If $p\sim1$ and η_* is a unit, apply (RFN18a); the cost is $(a_R/f_c)(1+u^2/x)\in\mathfrak{n}$, using that j^2B is a unit. If $p\sim1$ and $\eta_*\in\mathfrak{m}_c$, (RFN3d) gives

$$\frac{a_R}{M_eyx}=O\Big(\frac{T(1+1/n)}{j^2}v^2J^2\Big)=O(1), \tag{RFN24b}$$

because $(1+1/n)/j^2=O(1)$ in this source case. The other two conditions in (RFN18d) follow from (RFN24a). The exact budget (RFN18c) therefore buys this row at infinitesimal cost as well. These estimates treat the whole square $(\alpha-BTCHu^2)^2$; there is no truncation at a testing degree.

Full Gram, legal roots, and restoration. After the normalizations, the entire remaining mixed hard term is

$$2(\alpha+\tfrac12)uv\sqrt{TH(1-H)}\left(\frac{\widehat{N}}{\sqrt{n+h_0}}-c_bb\right)\widehat{M}.$$

Its coefficients relative to $\widehat{N}$, $\widehat{M}$, and, when present, $\sqrt{M_e}b$ are infinitesimal; (RFN16) controls the shift. All other remaining normal couplings were allocated above. Choose a fixed sufficiently large radial pivot, and then purchase all the finitely many free helpers. Their scalar costs lie in $\mathfrak{n}$, so fixed positive fractions of (RFN17) survive. The complete bounded Gram has strict positive real residue: the $\widehat{N}$ pivot is one, the $\widehat{M}$ pivot has residue $\overline{k}/2$, the retained scalar pivots are positive, and the only bounded radial off-diagonals have been absorbed by the chosen real radial pivot.

The separated strict bounded-Gram lemma now gives a finite sum of squares in $\mathcal{T}$. Its row entries need not be bounded. All budget remainders are explicit finite separated sums of squares, all source-only completions use positive pure source denominators, and all state-coordinate inverses are pure testing. Restore the whole squares (RFN15), the retained $nv^2(N_0+pb)^2$ square, the scalar budget remainders, the $mpD_u^2N_0^2$ packet, and the pure testing factor D_u. This certifies the complete original parent (RFN4). Every bounded coefficient used in the argument depends only on the source and on u,v. The arbitrary state ratio z remains inside retained rows throughout. Hence the proof has no bounded-state restriction, and it includes literal $n=0$ without any limiting argument.

Audit boundary. The exact checker validates the complete parent, both full normals, all hard coefficients, the actual source cancellations, the scalar circuit, and the finite Gram/current budget identities. The source bounds and the signed scalar allocation are the separately stated RS input; the analytical argument above proves the full normal lift from that input. A separate state/sign proof and a separate outer angular join remain necessary for a whole-source all-testing claim. No explicit global LMI, optimized lift size, or external formal proof is claimed here.

Chapter 21

Reconstructed R2 source entry and full two-packet extraction

Component p667905169990
Source: REC/weak_c2/r2_source_entry_reconstructed.tex
SHA256: 6679051699909c37d8a3ec490c67de5b13db3076535d9ea4e35f4ad9c917db8f
Inventory use: RER1-RER8 | Actual original R2 source, all openings, unchanged coefficient formulas and RER1-7 bounds

21.1 Reconstructed R2 source entry and full two-packet extraction

Provenance and scope. This is a fresh mathematical reconstruction of the retained RE1–RE8 source-entry formulas. The original late-R21 file is absent. The identities are checked again against the unchanged native source formulas, and the boundedness proof is given in full. This lemma is structural: it supplies neither a scalar contact conductor nor an all-testing R2 certificate. In particular, restoring this lemma does not restore the missing thin-source, normal, or outer-chart proofs.

Use the original weak-C2 native source (a, B, t, r, h), with $0 < r, h < 1$, $a, B, t > 0$, the original strict source gaps, and $n \geq 0$. In R2, t is positive unbounded while

$$B, \qquad z = aB, \qquad y = tB$$

are positive infinitesimals. There is no bound on a and no lower bound on r, h, $1-h$, or $e = 1-r$. Put $l = 1-h$ and $S = 1/t$. Define the actual native arrays

$$c = 1 + zr, \quad s_c = 1 + B(l + ar), \quad R = 1 + hz, \quad g = R - ae > 0, \quad j_D = c - l/t > 0,$$

$$\begin{aligned} \Delta &= a + (1+z) + (1+z)B + 2z(1+z)r + a^3B^2r^2 + 2a^2Br \\ &\quad + tB\{1 + a + z + a^2Br(2-r)\}, \\ M &= (1+z)(1+ar)(h+t) - a(1+y)e\,l. \end{aligned}$$

The six weights used throughout this section are exactly

$$\begin{aligned} n &= \frac{(1+ar)cs_c - r\Delta}{2r\Delta}, & m &= \frac{a^2(1+y)ecs_c}{2g\Delta}, \\ C &= \frac{cM}{2l\Delta}, & A &= \frac{tc(1+ar)s_c}{2h\Delta}, \\ D &= \frac{(1+z)(1+ar)(1-atBr)s_c}{2tj_D\Delta}, & \alpha &= \frac{B^2(1+ar)M}{2(1-hB)\Delta}. \end{aligned} \tag{RER1}$$

All denominators are pure source. The original positivity assumptions are retained without adding a rate comparison.

Physicality controls the apparently unbounded source scale. The exact numerator identity is

$$(1+ar)(1+zr)\{1+B(l+ar)\}-r\Delta = eQ-BK, \tag{RER2}$$

where

$$K=(a+1)(h+t)(1+z),$$

$$Q=1+B+y+z(h+t+2r)+z^2\{t(1+er)+h(1+r)\}+Bz(h+t). \tag{RER3}$$

Every term of Q is nonnegative. Since $r\Delta>0$, original physicality is exactly $eQ\geq BK$, including equality when $n=0$. Put $\kappa=ay=tz$. Before assuming that κ is bounded, direct termwise inspection of (RER3), using B,y,z infinitesimal, gives $Q\leq C_0(1+\kappa)$ for a fixed positive real C_0. The strict gap $g>0$ gives $ae<R\sim 1$, so multiplication of the physicality inequality by a yields

$$\kappa(a+1)(1+hS)(1+z)\leq aeQ<RQ\leq C_1(1+\kappa) \tag{RER4}$$

for a fixed positive real C_1.

If a is bounded, $\kappa=ay$ is infinitesimal. If a is unbounded, move $C_1\kappa$ to the left in (RER4). Since $a+1-C_1\geq (a+1)/2$, one obtains $\kappa\leq 2C_1/(a+1)$, again infinitesimal. In either case, (RER4) also bounds $(a+1)\kappa$. Returning to (RER3) and then to $eQ\geq BK$ proves

$$ay\in \mathrm{I},\qquad a(a+1)y=O(1),\qquad Q\sim 1,\qquad \frac{(a+1)y}{e}=O(1). \tag{RER5}$$

No division by n occurred; the argument includes the face $n=0$.

Uniform source normalizations. Put

$$k=\frac{1+ar}{1+a},\qquad p=\frac{ae}{R}\in(0,1),\qquad \mu=m(1-p)=\frac{mg}{R},\qquad \delta_n=nr.$$

The quantity k is a positive bounded unit. Indeed $k\leq 1$; if $a\leq 1$, then $k\geq 1/(1+a)\geq 1/2$. If $a>1$, use $ae<R=1+hz$ to get $k>(a-hz)/(1+a)$, which has a positive real lower bound. Direct termwise division of the displayed polynomial Δ by $1+a$ gives $\Delta/(1+a)\sim 1$. Also

$$\frac{M}{t(1+ar)}\sim 1:$$

the first term has ratio $(1+z)(1+hS)\sim 1$, while the absolute ratio of the second is at most $R(1+y)S/(1+ar)$ and hence is infinitesimal. Since $c,s_c,j_D,1-hB\sim 1$, (RER1) now gives

$$\frac{SC\,l}{k}\sim\frac{1}{2},\quad \frac{SA\,h}{k}\sim\frac{1}{2},\quad \frac{D}{Sk}\sim\frac{1}{2},\quad \frac{\alpha}{Sy^2(1+a)k^2}\sim\frac{1}{2}. \tag{RER6}$$

For the D relation use $atB=ay=\kappa\in \mathrm{I}$ from (RER5). Thus all original h and l openings are present.

Two exact bounded products remove the remaining nonuniform source poles:

$$mg=\frac{a^2e(1+y)cs_c}{2\Delta},\qquad \delta_n=\frac{eQ-BK}{2\Delta}. \tag{RER7}$$

The quantities mg, mg/a, mg/a^2, μ, δ_n, and nr^2 are bounded. For example $a^2e/(1+a)\leq Ra/(1+a)=O(1)$, while $0\leq\delta_n\leq eQ/(2\Delta)=O(1)$. These estimates do not imply that m or n themselves are bounded.

Exact extraction from the entire parent. Let testing u, v, W, X, Y, Z be arbitrary, independently of the source; put $D_u = 1 + u^2$, $D_v = 1 + v^2$. The complete original quarter-parent is

$$\begin{aligned}\mathcal{P} = {} & m(vW - X + uvY - uZ)^2 + nD_uD_vX^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ & + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2.\end{aligned}$$

Define the mixed rows

$$q_a = v(W + uY), \qquad N = X + uZ - pq_a, \qquad \xi = X + ruZ.$$

Then the following identity retains both actual positive packets, without setting either normal to zero:

$$\mathcal{P} = mpN^2 + nD_uD_v\xi^2 + G, \tag{RER8}$$

where

$$\begin{aligned}G = {} & \mu\{N^2 - 2q_aN + (1 - p)q_a^2\} + \delta_nD_uD_v(-2uZX - ru^2Z^2) \\ & + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ & + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2.\end{aligned}$$

It follows simply by expanding $m\{(1 - p)q_a - N\}^2 - mpN^2$ and $n(X^2 - \xi^2) = nr(-2uZX - ru^2Z^2)$. Both retained packets are finite separated SOS: their source weights are nonnegative, D_uD_v is pure testing, and N, ξ are allowed mixed rows. When $n = 0$ the second packet vanishes literally. The exact identity and all estimates above are the reconstructed structural conclusion; no positivity conclusion about G is asserted.

Chapter 22

A simultaneous full-D/full-Y identity and an extra-row conductor

Component `pa73a17051258`
Source: `R17/definitions/c1_complete_joint_frame.tex`
SHA256: `a73a170512582634bf9285c08110515d2f736dda21eada108e1caa3db50c0032`
Inventory use: JF1–JF5, definitions, full-parent derivation | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

22.1 A simultaneous full-D/full-Y identity and an extra-row conductor

Scope and status. This is an independently reconstructed exact identity for the complete original C1 parent, not a certificate admitting C14.3b. It retains the same stationary source, $\Lambda = n = 0$, $\beta \geq 0$, all j and $1 - r$ openings, independent source/testing fields, and the remaining local v/u-infinitesimal testing chart. No mixed inverse or comparison of source and testing rates is used below.

Write $\mathcal{P} = F/4$, initially at $Z = 1$. Set

$$a = 1/\zeta, \quad t = \eta, \quad y = tx/a, \quad U = u^2, \quad V = v^2, \quad B = \beta + \tfrac{1}{2}, \quad D_p = D + \tfrac{1}{2},$$

$$E = tuW - v, \quad Q_t = Y - rv/t, \quad P = E/t - Q_t,$$
$$\xi = X + hu, \quad R_D = v(U - x), \quad T_D = R_D - uv\xi/h.$$

Thus $Q_t = rE/t + \nu$ in the R15 notation. Put

$$q = \alpha + (m + B)(1 - h)^2, \quad \mu = m - a^2 q, \quad \lambda = \mu/(tax), \quad H = avW - u - a\xi,$$

$$F_0 = V - y, \quad F_b = F_0 + vE, \quad F_t = F_0 + (t/r)vQ_t, \quad T_0 = A + \alpha + \tfrac{1}{2}, \quad T_m = T_0 + m, \quad e_4 = r^2 T_m/t^2.$$

Here $F_t = \widetilde{F}$ and $F_b = \overline{F}$. Define pure-source coefficients

$$b = Ar/t - m(1 - h) - \alpha - \tfrac{1}{2}, \qquad b' = b + e_4 yt/(2r),$$

$$\kappa = 2[B(1 - h) - D_p j/t - mr/t], \qquad \gamma = 2b'/h + 2D_p j/r - 2tB(1 - h)/r.$$

They satisfy $b' = O(x)$, $t\kappa \sim -1$ and $\gamma \sim 1$.

Complete parent, with no omitted hard row. Let

$$L = UV + (ax/t)V^2/U + tx^2/a - 3xV,$$

$$\mathscr{C} = \lambda L + \frac{\mu V}{t^2 U} E^2 + 2\left(\frac{\mu v^3}{t^2 U} - \lambda xv\right) E.$$

The exact full-parent identity is

$$\begin{aligned}\mathcal{P} &= qH^2 + \mathscr{C} + CP^2 + AQ_t^2 + D_p(P - hT_D)^2 + 2b'Q_tT_D \\ &\quad + \mathscr{R} + e_4UF_t^2 - e_4yu(H + a\xi)F_t \\ &\quad + AUQ_t^2 + B(1+V)\xi^2 - 2(\mu/a)H\xi - \mu\xi^2 \\ &\quad + \kappa uF_t\xi + \gamma uvQ_t\xi,\end{aligned} \tag{JF1}$$

where the entire joint radial row is

$$\mathscr{R} = \frac{C}{t^2}(F_b - rF_t)^2 + \frac{r^2T_0}{t^2}F_t^2 + \left(\frac{2mr}{t^2} + e_4x\right)F_bF_t. \tag{JF2}$$

In particular all $mVEQ_t/t$ terms are included in JF2; they are not separate errors. The $mUVQ_t^2$ contribution is included in the complete $e_4UF_t^2$ row, so it cannot also be retained independently.

A short independent way of certifying JF1 is useful. Without imposing any source formulas, the difference between the original parent and its displayed right side is exactly a linear combination of only

$$vE,\ vQ_t,\ u\xi,\ UV,\ V,\ 1.$$

The displayed expression has zero value and all five first derivatives at the original contact $U = x, V = y, E = Q_t = \xi = 0$. Its six coefficients therefore vanish by the six original stationary contact balances. The checker establishes this polynomial identity, including the value and every normal, before applying contact. This uses the complete original parent rather than a truncated model.

One strict radial pivot, with its exact companion. Set $\rho = m + rT_mx/2$. Then JF2 has the pure-source completion

$$\mathscr{R} = \frac{C}{t^2}[F_b - r(1 - \rho/C)F_t]^2 + \frac{r^2}{t^2}[T_0 + 2\rho - \rho^2/C]F_t^2. \tag{JF3}$$

Here $\rho = O(x)$ and $T_0 + 2\rho - \rho^2/C \sim 1/2$; all denominators are pure source and this remains uniform at both source openings. Both F_t/t and F_b/t have bounded allocations on JF3. This does not provide an independent strict F_0/t pivot. The non-small remaining coefficient is precisely $\gamma uvQ_t\xi$. In particular a source split making $x/\sqrt{t}$ bounded does not remove it, because $\gamma \sim 1$.

The extra circuit row participates in the leading cancellation. Use $c^2 = a/t$, $k^2 = x$, $p = ckv/u$, and homogenize now in Z:

$$J_0 = uv - x/c, \quad R_c = kv(tcW - Z), \quad R = \sqrt{\lambda}\,tk(cv - u)W, \quad \tau = tcW - Z.$$

The generic exact circuit completion is

$$\mathscr{C} = \lambda J_0^2Z^2 + R^2 + \underbrace{\frac{2\lambda k}{p}R_c^2}_{\mathcal{T}^2} - \lambda xE^2.$$

The full-D conductor is

$$R_D = uJ_0Z + kR_c - (x/c)(H + a\xi), \qquad H + a\xi = avW - uZ. \tag{JF4}$$

The additional exact polarization is

$$E(H + a\xi) = uv\tau^2 - t(u - cv)^2WZ. \tag{JF5}$$

Consequently the extra circuit row and the principal cross left after using JF4 satisfy

$$\mathcal{T}^2 + \kappa h(x/c)E(H + a\xi) = \frac{x}{c}(2\lambda + \kappa h)uv\tau^2 - \kappa ht\frac{x}{c}(u - cv)^2WZ. \tag{JF6}$$

This cancels the previously non-small E–angular Gram coupling. It is not itself an SOS decomposition: the independent native-source calculation in `../small_eta/check_circuit_remainder_source.py` gives $2\lambda + \kappa h < 0$ of order x/t in this chart. Also, the sign of WZ must be retained. The second summand is a literal SOS on the pure-testing sign chart $WZ \geq 0$ because $-\kappa > 0$, but not on the opposite sign chart.

An exact allocation of the negative small extra-row fraction. There is nevertheless a useful conductor:

$$x\frac{k}{p}R_c^2 = UR_c^2 - kuJ_0R_c\tau. \tag{JF7}$$

If the testing state is normalized with Z a unit and W bounded, then $\tau/Z = tc(W/Z) - 1$ is bounded. Thus the right side is an infinitesimal Gram on $\sqrt{\lambda}R_c$ and $\sqrt{\lambda}J_0Z$, with coefficients U and $ku(\tau/Z)$. Since $\sqrt{\lambda}R_c = R + \sqrt{\lambda}kE$, its additional coefficient on $E/\sqrt{t}$ is $\sqrt{\lambda t}\,k = O(k)$. Therefore a negative $O(x)$ fraction of $\mathcal{T}^2$ has a bounded infinitesimal allocation on this state chart. No bound on p is used. The companion identity

$$F_b/t = (W/Z)J_0Z + (x/a)\tau \tag{JF8}$$

may supply a strict xZ radial pivot in a simultaneous state-chart reassembly, since $\tau/Z \sim -1$ there. All divisions in JF7–JF8 are pure testing. Neither identity is an authorization to assume a strict $E/\sqrt{t}$ pivot after the leading source block has become semidefinite. That final simultaneous allocation is still required.

A direct conductor on the normalized Z-unit state chart. There is a simpler route on this state chart that does not perform JF4–JF6. Put $e_E = E/\sqrt{t}$ and retain the original strict source completion of the first line of JF1. Since $Q_t = E/t - P$ and $vZ = tuW - E$,

$$uvQ_t\xi = \sqrt{t}\,U(W/Z)e_E\xi - u(\xi/Z)e_E^2 - uvP\xi. \tag{JF9}$$

When the testing state is normalized with Z a unit, W/Z and X/Z are bounded and $\xi/Z = X/Z + hu$ is bounded. Thus all three coefficients in JF9 are infinitesimal in the bounded tensor ring. The last uses the bounded pure-source allocation of P on the completed source roots. The first line of JF1 has source Schur pivot $tS \sim 1/2$, and $E/\sqrt{t}$ has a bounded allocation there, including after the full D_p shift. Consequently JF9 removes the remaining γ coupling on this state chart without producing a semidefinite source frame. The checker verifies JF9 exactly. The accompanying complete state-chart admission, if used, must still list the retained positive rows and the strict angular block; JF9 alone does not cover states with normalized Z infinitesimal.

Receipts and audit conclusion. `check_joint_frame.py` has sixteen exact checks. They establish JF1 against the complete original parent through precisely its six stationary contact balances, JF3–JF8, and the exact surviving γ coupling. This resolves the double counting of the low and high full-Y radial rows and gives a new extra-row conductor. It does not establish the remaining strict frame, does not admit a new all-tests source subdomain, and does not close C14.3b.

Chapter 23

A uniform native C1 cap and an ordinary-angle admission

Component `p42ee07a067c1`
Source: `R17/definitions/c1_source_formulas.tex`
SHA256: `42ee07a067c1a1a2267d850dcf4d29bf0cb3d3e15dd090c8a588f757033e52c6`
Inventory use: NC1–NC7 | NC1–NC7; complete actual-unit/unit theorem | NC1-NC4

23.1 A uniform native C1 cap and an ordinary-angle admission

This section uses the original canonical C1 source, not an enlarged formal parameter family. Thus $\lambda = n = 0$, $d > 0$, $m, C, A, \alpha > 0$, $D, \beta \geq 0$ with at most one zero, and the source has its actual finite noncommuting stationary contact. Use the inherited global sign entry and normalize $C = 2$. The exact full-parent Alice reflection permits $0 < k \leq 1$. Put

$$x = k^2, \quad r = 1 - a\eta, \quad \theta = t\zeta, \quad h = 1 - \theta, \quad y = v_c^2 = x\eta\zeta,$$

$$K = 1+x,\ R = 1+xr,\ T = 1+xh,\ G = h+\zeta, \quad M = R-t,\ J = T-a,\ E = 1-x\eta h,\ Q = 1-x\zeta r. \tag{NC1}$$

The inherited exact signs give $0 < r, \theta < 1$, $E, Q, M, J > 0$ and $0 < a, t < 2$. There is no size assumption on η, ζ, e or a positive lower bound on any opening.

Three short positive source identities. The original coquotient polynomials obey

$$\begin{aligned} P_4 &= TE + x(1-r)G, \\ P_\alpha &= Rh(1+y) + \zeta ME, \\ P_d &= p_0 + \eta p_1, \qquad p_0 = 1 + 2xh + rx^2G + x\zeta, \quad p_1 = x\{K\zeta + xh\theta\}. \end{aligned} \tag{NC2}$$

These are obtained by substituting $a\eta = 1 - r$ and $t\zeta = \theta$ in the unchanged seven native coquotients. In particular $P_d \geq 1$. The original nonnegative D, β are precisely the inequalities

$$B_5 := K(1+y)(r+\eta) - (\eta T - 1 + r)P_d \geq 0, \qquad B_\beta := Kx(r+\eta)G - \theta P_d \geq 0. \tag{NC3}$$

The exact relative coefficients are

$$\begin{aligned} \frac{C}{d} &= \frac{K(r+\eta)E}{2(1-r)P_d}, & \frac{A}{d} &= \frac{\eta P_4}{2rP_d}, \\ \frac{m}{d} &= \frac{Kx(r+\eta)h(1+y)}{2M\zeta P_d}, & \frac{\alpha}{d} &= \frac{x^2\zeta(r+\eta)P_\alpha}{2QP_d}, \\ \frac{D}{d} &= \frac{B_5}{2J\eta P_d}, & \frac{\beta}{d} &= \frac{B_\beta}{2\theta P_d}. \end{aligned} \tag{NC4}$$

All denominators have their original positive signs; no denominator is silently declared a unit.

Lemma 23.1 (A uniform native gap)**.** *On every physical native C1 contact oriented by* $0 < k \le 1$*,*

$$E \ge \frac{1}{18}. \tag{NC5}$$

This includes the literal boundaries $D = 0$ *or* $\beta = 0$*.*

Proof. Define

$$\mathcal{A} = \theta T + K\zeta > 0, \qquad \mathcal{C} = RG(1 + rxh),$$

$$\begin{aligned}\mathcal{B} = {} & h^2 r x^2 + hrx(2-h) + h^2 x + hrx^2\zeta + hx\zeta + hx + h \\ & + rx\zeta + x\zeta + 2\zeta + 1.\end{aligned}$$

Direct multiplication of (NC2)–(NC3), using $x\eta h = 1 - E$, gives the polynomial identity

$$hB_5 = -\mathcal{A}E^2 + \mathcal{B}E - \mathcal{C}. \tag{NC6}$$

Every summand in $\mathcal{B}$ is nonnegative. Since $0 < x, r, h \le 1$ and $h(2-h) \le 1$, its six terms without ζ sum to at most 6 and the coefficients of ζ sum to at most 6. Hence $\mathcal{B} \le 6(1+\zeta)$. The native inequality $\theta = t\zeta < 2\zeta$ gives $1 < h + 2\zeta$, so $1 + \zeta < 3G$. Also $\mathcal{C} \ge G > 0$. From $B_5 \ge 0$ and (NC6),

$$G \le \mathcal{C} \le \mathcal{B}E \le 18GE.$$

Cancel the positive pure source G. This proves (NC5) without inverting an opening or assuming any of η, ζ bounded. □

An elementary division-free elimination. Keep p_0, p_1 from (NC2), and set

$$\begin{aligned} a_2 &= Kx\zeta - Tp_1, & a_1 &= K(1 + x\zeta r) - Tp_0 + (1-r)p_1, & a_0 &= Kr + (1-r)p_0, \\ b_1 &= KxG - \theta p_1, & b_0 &= KxrG - \theta p_0. \end{aligned}$$

Then $B_5 = a_2\eta^2 + a_1\eta + a_0$ and $B_\beta = b_1\eta + b_0$. Expanding these two expressions, or just eliminating their quadratic and linear terms successively, gives

$$\begin{aligned} & b_1^2 B_5 - a_2 B_\beta^2 - (b_1 a_1 - 2a_2 b_0) B_\beta \\ & \quad = b_1^2 a_0 - b_1 a_1 b_0 + a_2 b_0^2 \\ & \quad = x^2 h^2 K^2 R G^2 \{1 + 2xh + rx^2 h^2 + (1-r)x\zeta\}. \end{aligned} \tag{NC7}$$

The last equality is a short polynomial identity in x, r, h, ζ; all five of its input coefficients are printed above. No inverse of b_1, no resultant premise, and no localization is used. For reference, $b_1 = xh\{K(1+\zeta) - x\theta^2\} > 0$ as well.

Theorem 23.2 (Actual unit/unit C1 contacts enter the five-event exit)**.** *If both actual C1 half-angles* k, v_c *are positive bounded units, the original source has five consecutive events bounded below relative to* d*. Consequently its entire original parent has the inherited all-tests tensor-SOS certificate. All upper event scales and the retained zero-coefficient boundaries are allowed.*

Proof. After the exact Alice orientation, x, y are bounded positive units. Thus $\eta\zeta = y/x$ is a bounded positive unit. If η were unbounded, ζ would be infinitesimal. Then $\theta = t\zeta$ would be infinitesimal, contradicting $x\eta(1-\theta) < 1$. If ζ were unbounded, η would be infinitesimal and $r = 1 - a\eta$ would have residue one, contradicting $x\zeta r < 1$. Therefore η, ζ are both bounded positive units.

All polynomials in (NC2)–(NC3) now have bounded coefficients and bounded arguments. If h were infinitesimal, the exact specialization

$$B_\beta\big|_{\theta=1} = -1 - x\zeta(1-r) \le -1$$

would contradict $B_\beta \geq 0$. Thus h is a positive bounded unit. The uniform cap gives that E is a unit. Equations (NC2) make P_4, P_α, P_d positive bounded units: use $P_4 \geq E$, $P_\alpha \geq Rh(1+y)$, and $P_d \geq 1$.

The first four equations of (NC4) now show that C, A, m, α are all strong relative to d. Their denominators are positive bounded quantities; they may be infinitesimal. Suppose both D/d and β/d were infinitesimal, including a possible literal zero. Their last two bounded denominator products in (NC4) would make B_5, B_β infinitesimal. Every coefficient on the left of (NC7) is bounded, so that side would be infinitesimal. Its right side is a positive unit, since x, h, K, R, G are units and the last bracket is at least one. This is a contradiction.

Thus D or β is strong. The original cycle $(\lambda, D, C, A, \alpha, m, \beta, n)$ has respectively the five consecutive strong events (D, C, A, α, m) or (C, A, α, m, β). Apply the inherited five-consecutive-event theorem, whose stated scope allows arbitrary upper weights and every independent testing chart. No independent test is identified with an actual source angle in this source-entry argument. □

Exact intersection with the revision-3 residual rows. The new all-tests theorem closes the actual (U, U) subcase of **C1.5**. It does not close the whole C1.5 row. It has no intersection with C1.1 or C1.4, where k is infinitesimal. In C1.1, the inherited $q = x\eta \leq 1/2 + \mathrm{Inf}$ bound and ζ infinitesimal already force $v_c^2 = q\zeta$ infinitesimal. The uniform cap (NC5), unlike the new all-tests consequence, applies to every original C1.1, C1.4 and C1.5 source. All statements may be intersected with positive infinitesimal δ_{tot} and the complement of every earlier admission without altering their proofs. No remaining typed intersection is asserted nonempty.

Verification scope. The accompanying checker reconstructs P_4, P_α, P_d from the original native formulas and checks (NC2), (NC6), (NC7), all six ratios in (NC4), and the $\theta = 1$ specialization exactly. It corroborates these sixteen short identities. The ordered-field bounds and invocation of the full-parent five-event theorem are the mathematical proof above; a checker PASS alone is not an all-tests admission. The root's separate scoped mathematical review found no blocking gap.

Chapter 24

A five-root local certificate on an A-unbounded weak-C2 subchart

Component `p9ec3d6bbd717`
Source: `R17/definitions/weak_c2_complete_WA_frame.tex`
SHA256: `9ec3d6bbd7170be3f5d3cc22ae2d244ff93c9445f41550c5c9369804dab0bbcd`
Inventory use: WA3-WA8 and WA10

24.1 A five-root local certificate on an A-unbounded weak-C2 subchart

Revision 10 scope update. The inherited boundary extension in `inherited_r16/Q222_Revision13/advances/weak_A_boundary/extension.tex` extends this admission to z/B positive infinitesimal with all other source hypotheses retained. The unit-ratio restriction below records the original revision-9 theorem.

Theorem 24.1. *Retain the original physical C2 source, its exact stationary contact, all strict source gaps, and $n \geq 0$. Normalize $d = 1$. Suppose B, z, y are positive infinitesimals, $r, G/B, j, y/B, L/B$ are positive bounded units, and H/y is infinitesimal. Impose in addition*

$$z/B \text{ and } 1 - r \text{ are positive bounded units.} \tag{WA1}$$

Then the entire original parent is a finite tensor sum of squares for arbitrary independent testing half-angles u, v that are infinitesimal or zero, including either sign and complex states. There is no comparison of a testing scale with B, H, z, y. The total-scale parameter δ_{tot} is positive infinitesimal. This is a local testing certificate on the stated source subchart; it does not assert that its intersection with the final residual complement is nonempty, or supply an unproved remote-testing join.

No membership in the residual complement is needed in this proof. When applied to the original residual problem, its existing complement condition is still intersected with the source chart; no nonemptiness of that intersection is inferred.

The original omitted event coefficients remain $\lambda = \beta = 0$. Here $B = H + L$, and the actual contact in the positive gauge is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(\sqrt{z}, \sqrt{y}; \frac{B}{\sqrt{zy}}, -r\sqrt{z}, \frac{H}{\sqrt{y}}, 1\right).$$

The source ratio $\ell = B/z$ below is not an event coefficient.

The source pivots survive the small H opening. Write $h = H/B$, $\kappa = B/y$, $\ell = B/z$. Both κ and ℓ are positive bounded units, and h is infinitesimal. The original formulas imply that n, m, C, D, α are bounded, that m, C, D are positive bounded units, and that Ah is a positive bounded unit. Thus A is unbounded and $Ah^2, \sqrt{A}\,h, A^{-1/2}$ are infinitesimal. No bound on the relative rates of h and B is used.

Here is an explicit check of the radial pivot. With $T_c = B + zr$, the original formula is

$$\frac{\alpha}{B^2} = \frac{T_c}{\delta}\frac{M/B^2}{2(1-H)}, \qquad M - yG = H(1+z)T_c + z\{yj + rL(1+y)\} > 0. \tag{WA2}$$

Consequently $M/B^2 \geq (y/B)(G/B)$ is bounded below by a positive real, and it is bounded above by the original formula for M and the unit ratios in (WA1). Since T_c/δ is a positive bounded unit, α/B^2 is a positive bounded unit uniformly as $H/y \to 0$. Likewise

$$m = \frac{z^2(1+y)(1-r)(1+zr)(1+zr+L)}{2G\delta}$$

is a positive bounded unit under (WA1); the local residue argument uses only $n \geq 0$. Finally A is the only unbounded retained coefficient, so $\delta_{\mathrm{tot}} = 1/(1+n+m+C+A+D+\alpha)$ is positive infinitesimal. These arguments preserve all original coefficient and source-gap hypotheses.

Five homogeneous contact roots. Put $U = u^2$, $V = v^2$, and define

$$\xi = X + ruZ, \qquad q = \sqrt{A}\,(Y - huW), \qquad R_1 = uW - \kappa vZ, \quad R_2 = vW - \ell uZ, \quad T = uvW - BZ. \tag{WA3}$$

Every root vanishes at the actual source contact. Inverses and radicals in their coefficients are pure source. The large A packet is retained in q; no positive packet is subtracted from the parent and no physicality hypothesis is transferred to a modified remainder.

Use the following source coefficients for the exact scalar graph:

$$\begin{aligned}
a &= (C + D + \tfrac{1}{2})(1-h)^2 + Ah^2, & a_4 &= Ah^2,\\
b_2 &= 2mh + C(1-h)^2 + (A + \alpha + \tfrac{1}{2})h^2, & b_4 &= h^2(m + A + \alpha + \tfrac{1}{2}),\\
c &= \alpha, & c_2 &= nr^2 + (m + \tfrac{1}{2})(1-r)^2 + \alpha,\\
c_4 &= nr^2, & d_2 &= (n + \tfrac{1}{2})r^2 + 2Dr,\\
d_4 &= (n + D + \tfrac{1}{2})r^2.
\end{aligned}$$

Thus the graph on $\xi = q = 0$ is

$$\begin{aligned}
F_g/4 = {} & [aU + a_4U^2 + (m + b_2U + b_4U^2)V]W^2\\
& + [c + c_2U + c_4U^2 + (D + d_2U + d_4U^2)V]Z^2 - 2uv(e_0 + e_1U)WZ,
\end{aligned}$$

where e_0, e_1 are exactly the original coefficients in (WS3). Set

$$\begin{gathered}
\beta_s = \frac{Bd_4 + \kappa c_4}{\kappa\ell^2}, \qquad \gamma_s = b_4 + \frac{d_4}{\kappa\ell},\\
j_0 = \frac{Bd_4 - a_4\kappa^2\ell + d_2\ell}{2\kappa\ell^2}, \qquad j_1 = \frac{d_4}{2\kappa\ell}, \qquad J_s(U) = j_0 + j_1U,\\
f_s = \frac{1}{2}\left(b_2 - \frac{c_4}{\ell^2} - \frac{c}{B^2} - \frac{Bd_4}{\kappa\ell^2}\right).
\end{gathered} \tag{WA4}$$

The four original scalar contact equations (value and the three logarithmic derivatives in $u, v, W/Z$) give the exact identity

$$F_g/4 = (R_1, R_2, T)\,\mathsf{S}\,(R_1, R_2, T)^T, \qquad \mathsf{S} = \begin{pmatrix} a + a_4U & uvJ_s & -vJ_s\\ uvJ_s & m + \beta_sU & u(f_s - j_1U)\\ -vJ_s & u(f_s - j_1U) & c/B^2 + \gamma_sU \end{pmatrix}. \tag{WA5}$$

This is a different scalar identity from the excluded polynomial two-block splitter (JG7). In particular, the last diagonal retains the full radial mass α/B^2 and the Gram depends on the independent testing variables. All source coefficients in (WA4) are bounded; no uncancelled $1/B$ occurs.

Both full first normals and the complete hard block. For the ξ normal define

$$\begin{gathered} p_0 = \frac{D + r(n+1/2)}{\kappa}, \qquad p_1 = \frac{r(n+D+1/2)}{\kappa}, \qquad q_1 = \frac{nr + Bp_1}{\ell}, \\ f_x = -mh + (D+\tfrac{1}{2})(1-h) - p_0 - q_1, \\ \mathsf{L}_x = \big(uv(p_0 + p_1 U),\ -m + q_1 U,\ u(f_x - p_1 U)\big). \end{gathered} \tag{WA6}$$

For the unnormalized $\eta = Y - huW$ normal set

$$\begin{gathered} p_y = -(C + D + \tfrac{1}{2})(1-h) + Ah, \qquad p'_y = Ah, \\ k_v = -C(1-h) + Ah + m + (\alpha + \tfrac{1}{2})h, \qquad k_{uv} = h(A + m + \alpha + \tfrac{1}{2}), \\ k_z = -m(1-r) + r(D + \tfrac{1}{2}) - \alpha - \tfrac{1}{2}, \\ q_y = \frac{-k_z - \kappa p'_y - Bk_{uv}}{\ell}, \qquad f_y = k_v - q_y, \\ \mathsf{L}_y = (p_y + p'_y U,\ uvq_y,\ v(f_y + k_{uv} U)). \end{gathered} \tag{WA7}$$

The two remaining original source stationarity equations, in X, Y, give these exact representations of the complete first normals. No first-normal term is discarded. Define also

$$\begin{aligned} h_{xx} &= n(1+U)(1+V) + m + (D+\tfrac{1}{2})UV + \tfrac{1}{2}(1+V), \\ h_{xy} &= -uv(m + D + \tfrac{1}{2}), \\ h_{yy} &= A(1+U)(1+V) + C(1+V) + D + \tfrac{1}{2} + mUV + (\alpha + \tfrac{1}{2})(1+U)V. \end{aligned}$$

Then the *entire original four-state parent* is exactly

$$\begin{gathered} F/4 = \mathcal{R}^T \mathsf{G}(u,v)\mathcal{R}, \qquad \mathcal{R} = (\xi, q, R_1, R_2, T)^T, \\ \mathsf{G} = \begin{pmatrix} h_{xx} & h_{xy}/\sqrt{A} & \mathsf{L}_x \\ h_{xy}/\sqrt{A} & h_{yy}/A & \mathsf{L}_y/\sqrt{A} \\ \mathsf{L}_x^T & \mathsf{L}_y^T/\sqrt{A} & \mathsf{S} \end{pmatrix}. \end{gathered} \tag{WA8}$$

Block notation in (WA8) has sizes $1, 1, 3$. Every coefficient of this finite polynomial matrix in u, v is bounded and pure source. The checker first expands the original signed core and all six original event packets to verify the complete hard block and both normal rows. It then verifies (WA5)–(WA7) from the corresponding four plus two contact equations. Thus (WA8) is not merely a graph or Hessian certificate.

Strict residue and a legal separated Gram lift. For infinitesimal independent u, v, the residue of (WA8), in the printed root order, is

$$\overline{\mathsf{G}} = \begin{pmatrix} \bar{n} + \bar{m} + 1/2 & 0 & 0 & -\bar{m} & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & \bar{C} + \bar{D} + 1/2 & 0 & 0 \\ -\bar{m} & 0 & 0 & \bar{m} & 0 \\ 0 & 0 & 0 & 0 & \overline{\alpha/B^2} \end{pmatrix} \succ 0. \tag{WA9}$$

Indeed the nontrivial 2×2 determinant is $\bar{m}(\bar{n} + 1/2) > 0$, and (WA2) supplies the last positive pivot. The q couplings disappear because $A^{-1/2}$ is infinitesimal while Ah is bounded. Every other omitted entry has a factor u, v, or uv.

For completeness, strict bounded residue here licenses a finite tensor SOS without a mixed Cholesky or inverse. Let $\mathsf{G}_0 = \mathsf{G}(0,0)$, a positive definite pure-source matrix whose inverse is bounded. Whiten by its pure-source square root and write the resulting matrix as

$$I + \sum_{j=1}^{N} H_j t_j,$$

where each H_j is a bounded pure-source symmetric matrix and each t_j is a nonconstant pure-testing monomial in u, v, hence infinitesimal. Split I into N equal copies and orthogonally diagonalize each H_j purely in the source field. A resulting scalar multiplier is $1/N + \ell t_j$, with ℓ bounded. Choose a positive real $R > N|\ell|$ and put $a = N\ell/R$, $b = Rt_j$. Then $|a| < 1$, $|b| < 1$, and

$$\frac{1}{N} + \ell t_j = \frac{1}{2N}\{(1+a)(1+b) + (1-a)(1-b)\}. \tag{WA10}$$

Each product is the square of a pure-source square root times a pure-testing square root. This supplies finitely many legal tensor roots for (WA8). Only finitely many source eigenvalues occur, so one real R can be chosen for all of them. If $N = 0$ the source Gram itself is already sufficient.

This construction works for either sign and for zero u or v, and permits all relative source/testing rates. Applying the real identity separately to the real and imaginary state parts proves the complex-state assertion. It uses pure-testing square roots of $1 \pm Rt_j$, and consequently makes no claim to a global polynomial SOS in the testing variables. There is therefore no conflict with the obstruction proved on the physical SO1 family.

Exact scope of the gain. The result closes the local both-infinitesimal testing problem on (WA1), with all four original state coordinates retained and with no centered-packet subtraction. It does not yet cover the allowed source openings z/B infinitesimal or unbounded, or $1 - r$ infinitesimal: in the latter case the R_2 pivot in (WA9) can lose residue. A remote-testing theorem must still be joined before this local certificate is called an all-tests source admission. No assertion of nonemptiness of the final residual intersection is made. The accompanying checker verifies thirteen exact identities; the source bounds and finite lift above are the positivity proof.

Chapter 25

R9 source data and the finite coefficient calculation

Component `pd709027c7805`
Source: `R17/definitions/weak_c2_source_formulas.tex`
SHA256: `d709027c7805a97b112dd559ec1ac3a749c903cad55d59d6be0fab43ebd5227d`
Inventory use: complete original source coefficients; scalar and all first-normal arrays | complete original formulas and all first-normal arrays

25.1 R9 source data and the finite coefficient calculation

This appendix fixes all coefficient arrays used in the local theorem. It can be read with the original physical-source entry in the supplied R13 frontier audit. Every displayed formula is source-only. Put $r = 1 - e$, $z = aB$, $y = tB$, and

$$c_a = 1 + aBr, \quad S_c = 1 + B(1 - h + ar), \quad g = 1 + a(hB + r - 1), \quad j = c_a - (1 - h)/t,$$

$$\begin{aligned}\Delta &= a + (1 + aB) + (1 + aB)B + 2aB(1 + aB)r + a^3B^2r^2 + 2a^2Br \\ &\quad + tB\{1 + a + aB + a^2Br(2 - r)\}, \\ M &= (1 + aB)(1 + ar)(h + t) - a(1 + tB)e(1 - h).\end{aligned}$$

The physical stationary event coefficients are

$$\begin{aligned} n &= \frac{(1 + ar)c_aS_c - r\Delta}{2r\Delta}, & m &= \frac{a^2(1 + tB)ec_aS_c}{2g\Delta}, \\ C &= \frac{c_aM}{2(1 - h)\Delta}, & Ah &= \frac{tc_a(1 + ar)S_c}{2\Delta}, \\ D &= \frac{(1 + aB)(1 + ar)(1 - atBr)S_c}{2tj\Delta}, & \alpha &= \frac{B^2(1 + ar)M}{2(1 - hB)\Delta}. \end{aligned}$$

On R9, B, a, h, e are positive infinitesimals and t, g, j are positive units. In particular $t_0 > 1$. The exact physical constraint $n \geq 0$ is retained, including equality. The unit common denominator

$$\mathcal{D} = 2rg(1 - h)(1 - hB)(t - 1 + h + aBtr)\Delta$$

has residue $2(t_0 - 1) > 0$.

Scalar arrays. The notation Ah in these formulas means the product of A and h.

$$\begin{aligned}
q_s &= nr^2 + (m+\tfrac{1}{2})e^2, \qquad q = q_s + \alpha,\\
a_w &= (C+D+\tfrac{1}{2})(1-h)^2 + Ah\,h,\\
b_2 &= 2mh + C(1-h)^2 + Ah\,h + (\alpha+\tfrac{1}{2})h^2,\\
b_4 &= h^2(m+\alpha+\tfrac{1}{2}) + Ah\,h,\\
d_2 &= (n+\tfrac{1}{2})r^2 + 2Dr, \qquad d_4 = (n+D+\tfrac{1}{2})r^2,\\
e_0 &= me + D(1-h) + \alpha h + \tfrac{1}{2},\\
e_1 &= mhe + r(D+\tfrac{1}{2})(1-h) + (\alpha+\tfrac{1}{2})h,\\
c_F &= (b_2 - a^2nr^2)/t^2, \qquad \lambda = (m-a^2q)/(ta^2B),\\
f_{A0} &= Ah\,h/t^2 + d_2 - 2e_1/t + 2anr^2/t,\\
f_{C0} &= (\alpha - c_Ft^2B^2)/(taB^2),\\
f'_G &= -\{a_w/t^2 + D - 2e_0/t + 2aq/t + 2c_FtB\}/(aB).
\end{aligned}$$

The potentially singular a, B factors cancel in the combinations used in RC1. One direct proof is to clear $\mathcal{D}$ and the printed monomials and expand; all three resulting numerator polynomials are zero. Equivalently, the original value and two first derivatives at contact give precisely RC1. Thus RC1 is an exact identity, including arbitrarily small a and literal $n = 0$.

The needed size statements also follow by finite monic division. The R14 regularity calculation gives $f_B^{\mathrm{old}} = (m - a^2q_s)/(ta^2B) \sim 1/2$ jointly at $B = a = h = 0, r = 1$. Since

$$\lambda = f_B^{\mathrm{old}} - \frac{\alpha}{tB}, \qquad \frac{\alpha}{tB} = \frac{B}{t}\frac{\alpha}{B^2}, \qquad \frac{\alpha}{B^2} \sim t_0/2,$$

we obtain $\lambda \sim 1/2 > 0$ without any division by n or e. Likewise $c_F \sim 1/(2t_0)$, $d_4 \sim t_0/[2(t_0-1)]$, $b_4 \sim 0$. Direct substitution gives $C \sim Ah \sim t_0/2$, $D \sim 1/[2(t_0-1)]$ and $m, n, \alpha \sim 0$. These are positive-unit denominators and polynomial cancellations, rather than asymptotic replacements in the proof identity.

All first-normal arrays. For the angular weight $q = q_s + \alpha$ define

$$\begin{aligned}
K'_X &= -nr + (m+\tfrac{1}{2})e - aq,\\
K'_E &= a_w/t^2 - e_0/t + aq/t,\\
K_Y &= \{-(C+D+\tfrac{1}{2})(1-h) + Ah\}/t + D - \alpha,\\
x_3 &= 2\{(D+\tfrac{1}{2})[(1-h)/t - 1] + e/2 - nr(1+a/t) - mh/t\},\\
x_5 &= -2r(n+D+\tfrac{1}{2}),\\
a_E &= 2Ah\,h/t^2 - 2e_1/t, \qquad b_E = 2b_2/t^2, \qquad c_E = 2b_4/t^2,\\
a_Y &= 2\{Ah/t - me + r(D+\tfrac{1}{2}) - \alpha - \tfrac{1}{2}\},\\
b_Y &= 2\{-C(1-h) + Ah + m + (\alpha+\tfrac{1}{2})h\}/t,\\
c_Y &= 2\{Ah + h(m+\alpha+\tfrac{1}{2})\}/t.
\end{aligned}$$

Every array is bounded on R9; the only nonzero cross-term residue used in the final Gram is $a_E \to -1/(t_0-1)$. Direct differentiation of the original parent after the printed state substitution gives RC7–RC9. Their source contact balances RC6 follow by the same common-denominator calculation as RC1. These full-parent identities are also checked by reconstructing the parent independently in the short audit script.

Inherited input needed for the all-tests join. The R13 frontier audit explicitly extends the remote certificate to R9 with a infinitesimal. Its exact homogeneous endpoint identities cover the complement of both angles infinitesimal. The new local roots have no testing inverse and include

the zero axes directly; the original two sign gauges are retained. No R14 testing-ratio split is needed by the new local certificate. The provenance map names the unchanged remote, frontier, source and bounded-Gram modules used here. The complete global routing proof still depends on the unresolved source branches in the checklist.

Chapter 26

An exact beta-shared normalization for the two eta-large C1.4 charts

Component `p10ee3d96a3f1`
Source: `R13/advances/c1_eta_large_source/beta_shared_source_normalization.tex`
SHA256: `10ee3d96a3f15b4d3af3bd76a66d0544a866543c5f2b2d78e4f1b03c03b55b09`
Inventory use: BE1-BE9 original large-L source bounds | BE1–BE10 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

26.1 An exact beta-shared normalization for the two eta-large C1.4 charts

This section proves an exact source reduction, not an all-tests admission. It applies simultaneously to C14.1 and C14.2: the original native C1 source has $x = k^2$ positive infinitesimal, $L = x\zeta$ bounded below by a positive real, and η unbounded. All original physicality and residual-complement conditions are retained. We normalize $d = 1$. The conclusion includes every positive source opening and the literal faces $D = 0$ and $\beta = 0$. No testing variable occurs in the source normalization.

A uniform strengthening of the source gap. Write $\sigma = \eta^{-1}$, $\theta = 1 - h$, $K = 1 + x$, and $e = xh/\sigma = 1 - E$. The exact coefficients in (LH1), with no size specialization, give

$$B_5 = -xh\{KL + x\theta(1 + xh)\}\eta^2 + xb_1\eta + a_0,$$

where

$$\begin{aligned} b_1 = {} & -Lhrx - Lh - Lr + L - h^2rx^2 + h^2rx - 3h^2x \\ & - 2hrx + hx - 3h + 1, \\ a_0 = {} & Kr + (1 - r)\{1 + L + 2xh + rxL + rx^2h\}. \end{aligned}$$

Since $0 < r, h < 1$, $Lr < 1$, and L is bounded below, b_1/L and a_0/L are bounded uniformly in all openings; the second is positive. Dividing the inequality $B_5 \geq 0$ by $L\eta > 0$ gives

$$e\{K + x\theta(1 + xh)/L\} \leq x(b_1/L) + \sigma(a_0/L) \leq O(x) + O(\sigma). \tag{BE1}$$

The bracket is at least one. Therefore $e \in \mathrm{Inf}_{>0}$ and $E \sim 1$ on both eta-large charts. This proof does not assume x/σ bounded, and does not infer that xh/σ^2 is bounded.

Two exact identities and their bounded coefficients. Set

$$z = L^{-1}, \quad b = Lr, \quad Q = 1 - b, \quad R = 1 + xr, \quad M = R - xz\theta, \quad J = 1 + xh - \sigma(1 - r).$$

Here $z > 0$ is bounded, $0 < b, r < 1$, and $r = bz$. The opening-cleared denominator is

$$P = \sigma z P_d = \sigma + K + \sigma z(1 + bx + 2xh) + \sigma b x^2 h z^2 + x^2 h\theta z.$$

In particular $P, M, J \sim 1$. Introduce

$$H_0 = 1 + z - r = 1 + zQ, \qquad B = P + xz(J - \sigma)$$

and

$$\mathcal{H} = RH_0 + z\left[(1-r)xh(2+xr) - e\{2 + x(1 + h + r\theta)\} + Ke^2\right]. \tag{BE2}$$

Then $B \sim 1$, and $\mathcal{H} - H_0 \in \mathrm{Inf}$, while $H_0 \geq 1$ is bounded. Thus B and $\mathcal{H}$ are positive bounded units. The original NC coefficients satisfy exactly

$$2\theta P\beta = hB - \sigma H_0, \tag{BE3}$$

$$2P(JD + x\theta\beta) = \sigma^2\mathcal{H}. \tag{BE4}$$

To verify these without any cancellation of an opening, use

$$N_D = K(1+\sigma z)(1+\sigma r) - JP = 2JPD, \quad N_\beta = K(1+\sigma r)(1+xhz) - \theta P = 2\theta P\beta.$$

Expansion gives $N_\beta = hB - \sigma H_0$. The identity $N_D + xN_\beta = \sigma^2\mathcal{H}$ follows on inserting $\sigma e = xh$. These are polynomial identities after multiplication by σ^2; no resultant, opening inverse, or approximation is used. As an immediate consequence,

$$D = O(\sigma^2), \qquad x\theta\beta = O(\sigma^2), \qquad JD + x\theta\beta \asymp \sigma^2. \tag{BE5}$$

In particular the collapsing D pivot cannot be given a fixed positive unit cap in this source chart. The estimate is stronger than bounding D by $O(\sigma^2 + x\sigma)$ before using original beta nonnegativity.

A source sphere retaining both coefficient boundaries. All radicals below are pure source. Define

$$d_0 = \sqrt{\frac{2PJD}{\sigma^2\mathcal{H}}}, \qquad b_0 = \sqrt{\frac{2Px\theta\beta}{\sigma^2\mathcal{H}}}.$$

Then exactly

$$d_0, b_0 \geq 0, \quad d_0^2 + b_0^2 = 1, \quad D = \frac{\sigma^2\mathcal{H}}{2PJ}d_0^2, \quad \beta = \frac{\sigma^2\mathcal{H}}{2Px\theta}b_0^2. \tag{BE6}$$

No inverse of D, β, d_0, or b_0 occurs. In particular $D = 0$ corresponds to $(d_0, b_0) = (0, 1)$, and $\beta = 0$ to $(1, 0)$. Equation (BE3) now becomes the exact positive decomposition

$$\frac{h}{\sigma} = \frac{H_0}{B} + \frac{\sigma}{x}\frac{\mathcal{H}}{B}b_0^2. \tag{BE7}$$

It displays, rather than discards, the possibly unbounded factor h/σ. The latter is bounded below by a positive unit, but may be unbounded when σ/x is unbounded.

Put

$$\kappa = \frac{K(1+\sigma r)(1+\sigma z)}{2MP} \sim \frac{1}{2}, \qquad m_0 = \kappa x^2 z\frac{H_0}{B}, \qquad m_1 = \kappa x\sigma z\frac{\mathcal{H}}{B}.$$

The original m coefficient has the exact split

$$m = m_0 + m_1 b_0^2, \qquad m_0 \asymp x^2 z, \quad m_1 \asymp x\sigma z. \tag{BE8}$$

Thus the same nonnegative source coordinate b_0 occurs in the beta packet and in the extra part of m. This is an exact beta-aware replacement for an assumed bounded $\mu = h/\sigma$. All factors hidden in the two comparisons in (BE8) are bounded positive units uniformly in the original openings. For the other three positive coefficients the opening-cleared products are also units:

$$\frac{C(1-r)}{z} \sim \frac{1}{2}, \qquad Ar \sim \frac{H_0}{2}, \qquad \alpha Qz \sim \frac{1}{2}. \tag{BE9}$$

Indeed their exact expressions are those in the original large-L source chart, which are algebraic for bounded z as well; use (BE1). For example $Ar = [z(1+xh)E + (1-r)(1+xhz)]/(2P)$. No lower bound on r, $1-r$, Q, or θ was imposed.

Exact full-parent preservation. For arbitrary independent reciprocal testing coordinates u, w, put $D_u = 1 + u^2$, $D_w = 1 + w^2$ and

$$\begin{aligned} M_w &= W - wX + uY - uwZ, \quad R_w = uW - Y, \\ J_w &= uwW + uX - wY - Z, \quad B_w = X + uZ, \\ K_w &= J_w^2 + D_u(Y - wZ)^2 + D_w B_w^2 - D_u D_w Z^2. \end{aligned}$$

The entire original reciprocal parent is exactly

$$\begin{aligned} F_w = {} & 4m_0 M_w^2 + 4m_1(b_0 M_w)^2 + 4CD_w R_w^2 + 4AD_u D_w Y^2 \\ & + \frac{2\sigma^2\mathcal{H}}{PJ}(d_0 J_w)^2 + 4\alpha D_u(Y - wZ)^2 + \frac{2\sigma^2\mathcal{H}}{Px\theta} D_w(b_0 B_w)^2 + 2K_w. \end{aligned} \tag{BE10}$$

This is an equality in the complete four-state quadratic form. It therefore preserves every first state normal and every hard quadratic coupling; it is not a restriction to a contact line or a leading form. Every new coefficient or radical is pure source, and all D_u, D_w factors expand as finite testing sums of squares. The identity is polynomial in u, w, W, X, Y, Z and therefore includes all signs and every independent testing rate. It holds at literal $u = 0$ and $w = 0$. Bihomogenizing u and w gives the corresponding identity at their other literal projective endpoints, without division by a testing coordinate. Applying the real identity to real and imaginary state parts gives the complex-state identity as well.

The source sphere, (BE7), and (BE8) do not make $2K_w$ nonnegative. They leave a precise full-parent allocation problem: use the common b_0 packet in (BE10) together with the d_0 packet, uniformly through $d_0 = 0$ and $b_0 = 0$. Consequently neither C14.1 nor C14.2 is admitted by this section. The sigma-unit six-root proof's positive-unit D pivot cannot be imported unchanged. No claim of nonemptiness of any residual-complement source domain is made.

Verification scope. The accompanying narrow checker verifies (BE1)'s polynomial input, (BE3), (BE4) in both reciprocal and native notation, the exact m split, the opening-cleared coefficient products, and (BE10) as the complete quadratic polynomial, including its bihomogenization. The asymptotic deductions use the ordered-field inequalities above; checker success alone is not an all-tests certificate.

Targeted full-normal extension and the exact null direction. Here is a test of the proposed normalization on the normal block, with no source opening omitted. At $w = 0$, put $X = N_x - uhZ$ and $t = u^2$. The original parent splits into its manifestly nonnegative (W, Y) block and the following complete (N_x, Z) block:

$$\begin{aligned} Q_{xz} = {} & [4Dt + 4\beta + 2(1 + t)]N_x^2 \\ & + 4u[-2D(1 + th) + 2\beta\theta - h(1 + t)]N_x Z \\ & + [4D(1 + th)^2 + 4\beta t\theta^2 + 2t(1 + t)h^2]Z^2. \end{aligned} \tag{BE11}$$

At the literal endpoint $u = w = 0$ this is $(4\beta + 2)X^2 + 4DZ^2$. Thus when $D = 0$, the vector $(W, X, Y, Z) = (0, 0, 0, 1)$ is an *exact* null direction of the full parent. The old six-root argument cannot impose a strictly positive pivot on σN_z at this face. This is an obstruction to that proof architecture, not to positivity; $w = 0$ itself already has an elementary SOS.

The sphere suggests the legal replacement roots

$$\Gamma = \sigma Z, \qquad Z_d = d_0\Gamma, \qquad Z_b = b_0(u/k)\Gamma, \qquad Z_e = \sqrt{e}\,(u/k)\Gamma.$$

Set $p = \sigma b_0/k$. Equation (BE7) implies p bounded, since $hB = \sigma H_0 + \mathcal{H}p^2$. Also

$$\frac{xh^2}{\sigma^2} = \frac{eH_0}{B} + \frac{\mathcal{H}h}{B} b_0^2.$$

Consequently the entire Z^2 line of (BE11) equals

$$\frac{2\mathcal{H}}{PJ}(1+th)^2 Z_d^2 + 2\mathcal{H}\left\{\frac{\theta}{P} + \frac{(1+t)h}{B}\right\} Z_b^2 + \frac{2(1+t)H_0}{B} Z_e^2. \tag{BE12}$$

For u infinitesimal each displayed coefficient has a positive-unit residue, including $\theta \to 0$, because $\theta + h = 1$. To retain the first normal exactly, define

$$A_x = \frac{H_0}{B}\left\{\frac{xz(J-\sigma)}{P} - t\right\} - \frac{H_0}{P},$$

$$c_d = 4ud_0\left\{A_x - \frac{\sigma\mathcal{H}}{PJ}(1+th)\right\},$$
$$c_b = 4kb_0\left\{A_x + \frac{\mathcal{H}}{B}\frac{\sigma z(J-\sigma)}{P}\right\} - 4tp\frac{\mathcal{H}}{B}.$$

The middle line of (BE11) is exactly $c_d N_x Z_d + c_b N_x Z_b$. Both coefficients are infinitesimal when u is infinitesimal, through every source opening and either sphere endpoint. Extracting $4(\beta - \beta_*)N_x^2$ for any bounded $0 \le \beta_* \le \beta$ leaves a bounded Gram on (N_x, Z_d, Z_b, Z_e) with strictly positive diagonal residue. Hence this normal block has the standard bounded positive-Gram lift. No inverse of d_0, b_0, u, or θ enters these coefficients. At $d_0 = 0, u = 0$ all three Z roots vanish, as the exact null direction requires. At $b_0 = 0$, the Z_b root and its cross coefficient vanish.

For the full $w \neq 0$ contact conductor the corresponding proposed roots are $d_0(\sigma Z - uwW)$, $b_0(u/k)(\sigma Z - uwW)$, and $\sqrt{e}\,(u/k)(\sigma Z - uwW)$, together with the existing radial root and angular roots. Equations (BE11)–(BE12) verify the formerly collapsed normal direction and its first-normal coupling, but do not allocate the scalar contact polynomial and the w-dependent radial couplings. Their *joint* allocation is the remaining step for this proposed frame. No positivity claim for that uncomputed Gram is made here, and no endpoint already known to be elementary is counted as a new source-domain admission.

Chapter 27

C1.4: unbounded $L = x\zeta$, bounded-unit $\sigma = 1/\eta$

Component `pd4b92bf7de71`
Source: `R13/advances/c1_large_l_eta_unit/large_l_unit.tex`
SHA256: `d4b92bf7de710d88bc5573c42f6166cc28551992da3378645013dea8c9352474`
Inventory use: whole theorem and raw/capped full-parent proof | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

27.1 C1.4: unbounded $L = x\zeta$, bounded-unit $\sigma = 1/\eta$

This section proves this full original-parent subdomain, retaining every state normal and every independent source/testing rate. It makes no claim for infinitesimal σ. The inherited common remote theorem leaves u bounded and $w = 1/v$ infinitesimal; both u infinitesimal and u a bounded unit are covered below. All calculations use the exact NC source, with $d = 1$, $n = \lambda = 0$, and its original nonnegative D, β faces.

Source chart and bounded products. Put $x = k^2$, $z = q^2 = L^{-1}$, $r = bz$, $Q = 1 - b$, $\theta = 1 - h$, $K = 1 + x$, $R = 1 + xbz$, $M = R - xz\theta$, and

$$J = 1 + xh - \sigma(1 - bz), \qquad E = 1 - xh/\sigma.$$

Here x, z are positive infinitesimals, $0 < b < 1$, and σ is a positive bounded unit. The NC source gives exactly

$$P = \sigma + K + \sigma z(1 + bx + 2xh) + \sigma bx^2hz^2 + x^2h\theta z,$$
$$V_0 = z(1 + xh)E + (1 - bz)(1 + xhz),$$
$$C = cz, \qquad c = \frac{K(1 + \sigma bz)E}{2(1 - bz)P},$$
$$m = \mu x^2 z, \qquad \mu = \frac{K(1 + \sigma bz)h(1 + \sigma z)}{2\sigma MP},$$
$$A = a/z, \qquad a = \frac{V_0}{2bP},$$
$$\alpha = \ell/z, \qquad \ell = \frac{(1 + \sigma bz)[(x/\sigma)Rh(1 + \sigma z) + ME]}{2QP},$$
$$D = \frac{K(1 + \sigma z)(1 + \sigma bz)}{2JP} - \frac{1}{2}, \qquad \beta = \frac{K(1 + \sigma bz)(1 + xhz)}{2\theta P} - \frac{1}{2}.$$

In particular $c, \mu, ab, \ell Q, DJ, \beta\theta$ are bounded, $c, \mu, ab, \ell Q, DJ$ are positive units, and $ab - \ell Q \in z\mathcal{O}$. At $x = z = 0$ their residues are

$$c_0 = \frac{1}{2(1+\sigma)}, \quad \mu_0 = \frac{h}{2\sigma(1+\sigma)}, \quad (ab)_0 = (\ell Q)_0 = \frac{1}{2(1+\sigma)},$$
$$(DJ)_0 = \frac{\sigma^2}{2(1+\sigma)}, \quad (\beta\theta)_0 = \frac{(1+\sigma)h - \sigma}{2(1+\sigma)}.$$

Source nonnegativity gives $0 < \sigma \le 1$ and $h \ge \sigma/(1+\sigma)$ at residue. Thus h is a positive unit. The openings b, Q, J, θ remain independent; $\beta = 0$ is retained. The face $D = 0$ cannot occur here since DJ is a positive unit.

Write $D_u = 1 + u^2$, $D_w = 1 + w^2$ and

$$M_w = W - wX + uY - uwZ, \quad R_w = uW - Y, \quad J_w = uwW + uX - wY - Z, \quad B_w = X + uZ,$$
$$K_w = J_w^2 + D_u(Y - wZ)^2 + D_w B_w^2 - D_u D_w Z^2.$$

The complete reciprocal parent is

$$F_w = 4mM_w^2 + 4czD_wR_w^2 + 4(a/z)D_uD_wY^2 + 4DJ_w^2 + 4(\ell/z)D_u(Y - wZ)^2 + 4\beta D_w B_w^2 + 2K_w.$$

Choose bounded positive-unit caps $a_* \le a$, $\ell_* \le \ell$, $D_* \le D$, and a bounded nonnegative cap $\beta_* \le \beta$. Keep as nonnegative squares the four centered excesses

$$4(a - a_*)D_uD_wN_y^2, \quad 4(\ell - \ell_*)D_uN_y^2,$$
$$4(D - D_*)(J_w + JZ)^2, \quad 4(\beta - \beta_*)D_wN_x^2,$$
$$N_x = X + uhZ, \qquad N_y = (Y - bwZ)/q.$$

These roots vanish at the exact source contact

$$u = k, \quad w = q\sqrt{\sigma}, \quad (W, X, Y, Z) = (\sqrt{\sigma}/(kq), -kh, bq\sqrt{\sigma}, 1).$$

Set $d_J = (D - D_*)J$ and $b_\theta = \beta\theta$. Their coefficients and the centered excesses preserve the original parent.

Bounded full-normal remainder. Define $G = wZ/q$, $R_h = G - uqW$ and

$$c_N = 8D_u[abD_w - \ell Q], \qquad c_G = 4D_u[ab^2D_w + \ell Q^2].$$

The A, α remainder is $4D_u(a_*D_w + \ell_*)N_y^2 + c_NN_yG + c_GG^2$. Here $c_N \in (z, w^2)\mathcal{O}$ and c_G is a positive unit. Keep $c_GR_h^2 + c_NN_yR_h$ in the bounded Gram and use $G^2 - R_h^2 = 2uwWZ - u^2zW^2$. The rest is the bounded polynomial

$$\begin{aligned} G_0 = {} & 4D_u(a_*D_w + \ell_*)N_y^2 + c_NuqWN_y + c_G(2uwWZ - u^2zW^2) \\ & + 4mM_w^2 + 4czD_wR_w^2 + 4D_*J_w^2 - 8d_JZJ_w - 4d_JJZ^2 \\ & + 4\beta_*D_wB_w^2 + 8(b_\theta - \beta_*\theta)uD_wZB_w - 4(b_\theta - \beta_*\theta)u^2\theta D_wZ^2 + 2K_w. \end{aligned}$$

All its coefficients are bounded. Put $N_z = Z - uwW/\sigma$ and substitute $Z = N_z + uwW/\sigma$, $X = N_x - uhZ$, $Y = qN_y + bwZ$. For $t = u^2$, $\xi = w^2$, define

$$T_0 = 1 + th + b\xi, \quad M_0 = 1 + t(h - Q)\xi/\sigma, \quad R_f = 1 - b\xi/\sigma, \quad J_f = 1 - T_0/\sigma.$$

In the formulas below c_N, c_G mean their even-polynomial expressions. The exact expansion is

$$G_0 = \phi W^2 + wL_xWN_x + uqL_yWN_y + uwL_zWN_z + H(N_x, N_y, N_z),$$

where

$$\begin{aligned}
\phi &= 4\mu x^2 z M_0^2 + 4cz(1+\xi)tR_f^2 + c_G t(2\xi/\sigma - z)\\
&\quad + 4D_* t\xi J_f^2 - 8d_J t\xi J_f/\sigma - 4d_J Jt\xi/\sigma^2 + 4b_\theta\theta(1+\xi)t^2\xi/\sigma^2\\
&\quad + 2[t\xi J_f^2 + (1+t)tQ^2\xi^2/\sigma^2 + (1+\xi)t^2\theta^2\xi/\sigma^2 - (1+t)(1+\xi)t\xi/\sigma^2],\\
L_x &= -8\mu x^2 z M_0 + t[(8D_*+4)J_f - 8d_J/\sigma + (8b_\theta + 4\theta)(1+\xi)/\sigma],\\
L_y &= 8\mu x^2 z M_0 - 8cz(1+\xi)R_f + c_N - (8D_*+4)\xi J_f + 8d_J\xi/\sigma - 4(1+t)Q\xi/\sigma,\\
L_z &= 8\mu x^2 z(h-Q)M_0 - 8cz(1+\xi)bR_f + 2c_G - (8D_*+4)T_0 J_f - 8d_J J_f + 8d_J(T_0 - J)/\sigma\\
&\quad + 8b_\theta\theta(1+\xi)t/\sigma + 4(1+t)Q^2\xi/\sigma + 4(1+\xi)t\theta^2/\sigma - 4(1+t)(1+\xi)/\sigma.
\end{aligned}$$

For a compact explicit specification of the entire quadratic block, put

$$\begin{gathered}
m_N = -wN_x + uqN_y + uw(h-Q)N_z, \quad r_N = -qN_y - bwN_z,\\
j_N = uN_x - wqN_y - T_0 N_z, \quad b_N = N_x + u\theta N_z.
\end{gathered}$$

Then every quadratic normal coefficient is retained by the finite formula

$$\begin{aligned}
H &= 4D_u(a_* D_w + \ell_*)N_y^2 + 4mm_N^2 + 4czD_w r_N^2 + (4D_*+2)j_N^2 - 8d_J N_z j_N - 4d_J J N_z^2\\
&\quad + (4\beta_*+2)D_w b_N^2 + 8(b_\theta - \beta_*\theta)uD_w N_z b_N - 4(b_\theta - \beta_*\theta)u^2\theta D_w N_z^2\\
&\quad + 2D_u(qN_y - QwN_z)^2 - 2D_u D_w N_z^2.
\end{aligned}$$

The exact source substitutions give, at $(t,\xi) = (x, \sigma z)$,

$$\phi = \phi_t = \phi_\xi = L_x = L_y = L_z = 0.$$

Moreover $\phi(0,\xi) = 4\mu x^2 z$, $L_x(0,\xi) = -8\mu x^2 z$, and $\phi(t,\sigma z) \in z\mathcal{O}[t]$. The checker verifies all six source jets and these supports separately.

Exact six-root lift in the critical strip u infinitesimal. Monic division, first by $\xi - \sigma z$ and then by $t - x$, gives bounded polynomial coefficients satisfying

$$\begin{aligned}
\phi &= z(t-x)^2 A + t(t-x)(\xi - \sigma z)B + t(\xi - \sigma z)^2 C,\\
L_x &= (t-x)(tA_x + zB_x) + t(\xi - \sigma z)C_x,\\
L_y &= (t-x)A_y + (\xi - \sigma z)B_y,\\
L_z &= (t-x)A_z + (\xi - \sigma z)B_z.
\end{aligned}$$

For example, the constant coefficient of $L_x(t,\sigma z)/(t-x)$ equals $8\mu xz$, giving its (t,z) support. No small or mixed inverse enters these monic quotients. Also $A(0) = 4\mu$. Take the six finite separated roots

$$N_x, \quad N_y, \quad N_z, \quad R_h, \quad A_\theta = q(t-x)W, \quad V = uw(t-x)W.$$

The complete scalar lift follows from

$$\begin{aligned}
z(t-x)^2 W^2 &= A_\theta^2,\\
t(t-x)(\xi - \sigma z)W^2 &= u\sigma A_\theta R_h - \sigma V N_z,\\
t(\xi - \sigma z)^2 W^2 &= \sigma^2(qR_h - wN_z)^2.
\end{aligned}$$

All first normals lift by the following finite list, which includes their whole coefficient support:

$$\begin{aligned}
wt(t-x)W &= uV, & wz(t-x)W &= qwA_\theta,\\
wt(\xi - \sigma z)W &= u\sigma qwR_h - u\sigma w^2 N_z, & &\\
uq(t-x)W &= uA_\theta, & uq(\xi - \sigma z)W &= \sigma z R_h - \sigma qwN_z,
\end{aligned}$$

$$uw(t-x)W = V, \qquad uw(\xi-\sigma z)W = \sigma q w R_h - \sigma w^2 N_z.$$

Together with the displayed H these give an exact bounded six-root Gram, not merely a restriction to the contact line. At residue its first five diagonal pivots are respectively

$$2+4\beta_*, \quad 4(a_*+\ell_*), \quad 4[D_*+d_J(1+\sigma)], \quad \frac{2}{1+\sigma}, \quad 4\mu_0.$$

They are positive units through all allowed openings. The only potentially nonzero remaining cross entry is $N_z V$. The exact relation

$$V^2 = \sigma t A_\theta^2 + u\sigma(t-x)A_\theta R_h - \sigma(t-x)VN_z$$

introduces an arbitrarily large finite positive V^2 pivot with only infinitesimal changes to all other entries. Its real Schur complement therefore makes the residue strictly positive definite. The exact bounded Gram is positive definite by the standard finite-dimensional local criterion. All excess squares are nonnegative with separated coefficients.

Away from the angular contact: u a bounded unit. The same source chart has a positive-unit angular gap at every such real residue. A convenient raw finite frame is $E_t = qW, F_t = Y/q, G_t = wZ/q, R_t = wW, X, Z$; products obey $E_tG_t = R_tZ$. The residue parent divided by $2/(1+\sigma)$ is exactly

$$u^2E_t^2 + D_u\{F_t^2/b + (F_t-G_t)^2/Q\} + \frac{(uR_t+uX-Z)^2}{J_0} + \frac{(X+uZ)^2}{\theta} - (1+\sigma)D_uZ^2, \quad J_0 = 1-\sigma.$$

Eliminate the positive F_t and X directions and allocate the common product between the remaining two quadratic blocks. Up to a common positive factor the two blocks have diagonal products AB, CD_0 and total off-diagonal magnitude E_0, with $U = u^2 > 0$ and

$$\delta = 1-\sigma+\theta U, \qquad A = U, \quad B = 1+U, \quad C = U/\delta,$$
$$D_0 = (1+U)[(1+U)/\delta - (1+\sigma)], \quad E_0 = \sqrt{U}(1+U)/\delta.$$

Both diagonal products are strictly positive, since $1+U-(1+\sigma)\delta = \sigma^2 + U[(1+\sigma)h-\sigma] > 0$. All square roots in this sentence are real-residue quantities used to choose real allocation constants, not mixed tensor radicals. The required strict inequality is $E_0 < \sqrt{AB}+\sqrt{CD_0}$. Set $N = U(2\sigma-\theta U)$. Directly,

$$E_0^2 - AB - CD_0 = (1+U)N/\delta, \quad N^2 - 4U^2[1+U-(1+\sigma)\delta] = U^3(\theta^2U-4h).$$

If $N \le 0$ strictness is immediate. If $N > 0$, then $\theta^2U < 2\sigma\theta \le 2h < 4h$, again giving strictness. The single-opening faces follow by their constrained limits. At the only double opening $J = \theta = 0$, so $\sigma = h = 1$, the exact limit is $E_0 - \sqrt{CD_0} \longrightarrow \sqrt{U}$, whereas $AB - U = U^2 > 0$. Here is a full-frame square completion which also justifies those cap limits. For clarity first take $u > 0$ and select a finite real ρ in the nonempty interval

$$\max\{-\sqrt{AB}, E_0-\sqrt{CD_0}\} < \rho < \min\{\sqrt{AB}, E_0+\sqrt{CD_0}\}.$$

Its nonemptiness follows from the strict allocation inequality above. Put

$$r_\rho = D_u/u - \rho\delta/U, \qquad s_\rho = D_u[2\rho/u-(1+\sigma)] - \rho^2\delta/U > 0.$$

The preceding complete residue form equals

$$UE_t^2 + D_uG_t^2 - 2\rho E_tG_t + \frac{D_u}{bQ}(F_t-bG_t)^2$$
$$+ \frac{\delta}{J_0\theta}\left[X + \frac{U\theta R_t + u(J_0-\theta)Z}{\delta}\right]^2 + \frac{U}{\delta}(R_t-r_\rho Z)^2 + s_\rho Z^2.$$

The equality uses only $E_tG_t = R_tZ$ and is checked exactly. For $u < 0$ take the correspondingly signed ρ; all sign choices are real constants in this allocation. All triangular minimizer coefficients are bounded through every opening because U is a unit and $\delta = J_0 + U\theta$. Their positive hard weights stay bounded below. At the double face choose $|u| < |\rho| < \sqrt{U(1+U)}$, with matching sign; then s_ρ has a positive-unit limit. This proves uniform coercivity of the constrained residue, not just determinant positivity.

For completeness the exact centered cap rows in the raw frame are

$$F_t - bG_t, \quad uR_t + uX - qwF_t + (J-1)Z, \quad X + uhZ.$$

Only one of b, Q can tend to zero. The other two rows have hard parts on (R_t, X) with determinant u, a unit; each single row has a unit hard entry. Their simultaneous kernel is solved exactly by

$$F_t = bG_t, \qquad X = -uhZ, \qquad R_t = \{(u^2h - J + 1)Z + qwbG_t\}/u.$$

More generally, prescribed changes (e_F, e_D, e_β) in these three rows are produced by $\Delta F_t = e_F$, $\Delta X = e_\beta$, $\Delta R_t = (e_D - ue_\beta + qwe_F)/u$. This is a bounded separated right inverse: its only inversion is the pure-testing unit u. For the finite-cap existence argument fix u at its ordinary real residue, so the same inverse is an ordinary real finite-dimensional right inverse. No order on the tensor product is used. It lifts the limiting constrained kernel to the exact centered kernel while keeping every opening independent. The bounded residual form restricted to that kernel is the strictly positive constrained form just computed. An elementary block Schur complement then provides finite sufficiently large caps on every divergent row; bounded rows retain their full coefficients. This is the finite-cap lemma used in the UH proof. It yields a strictly positive bounded full-frame Gram. Exact coefficients differ from their residues infinitesimally, completing this strip. All divisions by δ and radicals in this paragraph belong only to the ordinary real residue calculation and its face limits. The exact tensor Gram uses the polynomial raw frame, real allocation constants, and finite source caps; it introduces no mixed inverse or radical.

Consequently this section closes the L unbounded, σ bounded-unit subdomain of C1.4 under the original parent, including all state normals, all independent opening rates, and $\beta = 0$. It does not close the σ infinitesimal subdomain or claim a whole-row count change.

Chapter 28

Small L, bounded η: exact source openings and the remote complement

Component `p6d2f659c0fde`
Source: `R13/advances/c1_lsmall_eta_bounded/remote_reduction.tex`
SHA256: `6d2f659c0fde22063ee3393474686230de529c0a1b18e3b727263ffa9d28ad83`
Inventory use: SB1–SB12 and full remote proof | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

28.1 Small L, bounded η: exact source openings and the remote complement

Work in the original C1 source, with $d = 1$, $x \in \mathrm{Inf}_{>0}$, ζ noninfinitesimal, $L = x\zeta \in \mathrm{Inf}_{>0}$, and η bounded. Native $r = 1 - a\eta$, $h = 1 - \theta$ remain strictly positive. This section admits every original projective test except possibly u, v both infinitesimal. It does not admit the remaining two-soft local chart.

Exact source restrictions, including the total-scale intersection. Put $\rho = r + \eta$ and $j = \eta J = \eta(1 + xh) - 1 + r > 0$. Since $a < 1 + xh$, ρ is a positive bounded unit. The exact beta numerator is

$$2\beta\theta P_d = K\rho(xh + L) - \theta P_d. \tag{SB1}$$

It implies $\theta = O(x + L) = O(L)$, since $x/L = 1/\zeta$ is bounded. Consequently

$$h, M, E, Q, P_d, P_4 \sim 1, \qquad B_5 \sim 1. \tag{SB2}$$

For the last assertion, use the exact expression $B_5 = K(1+L\eta)\rho - jP_d$: its residue is $r+\eta-(\eta-1+r) = 1$. Thus original $D = 0$ cannot occur on this source face; literal $\beta = 0$ remains included. The original formulas, with openings canceled before taking residues, give uniformly

$$\begin{gathered} C(1 - r) \sim \rho/2, \quad Ar/\eta \sim 1/2, \quad Dj \sim 1/2, \\ mL/x^2 \sim \rho/2, \quad \alpha/[L(L + x)] \sim \rho/2, \quad (2\beta + 1)\theta/(x + L) \sim \rho. \end{gathered} \tag{SB3}$$

In particular C, D are strong, m, α weak, and A is strong precisely when η is a unit. The four exact large-weight alternatives are

weight unbounded	equivalent opening
C	$1 - r \in \mathrm{Inf}$
D	$j \in \mathrm{Inf}$
A	$r \in \mathrm{Inf}$
β	$\theta/(x + L) \in \mathrm{Inf}$.

(SB4)

For the A equivalence, $r \in \mathrm{Inf}$ forces η to be a unit because $j > 0$; if r is a unit, η/r is bounded. All coefficients are bounded exactly when the four quantities in the right column of (SB4) are positive bounded units. That entire interior is already admitted by the positive-total-scale theorem. After intersection with $\delta_{\mathrm{tot}} \in \mathrm{Inf}_{>0}$, at least one of (SB4) therefore occurs. These are exhaustive necessary source alternatives, not assertions of their independent realizability.

The unit-η remote tests: two complete three-state blocks. Suppose first η is a unit, with real residue $e > 0$, and write $r_0 = \operatorname{res} r \in [0, 1]$. For an interior opening, the source residues are

$$C_0 = \frac{r_0 + e}{2(1 - r_0)}, \quad A_0 = \frac{e}{2r_0}, \quad D_0 = \frac{1}{2(e + r_0 - 1)}. \tag{SB5}$$

Retain the original m, α, β packets throughout. Bounded selected weights retain their actual coefficients; unbounded ones are capped at sufficiently large finite real numbers and their excesses retained as exact squares.

For u a bounded nonzero unit and v infinitesimal, use the full raw frame

$$p = uW, \quad Y, \quad R = vZ, \quad B_t = X + uZ.$$

The exact core is (CE2), and the D event is $J = p - Y + uvB_t - D_u R$. Every Gram coefficient is bounded after capping, and no state normal is omitted. Set $U = \operatorname{res} u^2 > 0$. The residue, divided by four, is $B_t^2/2$ plus

$$\begin{aligned} f_U = {} & C_0(p - Y)^2 + A_0(1 + U)Y^2 + D_0\{p - Y - (1 + U)R\}^2 \\ & + \tfrac{1}{2}\{(p - Y)^2 + (1 + U)UR^2 - 2(1 + U)pR\}. \end{aligned} \tag{SB6}$$

Its first two principal minors and determinant are, respectively,

$$\begin{aligned} \Delta_1 &= \frac{e(e + r_0)}{2(1 - r_0)j_0}, \qquad \Delta_2 = \frac{e^2(1 + U)(e + r_0)}{4r_0(1 - r_0)j_0}, \\ \Delta_3 &= \frac{Ue(1 + U)^2(e + r_0)^2}{8r_0(1 - r_0)j_0} > 0, \qquad j_0 = e + r_0 - 1. \end{aligned} \tag{SB7}$$

For u infinitesimal and v a bounded nonzero unit, use the full frame $(p = uW, X, Y, R = vZ)$. The same exact parent has residue, divided by four, $(1 + V)X^2/2$ plus

$$\begin{aligned} f_V = {} & C_0(1 + V)(p - Y)^2 + A_0(1 + V)Y^2 + D_0(p - Y - R)^2 \\ & + \tfrac{1}{2}\{(p - Y)^2 + VY^2 - 2pR\}, \qquad V = \operatorname{res} v^2 > 0. \end{aligned} \tag{SB8}$$

The leading two minors are positive, and

$$\det f_V = \frac{V(1 + V)(e + r_0)^2}{8r_0(1 - r_0)j_0} > 0. \tag{SB9}$$

These arguments include every opening in (SB4). To see explicitly why the finite cap remains valid, impose the kernels of the divergent events and inspect the remaining two-state Gram. The determinants are

opening	$\det(f_U\|_{\ker})$	$\det(f_V\|_{\ker})$
$r_0 = 0,\ e > 1$	$\dfrac{Ue^2(1 + U)}{4(e - 1)}$	$\dfrac{Ve}{4(e - 1)}$
$r_0 = 1$	$\dfrac{U(1 + U)^2(e + 1)}{4}$	$\dfrac{V(e + 1)}{4e}$
$j_0 = 0,\ 0 < r_0 < 1$	$\dfrac{U(1 + U)^2}{4r_0}$	$\dfrac{V(1 + V)}{4r_0(1 - r_0)}$.

(SB10)

The respective kernels are $Y = 0$, $p = Y$, and $p - Y = (1 + U)R$ for f_U or $p - Y = R$ for f_V. All first pivots are positive. At the only double opening, $r_0 = j_0 = 0, e = 1$, the two kernels leave the positive

scalar $U(1+U)R^2/2$, respectively $VR^2/2$. There are no other multiple openings since $e>0$. The elementary finite-penalty Schur lemma therefore supplies finite real caps simultaneously for every divergent selected event.

For completeness the remaining projective ends also use original homogeneous degree. If u is unbounded, (CE4)–(CE5) for bounded v, and (CE7)–(CE8) for unbounded v, are strict with any positive finite C, A caps. If v is unbounded and u bounded, put $w=1/v$ and use $(p=uW, X, Y, Z)$. The exact D event is $w(p-Y)+uX-Z$. At $w=0$ the core is $(1+u^2)(X^2+Y^2)$, while the three caps give $C(p-Y)^2+A(1+u^2)Y^2+D(uX-Z)^2$; these are strict on all four roots, including $u=0$. Each term suppressed in this residue has a bounded infinitesimal coefficient in this same full frame. Literal $w=0$ is therefore included. The both-unit tests are covered by the inherited noncommuting-residue theorem. Together these statements exhaust the complement of $u, v \in \mathrm{Inf}$.

The infinitesimal-η remote tests. Here $r\sim 1$, C, D are unbounded, and $y=L\eta$ is infinitesimal. Define the pure-source harmonic coefficient $\gamma = yCD/(C+D)$. Uniformly through the a, J openings,

$$\gamma \sim L/2, \qquad A/y \sim 1/(2L), \qquad \gamma A/y \sim 1/4, \qquad y/\gamma \sim 2\eta \in \mathrm{Inf}\,. \tag{SB11}$$

The strong-II remote proof (RR3)–(RR6), which uses only C, D unbounded, covers every noninfinitesimal v and every u. For u noninfinitesimal and v infinitesimal, use (RR7)–(RR9) with zero alpha cap and $k=-1$. In its notation let $a=u/\sqrt{1+u^2}$, $b=1/\sqrt{1+u^2}$, $\ell = vY$, $x_t = bX$, $R_0 = x_t + a(Z-\ell)$, and $q=\sqrt{y}Y$. Then

$$M_{-1} = \tfrac{1}{2}R_0^2 + \tfrac{1}{2}(b\ell)^2, \qquad v(Z-\ell) = avR_0 - bv(b\ell) - b\sqrt{y}\,V_h, \tag{SB12}$$

where $V_h = v(aX-bZ)/\sqrt{y}$. Normalize the harmonic pair as $P_h = \sqrt{\gamma}V_h$, $Q_h = \sqrt{A/y}\,q$. Its residue determinant is $1-b_0^2>0$ by (SB11), where $b_0 = \operatorname{res} b < 1$; the off-diagonal entry is b_0. The sole negative term in (RR9), $-v^2(Z-\ell)^2/2$, is an infinitesimal Gram perturbation of $(R_0, b\ell, P_h)$ by (SB12), since $\sqrt{y/\gamma}$ is infinitesimal. Thus the full four-root Gram is strict. Retain the original positive $\alpha(Z-\ell)^2$, the C1 beta event, and every other unused packet. The homogeneous endpoint $u=\infty$ has $a=\pm 1, b=0$ and is included; literal $v=0$ also causes no change.

The actual remaining obstruction. Only the simultaneous infinitesimal-test chart remains, intersected with (SB4) and all previous admissions. It has two soft state directions, not the single soft ray of the admitted L-unit proof. Indeed the original parent at $u=v=0$ is exactly

$$F = (4m+4\beta+2)X^2 + (4C+4A+4D+2)Y^2 + 4\alpha Z^2. \tag{SB13}$$

Since m, α are weak, the two W, Z directions remain soft after the hard X, Y normals are normalized. On the contact graph $X=-huZ$, $Y=ruW$, the exact two-soft scalar is (WS2)–(WS3) with $n=0$, the WS letters h, r replaced by native r, h, and the additional positive term $\beta\theta^2U(1+V)Z^2$ in $F_g/4$. Its two negative monomials are still $-2uv(b_0+b_1U)WZ$. No strict finite Gram on the existing hard roots supplies the missing W, Z pivots. A weighted contact circuit and its complete first-normal allocation are still needed. Neither the exact source alternatives nor the remote theorem is presented as that local completion.

Chapter 29

A C1 extension: L a unit and η infinitesimal

Component `p080aa62de16d`
Source: `R13/advances/c1_lunit/lunit_eta_small.tex`
SHA256: `080aa62de16d01bccf6ca074a21eee4a3010f088ec44dad2749c905e46d87ebd`
Inventory use: LC1–LC7 and local/remote join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

29.1 A C1 extension: L a unit and η infinitesimal

Theorem 29.1. *The entire original C1 source subdomain*

$$x = k^2 \in \mathrm{Inf}_{>0}, \qquad L = x\zeta \text{ a positive bounded unit}, \qquad \eta \in \mathrm{Inf}_{>0}$$

has a full-parent tensor-SOS certificate for every independent projective testing pair. All original openings and the retained boundaries $D = 0$ or $\beta = 0$ remain allowed whenever they occur in this source subdomain.

The proof is an exact adapter to the strong-II local and remote certificates. Its additional event is retained with a centered square; it is not deleted from the parent.

The native source in the shared contact geometry. Keep the native notation of (NC1)–(NC4), but in this section denote its r, h, a by r_n, h_n, a_n to avoid collisions. Normalize $d = 1$. The exact native contact with $Z_c = 1$ is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(\sqrt{x}, \sqrt{x\eta\zeta}; \sqrt{\zeta/\eta}, -\sqrt{x}\, h_n, \sqrt{x}\, r_n \sqrt{\zeta/\eta}, 1\right).$$

It therefore has the same contact geometry used in (LII2), with

$$z = x, \quad y = L\eta, \quad B = L, \quad H = Lr_n = B - a_n y, \quad r = h_n = 1 - \theta, \quad t = a_n, \quad j = 1 + zr - t = J_n. \tag{LC1}$$

In particular z, y are positive infinitesimals. Here $r_n \sim 1$, $0 < Lr_n < 1$, and hence $b_0 := \operatorname{res} B$ satisfies $0 < b_0 \le 1$. We use b_0 only for this source residue.

Set, as in the local strong-II proof,

$$c = Cyt, \quad d_p = Dyj, \quad \mu = m/z^2, \quad a_A = A/y, \quad \sigma = \alpha(1 - H), \quad \tau = \beta(1 - r) = \beta\theta.$$

These are bounded source quantities, and c, d_p, μ, a_A, σ are positive units. Writing $r_0 = \operatorname{res} r$, exact substitution in (NC2)–(NC4), with opening factors canceled first, gives

$$\begin{aligned} c, d_p &\sim \frac{b_0}{2(1+b_0)}, \qquad a_A \sim \frac{1}{2b_0(1+b_0)}, \\ \mu &\sim \frac{r_0}{2b_0(1+b_0)}, \qquad \sigma \sim \frac{b_0^2}{2(1+b_0)}, \\ \operatorname{res}\tau &= \frac{b_0}{2(1+b_0)} - \frac{1-r_0}{2}. \end{aligned} \tag{LC2}$$

Since $\tau \geq 0$, the last identity implies $r_0 \geq 1/(1+b_0) > 0$, proving the asserted unit lower bound for μ. Also $t + j = 1 + zr \sim 1$, and c, d_p being units proves C, D unbounded uniformly through either opening $t = 0$ or $j = 0$ in residue. Thus $D = 0$ is in fact absent from this specific subdomain by the exact source implications; it has not been excluded by an additional hypothesis. Literal $\beta = 0$ is included.

The source-harmonic quantity $yCD/(C + D)$ and the cap behavior of α have precisely the residues (RR1)–(RR2), with $h = b_0$. The remote proof (RR3)–(RR13) therefore applies unchanged, retaining the extra positive packet $4\beta D_v(X + uZ)^2$ throughout. This covers every test outside the local region where u, v are both infinitesimal, including the original projective endpoints.

The additional event and its exact local remainder. For the local region use exactly the graph, roots, harmonic pair, and caps (LII2)–(LII4), with the source quantities (LC1)–(LC2). Put

$$R = v(U - z)W, \qquad E_\beta = \xi + u\zeta + R/B = (X + uZ) - \frac{z(1-r)v}{B}W. \tag{LC3}$$

This root vanishes at the actual contact. Retain the entire positive packet $4\beta D_v E_\beta^2$, in addition to the first, second and fourth packets of (LII5); there is no n packet since $n = 0$. The exact remainder from the β event is

$$\begin{aligned} \Delta_\beta = {} & \frac{8\tau z}{B} D_v v W(\xi + u\zeta) \\ & + \frac{4\tau z}{B^2} D_v V\{2U - z(1+r)\}W^2. \end{aligned} \tag{LC4}$$

All its coefficients are bounded even if β is unbounded: only $\tau = \beta(1 - r)$ occurs. The exact full remainder is (LII6) with its n line deleted, plus (LC4). Thus no state normal, boundary coefficient, or first contact jet has been omitted.

The $R_h W$ gauge (LII7) is unchanged, because (LC4) contains no R_h. The coefficient representation (LII8), its exact contact and first-normal vanishings (LII9)–(LII10), and its bounded monic divisions (LII12)–(LII13) all remain valid. Explicitly, (LC4) adds $8\tau z(1 + V)/B$ to P_ξ and P_ζ, adds zero to P_N, and adds its printed scalar line to ϕ. Hence the essential support $P_\xi \in (U, z)$ and $\deg_U \phi \leq 2$ remain literal polynomial statements.

The only changed data in (LII11) are

$$\begin{aligned} d_s &= \frac{4z^2}{B^2}\left\{\mu(B + zr)^2 + \frac{r^2}{2} - \tau(1+r)\right\}, \\ e_s &= \frac{4z^2}{B^2}\left\{\frac{r^2}{2} - \tau(1+r)\right\}. \end{aligned} \tag{LC5}$$

Both quotients by z^2 are bounded; e_s need not be positive. Its contribution to the final Gram has an infinitesimal coefficient because it is multiplied by V. The decisive exact residue cancellation, using (LC2), is

$$\begin{aligned} \operatorname{res}\left\{\mu B^2 + \frac{r^2}{2} - \tau(1+r)\right\} &= \frac{b_0 r_0}{2(1+b_0)} + \frac{r_0^2}{2} - \left\{\frac{b_0}{2(1+b_0)} - \frac{1-r_0}{2}\right\}(1+r_0) \\ &= \frac{1}{2(1+b_0)}. \end{aligned} \tag{LC6}$$

Consequently

$$\operatorname{res}(d_s/z^2) = \frac{2}{b_0^2(1+b_0)}, \qquad \operatorname{res} A_\xi = \frac{4}{b_0(1+b_0)}. \tag{LC7}$$

The second equality is unchanged from (LII18): the new P_ξ term is a bounded multiple of z independent of U, so it has zero U-quotient residue. The four hard-normal residues (LII17) are unchanged as well.

Apply the exact allocations (LII15) to all the coefficients, including (LC4). They give the same bounded six-root Gram $(N, \xi, \zeta, R_h, R, T)$ as in (LII16). Its finite (ξ, R) block is exactly (LII19), with $h = b_0$, and has strictly positive Schur complement $2/[b_0(1+b_0)^2]$. The same exact conductor (LII20) supplies an arbitrarily large real T^2 pivot while changing every other residue entry by zero. The complete bounded Gram is strict. Restoring the four retained packets, including $4\beta D_v E_\beta^2$, proves the local tensor SOS for all independent rates and literal affine axes. The remote admission above completes the theorem.

Verification and exact frontier effect. The accompanying checker reconstructs the entire original C1 parent from this remainder and all four retained packets, verifies the new centered-event identity, the unchanged contact-ideal support, every new source residue, the finite block, and the conductor. Stationarity itself is the unchanged native source identity, not an added restriction. The result removes the whole L-unit, η-infinitesimal subdomain from C1.4. It does not assert the other L-unit source regimes or the entire C1.4 row.

Chapter 30

A sharper source bound on the remaining L-unit, η-large face

Component `p57d751e20564`
Source: `R13/advances/c1_lunit_eta_large/source_frontier.tex`
SHA256: `57d751e20564de6ea20a95078481d09cfe4389c6fb59ba65ada8a88eb9fef362`
Inventory use: opening-cleared source formulas | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

30.1 A sharper source bound on the remaining L-unit, η-large face

This section proves a source restriction and a leading quadratic identity. It does not claim the remaining full-parent admission. Use the unchanged native notation with $x \in \mathrm{Inf}_{>0}$, $L = x\zeta$ a positive unit, and η unbounded. Write $\theta = 1 - h$ and $K = 1 + x$.

In the exact polynomial $B_5 = a_2\eta^2 + a_1\eta + a_0$, substitution $\zeta = L/x$ gives

$$a_2 = -xh\{KL + x\theta(1 + xh)\}, \qquad a_1 = xb_1, \tag{LH1}$$

where

$$\begin{aligned} b_1 = {} & -Lhrx - Lh - Lr + L - h^2rx^2 + h^2rx - 3h^2x \\ & - 2hrx + hx - 3h + 1, \end{aligned}$$

and

$$a_0 = Kr + (1 - r)\{1 + L + 2xh + rxL + rx^2h\}.$$

Both b_1 and a_0 are bounded uniformly in every source opening, while $KL + x\theta(1 + xh)$ has a positive-unit lower bound. The original $B_5 \geq 0$ inequality therefore implies

$$x\eta h \leq O(x) + O(\eta^{-1}) \in \mathrm{Inf}, \qquad \boxed{E = 1 - x\eta h \sim 1.} \tag{LH2}$$

This is stronger on this face than the global bound $E \geq 1/18$. It does not require $x\eta$ bounded. In particular it is not legitimate to infer $x\eta^2 h = O(1)$ from (LH2); the exact available bound is $x\eta^2 h = O(x\eta) + O(1)$.

The corresponding opening-cleared leading source weights are

$$C_0 = \frac{1}{2L(1 - r)}, \qquad A_0 = \frac{1 + L(1 - r)}{2Lr}, \qquad \alpha_0 = \frac{L}{2(1 - Lr)}. \tag{LH3}$$

The ratios of the actual weights to these positive expressions tend to one. In reciprocal Bob coordinates put $H_t = wZ$ and $P = uW$. The leading three-state quadratic has the exact identity

$$\begin{aligned} & C_0(P - Y)^2 + A_0Y^2 + \alpha_0(Y - H_t)^2 + \tfrac{1}{2}Y^2 - PH_t \\ & \quad = C_0\{P - Y - L(1 - r)H_t\}^2 + (A_0 + \alpha_0 + \tfrac{1}{2})\{Y - LrH_t\}^2. \end{aligned} \tag{LH4}$$

All radical or inverse coefficients in this identity are pure source. Its two roots vanish at the exact contact to leading order. The original m, D, β packets, the small corrections to (LH3), and every normal coupling still require an exact simultaneous allocation. Neither (LH2) nor (LH4) is a tensor-SOS certificate for that remaining source face.

Why the completed C2 IH finite block cannot be copied unchanged. The common contact geometry is still exact, but its source pivot hypotheses change. To make this failure concrete, take a positive pure-source infinitesimal s and the native rational source

$$L = \tfrac{1}{2}, \qquad x = s^2, \qquad \eta = s^{-1}, \qquad r_n = s, \qquad h_n = 4s, \qquad \theta = 1 - 4s. \tag{LH5}$$

All native strict gaps are positive for sufficiently small s. Opening-cleared substitution into the exact formulas gives

$$C \sim 1, \quad sA \sim \tfrac{3}{2}, \quad D/s^2 \sim \tfrac{3}{2}, \quad m/x^2 \sim 4, \quad \alpha \sim \tfrac{1}{4}, \quad \beta/s \sim \tfrac{1}{2}. \tag{LH6}$$

Thus this diagnostic also has an unbounded original coefficient; it is not removed merely by the all-coefficients-bounded argument.

Write $B = L$, $H = Br_n$, $r_g = h_n$, $z = x$, $y = B\eta$. In the C1 version of the IC7 scalar bookkeeping, the beta event adds $4\beta(1 - r_g)^2$ to its mixed coefficient K. Explicitly set

$$\begin{aligned} K_1 &= 4AH^2 + 8DHr_g + 4\alpha(1 - H)^2 + 4m(H + r_g - 1)^2 \\ &\quad + 2(H + r_g - 1)^2 + 4\beta(1 - r_g)^2, \\ k_1 &= K_1 - 4B^2 m/z^2 + 8AH^2/y. \end{aligned} \tag{LH7}$$

The same exact substitution yields $k_1 \to -1$. Consequently $k_1/(Bs)$ is unbounded, whereas the completed strong-C2 IH finite block requires this coefficient to be bounded in its sT normalization. The C2 premise $r_g/s \le c_s$ also fails: here $r_g/s = 4$ and $c_s = (1 + B - H)/B \to 3$. This exhibits the precise failed transfer premise. It is not a counterexample to positivity or a proof that a generic remaining source is nonempty; this individual rational curve has its separate inherited admission. A new beta-aware soft normalization or circuit is required before that uniform C2 argument can be applied.

Chapter 31

C1 with both L and η units

Component p686209b87229
Source: R13/advances/c1_lunit_eta_unit/eta_unit.tex
SHA256: 686209b8722928e9065b28ff8b0caeb4dcb38c543c4ff8ae1dc50ddf91744b07
Inventory use: LU1–LU12 and cyclic remote join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

31.1 C1 with both L and η units

Theorem 31.1. *For the original C1 source with $x \in \mathrm{Inf}_{>0}$ and both $L = x\zeta$ and η positive bounded units, the entire original parent is tensor-SOS for every independent projective testing pair. All original positive openings and literal $\beta = 0$ are included.*

The exact source formulas force D strong on this subdomain, so $D = 0$ is absent by source implication. No opening is removed by an extra hypothesis, and the proof does not invoke a table or atlas.

The source and the only testing strip left by four events. Use the exact shared geometry (LC1), and write $\rho = r_n$ for the native shape parameter and $r = 1 - \theta$ for the geometric one. Thus

$$B = L, \quad y = B\eta, \quad H = B\rho = B - yt, \quad t = (1-\rho)/\eta, \quad j = 1 + zr - t, \quad z = x \in \mathrm{Inf}_{>0}\,.$$

Here B, y, η are units. For the boundary coefficient formulas below set

$$P = 1 + B + y, \qquad P_4 = 1 + B(1-\rho), \qquad S = H + y = P - P_4. \tag{LU1}$$

All three quantities are positive units, uniformly through $\rho \to 0$, $1 - \rho \to 0$, $j \to 0$, and $1 - H \to 0$. The native formulas imply that D, C, A, α are strong. One forward cyclic relabeling sends these four consecutive events to C, A, α, m and sends testing coordinates to $(u', v') = (v, 1/u)$. The full-parent four-consecutive theorem (CE11)–(CE16) therefore covers every test except

$$u \in \mathrm{Inf}, \qquad v \text{ a bounded nonzero unit.} \tag{LU2}$$

The cycle is the inverse of (CE9)–(CE10), so its literal endpoint scope is the original homogeneous parent's scope.

It remains to prove (LU2). All roots and coefficients used below are valid for both testing signs, including literal $u = 0$.

A bounded full-parent remainder with every opening retained. Use the graph, harmonic roots and gauge (LII2)–(LII8), with $k = \sqrt{y}$ now a source unit. The bounded source data are

$$c = Cyt, \quad d_p = Dyj, \quad \mu = m/z^2, \quad \sigma = \alpha(1-H), \quad \tau = \beta(1-r), \quad A_H = AH.$$

They are bounded; c, d_p, μ, σ, A_H are units. In residue, with the boundary symbols of (LU1), their exact values are

$$c = \frac{S}{2P}, \quad d_p = \frac{BP_4}{2P}, \quad \mu = \frac{Sr(1+y)}{2B^2P}, \quad \sigma = \frac{BS}{2P}, \quad A_H = \frac{yP_4}{2P}, \quad \tau = \frac{S}{2P} - \frac{1-r}{2}. \tag{LU3}$$

These are residue identities, not substitutions of source and testing fields. They are obtained after multiplying away every possibly small opening in the exact native formulas. Since $\tau \geq 0$, $r \geq P_4/P$ in residue and is a positive unit. Also $t + j \sim 1$, so $\ell = y(C + D)$ has a positive lower bound, $q = D/(C + D)$ is bounded in $[0, 1]$, and $g = yCD/(C + D)$ is a positive unit.

Choose positive bounded caps $\ell_* \leq \ell$, $A_* \leq A$, and $\alpha_* \leq \alpha$. Choose a bounded cap $0 \leq \beta_* \leq \beta$ as specified below. Retain the five contact-centered positive packets

$$\begin{gathered} 4(\ell - \ell_*)N^2, \qquad 4CV\eta_t^2, \qquad 4(A - A_*)D_uD_v\eta_t^2, \qquad 4(\alpha - \alpha_*)D_u(v\eta_t - \zeta)^2, \\ 4(\beta - \beta_*)D_v(\xi + u\zeta + R/B)^2, \qquad \eta_t = k(N + qR_h), \quad R = v(U - z)W. \end{gathered} \tag{LU4}$$

The exact remainder is (LII6), with its n line deleted, its A packet replaced by

$$4A_*D_uD_v\eta_t^2 + \frac{8A_H}{B}D_uD_vuW\eta_t + \frac{4A_HH}{B^2}D_uD_vUW^2, \tag{LU5}$$

and with (LC4) plus $4\beta_*D_v(\xi + u\zeta + R/B)^2$ added. This specifies every coefficient of the full remainder. Only bounded source products occur, even at an unbounded original weight.

The same dependent-root gauge and exact contact equations give (LII9)–(LII13). The new constant scalar coefficients are

$$\frac{d_s + e_sV}{z^2} = \frac{4}{B^2}\left[\mu(B + zr)^2 + \left\{\frac{r^2}{2} - \tau(1 + r) + \beta_*\right\}(1 + V)\right]. \tag{LU6}$$

The support $P_\xi \in (U, z)$ and scalar degree at most two in U remain exact. Apply all of (LII15). As v is a unit, eliminate T by the exact pure-factor identity

$$T = (uk/v)R. \tag{LU7}$$

Its coefficient is a bounded infinitesimal tensor. Thus the entire remainder has a bounded Gram on (N, ζ, R_h, ξ, R). No source/testing ratio or mixed inverse has been introduced. We now verify its complete residue.

A three-square identity supplies all three state/angle normals. The following boundary computation is an exact rational identity before taking any opening limit. Define

$$\begin{aligned} C_0 &= \frac{\rho + \eta}{2(1 - \rho)P}, & A_0 &= \frac{\eta P_4}{2\rho P}, \\ D_0 &= \frac{P_4}{2(\eta - 1 + \rho)P}, & \alpha_0 &= \frac{B^2(\rho + \eta)}{2(1 - B\rho)P}. \end{aligned}$$

These may have unbounded openings; only the final capped identity below is used as a bounded Gram. For an indeterminate p, put

$$e = Y - \rho p, \qquad f = Z - vp/B, \qquad T_v = (V - y)p.$$

The entire leading three-state form is exactly

$$\begin{aligned} &4C_0D_v(p - Y)^2 + 4A_0D_vY^2 + (4D_0 + 2)(p - Y - vZ)^2 + (4\alpha_0 + 2)(vY - Z)^2 - 2D_vZ^2 \\ &\quad = \frac{4D_0}{B^2}(T_v + Be + Bvf)^2 + 4\alpha_0(f - ve)^2 + 4(C_0 + A_0 + \tfrac{1}{2})D_ve^2. \end{aligned} \tag{LU8}$$

This identity holds for every v, not only near the actual Bob angle. In the contact allocation at $z = u = 0$, $e = k(N + qR_h)$ and $T_v = -B(kR_h + v\zeta)$. Subtract the residue contributions of the retained packets (LU4). The harmonic identity then gives the complete three-normal block

$$\boxed{4\ell_* N^2 + 4gR_h^2 + 4(A_* + \tfrac{1}{2})D_v y(N + qR_h)^2 + 4\alpha_*\{\zeta - vk(N + qR_h)\}^2.} \tag{LU9}$$

It has strictly positive residue on (N, ζ, R_h) because ℓ_*, g, α_* are positive units. No normal has been suppressed. Every rational cancellation used to obtain (LU9) has bounded source products on both sides, so the identity extends continuously through every opening listed after (LU1).

The remaining two-root block and a finite cap. There is no finite residue coupling between these three roots and (ξ, R). Such terms carry u, z, or the coefficient in (LU7). For completeness, the exact first-normal support gives this block without dropping a term. To first order in (U, z) write $P_\xi = UA(V) + zZ(V)$; both coefficient functions are affine in V, and

$$Z(V) = \frac{D_v}{B}(-4r + 8\tau - 8\beta_*).$$

The exact contact equation gives $A(y) + Z(y) = 0$ in residue. After the first allocation of (LII15), the coefficient of ξR is therefore $A(V) - (V - y)(A' + Z') = -Z(V)$. Using (LU3) and (LU6), the complete block is

$$M_{\xi R} = \begin{pmatrix} 2D_v(1 + 2\beta_*) & \dfrac{2D_v}{B}(P_4/P + 2\beta_*) \\ \dfrac{2D_v}{B}(P_4/P + 2\beta_*) & \dfrac{2}{B^2}\{(P_4/P + 2\beta_*)D_v - (S/P)r(V - y)\} \end{pmatrix}. \tag{LU10}$$

Here all letters denote real residues when positivity is assessed.

If β is bounded, take $\beta_* = \beta$. The native relation $(1 - r)(2\beta + 1) = S/P$ follows from (LU3), and (LU10) has determinant

$$\frac{4SD_v}{B^2 P}(1 + 2\beta)r(1 + y) > 0. \tag{LU11}$$

Its first diagonal entry is positive, including $\beta = 0$. If β is unbounded, take β_* to be a sufficiently large positive real cap. Increasing it adds the positive square $4\beta_* D_v(\xi + R/B)^2$. On its kernel $\xi = -R/B$, the uncapped block equals

$$\frac{2S}{B^2 P}\{1 + ry + (1 - r)V\}R^2 > 0. \tag{LU12}$$

The elementary real two-by-two Schur argument therefore supplies a finite cap making (LU10) strictly positive. Unboundedness of the original β permits that cap.

Together (LU9)–(LU12) prove strict positive residue of the full five-root Gram. The bounded Gram lemma gives its tensor SOS; restoring (LU4) proves (LU2). The cyclic four-event complement completes every testing chart and original projective endpoint.

Verification scope. The checker verifies the exact three-square state identity, all three capped normals, both explicit two-root positivity formulas, the scalar coefficient and support assertions, and prints the full extracted first-normal and scalar polynomials. The exact source contact and monic contact divisions are the unchanged full-parent argument of (LII7)–(LII15), with the bounded modifications (LU4)–(LU6). This theorem closes the L-unit, η-unit subdomain of C1.4; the η-unbounded subdomain is not included.

Chapter 32

The q-infinitesimal C1.1 source at both small testing angles

Component `pc376284fc7ae`
Source: `R13/advances/c1_small/q_inf_local_completion.tex`
SHA256: `c376284fc7aed6d1b9b713584cdd9ca01450ffdf24be8a5982dee16ad8684346`
Inventory use: QI1–QI10 and complete restoration | QI1-QI10 | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

32.1 The q-infinitesimal C1.1 source at both small testing angles

Theorem 32.1. *On the original C1.1 source with $q = k^2\eta$ infinitesimal, the entire original parent is tensor SOS whenever both independent testing half-angles u, v are infinitesimal. All original opening rates and $\beta = 0$ are retained. The boundary $D = 0$ is empty here because D_b has residue $1/2$.*

This statement includes every state; it does not yet assert the ordinary or unbounded v testing charts.

Use the preceding source notation and put $c = \sqrt{q}$, $e = c/g$, $h = H/R$, and $E_* = h - v$. Work first with $Z = 1$ and $u, v, h > 0$. Define

$$A = ch, \quad B = cL/R, \quad C = uv/b, \quad D = qg, \quad \Gamma = (ABCD)^{1/4}, \quad d_i = A_i - \Gamma.$$

Then

$$\Gamma^2 = qhv, \qquad BC = ev^2h, \qquad uh = uE_* + bC. \tag{QI1}$$

Here $h = u(W + uY)/(Rb^2)$ is separated into a positive pure-source factor and a positive pure-testing factor. Every radical below is therefore legal; no mixed inverse or order comparison is used. The exact original inequality $\beta \geq 0$ gives $t = O(e^2)$, while t is bounded. Hence t/e and $\sqrt{t/e}$ are bounded.

Exact scalar, with the D packet restored. Write $U = u^2, V = v^2$, $D_u = 1 + U, D_v = 1 + V$, and $a_t = 1 - \rho_0 + (t/R)V$. At both state normals zero the entire original scalar is

$$ah^2 + b_*hv + c_*v^2 + P_LL^2 + 2D_uC^2 + 4\alpha_bg^2D_u, \tag{QI2}$$

where

$$\begin{aligned} a &= 4D_vR^2\{c_b(1-\rho_0)^2 + a_b\rho_0^2/D_u\} + 2b^2R^2\{a_t^2 + V\rho_0^2/D_u\} \\ &\quad + 4\alpha_bg^2b^4VR^2\rho_0^2/D_u + 4D_bb^4R^2a_t^2, \\ b_* &= -4R\{D_u(1 + tV/R) - U\rho_0\} - 8\alpha_bg^2b^2R\rho_0 - 8D_bb^2Ra_tD_u, \\ c_* &= 4D_bD_u^2. \end{aligned}$$

All these rational functions have bounded source coefficient arrays; only D_u and positive pure-source units occur in their denominators. The r openings cancel exactly as in (SC12). In particular

$$D_b = \frac{N_D}{2JP_d}, \qquad N_D = \frac{K(1+qg^2)(1+b^2r) - JP_d}{b^2}$$

has a polynomial numerator: the numerator before division vanishes identically at $b^2 = 0$, where $P_d = 1 + qg^2$. This is a finite monic division, so it introduces no unbounded source coefficient array. Also α_b/q^2, h_2/q, and h_b/q have explicit bounded arrays from their source formulas. These observations justify the q coquotients below uniformly through both source openings.

Use formal specialization at $q = 0$ only to construct finite polynomial coquotients; it is not a new physical source. Put $z = b^2$, $J_0 = 1 - z(1-r)$, and $S = 1 - r + tV$. The three specialized coefficients are

$$\begin{aligned} a_0 &= 2D_v\{(1+zr)(1-r) + r/D_u\} + 2z\{S^2 + Vr^2/D_u\} + 2z^2S^2/J_0, \\ b_0 &= -4\{D_u(1+tV) - Ur\} - 4zSD_u/J_0, \\ c_0 &= 2D_u^2/J_0. \end{aligned}$$

Define $\widetilde{b}_0 = b_0 + 4tD_uV$, and the bounded finite coquotients

$$X = a_0 + \widetilde{b}_0 + c_0 = UX_U + VX_V, \qquad Y = -\widetilde{b}_0 - 2c_0 = UY_U + VY_V. \tag{QI3}$$

For definiteness, first subtract the value at $U = 0$, divide by U, and then divide that value by V. Their two V coquotients are

$$X_V = 2 + 2zr(1-2t) + \frac{2z(1+zr)t^2V}{J_0}, \qquad Y_V = \frac{4zt}{J_0}. \tag{QI4}$$

Thus $X_V = 2 + \varepsilon_V$, $\varepsilon_V \in B_{\mathrm{Inf}}$; X_U, Y_U have bounded arrays. Likewise write $a = a_0 + qa_1$, $b_* = b_0 + qb_1$, $c_* = c_0 + qc_1$ by finite numerator division. All a_1, b_1, c_1 have bounded arrays. Direct source specialization gives

$$\overline{a_1 - c_1} = 2, \qquad \overline{b_1 + 2c_1} = -8, \qquad \overline{R^2P_L/q} = 2, \qquad \overline{4\alpha_bD_u/q^2} = 2. \tag{QI5}$$

The following is the exact scalar identity to be allocated:

$$\begin{aligned} (\mathrm{QI2}) = {} & (c_0 + qc_1)E_*^2 + 2D_uC^2 + X_U(uE_* + bC)^2 + Y_Uu(uE_* + bC)E_* \\ & + X_V(vh)^2 + Y_Vv(vh)E_* - 4(tv/e)D_uBC \\ & + (a_1 - c_1)A^2 + (b_1 + 2c_1)\Gamma^2 + (R^2P_L/q)B^2 + (4\alpha_bD_u/q^2)D^2. \end{aligned} \tag{QI6}$$

It uses only (QI1) and $qv^2 = 2\Gamma^2 - A^2 + qE_*^2$. In particular the apparently dangerous negative t term is already an infinitesimal-coefficient BC term, since v is infinitesimal.

Two extra contact squares and their finite first jets. Set

$$F_* = vh - gD, \quad R_F = \frac{\sqrt{g\Gamma}}{c}(\Gamma - D), \quad V_0 = cv, \quad R_E = \sqrt{\Gamma}(\sqrt{A} - \sqrt{V_0}).$$

All four roots vanish at actual contact. Exact multiplication gives

$$\begin{aligned} (vh)^2 &= F_*^2 + 2R_F^2 + 2gF_*d_4 + g^2D^2 - 4g^2\Gamma d_4, \\ c\Gamma E_* &= 2\Gamma d_1 - R_E^2, \\ \Gamma F_* &= -2g\Gamma d_4 + R_F^2/g. \end{aligned} \tag{QI7}$$

The root R_E receives a strict real square pivot from E_*^2:

$$qE_*^2 = (A - V_0)^2 = 4R_E^2 + (\sqrt{A} - \sqrt{V_0})^4. \tag{QI8}$$

Reserve this fraction q of the strict E_* pivot; the remaining coefficient still has positive real residue. The leading $2(vh)^2$ in (QI6), by (QI7), gives strict F_* and R_F pivots, with only infinitesimal quadratic corrections and the printed first-jet term. The remaining leading angular form is $2(\sum A_i^2 - 4\Gamma^2)$, with the strict finite toric square list from (SC16).

For precision, the only additional bare-Γ normal couplings left by (QI6), apart from ordinary toric first jets, have the form $\Gamma E_* a_* + \Gamma F_* b'_*$ with

$$a_* = bu(2X_U + Y_U) + Y_V vg, \qquad b'_* = 2\varepsilon_V g.$$

At contact $u = bc, v = cg$ these satisfy

$$a_{*0}/c = b^2(2X_{U0} + Y_{U0}) + Y_{V0}g^2 \in B_{\mathrm{Inf}}, \qquad b'_{*0}/g = 2\varepsilon_{V0} \in B_{\mathrm{Inf}}.$$

Their differences from contact have finite bounded monic Taylor arrays. The conductors (SC18) allocate those differences against E_*, F_* with infinitesimal coefficients. Apply (QI7) to their constant terms. What remains is a bounded infinitesimal toric quadratic plus $\Gamma \sum \mu_i d_i + \Gamma^2 \sigma$. The exact scalar value and three original first jets imply its contact value and first-jet balances. The same two finite divisions as in (SC17)–(SC20), or the Hadamard contact-lift identity, allocate it on the strict toric root list. No infinite series or unsupported ideal-membership assertion is used: (QI3), the q coquotients, and the two displayed contact coefficient ratios are the finite certificates needed for the additional roots. Every coefficient function in these Taylor steps depends only on u, v. Thus only the first two conductors in (SC18)–(SC20) are used; the third state-coordinate jet supplies the inactive Hadamard balance. No bound on $(W + uY)/R$ after Z normalization is assumed.

Both complete state normals. Use $\widehat{N} = \sqrt{H_0}N$, $H_0 = 4h_2/R + 4h_b/t$, as in the preceding uniform-normal lemma. Add back the centered D packet from (SC10), including its normals. Its scalar root is exactly

$$J_* = (T + b^2 tV)h - D_u v = E_* + b^2 c(1 - tg^2)A - buC + b^2 tvF_* + b^2 tvgD. \tag{QI9}$$

Its full root is $J_* + ubvgN - b^2 N_y/\sqrt{s}$. All additional hard coefficients are infinitesimal after the $\widehat{N}$ normalization.

For the first N normal in (SC14), its H term is a bounded infinitesimal multiple of $uh = uE_* + bC$; its other terms are infinitesimal multiples of B, C. For the harmonic normal, multiply its H coefficient by $\sqrt{s}$. After the exact opening cancellations, its value at $q = U = V = 0$ is zero. Finite numerator coquotients therefore give

$$\sqrt{s}\,[H\text{-coefficient in } \mathcal{L}_y] = qK_q + UK_U + VK_V \tag{QI10}$$

with bounded arrays. Use

$$qH = RcA, \quad u^2 H = Ru(uE_* + bC), \quad v^2 H = Rv(F_* + gD).$$

The two remaining terms of $\mathcal{L}_y$ have infinitesimal coefficients on C, D since α_b/q^2 is bounded. Thus both complete first normals, including (QI9), are infinitesimal combinations of E_*, F_*, A, B, C, D. Their exact contact zeros make the coefficient sum of A, B, C, D zero at contact. One finite monic division and (SC18) allocate these on the same strict square list. The hard block has residue $\mathrm{diag}(1, 4)$ and all its other entries have zero residue. This proves a bounded positive residue Gram for the entire original parent, after restoring all retained positive packets.

States and signs. The restriction $Z = 0$ is positive by the original pure-testing state restriction. For $Z \neq 0$, divide the state by the pure-testing Z; its size is immaterial, since every coefficient array above depends only on u, v. On the negative-core chart, exact sign gauges give $u, v > 0$ and the inherited pure-testing Schur identity forces $W + uY > 0$. Hence $h > 0$, as required. The other core-sign chart and literal testing axes are direct original exits. Complex states split into real and imaginary parts. This completes the asserted small-u, small-v theorem.

Chapter 33

A uniform hard normal for the q-infinitesimal branch

Component `p7d43f3fccb7d`
Source: `R13/advances/c1_small/q_inf_normal_advance.tex`
SHA256: `7d43f3fccb7df6ddf6dbf56bd99ee3f0dcb0fa6417aec7de68b9b76c63b4a7ee`
Inventory use: entire exact normal construction, including H0 lower bound | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

33.1 A uniform hard normal for the q-infinitesimal branch

This intermediate lemma by itself is not an admission of the full q-infinitesimal C1.1 source. The following two completion sections supply the scalar allocation and remote joins. Keep (SC1)–(SC3), but do not discard the whole second paired packet. Its addition to the N^2 coefficient in (SC3) gives exactly

$$\frac{4}{R}(h_2 + h_b D_v) + \frac{4h_b M}{tR} D_v = H_0 + \frac{4h_b}{t} v^2, \qquad H_0 = \frac{4h_2}{R} + \frac{4h_b}{t}.$$

Retain the last summand as the positive square $4h_b(vN)^2/t$, and use the pure-source normalization

$$\widehat{N} = \sqrt{H_0}\, N.$$

The exact original inequality $\beta \geq 0$ gives $h_b/t \geq g^2/2$. The explicit formula for h_2 gives h_2/q a positive bounded unit, uniformly through both openings. Consequently $H_0 \geq c_0(q + g^2)$ for some positive real c_0. This argument also permits $\beta = 0$ literally. It does not require $e = \sqrt{q}/g$ or its reciprocal to be bounded.

The coefficient of $\widehat{N}^2$ from this source part is exactly one. Its remaining contribution $2g^2u^2v^2/H_0$ is bounded and infinitesimal on the small-angle testing chart. Moreover

$$\frac{g}{\sqrt{H_0}}, \quad \frac{\sqrt{q}}{\sqrt{H_0}}$$

are bounded. Since $h_b - h_2 = O(qg^2)$, the complete first normal (SC14), after this normalization, is an infinitesimal-coefficient linear combination of B, H, V for u, v infinitesimal. The first coefficient is bounded because

$$\frac{h_b D_v - h_2}{\sqrt{qH_0}} = O(g^2 + v^2),$$

and the other two coefficients are bounded by constant multiples of buv and v. The hard cross coefficient with N_y is likewise infinitesimal.

The issue left by this lemma is scalar and angular: H is unweighted in this statement, whereas the four-term circuit controls $\sqrt{q}H$. One must retain a strict D-based contact normal and allocate

every unweighted H coefficient on that normal and the circuit. The naive full extraction (SC10) fails in this branch: its raw P_H satisfies the exact formal specialization

$$P_H\big|_{q=0,\,v=0} = -\frac{2ru^2}{1+u^2},$$

so P_H/q need not be bounded. Thus the corrected hard-normal normalization removes the former $g/\sqrt{q}$ obstruction. The later protected-square lift establishes the missing scalar allocation.

Chapter 34

The other Bob testing charts for q infinitesimal

Component `pd2ecc6ac1891`
Source: `R13/advances/c1_small/q_inf_remote_completion.tex`
SHA256: `d2ecc6ac1891279167b66f70a02259edbbec8f2f424b13387bbb0042e5a390b4`
Inventory use: QR1–QR3 and all homogeneous endpoint frames | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

34.1 The other Bob testing charts for q infinitesimal

Theorem 34.1. *The original C1.1 source with $q = k^2\eta$ infinitesimal is tensor SOS at every independent test. Together with the q-unit theorem this closes the entire original C1.1 source.*

The inherited C, A-large admission already handles u not infinitesimal. The preceding local theorem handles u, v infinitesimal. Here are the two remaining v charts. Every positive opening rate and the literal permitted coefficient boundary is kept.

Ordinary v: the full normal residue is strict. Let v be a positive bounded unit, use $Z = 1$ on the negative-core chart, and retain the roots of (QI1). Normalize N by H_0 as before, but now restore the positive packet $4h_b(vN)^2/t$. Write

$$\tau = \operatorname{res} t, \qquad \rho = \operatorname{res}(t/e^2), \qquad a^2 = \frac{\tau}{2(1+\tau)}, \quad \gamma^2 = \frac{\rho}{2(1+\tau)}.$$

The original $\beta \geq 0$ equation implies $0 \leq \rho \leq 1$; $0 \leq \tau \leq 1$ follows from $M = R - t > 0$ and $R \sim 1$. The entire residue on $(\widehat{N}, B, C)$ has diagonal entries

$$1 + \frac{v^2}{1+\tau}, \qquad 2(1+\tau v^2), \qquad 2,$$

and polynomial cross coefficients

$$4av^2, \qquad -4\gamma v, \qquad -4\sqrt{\tau\rho}\, v.$$

Its exact square decomposition is

$$\begin{aligned} &2(C - \gamma v\widehat{N} - \sqrt{\tau\rho}\, vB)^2 + \widehat{N}^2 + 2B^2 \\ &\quad + (1-\rho)v^2 \left(\frac{\widehat{N}}{\sqrt{1+\tau}} + \sqrt{2\tau}\, B \right)^2 . \end{aligned} \tag{QR1}$$

Thus it is strictly positive definite, also at $\rho = 1$, the possible $\beta = 0$ endpoint, and at $\tau = 0$ or 1. The other residue blocks are exactly

$$2E_*^2 + 2(vh)^2 + 4D_v N_y^2.$$

There are no nonzero leading cross entries with these blocks.

For completeness, the boundedness check uses the entire raw scalar (QI2) and both full normal formulas (SC14), with (QI9) restored. Terms containing uh use $uh = uE_* + bC$; all their coefficients are infinitesimal. The N–B and N–C coefficients with nonzero residue are precisely those retained in (QR1). The harmonic normal has infinitesimal coefficient on h for bounded v: after the opening cancellation its coefficient lies in (b^2, q, u^2) with bounded arrays on this chart. Its other coefficients are infinitesimal as well. All remaining source-constant terms and weak monomials enter the same strict Gram through the exact bounded pure-testing identities

$$h = (vh)/v, \qquad 1 = ((vh) - vE_*)/v^2.$$

For example $A = ch$ and $D = qg$ are infinitesimal-coefficient combinations of these two roots. Consequently the actual full Gram has the positive residue just displayed. Restore all retained packets. The same sign, $Z = 0$, and complex-state joins as in the local theorem apply; no restriction on the size of the normalized state was used.

Unbounded v: a five-root endpoint frame. Put $w = 1/v$, so w is infinitesimal, and multiply the original parent by w^2. The exact source formulas give positive real constants μ, d_0 with

$$C/d \geq \mu/b^2, \qquad A/d \geq \mu/b^2, \qquad D/d \geq d_0 b^2.$$

For instance sufficiently small fixed constants below $1/4$ work, since $b^2C/d, b^2A/d$ are uniformly bounded below and $D/(db^2) \sim 1/2$. Retain all excesses and the entire m, α, β packets. No assumption on e is needed. Use the five roots

$$P = uW/b, \quad Q = Y/b, \quad X, \quad J_b' = bZ, \quad V_b = uwZ.$$

The homogenized signed core is exactly

$$\begin{aligned} K = {} & V_b^2 - 2buV_bQ - 2wPJ_b' + 2wV_bX + b^2w^2P^2 + 2buwPX - 2b^2w^2PQ \\ & + (D_u + w^2)(X^2 + b^2Q^2) - 2buwXQ. \end{aligned} \tag{QR2}$$

The three selected event packets are

$$4\mu D_w(P - Q)^2 + 4\mu D_u D_w Q^2, \qquad 4d_0(b^2wP + buX - b^2wQ - J_b')^2. \tag{QR3}$$

Combining these with $2K$, the bounded Gram has strictly positive residue

$$4\mu\{(P - Q)^2 + Q^2\} + 4d_0(J_b')^2 + 2X^2 + 2V_b^2.$$

All other displayed coefficients have zero residue. This is a full state-quadratic certificate, not a restriction to a contact graph. It includes the homogeneous endpoint $w = 0$ directly and uses no inverse involving independent source and testing variables.

The infinitesimal, unit, and unbounded v charts exhaust all tests remaining after the inherited C, A admission. Hence the entire q-infinitesimal C1.1 source, and consequently C1.1 itself, is closed.

Chapter 35

C1.1 with $k^2\eta$ a unit: a complete contact certificate

Component `p95350f69a353`
Source: `R13/advances/c1_small/q_unit_completion.tex`
SHA256: `95350f69a35347503869de108b3124c5ad72b0daadde632ce44dcbb244e890df`
Inventory use: SC10–SC20 and complete full-normal/contact proof | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

35.1 C1.1 with $k^2\eta$ a unit: a complete contact certificate

Theorem 35.1. *On the original C1.1 source, suppose $q = k^2\eta$ is a positive bounded unit. The entire original parent is tensor SOS at every independent projective test and every real or complex state. All positive opening rates and the permitted literal boundary $D = 0$ are included.*

The proof uses the exact full quadratic (SC3), not its leading scalar. The q-infinitesimal part of C1.1 is not included in this theorem.

One more centered packet. Use the variables of (SC1)–(SC5), and put $s_D = J/R$. Retain the additional nonnegative square

$$4D_b(J_b + s_D H)^2. \tag{SC10}$$

It vanishes at actual contact, where $H = RvZ$ and $J_b = -JvZ$. Its removal replaces $4D_bJ_b^2$ in (SC3) exactly by

$$-8D_b s_D H J_b - 4D_b s_D^2 H^2.$$

In particular it removes the entire standalone $(vZ)^2$ term. This step allows $D_b = 0$ literally and does not divide by D_b.

Set $Z = 1$ on the remaining state chart. Put $c = \sqrt{q}$ and $w = (W + uY)/R$, and use the four separated positive monomials

$$A = \frac{cuw}{b^2}, \quad B = \frac{cvw}{bg}, \quad C = \frac{uv}{b}, \quad D = c^2 g, \quad \Gamma = (ABCD)^{1/4}. \tag{SC11}$$

Their exact contact is $(u, v, w) = (k, cg, bg)$, with $k = bc$, and $A = B = C = D$ there. Notice that $H = RA/c$, $L = RB/c$, $V = C$, $g = D/q$, and $Hv = R\Gamma^2/q$.

Here are explicit formulas for the entire scalar at $N = N_y = 0$, after (SC10). Write $a_t = 1-\rho_0+pv^2$. Define

$$\begin{aligned} P_H &= 4D_v\{c_b(1-\rho_0)^2 + a_b\rho_0^2/D_u\} + 2b^2\{a_t^2 + v^2\rho_0^2/D_u\} \\ &\quad + 4\alpha_b g^2 b^4 v^2 \rho_0^2/D_u - 8D_b s_D b^2 a_t - 4D_b s_D^2, \\ P_L &= 4[h_2(1-p) + h_b p D_v]/R, \end{aligned}$$

$$P_{Hv} = -4[D_u(1+pv^2) - u^2\rho_0] - 8\alpha_b g^2 b^2 \rho_0 + 8D_b s_D D_u.$$

The exact scalar is

$$f = f_A A^2 + f_B B^2 + f_C C^2 + f_D D^2 - f_\Gamma \Gamma^2, \tag{SC12}$$

where

$$f_A = R^2 P_H/q, \quad f_B = R^2 P_L/q, \quad f_C = 2D_u, \quad f_D = 4\alpha_b D_u/q^2, \quad f_\Gamma = -RP_{Hv}/q.$$

These are rational functions of u, v over bounded source coefficients, with only positive-unit denominators. For example,

$$c_b(1-\rho_0)^2 = \frac{K(1+b^2 r)E(1-r)}{2P_d R^2}, \qquad a_b \rho_0^2 = \frac{P_4 K^2 r}{2P_d R^2}.$$

Thus neither r opening survives in their denominators. For u, v, b, g infinitesimal their residues are

$$\overline{f}_A = \overline{f}_B = \overline{f}_C = \overline{f}_D = 2, \qquad \overline{f}_\Gamma = 8. \tag{SC13}$$

This uses the exact source residues in (SC2), including $\overline{D}_b = (1-2\overline{q})/2$. It does not assume a positive lower bound on D_b.

Here is an explicit finite denominator certificate used below. Every function just displayed is $P(u,v)/(1+u^2)$ for a polynomial P with bounded pure-source coefficient array. For a monomial use

$$x^m - x_0^m = (x - x_0) \sum_{j=0}^{m-1} x^{m-1-j} x_0^j.$$

Applying this identity to each monomial supplies finite first and second difference arrays with bounded source coefficients. For the denominator use only the exact identity

$$\frac{1}{1+u^2} - \frac{1}{1+k^2} = -\frac{(u-k)(u+k)}{(1+u^2)(1+k^2)}.$$

Thus the first and second finite Taylor arrays about (k, cg) are bounded and use only pure-testing $1+u^2$ and pure-source $1+k^2$ denominators, both units. The same construction after $u = U^2, v = V^2$ gives the half-variable arrays below. This is a finite polynomial identity construction, not an analytic series.

Every first normal and the complete hard block. Let $\delta_\rho = \rho_0 - \rho$. The full first-normal coefficients of the remaining quadratic, with no normal suppressed, are

$$\begin{aligned}
\mathcal{L}_N &= \frac{8}{R}(h_b D_v - h_2)L + 4buvg(a_t - 2D_b s_D)H - 4D_u vgV, \\
\mathcal{L}_y &= 8D_v\sqrt{s}\,\delta_\rho H \\
&\quad + \frac{H}{\sqrt{s}}\left[-\frac{8D_v a_b u^2 \rho_0}{D_u} + 8D_b s_D b^2 + \frac{8\alpha_b g^2 b^4 v^2 \rho_0}{D_u} - 4b^2 a_t + \frac{4b^2 v^2 \rho_0}{D_u}\right] \\
&\quad + \frac{4bu}{\sqrt{s}}V - \frac{8\alpha_b g b^2 v}{\sqrt{s}}(gZ).
\end{aligned} \tag{SC14}$$

Each is a bounded infinitesimal-coefficient linear combination of A, B, C, D. At actual contact it is zero exactly, since the source is a stationary zero and every removed packet vanishes there. The coefficients of N^2, N_y^2, NN_y are respectively

$$\begin{aligned}
H_{NN} &= 4(h_2 + h_b D_v)/R + 2g^2 u^2 v^2, \\
H_{yy} &= 4D_v\left(1 - \frac{a_b u^2}{D_u s}\right) + \frac{4\alpha_b g^2 b^4 v^2}{D_u s} + \frac{2b^2}{s}\left(1 + \frac{v^2}{D_u}\right), \\
H_{Ny} &= -4buvg/\sqrt{s}.
\end{aligned} \tag{SC15}$$

Thus the hard Gram has residue $\mathrm{diag}(4q, 4)$. In (SC15), H_{Ny} denotes the polynomial coefficient of NN_y, not the off-diagonal matrix entry.

A finite geometric-mean square lift. Here is a detailed lift for the particular raw module (SC12). It is not a claim about arbitrary contact-vanishing polynomials. Define $\alpha_i = A_i - \Gamma$, with $(A_1, A_2, A_3, A_4) = (A, B, C, D)$, and put

$$T_4 = \sum_i A_i^2 - 4\Gamma^2.$$

A fixed positive Gram on the following finite roots represents T_4:

$$\alpha_i, \qquad Q_{ij|kl} = \sqrt{A_i A_j} - \sqrt{A_k A_l}, \qquad \sqrt{\Gamma}(\sqrt{A_i} - \sqrt{A_j}), \qquad \sqrt{\Gamma}((A_i A_j)^{1/4} - (A_k A_l)^{1/4}). \tag{SC16}$$

Use all four α_i, all three pair partitions, and all six unordered differences in the third group. To verify strictness, average positive multiples of the symmetric circuit decomposition in (SC9) and

$$T_4 = \sum_i \alpha_i^2 + 2\Gamma\left(\sum_i A_i - 4\Gamma\right).$$

over the three decompositions

$$\sum_i A_i - 4\Gamma = (\sqrt{A_i} - \sqrt{A_j})^2 + (\sqrt{A_k} - \sqrt{A_l})^2 + 2((A_i A_j)^{1/4} - (A_k A_l)^{1/4})^2.$$

The additional differences $A_i - A_j$ in the first decomposition are $\alpha_i - \alpha_j$ and give nonnegative corrections to the already strict α block.

Write $\epsilon_i = f_i - 2$, $\epsilon_\Gamma = f_\Gamma - 8$, and $\sigma = \sum_i \epsilon_i - \epsilon_\Gamma$. Then

$$f - 2T_4 = \sum_i \epsilon_i \alpha_i^2 + 2\Gamma \sum_i \epsilon_i \alpha_i + \Gamma^2 \sigma. \tag{SC17}$$

All ϵ have zero residue. The exact contact value implies $\sigma(k, cg) = 0$. Clear the pure-testing denominator D_u if needed; monic division gives $\sigma = (u-k)\sigma_u + (v-cg)\sigma_v$ with bounded coefficients. For the monomials (SC11), the exact conductors are

$$\begin{aligned} \Gamma(u-k) &= \sqrt{uk}\, Q_{13|24}, \\ \Gamma(v-cg) &= \sqrt{vcg}\, Q_{23|14}, \\ \Gamma(w-bg) &= \sqrt{wbg}\, Q_{12|34}. \end{aligned} \tag{SC18}$$

Thus the last term in (SC17) becomes Γ times a linear combination of the Q roots with infinitesimal bounded coefficients. The exact identity

$$Q_{ij|kl} - \tfrac{1}{2}(A_i + A_j - A_k - A_l) = -\tfrac{1}{2}(\sqrt{A_i} - \sqrt{A_j})^2 + \tfrac{1}{2}(\sqrt{A_k} - \sqrt{A_l})^2 \tag{SC19}$$

changes these terms into infinitesimal quadratic corrections on (SC16) and $\Gamma \sum_i \mu_i \alpha_i$, where each μ_i is infinitesimal and is rational in $\sqrt{u}, \sqrt{v}$ with bounded coefficient arrays and positive-unit denominators.

The exact three contact jets imply that the four source values $\mu_i(k, cg)$ coincide. Indeed the differential vectors of α_i/Γ at contact are the four rows

$$(1/2, -1/2, 1/2), \quad (-1/2, 1/2, 1/2), \quad (1/2, 1/2, -1/2), \quad (-1/2, -1/2, -1/2)$$

in logarithmic coordinates $(u/k, v/(cg), w/(bg))$. Their sole linear relation is the sum of all four rows. Cancelling the nonzero source factors in this equality establishes coincidence; no inverse of a small number enters a Gram coefficient. Call the common infinitesimal value μ_*. The common term is

$$\mu_* \Gamma \sum_i \alpha_i = \mu_* \Gamma \left(\sum_i A_i - 4\Gamma\right),$$

an infinitesimal correction on the explicitly displayed variance roots. For each remaining $\mu_i - \mu_*$, monic division in $\sqrt{u}, \sqrt{v}$ has bounded coquotients. Apply the half-variable conductors

$$\begin{aligned} \Gamma(\sqrt{u} - \sqrt{k}) &= (uk)^{1/4}\sqrt{\Gamma}((AC)^{1/4} - (BD)^{1/4}), \\ \Gamma(\sqrt{v} - \sqrt{cg}) &= (vcg)^{1/4}\sqrt{\Gamma}((BC)^{1/4} - (AD)^{1/4}). \end{aligned} \tag{SC20}$$

Their coefficient factors are infinitesimal. Multiplying by the remaining α_i proves that the entire scalar correction in (SC17), not merely its low-order jet, is a bounded zero-residue quadratic form on (SC16). This uses only two finite monic divisions. The boundedness of their coefficient arrays follows from the explicit rational functions (SC12), whose test denominators are powers of D_u, and from the displayed square-root substitutions; bounded function values alone are not being used as a substitute for this fact.

The same argument for a first normal is shorter. If $\mathcal{L} = \sum_i a_i A_i$, write $\mathcal{L} = \sum_i a_i\alpha_i + \Gamma\sum_i a_i$. Its exact contact zero gives $\sum_i a_i(k, cg) = 0$; one monic division and (SC18) allocate it on (SC16), with infinitesimal coefficients. Apply this to both complete expressions (SC14). Together with (SC15), this yields a bounded strictly positive-residue Gram for the entire original remainder. Restore (SC10), the paired packets, and all positive pure-source and pure-testing normalizations.

Exhaustion and legal radicals. For q a unit, $e^2 = q/g^2$ is unbounded. The exact adjacent-event rotations and the W-unit state argument (SC8) therefore reduce every unadmitted negative-core test to u, v, W infinitesimal and Z a unit. Literal axes are direct original-core exits. Use exact sign gauges and normalize $Z = 1$. The inherited pure-testing Schur identity forces $W + uY > 0$ on the remaining negative-core chart: with $w_t = (W + uY)/D_u$, the eliminated scalar is

$$\frac{uvD_uD_v}{\mathcal{D}}(D_uuvw_t^2 - 2D_uD_vw_t + uv), \qquad \mathcal{D} = u^2v^4 + u^2v^2 + u^2 + v^4 + 2v^2 + 1 > 0,$$

which is nonnegative when $w_t \le 0$; its eliminated two-square matrix is strictly positive. Hence $w = (W + uY)/R > 0$. Every monomial in (SC11), and every product under a radical in (SC16)–(SC20), factors into a positive pure-source quantity and a positive pure-testing quantity. The occurrence of R in w is a positive pure-source unit, not a mixed denominator. All test coefficients remain bounded because u, v, w are infinitesimal. If the general toric contact lemma is used, its third coordinate is literally the pure-testing $W + uY$, with pure-source contact Rbg; this direction is inactive in every coefficient function. Complex state evaluations split into real and imaginary parts. This supplies every independent test and proves the theorem.

Audit scope. The parent and the scoped toric-lift reviewer independently reviewed the finite lift, bounded rational arrays, full normal block, legal radicals, and joins within the same agent system. The thirty-nine exact checks verify the displayed whole-polynomial and source identities. A further six exact polynomial checks verify the actual source contact value, all three scalar first jets, and both full normal contact zeros directly from (NC1)–(NC4). Their explicit denominator-cleared numerators are printed in `check_q_unit_contact.py`; they confirm the original stationary-zero premise without sampling. No numerical table is a premise of this argument.

Chapter 36

C1.1: an opening-uniform full normal and a new angular circuit

Component `p9afbe73899af`
Source: `R13/advances/c1_small/small_u_harmonic_reduction.tex`
SHA256: `9afbe73899afc5182fd991d7b285b98dae466d7ecca55667cf04a3e5e251a889`
Inventory use: SC1–SC9; exact source cancellation, full-parent reduction and remote/state proof | SC2-SC20 | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

36.1 C1.1: an opening-uniform full normal and a new angular circuit

This section concerns the original C1.1 source, not an enlarged family. All coefficients are divided by $d > 0$. Put

$$b = \eta^{-1/2}, \quad g = \sqrt{\zeta}, \quad q = k^2\eta, \qquad x = b^2 q, \quad a = b^2(1-r). \tag{SC1}$$

Here b, g are positive infinitesimals, $q > 0$ is bounded, and $\overline{q} \le 1/2$. The new bounded two-normal reduction below is asserted for q a positive unit. The exact polynomial identities remain valid more generally, but their indicated boundedness argument would not control $g/\sqrt{q}$ when q is infinitesimal. No all-tests closure is inferred from the reduction.

Keep $K, R, T, M, J, E, Q, P_4, P_d, P_\alpha$ from (NC1)–(NC2), and set

$$\begin{aligned} c_b &= b^2 C/d = \frac{K(1+b^2r)E}{2(1-r)P_d}, & a_b &= b^2 A/d = \frac{P_4}{2rP_d}, \\ D_b &= D/(db^2), & \alpha_b &= \alpha/(db^2g^2), \\ h_2 &= \frac{Kq(1+b^2r)(1-tg^2)(1+qg^2)}{2P_d}, & h_b &= \frac{Kq(1+b^2r)[1+(1-t)g^2]}{2P_d}. \end{aligned}$$

Thus $m/d = h_2/(g^2M)$ and $\beta/d + 1/2 = h_b/(g^2 t)$. The exact relation

$$h_b - h_2 = \frac{Kq(1+b^2r)Eg^2}{2P_d} \tag{SC2}$$

removes the apparent opening dependence from the first normal of the paired m, β currents. The quantities D_b, α_b, h_2, h_b are bounded; α_b, h_2, h_b are units in the stated q-unit domain. Their residues are respectively $(1-2q)/2, q^2/2, q/2, q/2$, with bounded source letters on the right understood at residue.

Exact full-parent reduction. For bounded affine tests define

$$Q_b = (W + uY)/b, \quad Y_b = Y/b^2, \quad p = t/R, \quad N = \frac{(X+uZ)/b - pvQ_b}{g},$$

$$H = uQ_b/b, \quad L = vQ_b/g, \quad V = uvZ/b, \quad B_b = gN + pvQ_b, \quad D_u = 1 + u^2, \quad D_v = 1 + v^2.$$

All roots use only pure-source inverses. Retain the two nonnegative original packets

$$\frac{4h_2 t}{MR}N^2, \qquad \frac{4h_b M}{tR}D_v N^2.$$

The complete original parent divided by db^2, minus these packets, is exactly

$$\begin{aligned} G &= \frac{4}{R}\{(h_2 + h_b D_v)N^2 + 2(h_b D_v - h_2)LN + [h_2(1-p) + h_b p D_v]L^2\} \\ &\quad + 4D_v\{c_b(H - D_u Y_b)^2 + a_b D_u Y_b^2\} + 4D_b J_b^2 + 4\alpha_b g^2 D_u(vb^2 Y_b - Z)^2 + 2K_b, \qquad \text{(SC3)} \\ J_b &= b^2(H - D_u Y_b) + ubvB_b - D_u vZ, \\ K_b &= b^2(H - D_u Y_b)^2 + 2buv(H - D_u Y_b)B_b + u^2 v^2 B_b^2 \\ &\quad + D_u v^2 b^2 Y_b^2 + D_u V^2 - 2D_u V(Q_b + vB_b) + 2D_u u^2 vZY_b. \end{aligned}$$

These are entire quadratic polynomials in the state; no restriction of a state normal has been made. In particular, (SC3) is a direct input for collecting both complete first normals and their full hard quadratic, rather than an identity solely on a contact graph.

A harmonic normal removes both r openings. Put

$$s = c_b + a_b, \quad \rho = c_b/s, \quad h = c_b a_b/s, \quad \rho_0 = Kr/R, \qquad N_y = \sqrt{s}\,(D_u Y_b - \rho_0 H).$$

Then the complete C, A line in (SC3), divided by $4D_v$, equals

$$\left[N_y + \sqrt{s}(\rho_0 - \rho)H\right]^2 + hH^2 - \frac{a_b u^2}{D_u}\left(\frac{N_y}{\sqrt{s}} + \rho_0 H\right)^2. \qquad \text{(SC4)}$$

The inverse $1/D_u$ is pure testing, and every coefficient in (SC4) is bounded uniformly through either $r \to 0$ or $r \to 1$. Indeed s is bounded below, h is a positive unit, and a_b/s, $a_b\rho_0/\sqrt{s}$ and $a_b\rho_0^2$ are bounded. There is no mixed harmonic denominator.

Here is the exact source cancellation that also centers the normal without an opening loss. Define

$$S_0 = Kr(1 + b^2 r)E + (1 - r)P_4,$$

$$T_0 = q^2(1 - tg^2)^2 + qrg^2 + 2qtg^2 - qg^2 - 2q + r.$$

Then

$$\rho = \frac{Kr(1 + b^2 r)E}{S_0}, \qquad Kr - \rho R = -\frac{b^2 r(1 - r)KT_0}{S_0}, \qquad h = \frac{K(1 + b^2 r)EP_4}{2P_d S_0}. \qquad \text{(SC5)}$$

Both S_0 and the other denominators are positive units. Also $s \asymp [r(1-r)]^{-1}$, so $\sqrt{s}(\rho_0 - \rho)$ is infinitesimal without a lower bound on either opening. At the actual contact,

$$(u, v) = (b\sqrt{q}, \sqrt{q}g), \quad Q_b = RgZ, \quad Y_b = r\sqrt{q}gZ,$$

and both N, N_y vanish exactly. Thus (SC3)–(SC5) remove every unbounded source weight while retaining the actual contact and every state normal.

The leading angular circuit. For u, v infinitesimal, the leading scalar after the harmonic state normal has been removed is twice

$$(1 - q)H^2 + qL^2 + V^2 + (1 - 2q)(vZ)^2 + q^2(gZ)^2 - 2VQ_b.$$

Here the leading relation $H(vZ) = VQ_b$ is exact for the displayed roots. It splits as

$$(1 - 2q)(H - vZ)^2 + qH^2 + qL^2 + V^2 + q^2(gZ)^2 - 4qVQ_b. \qquad \text{(SC6)}$$

The final four positive terms form one AM–GM circuit, since

$$(\sqrt{q}H)(\sqrt{q}L)V(qgZ) = (qVQ_b)^2.$$

On the positive negative-core chart its radical factors separate between the source and testing fields. This proves positivity of the leading scalar, including $q = 1/2$, where its separate binomial drops out. The leading hard normal coefficients are $4q$ on N and 4 on N_y. This organizing form alone is not yet a certificate for all the higher terms in (SC3).

A useful exact conductor can be stated without comparing any source and testing rate. Normalize $Z = 1$ on its pure-testing state chart, put $c = \sqrt{q}$, $w = (W + uY)/R$, and now use the contact-exact roots

$$H_0 = uw/b^2,\ L_0 = vw/(bg),\ V_0 = uv/b,\ D_0 = c^2 g, \quad A_0 = cH_0,\ B_0 = cL_0,\ C_0 = V_0.$$

Their contact is $(u, v, w) = (bc, cg, bg)$. Set

$$E_0 = H_0 - v, \quad S_v = A_0 - D_0 - cE_0, \quad S_u = C_0 - D_0 - S_v, \quad S_w = B_0 - D_0 - S_v.$$

Direct multiplication gives

$$\begin{aligned} H_0(u - bc) &= (u - bc)E_0 + bS_u, \\ H_0(v - cg) &= (v - cg)E_0 + (v/c)S_v, \\ H_0(w - bg) &= (w - bg)E_0 + (bg/c)S_w. \end{aligned} \tag{SC7}$$

Only the positive-unit inverse $1/c$ occurs. These identities lift contact-vanishing coefficients times H_0 to the displayed root module with bounded coefficients on the small pure-testing chart. They do not alone prove a full scalar square lift: a first-jet syzygy among the redundant angular roots still has to be allocated, particularly when the coefficient of E_0^2 vanishes at $q = 1/2$.

Exact small-cost angular pivots. The four-term circuit has the exact symmetric decomposition

$$\sum_{i=1}^{4} A_i^2 - 4\sqrt{A_1A_2A_3A_4} = \frac{1}{3}\sum_{i<j}(A_i - A_j)^2 + \frac{2}{3}\sum_{\{ij\}|\{kl\}}(\sqrt{A_iA_j} - \sqrt{A_kA_l})^2,$$

where the second sum is over the three pair partitions. Use $(A_1, A_2, A_3, A_4) = (A_0, B_0, C_0, D_0)$ above and assume the positive chart $u, v, w > 0$. Then the following identities are exact:

$$\begin{aligned} c\sqrt{uw}\,E_0 &= \sqrt{uw}(A_0 - D_0) + b\sqrt{cg}(\sqrt{A_0D_0} - \sqrt{B_0C_0}), \\ \sqrt{vw}(u - bc) &= \frac{b^{3/2}}{\sqrt{c}}(\sqrt{A_0C_0} - \sqrt{B_0D_0}), \\ \sqrt{uw}(v - cg) &= b\sqrt{g/c}(\sqrt{B_0C_0} - \sqrt{A_0D_0}), \\ \sqrt{uv}(w - bg) &= \frac{b^{3/2}\sqrt{g}}{c}(\sqrt{A_0B_0} - \sqrt{C_0D_0}). \end{aligned} \tag{SC9}$$

The coefficient factors on the right are infinitesimal; all radicals separate between the source and testing fields. Consequently any fixed positive real multiples of the squares of the four left-hand roots can be inserted as additional positive Gram pivots at an infinitesimal cost in the displayed strict circuit Gram. To see this without a limiting assertion, write the four identities as $P = TZ$, with Z the displayed circuit roots and every entry of T in the bounded maximal ideal. Then $Z^TGZ = \tau P^TP + Z^T(G - \tau T^TT)Z$, and the second matrix retains the strictly positive real residue of G for every fixed real $\tau > 0$. This is a genuine full-rates Gram mechanism. It still requires showing that the entire first-normal and scalar remainders of (SC3) allocate on these weighted roots; the identities do not make that last membership automatic.

A quantified remote improvement when e is unbounded. The source identity $e^2 = q/g^2$ and (NI3) imply that m, β are both unbounded when e is unbounded. Three inverse-cycle rotations send this adjacent pair to C, A and transform tests as $(u, v) \mapsto (v, 1/u)$. Therefore the adjacent-event theorem covers all tests with v not infinitesimal. Combined with the previous C, A admission, only u, v infinitesimal remain in this e type. This includes every q-unit source considered above.

There is also a complete state-chart admission there. Normalize the pure-testing state by a coordinate of maximal absolute value. If W is a unit, cap C, A, m, β at a sufficiently large positive real μ, retain all excesses, and leave every other event as its original nonnegative square. At $u = v = X = Y = 0$, the quadratic part in (u, v, X, Y) is

$$2(uW-Y)^2+2(X+uZ)^2-4uvWZ+4\mu\{(uW-Y)^2+Y^2+(vW-X-uZ)^2+(X+uZ)^2\}. \quad \text{(SC8)}$$

The event part is strictly positive for every real $W \neq 0$ and bounded real Z: its zero equations give successively $Y = u = X = v = 0$. A sufficiently large finite μ therefore absorbs the entire signed quadratic core. The exact remainder has degree at least three in these four small roots, with bounded coefficients in W, Z, so it allocates with zero residue. This proves the entire original parent on this chart. If the core has negative sign and W is infinitesimal, then Z is a unit, since otherwise the core residue forces $X = Y = 0$, contrary to maximal-coordinate normalization. Thus the remaining state chart has $Z = 1$, W infinitesimal, and both testing half-angles infinitesimal. This is a full quantified narrowing, not closure of C1.1.

Completion and exact outstanding obligation. The following section supplies a finite scalar square lift and both complete first normals for the q-unit C1.1 subdomain, including $D = 0$. It uses a contact-centered D packet followed by the geometric mean conductor; an unweighted E_0^2 perturbation by itself is not controlled by the circuit uniformly at arbitrary relative rates. For q infinitesimal, the uniform normal and protected contact lift in Sections 32.1–34.1 resolve the additional $g/\sqrt{q}$ issue and all remote tests. No assumption of bounded e or $1/e$ is imposed. The original checker verifies thirty-nine whole-polynomial and source identities; the later modules include their own exact checks and scoped review.

Chapter 37

An interior small-L C1 source: a full three-circuit closure

Component `pe3ef590c01d6`
Source: `R13/advances/c1_small_interior/interior_closure.tex`
SHA256: `e3ef590c01d68ef36079d3024a86c5d75167e0b3ae52d689edad8ce6b1da744b`
Inventory use: IC7–IC8, IC11–IC19 and finite arrays | IC11-IC21 only | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

37.1 An interior small-L C1 source: a full three-circuit closure

Theorem 37.1. *Use the original physical native C1 source and all earlier intersections. Suppose $x = k^2$ is infinitesimal, and*

$$\eta,\ \zeta,\ r,\ 1-r,\ j_0 = \eta + r - 1 \quad \text{are positive bounded units.} \tag{IC1}$$

Then the full original parent is tensor SOS for every independent projective test and every state. No source/testing rate comparison is used. This is a subdomain of C14.3; it does not assert the remaining small-L source boundaries or C14.4.

Normalize $d = 1$, set $\rho = \eta + r$, $y = x\eta\zeta$, and retain all original packets. Here $L = x\zeta$ is infinitesimal. The source formulas (NC1)–(NC4) and (SB1)–(SB4) give

$$\begin{gathered} m/x \sim \rho/(2\zeta), \qquad \alpha/x^2 \sim \rho\zeta(1+\zeta)/2, \\ C = C_0 + O(x), \quad A = A_0 + O(x), \quad D = D_0 + O(x), \\ C_0 = \frac{\rho}{2(1-r)}, \quad A_0 = \frac{\eta}{2r}, \quad D_0 = \frac{1}{2j_0}. \end{gathered} \tag{IC2}$$

If the total scale is a unit, the earlier positive-total-scale theorem already applies. On the residual total-scale-infinitesimal intersection, all original coefficients except β are bounded, so β is unbounded. Its exact numerator implies

$$\theta = x\pi, \quad \pi \in \mathrm{Inf}_{>0}, \qquad \beta\theta/x = O(1), \quad \beta\theta^2/x^2 = O(\pi). \tag{IC3}$$

Thus $\beta = 0$ and $D = 0$ are absent on this particular residual source subdomain, by source implication. No permitted boundary is removed from a larger claimed theorem. All source denominators used below are products of positive bounded units from (IC1), M, J, Q, P_d, and their source specializations. The apparent x and x^2 denominators cancel by finite polynomial numerator division, as specified below.

The existing theorem in Section 28.1, equations (SB6)–(SB12), covers the original source x, L infinitesimal, ζ noninfinitesimal, and η bounded on the complement of u, v both infinitesimal, with every homogeneous endpoint. Its source domain contains (IC1). Literal testing axes are direct: the

half core is $D_v(X^2+Y^2)/2$ at $u=0$ and $\{(uW-Y)^2+(X+uZ)^2\}/2$ at $v=0$, and every other packet is nonnegative. The exact transformations $(u,Y,Z)\mapsto(-u,-Y,-Z)$ and $(v,X,Z)\mapsto(-v,-X,-Z)$ preserve the original parent, so work on the positive testing sign chart and restore all signs afterwards. For complex states apply the real identity to the real and imaginary parts. First work at $Z\neq 0$, divide the state by Z, and set $Z=1$; no bound on W is imposed. Afterward multiply the exact identity by the original Z^2. Its state-normal roots become the homogeneous linear forms displayed below and its scalar circuit roots become Z times the roots at $Z=1$. No Z inverse remains. The resulting identity is polynomial in the original state coordinates and hence also holds at $Z=0$.

A bounded-ratio frame retaining all three state normals. First suppose the pure-testing ratio $t=v/u$ is bounded, possibly infinitesimal. Define

$$\begin{gathered} E=\eta uW-v,\quad b=\frac{r(1-r)}{\eta(\eta+2r-1)},\quad N=Y-ruW-bE,\\ \xi=X+hu,\qquad N_x=\sqrt{\beta}\,\xi,\quad h=1-\theta. \end{gathered} \tag{IC4}$$

The state change is exact and uses only the pure-testing inverse u: $W=(v+E)/(\eta u)$, $Y=r(v+E)/\eta+bE+N$, $X=-hu+\xi$. Every normal vanishes at the actual source contact. Its full hard quadratic has the strictly positive diagonal residue

$$(E,N,N_x):\qquad \operatorname{diag}\left(\frac{\rho}{2\eta(\eta+2r-1)},\frac{\eta(\eta+2r-1)}{2r(1-r)j_0},1\right). \tag{IC5}$$

This is a statement about the complete original hard block. At bounded t, its remaining terms are bounded with zero residue: the potentially new mt^2 coefficient is infinitesimal, and normalizing ξ makes all its other hard entries infinitesimal. No state normal has been restricted or dropped.

The exact scalar, with the negative defect absorbed. Set $U=u^2, V=v^2$, $s=\sqrt{\eta\zeta}$, $k=\sqrt{x}$, and introduce the positive separated roots

$$A_t=uv,\quad B_t=\frac{kv^2}{su},\quad C_t=xs,\quad G=kv,\qquad F_v=V-y. \tag{IC6}$$

They satisfy $A_tB_tC_t=G^3$ and coincide at actual contact $(u,t)=(k,s)$. The pure-testing pole in B_t is inside the root; none of the coefficient arrays below has a negative power of t.

Use A_w, A_z, b_0, b_1 from (SN1)–(SN2), including the beta term. The scalar at $E=N=\xi=0$ is exactly

$$\mathcal{S}=\frac{VA_w}{\eta^2U}+A_z-\frac{2V}{\eta}(b_0+b_1U)=c_FF_v^2+f_AA_t^2+f_BB_t^2+f_CC_t^2-f_GG^2. \tag{IC7}$$

Here the complete coefficients are

$$\begin{aligned} c_F&=\frac{C(1-r)^2+Ar^2+(\alpha+\frac12)r^2+2mr}{\eta^2},\\ a_{UV}&=\frac{Ar^2}{\eta^2}+\frac{h^2}{2}+2Dh+\beta\theta^2-\frac{2b_1}{\eta},\\ a_{UUV}&=h^2(D+\tfrac12),\qquad a_{UVV}=\frac{r^2(m+A+\alpha+\frac12)}{\eta^2},\\ q_V&=\frac{C(1-r)^2+Ar^2+(D+\frac12)(1-r)^2}{\eta^2}+D-\frac{2b_0}{\eta},\\ q_U&=\alpha+(m+\beta+\tfrac12)\theta^2,\\ f_A&=a_{UV}+a_{UUV}U+a_{UVV}V,\qquad f_B=\frac{m\zeta}{\eta x},\\ f_C&=\frac{\alpha-c_Fy^2+q_UU}{\eta\zeta x^2},\qquad f_G=-\frac{q_V+2c_Fy}{x}. \end{aligned} \tag{IC8}$$

Thus the source-negative quadratic base has already been combined with the original m, α, and higher-angle terms. The residues are

$$c_F \sim \frac{\rho}{2\eta^2}, \qquad f_A, f_B, f_C \sim \lambda = \frac{\rho}{2\eta}, \qquad f_G \sim 3\lambda. \tag{IC9}$$

For example the leading scalar is exactly

$$\frac{\rho}{2\eta^2}(V-y)^2 + \frac{\rho}{2\eta}\left\{UV + \frac{xV^2}{\eta\zeta U} + \eta\zeta x^2 - 3xV\right\}. \tag{IC10}$$

The braces are the three-term AM–GM circuit $A_t^2 + B_t^2 + C_t^2 - 3G^2$, not a standalone quadratic PSD assertion.

Finite scalar Taylor arrays. Let $\epsilon_i = f_i - \lambda$ and $\epsilon_G = f_G - 3\lambda$, $h_c = \epsilon_A + \epsilon_B + \epsilon_C - \epsilon_G$. Equations (IC8) show that $h_c = h_{00} + aU + cV$ with bounded pure-source a, c. Exact contact value gives

$$h_c = a(u^2 - k^2) + c(u^2t^2 - k^2s^2). \tag{IC11}$$

For $d_u = u - k, d_t = t - s$, its finite second Taylor identity is

$$h_c = 2k(a + cs^2)d_u + 2k^2sc\,d_t + (a + ct^2)d_u^2 + 2kc(t+s)d_ud_t + k^2c\,d_t^2. \tag{IC12}$$

The coefficient of d_t^2 is infinitesimal. For any affine function $f = f_{00} + a_fU + c_fV$ among (IC8), the exact first difference is

$$f - f(k,s) = (u-k)(u+k)(a_f + c_ft^2) + (t-s)k^2(t+s)c_f. \tag{IC13}$$

These are the required bounded u arrays and infinitesimal t arrays; no appeal to unspecified Taylor membership is needed. The two exact first-jet balances are

$$2(\epsilon_{A0} - \epsilon_{C0}) + kh_{c,u0} = 0, \quad 2(\epsilon_{B0} - \epsilon_{C0}) + sh_{c,t0} = 0. \tag{IC14}$$

They follow from the original stationary contact, and are also checked directly using (NC4). The extra square $c_FF_v^2$ has zero value and first jets there. Apply the accompanying three-circuit contact-lift lemma with the exact conductors

$$G(u-k) = k(A_t - G), \qquad G(t-s) = s(B_t - G). \tag{IC15}$$

This gives a strictly positive residue Gram on the finite circuit root list, together with the strict F_v pivot.

Explicit finite allocations of every first normal. Use f_w, f_z, k_w, k_z in (SN2), and define the polynomials

$$\begin{aligned} P_Y(U,V) &= 2(k_w/\eta + k_z), \\ P_E(U,V) &= \frac{2}{\eta^2}(A_w/U - mV/U) - \frac{2}{\eta}(b_0 + b_1U) + bP_Y(U,V). \end{aligned} \tag{IC16}$$

The quotient $A_w/U - mV/U$ is polynomial, as its displayed numerator shows. For either polynomial P use the exact divisions

$$P = P_{00} + UP_U + VP_V, \quad P_U = \frac{P - P(0,V)}{U}, \quad P_V = \frac{P(0,V) - P(0,0)}{V}. \tag{IC17}$$

All arrays are bounded. The source formulas give, by finite numerator division, $P_{Y,00}/x, P_{E,00}/x$ bounded. The complete first normals of the original $F/4$ in the frame (IC4) are exactly

$$\begin{aligned} \mathcal{L}_N &= vP_Y, \\ \mathcal{L}_E &= 2mvt^2/\eta^2 + vP_E, \\ \mathcal{L}_x &= \frac{1}{\sqrt{\beta}}\left[2u\theta(m + \beta + \tfrac{1}{2}) - \frac{2m}{\eta}ut^2(1 + rU) + 2uVZ_X\right], \\ Z_X &= (\beta + \tfrac{1}{2})\theta - (D + \tfrac{1}{2})(1 + hU) + (D + \tfrac{1}{2})(1 - r)/\eta. \end{aligned} \tag{IC18}$$

These identities are independently checked by differentiating the entire original parent after the full state substitution (IC4).

Their actual root allocations are

$$\begin{aligned} vP &= \frac{P_{00}}{k}G + u(P_U + t^2 P_V)A_t, \\ \frac{2m}{\eta^2}vt^2 &= \frac{2mst}{\eta^2 k}B_t, \\ \mathcal{L}_x &= \frac{2u\theta(m+\beta+\frac{1}{2})}{\sqrt{\beta}\,xs}C_t - \frac{2ms(1+rU)}{\eta k\sqrt{\beta}}B_t + \frac{2vZ_X}{\sqrt{\beta}}A_t. \end{aligned} \tag{IC19}$$

Every coefficient in (IC19) is infinitesimal. Its u first-difference array is bounded and its t array is infinitesimal: the coefficient arrays are polynomials in u, t with bounded source arrays, and each term depending on t contains u, k, or $1/\sqrt{\beta}$; evaluation at $u = k$ retains such an infinitesimal factor. There are no negative t powers. The source ratios needed here are precisely

$$m/x, \quad P_{Y,00}/x, \quad P_{E,00}/x, \quad \theta(m+\beta+\tfrac{1}{2})/x, \quad \beta\theta/x, \tag{IC20}$$

all bounded by canceled finite numerator formulas. The exact original contact makes each complete normal zero at $(u,t) = (k,s)$. For a normal $\ell = \sum b_i A_i + b_G G$, put $B = \sum b_i + b_G$, and use the explicit monic divisions

$$B = (u-k)\frac{B(u,t) - B(k,t)}{u-k} + (t-s)\frac{B(k,t) - B(k,s)}{t-s}. \tag{IC21}$$

The first quotient is bounded and the second infinitesimal by the polynomial arrays just given. Replacing A_i by $A_i - G$ and applying (IC15) allocates each complete normal with infinitesimal coefficients on the same strict circuit roots. Joining (IC5), (IC7), and (IC19) therefore gives a bounded positive-residue Gram for the entire original parent in every bounded-v/u testing chart, including infinitesimal v/u and arbitrary state size.

The reciprocal testing-ratio chart. It remains to consider v/u unbounded, so u/v is infinitesimal. Return to homogeneous state coordinates and set

$$p = uW,\ q = vZ,\ E = \eta p - q, \quad Y = rp + bE + N, \quad B = X + uZ = \xi + \theta u Z.$$

Use the complete raw root list

$$E,\ N,\ N_x = \sqrt{\beta}\xi, \quad P = \sqrt{m}\,vW, \quad Q = \sqrt{\alpha}Z, \quad R_t = u^2 W, \quad S_t = uvW. \tag{IC22}$$

No independence of these roots is required. The exact source comparison error in the quadratic base is a quadratic with $O(x)$ source coefficients in (p, Y, q). It has zero residue on (IC22), since

$$\sqrt{x}\,p = \sqrt{x/m}\,(u/v)P, \quad Y = rp + bE + N, \quad q = \eta p - E, \tag{IC23}$$

and x/m is a positive bounded unit. The coefficient u/v is pure testing, not a mixed source/testing comparison. The comparison base itself is exactly $H_0 N^2 + \gamma E^2$, with the positive values in (IC5).

For completeness, with $B = 0$ the exact higher-angle part of the C, A, D packets and half core, after the quadratic base is removed, is

$$\begin{aligned} &U\{AY^2 - 2Dq(p - Y - q) + q^2/2 - pq\} \\ &\quad + V\{C(p-Y)^2 + (A+\tfrac{1}{2})Y^2\} + (A+\tfrac{1}{2})UVY^2 + (D+\tfrac{1}{2})U^2 q^2. \end{aligned} \tag{IC24}$$

Upon inserting $Y = rp + bE + N$, $q = \eta p - E$, its residue on (R_t, S_t) is strictly diagonal:

$$\kappa_R = \frac{\rho\eta}{2}, \qquad \kappa_S = \frac{\rho}{2}. \tag{IC25}$$

All other entries carry u, v, U, or V. For example $UpE = uR_tE$, $VpE = vS_tE$, $U^2p^2 = UR_t^2$, and $UVp^2 = US_t^2$. These identities allocate every term of (IC24) without an inverse or a rate comparison.

The terms involving B are equally explicit. Relative to $B = 0$, the D packet adds $2DuvB(p - Y - D_uq) + DUVB^2$, and the half core adds $uvB(p - Y - D_uq) + (D_v + UV)B^2/2$. Here

$$B = \frac{N_x}{\sqrt{\beta}} + \frac{\theta u}{\sqrt{\alpha}}Q, \quad uvp = uS_t,$$

so their Gram coefficients are infinitesimal. The three remaining original packets keep their entire roots:

$$\begin{aligned} \sqrt{m}\, M &= P + \sqrt{m}\, uvY - \sqrt{m}\, B, \\ \sqrt{\alpha}\sqrt{D_u}(vY - Z) &= \sqrt{D_u}\{\sqrt{\alpha}(rS_t + v(bE + N)) - Q\}, \\ \sqrt{\beta}\sqrt{D_v}B &= \sqrt{D_v}\{N_x + \sqrt{\beta\theta^2/\alpha}\, uQ\}. \end{aligned} \tag{IC26}$$

Each is its leading root $P, -Q, N_x$ plus an infinitesimal-coefficient combination of (IC22). Indeed $\theta/\sqrt{\alpha} = O(\pi)$, $\beta\theta^2/\alpha = O(\pi)$, and m, α are infinitesimal. Thus the complete original parent has a bounded Gram on (IC22) with strictly positive diagonal residue

$$(\gamma, H_0, 1, 1, 1, \kappa_R, \kappa_S). \tag{IC27}$$

This proves the reciprocal-ratio chart. Together with the bounded-ratio chart and the inherited remote theorem it proves the stated all-tests interior source theorem.

Verification and exact scope. The accompanying checker verifies (IC7)–(IC8), every full first normal by original-parent differentiation, the complete hard residues, all canceled source coquotients and contact jets, the finite Taylor identity, and the exact remote-ratio higher-angle identity and both strict pivots. Its denominator report specializes each canceled source denominator at $x = 0$ and records nonzero signed products of positive-unit factors from (IC1); no inverse opening or negative testing-ratio power is hidden in the proof. The ordered-field bounds and the full original remote join are the proof above. This theorem removes exactly the source subdomain (IC1), after the original total-scale intersection; it does not remove infinitesimal η, unbounded ζ, or an infinitesimal r, $1 - r$, or j_0 opening.

Chapter 38

Small L: a complete native normal identity and the exact soft defect

Component `pc15b7caff0a3`
Source: `R13/advances/c1_small_l_native/native_reduction.tex`
SHA256: `c15b7caff0a388deb1a68d7cd796d110161a9afb8573679bddea21cfd5eac5b5`
Inventory use: SN1–SN2 exact full-parent scalar/normal coefficients | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

38.1 Small L: a complete native normal identity and the exact soft defect

This is an exact reduction, not a new admission. Use the unchanged original C1 source, $d = 1$, $x, L = x\zeta \in \mathrm{Inf}_{>0}$, ζ noninfinitesimal, and first suppose η bounded. Retain all source openings, all independent testing rates, and literal $\beta = 0$. The original $D = 0$ face is empty here by (SB2). Only both infinitesimal testing angles remain by the existing remote theorem. No new conclusion on that local chart is asserted below.

Every original state normal, including the beta packet. Put $U = u^2, V = v^2, D_u = 1 + U, D_v = 1 + V$, and use native $r, h = 1 - \theta$:

$$X = -huZ + \xi, \qquad Y = ruW + \nu.$$

Write $F_g/4 = A_w W^2 + A_z Z^2 - 2uv(b_0 + b_1 U)WZ$, with (A_w, A_z, b_0, b_1) from (WS2)–(WS3), setting $n = 0$, replacing the WS letters (h, r) by native (r, h), and adding $\beta\theta^2 U D_v$ to A_z. Then the entire original parent is exactly

$$\begin{aligned} F/4 = F_g/4 &+ 2\xi(vWf_w + uZf_z) + 2\nu(uWk_w + vZk_z) \\ &+ H_{xx}\xi^2 + 2H_{xy}\xi\nu + H_{yy}\nu^2. \end{aligned} \tag{SN1}$$

where

$$\begin{aligned} f_w &= -m(1 + rU) + (D + \tfrac{1}{2})(1 - r)U, \\ f_z &= m\theta - (D + \tfrac{1}{2})V(1 + hU) + (\beta + \tfrac{1}{2})D_v\theta, \\ k_w &= mV(1 + rU) - CD_v(1 - r) + AD_uD_vr - (D + \tfrac{1}{2})(1 - r) + (\alpha + \tfrac{1}{2})D_urV, \\ k_z &= -m\theta U + (D + \tfrac{1}{2})(1 + hU) - (\alpha + \tfrac{1}{2})D_u, \\ H_{xx} &= m + \beta D_v + (D + \tfrac{1}{2})UV + D_v/2, \\ H_{yy} &= mUV + CD_v + AD_uD_v + D + \tfrac{1}{2} + (\alpha + \tfrac{1}{2})D_uV, \\ H_{xy} &= -(m + D + \tfrac{1}{2})uv. \end{aligned} \tag{SN2}$$

There is no discarded normal or excess packet in this identity. Its two exact first-normal contact equations are

$$\zeta f_w(x,y) + f_z(x,y) = 0, \qquad k_w(x,y) + \eta k_z(x,y) = 0, \qquad y = x\eta\zeta. \tag{SN3}$$

They follow by substituting (NC4), not by setting a testing angle equal to a source angle in the desired tensor assertion.

The beta opening creates no unbounded product in its first normal:

$$2\beta\theta P_d = K(r+\eta)(xh+L) - \theta P_d, \qquad \beta\theta = O(L), \quad \beta\theta^2 = O(L^2). \tag{SN4}$$

In particular, after the pure-source normalization $\widehat{\xi} = \sqrt{1+\beta}\,\xi$, its cross coefficient is $\beta\theta/\sqrt{1+\beta} = O(L)$, including every unbounded beta opening. This does not by itself allocate the other normals.

The exact common-radial specialization $U = \lambda x, V = \lambda y, W/Z = \sqrt{\zeta/\eta}$ obeys

$$\frac{F_g}{4Z^2} = (\lambda-1)^2(a_{\mathrm{rad}}\lambda + \alpha), \quad a_{\mathrm{rad}} = xL^2r^2(m+A+\alpha+\tfrac{1}{2}) + x^2yh^2(D+\tfrac{1}{2}) > 0. \tag{SN5}$$

The beta contribution is included; its degree is at most two in λ, so it contributes through the two contact balances. This is only a scalar source slice.

An opening-safe reference binomial. Set $e = \eta T$, $T = 1 + xh$, so that $j = e + r - 1 = \eta J > 0$ exactly. This choice is necessary: $\eta + r - 1$ itself may be negative at an original J opening. Define positive source-only comparison weights

$$C_* = \frac{r+e}{2(1-r)}, \quad A_* = \frac{e}{2r}, \quad D_* = \frac{1}{2j}, \qquad H_* = \frac{e(j+r)}{2r(1-r)j}.$$

For independent indeterminates p, t, Y there is the exact identity

$$\begin{aligned} &C_*(p-Y)^2 + A_*Y^2 + D_*(p-Y-t)^2 + \tfrac{1}{2}(p-Y)^2 - pt \\ &= H_*\left(Y - \frac{r(e+r)}{j+r}p + \frac{r(1-r)}{e(j+r)}t\right)^2 + \frac{e+r}{2e(j+r)}(ep-t)^2. \end{aligned} \tag{SN6}$$

The relevant separated roots are $p = uW$, $t = vZ$, and $\sqrt{e}\,uW - vZ/\sqrt{e}$. The latter also remains legitimate when η is infinitesimal; no mixed inverse is introduced. All original openings remain positive factors. Directly from (NC4),

$$\frac{C}{C_*} = \frac{K(r+\eta)E}{(r+\eta T)P_d} \sim 1, \quad \frac{A}{A_*} = \frac{P_4}{TP_d} \sim 1, \quad \frac{D}{D_*} = \frac{B_5}{P_d} \sim 1. \tag{SN7}$$

These are uniform relative comparisons, not a license to replace the actual packets in a semidefinite residue argument.

The actual quadratic base has a strictly negative soft determinant. For the actual coefficients set

$$H = C + A + D + \tfrac{1}{2}, \quad a = \frac{A(C+D+\frac{1}{2})}{H}, \quad c = \frac{D(C+A+\frac{1}{2})}{H}, \quad b = \tfrac{1}{2} + \frac{AD}{H}.$$

Exact completion of the same quadratic base gives its full hard normal and soft form

$$H\left(Y - \frac{C+D+\frac{1}{2}}{H}p + \frac{D}{H}t\right)^2 + ap^2 + ct^2 - 2bpt. \tag{SN8}$$

Its determinant is

$$ac - b^2 = \frac{8ACD - 4AD - 2A - 2C - 2D - 1}{8H}. \tag{SN9}$$

The unchanged native source has the following exact factorization:

$$8ACD - 4AD - 2A - 2C - 2D - 1 = -\frac{\eta(r+\eta)^2 K P_4\,\Delta}{r(1-r)jP_d^3}, \tag{SN10}$$

where

$$\Delta = R(\eta K + R)L^2 + (2 + xA_1)L + x(4h - 1 + xA_0), \qquad R = 1 + xr, \tag{SN11}$$

and the bounded polynomial arrays are

$$\begin{aligned} A_1 = {} & -\eta^2h^2x^2 - \eta^2h^2x - 2\eta h^2rx^2 - 2\eta h^2x + 2\eta hrx^2 + 3\eta hx + \eta h \\ & + 2hr^2x^2 + 6hrx + 4h + 2r, \\ A_0 = {} & \eta^2h^4x^2 - \eta^2h^3x^2 + \eta^2h^3x - \eta^2h^2x - 2\eta h^3rx^2 - 4\eta h^3x \\ & + \eta h^2rx^2 - \eta h^2rx + 2\eta h^2x - 4\eta h^2 + 2\eta h + h^2r^2x^2 + 4h^2rx + 4h^2 + 2hr. \end{aligned}$$

Since (SB1) gives $h \sim 1$, all arrays are bounded and

$$\Delta/(2L + 3x) \sim 1. \tag{SN12}$$

Indeed the L^2, xL, and x^2 errors divided by $L + x$ are infinitesimal, independently of the relative x, L rate. Every factor outside the minus sign in (SN10) is positive. Thus the actual base in (SN8) has a strictly indefinite soft block through every opening. The rank-one reference (SN6) is therefore insufficient even at the exact quadratic level. A weighted circuit must absorb the precise defect (SN10), using the original m, α, higher-angle terms, and the complete normals (SN1)–(SN2). Neither scalar leading PSD nor the radial source slice proves that allocation.

Unbounded eta and exact scope. The full identities (SN1)–(SN5), (SN8)–(SN11) are algebraic and also hold for unbounded η. The bounded-array estimate (SN12) and comparisons (SN7) above are only asserted for bounded η. The available exact canonical C1 stabilizer (AS5) sends k to $1/k$, and hence does not send this small-k face into the stated C1.1 theorem, which also assumes k infinitesimal. No other exact source-domain mapping has been established here. The original C1.1 proof cannot be imported merely because both faces have large C, A and weak D, α.

The checker verifies the entire parent, both complete first normals, the radial identity, all comparisons and the exact source defect. It records no new local or all-tests admission. C14.3 retains the both-infinitesimal testing chart, and C14.4 retains its previously specified infinitesimal-u charts, including all independent rates, openings and allowed boundaries. All earlier remote and homogeneous endpoint admissions remain intact.

Chapter 39

C1 UH: a homogeneous biquadratic conductor

Component `pf7ffa921793c`
Source: `R13/advances/c1_uh/uh_all_tests.tex`
SHA256: `f7ffa921793c0b1b432463bba3a4de95659a1186ce3299ca64731620b8b2ed33`
Inventory use: UH1–UH20 and complete source-boundary finite-cap proof | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

39.1 C1 UH: a homogeneous biquadratic conductor

All coefficients below are divided by the original positive pure-source d. We treat precisely (NI9) and (NI10), with no alteration of the native source (NC1)–(NC7). The revision-6 common-event theorem already covers all tests except positive unit u and $w = 1/v$ infinitesimal. Put $x = k^2$, $K = 1 + x$, $\sigma = 1/\eta$, $z = 1/\zeta = q^2$, $b = r/z$, $H = h/\sigma$, and retain $\sigma > 0$ (it need not be a unit). On (NI9), σ is infinitesimal and z is a unit; on (NI10), z is infinitesimal and σ is bounded. Throughout, k, H are positive units, b, z, σ are bounded, $Q = 1 - xb > 0$, and

$$r = zb, \quad R = 1+xzb, \quad M = R-z+\sigma zH, \quad J = 1+\sigma(xH-1+zb), \quad \theta = 1-\sigma H, \quad E_s = 1-xH.$$

The exact source polynomials, all with bounded arguments, are

$$\begin{aligned} P &= xK + \sigma x + \sigma z + \sigma z b x^2 + x^2\sigma z H(1-\sigma H) + 2x\sigma^2 z H + \sigma^2 z^2 b x^2 H, \\ V &= z(1+x\sigma H)E_s + x(1-zb)(1+\sigma z H), \\ B_\alpha &= R\sigma z H + RxH + ME_s. \end{aligned}$$

Writing $m = m_{\mathrm{old}}/z$, $c = C/z$, $a = Az$, $\ell = \alpha z$,

$$c = \frac{K(1+\sigma zb)E_s}{2(1-zb)P}, \quad m = \frac{Kx(1+\sigma zb)H(\sigma z + x)}{2MP}, \quad a = \frac{V}{2bP}, \quad \ell = \frac{x^2(1+\sigma zb)B_\alpha}{2QP}. \qquad \text{(UH1)}$$

Define the polynomial quotients

$$N_D = \frac{K(\sigma z + x)(1+\sigma zb) - JP}{\sigma}, \qquad N_\beta = \frac{Kx(1+\sigma zb)(1+\sigma zH) - \theta P}{\sigma}.$$

Then $D = \sigma N_D/(2JP)$ and $\beta = \sigma N_\beta/(2\theta P)$, exactly.

The bounded full parent. Let $D_u = 1 + u^2$, $D_w = 1 + w^2$. Its six event amplitudes are

$$M_w = W - wX + uY - uwZ,\ R_w = uW - Y,\ Y,\ J_w = uwW + uX - wY - Z,\ Y - wZ,\ B_w = X + uZ.$$

Thus the exact parent is

$$F = 4mzM_w^2 + 4czD_wR_w^2 + 4(a/z)D_uD_wY^2 + 4DJ_w^2 + 4(\ell/z)D_u(Y - wZ)^2 + 4\beta D_wB_w^2 + 2K_w, \quad \text{(UH2)}$$

where, without any restriction to a state ray,

$$\begin{aligned} K_w = {} & u^2w^2Z^2 - 2u^2wYZ - 2uwWZ + 2uw^2XZ + u^2w^2W^2 + 2u^2wWX - 2uw^2WY \\ & + (D_u + w^2)(X^2 + Y^2) - 2uwXY. \end{aligned}$$

For a potentially unbounded weight choose a finite cap, keeping bounded weights unchanged. Subtract and retain the following nonnegative centered excesses:

$$\begin{aligned} & 4(m - m_*)[qM_w - kM(wZ/q)]^2, \\ & 4(c - c_*)D_w[qR_w - x(1 - r)(wZ/q)]^2, \\ & 4(a - a_*)D_uD_w[(Y - xbwZ)/q]^2, \quad 4(\ell - \ell_*)D_u[(Y - xbwZ)/q]^2. \end{aligned} \quad \text{(UH3)}$$

The first line uses qM_w and the second qR_w; thus these are exactly excesses of the scaled weights in (UH1). Their remainders are bounded because $mM, c(1 - r), ab, \ell Q$ are bounded. When σ is a unit, additionally retain

$$4(D - D_*)(J_w + JZ)^2, \qquad 4(\beta - \beta_*)D_w(X + uhZ)^2. \quad \text{(UH4)}$$

Here DJ and $\beta\theta$ are bounded. When σ is infinitesimal, no (UH4) cap is needed: $D, \beta = O(\sigma^2)$, as proved below. Denote the complete remainder by G. Every retained row vanishes at the actual contact

$$u = k, \quad w = \frac{q\sqrt{\sigma}}{k}, \qquad (W, X, Y, Z) = (\frac{\sqrt{\sigma}}{q}, -kh, kbq\sqrt{\sigma}, 1). \quad \text{(UH5)}$$

No comparison between independent source and test rates occurs.

Exact homogeneous conductor, including every first normal. Put $h = \sigma H$, $\delta = u - k$, and use the six roots

$$\begin{aligned} N_x = X + khZ, \quad N_y = (Y - kbq^2W)/q, \\ U = wW - \sigma Z/k, \quad N = (wZ - q^2W/k)/q, \\ S = q\delta W, \quad T = \sigma\delta Z. \end{aligned} \quad \text{(UH6)}$$

Substitute $X = N_x - khZ$, $Y = qN_y + kbq^2W$ in the entire bounded remainder G. Its complete quadratic normal block is the coefficient collection in (N_x, N_y). Its scalar and first-normal coefficients have exactly the following supports:

$$\begin{aligned} G_{00} &= \frac{A}{q^2}w^2Z^2 + Bw^2W^2 + C\sigma^2Z^2 + D_1wWZ + Eq^2W^2, \\ G_{10} &= A_xwW + B_xZ + C_xw^2Z, \\ G_{01} &= q(A_y + B_yw^2)W + C_ywZ/q. \end{aligned} \quad \text{(UH7)}$$

All eleven displayed coefficient families and the quadratic normal coefficients are bounded polynomials in u, with bounded source coefficients; B_x is divisible by σ. In the unit-σ case divisibility means a bounded quotient, not a new polynomial identity. Formulas below specify every coefficient before caps, and (UH3)–(UH4) specify their exact changes. No state normal is omitted.

Set $s = \sigma/k$, $a_0 = q^2/k$ and

$$L = 2A/k + 2B\sigma/k + D_1, \quad J_0 = C - B/k^2, \quad K_0 = A/k^2 + 2B\sigma/k^2 + D_1/k + E.$$

The scalar is identically

$$AN^2 + BU^2 + qLWN + \sigma^2J_0Z^2 + q^2K_0W^2. \quad \text{(UH8)}$$

Exact source contact and all stationary jets give

$$L(k) = J_0(k) = K_0(k) = 0, \qquad K_0'(k) = -\sigma J_0'(k),$$
$$B_x(k) + s[A_x(k) + a_0 C_x(k)] = 0, \qquad A_y(k) + C_y(k)/k + s a_0 B_y(k) = 0. \tag{UH9}$$

These hold separately for every centered excess, hence for every cap. Write the monic divisions

$$L = \delta L_1, \quad J_0 = \delta j + \delta^2 J_2, \quad K_0 = -\sigma\delta j + \delta^2 K_2,$$
$$B_x + s(A_x + a_0 C_x) = \sigma\delta H_x, \qquad A_y + C_y/k + s a_0 B_y = \delta H_y.$$

All quotients are bounded. The scalar and full first normals become

$$\begin{aligned} G_{00} &= AN^2 + BU^2 + (L_1 + k\sigma j)SN - kjTU + J_2 T^2 + K_2 S^2, \\ G_{10} &= (A_x + a_0 C_x)U + C_x q w N + H_x T, \\ G_{01} &= (C_y + s q^2 B_y)N + q B_y w U + H_y S. \end{aligned} \tag{UH10}$$

Together with the unchanged full quadratic normal block, this is an exact bounded six-root Gram for G, not a scalar restriction. The only root relation needed for strict completion is

$$T^2 = \sigma S^2 + k\sigma\delta SN - k\delta TU. \tag{UH11}$$

Its verification is one multiplication using (UH6). Adding any bounded multiple of its zero changes no parent. No mixed inverse, root, order, or source/test identification occurs in (UH6)–(UH11).

For the cap argument it is useful to print every centered row directly in this same frame. In the order $m, c, (a, \ell), D, \beta$, they are

$$\begin{aligned} & uzN_y - (kR + z\delta)N - qwN_x - (zQ/k)S, \\ & S - zN_y - x(1-r)N, \qquad N_y - xbN, \\ & (u - kbz)U + uN_x - qwN_y + (E_s/k)T, \qquad N_x + HT. \end{aligned} \tag{UH20}$$

These are exact identities before taking any residue.

Explicit coefficient input. For (UH2) before caps put $j_u = u - kbz$, $p_u = 1 + ukbz$, $t_u = kh - u$. The full scalar coefficients in (UH7) are

$$\begin{aligned} A &= 4mz^2 t_u^2 + 4\ell D_u + (4\beta + 2)z t_u^2, \\ B &= (4cz + 4D + 2)j_u^2 + 4aD_u x b^2 z, \\ C &= [4D(1 + ukh)^2 + 4\beta(u - kh)^2 + 2D_u x h^2]/\sigma^2, \\ D_1 &= 8mzp_u t_u - 8Dj_u(1 + ukh) - 8\ell D_u kb - 4u^2 k(h + bz) - 4u + 4uxhbz, \\ E &= 4mp_u^2 + 4cj_u^2 + 4(a + \ell)D_u x b^2 + 2D_u x b^2 z. \end{aligned}$$

The first normals are

$$\begin{aligned} A_x &= -8mzp_u + (8D + 4)uj_u, \\ B_x &= -8Du(1 + ukh) + 8\beta(u - kh) - 4D_u kh, \\ C_x &= -8mzt_u + (8\beta + 4)(u - kh), \\ A_y &= 8mzup_u - 8czj_u + 8(a + \ell)D_u kb + 4D_u kbz, \\ B_y &= -8czj_u + 8aD_u kb - (8D + 4)j_u, \\ C_y &= 8mz^2 ut_u - 8\ell D_u + 8Dz(1 + ukh) + 4uzt_u. \end{aligned}$$

The coefficients of $N_x^2, N_y^2, N_x N_y$ are respectively

$$4mzw^2 + 4Du^2 + 4\beta D_w + 2(D_u + w^2),$$
$$4mz^2u^2 + 4cz^2 D_w + 4aD_u D_w + 4\ell D_u + 4Dzw^2 + 2z(D_u + w^2), \qquad -8mzuqw - 8Duqw - 4uqw. \tag{UH12}$$

Here products such as Du mean $D\,u$, whereas $D_u = 1 + u^2$.

Infinitesimal σ: source collapse and strict residue. The exact polynomial $P_m = (N_D + xN_\beta)/\sigma$ is bounded. At $\sigma = 0$, $N_D = -xN_\beta$ and

$$N_\beta = x(K+z)H + bxz - x - z.$$

Nonnegativity of D, β forces

$$H = H_0 + O(\sigma), \quad H_0 = \frac{x + zQ}{x(K+z)}, \qquad E_0 = 1 - xH_0 = \frac{R}{K+z}. \tag{UH13}$$

Indeed $JD + x\theta\beta = \sigma^2 P_m/(2P)$ and J, θ are units. Thus $D, \beta = O(\sigma^2)$, including either literal zero. Moreover

$$(D + x\beta)/\sigma^2 \longrightarrow KH_0E_0/2 > 0.$$

The other limiting coefficients are

$$c_0 = \frac{R}{2x(1-zb)(K+z)}, \quad m_0 = \frac{xH_0}{2(1-zQ)}, \quad a_0^{\mathrm{wt}} = \frac{H_0}{2b}, \quad \ell_0 = \frac{xR}{2Q(K+z)}. \tag{UH14}$$

At a matched angle $u = k$, the residue Gram (UH10) splits into (N_y, N, S) and (N_x, U, T), because their only off-block coefficients contain w. Before caps, the (N_y, N) block is positive definite; its angular Schur complement is

$$\Gamma = 8RH_0/K > 0. \tag{UH15}$$

The (N_x, U) block is precisely

$$2K(N_x + xH_0U)^2 + 2KxH_0E_0U^2. \tag{UH16}$$

The remaining T couplings are bounded. As σ, δ have zero residue, (UH11) permits an arbitrarily large positive T^2 pivot without changing any other residue coefficient. This proves a strict full six-root residue Gram. For $u^2 \neq x$, use the full raw frame $(E_t, F_t, G_t, R_t, X, T_0) = (qW, Y/q, wZ/q, wW, X, \sigma Z)$. Its leading first block is

$$\begin{aligned} &4m_0[E_t + uz(F_t - G_t)]^2 + 4c_0(uE_t - zF_t)^2 + 4a_0^{\mathrm{wt}} D_u F_t^2 + 4\ell_0 D_u (F_t - G_t)^2 \\ &\quad + 2z\{u^2(G_t - F_t)^2 + F_t^2\} - 4uE_tG_t. \end{aligned}$$

Put $v_0 = u^2$ and $M_0 = 1 - zQ$. In $(F_t, F_t - G_t)$ the hard block is diagonal, with weights

$$B_h = \frac{2(x+zQ)[v_0(1-r)+R]}{bx(1-r)(K+z)}, \qquad L_h = \frac{2R[v_0(x+zQ)+xM_0]}{Q(K+z)M_0}.$$

The exact E_t Schur complement is

$$\frac{2(v_0 - x)^2(1+v_0)RH_0}{[v_0(1-r)+R][v_0(x+zQ)+xM_0]}. \tag{UH17}$$

It has a positive-unit limit through every opening when $v_0 \neq x$. The hard minimizers are

$$\frac{F_t}{E_t} = \frac{ubK}{v_0(1-r)+R}, \qquad \frac{F_t - G_t}{E_t} = \frac{-uKQ}{v_0(x+zQ)+xM_0},$$

also bounded through every opening. The other raw blocks are $2\{u^2R_t^2 + 2u^2R_tX + D_uX^2\}$ and $4(D + u^2\beta)T_0^2/\sigma^2$, both strictly positive. This gives the full bounded away-angle lift, without a determinant-only claim or any omitted first state normal.

Unit σ: angular Schur sign. Here (NI10) gives $z = 0$ at residue. Put $h = \sigma H$, $J = 1+\sigma(xH-1)$, $\theta = 1 - \sigma H$. The source signs imply

$$0 < \sigma < K, \qquad \frac{1}{K+\sigma} \le H \le \frac{x+\sigma}{x(K+\sigma)}. \tag{UH18}$$

These inequalities are stated on the interior; its closure includes $\sigma = K$ when $J = \theta = 0$. The limits are

$$m = \frac{KxH}{2(K+\sigma)}, \quad c = \frac{K(1-xH)}{2x(K+\sigma)}, \quad a = \frac{1}{2b(K+\sigma)}, \quad \ell = \frac{x}{2Q(K+\sigma)},$$
$$D = \frac{K}{2J(K+\sigma)} - \tfrac{1}{2}, \quad \beta = \frac{K}{2\theta(K+\sigma)} - \tfrac{1}{2}.$$

The two hard blocks in (UH10) are positive definite. Their angular Schur coefficients γ_S, γ_T have no ST term and satisfy

$$\gamma_S + \sigma\gamma_T = \frac{-2p(H)}{\sigma(K+\sigma)} > 0, \qquad p(H) = Kx(K+4\sigma)H^2 - 2K(x+2\sigma)H + x. \tag{UH19}$$

Convexity of p and (UH18) prove strict negativity: its endpoint values are respectively

$$\frac{-\sigma(3\sigma x + 4\sigma + 4x + 4)}{(K+\sigma)^2}, \quad \frac{-\sigma(4\sigma x + 3\sigma + 4x^2 + 4x)}{x(K+\sigma)^2}.$$

Use (UH11) to distribute this positive total between S^2, T^2; for example choose its multiple so both resulting Schur coefficients are positive. This yields the required strict six-root Gram.

For completeness, angles $u^2 \ne x$ are strictly positive as well. Eliminating the two hard state variables from the leading form, multiplying by the positive source factor $(K+\sigma)/2$, and writing $U_0 = u^2$, $P_0 = (w/q)(Z/q)$, $T_0 = (Z/q)^2$ gives a positive Laurent function

$$A + BP_0^2 + CP_0^2/T_0 + D_0T_0 - 2EP_0,$$

where

$$A = KxH + K(1-xH)U_0/x, \; B = x(1+U_0), \; C = \frac{KU_0}{J+\theta U_0},$$
$$D_0 = (1+U_0)\left[\frac{K(1+U_0)}{J+\theta U_0} - (K+\sigma)\right], \quad E = \frac{K\sqrt{U_0}(1+U_0)}{J+\theta U_0}.$$

The strict criterion is $E < \sqrt{AB} + \sqrt{CD_0}$. Squaring only to analyze these positive real residue quantities gives

$$(E^2 - AB - CD_0)^2 - 4ABCD_0 = \frac{K^2(1+U_0)^2}{(J+\theta U_0)^2}(U_0 - x)^2 Q_2(U_0).$$

The quadratic Q_2 has positive constant $H^2x^2J^2$, positive leading coefficient $\theta^2(1-xH)^2$, and $Q_2(x) = xp(H) < 0$. Explicitly, with $E_s = 1 - xH$,

$$Q_2(U_0) = E_s^2\theta^2U_0^2 + 2E_s\{x\theta^2 + E_s\theta J + (K+\sigma)\theta - 2K\}U_0 + x^2H^2J^2.$$

The concave quadratic $U_0(K+\sigma) - [x^2H + (1-xH)U_0](J+\theta U_0)$ is negative at zero and infinity and positive at x; at each of its two zeros the preceding discriminant is negative. Consequently its positive interval lies inside $Q_2 < 0$. This proves the criterion for all $U_0 \ne x$. Sign charts with $P_0 \le 0$ are immediate. These square roots are an analysis of real residues only and are not tensor roots or mixed localizations.

This inequality is directly a bounded Gram allocation, not a comparison of mixed ratios. Before choosing projective coordinates, put $E_t = qW$, $G_t = wZ/q$, $R_t = wW$. The exact relation $E_tG_t = R_tZ$ permits distributing the cross term between the two quadratic blocks (E_t, G_t) and

(R_t, Z). The strict inequality above is exactly the condition that both 2×2 blocks can be made positive definite by a finite real allocation. It therefore handles every independent rate of q, w. At $J = 0$ or $\theta = 0$, the root-interval argument permits one endpoint at zero or infinity. At the double face $J = \theta = 0$, necessarily $\sigma = K$, $H = 1/K$. As $J + \theta U_0 \to 0$,

$$E - \sqrt{CD_0} \longrightarrow K\sqrt{U_0}, \qquad AB - K^2 U_0 = (U_0 - x)^2.$$

Thus the constrained allocation remains strictly positive away from $U_0 = x$, including this double face without dividing by a vanishing residue denominator.

Opening faces, tensor conclusion, and exact scope. The finite-cap argument is applied to the full Gram above. For σ infinitesimal the possible diverging rows have openings $1 - zb$, $1 - zQ$, b, Q; at most two can diverge together. The pairs (c, m), (m, a), (c, ℓ) have independent hard rows; other pairs are excluded by $0 < zb, xb < 1$ and $1 - zQ > 0$. In the (N_y, N) columns of (UH20), their determinants have absolute values kzK, k, x, respectively; each is a unit on its compatible face. The hard block is uniformly coercive, since it contains positive-unit multiples of N_y^2 and $(N_y - N)^2$. Its minimizing hard coordinates for fixed angular S are explicitly

$$\frac{N_y}{S} = \frac{b(2bxz - x + 1)}{K}, \qquad \frac{N}{S} = \frac{4bxz - x - 2z + 1}{xK}.$$

They are bounded through every opening. Thus the limit of the uncapped Schur complement is the Schur complement restricted to the diverging centered rows' kernel. The quotients (UH15)–(UH16) stay positive units there. The elementary finite-dimensional penalty lemma now applies: a bounded symmetric form strictly positive on the common kernel becomes strictly positive after sufficiently large finite positive multiples of these rows' squares are added. This proves the finite-cap claim, with the entire angular right-hand sides in (UH20) retained. Away from $u^2 = x$, (UH17) and its printed bounded hard minimizers give the same constrained strictness directly.

For unit σ, only b, Q, J, θ can be openings. The b, Q rows coincide in the hard direction $N_y - xbN$ and cannot both diverge. The J, θ rows on (N_x, U) are independent even if both vanish; their determinant is the unit u. Formula (UH19) remains a positive unit through these faces. More explicitly, the two hard blocks dominate positive-unit multiples of $N_y^2 + (N_y - N)^2$ and $N_x^2 + (N_x + U)^2$. Their angular minimizers are

$$\frac{N_y}{S} = \frac{b(1 - HxK)}{K}, \quad \frac{N}{S} = \frac{1 - HxK}{xK},$$

$$\frac{N_x}{T} = \frac{\sigma H^2 xK - HxK - \sigma H(2x + 1) + x}{\sigma K}, \quad \frac{U}{T} = \frac{HxK(K + \sigma) - xK - \sigma}{x\sigma K}.$$

Their D and β row defects divided by T are respectively $k(HK - 1)J/(\sigma K)$ and $-x\theta(HK - 1)/(\sigma K)$. Hence every relevant constrained limit exists with bounded minimizers; the same penalty argument applies. On the angular relation $T^2 = \sigma S^2$, its strictly positive Schur value is (UH19), and after taking finite caps the bounded zero (UH11) distributes that value into two strict angular pivots. The away-angle allocation and its single/double-face limits above give the same conclusion there. All removed excesses remain exact positive source/test squares. No positive lower bound on an original opening has been imposed.

A strict bounded residue Gram lifts by the inherited tensor Gram lemma. Every root in (UH6) is a finite sum of separated source and testing factors. After restoring the retained squares this certifies the complete original parent, including $D = 0$ or $\beta = 0$. Together with revision-6 remote tests, this closes both (NI9) and (NI10). Those are the entire actual UH portion of C1.5. The already proved UU and UI portions and exact source stabilizers therefore close the entire C1.5 row, subject to the exact checks and mathematical review recorded alongside this section.

Chapter 40

A complete local strong-II Gram with independent testing rates

Component `pe16499ccc4c6`
Source: `R13/advances/c2_strong_local/local_strong_ii.tex`
SHA256: `e16499ccc4c6e225c37f6738a69f33eee35bd3011807dfbaf3b930149d79d28a`
Inventory use: LII2–LII20 exact frame, contact divisions and allocation | Complete strong-II local current, exact scalar and every state normal, strict Gram | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

40.1 A complete local strong-II Gram with independent testing rates

Theorem 40.1 (Strong-II local allocation)**.** *Let the positive C2 source satisfy $z, y \in \mathrm{Inf}_{>0}$ and let $B = H + L$ be a positive unit. Keep the exact source coefficients of (GE1), with $n \geq 0$ and the remaining displayed weights positive. Then the original C2 parent belongs to $\sum (K_c \otimes_{\mathbb{R}} K_t)^2$ whenever both independent affine testing coordinates u, v are infinitesimal, including either literal testing axis. There is no comparison hypothesis between u^2, z, between v^2, y, or between the two source rates. All openings of r, L/y, $1 + zr - L/y$, and $1 - H$ are retained, as is literal $n = 0$.*

This theorem is a full-parent local result. The remote testing charts and the original projective endpoints are separate obligations; they must be joined before the strong II/HH checklist row is admitted. All identities below are polynomial in testing coordinates. Inverses are taken only in the pure source field, and the final step uses the bounded Gram strict-real-residue lemma already used in the completed local UI certificate. In particular no ordering or inverse of the mixed tensor is used.

Source units and notation. Write $U = u^2, V = v^2, k = \sqrt{y}$ and

$$a_c = 1 + zr, \quad t = L/y, \quad j = a_c - t, \quad s = a_c/B, \quad c = CL, \quad d = Dyj.$$

The source entry gives bounded t, j, r, positive units $B, c, d, m/z^2, A/y$, bounded nr and $\alpha(1 - H)$, and strong α. These facts include the arbitrary openings $t \to 0$, $j \to 0$, $r \to 0$, and $1 - H \to 0$. They follow directly from the positive source formulas and were proved in (DI7). Let $h = \mathrm{st}\, B$; then $0 < h \leq 1$ and, with $r_0 = \mathrm{st}\, r$,

$$\begin{gathered} c_0 = d_0^{\mathrm{pair}} = \frac{h}{2(1+h)}, \quad \mathrm{st}(m/z^2) = \frac{1 - r_0}{2h^2(1+h)}, \quad \mathrm{st}(A/y) = \frac{1}{2h(1+h)}, \\ \mathrm{st}(nr) = \frac{1}{2(1+h)} - \frac{r_0}{2}, \qquad \mathrm{st}\{\alpha(1-H)\} = \frac{h^2}{2(1+h)}. \end{gathered} \tag{LII1}$$

These residues follow directly from (GE1): $L \to 0$, $H \to h$, $a_c \to 1$, $G \to h$, $\delta \to h(1+h)$, and the source numerator $M \to h^2$. Substitution in the six rational coefficient formulas gives (LII1),

including the identity $\mathrm{st}(nr) = 1/[2(1+h)] - r_0/2$ without division by n. Define the source quantities (with q, g bounded)

$$\ell = y(C+D) = c/t + d/j, \qquad q = \frac{D}{C+D} = \frac{dt}{cj+dt}, \qquad g = \frac{yCD}{C+D} = \frac{cd}{cj+dt}.$$

Here $0 < q < 1$, g is a positive unit with residue $h/[2(1+h)]$, and ℓ has a positive real lower bound, although it may be unbounded. Indeed $cj + dt$ is a unit because $t + j = a_c$ is a unit. Choose positive real $\ell_* \le \ell$ and $\alpha_* \le \alpha$.

The state roots and the exact C,D completion. Use the source-fixed parity graph

$$X = -\frac{zrv}{B}W + \xi, \qquad Y = \frac{Hu}{B}W + \eta, \qquad Z = \frac{uv}{B}W + \zeta. \tag{LII2}$$

It agrees with the actual kernel at $U = z, V = y$ and is a change of all four homogeneous state coordinates with determinant one. Set

$$Q_n = u\xi - \zeta, \quad K = ukW, \quad R_h = \frac{vQ_n - su(V-y)W}{k}, \qquad N = \frac{\eta}{k} - qR_h. \tag{LII3}$$

Thus $\eta = k(N + qR_h)$. Every root in (LII3) vanishes at the actual contact. If $Q = uW - Y$ and $J = Q + v(uX - Z)$, then

$$Q/k = (t/B)K - N - qR_h, \qquad J/k = (1-q)R_h - N - (j/B)K.$$

Consequently the exact weighted completion is

$$\begin{aligned} 4CQ^2 + 4DJ^2 &= 4\ell N^2 + 4gR_h^2 + \frac{8(d-c)}{B}KN - 8gsKR_h + \frac{4(ct+dj)}{B^2}K^2, \\ 4CVQ^2 &= 4CV\eta^2 + \frac{4ct}{B^2}VK^2 - \frac{8c}{B}VK(N + qR_h). \end{aligned} \tag{LII4}$$

There is no estimate in these identities. Retain the following four nonnegative, contact-centered packets outside the remainder:

$$4(\ell - \ell_*)N^2, \qquad 4CV\eta^2, \qquad 4n(1+U)(1+V)\xi^2, \qquad 4(\alpha - \alpha_*)(1+U)(v\eta - \zeta)^2. \tag{LII5}$$

Call the exact remainder G. Its coefficients are bounded after writing $\eta = k(N + qR_h)$ and using (LII4). For completeness, its entire formula is

$$\begin{aligned} G = {} & 4\ell_* N^2 + 4gR_h^2 + \frac{8(d-c)}{B}KN - 8gsKR_h \\ & + \frac{4(ct+dj)}{B^2}K^2 + \frac{4ct}{B^2}VK^2 - \frac{8c}{B}VK(N + qR_h) \\ & + 4n(1+U)(1+V)\left\{\frac{z^2r^2V}{B^2}W^2 - \frac{2zrv}{B}W\xi\right\} \\ & + 4\alpha_*(1+U)(v\eta - \zeta)^2 + 4\alpha(1+U)\left\{\frac{(1-H)^2UV}{B^2}W^2 - \frac{2(1-H)uv}{B}W(v\eta - \zeta)\right\} \\ & + 4mM^2 + 4A(1+U)(1+V)Y^2 \\ & + 2\{J^2 + (1+U)(vY - Z)^2 + (1+V)(X + uZ)^2 - (1+U)(1+V)Z^2\}, \end{aligned} \tag{LII6}$$

where $M = vW - X + uvY - uZ$ and (LII2) is used in the last line. Every potentially unbounded original coefficient occurs here only through nr, $\alpha(1-H)$, or the bounded quantities in (LII1). No division by a testing coordinate has been introduced.

One exact gauge removes the dependent-root first normal. In the formal bounded polynomial (LII6), the coefficient of R_hW is $uk\rho(U,V)$ for a bounded source polynomial ρ. Replace this entire term using the exact identity

$$KR_h = uvWQ_n - sU(V-y)W^2. \tag{LII7}$$

Call the resulting polynomial $\widetilde{G}$. It is equal to G on the roots (LII3), and has no R_hW term. This gauge is useful because R_h is a dependent root: it permits the state and angular contact equations to be read as formal coefficient vanishings.

More precisely, by parity write uniquely

$$\widetilde{G} = Q_0(N,\xi,\zeta,R_h) + W\{vP_\xi(U,V)\xi + uvP_\zeta(U,V)\zeta + ukP_N(U,V)N\} + \phi(U,V)W^2, \tag{LII8}$$

where all coefficients are bounded source polynomials and Q_0 is quadratic in the four displayed roots. Formula (LII6), followed by the one replacement (LII7), is an exact coefficient-extraction specification for every entry in (LII8); the accompanying checker also prints them without suppression.

The retained packets (LII5) vanish together with their first jets at the actual contact. The exact original source state equations thus give

$$P_\xi(z,y) = P_\zeta(z,y) = P_N(z,y) = 0. \tag{LII9}$$

Indeed the physical η derivative first forces the N coefficient to vanish; the physical ξ,ζ derivatives then force their formal coefficients to vanish because the R_hW coefficient was removed. The exact original contact and the two angular derivatives now give

$$\phi(z,y) = \phi_U(z,y) = \phi_V(z,y) = 0. \tag{LII10}$$

This uses the six exact original source substitutions already checked in R6. It imposes no new relation between source and testing fields.

Bounded division and the complete angular allocation. There are two additional elementary support facts in (LII8):

$$P_\xi \in (U,z), \qquad \deg_U \phi \le 2, \qquad \phi(0,V) = V(d_s + e_sV), \tag{LII11}$$

where

$$d_s = \frac{4}{B^2}\{m(B+zr)^2 + (n+\tfrac12)z^2r^2\}, \qquad e_s = \frac{4}{B^2}(n+\tfrac12)z^2r^2.$$

Both d_s/z^2 and e_s/z^2 are bounded. The first support assertion is literal bounded-coefficient membership: setting $U = z = 0$ in the extracted polynomial gives zero. It is also clear before expansion: all ξW terms not supplied by the gauge contain U, m, nzr, or zr; those supplied by the gauge contain U.

Monic division using (LII9) therefore gives bounded polynomials

$$\begin{aligned} P_\xi &= (U-z)A_\xi + z(V-y)B_\xi, \\ P_\zeta &= (U-z)A_\zeta + (V-y)B_\zeta, \\ P_N &= (U-z)A_N + (V-y)B_N. \end{aligned} \tag{LII12}$$

For the first line, evaluate P_ξ at $U = z$: the support assertion supplies the factor z with bounded quotient, and its value at $V = y$ is zero. All other divisions are by the monic polynomials $U - z$ or $V - y$. In particular none creates an inverse of a small source or a testing difference.

The scalar has the exact bounded decomposition

$$\phi = \frac{d_s + e_sV}{z^2}V(U-z)^2 + U(U-z)(V-y)A_s(V) + U(V-y)^2C_s(V). \tag{LII13}$$

Here is a constructive proof, including boundedness. Write $\phi = a(V)U^2 + b(V)U + V(d_s + e_sV)$ and subtract the first term of (LII13). The remaining U^2, U coefficients a_1, b_1 vanish at $V = y$ by the value and U derivative in (LII10). Divide them by $V - y$ to obtain A_s, B_s. The V derivative gives $zA_s(y) + B_s(y) = 0$, so set $C_s = (B_s + zA_s)/(V - y)$. All three divisions are monic divisions of bounded polynomials. This proves (LII13) exactly, not asymptotically.

Introduce the two legal angular roots

$$R = v(U - z)W, \qquad T = uk(U - z)W. \tag{LII14}$$

The following identities allocate every term in (LII12)–(LII13):

$$\begin{aligned} zv(V - y)W &= \frac{uVQ_n - uvkR_h}{s} - (V - y)R, \\ uv(V - y)W &= \frac{VQ_n - vkR_h}{s}, \\ uk(V - y)W &= \frac{vkQ_n - yR_h}{s}, \\ U(U - z)(V - y)W^2 &= -\frac{R_hT - uRQ_n}{s}, \\ U(V - y)^2W^2 &= \frac{VQ_n^2 - 2vkR_hQ_n + yR_h^2}{s^2}. \end{aligned} \tag{LII15}$$

Together with $v(U - z)W = R$, $uv(U - z)W = uR$, and $uk(U - z)W = T$, these are an explicit complete allocation of the full remainder as a bounded symmetric Gram in

$$(N, \xi, \zeta, R_h, R, T). \tag{LII16}$$

The original first state normals are all present. The scalar restriction has not been used as a substitute for them.

The finite block and the missing pivot. The residue of Q_0 on the first four roots is

$$\operatorname{diag}(4\ell_*, 2, 4\alpha_*, 4g_0). \tag{LII17}$$

For example the signed core at $W = 0$ is exactly $2(\xi + u\zeta - uv\eta)^2 + 2(1 + U)V\xi^2 + 2(1 + V)\eta^2$; all its other entries vanish in the present residue. The extra terms in the first three lines of (LII15) have infinitesimal Gram coefficients except for $A_\xi \xi R$ and the allowed $A_N NT$. Likewise the last line has only infinitesimal coefficients, and the fourth line has one potentially finite R_hT entry. Thus, apart from entries in the last row and column T, the sole finite off-diagonal entry is between ξ and R.

The exact coefficient extraction yields

$$\operatorname{st} A_\xi = \frac{4}{h(1 + h)}, \qquad \operatorname{st}(d_s/z^2) = \frac{2}{h^2(1 + h)}. \tag{LII18}$$

For a short independent check of the first value, the unscaled original first-normal coefficient on the $V = y$ slice is

$$-4nzr(1 + y) + 4m(1 - H) - (4D + 2)yj + 2(1 + y) \longrightarrow 2 - 4d_0^{\text{pair}} = \frac{2}{1 + h}.$$

Its full multiplier is $2/B$. Passing from the original normal to (LII8) changes this U-quotient only by a bounded multiple of U or z, so its residue is unchanged. The second value follows from (LII1):

$$\operatorname{st}\{m(B + zr)^2/z^2 + (n + \tfrac{1}{2})r^2\} = \frac{1 - r_0}{2(1 + h)} + \frac{r_0}{2(1 + h)} = \frac{1}{2(1 + h)}.$$

Hence the finite (ξ, R) Gram block is exactly

$$\begin{pmatrix} 2 & \dfrac{2}{h(1+h)} \\ \dfrac{2}{h(1+h)} & \dfrac{2}{h^2(1+h)} \end{pmatrix}, \qquad \text{Schur complement} = \frac{2}{h(1+h)^2} > 0. \tag{LII19}$$

This calculation retains $r_0 = 0$, $n = 0$, and $h = 1$.

Finally the exact toric conductor is the zero identity

$$\boxed{T^2 - UR^2 - \frac{U-z}{s} R_h T + \frac{u(U-z)}{s} R Q_n = 0.} \tag{LII20}$$

Add any positive real multiple Λ of this zero identity to the Gram. Its only nonzero residue contribution is ΛT^2. The five-by-five block excluding T is strictly positive by (LII17)–(LII19); all entries incident to T are bounded. Choose a sufficiently large real Λ by the ordinary real Schur complement. The entire six-by-six residue Gram is then strictly positive. The bounded Gram lemma gives a tensor SOS for G, and adding the four packets (LII5) proves the theorem.

Quantifiers and endpoint discipline. No proof step divides by u, v, $U - z$, or $V - y$: parity quotients and monic polynomial divisions are coefficient extractions whose final identities are polynomial. Thus the two literal affine axes are included. The source inverses $1/B$, $1/k$, and the source packet square roots are legal even when their values are large; all coefficients of the final Gram are bounded. The source ratios $t, j, r, 1 - H$ have not been bounded away from zero. The new roots raise testing degree, so the local certificate is not homogenized to assert the original literal projective endpoints. Those are joined by the separate direct-parent remote certificate.

Checkable scope. `check_local_strong_ii.py` checks (LII4), (LII7), (LII15), (LII20), the complete bounded remainder's parity and support, its scalar $U = 0$ value, its hard residue, and all three values in (LII18)–(LII19). It prints the exact extracted polynomials in `extracted_coefficients.txt`. Source contact and stationarity are inherited from the six exact original source checks in `analysis_atlas/oct19_r6_c2_double/check_common_ii.py`; no additional generic source substitution is claimed here.

Chapter 41

The entire remote complement of strong C2 II

Component `p590112f102bd`
Source: `R13/advances/c2_strong_remote/strong_ii_remote.tex`
SHA256: `590112f102bd24dd613da1a20c6032dec181049b2680e6aabe8c26b6f35a30c4`
Inventory use: RR3–RR13 exact full-parent identities | Complete original homogeneous remote parent and all state/projective endpoints | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

41.1 The entire remote complement of strong C2 II

This section is a full-parent tensor-SOS admission. It applies to the original C2 positive chart with z, y infinitesimal and $B = H + L$ a positive unit, and to every testing pair for which at least one testing angle is not infinitesimal. Literal testing axes and projective infinities are included in the assertions below. The local case in which both testing angles are infinitesimal is not asserted here.

Source constants, including all openings. Put $t = L/y$, $j = 1 + zr - t$, and write $h = \operatorname{res} H$. Then $0 < h \le 1$, $t + j = 1 + zr \sim 1$, and the exact source formulas give

$$Cyt \sim Dyj \sim \frac{h}{2(1+h)}, \qquad \frac{A}{y} \sim \frac{1}{2h(1+h)}.$$

Consequently C, D are unbounded, and the positive pure-source quantity

$$\gamma = \frac{yCD}{C+D} = \frac{(Cyt)(Dyj)}{(Cyt)j + (Dyj)t} \sim \frac{h}{2(1+h)}. \tag{RR1}$$

These statements are uniform when either t or j tends to zero. If $h < 1$, then

$$\alpha \sim \frac{h^2}{2(1-h^2)};$$

if $h = 1$, then α is unbounded. Thus, for every real $0 < h_0 < h$, the positive real number

$$\beta_0 = \frac{h_0^2}{2(1-h_0^2)} < \alpha \tag{RR2}$$

is an admissible cap. No lower bound on $r, j, t, 1 - H$ is imposed, and $n = 0$ is retained.

Homogeneous testing coordinates. For finite u, v define

$$a = \frac{u}{\sqrt{1+u^2}}, \quad b = \frac{1}{\sqrt{1+u^2}}, \qquad c = \frac{v}{\sqrt{1+v^2}}, \quad d = \frac{1}{\sqrt{1+v^2}}.$$

Use the same homogeneous formulas at infinity, with $(a,b) = (1,0)$ or $(c,d) = (1,0)$. These are the original parent's homogeneous endpoint values. In particular no raised-degree graph is being specialized at an endpoint. Put

$$E = aW - bY, \quad K = aX - bZ, \quad L = bX + aZ, \quad J = dE + cK, \quad Q = cY - dZ.$$

The original parent divided by $4(1+u^2)(1+v^2)$ is exactly

$$nX^2 + m\{bcW - bdX + acY - adZ\}^2 + CE^2 + AY^2 + (D+\tfrac{1}{2})J^2 + (\alpha+\tfrac{1}{2})Q^2 + \tfrac{1}{2}(L^2 - Z^2). \quad \text{(RR3)}$$

At projective endpoints this equation means the original homogeneous form, not a division by an infinite number. All radicals and normalizations used here belong to a single testing field.

Proposition 41.1 (All tests with v not infinitesimal)**.** *For every such v, including $v = \infty$, and every u, including $u = 0, \infty$, the original full parent is tensor-SOS.*

Proof. Here c is a unit. Set $P = bL$, $R_1 = \sqrt{C}E$, and $R_2 = \sqrt{D}J$. Orthogonality gives

$$L^2 - Z^2 = P^2 + 2aKP - b^2K^2, \qquad K = \frac{R_2}{c\sqrt{D}} - \frac{dR_1}{c\sqrt{C}}. \quad \text{(RR4)}$$

Both coefficients on the right are infinitesimal bounded tensors: $1/c$ is bounded in the testing field, while $1/\sqrt{C}$ and $1/\sqrt{D}$ are infinitesimal in the source field. Therefore

$$R_1^2 + R_2^2 + \tfrac{1}{2}P^2 + aKP - \tfrac{1}{2}b^2K^2$$

has a bounded Gram with residue $\mathrm{diag}(1,1,1/2)$. It is tensor-SOS by the strict bounded-Gram lemma. Every other term of (RR3), namely the retained n, m, A, α packets and $J^2/2 + Q^2/2$, is already SOS. This uses all four state coordinates. □

An exact harmonic identity for the remaining remote strip. Now let v be infinitesimal, possibly zero, and let u not be infinitesimal, possibly infinity. Thus a is a unit. Write $D_v = 1 + v^2$, and put

$$s = C + D, \quad \theta = D/s, \quad N = \sqrt{s}(E + \theta vK), \quad V = \frac{vK}{\sqrt{y}}, \quad q = \sqrt{y}Y.$$

No comparison between v and $\sqrt{y}$ is made. The roots may be unbounded; their Gram coefficients below are bounded. The exact pure-source harmonic identity is

$$CD_vE^2 + D(E + vK)^2 = N^2 + \gamma V^2 + Cv^2E^2. \quad \text{(RR5)}$$

Also set $x = bX$, $z_0 = Z$, and $\ell = vY$. The single toric relation needed below is the polynomial identity

$$ax\ell - b^2z_0\ell = bVq. \quad \text{(RR6)}$$

For a real cap $0 < \beta_0 < \alpha$, the original parent divided by $4(1+u^2)$ is the sum of the explicitly positive packets

$$\begin{aligned} nD_vX^2 + m\{b(vW - X) + a(vY - Z)\}^2 + N^2 + Cv^2E^2 \\ + \tfrac{1}{2}(E + vK)^2 + (\alpha - \beta_0)(\ell - z_0)^2 \end{aligned} \quad \text{(RR7)}$$

and the following exact bounded quadratic form:

$$G = \gamma V^2 + \frac{A}{y}D_vq^2 + (\beta_0 + \tfrac{1}{2})(\ell - z_0)^2 + \frac{D_v}{2}(x^2 + 2axz_0 - b^2z_0^2). \quad \text{(RR8)}$$

This decomposition is valid also at $u = \infty$, by the original homogeneous convention. It has used no state graph or state inverse.

Proposition 41.2 (All tests with u not infinitesimal and v infinitesimal)**.** *The full original parent is tensor-SOS for all these tests, including $v = 0$ and $u = \infty$.*

Proof. For a real constant k define

$$M_k = (\beta_0 + \tfrac{1}{2})(\ell - z_0)^2 + \tfrac{1}{2}(x^2 + 2axz_0 - b^2 z_0^2) + k(ax\ell - b^2 z_0 \ell).$$

By (RR6), the exact form (RR8) equals

$$\boxed{G = D_v M_k + \gamma V^2 + \frac{A}{y} D_v q^2 - k D_v b V q - v^2(\beta_0 + \tfrac{1}{2})(\ell - z_0)^2.} \tag{RR9}$$

This is a full-parent identity; every state-normal term is present.

First suppose b has positive residue b_0. Since a is a unit, $0 < b_0 < 1$. Choose $0 < h_0 < h$ sufficiently close to h that

$$b_0(1 + h - h_0) < 1,$$

and take the admissible cap (RR2). There exists a positive real κ satisfying

$$\frac{1}{1 + b_0 h_0} < \kappa < \min\left\{\frac{1}{1 - b_0 h_0}, \frac{1}{b_0(1 + h)}\right\}. \tag{RR10}$$

Set $k = -\kappa$. The Gram of M_k on (x, z_0, ℓ) has first pivot $1/2$, second Schur pivot β_0, and determinant

$$\det M_k = \frac{2\beta_0 b^2 k^2 - (2\beta_0 + 1)(k + 1)^2}{8}. \tag{RR11}$$

Its residue is strictly positive by the first two inequalities in (RR10), because $h_0^2 = 2\beta_0/(2\beta_0 + 1)$. The Gram on (V, q) in (RR9) has positive diagonal residues and determinant residue

$$\frac{1}{4}\left\{\frac{1}{(1 + h)^2} - k^2 b_0^2\right\} > 0. \tag{RR12}$$

Thus both blocks have strictly positive residue. The final term in (RR9) is a bounded infinitesimal perturbation on (x, z_0, ℓ), so the combined bounded Gram remains strict and is tensor-SOS.

If b is infinitesimal, including $b = 0$, choose any positive real cap $\beta_0 < \alpha$ and take $k = -1$. There is now an exact three-square identity

$$M_{-1} = \tfrac{1}{2}\{x + a(z_0 - \ell)\}^2 + \beta_0(z_0 - \ell)^2 + \tfrac{1}{2}(b\ell)^2. \tag{RR13}$$

On these three roots and (V, q), (RR9) has positive diagonal residue; the only off-block term is $D_v b V q$, whose coefficient is infinitesimal. The negative perturbation changes only the $(z_0 - \ell)^2$ coefficient by an infinitesimal. Its Gram is again strict. Formula (RR7) restores all retained positive packets. The argument does not divide by u, v, b, or any state coordinate, so the asserted literal boundaries are included. □

Scope and join. The two propositions cover precisely the complement of the region where both testing angles are infinitesimal, for every strong-II source with B a unit. They preserve independent source and testing fields, arbitrary source openings, testing signs, and literal $n = 0$. Combined with a separately verified full-parent admission of that local region, they close the strong II/HH orbit by the exact C2 reflection. They do not close the B-infinitesimal weak face.

Chapter 42

Full hi retained contact module

Component `p4363feef60dd`
Source: `R13/advances/hi/full_hi_retained_contact_module.tex`
SHA256: `4363feef60ddab557436c46b295097f496817b78aa6339db887fd01d8b9fcac4`
Inventory use: Whole actual n>=0 HI scalar, both complete normals, every R0 opening and strict uniform hard/circuit residues

42.1 Full HI extension by the exact nonnegative n packet

The preceding contact lift extends to every original $n \geq 0$; no additional comparison of source and testing scales is required. Retain all notation in (N0M.2)–(N0M.5), except for the coefficient replacements specified below. To distinguish the original source data from the four toric roots, write A_{src} for the original coefficient A and $B_{\mathrm{src}} = H + yt$ for the original source B. Put

$$\nu = 2nR = \frac{(B_{\mathrm{src}} + R)(1+R)S_c - R\Delta}{\Delta}, \qquad h = H - \nu, \qquad \zeta = R + \nu.$$

The source quantity ν is nonnegative and bounded, even when R is infinitesimal and n is unbounded. Exact source identities are

$$\frac{2A_{\mathrm{src}}H}{qy} = \zeta, \qquad c - d = \frac{y}{2}\{\zeta(1+q) - t\}, \qquad \mu - a_* = \frac{q}{2}\{E - \zeta(1+y)\}. \tag{HI7.1}$$

In addition to all four opening-excess squares $\mathcal{E}$ in (N0M.3), retain the actual nonnegative square packet

$$\mathcal{N} = 4nd_pd_vx^2, \qquad x = X + \frac{R}{H}pvY = -gs + (H+R)TQ + v\sqrt{y}\,N_c - p\sqrt{q}\,N_m. \tag{HI7.2}$$

This packet is zero at the original source contact. Its exact difference from the original $4nd_pd_vX^2$ is

$$\begin{aligned} 4nd_pd_v(X^2 - x^2) = {} & 4\nu d_pd_vg\,TsQ - 2\nu(2H+R)d_pd_vT^2Q^2 \\ & - 4\nu d_pd_vTQ(v\sqrt{y}\,N_c - p\sqrt{q}\,N_m). \end{aligned}$$

Every coefficient on the right is bounded. The unbounded coefficient n occurs only in the retained positive square.

The full exact identity (N0M.4) therefore holds with $F - \mathcal{E}$ replaced by $F - \mathcal{E} - \mathcal{N}$, with the same h_c, h_m, f_A, f_B, and with

$$\begin{aligned} C_s &= 4d_v\left[y\{\zeta(1+q) - d_pg\} + \frac{2dt}{S_c}\delta_v\right], \\ M_s &= 4d_p\left[q\{d_vg - \zeta(1+y)\} - \frac{2\mu E}{S_c}\delta_p\right], \\ \beta_c &= 4h\sqrt{y}\,vd_pd_v, \qquad\qquad \beta_m = -4h\sqrt{q}\,pd_pd_v, \end{aligned}$$

$$f_C = 2\{H^2 - \nu(2H + R)\}d_p d_v, \qquad f_D = 2H(R + \nu)d_p d_v,$$
$$e = 4hd_p d_v g. \tag{HI7.3}$$

In this notation the complete first-normal couplings are $N_c(C_s B + \beta_c C) + N_m(M_s A + \beta_m C)$, and the graph is $\sum_i f_i A_i^2 - eG^2$ on the positive sign chart. Every coefficient is a polynomial in p, v with bounded source coefficients. The normal coupling coefficients are infinitesimal. Their exact contact sums vanish.

Let $r = \bar{R} \in [0, 1]$. The original source formulas give

$$\bar{\nu} = \frac{1 - r}{3 + r}, \qquad \bar{c} = \bar{d} = \bar{\mu} = \bar{a}_* = \frac{1 + r}{2(3 + r)}.$$

Consequently all four graph coefficients have the same strictly positive residue

$$\bar{f}_A = \bar{f}_B = \bar{f}_C = \bar{f}_D = \lambda(r) = \frac{2(1 + r)^2}{3 + r} \in [2/3, 2], \qquad \bar{e} = 4\lambda(r). \tag{HI7.4}$$

The two state-normal diagonal coefficients have residue $4/(3+r) \in [1, 4/3]$. These statements remain valid at $r = 0$, where n can be unbounded.

All exact contact jets (N0M.6) hold unchanged for the new coefficients; also $C_s^* + \beta_c^* = M_s^* + \beta_m^* = 0$. They can either be obtained by subtracting the contact-vanishing packets from the original parent, or checked directly using (HI7.1). The accompanying checker verifies them as rational identities for the complete source, without imposing $n = 0$ and without replacing R by r.

Choose $k = (f_C^* + f_D^*)/2$, now a positive bounded source unit with residue $\lambda(r)$. The exact half-angle contact lift (N0M.8)–(N0M.9) then applies verbatim. The coefficient errors are infinitesimal bounded tensors, the finite circuit supplies strictly positive pivots on every conductor root, and the state-normal block has a strictly positive scalar residue. On the opposite-sign or zero-state charts the direct strict raw Gram argument applies as before. The strict bounded-Gram lemma supplies an exact tensor SOS for $F - \mathcal{E} - \mathcal{N}$; adding the five retained positive packets gives an exact SOS for the unchanged full original parent throughout the critical HI testing corner.

No inversion or order on a mixed tensor is used. The only radicals of state expressions are taken in the pure-testing real closed field after a pure-testing sign choice. The result is a tensor SOS at each testing state, not a claim of an SOS with the original polynomial degree in the state amplitudes. Together with the already established remote tests and literal axes, this supplies the remaining strong C2 HI admission.

Exact verification. `check_full_hi_contact_module.py` first reuses the 18 exact normal-module checks, then verifies the n-packet identity, all new source and contact identities, and the common-residue formulas. The finite-radical bounded lift is proved above and in the independently reviewed four-term contact-lift lemma.

Chapter 43

Retained contact module

Component `p888d9f79ba3b`
Source: `R13/advances/hi/retained_contact_module.tex`
SHA256: `888d9f79ba3b02bd18fe6e06974a16d9ff598ff433f31d3dd360c7d9e4fe2407`
Inventory use: N0M exact bounded retained remainder, four-monomial circuit, full half-angle conductor and complete normals; only construction imported by full-n theorem

43.1 Literal $n = 0$ HI: a retained four-term contact module

All source and testing variables in this section retain their original meanings. In particular $q, y, p, v > 0$ are independently infinitesimal; no comparison between a source scale and a testing scale is made. Write

$$d_p = 1+p^2, \quad d_v = 1+v^2, \quad \delta_p = p^2-q, \quad \delta_v = v^2-y, \quad S_c = 1+R+yt = j+t(1+y) = K+E(1+q).$$

Use bounded source units

$$c = Cyt, \qquad d = Dyj, \qquad \mu = mqK, \qquad a_* = \alpha qE.$$

At literal $n = 0$, the exact source formulas give

$$2AH/(qy) = R, \qquad \bar{R} = 1, \tag{43.1}$$

$$c - d = \frac{y}{2}\{R(1+q) - t\}, \qquad \mu - a_* = \frac{q}{2}\{E - R(1+y)\}. \tag{N0M.1}$$

The four starred weights have residue $1/4$, even when any positive opening tends to zero. The last two identities follow from the rational source formulas; before imposing $n = 0$ their respective right sides are $y\{(2n+1)R(1+q) - t\}/2$ and $q\{E - (2n+1)R(1+y)\}/2$.

A common state with exact contact. Set

$$A_0 = W - pY, \qquad B_0 = vY - Z, \qquad S = X - vW - pZ + pvY, \qquad s = -S/S_c, \qquad Q = \frac{\sqrt{qy}}{H}Y.$$

Then $J = d_v A_0 + vS$ and $M = d_p B_0 - pS$. Define

$$N_c = \frac{A_0 - tvs}{\sqrt{y}}, \qquad N_m = \frac{B_0 + Eps}{\sqrt{q}}, \qquad P = \frac{p}{\sqrt{q}}, \quad V = \frac{v}{\sqrt{y}}, \quad T = PV,$$

and the additional contact normals

$$N_D = d_v N_c + t\delta_v V s, \qquad N_M = d_p N_m - E\delta_p P s. \tag{N0M.2}$$

At the actual source contact, $P = V = 1$, $s = Q = \sqrt{q}$, and $N_c = N_m = N_D = N_M = 0$. All definitions are tensor identities; neither P nor V is assumed bounded.

Retain all singular opening excesses. The following four positive squares are retained in the parent:

$$\mathcal{E} = \frac{4cjd_v}{tS_c}N_c^2 + \frac{4dt(1+y)}{jS_c}N_D^2 + \frac{4\mu E(1+q)}{KS_c}N_M^2 + \frac{4a_*Kd_p}{ES_c}N_m^2. \tag{N0M.3}$$

Their coefficients are products of positive pure-source and pure-testing factors, so the displayed terms are actual tensor squares. They need not be bounded. No excess is discarded before collecting the remainder.

Put $g = S_c - tv^2 - Ep^2$. The unchanged full parent has the exact identity

$$F - \mathcal{E} = h_cN_c^2 + h_mN_m^2 - 4\sqrt{qy}\,pvd_pd_vN_cN_m + N_c\{C_sVs + 4H\sqrt{y}\,vd_pd_vTQ\} + N_m\{M_sPs - 4H\sqrt{q}\,pd_pd_vTQ\} + \mathcal{G}, \tag{N0M.4}$$

where

$$\begin{aligned} h_c &= \frac{4d_v}{S_c}\{c(1+y) + dd_v\} + 2yd_v(1+v^2+p^2v^2),\\ h_m &= \frac{4d_p}{S_c}\{\mu d_p + a_*(1+q)\} + 2qd_p(1+p^2+p^2v^2),\\ C_s &= 4d_v\left[y\{R(1+q) - d_pg\} + \frac{2dt}{S_c}\delta_v\right],\\ M_s &= 4d_p\left[q\{d_vg - R(1+y)\} - \frac{2\mu E}{S_c}\delta_p\right]. \end{aligned}$$

The two diagonal coefficients have residue one. The normal–normal off-diagonal coefficient and all four raw coefficients in the first normal line are infinitesimal bounded tensors. In particular no inverse opening occurs in (N0M.4).

The scalar graph is

$$\mathcal{G} = f_AP^2s^2 + f_BV^2s^2 + f_CT^2Q^2 + f_DQ^2 - eTsQ, \tag{N0M.5}$$

with

$$\begin{aligned} f_A &= 4\left[\mu K + a_*Ed_p - 2\mu E\delta_p + \frac{\mu E^2}{S_c}\delta_p^2\right] + 2qd_v(E\delta_p - K + tv^2)^2 - 2E^2qv^2d_p,\\ f_B &= 4\left[ctd_v + dj - 2dt\delta_v + \frac{dt^2}{S_c}\delta_v^2\right] + 2y(t\delta_v - j)^2,\\ f_C &= 2H^2d_pd_v, \qquad f_D = 2HRd_pd_v, \qquad e = 4Hd_pd_vg. \end{aligned}$$

All five are polynomials in p, v with bounded source coefficients. Their residues are, respectively, $2, 2, 2, 2, 8$. This assertion is uniform over every opening permitted by the exact HI source entry.

The exact source contact jets. A star denotes $p = \sqrt{q}$, $v = \sqrt{y}$. Set $\sigma = f_A + f_B + f_C + f_D - e$. The rational source formulas give

$$\begin{aligned} f_A^* + f_B^* &= f_C^* + f_D^* = e^*/2,\\ q^{1/4}(\partial_{\sqrt{p}}\sigma)^* &= -4(f_A^* - f_D^*),\\ y^{1/4}(\partial_{\sqrt{v}}\sigma)^* &= -4(f_B^* - f_D^*). \end{aligned} \tag{N0M.6}$$

Also

$$C_s^* + 4Hy(1+q)(1+y) = 0, \qquad M_s^* - 4Hq(1+q)(1+y) = 0. \tag{N0M.7}$$

These are actual-contact identities, not residue identities. The checker verifies the only source-dependent ones modulo the exact numerator of n, rather than replacing R by its residue.

Finite contact lift of the graph and both state normals. Here the SOS is a tensor SOS at each original pure-testing state. Its roots need not be linear polynomials in W, X, Y, Z. If $sQ > 0$, a common sign change permits $s, Q > 0$. This is a pure-testing sign choice, because sQ is a positive source factor times $-SY$. Put

$$A = Ps, \quad B = Vs, \quad C = TQ, \quad D = Q, \quad G = (ABCD)^{1/4} = \sqrt{TsQ}, \quad E_i = A_i - G.$$

Every A_i and G is a positive decomposable tensor. Thus every radical below is a product of a pure-source radical and a pure-testing radical. The basic circuit is

$$L = A^2 + B^2 + C^2 + D^2 - 4G^2 = \sum_i E_i^2 + 2G \sum_i E_i. \tag{N0M.8}$$

For each pairing $ij|k\ell$,

$$G \sum_i E_i = G(\sqrt{A_i} - \sqrt{A_j})^2 + G(\sqrt{A_k} - \sqrt{A_\ell})^2 + 2G\{(A_iA_j)^{1/4} - (A_kA_\ell)^{1/4}\}^2.$$

Moreover,

$$\begin{aligned} L &= (A - C)^2 + (B - D)^2 + 2(\sqrt{AC} - \sqrt{BD})^2, \\ (r - r')^2 &= (r - G)^2 + (r' - G)^2 + 2G(\sqrt{r} - \sqrt{r'})^2 \quad (r = \sqrt{AC},\ r' = \sqrt{BD}), \end{aligned}$$

and the analogous identities hold for $BC|AD$. A finite average of these decompositions gives strictly positive constant diagonal pivots simultaneously on the E_i, the variance roots in (N0M.8), and

$$F_p = \sqrt{AC} - G, \qquad F_v = \sqrt{BC} - G.$$

The exact half-angle conductors are

$$G(\sqrt{p} - q^{1/4}) = q^{1/4}F_p, \qquad G(\sqrt{v} - y^{1/4}) = y^{1/4}F_v. \tag{N0M.9}$$

Their coefficients are pure-source infinitesimals.

For completeness, the first-jet cancellation is explicit. Choose the positive source unit $k = (f_C^* + f_D^*)/2 \sim 2$. Taylor-expand the bounded coefficient polynomials in $z = \sqrt{p}, w = \sqrt{v}$ around $z_0 = q^{1/4}, w_0 = y^{1/4}$. All divided-difference coefficients remain bounded, since these are monic polynomial divisions by pure-source centers. Explicitly each monomial has the finite identity

$$z^i w^j = \sum_{a=0}^{i} \sum_{b=0}^{j} \binom{i}{a} \binom{j}{b} z_0^{i-a} w_0^{j-b} (z - z_0)^a (w - w_0)^b.$$

Collecting total degrees zero, one, and at least two is the required exact Taylor certificate. Its coefficients are bounded because all exponents are nonnegative and all source coefficient denominators are positive source units. No analytic remainder or mixed division enters. After (N0M.9), every term in $\mathcal{G} - kL$ is an infinitesimal bounded quadratic in the listed roots, except possibly

$$G\left[2\sum_i (f_i^* - k)E_i + z_0\sigma_z^* F_p + w_0\sigma_w^* F_v\right].$$

By (N0M.6) this equals

$$\begin{aligned} 2(f_D^* - k)G \sum_i E_i &- \frac{z_0\sigma_z^*}{2}\{\sqrt{G}(\sqrt{A} - \sqrt{C})\}^2 \\ &- \frac{w_0\sigma_w^*}{2}\{\sqrt{G}(\sqrt{B} - \sqrt{C})\}^2. \end{aligned}$$

All three coefficients are source infinitesimals, and all their roots already have strict pivots in the finite averaged circuit.

Finally the N_c coupling in (N0M.4) is $N_c(C_sB + \beta_c C)$, where $\beta_c = 4H\sqrt{y}\, v d_p d_v$ is infinitesimal. The sum $C_s + \beta_c$ vanishes at the source contact by (N0M.7). Writing $B = G + E_B$, $C = G + E_C$, monic-dividing that coefficient sum in $z - z_0, w - w_0$, and applying (N0M.9) expresses this entire coupling by infinitesimal bounded entries on N_c, E_B, E_C, F_p, F_v. The N_m coupling is identical, using A, C and $\beta_m = -4H\sqrt{q}\, p d_p d_v$. Thus the two state normals have an identity-residue block, and the graph has the strict circuit block. The established strict bounded-Gram absorption lemma now gives an actual tensor SOS of $F - \mathcal{E}$.

If $sQ < 0$, the graph is instead a strict bounded diagonal form on the four raw roots and $\sqrt{T|sQ|}$, with residue coefficients $2, 2, 2, 2, 8$. The first-normal couplings already have infinitesimal raw coefficients. If $sQ = 0$, simply omit the last root. The same bounded-Gram lemma applies directly. Consequently this argument covers literal $n = 0$ throughout the critical HI testing corner, at arbitrary independent source/testing rates and through all positive source openings. It does not assert the corresponding $n > 0$ extension.

Verification scope. `check_retained_contact_module.py` checks the pair excess identities, the complete signed core, the four-term graph, the exact source differences, and all contact-jet identities. Its 18 checks are exact symbolic equalities. The finite-radical lift and the uniform boundedness argument are the analytic argument above, not conclusions drawn from numerical sampling.

Chapter 44

Beta-positive extension and the exact remaining local source branch

Component p909aa5765721
Source: R13/advances/revision13_c1_eta/beta_positive_extension.tex
SHA256: 909aa5765721968d80341f4fb77f53afd99b693ca6386f3c6f0b2ef839d2aa0a
Inventory use: Bounded-Mbeta and bounded-H unit-u admissions | JB1–JB8 and complete bounded-Mbeta theorem | bounded-Mbeta whole-source theorem

44.1 Beta-positive extension and the exact remaining local source branch

Retain the same original C14.1 source as JR1–JR8, with x, z, σ positive infinitesimals, but allow $\beta \geq 0$. Write

$$b_\theta = \beta\theta, \qquad H = h/\sigma, \qquad M_\beta = b_\theta/\sigma^2.$$

This section proves two precise extensions. First, if M_β is bounded, the *entire source domain is admitted for every test.* This includes the beta-zero theorem. Second, if H is bounded but M_β is unbounded, every bounded nonzero-unit u is admitted at the residual infinitesimal-w tests. On this second source domain only u infinitesimal remains. Neither statement assumes d_0 a unit: the second includes $D = 0$. The H-unbounded domain and C14.2 are not admitted here.

The generic joint radial identity. The exact six-root identity JR5 continues to hold if its coefficients are given the following beta corrections:

$$\begin{aligned} k_1 &\mapsto k_1 + 4b_\theta\theta, \qquad R_x \mapsto R_x + 8b_\theta, \\ Z_0 &\mapsto Z_0 + \frac{4b_\theta\theta t}{\sigma^2}, \qquad Z_x \mapsto Z_x + 8ub_\theta/\sigma, \\ S &\mapsto S + 4b_\theta\theta/\sigma^2. \end{aligned} \tag{JB1}$$

Here the uncorrected formulas on the right are the polynomials in JR2–JR3 evaluated at the actual h, D, m, p, l, not at a beta-zero source. The scalar B remains its full monic quotient. The additional retained centered excess is $4\beta D_w N_x^2$. The complete generic coefficient is, explicitly,

$$S = \frac{2\{h^2(2D+1)(t+x) + 4Dh - 2bp\sigma^2 z + 2b_\theta\theta + h^2\}}{\sigma^2}. \tag{JB2}$$

All four cancellations JR4 are exact with these changes. They were checked symbolically with b_θ free. Thus no radial pole remains unallocated in this extension.

The source estimates needed by the extension. The source equation BE3 gives exactly

$$H = H_0/B + \frac{2P}{B}\frac{b_\theta}{\sigma}. \tag{JB3}$$

Consequently M_β bounded implies $H \sim 1$ and $\theta \sim 1$. The sphere budget then implies $D/\sigma^2 \sim 1/2$, since $xb_\theta/\sigma^2 = xM_\beta$ is infinitesimal. Also $\mu \sim 1/2$.

More generally, suppose only H bounded. Then $\operatorname{res} H \geq 1$; no exact inequality $H \geq 1$ is imposed. One has $\theta \sim 1$, and b_θ/σ is bounded. Write

$$d = \operatorname{res}(D/\sigma^2), \qquad c = \operatorname{res}(xb_\theta\theta/\sigma^2).$$

BE4–BE6 give $d, c \geq 0$ and $d + c = 1/2$. Formula JB3 gives $\operatorname{res}(b_\theta/\sigma) = (\operatorname{res} H - 1)/2$. The bounded source formulas also give $\mu \sim H/2$, $p, l \sim 1/2$, $C = O(z)$, and $p - l \in z\mathcal{O}_{\text{source}}$. These statements retain both source-sphere endpoints and both b, Q openings.

An exact square joining the beta pole to its angular reserve. Suppose first u is a bounded nonzero unit, so that $1/u$ is an allowed bounded pure-testing inverse. Put

$$a_\beta = b_\theta\theta, \qquad C_\beta = 4a_\beta/\sigma^2, \qquad T_c = t\Gamma + V, \qquad P_\beta = uZ - (x/\sigma)wW.$$

The identity $T_c = \sigma u P_\beta$ has no u denominator. Adding (C_β/t) times JR6's zero, and completing the resulting T_c square, gives the exact entire-parent identity

$$G = 4a_\beta P_\beta^2 + \widehat{G}, \tag{JB4}$$

where G is the complete generic JR5 and

$$\begin{aligned}\widehat{G} = G &- C_\beta t\Gamma^2 - (C_\beta + C_\beta x/t)\Gamma V - C_\beta\sigma A_\theta^2 \\ &- C_\beta\sigma(t - x)A_\theta R/u.\end{aligned} \tag{JB5}$$

Every coefficient of $\widehat{G}$ is bounded when H is bounded and u a unit. The only apparent angular reserve subtraction is $4a_\beta/\sigma$. Its exact source comparison is

$$\mu - a_\beta/\sigma = \kappa H_0/B + \frac{b_\theta}{\sigma}\left(\frac{2\kappa P}{B} - \theta\right), \tag{JB6}$$

where κ is the BE8 source factor. The bracket is infinitesimal, so this difference has residue $1/2$. Thus the beta pole is paid for by the actual shared m packet. No leading coefficient is substituted in the exact identity JB4.

Bounded M_β: all residual tests. When M_β is bounded and u a unit, the residue of $\widehat{G}$ is exactly the beta-zero residue in JR7 before the finite V pivot: $H \sim 1$, $d = 1/2$, $c = 0$. Add the same pure-testing pivot $\lambda(t) = 2(t^2 + t + 1)/(1 + t)^2$ and apply JR8. Restore the positive square in JB4 and every centered excess. This covers every unit u.

When u is infinitesimal, use the generic six-root form directly, without $1/u$. Its residue has five positive diagonal pivots equal to two, with the only cross term

$$(2 + 4M_0)\Gamma V, \qquad M_0 = \operatorname{res} M_\beta \geq 0.$$

Add JR6's zero with the bounded pure-source coefficient $\lambda = (1 + 2M_\beta)^2$. All its non-$V^2$ corrections have zero residue. The (Γ, V) Schur complement is $(1 + 2M_0)^2/2 > 0$. This includes literal $u = 0$ and either sign. The exact remote join is the CE9–CE10 and CE20–CE21 join printed after JR8; it applies unchanged here because D, β were unused. This proves the full all-tests admission under the sole extra pure-source condition $\beta\theta/\sigma^2$ bounded.

Unbounded M_β, bounded H: all unit-Alice tests. For any real nonzero residue u_0, set $t_0 = u_0^2$. In the notation of JB3, and writing H now for its finite real residue, the only nontrivial $\widehat{G}$ residue block is

$$\begin{aligned} Q = {} & 2(1+t_0)N_x^2 + [4d + 2H^2 t_0(1+t_0)]\Gamma^2 \\ & - 4u_0(1+Ht_0)N_x\Gamma - 4Hu_0 N_x V \\ & + [2H^2(1+t_0) - 4c/t_0]\Gamma V. \end{aligned} \tag{JB7}$$

Its angular (A_θ, R) block is the strict matrix from JR7 and its N_y pivot is $2(1+t_0)$. All other residue cross entries vanish. The beta square in JB4 equals $(C_\beta/t)T_c^2$, with C_β unbounded. On its exact residue kernel $V = -t_0\Gamma$, JB7 reduces to

$$2(1+t_0)N_x^2 - 4u_0 N_x\Gamma + 2\Gamma^2, \qquad \det = 4. \tag{JB8}$$

Here the cancellation uses exactly $d + c = 1/2$. It remains strict at $d = 0$ and $c = 0$. The row $T_c = V + t\Gamma$ has a bounded separated right inverse, obtained by changing V alone. The elementary finite penalty lemma therefore chooses a positive real K such that $\widehat{G} + (K/t)T_c^2$ has strictly positive residue. This is possible because $t_0 > 0$. The exact retained excess is $((C_\beta - K)/t)T_c^2$: $C_\beta - K$ is a positive pure-source coefficient, and its square root separates from the pure-testing $1/\sqrt{t}$. No ordering of C_β/t minus a real number is used. This proves the entire parent for all bounded nonzero-unit u, with infinitesimal or zero w, all signs and complex states.

The new exact remaining obstruction on the bounded-H source branch is therefore M_β unbounded and u infinitesimal. JB5 has x/u and x/u^2 terms there, so its bounded-Gram argument cannot be imported. The square P_β itself is still legal and has no u inverse; a joint allocation with the remaining angular terms is required to remove those coefficient denominators without comparing independent source and testing rates. No such allocation is claimed.

Denominator and boundedness audit. Every source inverse in the roots or exact coefficients is a pure-source inverse. On the large-L, bounded-H domains, P, J, B and the original geometry coefficient $M = 1 + xbz - xz\theta$ are units; $1 - r$ and θ are units as well. Possible b, Q inverses occur only in the original source weights and the retained positive centered excesses, while their opening-cleared products p, l are bounded. The normalized quantities H, D/σ^2, b_θ/σ, and $xb_\theta\theta/\sigma^2$ are bounded by JB3 and the BE sphere. The corrections in JB5 therefore leave bounded coefficients: the remaining Γ^2 terms involve D/σ^2, Dh/σ^2 and H^2; the ΓV coefficient involves those same bounded source products and $(xb_\theta\theta/\sigma^2)/t$; the angular-radial coefficient involves $b_\theta\theta/\sigma$ times $(t-x)/u$. The factors $1/u, 1/t$ are used only when u is a testing unit. All $u = 0$ and infinitesimal-u assertions for bounded M_β instead use the unaltered generic six-root identity and its bounded pure-source pivot $(1 + 2M_\beta)^2$. The everywhere bounded testing denominator $1 + t = 1 + u^2$ in JR8 is also pure testing. No mixed inverse, mixed radical, or source/testing rate comparison occurs in either admitted domain or its remote join.

Chapter 45

A complete joint radial allocation on the literal beta-zero face

Component `pb1104217f67b`
Source: `R13/advances/revision13_c1_eta/joint_radial_beta_zero.tex`
SHA256: `b1104217f67bbdd2b1f875599ad2951b855cc402f2ba9f112d1e42c0e9cd24ca`
Inventory use: CE9-CE10/CE20-CE21 literal, sign and remote join | JR1–JR8 and exact CE remote join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

45.1 A complete joint radial allocation on the literal beta-zero face

Quantified domain and conclusion. Retain the original physical C1.4 source and all its residual-complement conditions. Suppose $x = k^2$, $z = L^{-1}$ and $\sigma = \eta^{-1}$ are positive infinitesimals, $0 < b < 1$, and the original coefficient $\beta = 0$. There are no further source hypotheses. In particular $h/\sigma \sim 1$ and $D/\sigma^2 \sim 1/2$ follow from BE3–BE7, rather than being imposed. The conclusion below covers every bounded independent real test u, every infinitesimal or zero independent reciprocal test w, all their signs, and real or complex states. The previously proved remote complement supplies the other tests on this large-L source chart. Thus this proves the literal $\beta = 0$ part of C14.1. The adjoining Section 44.1 extends the actual admission to all sources with $\beta\theta/\sigma^2$ bounded; the present result is its beta-zero corollary and supplies the residue calculation used there. C14.2 is not included.

The calculation joins the entire radial scalar quotient with every radial first normal *before* deciding boundedness. Their exact cancellations remove the BR3 pole, including its separate $8(ab)b/\sigma$ summand through $Q = 1 - b$ infinitesimal.

Notation and the original contact identities. Use the complete reciprocal parent of BE10 and the large-L source notation. Thus $m = \mu x^2 z$, $C = cz$, $A = a/z$, and $\alpha = \ell/z$. In expressions below $p = ab$ and $l = \ell Q$ are the two opening-cleared products; p is not the source parameter b. Set $t = u^2$, $\xi = w^2$, $\xi_0 = \sigma z$, $\delta = \xi - \xi_0$, $g = h - Q$, $D_u = 1 + t$, $D_w = 1 + \xi$, and $q = \sqrt{z}$. The source identities established in the handoff give

$$\phi = \phi_t = \phi_\xi = L_x = L_y = L_z = 0 \quad \text{at } (t, \xi) = (x, \xi_0),$$

as well as $\phi(0, \xi) = 4m$ and $L_x(0, \xi) = -8m$. Here ϕ, L_x, L_y, L_z are the *complete* EN2 polynomials, with $D_* = D$, $\beta_* = 0$ and $\beta = 0$.

Choose the fixed caps $a_* = \ell_* = 1/4$. They are smaller than the actual a, ℓ, since $p, l \sim 1/2$ and $0 < b, Q < 1$. Retain the two exact nonnegative centered excesses

$$E_{\text{cap}} = 4(a - a_*)D_u D_w N_y^2 + 4(\ell - \ell_*)D_u N_y^2.$$

Define six separated roots

$$\begin{aligned} N_x &= X + uhZ, \qquad N_y = (Y - bwZ)/q, \\ \Gamma &= \sigma Z - uwW, \qquad R = wZ/q - uqW, \\ A_\theta &= q(t-x)W, \qquad V = uw(t-x)W. \end{aligned} \tag{JR1}$$

No test/source quotient is used in their definition. Their inverses and radicals are pure source.

Exact quotient specification. The contact identities give the exact divisions

$$\begin{aligned} \phi &= z(t-x)^2 A + t(t-x)\delta B + t\delta^2 C_r, \\ L_x &= (t-x)(tA_x + zB_x) + t\delta C_x, \\ L_y &= (t-x)A_y + \delta B_y, \\ L_z &= (t-x)A_z + \delta B_z. \end{aligned}$$

All quotients are monic polynomial divisions in one testing variable. Their complete support can equivalently be specified by

$$\begin{aligned} A = 4\mu + \frac{2h^2(2D+1)}{\sigma}t, \qquad A_x = -\frac{4h(2D+1)}{\sigma}, \qquad B_x = 8\mu x, \\ A_y = 8(p-l) + 4z\{(1+2m)g + 2Dh + 2p\sigma\}, \\ S := A_z/\sigma - B = S_0 + \frac{2h^2(2D+1)}{\sigma^2}(t-x). \end{aligned} \tag{JR2}$$

The same monic divisions give the much shorter exact expression

$$S_0 = \frac{2\{2h^2(2D+1)x + 4Dh - 2bp\sigma^2 z + h^2\}}{\sigma^2}. \tag{JR2a}$$

Thus $S_0 \sim 2$ follows immediately from $h/\sigma \sim 1$ and $D/\sigma^2 \sim 1/2$. This removes any dependency on the longer R10 reduction modulo the original beta numerator. The source contact jets are still needed for the quotient remainders to vanish. For a denominator-free quotient specification, C_r is obtained by dividing ϕ twice by $\xi - \xi_0$ and then dividing the quotient by t; B is the quotient of $t^{-1}\phi_\xi(t, \xi_0)$ by $t - x$; and A_z is the quotient of $L_z(t, \xi_0)$ by $t - x$. These divisions introduce no mixed inverse.

The remaining exact coefficients are most compactly written as

$$\begin{aligned} K_r := \sigma^2 C_r &= k_0 + k_1 t + k_2 \xi, \\ k_0 &= 2b\{(2C+1)b + 4D + 4p\sigma + 2(2C+2D+1)\sigma(bz-1)\}, \\ k_1 &= 2(1+2m)g^2 + 8Dhb + 8pb\sigma, \\ k_2 &= 2b^2(2C+2D+1), \\ R_x := \sigma C_x &= -4\{2Db + (1+2m)g\}, \\ R_y := \sigma B_y &= 4\{t[2Dh + 2p\sigma + (1+2m)g] + b(2C+2D+1)(\xi + \sigma z) \\ &\qquad + (2C+1)(b-\sigma) + 2D(1-\sigma) + 2p\sigma\}, \\ E_r &= 2(C+D)(bz-1) + 2p + bz - 1, \qquad R_\gamma = -4(E_r + 2pt), \\ Z_0 &= \frac{4D + 2t(4Dh + h^2) + 2t^2h^2(2D+1)}{\sigma^2}, \\ Z_x &= -\frac{4u}{\sigma}\{2D + h + th(2D+1)\}. \end{aligned} \tag{JR3}$$

In particular all of these coefficients are bounded when u is bounded and w infinitesimal. The σ^{-1} and σ^{-2} in Z_0, Z_x are controlled by the *proved pure-source* bounds on h, D.

The joint cancellation, including all first normals. The exact full normal block EN3–EN4 obeys

$$\begin{aligned} H_z/\sigma^2 + \xi C_r - (\xi/\sigma)B_z &= Z_0,\\ B_z - 2\sigma C_r &= bR_\gamma,\\ \frac{4}{\sigma}\{2DT_{\mathrm{test}} + bD_w(1+2C) + tg(1+2m)\} - B_y &= R_\gamma,\\ \frac{4u}{\sigma}\{-2DT_{\mathrm{test}} - hD_u - (1+2m)\xi g\} - u\xi C_x &= Z_x. \end{aligned} \tag{JR4}$$

Here $T_{\mathrm{test}} = 1 + th + b\xi$. These are polynomial identities after clearing pure-source powers of σ. No coefficient is discarded: K_r contains the full BR3 term, and the last three lines keep its radial couplings. The cancellation explains why a scalar-only extraction of BR1 followed by boundedness checking cannot see this certificate.

Put

$$\begin{aligned} H_x &= 2(1+t+\xi) + 4Dt + 4m\xi,\\ H_y &= 4D_u(a_*D_w + \ell_*) + 4mtz + 4CzD_w + (4D+2)\xi z + 2D_u z,\\ c_G &= 4D_u\{pbD_w + lQ\}, \qquad c_N = 8D_u\{pD_w - l\}. \end{aligned}$$

The *entire original parent*, without restriction to a state subspace, is exactly $F_w = E_{\mathrm{cap}} + G$, where

$$\begin{aligned} G = {} & H_xN_x^2 + H_yN_y^2 + Z_0\Gamma^2 + (c_G + zK_r)R^2 + AA_\theta^2\\ & - 4uqw(2m+2D+1)N_xN_y + Z_xN_x\Gamma + qwR_\gamma N_y\Gamma + qwbR_\gamma R\Gamma\\ & + uqwR_xN_xR + (c_N + zR_y)N_yR + qwB_xN_xA_\theta + uA_yN_yA_\theta\\ & + uA_xN_xV + u\sigma BA_\theta R + S\Gamma V. \end{aligned} \tag{JR5}$$

For example, the scalar radial conductor is $t\delta^2W^2 = \sigma^2(qR - w\Gamma/\sigma)^2$; the radial Z first normal is $uw\delta W(\Gamma/\sigma) = qwR\Gamma - (\xi/\sigma)\Gamma^2$. Their joint use gives the first two identities of JR4. The other conductors are those of the verified large-L proof with $N_z = \Gamma/\sigma$. These identities prove JR5 exactly and include every first normal and every hard quadratic coupling.

One bounded positive Gram for every bounded Alice test. All entries of JR5 are bounded separated tensors. Indeed

$$h/\sigma \sim 1, \quad D/\sigma^2 \sim 1/2, \quad \mu \sim 1/2, \quad C = O(z), \quad p, l \sim 1/2, \quad p - l \in z\mathcal{O}_{\mathrm{source}},$$

and $\sigma B = A_z - \sigma S \sim 4$ uniformly for bounded u. The displayed formula for A_z obtained from L_z has only the bounded terms h^2/σ, Dh/σ, p, l, and bounded multiples of z; hence this last assertion does not order any mixed rate.

For completeness, the possibly needed opening-uniform divisibility $p - l \in z\mathcal{O}_{\mathrm{source}}$ follows directly from the exact source formulas, not from a residue subtraction. Write $H = h/\sigma$. The two numerators over their common unit denominator $2P$ are

$$\begin{aligned} U_A &= z(1+x\sigma H)(1-xH) + (1-bz)(1+x\sigma Hz),\\ U_\ell &= (1+\sigma bz)\big[xH(1+xbz)(1+\sigma z) + \{1+xbz - xz(1-\sigma H)\}(1-xH)\big]. \end{aligned}$$

They are bounded polynomials in the source letters and both equal one at $z = 0$. Consequently $(U_A - U_\ell)/z$ is a bounded polynomial. Neither numerator has a b or Q denominator. This proves the claimed divisibility through both openings and is the source argument in EN6's adjoining radial paragraph.

The exact syzygy

$$V^2 = \sigma tA_\theta^2 + u\sigma(t-x)A_\theta R - (t-x)V\Gamma \tag{JR6}$$

allows the addition of zero times the pure-testing function

$$\lambda(t) = \frac{2(t^2+t+1)}{(1+t)^2} > 0.$$

Specifically add $\lambda[V^2 - \sigma t A_\theta^2 - u\sigma(t-x)A_\theta R + (t-x)V\Gamma]$ to JR5. There is no source/test denominator in λ.

At any real residue u_0, write $t_0 = u_0^2$. The N_y pivot is $2(1+t_0)$. The (A_θ, R) block is

$$\begin{pmatrix} 2 & 2u_0 \\ 2u_0 & 2(1+t_0) \end{pmatrix}, \qquad \det = 4.$$

The remaining (N_x, Γ, V) block is

$$\begin{pmatrix} 2(1+t_0) & -2u_0(1+t_0) & -2u_0 \\ -2u_0(1+t_0) & 2(1+t_0+t_0^2) & 1+t_0+\lambda(t_0)t_0/2 \\ -2u_0 & 1+t_0+\lambda(t_0)t_0/2 & \lambda(t_0) \end{pmatrix}. \tag{JR7}$$

Its first two leading pivots are positive, with leading 2×2 determinant $4(1+t_0)$. The last Schur complement is exactly

$$\lambda(1+t_0^2) + \frac{2t_0^2}{1+t_0} - \frac{1}{2}(1+t_0+\lambda t_0/2)^2 = \frac{4t_0+3}{2(1+t_0)^4} > 0. \tag{JR8}$$

Thus the residue Gram is strictly positive definite for every finite real u_0, including zero and both signs. Crucially it does not depend on the residue of b: the result remains strict through $b \to 0$ and $Q \to 0$. The finite bounded positive-Gram criterion already used by the project then gives the exact legal tensor-SOS.

Boundaries and scope. JR1–JR6 are identities, including $u = 0$ and $w = 0$; no root is inverted at a zero. Their sign coverage is literal. Apply the real identity to the real and imaginary parts of the state for complex states. The exact remote join is the common-parent theorem in `baseline/advances/c1_common/common_remote.tex`, Section `sec:r6-c1-common-remote`. First, A, α are unbounded on this source, so its adjacent-event theorem transported by (CE9)–(CE10) covers every v that is not projectively unbounded. For projectively unbounded v and projectively unbounded u, put $p_t = 1/u$ and apply the complete raw frame (CE20), whose strict residue is (CE21). Its exact source premises (CE17), without the unneeded m lower bound in this p_t chart, hold with $q = \sqrt{z}$: $C \sim z/2$, $A \geq 1/(4z)$, and $\alpha \geq 1/(4z)$. The C lower bound may likewise be chosen as $z/4$. The unselected D, β events are retained. The identities include literal $p_t = 0$, $w = 0$ and all signs. These two remote cases and JR1–JR8 exhaust all projective tests on the printed source face. No source domain with L a unit, or $\beta > 0$, is admitted by this proof.

Corroboration. The accompanying focused checker verifies the complete joint polynomial allocation, the four cancellations, every conductor, the short exact S coefficient, and the rational Schur-complement identity. The original six contact identities are explicitly identified imported results, not silently replaced by numerical residue checks.

Chapter 46

Small-L, unbounded eta: uniform source weights

Component p66922bb7f235
Source: R13/advances/revision13_c1_small_l/eta_large_bounds.tex
SHA256: 66922bb7f235f4ea968033105881fd80854a5e65dc0223bf372d0969701ccc0a
Inventory use: ES source bounds | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

46.1 Small-L, unbounded eta: uniform source weights

This is a source theorem, not an additional all-tests admission. Use the original physical C1 source with $x, L = x\zeta > 0$ infinitesimal, ζ noninfinitesimal and η unbounded. Put $\sigma = 1/\eta$, $S = \sigma + L$, and retain the original $0 < r, h < 1$, $\theta = 1 - h$, $M > 0$, and $D, \beta \geq 0$. The native gap gives $E = 1 - xh/\sigma \geq 1/18$. Set

$$e = \frac{xh}{\sigma} = 1 - E, \quad P = \sigma P_d, \quad J = 1 + xh - \sigma(1 - r), \quad H_0 = 1 + L(1 - r), \quad B = P + x(J - \sigma),$$

$$\mathcal{H} = RH_0 + (1 - r)xh(2 + xr) - e\{2 + x(1 + h + r\theta)\} + Ke^2, \quad R = 1 + xr, \quad K = 1 + x. \tag{ES1}$$

No $E \sim 1$ assumption is made here. The original source formulas give the two exact identities

$$2\theta P\beta = hB - \sigma H_0, \qquad 2P(JD + x\theta\beta) = \sigma^2\mathcal{H}. \tag{ES2}$$

They are the beta-shared identities with the reciprocal-L factors cleared. In particular their use does not require bounded $1/L$.

The positive denominator is exactly

$$P = \sigma(1 + L + 2xh + rxL + rx^2h) + KL + x^2h\theta. \tag{ES3}$$

Since $x/L = 1/\zeta$ is bounded and $x^2h = x\sigma e$, all terms beyond $\sigma + L$ in (ES3), divided by S, are infinitesimal. Consequently

$$\frac{P}{S} \sim 1, \qquad J \sim 1, \qquad \frac{B}{S} - (1 + x/S) \in \mathrm{Inf}, \qquad H_0 \sim 1, \qquad \mathcal{H} - E^2 \in \mathrm{Inf}\,. \tag{ES4}$$

Here x/S is bounded. Both B/S and $\mathcal{H}$ are positive bounded units, uniformly through all original openings; for the latter use the actual bound $E \geq 1/18$.

Define nonnegative pure-source sphere coordinates

$$d_0^2 = \frac{2PJD}{\sigma^2\mathcal{H}}, \qquad b_0^2 = \frac{2Px\theta\beta}{\sigma^2\mathcal{H}}.$$

Then

$$d_0^2 + b_0^2 = 1, \qquad D = \frac{\sigma^2 \mathcal{H}}{2PJ} d_0^2, \qquad x\theta\beta = \frac{\sigma^2 \mathcal{H}}{2P} b_0^2. \tag{ES5}$$

Thus $D = O(\sigma^2/S) = O(\sigma)$ and $x\theta\beta = O(\sigma^2/S)$, including either literal coefficient boundary. The first identity of (ES2) becomes

$$eB = xH_0 + \sigma \mathcal{H} b_0^2. \tag{ES6}$$

Put $\kappa = x/S$ and $\nu = \sigma/S$. Multiplying $e(B/S) = \kappa H_0 + \nu \mathcal{H} b_0^2$ by e and using $e\nu = \kappa h$ gives

$$e^2(B/S) = \kappa\{eH_0 + h\mathcal{H} b_0^2\}. \tag{ES7}$$

All factors in braces are bounded and nonnegative. Hence $e^2 = O(x/S)$, and (ES6) also gives $x/S = O(e)$. In particular ζ unbounded implies e infinitesimal, since $x/S \le 1/\zeta$. The unit-ζ cases may have e a positive bounded unit; (SN7) and (SN12) are not imported.

The other original coefficients have the opening-cleared bounds

$$\frac{CS(1-r)}{E} \sim \tfrac{1}{2}, \qquad \frac{ASr}{E} \sim \tfrac{1}{2}, \qquad m = \frac{K(1+\sigma r)eS}{2M\zeta P}, \tag{ES8}$$

$$\alpha = \frac{(1+\sigma r)\{LReS + L^2 ME\}}{2QP} = O(L), \qquad Q = 1 - Lr \sim 1. \tag{ES9}$$

For A use $P_4 = (1+xh)E + (1-r)(xh+L) \sim E$. The bound on α follows because $0 < M < R$ is bounded and $L/S \le 1$. Thus C, A are unbounded while D, α are infinitesimal on the entire stated source domain. No bound on m is asserted through $M = 0$, and no bound on beta is asserted through $\theta = 0$.

Finally put $\mu = K(1+\sigma r)S/(2M\zeta P) > 0$. The original m packet has the exact shared-sphere split

$$m = m_0 + m_1 b_0^2, \qquad m_0 = \frac{\mu x H_0}{B}, \qquad m_1 = \frac{\mu\sigma\mathcal{H}}{B}. \tag{ES10}$$

The remaining source opening M is displayed explicitly. These identities provide valid weights for C14.4 and the boundedness step missing from its earlier checklist. A full-parent allocation for every residual infinitesimal-u test is still required. The narrow checker verifies (ES2), (ES3), (ES6)–(ES10) algebraically; the inequalities above are the ordered-field argument. No mixed inverse, mixed order, or source/testing rate comparison is used.

Chapter 47

The actual-j C1 opening: centering the whole divergent event

Component `p48da253a150f`
Source: `R13/advances/revision13_c1_small_l/j_opening_closure.tex`
SHA256: `48da253a150f6b5a5385852f6d99ba5b1ba2fa1d506d1c4e44df4e13c80ec599`
Inventory use: JO1–JO21 and all-test join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

47.1 The actual-j C1 opening: centering the whole divergent event

Theorem 47.1. *Use the original physical native C1 source with $d = 1$, $x = k^2 > 0$ infinitesimal, and*

$$\eta,\ \zeta,\ r,\ 1-r \quad \text{positive bounded units}, \qquad j = \eta(1+xh) - 1 + r \quad \text{positive infinitesimal.} \tag{JO1}$$

Then the complete original parent is tensor SOS for every independent projective test and every state. Literal $\beta = 0$ is retained. There is no assumption that $j_0 = \eta + r - 1$ is positive: $j_0 = 0$ and a slightly negative j_0 are included. Simultaneous infinitesimal r and j are outside this theorem.

All scalar and Gram identities in this section use the normalized parent $F/4$; multiplication of every square root by 2 restores F.

The divergence $D \asymp 1/j$ must be centered by its full event, including its testing-angle dependence. A normal centered only by its quadratic limit would lose control when $j \ll x$. The following exact frame removes that problem before any residue is taken.

Source bounds and contact-exact coordinates. Write $\theta = 1 - h = x\pi$, where $\pi > 0$ is bounded by (SB1), $B_x = \beta + 1/2 \geq 1/2$, and $y = x\eta\zeta$. Put

$$R_0 = 1 + j - \eta = r + \eta x h, \qquad g(U) = R_0 - \eta h U, \qquad U = u^2,\ V = v^2. \tag{JO2}$$

Thus $g(x) = r$ exactly. The native source formulas, with $r = 1 + j - \eta(1 + xh)$, imply

$$\begin{gathered} C, A = O(1), \quad Dj = \tfrac{1}{2} + O(x), \quad m/x \sim 1/(2\zeta), \quad \alpha/x^2 \sim \zeta(1+\zeta)/2, \\ B_x\theta/x = K(\eta + r)(h + \zeta)/(2P_d), \qquad \theta/\sqrt{\alpha}, \quad B_x\theta^2/\alpha = O(1). \end{gathered} \tag{JO3}$$

In particular $D > 0$, so its square root and inverse are pure source and legitimate. Set

$$\begin{gathered} p = uW, \quad q = vZ, \quad E_t = \eta p - q, \qquad N_d = \sqrt{D}\,(J_t + jp), \\ J_t = uW + uvX - Y - vZ, \qquad N_x = \sqrt{B_x}(X + huZ), \quad n = N_d/\sqrt{D}, \quad \xi = N_x/\sqrt{B_x}. \end{gathered} \tag{JO4}$$

The exact inverse state formulas are polynomial and homogeneous:

$$Y_0 = (1+j)p - q - n, \qquad Y = Y_0 - hUq + uv\xi, \qquad X = -huZ + \xi, \qquad J_t = n - jp. \tag{JO5}$$

All three normals vanish at the actual source contact. At their zero graph, $q = \eta p$ and $Y = g(U)p$, so the original native contact graph is recovered at $U = x$ without changing any source contact jet.

For later use introduce positive reference weights

$$\widetilde{C} = \frac{1+j}{2(\eta - j)}, \quad \widetilde{A} = \frac{\eta}{2R_0}, \quad \widetilde{D} = \frac{1}{2j}.$$

Since $R_0, 1 - R_0, \eta$ are units, exact source numerator division gives

$$C - \widetilde{C} = O(x), \qquad A - \widetilde{A} = O(x), \qquad Dj - \tfrac{1}{2} = O(x). \tag{JO6}$$

The last assertion is deliberately about Dj, not D. The canceled denominators of (JO3),(JO6), at $x = j = 0$, contain only the unit factors $\eta, \zeta, 1 - \eta$ and constants. No comparison of j/x has been made.

The complete bounded hard block. The reference quadratic base has the exact identity

$$\begin{aligned} &\widetilde{C}(p - Y_0)^2 + \widetilde{A}Y_0^2 + \widetilde{D}(p - Y_0 - q)^2 + \tfrac{1}{2}(p - Y_0)^2 - pq \\ &= \widetilde{\gamma}E_t^2 + \widetilde{H}\{(1 - \widetilde{b})E_t - n\}^2, \\ \widetilde{H} &= \frac{\eta(j + R_0)}{2R_0(1 - R_0)j}, \quad \widetilde{b} = \frac{R_0(1 - R_0)}{\eta(j + R_0)}, \quad \widetilde{\gamma} = \frac{1+j}{2\eta(j + R_0)}. \end{aligned} \tag{JO7}$$

No p term survives. Furthermore

$$1 - \widetilde{b} = \frac{j(1+j)}{\eta(j + R_0)}, \qquad \widetilde{H}/D \sim 1. \tag{JO8}$$

On (E_t, N_d) this is a bounded hard Gram with diagonal residue $(1/(2\eta(1 - \eta)), 1)$ and vanishing off-diagonal residue. The actual-base error has coefficients $O(x)$ in (p, E_t, N_d). For its apparently singular part use

$$(D - \widetilde{D})(n - jp)^2, \quad \frac{D - \widetilde{D}}{D} = O(x), \quad \frac{(D - \widetilde{D})j}{\sqrt{D}} = O(x\sqrt{j}), \quad (D - \widetilde{D})j^2 = O(xj). \tag{JO9}$$

These are consequences of (JO3),(JO6) and hold for every relative source rate. All other weights in the base are bounded.

For bounded $t = v/u$ at $u, v > 0$ both infinitesimal, put $Z = 1$. Then $p = (v + E_t)/\eta$ in (JO5). The higher-angle terms have bounded infinitesimal hard coefficients, and the m hard coefficient mt^2 is infinitesimal. Combining beta with its half-core part gives the third hard pivot on N_x. Thus the full hard residue is

$$\operatorname{diag}(1/(2\eta(1 - \eta)), 1, 1). \tag{JO10}$$

The exact higher-angle identity printed below also proves these hard bounds directly; no state normal is suppressed here.

An exact scalar three-circuit with bounded polynomial coefficients. Use $s = \sqrt{\eta\zeta}$ and the same separated roots

$$A_t = uv, \quad B_t = \frac{kv^2}{su}, \quad C_t = xs, \quad G = kv, \quad F_v = V - y.$$

On the zero graph of (JO4), the exact scalar is

$$\mathcal{S} = c_F F_v^2 + f_A(U, V)A_t^2 + f_B B_t^2 + f_C(U)C_t^2 - f_G G^2, \tag{JO11}$$

where, with $g = g(U)$,

$$\begin{aligned}
c_F &= \frac{2mR_0 + C(1-R_0)^2 + (A+\alpha+\frac{1}{2})R_0^2}{\eta^2},\\
q_U &= \alpha + (m+B_x)\theta^2,\\
q_V &= \frac{C(1-R_0)^2 + AR_0^2 + (D+\frac{1}{2})j^2}{\eta^2} - \frac{2m\theta}{\eta} - \frac{(2\alpha+1)R_0}{\eta} - \tfrac{1}{2},\\
f_B &= \frac{m\zeta}{\eta x}, \quad f_C = \frac{\alpha - c_F y^2 + q_U U}{\eta\zeta x^2}, \quad f_G = -\frac{q_V + 2c_F y}{x},\\
f_A &= \frac{mV(g^2-2\eta h)}{\eta^2} - \frac{2m\theta g}{\eta} + C(1+V)\{2h(1-R_0)/\eta + h^2U\}\\
&\quad + \{A(1+V) + (\alpha+\tfrac{1}{2})V\}\{g^2/\eta^2 - 2R_0h/\eta + h^2U\}\\
&\quad - (2\alpha+1)(g/\eta - h) + B_x\theta^2 - \tfrac{1}{2}.
\end{aligned} \tag{JO12}$$

In particular f_A is an explicit bounded polynomial of degree at most two in U and one in V; the divergent D is absent. The only D factor elsewhere is $Dj^2 = O(j)$. These formulas follow by substituting $W = v/(\eta u)$, $X = -hu$, $Y = g(U)v/\eta$ in the entire parent; the new checker differentiates that entire substituted parent as well.

The source bounds and (JO12) give

$$c_F \sim 1/(2\eta^2), \qquad f_A, f_B, f_C \sim \lambda = 1/(2\eta).$$

Exact scalar contact gives $f_G = f_A(x,y) + f_B + f_C(x)$, proving that f_G is bounded and $f_G \sim 3\lambda$ without estimating a singular quotient q_V/x. Every nonconstant coefficient in f_A, f_C is bounded; for q_U/x^2 use (JO3).

For clarity the required finite Taylor step works for these higher polynomial coefficients too. Let $H(u,t) = f_A(u^2, u^2t^2) + f_B + f_C(u^2) - f_G$. It is a finite polynomial in u, t with bounded source arrays and $H(k,s) = 0$. Write $\Delta_u = u - k$, $\Delta_t = t - s$. A finite identity, obtained by monic polynomial division, is

$$H = H_u(k,s)\Delta_u + H_t(k,s)\Delta_t + H_{uu}\Delta_u^2 + H_{ut}\Delta_u\Delta_t + H_{tt}\Delta_t^2, \tag{JO13}$$

where one may take

$$\begin{aligned}
H_{uu} &= \frac{H(u,t) - H(k,t) - (u-k)H_u(k,t)}{(u-k)^2},\\
H_{ut} &= \frac{H_u(k,t) - H_u(k,s)}{t-s},\\
H_{tt} &= \frac{H(k,t) - H(k,s) - (t-s)H_t(k,s)}{(t-s)^2}.
\end{aligned}$$

These displayed divisions mean their finite polynomial quotients, not mixed inverses. The first two arrays are bounded and H_{tt} is infinitesimal: every t-dependent monomial of $H(k,t)$ contains k^2. The coefficient first differences similarly have a bounded u quotient and infinitesimal t quotient. Exact first angular stationarity supplies the two unchanged contact-jet balances of the three-circuit lemma. The chain rule is legitimate here because $g(x) = r$ and the original first state derivatives vanish at contact. Consequently (JO11) has the strict finite three-circuit Gram and the strict F_v pivot, including the complete soft defect.

Every complete first normal. The following bounded polynomials are explicit:

$$\begin{aligned}
P_Y/2 &= m\{V(1+gU)/\eta-\theta U\}-CD_v(1-g)/\eta+AD_uD_vg/\eta\\
&\quad+(D+\tfrac12)j/\eta+(\alpha+\tfrac12)D_u(gV/\eta-1),\\
P_E &= \frac{2mV}{\eta^2}\{g+1+j+g(1+j)U\}-\frac{2m\theta}{\eta}\{1+(1+j)U\}\\
&\quad-\frac{2CD_vj(1-g)}{\eta^2}+\frac{2AD_uD_vg(1+j)}{\eta^2}+\frac{2(D+\frac12)j^2}{\eta^2}\\
&\quad+\frac{2(\alpha+\frac12)D_u(1+j)}{\eta}(gV/\eta-1),\\
Z_X &= B_x\theta+P_Y/2-(D+\tfrac12)j/\eta.
\end{aligned}\tag{JO14}$$

Their only surviving D factors are Dj and Dj^2; D cancels entirely from Z_X. Direct full-parent differentiation gives

$$\begin{aligned}
\mathcal{L}_d &= -vP_Y/\sqrt{D},\\
\mathcal{L}_E &= 2mvt^2/\eta^2+vP_E,\\
\mathcal{L}_x &= \frac{1}{\sqrt{B_x}}\{2u\theta(m+B_x)-2mut^2(1+gU)/\eta+2uVZ_X\}.
\end{aligned}\tag{JO15}$$

No normal is missing. For $P=P_Y,P_E$ use the exact finite division $P=P_{00}+UP_U+VP_V$. Contact implies respectively

$$\begin{aligned}
P_{Y,00}/x &= -P_{Y,U}(x,y)-\eta\zeta P_{Y,V}(y),\\
P_{E,00}/x &= -2(m/x)\eta\zeta/\eta^2-P_{E,U}(x,y)-\eta\zeta P_{E,V}(y).
\end{aligned}\tag{JO16}$$

Thus these coquotients are bounded by finite formulas with no j inverse. Allocate vP by $vP=(P_{00}/k)G+u(P_U+t^2P_V)A_t$; allocate the additional $2mvt^2/\eta^2$ on B_t with coefficient $2mst/(\eta^2k)$. The X line allocates exactly on (C_t,B_t,A_t) with coefficients

$$\frac{2u\theta(m+B_x)}{\sqrt{B_x}xs},\quad -\frac{2ms(1+gU)}{\eta k\sqrt{B_x}},\quad \frac{2vZ_X}{\sqrt{B_x}}.\tag{JO17}$$

Every displayed coefficient is infinitesimal; for the first use $\theta(m+B_x)/x=O(1)$, and for the others use $m/x=O(1)$. The N_d row has the additional bounded factor $1/\sqrt{D}$. Each sum of coefficient functions vanishes at contact. Its finite u difference is bounded and its t difference at $u=k$ is infinitesimal, because every t dependence carries u or k. The two unchanged conductors therefore place every whole normal on the strict circuit roots with infinitesimal coefficients. Joining (JO10) proves the full parent on the bounded-ratio chart.

The reciprocal testing ratio: a complete polynomial remainder. For v/u unbounded, return to homogeneous state coordinates and use

$$E_t,\ N_d,\ N_x,\ P_m=\sqrt{m}\,vW,\ Q_\alpha=\sqrt{\alpha}Z,\ R_t=u^2W,\ S_t=uvW.\tag{JO18}$$

The entire base error from (JO9) has zero Gram residue on this list: its p^2 coefficient is $O(x)$ and $\sqrt{x}\,p=\sqrt{x/m}(u/v)P_m$. Every coefficient is a product of a bounded source factor and a bounded testing factor; u/v is pure testing.

Here is the full higher-angle identity used to control the other terms. Write $T_s=-hUq+uv\xi$, so $Y=Y_0+T_s$, with Y_0 in (JO5). After removing the actual quadratic base and the three entire positive packets mR_m^2, $\alpha D_u(vY-Z)^2$ and $B_xD_v(X+uZ)^2$, the exact remainder is

$$\begin{aligned}
\mathcal{R} &= C\{V(p-Y_0)^2-2D_v(p-Y_0)T_s+D_vT_s^2\}\\
&\quad+A\{(D_uD_v-1)Y_0^2+2D_uD_vY_0T_s+D_uD_vT_s^2\}\\
&\quad+\tfrac12D_uVY^2-UqY+(h-\tfrac12)Uq^2-uvq\xi.
\end{aligned}\tag{JO19}$$

This is a quadratic in the full state, with bounded coefficients C, A, h and no D. It is checked as a whole polynomial. Insert $q = \eta p - E_t$, $n = N_d/\sqrt{D}$, $\xi = N_x/\sqrt{B_x}$. Every term contains U, V or uv. Cross terms with p have infinitesimal coefficients on (JO18), using $Up = uR_t$, $Vp = vS_t$ and $uvp = uS_t$. Hard quadratic terms have zero residue. Terms $U^a V^b p^2$ are allocated on R_t^2 when $a > 0$, and on S_t^2 when $a = 0, b > 0$. Their two only nonzero residues, verified from (JO19), are

$$\kappa_R = \eta/2, \qquad \kappa_S = 1/2. \tag{JO20}$$

Every remaining such coefficient contains another U or V. These identities explicitly exhaust the scalar and every normal coupling of (JO19).

The three retained positive packets have leading roots $P_m, -Q_\alpha, N_x$. Their remaining root coefficients are infinitesimal because m, α are infinitesimal and $\theta/\sqrt{\alpha}$, $B_x\theta^2/\alpha$ are bounded:

$$\sqrt{m}\, R_m = P_m + \sqrt{m}\, uvY - \sqrt{m}(\xi + \theta u Z), \quad \sqrt{B_x D_v}(X + uZ) = \sqrt{D_v}\{N_x + \sqrt{B_x\theta^2/\alpha}\, uQ_\alpha\}.$$

The alpha packet is $\sqrt{D_u}(\sqrt{\alpha}\, vY - Q_\alpha)$. Thus the complete parent has a bounded Gram on (JO18) with strictly positive diagonal residue

$$(1/(2\eta(1-\eta)), 1, 1, 1, 1, \eta/2, 1/2). \tag{JO21}$$

This proves the reciprocal-ratio chart and completes the local proof.

All-tests join and receipt. The inherited small-L, bounded-η remote theorem applies to (JO1), including the exact original source openings and literal beta zero. It covers the complement of both-infinitesimal testing angles. The original exact sign gauges, literal axes, homogeneous state extension and real/imaginary decomposition apply as in IC1 and RO. All coefficients and radicals used above are pure-field quantities; no mixed inverse or comparison of source and testing rates occurs.

The two new checkers record 25 and 4 exact assertions. They verify the original full-parent substitution, every complete first normal, the scalar identity, all newly used source denominator factors, the reference hard cancellation, the whole reciprocal higher-angle identity and both strict soft pivots. The finite polynomial bounds and full-rates circuit allocation are the proof above, rather than an inference from those checks. This admits precisely (JO1) and its exact symmetry images. It does not remove the simultaneous r, j opening or any unbounded-ζ or infinitesimal-η source.

Chapter 48

Small-L C1: a weighted closure through both r openings

Component p3df6e6bdd8f4
Source: R13/advances/revision13_c1_small_l/r_opening_closure.tex
SHA256: 3df6e6bdd8f4d43ff45c57f4e6b67343d30198200f2f2dff1a3e30e115b47d90
Inventory use: RO1–RO14 and all-test join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

48.1 Small-L C1: a weighted closure through both r openings

Theorem 48.1. *Use the original physical native C1 source, normalized by $d = 1$, and retain all original intersections. Suppose that $x = k^2$ is positive infinitesimal and*

$$\eta, \quad \zeta, \quad j_0 = \eta + r - 1 \quad \text{are positive bounded units.} \tag{RO1}$$

Then the complete original parent is tensor SOS at every independent projective test and every state. The original $0 < r < 1$ may approach either endpoint at any positive rate, and the literal $\beta = 0$ face is included. No lower positive real bound on r, $1 - r$, or β is imposed. This extends the admitted IC1 domain; it does not cover an infinitesimal j_0, infinitesimal η, or unbounded ζ.

All scalar and Gram identities in this section use the normalized parent $F/4$; multiplication of every square root by 2 restores F.

The proof reuses the unchanged exact scalar and full-normal identities (IC7)–(IC8), (IC16)–(IC19), and the three-circuit lemma. Its new content is a uniform weighting of the two divergent event openings, and a different X normal that retains bounded and zero beta.

Opening-cleared source arrays. Put $\rho = \eta + r$, $y = x\eta\zeta$, $h = 1 - \theta$, $c = C(1 - r)$ and $a = Ar$. The beta numerator (SB1) gives $\theta = x\pi$ with $\pi > 0$ bounded. We do not require π infinitesimal. The original source formulas give, uniformly for all $0 < r < 1$,

$$\begin{gathered} c = \rho/2 + O(x), \quad a = \eta/2 + O(x), \quad D = 1/(2j_0) + O(x), \quad h = 1 + O(x), \\ m/x \sim \rho/(2\zeta), \qquad \alpha/x^2 \sim \rho\zeta(1+\zeta)/2, \qquad B_x\theta/x = \frac{K\rho(h+\zeta)}{2P_d}, \quad B_x = \beta + \tfrac{1}{2} \geq \tfrac{1}{2}. \end{gathered} \tag{RO2}$$

These are finite canceled numerator identities, followed by residue evaluation. Every canceled denominator at $x = 0$ is a product of positive units from $\eta, \zeta, j_0, j_0 + r$ and ρ, with no factor r, $1 - r$, or π. In particular $D > 0$ on this domain; the original $D = 0$ face has empty intersection here, as already proved in (SB2). Both $\theta/\sqrt{\alpha}$ and $B_x\theta^2/\alpha$ are bounded, not necessarily infinitesimal.

The exact source formulas for the two cleared strong weights are

$$c = \frac{K\rho E}{2P_d}, \qquad a = \frac{\eta P_4}{2P_d}.$$

Their order-x differences in (RO2), and the following arrays, have opening-free finite canceled formulas:

$$\frac{P_{Y,00}}{x}, \quad \frac{P_{E,00}}{x}, \quad \frac{\theta(m+B_x)}{x}, \quad \frac{\alpha+(m+B_x)\theta^2}{x^2}, \quad \frac{q_V}{x}. \tag{RO3}$$

Here P_Y, P_E, q_V are precisely (IC16) and (IC8). To check why no unbounded event remains in these arrays, replace every occurrence of $C(1-r)^2$ by $c(1-r)$, every Ar^2 by ar, every $C(1-r)$ by c, and every Ar by a before division. No other occurrence of C or A survives in the scalar or first-normal polynomials. The accompanying new checker prints the denominators of (RO2)–(RO3), retains arbitrary bounded π, and checks both $r=0$ and $r=1$ after cancellation. This is a finite symbolic criterion for boundedness on the stated field domain, not a numerical endpoint approximation.

The normalized hard frame. Define the pure-source coefficients

$$b=\frac{r(1-r)}{\eta(j_0+r)}, \quad H_0=\frac{\eta(j_0+r)}{2r(1-r)j_0}, \quad \gamma=\frac{\rho}{2\eta(j_0+r)}.$$

Both $1/\sqrt{H_0}$ and $1/\sqrt{B_x}$ are bounded; H_0 itself may be unbounded. Put

$$E_t=\eta uW-vZ, \quad N_y=\sqrt{H_0}(Y-ruW-bE_t), \quad N_x=\sqrt{B_x}(X+huZ). \tag{RO4}$$

All three normals vanish at the actual source contact. For the comparison weights

$$C_0=\frac{\rho}{2(1-r)}, \quad A_0=\frac{\eta}{2r}, \quad D_0=\frac{1}{2j_0},$$

the exact quadratic base is

$$C_0(p-Y)^2+A_0Y^2+D_0(p-Y-q)^2+\tfrac{1}{2}(p-Y)^2-pq=H_0(Y-rp-b(\eta p-q))^2+\gamma(\eta p-q)^2. \tag{RO5}$$

Thus the divergent normal has already been normalized before taking residue. The cancellation factors needed for its entire hard block are

$$\frac{C_0}{H_0}=\frac{r\rho j_0}{\eta(j_0+r)}, \quad \frac{A_0}{H_0}=\frac{(1-r)j_0}{j_0+r}, \quad \frac{D_0}{H_0}=\frac{r(1-r)}{\eta(j_0+r)}. \tag{RO6}$$

For example, after $p=(v+E_t)/\eta$ and $Z=1$, the E_t coefficients in $p-Y$ and Y are respectively

$$\frac{1-r}{\eta}-b=\frac{(1-r)j_0}{\eta(j_0+r)}, \qquad \frac{r}{\eta}+b=\frac{r\rho}{\eta(j_0+r)}. \tag{RO7}$$

The strong C, A factors therefore cancel even on those complete first-state directions. For instance an error $\delta_c(p-Y)^2/(1-r)$, with $\delta_c=O(x)$, has a N_y^2 coefficient $\delta_c/((1-r)H_0)=O(x)$, an E_t^2 coefficient $O(x)(1-r)$, and an E_tN_y coefficient $O(x)/\sqrt{H_0}$. The A error has the same properties using rH_0 and the second formula in (RO7). The original higher-angle factors have the same cleared coefficients, multiplied by $U=u^2$ or $V=v^2$.

Suppose first that $u, v>0$ are infinitesimal and the pure-testing ratio $t=v/u$ is bounded. After the state substitution (RO4), the complete hard quadratic on (E_t, N_y, N_x) consequently has bounded entries and residue

$$\operatorname{diag}(\gamma, 1, 1). \tag{RO8}$$

Here γ is a positive bounded unit even when either r opening is infinitesimal. The additional mt^2 contribution is infinitesimal. The beta and half-core coefficients jointly give the third pivot because m/B_x is infinitesimal. These estimates concern the full hard block, not only its restriction to the contact graph.

The complete scalar and all first normals. Normalize $Z = 1$, homogenizing at the end as in IC1. Use the unchanged positive separated roots

$$A_t = uv, \quad B_t = \frac{kv^2}{su}, \quad C_t = xs, \quad G = kv, \quad F_v = V - y, \qquad s = \sqrt{\eta\zeta}.$$

The scalar at all three normals zero is exactly (IC7). Its coefficient residues are still

$$c_F \sim \frac{\rho}{2\eta^2}, \quad f_A, f_B, f_C \sim \lambda = \frac{\rho}{2\eta}, \quad f_G \sim 3\lambda. \tag{RO9}$$

They do not depend on the residue of π. Every coefficient of U, V in (IC8) is bounded after the clearing described above; for the additional term in f_C use $B_x\theta^2/x^2 = (B_x\theta/x)\pi$. The exact contact value and the two first-jet identities (IC11)–(IC14) are unchanged. Their finite arrays therefore obey precisely the hypotheses of the three-circuit lemma, with s a unit. It gives the same strict finite circuit Gram and the strict F_v pivot. In particular the negative soft defect (SN10) has been allocated by the entire scalar (IC7), not dropped by comparing only its quadratic base.

The complete first normals are (IC18), with the N line divided by $\sqrt{H_0}$ and the X line divided by $\sqrt{B_x}$ instead of $\sqrt{\beta}$. Their exact root allocations are (IC19) with those same replacements. Every coefficient remains infinitesimal: the N_y factors are bounded source factors times the old cleared polynomials; the X coefficient on C_t is

$$\frac{2u\theta(m + B_x)}{\sqrt{B_x}\, xs},$$

which is infinitesimal by (RO3), and its coefficients on B_t, A_t are respectively $O(k)$ and $O(v)$. In the latter coefficient $Z_X = B_x\theta - (D + \frac{1}{2})(1 + hU) + (D + \frac{1}{2})(1 - r)/\eta$ is bounded. The E_t line uses $m/x, P_{E,00}/x$ and bounded polynomial arrays, as in IC1. There is no negative power of t in any coefficient.

Each coefficient is polynomial in u, t with bounded pure-source arrays; every term depending on t carries u, k or $m/k = O(k)$. The exact monic differences (IC21) hence have bounded u quotient and infinitesimal t quotient after evaluation at $u = k$. The unchanged exact source contact cancels the bare G term in each complete normal. The conductors $G(u-k) = k(A_t - G)$ and $G(t-s) = s(B_t - G)$ therefore allocate all normals with infinitesimal coefficients on the strict circuit roots. Joining this allocation to (RO8) proves the entire parent on the bounded-ratio chart.

The reciprocal ratio chart. For v/u unbounded use the homogeneous root list

$$E_t, \quad N_y, \quad N_x, \quad P_m = \sqrt{m}\, vW, \quad Q_\alpha = \sqrt{\alpha}Z, \quad R_t = u^2W, \quad S_t = uvW. \tag{RO10}$$

The error of the actual quadratic base relative to (RO5) is an $O(x)$ quadratic in (p, E_t, N_y) after the opening cancellations (RO6)–(RO7). Every $\sqrt{x}\, p$ factor is allocated exactly by

$$\sqrt{x}\, p = \sqrt{x/m}\, (u/v) P_m,$$

with a bounded source unit x/m and an infinitesimal pure-testing ratio u/v. Thus the entire error has zero Gram residue on (RO10). All coefficients in the exact higher-angle identity (IC24) remain bounded after $Y = rp + bE_t + N_y/\sqrt{H_0}$. This follows term by term from (RO6), $b = O(r(1-r))$, $Ar = a$, and $C(1-r) = c$. Its only nonzero scalar residues are the two positive diagonal entries

$$\kappa_R = \rho\eta/2, \qquad \kappa_S = \rho/2. \tag{RO11}$$

No opening has been substituted before cancellation.

It remains essential to combine beta with its half-core part. Set $B_t^{\text{state}} = X + uZ$ and $J_0 = p - Y - D_u vZ$. Relative to $B_t^{\text{state}} = 0$, the exact D, beta and half-core increment is

$$2(D + \tfrac{1}{2})uvB_t^{\text{state}} J_0 + (D + \tfrac{1}{2})UV(B_t^{\text{state}})^2 + B_x D_v (B_t^{\text{state}})^2. \tag{RO12}$$

This whole-polynomial identity is checked in the new checker. Use

$$B_t^{\mathrm{state}} = \frac{N_x}{\sqrt{B_x}} + \frac{\theta u}{\sqrt{\alpha}} Q_\alpha .$$

The first two terms of (RO12) have zero Gram residue on (RO10); D is bounded, $1/\sqrt{B_x}$ and $\theta/\sqrt{\alpha}$ are bounded, and every coefficient carries u or v. The three remaining positive packets have their entire roots

$$\begin{aligned} \sqrt{m}\, R_m &= P_m + \sqrt{m}\, uvY - \sqrt{m}\, B_t^{\mathrm{state}}, \\ \sqrt{\alpha}\sqrt{D_u}(vY - Z) &= \sqrt{D_u}\{\sqrt{\alpha}[rS_t + v(bE_t + N_y/\sqrt{H_0})] - Q_\alpha\}, \\ \sqrt{B_x}\sqrt{D_v} B_t^{\mathrm{state}} &= \sqrt{D_v}\{N_x + \sqrt{B_x\theta^2/\alpha}\, uQ_\alpha\}. \end{aligned} \tag{RO13}$$

Each is its leading root plus an infinitesimal-coefficient combination of (RO10). The square root in the last line is pure source and bounded; its factor u supplies the needed infinitesimal even when beta is zero or bounded. The full parent therefore has a bounded Gram with strict diagonal residue

$$(\gamma, 1, 1, 1, 1, \kappa_R, \kappa_S). \tag{RO14}$$

This proves the remaining ratio chart.

All-tests join and exact scope. The earlier small-L, bounded-η remote theorem includes (RO1) with every original opening, so it covers the complement of u, v both infinitesimal. Literal axes have the elementary full-core identities recorded in IC1. Its exact sign gauges apply unchanged. Homogenizing the $Z = 1$ identity gives polynomial state identities, including $Z = 0$; applying them to real and imaginary parts covers complex states. All inverse and radical factors above are pure source or pure testing. This completes the claimed full-parent all-tests admission.

The newly removed source region is the portion of (RO1) outside IC1, namely either r or $1 - r$ infinitesimal, with all total-scale and earlier-complement intersections retained. Remaining C14.3 source configurations still include infinitesimal η, unbounded ζ, and nonunit or nonpositive j_0 while the actual $j = \eta(1 + xh) - 1 + r$ remains positive. No conclusion on those regions or on C14.4 follows from this theorem.

A local algebraic extension below beta zero. The preceding two local ratio-chart arguments have a stronger precise scope that is useful when another event is restored later. Assume the exact native stationary coefficient formulas (NC1)–(NC4), $x > 0$ infinitesimal, (RO1), $0 < r, h < 1$, and

$$B_x = \beta + \tfrac{1}{2} \geq b_* > 0 \quad \text{for some real } b_*. \tag{RO15}$$

Original source physicality and $\beta \geq 0$ are not required for this local assertion. Then the full C1-form parent is tensor SOS at all independent u, v both infinitesimal, including their literal zero axes. In particular $-1/2 < \beta < 0$ is allowed whenever $\beta + 1/2$ is a positive bounded unit.

Indeed $P_d \sim 1$ before any h estimate, since all its source arguments are bounded. The exact formula $B_x\theta/x = K\rho(h + \zeta)/(2P_d)$ has bounded right side, and (RO15) therefore implies θ/x bounded. All of (RO2)–(RO14) follows, with $1/\sqrt{B_x} \leq 1/\sqrt{b_*}$. In particular $m, \alpha > 0$, $C, A > 0$, and $D \sim 1/(2j_0) > 0$ on this domain. No remaining use of $\beta \geq 0$ occurs in the local scalar, first-normal or hard-block proof: its complete contribution is combined with the half-core as (RO12). At $u = 0$ the signed core contributes $D_v(X^2 + Y^2)/2$, so its X^2 part plus the beta packet is $B_x D_v X^2$. At $v = 0$ the same combination is $B_x(X + uZ)^2$ and the remaining half-core is $(uW - Y)^2/2$. Thus both axes are also exact SOS under (RO15). Only this local assertion is extended. The earlier all-tests remote admission is not imported for negative beta.

Chapter 49

The simultaneous r, j opening: a harmonic full-event frame

Component `pbf6de1d7328a`
Source: `R13/advances/revision13_c1_small_l/simultaneous_r_j_closure.tex`
SHA256: `bf6de1d7328a967852a61f39ed8fdc610c3ce8dd87434cac879ae2ee0f0fd2e8`
Inventory use: HO1–HO24 and all-test join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

49.1 The simultaneous r, j opening: a harmonic full-event frame

Theorem 49.1. *Use the original physical native C1 source, with $d = 1$ and $x = k^2 > 0$ infinitesimal. Suppose η, ζ are positive bounded units and r and the actual $j = \eta(1 + xh) - 1 + r$ are both positive infinitesimals. Then the complete original parent is tensor SOS for every independent projective test and every state. Literal $\beta = 0$ is retained. No relative rate among x, r, j or between source and testing parameters is imposed.*

All scalar and Gram identities in this section use the normalized parent $F/4$; multiplication of every square root by 2 restores F.

The essential change is to combine the whole positive A, D events before choosing the scalar graph. The weighted uvX term is retained inside a hard root. Solving for $p = uW$, rather than for $q = vZ$, then gives a polynomial inverse with a pure-source unit denominator.

Source bounds and the harmonic frame. Write $U = u^2$, $V = v^2$, $D_u = 1 + U$, $D_v = 1 + V$, $\theta = 1 - h = x\pi$, $B_x = \beta + 1/2$, and

$$T = 1 + xh, \qquad e = \eta T = 1 - r + j, \qquad y = x\eta\zeta. \tag{HO1}$$

The bounded-η source bound gives $\pi = O(1)$. Here $e, \eta, h, 1 - r \sim 1$. Exact native numerator cancellation gives

$$\begin{gathered} C - C_* = O(x), \qquad Ar - e/2 = O(x), \qquad Dj - 1/2 = O(x), \\ C_* = \frac{1+j}{2(1-r)}, \qquad A_* = \frac{e}{2r}, \qquad D_* = \frac{1}{2j}, \\ m/x \sim 1/(2\zeta), \qquad \alpha/x^2 \sim \zeta(1+\zeta)/2, \\ B_x\theta/x = O(1), \qquad q_U/x^2 = O(1), \quad q_U = \alpha + (m + B_x)\theta^2. \end{gathered} \tag{HO2}$$

The checker cancels the exact rational formulas before setting $x = r = j = 0$: every denominator in the seven displayed bounded coquotients factors there into constants and powers of the retained unit ζ. This proves boundedness at arbitrary simultaneous source rates, rather than just nonvanishing at

a selected boundary. In particular $A, D > 0$ and both are unbounded. Also $\theta/\sqrt{\alpha}$ and $B_x\theta^2/\alpha$ are bounded, including $\beta = 0$.

Define pure-source positive quantities

$$\mathsf{H} = A + D, \qquad \rho = \frac{D}{\mathsf{H}}, \qquad \mathsf{K} = \frac{AD}{\mathsf{H}}, \qquad \delta = \frac{Ar + Dj}{\mathsf{H}}. \tag{HO3}$$

Both H and K are unbounded, $0 < \rho < 1$, and $\mathsf{K}^{-1} = A^{-1} + D^{-1} = O(r + j)$. Put $p = uW$, $q = vZ$, $J_0 = p + uvX - q$, and use the full-event roots

$$\begin{aligned} E &= \sqrt{\mathsf{K}}\{J_0 - (r - j)p\}, \\ N &= \sqrt{\mathsf{H}}\{Y - \rho J_0 - \delta p\}, \\ N_x &= \sqrt{B_x}(X + huZ), \qquad \ell = E/\sqrt{\mathsf{K}}, \quad n = N/\sqrt{\mathsf{H}}, \quad \xi = N_x/\sqrt{B_x}. \end{aligned} \tag{HO4}$$

All three vanish at the actual source contact. Their exact inverse is

$$\begin{aligned} p &= \frac{(1 + hU)q - uv\xi + \ell}{e}, \quad Y = rp + \rho\ell + n, \qquad X = -huZ + \xi, \\ J_t &= -jp + (1 - \rho)\ell - n, \qquad J_t = p + uvX - Y - q. \end{aligned} \tag{HO5}$$

The denominator e is pure source. In particular no inverse of $1 + hU$ occurs, even if the testing and source rates are unrelated. The useful exact weighted identities are

$$\frac{A\rho^2}{\mathsf{K}} = \rho, \qquad \frac{A}{\mathsf{H}} = 1 - \rho, \qquad \frac{D(1 - \rho)^2}{\mathsf{K}} = 1 - \rho, \qquad \frac{D}{\mathsf{H}} = \rho. \tag{HO6}$$

The complete hard block and its reference cancellation. The A, D base has the exact identity

$$\begin{aligned} AY^2 + DJ_t^2 = N^2 + E^2 + 2p\left\{\frac{Ar + Dj}{\sqrt{\mathsf{H}}}N + \sqrt{\mathsf{K}}(r - j)E\right\} \\ + (Ar^2 + Dj^2)p^2. \end{aligned} \tag{HO7}$$

Both displayed cross coefficients and the last coefficient are infinitesimal: the squared E cross coefficient is $O(r + j)$ and the squared N cross coefficient is $O(\mathsf{H}^{-1})$. For bounded testing ratio $t = v/u$, (HO5) has bounded coefficients on the hard roots, and the only potentially large m coefficient is mt^2/K, which is infinitesimal. The higher A packet is controlled by (HO6); for example

$$\frac{Ar^2}{\mathsf{K}} = r^2 + \frac{Ar}{Dj}rj = O((r + j)^2), \qquad \frac{Ar\rho}{\mathsf{K}} = r. \tag{HO8}$$

Consequently its hard coefficients carry an additional U or V. The bounded C and half-core packets have infinitesimal coefficients on E, N, and combining the beta packet with its half-core part gives the third hard pivot. The full hard residue on (E, N, N_x) is $\mathrm{diag}(1, 1, 1)$.

A stronger exact base statement is needed for unbounded t. Set $\widetilde{q} = q - uvX = ep - \ell$, so that $J_t = p - Y - \widetilde{q}$. The reference quadratic form

$$\mathcal{B}_* = C_*(p - Y)^2 + A_*Y^2 + D_*(p - Y - \widetilde{q})^2 + \tfrac{1}{2}(p - Y)^2 - p\widetilde{q} \tag{HO9}$$

has no soft p coordinate after substituting $Y = rp + \rho\ell + n$ and $\widetilde{q} = ep - \ell$:

$$\mathcal{B}_* = (A_* + C_* + \tfrac{1}{2})(\rho\ell + n)^2 + D_*\{(1 - \rho)\ell - n\}^2. \tag{HO10}$$

This is an exact contact cancellation, checked before taking residues. Since $A_*/A = 1 + O(x)$ and $D_*/D = 1 + O(x)$, (HO6) shows that its bounded hard Gram on (E, N) has residue $\mathrm{diag}(1, 1)$. Replacing the reference weights by the actual C, A, D creates a quadratic error with coefficients $O(x)$ in (p, E, N). For the singular weights this follows from $(A - A_*)Y^2 = O(x)\,AY^2$ and $(D - D_*)J_t^2 = O(x)\,DJ_t^2$, using (HO6),(HO7); neither an x/r nor an x/j estimate is used.

The exact bounded-ratio scalar circuit. At $E = N = N_x = 0$ and $Z = 1$, let

$$w(U) = \frac{1 + hU}{e}, \qquad p = w(U)v, \quad Y = rw(U)v, \quad X = -hu. \tag{HO11}$$

Since $w(x) = 1/\eta$, this graph agrees exactly with the native source contact. The source contact value and all first state and angular jets are therefore inherited by the chain rule.

For compact scalar coefficients set $a = C(1-r)^2$, $b = Ar^2$, $d = (D+1/2)j^2$, $c = (\alpha+1/2)r^2$, and

$$L_A(U) = 1 + 2h + (2h + h^2)U + h^2U^2.$$

The complete scalar is exactly

$$\mathcal{S} = c_F(V - y)^2 + f_A(U, V)A_t^2 + f_B B_t^2 + f_C(U)C_t^2 - f_G G^2, \tag{HO12}$$

on the separated roots $A_t = uv$, $B_t = kv^2/(su)$, $C_t = xs$, $G = kv$, where $s = \sqrt{\eta\zeta}$. The coefficients are

$$\begin{aligned}
c_F &= \frac{2m(h + r) + a + b + c}{e^2},\\
q_V &= \frac{a + b + d}{e^2} - \frac{2m\theta}{e} - \frac{(2\alpha + 1)r}{e} - \tfrac{1}{2},\\
f_B &= \frac{m\eta\zeta}{e^2 x}, \qquad f_C = \frac{\alpha - c_F y^2 + q_U U}{\eta\zeta x^2}, \qquad f_G = -\frac{q_V + 2c_F y}{x},\\
f_A &= \frac{mV}{e^2}\{(h + r)^2 + 2hr + 2hr(h + r)U + h^2r^2U^2\} - \frac{2m\theta}{e}(h + r + hrU)\\
&\quad + \frac{a}{e^2}(2h + h^2U)(1 + V) + \frac{b}{e^2}L_A(U)(1 + V) + \frac{d}{e^2}(2h + h^2U) + \frac{c}{e^2}L_A(U)V\\
&\quad - \frac{(2\alpha + 1)r}{e}(1 + h + hU) + B_x\theta^2 - \tfrac{1}{2}.
\end{aligned} \tag{HO13}$$

The only opening weights here are Ar^2 and Dj^2, both infinitesimal. Thus f_A is a bounded polynomial of degree at most two in U and one in V, and all nonconstant coefficients of f_C are bounded by (HO2). Direct source reduction gives

$$c_F \sim \tfrac{1}{2}, \qquad f_A, f_B, f_C \sim \tfrac{1}{2}. \tag{HO14}$$

Exact scalar contact gives $f_G = f_A(x, y) + f_B + f_C(x) \sim 3/2$, without separately estimating q_V/x.

Apply the finite three-circuit lemma with $\lambda = 1/2$. The polynomial Taylor arrays are the monic quotients in (JO13), now applied to $f_A(u^2, u^2t^2) + f_B + f_C(u^2) - f_G$. Their u and mixed arrays are bounded; the t, t array and the coefficient t first differences are infinitesimal, since every t dependence at $u = k$ contains k^2. The two exact angular contact jets give the two circuit balances. This supplies the strict three-circuit Gram and the strict $V - y$ pivot for the entire scalar, including its negative soft term.

All complete first normals. The following formulas come from differentiating the entire original parent after (HO5), with $Z = 1$. Put $a_E = r/e + \rho$, $b_E = (1 - r)/e - \rho$, $c_E = 1 - \rho - j/e$. The bounded polynomials are

$$\begin{aligned}
P_N/2 &= m\{wV(1 + rU) - \theta U\} - CD_v(1 - r)w + AD_uD_vrw\\
&\quad + (D + \tfrac{1}{2})jw + (\alpha + \tfrac{1}{2})D_u(rwV - 1),\\
P_E/2 &= \frac{mV}{e^2}(h + r + hrU) - \frac{m\theta}{e} + ma_E\{wV(1 + rU) - \theta U\}\\
&\quad + CD_v(1 - r)wb_E + AD_uD_vrwa_E - (D + \tfrac{1}{2})jwc_E + (\alpha + \tfrac{1}{2})D_u(rwV - 1)a_E,\\
Z_X &= \frac{m\theta(1 + rU)}{e} + B_x\theta\\
&\quad - \frac{CD_v(1 - r)^2w + AD_uD_vr^2w + (D + \tfrac{1}{2})j^2w + (\alpha + \tfrac{1}{2})D_ur(rwV - 1)}{e}.
\end{aligned} \tag{HO15}$$

Their only opening-weight factors are Ar, Dj, Ar^2, Dj^2. Every coefficient is bounded uniformly in the source rates and $0 < \rho < 1$. The three complete first normals are

$$\begin{aligned}\mathcal{L}_N &= \frac{vP_N}{\sqrt{\mathsf{H}}},\\ \mathcal{L}_E &= \frac{2mvt^2/e^2 + vP_E}{\sqrt{\mathsf{K}}},\\ \mathcal{L}_x &= \frac{1}{\sqrt{B_x}}\left\{2u\theta(m+B_x) - 2mwut^2(1+rU)\left(1+\frac{V(1+rU)}{e}\right) + 2uVZ_X\right\}.\end{aligned} \tag{HO16}$$

For either $P = P_N, P_E$, use its exact finite monic decomposition $P = P_{00} + UP_U + VP_V$. State contact gives

$$\begin{aligned}P_{N,00}/x &= -P_{N,U}(x,y) - \eta\zeta P_{N,V}(y),\\ P_{E,00}/x &= -2(m/x)\eta\zeta/e^2 - P_{E,U}(x,y) - \eta\zeta P_{E,V}(y).\end{aligned} \tag{HO17}$$

Thus these coquotients are bounded without any new opening inverse. Allocate $vP = (P_{00}/k)G + u(P_U + t^2P_V)A_t$. The extra E term allocates on B_t with coefficient $2mst/(e^2k\sqrt{\mathsf{K}})$. The three X terms allocate respectively on C_t, B_t, A_t, with coefficients

$$\frac{2u\theta(m+B_x)}{xs\sqrt{B_x}}, \qquad \frac{-2msw(1+rU)\{1+V(1+rU)/e\}}{k\sqrt{B_x}}, \qquad \frac{2vZ_X}{\sqrt{B_x}}. \tag{HO18}$$

Every displayed allocation coefficient is infinitesimal. The coefficient sums vanish at contact by the exact state jets; their finite u first differences are bounded and their t first differences at $u = k$ are infinitesimal. This follows directly from the bounded polynomial arrays in (HO15): every t dependence carries u or k. The two unchanged circuit conductors therefore place every whole normal on the strict circuit roots with infinitesimal coefficients. Together with the complete hard block, this proves all bounded t, including infinitesimal t, without a missing state normal.

The reciprocal ratio: retain both soft directions. If v/u is unbounded, work homogeneously and use seven roots

$$E, \quad N, \quad N_x, \quad P_m = \sqrt{m}\,vW, \quad Q_\alpha = \sqrt{\alpha}Z, \quad R_t = uvZ, \quad S_t = uvW. \tag{HO19}$$

The base error following (HO10) has zero Gram residue on this list: use the exact separated identity $\sqrt{x}\,p = \sqrt{x/m}(u/v)P_m$. This absorbs every $O(x)$ coefficient in (p, E, N), since x/m is a source unit and u/v is a pure-testing infinitesimal.

After removing the actual quadratic base (HO9), with actual C, A, D, and the three entire positive packets mM_t^2, $\alpha D_u(vY - Z)^2$, $B_xD_v(X + uZ)^2$, the full remaining quadratic is exactly

$$\begin{aligned}\mathcal{R} = {} & CV(p-Y)^2 + A(U+V+UV)Y^2 + \tfrac{1}{2}D_uVY^2 - \theta UqY\\ & + (h-\tfrac{1}{2})Uq^2 + \tfrac{1}{2}h^2U^2q^2 - uv\xi\{Y + (1+hU)q\} + \tfrac{1}{2}UV\xi^2.\end{aligned} \tag{HO20}$$

This identity retains the whole D event in the quadratic base, so no coefficient containing a bare divergent D remains. It is checked as one complete homogeneous polynomial.

Here is a finite allocation of every term. The exact identity

$$up = \frac{(1+hU)R_t - u^2v\xi + uE/\sqrt{\mathsf{K}}}{e} \tag{HO21}$$

has bounded coefficients and uses no mixed inverse. The C packet has leading root $\sqrt{C}(1-r)S_t$; its other coefficients are $-\sqrt{C}\,\rho v/\sqrt{\mathsf{K}}$ and $-\sqrt{C}\,v/\sqrt{\mathsf{H}}$ on E, N, both infinitesimal. All three A higher packets have zero residue. For example,

$$\sqrt{A}\,uY = \sqrt{A}\,r(up) + \sqrt{\rho}\,uE + \sqrt{1-\rho}\,uN, \qquad \sqrt{A}\,vY = \sqrt{A}\,rS_t + \sqrt{\rho}\,vE + \sqrt{1-\rho}\,vN. \tag{HO22}$$

Here $\sqrt{A}\,r = O(\sqrt{r})$ and (HO8),(HO21) control every coefficient; the uvY packet is v times the first. The half-VY^2 term is smaller by the same identities. The $-\theta UqY$ term is $-\theta R_t(uY)$ and has zero residue. The next two terms are $(h-1/2)R_t^2$ and $h^2UR_t^2/2$. Finally, the cross term is $-\xi v(uY) - \xi v(1+hU)R_t$ and all its coefficients, as well as those of $UV\xi^2/2$, are infinitesimal. Thus this allocation exhausts the complete quadratic (HO20); its only nonzero soft residues are $1/2$ on each of R_t, S_t.

The three retained positive packets have leading roots $P_m, -Q_\alpha, N_x$:

$$\begin{aligned}\sqrt{m}\,M_t &= P_m + \sqrt{m}\,uvY - \sqrt{m}\,\xi - \sqrt{m/\alpha}\,\theta uQ_\alpha,\\ \sqrt{\alpha D_u}(vY - Z) &= \sqrt{D_u}(\sqrt{\alpha}\,vY - Q_\alpha),\\ \sqrt{B_x D_v}(X + uZ) &= \sqrt{D_v}\{N_x + \sqrt{B_x\theta^2/\alpha}\,uQ_\alpha\}.\end{aligned} \tag{HO23}$$

Every other coefficient here is infinitesimal by (HO2),(HO21), because $\theta\sqrt{m/\alpha} = O(\sqrt{x})$. The complete bounded Gram on (HO19) therefore has strictly positive diagonal residue

$$\operatorname{diag}(1,1,1,1,1,1/2,1/2). \tag{HO24}$$

This proves the reciprocal chart for the full parent and every state.

All-tests join and receipt. The inherited small-L, bounded-η remote theorem covers the complement of both-infinitesimal testing angles on this original physical source domain. Its exact source openings and literal beta zero remain available. The original exact sign gauges, literal axes, homogeneous state extension and real/imaginary decomposition join as in IC1, RO and JO. All roots, radicals and inverses above are pure-field or separated; every bound is a product of a source bound and a testing bound.

The separate checker records 28 exact assertions, including the full-parent scalar substitution, all three complete first normals, the two full-event maps, the harmonic completion, the exact reference kernel cancellation, the whole reciprocal remainder, its two strict soft pivots and the retained-unit denominator factorizations. Those checks support the finite bounds and circuit allocation proved above; they are not being used as a substitute for them. This theorem removes precisely the simultaneous positive infinitesimal r, j opening with unit η, ζ.

Corollary 49.2 (The entire unit-η,ζ small-L region)**.** *Within the original physical native small-L C1.4 source, suppose $x > 0$ is infinitesimal and η, ζ are positive bounded units. Then the full parent is tensor SOS at every independent projective test and every state, with all original source intersections retained.*

Proof. The positive actual gap $j = \eta(1+xh) - 1 + r$ is bounded. If j is a unit, then $j_0 = j - \eta xh$ is a positive unit, so RO applies, including either r opening. If j is infinitesimal, then $1 - r \sim \eta$ is a unit. A unit r is exactly the JO domain; an infinitesimal r is exactly the HO domain. These alternatives exhaust the stated source, require no comparison of j with x, and each supplies its complete local, remote and endpoint join. □

Chapter 50

The shared C1–C2 boundary and a positive source partition

Component `pc024f5300388`
Source: `R13/advances/revision13_source_overlap/source_overlap.tex`
SHA256: `c024f5300388b76ae18ed766c6b3dd4c9edc8b5ff3c2706e6d51805310e44fb8`
Inventory use: OP1-OP5

50.1 The shared C1–C2 boundary and a positive source partition

This section records an exact source-only identity. It does not by itself admit any new independent testing domain. Its purpose is to identify the same physical boundary in the two remaining source charts and to retain the nonnegative slack when that boundary opens.

Use the original weak-C2 variables $B = H + L$, $z, y > 0$ and $0 < r < 1$, with the unchanged source gaps and nonnegative n. Write $e = 1 - r$, $K = 1 + z$, $T_c = B + zr$, and let δ be the original positive C2 denominator. Define the positive source polynomial

$$P_n = B + B^2 + 2Bzr + BHz + yB(1 + z) + Hrz^2 + yz^2re. \tag{OP1}$$

Then the following identity holds on the entire original source:

$$\boxed{eP_n = (1 + z)(H + y)(B + zr) + 2r\delta n.} \tag{OP2}$$

Every factor in the first summand is positive, and the second summand is nonnegative, including literal $n = 0$. In particular the pure-source weights

$$q = \frac{(1 + z)(H + y)(B + zr)}{eP_n} > 0, \qquad s_n = \frac{2r\delta n}{eP_n} \geq 0 \quad \text{satisfy} \quad q + s_n = 1. \tag{OP3}$$

Both square roots are legitimate source-field elements. Neither weight is assumed to have nonzero residue.

Proof. The unchanged formula for n is

$$2r\delta n = (B + zr)(1 + zr)(1 + zr + B - H) - r\delta.$$

Substitution of the printed polynomial for δ, followed by collecting terms, gives OP2. This is a polynomial identity, so no source boundary is obtained by continuity and no opening has been inverted except the displayed positive pure-source factors. □

Exact correspondence of the source contacts. Use a subscript 1 for native C1 notation in this paragraph. The map

$$x_1 = z, \qquad \eta_1 = \frac{y}{B}, \qquad \zeta_1 = \frac{B}{z}, \qquad r_1 = \frac{H}{B}, \qquad \theta_1 = e, \tag{OP4}$$

with $k_1 = \sqrt{z}$ and the native parameter $\Delta_1 = B/(z\sqrt{y})$, gives

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(\sqrt{z}, \sqrt{y}; \frac{B}{\sqrt{zy}}, -r\sqrt{z}, \frac{H}{\sqrt{y}}, 1\right)$$

from the original native C1 contact formula. Thus it is the same contact, including all state coordinates. Direct substitution in the native positive polynomial P_d gives $BP_d = P_n$. The native C1 beta coefficient at this formal contact therefore obeys

$$\beta_1 = -\frac{r\delta n}{eP_n} = -\frac{s_n}{2}, \qquad \beta_1 + \tfrac{1}{2} = \frac{q}{2} > 0. \tag{OP5}$$

For $n = 0$, the original C2 parent has $\lambda = n = \beta = 0$. It is literally an original C1 parent, up to an optional positive normalization of C. The native source entry and uniqueness at stationary contact identify its coefficients with OP4. For $n > 0$, OP5 makes the formal C1 beta negative. Hence OP4 does *not* map that source to the nonnegative C1 event cone, and no physical C1 all-tests theorem may be imported there.

The revision-12 small-gap chart. Put $z = aB$, $y = tB$, $H = hB$, and $B = e\ell$, with the exact BS1/GR1 hypotheses. Here a, t are positive bounded units, h, e are infinitesimal, $t_0 > 1$, and ℓ is positive bounded. Canceling the pure-source factor e before taking residues gives

$$q_0 = \ell_0 t_0(1 + a_0), \qquad (s_n)_0 = 1 - \ell_0 t_0(1 + a_0). \tag{OP6}$$

This is consistent with the original constraint $n \geq 0$ and keeps its equality case. At $n = 0$, OP4 has

$$L_1 = x_1\zeta_1 = B, \quad \eta_1 = t, \quad \zeta_1 = 1/a, \quad r_1 = h, \quad (j_{0,1})_0 = t_0 - 1 > 0.$$

It therefore enters the weighted small-L C1 theorem in Section 48.1. This overlap is an exact reuse of a proved physical boundary, not a claim that its positive-n neighborhood is covered. All total-scale and earlier-complement intersections remain in force.

Chapter 51

A centered angular packet closes the physical small-gap C2 chart

Component p7c5e0b890910
Source: R13/advances/revision13_weak_local/centered_circuit_closure.tex
SHA256: 7c5e0b8909104814fe60809f554c8de3ac83eb9d7c92370f9ac3b02a69b396ce
Inventory use: WC1-WC28

51.1 A centered angular packet closes the physical small-gap C2 chart

Theorem 51.1. *Retain exactly the physical source hypotheses (BS1)–(BS3), the original stationary contact, every strict source gap, and the original positive infinitesimal total-scale intersection. Then the entire original C2 parent is finite tensor SOS for every independent testing pair with u, v infinitesimal or zero, every sign, and every real or complex state. In particular this includes $n = 0$, both possibilities for the bounded ratio $B/(1-r)$, and arbitrary relative rates of H/B and $1-r$.*

Joining this local assertion to (GR2)–(GR7) and its printed exact homogeneous endpoints gives an all-tests admission of precisely the source chart (BS1). No other weak-C2 source-gap opening is included.

Write $a = z/B$, $t = y/B$, $h = H/B$, $e = 1-r$. Thus a, t are positive bounded units with residues $\rho > 0$, $\tau > 1$, and B, e, h are positive infinitesimals. The event coefficients n, m, C, A, D, α have their unchanged original meanings; $\lambda = \beta = 0$. All identities below are for $F/4$.

The angular packet that stays centered at every opening. The decisive source quantities are

$$q_s = nr^2 + (m + \tfrac{1}{2})e^2 > 0, \qquad \widetilde{m} = m - a^2 q_s. \tag{WC1}$$

The exact original source formulas, and not a comparison of rates, give

$$\widetilde{m}/B \longrightarrow a_0^2 t_0/2 > 0. \tag{WC2}$$

More strongly $\widetilde{m}/B$ is jointly regular at $B = h = 0, r = 1$: its denominator has a positive-unit residue there. This remains true when e/B is unbounded. The two pieces of q_s must be kept together. Subtracting only the nr^2 piece leaves an uncontrolled e^2/B coefficient.

Here is a finite way to check all regularity assertions used below. Put

$$c_a = 1 + aBr, \quad S_c = 1 + B(1 - h + ar), \quad g = 1 + a(hB + r - 1), \quad j = c_a - (1-h)/t, \quad \Delta = \delta/B.$$

Use the original normalized formulas

$$\begin{aligned} n &= \frac{(1+ar)c_aS_c - r\Delta}{2r\Delta}, & m &= \frac{a^2(1+tB)ec_aS_c}{2g\Delta}, \\ C &= \frac{c_aM}{2(1-h)\Delta}, & Ah &= \frac{tc_a(1+ar)S_c}{2\Delta}, \\ D &= \frac{(1+aB)(1+ar)(1-atBr)S_c}{2tj\Delta}, & \alpha &= \frac{B^2(1+ar)M}{2(1-hB)\Delta}, \end{aligned} \tag{WC3}$$

where

$$M = (1+aB)(1+ar)(h+t) - a(1+tB)e(1-h)$$

and Δ is exactly the polynomial in (BS1). Every denominator divides

$$\mathcal{D} = 2rg(1-h)(1-hB)(t-1+h+aBtr)\Delta,$$

whose residue is $2(t_0-1)(1+a_0) > 0$. For an expression linear in n, m, C, Ah, D, α, clear $\mathcal{D}$ and its printed pure-source denominator. Monic division of the numerator by B, once or twice as printed, is exact. Evaluating the resulting polynomial quotient at $B = h = 0, r = 1$ gives the residues below. This supplies finite polynomial identities, rather than an appeal to asymptotic contact cancellation. For (WC2), the constant numerator vanishes identically at $B = 0$ for every h, r; at that face

$$n = \frac{e}{2r(1+a)}, \quad m = \frac{a^2e}{2(1-ae)(1+a)}, \quad m/a^2 = nr^2 + (m+\tfrac{1}{2})e^2.$$

The focused checker implements these finite divisions with $\mathcal{D}$ kept as a common unit denominator.

Bounded testing ratio and all three hard state directions. Take $u, v > 0$ and first suppose the pure-testing ratio $w = v/u$ is bounded, including infinitesimal w. At $Z \neq 0$ divide all states by Z and set $Z = 1$; the homogeneous restoration is described below. Set

$$E = tuW - v, \qquad N_y = (Y - huW)/\sqrt{h}, \qquad \xi = X + ru. \tag{WC4}$$

Equivalently

$$W = (v+E)/(tu), \quad Y = h(v+E)/t + \sqrt{h}\,N_y, \quad X = \xi - ru.$$

Only a pure-testing inverse and a pure-source square root occur. In this frame the complete hard quadratic has bounded coefficients and joint residue

$$(E, N_y, \xi): \qquad \operatorname{diag}\left(\frac{1}{2(t_0-1)}, \frac{t_0}{2}, \frac{1}{2}\right) \succ 0. \tag{WC5}$$

For example $A = Ah/h$ always occurs multiplied by the square of $h(v+E)/t + \sqrt{h}N_y$, so its normalized coefficients contain Ah, $\sqrt{h}\,Ah$, or hAh, and no negative power of h. The only testing-ratio dependence in the hard block is polynomial in bounded w, such as mw^2; all its coefficients are bounded. The displayed residue is obtained from the whole original four-state parent, not a restricted soft graph.

Exact scalar identity with two retained weighted roots. Retain the scalar coefficients from (WA3)–(WA5), writing the coefficient of UW^2 as a_w to avoid collision with $a = z/B$:

$$\begin{aligned} a_w &= (C+D+\tfrac{1}{2})(1-h)^2 + Ah\,h, \quad b_2 = 2mh + C(1-h)^2 + Ah\,h + (\alpha+\tfrac{1}{2})h^2, \\ b_4 &= h^2(m+\alpha+\tfrac{1}{2}) + Ah\,h, \quad d_2 = (n+\tfrac{1}{2})r^2 + 2Dr, \quad d_4 = (n+D+\tfrac{1}{2})r^2, \\ e_0 &= me + D(1-h) + \alpha h + \tfrac{1}{2}, \quad e_1 = mhe + r(D+\tfrac{1}{2})(1-h) + (\alpha+\tfrac{1}{2})h. \end{aligned} \tag{WC6}$$

Let $x = aB$, $y = tB$, $k = \sqrt{x}$, $s = \sqrt{t/a}$ and

$$A_t = uv, \quad B_t = \frac{kv^2}{su}, \quad C_t = xs, \quad G = kv, \quad F_v = V - y, \quad H_u = \frac{a}{t}\frac{v^2}{u} - u. \tag{WC7}$$

Then $A_tB_tC_t = G^3$, and all three circuit amplitudes equal G at the actual contact $u = k, w = s$. Both H_u and F_v vanish there. Define

$$\begin{aligned}
c_F &= (b_2 - a^2nr^2)/t^2,\\
q_V &= a_w/t^2 + D - 2e_0/t + 2aq_s/t,\\
f_A &= Ah\,h/t^2 + d_2 - 2e_1/t + 2anr^2/t + d_4U + b_4V/t^2,\\
f_B &= \widetilde{m}/(ta^2B),\\
f_C &= [\alpha - c_Fy^2 + \alpha U]/(taB^2),\\
f_G &= -(q_V + 2c_Fy)/(aB).
\end{aligned} \tag{WC8}$$

The entire scalar graph $E = N_y = \xi = 0$ is exactly

$$\begin{aligned}
\mathcal{S} = {} & q_sH_u^2 + nr^2u^2H_u^2 + c_FF_v^2\\
& + f_AA_t^2 + f_BB_t^2 + f_CC_t^2 - f_GG^2.
\end{aligned} \tag{WC9}$$

This is an unconditional polynomial identity after multiplying by the pure-testing square u^2. To see its cancellation directly,

$$H_u^2 = U - 2aV/t + a^2V^2/(t^2U).$$

Thus $q_sH_u^2$ removes the full q_sU coefficient and replaces m by $\widetilde{m}$ in the V^2/U coefficient. The second square removes nr^2U^2 and changes only the displayed V^2 and UV coefficients. Nothing has been discarded from the original parent, and source physicality is not transferred to a remainder.

The finite source divisions described after (WC3) give

$$\begin{aligned}
c_F \to 1/(2t_0), \quad f_A, f_B, f_C \to \tfrac{1}{2}, \quad f_G \to \tfrac{3}{2},\\
q_V/B \to -1 - 3a_0/2, \qquad \partial_U f_C \to (1 + a_0)/(2a_0).
\end{aligned} \tag{WC10}$$

All coefficient arrays in f_A, f_C are bounded; each is affine in U, V. The constant f_B, f_G have no testing dependence. These are joint regular reductions and contain no e/B or h/e assumption. The two positive weighted roots

$$R_q = \sqrt{q_s}\,H_u, \qquad R_n = \sqrt{n}\,ruH_u \tag{WC11}$$

remain inside the root list, including literal $n = 0$.

Finite contact lifting of the scalar circuit. Apply the already proved three-circuit contact-lift lemma with $\lambda = 1/2$ to the last four terms of (WC9), retaining the strict $c_FF_v^2$ pivot and the two squares (WC11). Its conductors are the exact separated identities

$$G(u - k) = k(A_t - G), \qquad G(w - s) = s(B_t - G). \tag{WC12}$$

The required finite Taylor arrays are precisely (IC11)–(IC14), with the present f_A, f_B, f_C, f_G and x, y, s. They apply because these functions are affine in $U = u^2, V = u^2w^2$, with bounded pure-source coefficients. Actual source contact gives all three value/first-jet balances: the squares removed in (WC9) have zero value and first derivative there. No Hessian of a modified physical parent is asserted. Consequently the scalar has a finite strict bounded-residue Gram on R_q, R_n, F_v and the finite three-circuit root list.

Every full first normal is allocated on the same roots. Use the complete graph normal coefficients f_w, f_z, k_w, k_z in (WA6)–(WA7), equivalently

$$\begin{aligned} f_w &= -m + [-mh + (D+\tfrac12)(1-h)]U, \\ f_z &= -nr(1+U)(1+V) + me - (D+\tfrac12)V(1+rU) + \tfrac12 e(1+V), \\ k_w &= -(C+D+\tfrac12)(1-h) + Ah + Ah\,U \\ &\quad + [-C(1-h) + Ah + m + (\alpha+\tfrac12)h]V + [Ah + h(m+\alpha+\tfrac12)]UV, \\ k_z &= D - \alpha + [-me + r(D+\tfrac12) - \alpha - \tfrac12]U. \end{aligned} \tag{WC13}$$

Put

$$P_Y = 2(k_w/t + k_z), \qquad P_E = 2(A_w/U - mV/U)/t^2 - 2(e_0 + e_1U)/t,$$

where A_w is the complete W^2 coefficient in (WS2). The exact first derivatives of $F/4$ in the frame (WC4) are

$$\begin{aligned} \mathcal{L}_{N_y} &= \sqrt{h}\, vP_Y, \\ \mathcal{L}_E &= 2mvw^2/t^2 + vP_E, \\ \mathcal{L}_\xi &= 2u\{w^2 f_w/t + f_z\}. \end{aligned} \tag{WC14}$$

These identities are verified by differentiating the entire original parent after (WC4), retaining all original packets.

Extract the following complete angular contributions:

$$\begin{aligned} \mathcal{L}_\xi &= -2a\sqrt{q_s}\, R_q + 2\sqrt{n}\, uR_n + \mathcal{R}_X, \\ \mathcal{L}_E &= 2a\sqrt{q_s}\,(w/t)R_q + \mathcal{R}_E, \\ \mathcal{L}_{N_y} &= \sqrt{h}\,\mathcal{R}_Y, \quad \mathcal{R}_Y = vP_Y. \end{aligned} \tag{WC15}$$

The displayed coefficients are infinitesimal because q_s, n are infinitesimal and u, w are bounded. No inverse of either weight appears. The exact residuals in (WC15) are finite polynomials in u, w of the following explicitly checkable form:

$$\mathcal{R}_\bullet = B\sum_{j=0}^{3} c_{\bullet j} uw^j + \sum_{i\ge 3,\, j\ge 1} d_{\bullet ij} u^i w^j, \tag{WC16}$$

where all source coefficients are bounded; only the monomials actually occurring in (WC13)–(WC15) are included. Their exact supports are

	Buw^j	u^iw^j $(i \ge 3)$
X	$j = 0, 2$	$(i,j) = (3,2), (5,2)$
E	$j = 1, 3$	$(i,j) = (3,1), (3,3), (5,3)$
Y	$j = 1$	$(i,j) = (3,1), (3,3), (5,3).$

(WC17)

Equation (WC16) is a finite numerator-division assertion, checked with (WC3) and the common unit denominator $\mathcal{D}$. It does not follow merely from normal vanishing. In particular the apparent uncentered e/B terms cancel in the uw^0, uw^2, uw^3 entries only after retaining all of q_s in (WC15).

For clarity, its complete arrays may be written without any coefficient-extraction convention. Put

$$\begin{aligned} K_X &= -nr + (m+\tfrac12)e - aq_s, \\ K_E &= a_w/t^2 - e_0/t + aq_s/t, \\ K_Y &= [-(C+D+\tfrac12)(1-h) + Ah]/t + D - \alpha, \\ x_3 &= 2\{(D+\tfrac12)[(1-h)/t - 1] + e/2 - nr(1+a/t) - mh/t\}, \\ x_5 &= -2r(n + D + \tfrac12). \end{aligned}$$

Then exactly

$$\begin{aligned}
\mathcal{R}_X &= 2K_X u - 2\widetilde{m}\,uw^2/t + x_3u^3w^2 + x_5u^5w^2,\\
\mathcal{R}_E &= 2K_E uw + 2\widetilde{m}\,uw^3/t^2 + (2Ah\,h/t^2 - 2e_1/t)u^3w + 2b_2u^3w^3/t^2 + 2b_4u^5w^3/t^2,\\
\mathcal{R}_Y &= 2K_Y uw + 2[Ah/t - me + r(D+\tfrac{1}{2}) - \alpha - \tfrac{1}{2}]u^3w\\
&\quad + 2[-C(1-h) + Ah + m + (\alpha+\tfrac{1}{2})h]u^3w^3/t\\
&\quad + 2[Ah + h(m+\alpha+\tfrac{1}{2})]u^5w^3/t.
\end{aligned} \tag{WC17a}$$

The only source quotients required for these arrays are $\widetilde{m}/B, K_X/B, K_E/B, K_Y/B$. Each has a unit denominator after exact monic division. Their numerators vanish at $B=0$ for every h, r, not merely at the joint corner. For example $m/a^2 = q_s$ and $-nr+(m+1/2)e = m/a$ on that face prove $K_X = 0$. Substituting (WC3) at $B=0$ gives $K_Y = 0$ directly; that equality implies $a_w = Ah + t(1-h)D$, which reduces $K_E = 0$ to $Ah/t - me - 1/2 + m/a = 0$. These are exact face identities with the same unit denominators, so the four polynomial divisions prove (WC16) for arbitrary independent source openings.

Every monomial in (WC16) now has an explicit legal circuit allocation:

$$\begin{aligned}
Bu &= uC_t/(as), & Buw &= (B/k)G,\\
Buw^2 &= (Bs/k)B_t, & Buw^3 &= (Bs/k)wB_t,\\
u^iw^j &= u^{i-2}w^{j-1}A_t \quad (i\ge 3, j\ge 1).
\end{aligned} \tag{WC18}$$

All resulting coefficients are infinitesimal. Their finite first divided differences in u are bounded, and the divided differences in w after setting $u=k$ are infinitesimal: every w-dependent coefficient retains u or k. Both properties follow directly from the displayed monomials, with no testing-ratio inverse. Each residual normal vanishes at the actual contact, because the original complete normal and H_u vanish there. The monic division (IC21), followed by (WC12), therefore expresses each full normal with infinitesimal coefficients on the same strict circuit roots. Together with (WC5), (WC9), and (WC15), this proves a finite strict bounded-residue Gram for the entire parent at every bounded w.

The reciprocal ratio uses the physical source scale bound. Suppose $w = v/u$ is unbounded, so $\omega = u/v$ is infinitesimal. Return to homogeneous states. Set

$$p = uW, \quad q = vZ, \quad E = tp - q, \quad Y = hp + \sqrt{h}N_y, \quad \xi = X + ruZ,$$

and use the complete root list

$$E,\ N_y,\ \xi, \quad P = \sqrt{m}\,vW, \quad Q = \sqrt{\alpha}Z, \quad R = u^2W, \quad S = uvW. \tag{WC19}$$

Here m/e and α/B^2 are positive bounded units. BS3 gives B/e and n/e bounded, hence n/m bounded. The pure-source comparison coefficients

$$C_* = \frac{t+h}{2(1-h)}, \qquad (Ah)_* = t/2, \qquad D_* = \frac{1}{2(t+h-1)}$$

satisfy

$$C - C_* = Bc_B + ec_e, \quad Ah - t/2 = Ba_B + ea_e, \quad D - D_* = Bd_B + ed_e, \tag{WC20}$$

with bounded pure-source coefficients. To check this finitely, clear the unit denominators in (WC3), subtract the numerator at $B=0$, divide that difference by B, and divide its remaining $B=0$ numerator by $r-1=-e$. The latter numerator vanishes identically at $r=1$ for every h. These are the exact finite divisions implemented by the focused checker.

The quadratic comparison base

$$C_*(p-Y)^2 + \frac{t}{2h}Y^2 + D_*(p-Y-q)^2 + \tfrac{1}{2}(p-Y)^2 - pq$$

is exactly

$$D_*E^2 - 2D_*\sqrt{h}\,EN_y + \frac{t(t+2h-1)}{2(1-h)(t+h-1)}N_y^2. \tag{WC21}$$

It has strict diagonal residue $1/[2(t_0-1)], t_0/2$. The comparison error has infinitesimal coefficients on (WC19):

$$\sqrt{e}\,p = \sqrt{e/m}\,\omega P, \qquad q = tp - E, \qquad Y = hp + \sqrt{h}N_y. \tag{WC22}$$

For the A term use the exact expansion $(Ah - t/2)Y^2/h = (Ah - t/2)[hp^2 + 2\sqrt{h}\,pN_y + N_y^2]$. Thus no e/h or h/e comparison is introduced.

With $X + uZ = 0$, the exact higher-angle part of the C, A, D packets and half core, after removing their quadratic base, is

$$\begin{aligned} &U\{AY^2 - 2Dq(p - Y - q) + q^2/2 - pq\} \\ &\quad + V\{C(p-Y)^2 + (A+\tfrac{1}{2})Y^2\} + (A+\tfrac{1}{2})UVY^2 + (D+\tfrac{1}{2})U^2q^2. \end{aligned} \tag{WC23}$$

Substitute (WC22). Its residue on (R, S) is strictly diagonal, with pivots $t_0^2/2, t_0/2$. Every other coefficient is bounded and infinitesimal: for instance $UpN_y = uRN_y$, $VpE = vSE$, $UVp^2 = US^2$. Terms with $A = Ah/h$ are regular by the same explicit Y expansion used after (WC22).

To restore the complete X dependence write

$$B_x = X + uZ = \xi + euZ = \xi + t(e/\sqrt{m})\omega^2 P - e\omega E. \tag{WC24}$$

Both added coefficients are infinitesimal. Relative to $B_x = 0$, the exact D packet and half core add

$$2(D+\tfrac{1}{2})uvB_x(p - Y - (1+U)q) + (D+\tfrac{1}{2})UVB_x^2 + \tfrac{1}{2}(1+V)B_x^2. \tag{WC25}$$

The last term supplies the strict ξ pivot $1/2$; all remaining Gram coefficients are infinitesimal, using $uvp = uS$. The remaining three original positive packets retain their full roots:

$$\begin{aligned} \sqrt{m}\,M &= P - \sqrt{m}\,B_x + \sqrt{m}\,uvY, \\ \sqrt{\alpha}\,(vY - Z) &= -Q + \sqrt{\alpha}hS + \sqrt{\alpha h}\,vN_y, \\ \sqrt{n}\,X &= \sqrt{n}\,\xi + r\sqrt{n}\,\omega E - rt\sqrt{n/m}\,\omega^2 P. \end{aligned} \tag{WC26}$$

Retain the original positive pure-testing D_u, D_v factors. The first two roots have leading entries $P, -Q$; every coefficient of the third is infinitesimal, including $n = 0$. No square or first normal was dropped. Hence the complete parent has a bounded Gram on (WC19) with strict diagonal residue

$$\operatorname{diag}\left(\frac{1}{2(t_0-1)}, \frac{t_0}{2}, \frac{1}{2}, 1, 1, \frac{t_0^2}{2}, \frac{t_0}{2}\right) \succ 0. \tag{WC27}$$

This proves the reciprocal ratio without comparing any source scale with u or v.

Restoration, verification, and exact admission. The exact sign gauges $(u, Y, Z) \mapsto (-u, -Y, -Z)$ and $(v, X, Z) \mapsto (-v, -X, -Z)$ preserve the entire original parent, including the nX^2 packet. Literal axes have the exact identities

$$\begin{aligned} (F/4)|_{u=0} &= nD_vX^2 + m(vW - X)^2 + (C + A + \tfrac{1}{2})D_vY^2 + D(Y + vZ)^2 \\ &\quad + \alpha(vY - Z)^2 + \tfrac{1}{2}D_vX^2, \\ (F/4)|_{v=0} &= nD_uX^2 + (m+\tfrac{1}{2})(X + uZ)^2 + (C + D + \tfrac{1}{2})(uW - Y)^2 \\ &\quad + AD_uY^2 + \alpha D_uZ^2. \end{aligned} \tag{WC28}$$

All displayed coefficients are nonnegative. Homogenize the bounded-ratio proof by multiplying its identity by Z^2: E, N_y, ξ become the homogeneous linear forms, and each scalar circuit root and H_u

is multiplied by Z. No Z inverse remains, so the identity extends to $Z = 0$. The same real identity applied to the real and imaginary state parts gives the complex-state assertion. The previously proved GR join supplies every remaining independent testing chart and projective endpoint on this exact source chart.

The new checker verifies the scalar identity (WC9), all three complete first normals by direct original-parent differentiation, every entry of the hard residue, the finite source divisions and normal supports, the exact reciprocal higher-angle identity, and both strict higher-angle pivots. These are focused checks of the new argument; no previous verification suite is replayed. The finite Gram/positivity proof is the argument above. It closes only the BS1 small-$1 - r$, z/B-unit source chart, including its printed strict gaps and $n = 0$; the other C2 source-gap complements remain separate obligations.

Chapter 52

Exact weak-C2 source complement after the small-gap admission

Component `pe2fc3cbe3819`
Source: `R13/advances/revision13_weak_local/frontier_audit.tex`
SHA256: `e2fc3cbe38194b492857ca85f59e74674af370627b314af72c93af4b4e1b8c64`
Inventory use: R9 source/remote extension WF4-WF7 and exact ordered source table

52.1 Exact weak-C2 source complement after the small-gap admission

This audit uses the original source alternatives (WII1)–(WII6) and precisely the proved WA, WE, SL, and WC admissions. The finalized WC proof and its checker are unchanged.

Common quantified source domain. Work in the canonical reflected strong-C/weak-α C2 II orientation. Retain the original physical source and stationary contact, every strict source gap, $n \geq 0$, $d = 1$, $B, z, y > 0$ infinitesimal, and the original positive-infinitesimal total-scale intersection. Every row below is further intersected with the complement of every inherited admission and every exact symmetry image. No row is asserted nonempty.

Use consistent notation

$$a = z/B, \quad t = y/B, \quad h = H/B, \quad l = 1 - h = L/B, \quad e = 1 - r, \quad g = G/B.$$

The positive bounded quantities h, l, r, e, g, j each have type I (infinitesimal) or U (positive bounded unit). The quantities $a, t > 0$ can additionally have type O (unbounded). Retain exactly

$$g = 1 - ae + hz > 0, \qquad j = 1 + zr - l/t > 0, \qquad B - zyr > 0$$

and every other original physical/contact condition. Shape entries are not independent choices.

Some useful exact implications are

$$\begin{gathered} ae < 1 + hz, \qquad e \in \mathrm{U} \Longrightarrow a \text{ bounded}, \qquad a \in \mathrm{O} \Longrightarrow e \in \mathrm{I}, \\ a \in \mathrm{I} \Longrightarrow g \sim 1, \qquad g \in \mathrm{I} \Longrightarrow ae \sim 1. \end{gathered} \tag{WF1}$$

Thus infinitesimal a is impossible when g is infinitesimal. For t a unit and h infinitesimal, l is a unit; j a unit implies $t_0 > 1$, whereas j infinitesimal implies $t_0 = 1$. The directed strong-C condition also retains the exclusion from (WII6): $r, g, H/y, j$ cannot all be infinitesimal.

Exactly what the current admissions remove. Call t, g, j units and h infinitesimal the regular-gap A opening. On this source l is a unit. If e is a unit, a is bounded by (WF1). A unit r is covered by WA for unit a and by WE for infinitesimal a; infinitesimal r is covered by SL. Therefore every e-unit source on this opening is admitted, with no omitted r-type. If e is infinitesimal, r is a unit and WC removes exactly the unit-a type. The exact remainder on this opening is

$$\boxed{e \in \mathrm{I}, \qquad a \in \mathrm{I} \text{ or } a \in \mathrm{O}.} \tag{WF2}$$

An ordered disjoint residual partition. With the common domain and all standing exclusions imposed, the remaining source shapes are the following mutually disjoint rows:

$$\begin{array}{r|l}
R1 & t \in \mathrm{I} \\
R2 & t \in \mathrm{O} \\
R3 & t, h \in \mathrm{U}, \quad l \in \mathrm{I} \\
R4 & t, h, l \in \mathrm{U}, \quad j \in \mathrm{I} \\
R5 & t, h, l, j \in \mathrm{U}, \quad r \in \mathrm{I} \\
R6 & t, h, l, j, r \in \mathrm{U}, \quad g \in \mathrm{I} \\
R7 & t \in \mathrm{U},\ h, g \in \mathrm{I}, \quad \text{excluding } r, j \in \mathrm{I} \\
R8 & t, g \in \mathrm{U}, \quad h, j \in \mathrm{I} \\
R9 & t, g, j \in \mathrm{U}, \quad h, e, a \in \mathrm{I} \\
R10 & t, g, j \in \mathrm{U}, \quad h, e \in \mathrm{I},\ a \in \mathrm{O}.
\end{array} \tag{WF3}$$

In R7 the excluded r, j-intersection is exactly the already removed two-weak case, because $H/y = h/t$ is infinitesimal.

No a-type is silently discarded. R1–R4 and R8 retain all three positive a-types subject to (WF1) and the original source constraints. R5 has bounded a, hence type I or U. R6–R7 have unit or unbounded a, with $ae \sim 1$. R9 and R10 specify the remaining a-type explicitly.

The unbounded original coefficients supplied by (WII3) are: C, D in R1; C, A in R2; C in R3; D in R4; n in R5; m in R6; A, m in R7; A, D in R8; and A in R9–R10. Additional n, m, D openings remain governed by the exact r, g, j conditions. In R1, A is weak; in R2, D is weak. Two unbounded adjacent coefficients alone are not an invocation of a different dominant-event theorem.

For exhaustiveness, first split t into its three types. At unit t, split h into infinitesimal or unit. For unit h, split l, then j, then r, then g. The all-unit remainder has positive-unit total scale by (WII4), hence is absent from the original residual intersection. These are R3–R6. For infinitesimal h, split g and then j, giving R7–R8. With both gaps units, (WF2) leaves exactly R9–R10. This proves the partition, without asserting nonemptiness after all inherited admissions and symmetry images are removed.

52.1.1 A remote reduction for the simultaneous small-e, small-a opening

Lemma 52.1. *On R9, every nonlocal independent testing chart is finite tensor SOS, with every original state component and literal projective endpoint retained. The only remaining testing problem is the full local parent with both u, v infinitesimal or zero. This is a remote reduction, not a new all-tests source admission.*

The exact normalized source (WE1) remains valid without a unit-a assumption. On R9, its polynomials satisfy

$$\Delta \sim 1, \qquad K = (1+a)(h+t)(1+aB) \sim t, \qquad Q \sim 1, \tag{WF4}$$

where Q is the exact positive polynomial in (BS2). These are joint reductions: B, a, h, e are infinitesimal, $r = 1 - e$, and t is a positive unit. The same exact source identity gives

$$2r\Delta n = eQ - BK, \qquad 0 < B/e \le Q/K, \qquad 0 \le n/e \le Q/(2r\Delta). \tag{WF5}$$

In particular n is infinitesimal, including literal $n = 0$.

Since $g = 1 + a(hB - e) \sim 1$ and $j = 1 + aBr - (1-h)/t \sim 1 - 1/t$ is a positive unit, $t_0 > 1$. Writing $c_a = 1 + aBr$, $S_c = 1 + B(1 - h + ar)$, and

$$M_* = (1 + aB)(1 + ar)(h + t) - a(1 + tB)e(1 - h),$$

the remaining exact source formulas yield

$$\begin{aligned}
\frac{m}{a^2 e} &= \frac{(1 + tB)c_a S_c}{2g\Delta} \longrightarrow \frac{1}{2}, \\
Ah &= \frac{tc_a(1 + ar)S_c}{2\Delta} \longrightarrow t_0/2, \\
C &= \frac{c_a M_*}{2(1 - h)\Delta} \longrightarrow t_0/2, \\
D &= \frac{(1 + aB)(1 + ar)(1 - atBr)S_c}{2tj\Delta} \longrightarrow \frac{1}{2(t_0 - 1)}, \\
\frac{\alpha}{B^2} &= \frac{(1 + ar)M_*}{2(1 - hB)\Delta} \longrightarrow t_0/2.
\end{aligned} \tag{WF6}$$

All denominators have positive-unit residues. Hence $m = O(a^2 e)$, $\alpha = O(B^2)$, $A^{-1/2} = \sqrt{h/(Ah)}$ is infinitesimal, and the complete coefficient conditions actually used by (GR2)–(GR7) hold:

$$n, m, \alpha, A^{-1/2} \in \mathrm{Inf}, \qquad C_0 = t_0/2, \qquad D_0 = 1/[2(t_0 - 1)]. \tag{WF7}$$

Those exact complete-parent identities contain no inverse a. Their small m packet is retained inside $\sqrt{m}$, and the n and α roots permit zero weights without division. Every printed strict residue determinant is consequently unchanged. The existing coefficient-bound and homogeneous-endpoint arguments therefore apply verbatim to every nonlocal test. This proves the lemma without replaying any unchanged algebraic checker. It does not import the local WC circuit.

The remaining local obstruction is a normalization issue. WC does not yet give the R9 local theorem. Its present scalar coordinates contain

$$s = \sqrt{t/a} \text{ unbounded}, \qquad a\partial_U f_C = \alpha/(tB^2) \longrightarrow \tfrac{1}{2}. \tag{WF8}$$

The unit-s hypothesis and bounded finite coefficient arrays of the three-circuit lift therefore fail at the new source boundary. There is no license to call U/a small: the source and testing fields are independent. In the WC reciprocal frame, the bound on n/m is also lost; (WF5)–(WF6) give only $n/m = O(a^{-2})$, so they do not control $\sqrt{n/m}(u/v)^2$ uniformly.

The older WE-rescaled WA frame has the same separate failure: scaling its angular root by $\sqrt{e}$ leaves nonzero-residue $J_s/(a\sqrt{e})$ and $f_s/(a\sqrt{e})$ coefficients. These are limitations of the stated frames, not non-SOS obstructions.

The smallest decisive new lemma is a contact lift with the reciprocal source slope $c = \sqrt{a/t}$ infinitesimal and source Bob radius $\sqrt{y}$ infinitesimal, retaining the whole centered mass $q_s = nr^2 + (m + 1/2)e^2$. It must give explicit finite conductors for the U/a opening and retain $\sqrt{n}\,uZ$, or an exact contact-centered replacement, as an additional weighted root instead of dividing it by $\sqrt{m}$. It must include both full soft states, every normal, $n = 0$, arbitrary h/a, and all independent testing ratios. No such lemma is proved here.

R10 is separate: a is unbounded, ae is bounded, and g remains a positive unit. Normalizing by the source product ae is a natural next calculation, but neither the unit-a WC theorem nor the R9 remote reduction supplies that source-domain admission.

Chapter 53

A three-monomial contact lift with one unit coordinate

Component p466b31015532
Source: R13/advances/three_circuit/three_circuit_lift.tex
SHA256: 466b31015532add0a1fe6fd3d0b6ce3b922c0dd5600800cd9a685450a6a0f9ef
Inventory use: complete finite contact-lift lemma | finite cubic SOS, conductors, scalar and normal lifts | C14_ ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

53.1 A three-monomial contact lift with one unit coordinate

Work in the same tensor ring $T = K_c \otimes_{\mathbb{R}} K_t$ and bounded subring B as throughout this package. Every inverse and radical below is of a pure-field or positive separated element. Let $k > 0$ be a pure-source infinitesimal and $s > 0$ a positive pure-source bounded unit. Let $u > 0$ be a pure-testing infinitesimal and $t > 0$ a positive pure-testing bounded element (possibly infinitesimal). Define

$$A_1 = u^2 t, \qquad A_2 = \frac{kut^2}{s}, \qquad A_3 = k^2 s, \qquad G = kut, \qquad d_i = A_i - G.$$

Then $A_1 A_2 A_3 = G^3$. Two exact conductor identities are

$$G(u - k) = k d_1, \qquad G(t - s) = s d_2. \tag{53.1}$$

They permit a unit testing ratio t without comparing a mixed tensor.

Put $Q = G \sum_i d_i$ and $L = \sum_i A_i^2 - 3G^2$. For $a_i = A_i^{1/3}$, the elementary cubic identity gives

$$\begin{aligned} Q &= \frac{1}{2} \sum_{1 \le i < j \le 3} \sum_{\ell=1}^{3} \{\sqrt{G a_\ell}(a_i - a_j)\}^2, \\ L &= \sum_i d_i^2 + 2Q. \end{aligned} \tag{53.2}$$

Thus L has a positive constant diagonal Gram on a finite list of three difference roots and nine cubic-identity roots. No assertion that every nonnegative tensor is SOS is used.

Exact finite lift. Let $\lambda \in B_c$ have positive real residue, and consider

$$F = \sum_{i=1}^{3} (\lambda + \epsilon_i) A_i^2 - (3\lambda + \epsilon_g) G^2, \qquad h = \sum_i \epsilon_i - \epsilon_g.$$

Write $\Delta_u = u - k$, $\Delta_t = t - s$. Assume explicit finite identities

$$\epsilon_i - \epsilon_{i0} = K_{iu} \Delta_u + K_{it} \Delta_t, \qquad \epsilon_i, \epsilon_{i0} \in B_{\mathrm{Inf}}, \tag{53.3}$$

$$h = b_u\Delta_u + b_t\Delta_t + \sum_{j,\ell\in\{u,t\}} H_{j\ell}\Delta_j\Delta_\ell, \qquad b_u, b_t \in K_c, \tag{53.4}$$

$$2(\epsilon_{10} - \epsilon_{30}) + kb_u = 0, \qquad 2(\epsilon_{20} - \epsilon_{30}) + sb_t = 0. \tag{53.5}$$

Require $K_{iu}, H_{uu}, H_{ut}, H_{tu} \in B$ and $K_{it}, H_{tt} \in B_{\mathrm{Inf}}$. The distinction is essential: the second conductor has a unit coefficient, so boundedness alone of the second-coordinate arrays does not suffice.

Then F is tensor SOS. Indeed, expanding $A_i = G + d_i$ and applying (53.1) and (53.5) yields the exact identity

$$\begin{aligned} F - \lambda L = {} & \sum_i \epsilon_i d_i^2 + 2\epsilon_{30} Q + 2\sum_i (kK_{iu} d_1 d_i + sK_{it} d_2 d_i) \\ & + k^2 H_{uu} d_1^2 + ks(H_{ut} + H_{tu}) d_1 d_2 + s^2 H_{tt} d_2^2. \end{aligned} \tag{53.6}$$

In detail, the two bare-G first-order terms are cancelled by $G^2 b_u \Delta_u = kb_u G d_1$ and $G^2 b_t \Delta_t = sb_t G d_2$. All remaining coefficients in (53.6) are bounded infinitesimals. The strict diagonal Gram for λL in (53.2) therefore remains strictly positive in real residue. The bounded-Gram lemma supplies a finite tensor SOS. This argument uses no power series or mixed inverse.

Normal extension. Suppose a coupling to a normalized hard state normal has the exact form $\ell = \sum_i c_i A_i$ with $c_i \in B_{\mathrm{Inf}}$, $\sum_i c_{i0} = 0$, and

$$\sum_i (c_i - c_{i0}) = J_u\Delta_u + J_t\Delta_t, \qquad J_u \in B, \quad J_t \in B_{\mathrm{Inf}}.$$

Then

$$\ell = \sum_i c_i d_i + kJ_u d_1 + sJ_t d_2$$

has infinitesimal coefficients on the same root list. It can be joined to a hard normal block with strictly positive real residue. Extra already-controlled scalar roots may be appended, provided their coupling coefficients are likewise infinitesimal. The source-specific arrays and hard-block bounds remain separate proof obligations.

Small-L coordinates. For $x = k^2$, $s = \sqrt{\eta\zeta}$ and $t = v/u$, this gives

$$(A_1, A_2, A_3, G) = \left(uv, \sqrt{\frac{x}{\eta\zeta}}\frac{v^2}{u}, \sqrt{\eta\zeta}\, x, \sqrt{x}\, v \right).$$

The lemma applies only when the stated pure-field size assumptions and all finite scalar and normal identities have been proved. A leading three-term AM–GM form alone does not close a source domain.

Chapter 54

A five-root local certificate on an A-unbounded weak-C2 subchart

Component p97700cec8dc8
Source: R13/advances/weak_A/five_root_local.tex
SHA256: 97700cec8dc883e38200b3d41d6f0aa314995bbda7be5fecd6f9ea8864e1fda0
Inventory use: WA1-WA10

54.1 A five-root local certificate on an A-unbounded weak-C2 subchart

Revision 10 scope update. Section 56.1 extends this admission to z/B positive infinitesimal with all other source hypotheses retained. The unit-ratio restriction below records the original revision-9 theorem.

Theorem 54.1. *Retain the original physical C2 source, its exact stationary contact, all strict source gaps, and $n \geq 0$. Normalize $d = 1$. Suppose B, z, y are positive infinitesimals, $r, G/B, j, y/B, L/B$ are positive bounded units, and H/y is infinitesimal. Impose in addition*

$$z/B \text{ and } 1 - r \text{ are positive bounded units.} \tag{WA1}$$

Then the entire original parent is a finite tensor sum of squares for arbitrary independent testing half-angles u, v that are infinitesimal or zero, including either sign and complex states. There is no comparison of a testing scale with B, H, z, y. The total-scale parameter δ_{tot} is positive infinitesimal. This is a local testing certificate on the stated source subchart; it does not assert that its intersection with the final residual complement is nonempty, or supply an unproved remote-testing join.

No membership in the residual complement is needed in this proof. When applied to the original residual problem, its existing complement condition is still intersected with the source chart; no nonemptiness of that intersection is inferred.

The original omitted event coefficients remain $\lambda = \beta = 0$. Here $B = H + L$, and the actual contact in the positive gauge is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(\sqrt{z}, \sqrt{y}; \frac{B}{\sqrt{zy}}, -r\sqrt{z}, \frac{H}{\sqrt{y}}, 1\right).$$

The source ratio $\ell = B/z$ below is not an event coefficient.

The source pivots survive the small H opening. Write $h = H/B$, $\kappa = B/y$, $\ell = B/z$. Both κ and ℓ are positive bounded units, and h is infinitesimal. The original formulas imply that n, m, C, D, α are bounded, that m, C, D are positive bounded units, and that Ah is a positive

bounded unit. Thus A is unbounded and $Ah^2, \sqrt{A}\,h, A^{-1/2}$ are infinitesimal. No bound on the relative rates of h and B is used.

Here is an explicit check of the radial pivot. With $T_c = B + zr$, the original formula is

$$\frac{\alpha}{B^2} = \frac{T_c}{\delta}\frac{M/B^2}{2(1-H)}, \qquad M - yG = H(1+z)T_c + z\{yj + rL(1+y)\} > 0. \tag{WA2}$$

Consequently $M/B^2 \geq (y/B)(G/B)$ is bounded below by a positive real, and it is bounded above by the original formula for M and the unit ratios in (WA1). Since T_c/δ is a positive bounded unit, α/B^2 is a positive bounded unit uniformly as $H/y \to 0$. Likewise

$$m = \frac{z^2(1+y)(1-r)(1+zr)(1+zr+L)}{2G\delta}$$

is a positive bounded unit under (WA1); the local residue argument uses only $n \geq 0$. Finally A is the only unbounded retained coefficient, so $\delta_{\mathrm{tot}} = 1/(1+n+m+C+A+D+\alpha)$ is positive infinitesimal. These arguments preserve all original coefficient and source-gap hypotheses.

Five homogeneous contact roots. Put $U = u^2$, $V = v^2$, and define

$$\xi = X + ruZ, \qquad q = \sqrt{A}\,(Y - huW), \qquad R_1 = uW - \kappa vZ, \quad R_2 = vW - \ell uZ, \quad T = uvW - BZ. \tag{WA3}$$

Every root vanishes at the actual source contact. Inverses and radicals in their coefficients are pure source. The large A packet is retained in q; no positive packet is subtracted from the parent and no physicality hypothesis is transferred to a modified remainder.

Use the following source coefficients for the exact scalar graph:

$$\begin{aligned}
a &= (C + D + \tfrac{1}{2})(1-h)^2 + Ah^2, & a_4 &= Ah^2,\\
b_2 &= 2mh + C(1-h)^2 + (A + \alpha + \tfrac{1}{2})h^2, & b_4 &= h^2(m + A + \alpha + \tfrac{1}{2}),\\
c &= \alpha, & c_2 &= nr^2 + (m + \tfrac{1}{2})(1-r)^2 + \alpha,\\
c_4 &= nr^2, & d_2 &= (n + \tfrac{1}{2})r^2 + 2Dr,\\
d_4 &= (n + D + \tfrac{1}{2})r^2.
\end{aligned}$$

Thus the graph on $\xi = q = 0$ is

$$\begin{aligned}
F_g/4 = {} & [aU + a_4U^2 + (m + b_2U + b_4U^2)V]W^2\\
& + [c + c_2U + c_4U^2 + (D + d_2U + d_4U^2)V]Z^2 - 2uv(e_0 + e_1U)WZ,
\end{aligned}$$

where e_0, e_1 are exactly the original coefficients in (WS3). Set

$$\begin{gathered}
\beta_s = \frac{Bd_4 + \kappa c_4}{\kappa\ell^2}, \qquad \gamma_s = b_4 + \frac{d_4}{\kappa\ell},\\
j_0 = \frac{Bd_4 - a_4\kappa^2\ell + d_2\ell}{2\kappa\ell^2}, \qquad j_1 = \frac{d_4}{2\kappa\ell}, \qquad J_s(U) = j_0 + j_1U,\\
f_s = \frac{1}{2}\left(b_2 - \frac{c_4}{\ell^2} - \frac{c}{B^2} - \frac{Bd_4}{\kappa\ell^2}\right).
\end{gathered} \tag{WA4}$$

The four original scalar contact equations (value and the three logarithmic derivatives in $u, v, W/Z$) give the exact identity

$$F_g/4 = (R_1, R_2, T)\,\mathsf{S}\,(R_1, R_2, T)^T, \qquad \mathsf{S} = \begin{pmatrix} a + a_4U & uvJ_s & -vJ_s\\ uvJ_s & m + \beta_sU & u(f_s - j_1U)\\ -vJ_s & u(f_s - j_1U) & c/B^2 + \gamma_sU \end{pmatrix}. \tag{WA5}$$

This is a different scalar identity from the excluded polynomial two-block splitter (JG7). In particular, the last diagonal retains the full radial mass α/B^2 and the Gram depends on the independent testing variables. All source coefficients in (WA4) are bounded; no uncancelled $1/B$ occurs.

Both full first normals and the complete hard block. For the ξ normal define

$$\begin{gathered} p_0 = \frac{D + r(n+1/2)}{\kappa}, \qquad p_1 = \frac{r(n+D+1/2)}{\kappa}, \qquad q_1 = \frac{nr + Bp_1}{\ell}, \\ f_x = -mh + (D+\tfrac{1}{2})(1-h) - p_0 - q_1, \\ \mathsf{L}_x = \big(uv(p_0 + p_1 U),\ -m + q_1 U,\ u(f_x - p_1 U)\big). \end{gathered} \tag{WA6}$$

For the unnormalized $\eta = Y - huW$ normal set

$$\begin{gathered} p_y = -(C + D + \tfrac{1}{2})(1-h) + Ah, \qquad p'_y = Ah, \\ k_v = -C(1-h) + Ah + m + (\alpha + \tfrac{1}{2})h, \qquad k_{uv} = h(A + m + \alpha + \tfrac{1}{2}), \\ k_z = -m(1-r) + r(D + \tfrac{1}{2}) - \alpha - \tfrac{1}{2}, \\ q_y = \frac{-k_z - \kappa p'_y - Bk_{uv}}{\ell}, \qquad f_y = k_v - q_y, \\ \mathsf{L}_y = (p_y + p'_y U,\ uvq_y,\ v(f_y + k_{uv} U)). \end{gathered} \tag{WA7}$$

The two remaining original source stationarity equations, in X, Y, give these exact representations of the complete first normals. No first-normal term is discarded. Define also

$$\begin{aligned} h_{xx} &= n(1+U)(1+V) + m + (D+\tfrac{1}{2})UV + \tfrac{1}{2}(1+V), \\ h_{xy} &= -uv(m + D + \tfrac{1}{2}), \\ h_{yy} &= A(1+U)(1+V) + C(1+V) + D + \tfrac{1}{2} + mUV + (\alpha + \tfrac{1}{2})(1+U)V. \end{aligned}$$

Then the *entire original four-state parent* is exactly

$$\begin{gathered} F/4 = \mathcal{R}^T \mathsf{G}(u,v)\mathcal{R}, \qquad \mathcal{R} = (\xi, q, R_1, R_2, T)^T, \\ \mathsf{G} = \begin{pmatrix} h_{xx} & h_{xy}/\sqrt{A} & \mathsf{L}_x \\ h_{xy}/\sqrt{A} & h_{yy}/A & \mathsf{L}_y/\sqrt{A} \\ \mathsf{L}_x^T & \mathsf{L}_y^T/\sqrt{A} & \mathsf{S} \end{pmatrix}. \end{gathered} \tag{WA8}$$

Block notation in (WA8) has sizes $1, 1, 3$. Every coefficient of this finite polynomial matrix in u, v is bounded and pure source. The checker first expands the original signed core and all six original event packets to verify the complete hard block and both normal rows. It then verifies (WA5)–(WA7) from the corresponding four plus two contact equations. Thus (WA8) is not merely a graph or Hessian certificate.

Strict residue and a legal separated Gram lift. For infinitesimal independent u, v, the residue of (WA8), in the printed root order, is

$$\overline{\mathsf{G}} = \begin{pmatrix} \bar{n} + \bar{m} + 1/2 & 0 & 0 & -\bar{m} & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & \bar{C} + \bar{D} + 1/2 & 0 & 0 \\ -\bar{m} & 0 & 0 & \bar{m} & 0 \\ 0 & 0 & 0 & 0 & \overline{\alpha/B^2} \end{pmatrix} \succ 0. \tag{WA9}$$

Indeed the nontrivial 2×2 determinant is $\bar{m}(\bar{n} + 1/2) > 0$, and (WA2) supplies the last positive pivot. The q couplings disappear because $A^{-1/2}$ is infinitesimal while Ah is bounded. Every other omitted entry has a factor u, v, or uv.

For completeness, strict bounded residue here licenses a finite tensor SOS without a mixed Cholesky or inverse. Let $\mathsf{G}_0 = \mathsf{G}(0,0)$, a positive definite pure-source matrix whose inverse is bounded. Whiten by its pure-source square root and write the resulting matrix as

$$I + \sum_{j=1}^{N} H_j t_j,$$

where each H_j is a bounded pure-source symmetric matrix and each t_j is a nonconstant pure-testing monomial in u, v, hence infinitesimal. Split I into N equal copies and orthogonally diagonalize each H_j purely in the source field. A resulting scalar multiplier is $1/N + \ell t_j$, with ℓ bounded. Choose a positive real $R > N|\ell|$ and put $a = N\ell/R$, $b = Rt_j$. Then $|a| < 1$, $|b| < 1$, and

$$\frac{1}{N} + \ell t_j = \frac{1}{2N}\{(1+a)(1+b) + (1-a)(1-b)\}. \tag{WA10}$$

Each product is the square of a pure-source square root times a pure-testing square root. This supplies finitely many legal tensor roots for (WA8). Only finitely many source eigenvalues occur, so one real R can be chosen for all of them. If $N = 0$ the source Gram itself is already sufficient.

This construction works for either sign and for zero u or v, and permits all relative source/testing rates. Applying the real identity separately to the real and imaginary state parts proves the complex-state assertion. It uses pure-testing square roots of $1 \pm Rt_j$, and consequently makes no claim to a global polynomial SOS in the testing variables. There is therefore no conflict with the obstruction proved on the physical SO1 family.

Exact scope of the gain. The result closes the local both-infinitesimal testing problem on (WA1), with all four original state coordinates retained and with no centered-packet subtraction. It does not yet cover the allowed source openings z/B infinitesimal or unbounded, or $1 - r$ infinitesimal: in the latter case the R_2 pivot in (WA9) can lose residue. A remote-testing theorem must still be joined before this local certificate is called an all-tests source admission. No assertion of nonemptiness of the final residual intersection is made. The accompanying checker verifies thirteen exact identities; the source bounds and finite lift above are the positivity proof.

Chapter 55

A direct all-tests join for the five-root A-unbounded subchart

Component `p0d4eedd6c7a4`
Source: `R13/advances/weak_A/five_root_remote.tex`
SHA256: `0d4eedd6c7a4226bd797b288d93901cec2bcdae6b3ffe11bc5e98a60609914b4`
Inventory use: WR1-WR7

55.1 A direct all-tests join for the five-root A-unbounded subchart

Revision 10 scope update. Section 56.1 extends this admission to z/B positive infinitesimal with all other source hypotheses retained. The unit-ratio restriction below records the original revision-9 theorem.

This section supplies a new direct join for exactly the source hypotheses (WA1). It does not import the strong-II remote theorem or any other weak-C2 remote admission. All source physicality, strict-gap and contact hypotheses from the local theorem remain in force. No membership in the residual complement is needed; applications to the original residual problem retain that intersection without asserting that it is nonempty.

Theorem 55.1. *Under the source hypotheses of the five-root local theorem, including the positive-unit assumptions on z/B and $1-r$, the entire original parent is tensor SOS for every independent projective testing angle pair and every real or complex projective state. Literal zero and infinite half-angles and all signed charts are included. This is an all-tests source-subdomain admission; it does not assert nonemptiness of its intersection with the previous residual complement.*

The limiting source is explicit and all four finite weights are strict. Put

$$\rho = \operatorname{st}(z/B) > 0, \qquad \tau = \operatorname{st}(y/B) > 0, \qquad r_0 = \operatorname{st}(r) \in (0,1).$$

The original unit gaps give $g_0 = 1-\rho(1-r_0) > 0$ and $j_0 = 1-1/\tau > 0$, hence $\tau > 1$. Exact reduction of the unchanged source formulas gives

$$\begin{aligned} n_0 &= \frac{1-r_0}{2r_0(1+\rho)}, & m_0 &= \frac{\rho^2(1-r_0)}{2g_0(1+\rho)}, \\ C_0 &= \frac{\tau(1+\rho r_0)-\rho(1-r_0)}{2(1+\rho)}, & D_0 &= \frac{1+\rho r_0}{2(1+\rho)(\tau-1)}. \end{aligned} \tag{WR1}$$

Every displayed coefficient is strictly positive. For C_0 the numerator equals $\tau g_0 + \rho(\tau-1) + \rho r_0 > 0$. In particular n is a positive bounded unit on this subchart; the stronger fact is used at one remote endpoint. The formulas are joint regular reductions under $B \to 0$, $H/B \to 0$, with the other source ratios tending to their residues; they do not compare the two infinitesimal rates. Also $\alpha \in \mathrm{Inf}$ and $A^{-1/2} \in \mathrm{Inf}$.

Normalize the hard state by $\widehat{Y} = \sqrt{A}\,Y$ in each remote chart. This is a pure-source state change, and no parent term is removed. On bounded testing charts its matrix has bounded pure-source polynomial coefficients; all couplings of $\widehat{Y}$ to W, X, Z have zero residue. Its hard diagonal residue is the positive angle factor multiplying AY^2. It remains to check the complete three-state residue on $Y = 0$.

Finite angles outside the local corner. For finite real angle residues u_0, v_0, write $U = u_0^2$, $V = v_0^2$, $D_u = 1 + U$, $D_v = 1 + V$. Let P_3 be the matrix of the following complete residue of $F/4$ on $Y = 0$:

$$\begin{aligned} F_3 = {} & n_0 D_u D_v X^2 + m_0(v_0 W - X - u_0 Z)^2 + C_0 D_v U W^2 \\ & + (D_0 + \tfrac{1}{2})(u_0 W + u_0 v_0 X - v_0 Z)^2 + \tfrac{1}{2} D_v (X + u_0 Z)^2 - \tfrac{1}{2} D_u V Z^2. \end{aligned} \tag{WR2}$$

No hard-state relaxation term remains in this residue, because the normalized $\widehat{Y}$ cross row is infinitesimal. Define the strictly positive real constants

$$a_* = \frac{D_0 m_0 (2n_0 + 1)}{2}, \quad c_* = \frac{C_0 (2m_0 + 1)(2D_0 + 2n_0 + 1)}{4}, \quad d_* = \frac{C_0 D_0 (2n_0 + 1)}{2}.$$

The exact residue determinant is

$$\det \mathsf{P}_3 = D_u D_v \left\{ a_* \left(V - \frac{\tau}{\rho} U \right)^2 + c_* U^2 V + d_* U V^2 \right\}. \tag{WR3}$$

The checker obtains this directly from the full three-state polynomial and (WR1), not from a numerical table. It is strictly positive whenever $U + V > 0$, including either axis. The restriction of (WR2) to $Z = 0$ is positive definite on (W, X): its positive n_0, m_0, C_0 event squares already have no common nonzero kernel when $U + V > 0$, and the core restriction is nonnegative. The real Schur criterion and (WR3) therefore give $\mathsf{P}_3 \succ 0$. Together with the hard $\widehat{Y}$ pivot, the entire four-state residue is strictly positive definite.

For independent bounded testing angles with these residues, expand the normalized full matrix around the real pair (u_0, v_0). Its nonconstant monomials in $u - u_0, v - v_0$ are pure-testing infinitesimals. The finite separated Gram construction (WA10) applies verbatim. If $u_0 = v_0 = 0$, use the local five-root theorem instead. Thus every bounded testing pair is covered.

One infinite angle. For u unbounded use $p = 1/u$ and the full homogeneous parent $p^2 F(1/p, v)$; at literal $u = \infty$ use its polynomial value at $p = 0$. If v is bounded, the complete three-state residue on $Y = 0$ is

$$C_0(1 + v_0^2) W^2 + n_0(1 + v_0^2) X^2 + (D_0 + \tfrac{1}{2})(W + v_0 X)^2 + (m_0 + \tfrac{1}{2}) Z^2. \tag{WR4}$$

It is positive definite for every finite real v_0, including $v_0 = 0$. The hard $\widehat{Y}$ pivot is $1 + v_0^2 > 0$. For v unbounded and u bounded, put $q = 1/v$ and use the polynomial parent $q^2 F(u, 1/q)$. Its complete three-state residue is

$$(m_0 + C_0 u_0^2) W^2 + (n_0 + \tfrac{1}{2})(1 + u_0^2) X^2 + D_0 (u_0 X - Z)^2, \tag{WR5}$$

again positive definite, with hard pivot $1 + u_0^2$. Both identities are verified directly against the original parent. The same bounded separated Gram lift applies in the infinitesimal variables $p, v - v_0$ or $u - u_0, q$. This includes literal infinity and literal zero of the other half-angle.

Both infinite angles: keep the exact null state. Put $p = 1/u$, $q = 1/v$, and work with the full polynomial parent $\widehat{F} = p^2 q^2 F(1/p, 1/q)$. Introduce the five roots

$$(W, X, \widehat{Y}, P, Q), \qquad \widehat{Y} = \sqrt{A}\,Y, \quad P = pZ, \quad Q = qZ.$$

Write $Y = A^{-1/2}\widehat{Y}$ in the following exact quadratic identity, interpreted as a polynomial in these five roots:

$$\begin{aligned}\widehat{F}/4 = {} & n(1+p^2)(1+q^2)X^2 + m(pW - pqX + Y - Q)^2 + C(1+q^2)(W - pY)^2 \\ & + (1+p^2)(1+q^2)\widehat{Y}^2 + D(qW + X - pqY - P)^2 + \alpha(1+p^2)(Y-Q)^2 \\ & + \tfrac{1}{2}\Big\{(qW + X - pqY - P)^2 + (1+p^2)(Y-Q)^2 + p^2(1+q^2)X^2 \\ & \qquad +2(1+q^2)XP - (1+q^2)P^2\Big\}.\end{aligned} \tag{WR6}$$

The checker verifies this as the complete original four-state parent after $P = pZ$, $Q = qZ$, not only to leading order. All matrix coefficients in (WR6) are bounded source polynomials in p, q. Its residue, in the printed root order, is

$$\begin{pmatrix} C_0 & 0 & 0 & 0 & 0 \\ 0 & n_0 + D_0 + 1/2 & 0 & -D_0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & -D_0 & 0 & D_0 & 0 \\ 0 & 0 & 0 & 0 & m_0 + 1/2 \end{pmatrix} \succ 0, \tag{WR7}$$

because the nontrivial determinant is $D_0(n_0 + 1/2) > 0$. The separated lift (WA10) therefore gives a finite tensor SOS for arbitrary independently infinitesimal p, q, of either sign or zero. At $p = q = 0$ both P, Q vanish, respecting the original exact null direction $W = X = Y = 0$; no positive Z pivot is invented.

Coverage and restoration. Every finite element of an ordered extension is either bounded or unbounded in absolute value. A bounded testing pair is covered by (WA1)–(WA10) if both residues vanish, and by (WR2)–(WR3) otherwise. A single unbounded coordinate is covered by (WR4) or (WR5); two unbounded coordinates are covered by (WR6)–(WR7). Literal projective endpoints use the polynomial chart values with $p = 0$ or $q = 0$. For nonzero reciprocal coordinates, restoring F multiplies the established identity by the pure-testing square p^{-2}, q^{-2}, or $(pq)^{-2}$; at literal infinity no division is performed. All four homogeneous state coordinates were retained in every chart, and splitting real and imaginary state parts gives the complex-state statement. Thus the join exhausts all independent testing types without a source/testing rate assumption or an imported strong-face theorem.

Corollary 55.2 (The polynomial obstruction family is nevertheless admitted)**.** *The entire two-parameter physical family (SO1),*

$$B = \varepsilon, \quad z = \varepsilon/2, \quad y = 2\varepsilon, \quad r = 9/10, \quad H = \varepsilon\eta, \quad L = \varepsilon(1-\eta),$$

with independent positive infinitesimals ε, η, is all-tests tensor SOS for the original full parent.

Proof. It has $z/B = 1/2$, $1 - r = 1/10$, $H/y = \eta/2$ infinitesimal, and $y/B, L/B, r, G/B, j$ positive bounded units. Its global physicality and all original source signs were proved in the obstruction module. It therefore satisfies every hypothesis of the present all-tests theorem. □

The strict $K_1 < 0$ and (JG13) failures and the nonexistence of a global polynomial SOS of its scalar graph remain correct. The local tensor roots use pure-testing radicals as in (WA10), so these facts are compatible. The whole family is now a proved admission and is removed from the residual complement. No claim that any other residual intersection is nonempty is implied.

Chapter 56

The small-ratio weak-C2 opening is also admitted for all tests

Component p296e824cb97a
Source: R13/advances/weak_A_boundary/extension.tex
SHA256: 296e824cb97a0923083aae8ef939c13988961e23d165a42f581ecd58ab6690b4
Inventory use: WE1-WE11

56.1 The small-ratio weak-C2 opening is also admitted for all tests

Theorem 56.1. *Retain the original physical C2 source, its exact stationary contact, all strict source gaps, $n \geq 0$, and the normalization $d = 1$. Suppose $B, z, y > 0$ are infinitesimal, $r, 1 - r, G/B, j, y/B, L/B$ are positive bounded units, and H/y is infinitesimal. If $a = z/B$ is positive infinitesimal, the entire original parent is a finite tensor SOS at every independent projective testing pair, for all real or complex states, including signed charts and literal zero and infinite half-angles. No comparison between source infinitesimal rates or between source and testing rates is imposed. In particular $\delta_{\mathrm{tot}} > 0$ is infinitesimal. This admission does not assert nonemptiness of its intersection with the previous residual complement.*

The original omitted event coefficients remain $\lambda = \beta = 0$. Here $B = H + L$, $h = H/B$, $t = y/B$, and the original contact is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = (\sqrt{z}, \sqrt{y}; B/\sqrt{zy}, -r\sqrt{z}, H/\sqrt{y}, 1)\,.$$

No residual-complement premise is used. The proof extends the exact six-contact full-parent Gram (WA3)–(WA8) and its finite separated lift (WA10); it does not use the disproved global polynomial splitter.

Exact cancellation, with no h/a restriction. Put $\Delta = \delta/B$, $M_* = M/B^2$, $\mu = m/a^2$ and $\alpha_* = \alpha/B^2$. The unchanged source formulas give

$$\begin{aligned}
c_a &= 1 + aBr, \quad S = 1 + B(1 - h + ar),\\
g = G/B &= 1 + a(hB + r - 1), \quad j = c_a - (1 - h)/t,\\
\Delta &= a + (1 + aB) + (1 + aB)B + 2aB(1 + aB)r\\
&\quad + a^3B^2r^2 + 2a^2Br + tB[1 + a + aB + a^2Br(2 - r)]'\\
M_* &= (1 + aB)(1 + ar)(h + t) - a(1 + tB)(1 - r)(1 - h),\\
n &= \frac{(1 + ar)c_aS - r\Delta}{2r\Delta}, \qquad \mu = \frac{(1 + tB)(1 - r)c_aS}{2g\Delta},\\
C &= \frac{c_aM_*}{2(1 - h)\Delta}, \quad Ah = \frac{tc_a(1 + ar)S}{2\Delta},\\
D &= \frac{(1 + aB)(1 + ar)(1 - atBr)S}{2tj\Delta}, \quad \alpha_* = \frac{(1 + ar)M_*}{2(1 - hB)\Delta}.
\end{aligned} \tag{WE1}$$

All displayed denominators are units at the joint corner $B = a = h = 0$; $j > 0$ a unit implies $t_0 = \mathrm{st}(t) > 1$. The subscript 0 denotes residue, and $0 < r_0 < 1$. Thus

$$n_0 = \frac{1 - r_0}{2r_0}, \quad \mu_0 = \frac{1 - r_0}{2}, \quad C_0 = \frac{t_0}{2}, \quad D_0 = \frac{1}{2(t_0 - 1)}, \quad (Ah)_0 = (\alpha_*)_0 = \frac{t_0}{2}. \tag{WE2}$$

These are positive; $m = a^2\mu$ is infinitesimal, A is unbounded, and $Ah^2, \sqrt{A}h, A^{-1/2}$ are infinitesimal. All other retained coefficients are bounded. Consequently $\delta_{\mathrm{tot}} = 1/(1 + n + m + C + A + D + \alpha)$ is positive infinitesimal, as required in the original source normalization.

The only new potentially unbounded local coefficient is $(b_2 - \alpha_*)/a$, where

$$b_2 = 2a^2\mu h + C(1 - h)^2 + (Ah)h + (B^2\alpha_* + 1/2)h^2.$$

At $a = 0$, with $d_* = 1 + B(1 + t)$, the exact values are

$$C = \frac{h + t}{2(1 - h)d_*}, \quad Ah = \frac{t[1 + B(1 - h)]}{2d_*}, \quad \alpha_* = \frac{h + t}{2(1 - Bh)d_*}.$$

Substitution gives $b_2 = \alpha_*$ identically for arbitrary B, h, t. Clearing the unit denominators in (WE1), the numerator of $b_2 - \alpha_*$ is therefore divisible by a in the polynomial ring. Its quotient has a unit denominator at $B = a = h = 0$. Hence

$$Q_* = (b_2 - \alpha_*)/a \quad \text{is bounded.} \tag{WE3}$$

This is a joint regularity argument, so it permits arbitrary h/a, including an unbounded ratio. The checker verifies the exact vanishing, polynomial divisibility, and nonzero denominator at the joint corner.

The complete local five-root Gram survives. Keep $\kappa = B/y$ and every coefficient and normal in (WA3)–(WA8), but set $\ell = 1/a$ and replace R_2 by

$$\widetilde{R}_2 = aR_2 = avW - uZ.$$

The entire full-parent matrix is the exact congruence

$$\widetilde{G} = EGE, \qquad E = \mathrm{diag}(1, 1, 1, a^{-1}, 1), \qquad \widetilde{\mathcal{R}} = (\xi, q, R_1, \widetilde{R}_2, T). \tag{WE4}$$

Its newly rescaled coefficients are

$$\begin{aligned}
m/a^2 &= \mu, \qquad \beta_s/a^2 = (Bd_4 + \kappa c_4)/\kappa,\\
J_s(U)/a &= \frac{aBd_4 - a_4\kappa^2 + d_2 + d_4U}{2\kappa}, \qquad f_s/a = \tfrac{1}{2}[Q_* - ac_4 - aBd_4/\kappa],\\
(-m + q_1U)/a &= -a\mu + (nr + Bp_1)U,\\
q_y/a &= -k_z - \kappa p'_y - Bk_{uv}.
\end{aligned} \tag{WE5}$$

All are bounded, by (WE1)–(WE3). The other coefficients are bounded as before; the complete second normal remains divided by $\sqrt{A}$. Thus no first normal or hard-state term is omitted. For independent infinitesimal or zero u, v the complete residue is

$$\overline{\widetilde{G}} = \operatorname{diag}(n_0 + 1/2, 1, C_0 + D_0 + 1/2, \mu_0, t_0/2) \succ 0. \tag{WE6}$$

In particular the previous ξ, R_2 cross entry becomes $-a\mu$ and has zero residue. Applying the pure-source and pure-testing finite lift (WA10) proves the full local statement.

Bounded tests with nonzero u residue. Normalize $\widehat{Y} = \sqrt{A}Y$ without centering and use the complete three-state residue (WR2), now with $m_0 = 0$ and (WE2). For $U = u_0^2$, $V = v_0^2$, its exact determinant is

$$\det P_3 = (1+U)(1+V)U[b_*U + c_*UV + d_*V^2], \tag{WE7}$$

where

$$b_* = \frac{n_0(2C_0 + 2D_0 + 1)}{4}, \quad c_* = \frac{C_0(2D_0 + 2n_0 + 1)}{4}, \quad d_* = \frac{C_0 D_0(2n_0 + 1)}{2}.$$

Every coefficient is positive. For $U > 0$, the restriction to $Z = 0$ is positive definite from the positive n_0, C_0 packets. The determinant therefore gives a positive definite complete residue, with the additional hard $\widehat{Y}$ pivot $(1+U)(1+V)$. The finite separated lift around (u_0, v_0) applies.

The missing finite-v and reciprocal-v charts. For u infinitesimal or zero and v a bounded unit, use the five roots

$$(S_*, R_*, X, \widehat{Y}, Z) = (aW, uW, X, \sqrt{A}Y, Z), \qquad Y = A^{-1/2}\widehat{Y}.$$

Writing $D_u = 1 + u^2$, $D_v = 1 + v^2$, the following is the entire original $F/4$, as a quadratic polynomial in those roots:

$$\begin{aligned} &nD_uD_vX^2 + \mu(vS_* - aX + auvY - auZ)^2 + CD_v(R_* - Y)^2 + D_uD_v\widehat{Y}^2 \\ &\quad + D(R_* + uvX - Y - vZ)^2 + \alpha D_u(vY - Z)^2 \\ &\quad + \tfrac{1}{2}\{(R_* + uvX - Y - vZ)^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_uD_vZ^2\}. \end{aligned} \tag{WE8}$$

The coefficients are polynomials in u, v. Their pure-source factors $a, \mu, n, C, D, \alpha, A^{-1/2}$ are bounded. At $u_0 = 0$, $v_0 \neq 0$, the separate pivots on $(S_*, X, \widehat{Y})$ are $\mu_0 v_0^2, (n_0 + 1/2)(1 + v_0^2), 1 + v_0^2$. The remaining block on (R_*, Z) is

$$\begin{pmatrix} C_0(1+v_0^2) + D_0 + 1/2 & -(D_0 + 1/2)v_0 \\ -(D_0 + 1/2)v_0 & D_0 v_0^2 \end{pmatrix}, \qquad \det = C_0 D_0 v_0^4 > 0. \tag{WE9}$$

Here $C_0D_0 - D_0/2 - 1/4 = 0$ exactly. Thus (WA10) applies in $u, v - v_0$, including literal $u = 0$; roots need not be independent for a positive definite Gram to give the asserted SOS.

For v unbounded or literally infinite, put $q_v = 1/v$ and use the polynomial homogeneous parent $q_v^2 F(u, 1/q_v)/4$. In the same five roots it is exactly

$$\begin{aligned} &nD_u(1+q_v^2)X^2 + \mu(S_* - aq_vX + auY - auq_vZ)^2 + C(1+q_v^2)(R_* - Y)^2 + D_u(1+q_v^2)\widehat{Y}^2 \\ &\quad + D(q_vR_* + uX - q_vY - Z)^2 + \alpha D_u(Y - q_vZ)^2 \\ &\quad + \tfrac{1}{2}\{(q_vR_* + uX - q_vY - Z)^2 + D_u(Y - q_vZ)^2 \\ &\qquad + (1+q_v^2)(X + uZ)^2 - D_u(1+q_v^2)Z^2\}. \end{aligned} \tag{WE10}$$

For independent infinitesimal or zero u, q_v its full residue is diagonal with pivots

$$(\mu_0, C_0, n_0 + 1/2, 1, D_0) > 0. \tag{WE11}$$

Its coefficients are bounded for the same reason as (WE8). The separated lift again applies, including $q_v = 0$ literally. The checker expands (WE8) and (WE10) back to all six original event packets and the complete signed core, and checks both residues.

Remaining endpoints and exhaustive coverage. If u is unbounded or infinite and v bounded, (WR4) is still positive definite with $m_0 = 0$: its positive coefficients are $C_0, n_0, D_0 + 1/2, 1/2$, and the hard pivot is $1 + v_0^2$. If v is unbounded or infinite and u is a bounded unit, (WR5) is positive definite because $C_0 u_0^2 > 0$, $n_0 + 1/2 > 0$, and $D_0 > 0$, with hard pivot $1 + u_0^2$. If both are unbounded or infinite, use the full homogeneous five-root identity (WR6). Its residue (WR7) is still strictly positive: the Q pivot becomes $1/2$, and the other nontrivial block determinant remains $D_0(n_0 + 1/2) > 0$. All these identities have bounded coefficients also at $m_0 = 0$.

For two bounded tests, both zero residues use (WE4)–(WE6); nonzero u residue uses (WE7); zero u and nonzero v residue uses (WE8)–(WE9). A single unbounded coordinate is covered by the preceding paragraph, except for unbounded v and zero u residue, which is (WE10)–(WE11). Two unbounded coordinates use (WR6)–(WR7). These cases exhaust every independent testing type. Literal projective endpoints use the exact polynomial chart value. Away from literal endpoints, restoration multiplies by a pure-testing square p^{-2}, q_v^{-2}, or $(pq_v)^{-2}$. No division is made at a literal endpoint. All four original homogeneous state coordinates are retained; real and imaginary parts give the complex-state result. The proof establishes the full-parent all-tests admission on this source opening, without a scalar-to-parent inference or a new physicality assumption.

The unit-ratio assumption can now be removed. Under the retained hypotheses, $H \in \mathrm{Inf}$ and $1 - r$ is a positive unit. Since $g = 1 - a(1 - r - H) > 0$, one has $0 < a < (1 - r - H)^{-1}$, so a is bounded. Every positive such a is either a unit, covered by (WA1), or infinitesimal, covered here. Their union therefore gives the same all-tests theorem without the additional z/B-unit assumption. This argument does not cover a vanishing $1 - r$ or any other omitted source gap.

Chapter 57

The small-gap weak-C2 source reduces to two infinitesimal tests

Component pbb649976abc6
Source: R13/advances/weak_small_gap_remote/remote_reduction.tex
SHA256: bb649976abc6626a60a6f4d950e998d5622f9596c136b333213ba13b2ab35af8
Inventory use: GR1-GR7

57.1 The small-gap weak-C2 source reduces to two infinitesimal tests

Theorem 57.1. *Under the source hypotheses of Section 58.1, the entire original parent is finite tensor SOS at every independent projective testing pair except possibly when both u, v are bounded with zero residue. All signs, literal projective endpoints, real and complex states, and literal $n = 0$ are included. This is a remote-testing reduction, not an all-tests source admission.*

Write $\rho = \mathrm{st}(z/B) > 0$ and $\tau = \mathrm{st}(y/B) > 1$. The exact source formulas give

$$n, m, \alpha \in \mathrm{Inf}_{\geq 0}, \quad A^{-1/2} \in \mathrm{Inf}_{>0}, \qquad C_0 = \tau/2, \quad D_0 = 1/[2(\tau - 1)]. \tag{GR1}$$

The source-scale lemma retains all nonnegative openings, including $n = 0$. Normalize only $\widehat{Y} = \sqrt{A}Y$ and put $Y = A^{-1/2}\widehat{Y}$ in every full-parent identity below. All remaining event coefficients are bounded. The strict-residue lift is (WA10), expanded around the relevant real testing residues; no mixed inverse or radical is used.

A six-root identity removes the second axis degeneracy. If u is a bounded unit or unbounded, use $p = 1/u$, which is bounded, and the polynomial homogeneous parent $F_p = p^2 F(1/p, v)$. Literal $u = \infty$ means $p = 0$. Suppose v is infinitesimal or zero. Put $D_p = 1 + p^2$, $D_v = 1 + v^2$, and use the six roots

$$N = \sqrt{n}X, \quad V_x = vX, \quad K = pX + Z, \quad R = \sqrt{\alpha}\, pX, \quad W, \quad \widehat{Y}.$$

The roots with zero source weight are simply zero; no division by n or α occurs. Define $J = W + D_p V_x - pY - pvK$. There is the exact complete-parent identity

$$\begin{aligned} F_p/4 = {} & D_p D_v N^2 + m(pvW + vY - K)^2 + CD_v(W - pY)^2 + D_p D_v \widehat{Y}^2 \\ & + DJ^2 + D_p[R + \sqrt{\alpha}(vY - K)]^2 \\ & + \tfrac{1}{2}\{J^2 + (1 - p^2v^2)K^2 \\ & \qquad + D_p[v^2Y^2 - 2vYK + 2pYV_x + 2pvKV_x - p^2V_x^2]\}. \end{aligned} \tag{GR2}$$

This identity follows by $Z = K - pX$ and $vX = V_x$ in the original signed core and all original event packets. In particular the m packet loses its apparent separate pX term exactly. The α packet is

retained in the extra root R rather than discarded. All coefficients in (GR2) are bounded pure-source polynomials in bounded p, v after the normalization of Y.

At $p_0 = \mathrm{st}(p)$, $v_0 = 0$, write $d_p = 1 + p_0^2$. The full six-root residue is

$$d_p(N^2 + R^2 + \widehat{Y}^2) + \tfrac{1}{2}K^2 + C_0W^2 + (D_0 + \tfrac{1}{2})(W + d_pV_x)^2 - \tfrac{1}{2}d_pp_0^2V_x^2. \tag{GR3}$$

Its last two-state block has positive first diagonal and determinant

$$d_pC_0(D_0 + \tfrac{1}{2}) > 0, \tag{GR4}$$

because $C_0(D_0 + 1/2) = (C_0 + D_0 + 1/2)/2$. Every other displayed pivot is strictly positive. Thus the full parent is covered for all bounded p and infinitesimal or zero v, including both signed finite p and literal $p = 0$.

The other axis retains the small m packet. For u infinitesimal or zero and v a bounded unit, use

$$(S, R_u, X, \widehat{Y}, Z) = (\sqrt{m}W, uW, X, \sqrt{A}Y, Z).$$

The entire identity (WE8) applies with its m packet written $(vS - \sqrt{m}X + \sqrt{m}\,uvY - \sqrt{m}\,uZ)^2$. Every other packet is unchanged. The complete residue has separate pivots $v_0^2, (1 + v_0^2)/2, 1 + v_0^2$ on $(S, X, \widehat{Y})$; its remaining (R_u, Z) block is

$$\begin{pmatrix} C_0(1 + v_0^2) + D_0 + 1/2 & -(D_0 + 1/2)v_0 \\ -(D_0 + 1/2)v_0 & D_0v_0^2 \end{pmatrix}, \qquad \det = C_0D_0v_0^4 > 0. \tag{GR5}$$

For v unbounded, put $q = 1/v$ and use the complete identity (WE10) with the same m-packet change. At $u = q = 0$ its five-root residue is diagonal with pivots

$$(1, C_0, 1/2, 1, D_0) > 0. \tag{GR6}$$

These bounded-coefficient identities also cover literal $u = 0$ and $q = 0$. All first normals and the full hard state block are retained.

The remaining remote charts are strict directly. For two bounded unit angles, the full three-state residue (WR2) with $n_0 = m_0 = 0$ has determinant, with $U = u_0^2$, $V = v_0^2$,

$$(1 + U)(1 + V)UV\left\{\frac{C_0(2D_0 + 1)}{4}U + \frac{C_0D_0}{2}V\right\} > 0. \tag{GR7}$$

Its restriction to $Z = 0$ is positive definite from the positive $C_0U(1 + V)W^2$ and $(1 + V)X^2/2$ terms. Including the normalized hard $\widehat{Y}$ pivot gives a strict full-parent Gram residue. For u unbounded and v a bounded unit, (WR4) has a strict (W, X) block from $C_0(1 + V)W^2 + (D_0 + 1/2)(W + v_0X)^2$, a Z pivot $1/2$, and a hard pivot $1 + V$. For v unbounded and u a bounded unit, (WR5) is strict: its W pivot is $C_0U > 0$ and its (X, Z) block is $(1 + U)X^2/2 + D_0(u_0X - Z)^2$. For both angles unbounded, the exact five-root identity (WR6) has residue (WR7) with $n_0 = m_0 = 0$. Its nontrivial block determinant is $D_0/2 > 0$, and the remaining pivots are $C_0, 1, 1/2$. The coefficients in each identity are still bounded under (GR1), so this use of the algebra does not import an incompatible source-domain admission.

These charts exhaust every test pair except both bounded with zero residue. Literal infinity uses polynomial chart values; restoration elsewhere uses only pure-testing squares. Splitting real and imaginary state parts gives the complex assertion. On the printed source chart, the exact remaining task is therefore the entire local parent with u, v independently infinitesimal or zero, retaining the physical constraint $(1 - r)Q \geq BK$ and every equality case. No comparison between a source scale and a testing scale has been introduced.

Chapter 58

The physical small-$1-r$ source has a forced scale bound

Component `p0c68f3aecc59`
Source: `R13/advances/weak_small_gap_scale/boundary_scale_lemma.tex`
SHA256: `0c68f3aecc59fba087d199c915d7c464be358b028e41022ad9704248c96e6860`
Inventory use: BS1-BS5

58.1 The physical small-$1-r$ source has a forced scale bound

Retain the original physical C2 source, its exact contact and strict gaps, with $n \geq 0$, $d = 1$, $B = H + L$, and $B, z, y > 0$ infinitesimal. Put

$$a = z/B, \quad t = y/B, \quad h = H/B, \quad e = 1 - r > 0.$$

Assume e, h infinitesimal and $a, t, G/B, j, L/B$ positive bounded units. Thus $\rho = \mathrm{st}(a) > 0$, $\tau = \mathrm{st}(t) > 1$, and r is a unit.

Lemma 58.1. *The exact original source identity is*

$$2r\Delta n = eQ - BK, \qquad \Delta = \delta/B, \tag{BS1}$$

where

$$\begin{aligned} K &= (1+a)(h+t)(1+aB), \\ Q &= 1 + B[1 + t + ah + 2ar + at] \\ &\quad + B^2\{a(h+t) + a^2[h(r+1) + t(1 + r - r^2)]\}. \end{aligned} \tag{BS2}$$

Both Q, K are positive bounded units, with residues $Q_0 = 1$, $K_0 = (1+\rho)\tau$, and $\Delta_0 = 1 + \rho$. Consequently source nonnegativity is exactly equivalent to

$$0 < B/e \leq Q/K, \qquad 0 \leq n/e \leq Q/(2r\Delta). \tag{BS3}$$

In particular B/e and n/e are bounded, regardless of the relative rate of h.

Proof. The original normalized source formula is

$$2r\Delta n = (1 + ar)(1 + aBr)[1 + B(1 - h + ar)] - r\Delta,$$

with

$$\begin{aligned} \Delta &= a + (1 + aB) + (1 + aB)B + 2aB(1 + aB)r + a^3B^2r^2 + 2a^2Br \\ &\quad + tB[1 + a + aB + a^2Br(2 - r)]. \end{aligned}$$

Subtracting its value at $r = 1$ and dividing by $1 - r$ gives the polynomial Q in (BS2); the value at $r = 1$ is $-BK$. Every term in Q is positive, since $0 < r < 1$ implies $1 + r - r^2 > 0$. The stated residues and inequalities follow. □

The literal $r = 1$ face is not an admissible anchor. At $r = 1$ and still $B > 0$, the exact identity becomes

$$n\big|_{r=1} = -\frac{B(h+t)(1+aB)}{2[1+B(1+2a+t)+aB^2(1+a+t)]} < 0. \tag{BS4}$$

Thus simply setting $r = 1$ exits the original nonnegative-source domain. An event-face theorem requiring $n \geq 0$ cannot be applied to that unmodified endpoint. This is a limitation of that endpoint reduction, not a refutation of any alternative full-parent proof.

What this repairs, and what remains unresolved. Let $\lambda_* = B/e$, which is bounded by (BS3). Its residue may be zero. Joint substitution $B = e\lambda_*$, $r = 1 - e$ gives

$$(n/e)_0 = \frac{1-\lambda_{*,0}(1+\rho)\tau}{2(1+\rho)}, \qquad (m/e)_0 = \frac{\rho^2}{2(1+\rho)} > 0. \tag{BS5}$$

For the scalar coefficient β_s in (WA4),

$$(\beta_s/e)_0 = \rho^2\left[\frac{1}{2(1+\rho)} + \frac{\lambda_{*,0}\tau}{2(\tau-1)}\right] > 0. \tag{BS6}$$

So source physicality removes the apparent diagonal-coefficient pole after scaling R_2 by $\sqrt{e}$. It does not remove the previously established cross-coefficient poles: the same exact source still forces

$$(j_0)_0 = \frac{\rho\tau(\tau+1)}{4(\tau-1)}, \quad (j_1)_0 = \frac{\rho\tau^2}{4(\tau-1)}, \quad (f_s)_0 = -\frac{\rho\tau}{4}.$$

Thus that fixed Gram still has unbounded $J_s/\sqrt{e}$ and $f_s/\sqrt{e}$. No new local full-parent certificate or all-tests admission is claimed here. The decisive result is the exact physical scale bound and the exclusion of the literal $r = 1$ nonnegative-source anchor. Source physicality and contact are retained as hypotheses throughout; no coefficient-sign argument is used to infer global physicality.

Chapter 59

One complete Gram for every bounded test on the small-r opening

Component p99372d145f17
Source: R13/advances/weak_small_r/local_small_r.tex
SHA256: 99372d145f17c3b661f5ddcadf1e9a3a21f60de2470ee6552a94cb44f19444d0
Inventory use: SL1-SL5

59.1 One complete Gram for every bounded test on the small-r opening

Theorem 59.1. *Retain the original physical C2 source, its exact stationary contact, all strict source gaps, and $d = 1$. Suppose $B, z, y, r > 0$ are infinitesimal; $a = z/B$ is positive bounded (a unit or infinitesimal); G/B, j, y/B, L/B are positive bounded units; and H/y is infinitesimal. Then the entire original four-state parent is a finite tensor SOS for arbitrary independent bounded testing half-angles u, v, of either sign, and for every real or complex state. There is no comparison of source or testing rates. The total-scale parameter δ_{tot} is positive infinitesimal. This covers all bounded tests; the projective endpoint join is a separate obligation. No residual-complement nonemptiness is claimed.*

Here $B = H + L$, $h = H/B$, $t = y/B$, $\kappa = 1/t$, $\ell = 1/a$; the omitted event coefficients remain $\lambda = \beta = 0$. The contact is the unchanged original contact

$$(\sqrt{z}, \sqrt{y}; B/\sqrt{zy}, -r\sqrt{z}, H/\sqrt{y}, 1).$$

No residual membership premise is used. All source physicality, contact and strict-gap hypotheses remain in force.

Both unbounded source coefficients can be retained. Use the exact normalized source formulas (WE1), whose denominators are regular at $B = h = r = 0$ after replacing n by $N = nr$. Explicitly, with $c_a = 1 + aBr$, $S = 1 + B(1 - h + ar)$ and $\Delta = \delta/B$,

$$N = \frac{(1 + ar)c_a S - r\Delta}{2\Delta}.$$

Let $\rho = \mathrm{st}(a) \in [0, 1)$ and $\tau = \mathrm{st}(t) > 1$. Direct joint reduction gives

$$\begin{aligned} N_0 &= \frac{1}{2(1+\rho)}, \qquad (m/a^2)_0 = \frac{1}{2(1-\rho^2)}, \qquad C_0 = \frac{\tau - \rho}{2(1+\rho)}, \\ D_0 &= \frac{1}{2(1+\rho)(\tau - 1)}, \qquad (Ah)_0 = \frac{\tau}{2(1+\rho)}, \qquad (\alpha/B^2)_0 = \frac{\tau - \rho}{2(1+\rho)} = C_0. \end{aligned} \tag{SL1}$$

Each displayed residue is strictly positive. Thus $n = N/r$ and $A = (Ah)/h$ are positive unbounded, with no comparison between them. In contrast nr is a unit and $nr^2 = Nr$ is infinitesimal; $\mu = m/a^2, C, D, Ah, \alpha/B^2$ are units, $m = a^2\mu$ is bounded, and α is infinitesimal. All normalized source denominators are units since $G/B \to 1-\rho > 0$, $j \to 1 - 1/\tau > 0$, and $\Delta \to 1+\rho > 0$. These are joint regular reductions, not iterated limits. In particular $\delta_{\mathrm{tot}} = 1/(1+n+m+C+A+D+\alpha) > 0$ is infinitesimal.

Exact congruence of the complete full-parent identity. Keep all source coefficients, the scalar block S, and both complete normal rows $\mathsf{L}_x, \mathsf{L}_y$ in (WA3)–(WA8), with $\ell = 1/a$. Normalize the first root and the second scalar root by pure-source factors:

$$\widehat{\xi} = \sqrt{n}(X + ruZ), \quad q = \sqrt{A}(Y - huW), \quad R_1 = uW - \kappa vZ, \quad \widetilde{R}_2 = avW - uZ, \quad T = uvW - BZ. \tag{SL2}$$

Every root still vanishes at the exact contact. If $E_s = \mathrm{diag}(1, a^{-1}, 1)$ and $\widehat{\mathcal{R}} = (\widehat{\xi}, q, R_1, \widetilde{R}_2, T)^T$, then the entire original parent has the exact identity

$$F/4 = \widehat{\mathcal{R}}^T \widehat{G} \widehat{\mathcal{R}}, \qquad \widehat{G} = \begin{pmatrix} h_{xx}/n & h_{xy}/\sqrt{nA} & \mathsf{L}_x E_s/\sqrt{n} \\ h_{xy}/\sqrt{nA} & h_{yy}/A & \mathsf{L}_y E_s/\sqrt{A} \\ E_s \mathsf{L}_x^T/\sqrt{n} & E_s \mathsf{L}_y^T/\sqrt{A} & E_s \mathsf{S} E_s \end{pmatrix}. \tag{SL3}$$

This is the pure-source congruence of (WA8) by $\mathrm{diag}(n^{-1/2}, 1, 1, a^{-1}, 1)$, so it retains the exact signed core, all event packets, both full normals and the entire hard block. It invokes the previously verified six original contact identities; no scalar-to-parent implication is used.

Here is a complete coefficient-bound check. In the scalar block, every occurrence of n is $nr^2 = Nr$. The exact rescaled formulas (WE5) show that $\beta_s/a^2, J_s/a, j_1/a$ are bounded (in fact infinitesimal). The only possible new pole is f_s/a. The exact source identity $b_2 - \alpha/B^2 = 0$ at $a = 0$ holds for arbitrary B, h, r, t and contains no inverse r. Its cleared numerator is divisible by a, and a canceled denominator reduces to $2(\rho-1)(\rho+1) \neq 0$ at $B = h = r = 0$, $0 \le \rho < 1$. The quotient numerator also vanishes identically at $B = h = r = 0$, for every a in this residue interval. Consequently $Q_* = (b_2 - \alpha/B^2)/a$ is infinitesimal uniformly, including joint r, a infinitesimal with arbitrary rates. Formula $f_s/a = [Q_* - ac_4 - aBd_4/\kappa]/2$ is therefore infinitesimal. All other rescaled scalar coefficients are bounded, with $m/a^2 = \mu$ a unit. In the first normal, the only potentially large terms occur through

$$p_0 = \frac{D + nr + r/2}{\kappa}, \quad p_1 = \frac{nr + r(D+1/2)}{\kappa}, \quad q_1 = \frac{nr + Bp_1}{\ell}.$$

They are bounded, as is $f_x = -mh + (D+1/2)(1-h) - p_0 - q_1$; therefore every coefficient of $\mathsf{L}_x E_s/\sqrt{n}$ is infinitesimal, since its middle entry before division by $\sqrt{n}$ is $(-m + q_1 U)/a = -a\mu + (nr + Bp_1)U$. The second normal depends only on bounded $m, C, D, \alpha, r, h, Ah$ and unit κ; its rescaled middle coefficient is bounded because $q_y/a = -k_z - \kappa p_y' - Bk_{uv}$. Thus all its coefficients become infinitesimal after division by $\sqrt{A}$. Finally, writing $U = u^2, V = v^2$, the complete hard entries are

$$\begin{aligned} h_{xx}/n &= (1+U)(1+V) + [m + (D+1/2)UV + (1+V)/2]/n, \\ h_{xy}/\sqrt{nA} &= -uv(m + D + 1/2)/\sqrt{nA}, \\ h_{yy}/A &= (1+U)(1+V) + [C(1+V) + D + 1/2 + mUV + (\alpha+1/2)(1+U)V]/A. \end{aligned} \tag{SL4}$$

No ratio n/A or A/n occurs. Thus $\widehat{G}$ is a finite polynomial matrix in u, v with bounded pure-source coefficients, uniformly for arbitrary independent source rates.

For arbitrary bounded testing u, v, put $u_0 = \mathrm{st}(u)$, $v_0 = \mathrm{st}(v)$, and $D_0^{\mathrm{test}} = (1+u_0^2)(1+v_0^2) > 0$. The scalar source coefficients $a_4, \beta_s/a^2, J_s/a, j_1/a, \gamma_s, f_s/a$ are all infinitesimal. Indeed $nr^2 = Nr \in \mathrm{Inf}$, $d_2, d_4, Ah^2 \in \mathrm{Inf}$, and $Q_* \in \mathrm{Inf}$ by the just-proved exact quotient identity. The complete normalized normal rows have infinitesimal coefficients. Consequently the entire Gram has the strict diagonal residue

$$\overline{\widehat{G}} = \mathrm{diag}(D_0^{\mathrm{test}}, D_0^{\mathrm{test}}, C_0 + D_0 + 1/2, \mu_0, C_0) \succ 0. \tag{SL5}$$

This is stronger than a both-infinitesimal testing statement. Evaluate the source polynomial Gram at the real pair (u_0, v_0); its pure-source inverse is bounded. Whiten there and expand in the pure-testing infinitesimals $u - u_0, v - v_0$. The finite separated lift (WA10) then applies verbatim, using only pure-source and pure-testing radicals. All bounded tests of either sign and zero are covered, independently of their rates relative to any source scale. Splitting real and imaginary state parts proves the complex-state conclusion. A separate join below handles every unbounded or literal infinite testing angle.

Chapter 60

All projective tests on the small-r weak-C2 source opening

Component p34609a6519a6
Source: R13/advances/weak_small_r/remote_small_r.tex
SHA256: 34609a6519a6ac418e41c31a89eeef0570026571f12f1742c0687d57f1f5e668
Inventory use: SR1-SR8

60.1 All projective tests on the small-r weak-C2 source opening

Theorem 60.1. *Retain the original physical weak-C2 source, exact stationary contact, all strict source gaps, $n \geq 0$, and $d = 1$. Suppose $B, z, y, r > 0$ are infinitesimal, $G/B, j, y/B, L/B$ are positive bounded units, and H/y is infinitesimal. Then the entire original parent is finite tensor SOS at every independent projective testing pair, with all signs, literal endpoints, and real or complex states. There is no comparison of independent source and testing rates.*

Write $a = z/B$, $t = y/B$, $h = H/B$. Positivity of $G/B = 1 - a(1 - r - H)$ makes a bounded; its unit gap gives $0 \leq \rho = \mathrm{st}(a) < 1$. Put $\tau = \mathrm{st}(t) > 1$. The preceding local lemma proves the source estimates, the exact full-parent five-root identity, and its strict residue for every bounded testing pair. In particular

$$\mu_0 = \frac{1}{2(1-\rho^2)}, \quad m_0 = \rho^2 \mu_0, \quad C_0 = \frac{\tau - \rho}{2(1+\rho)}, \quad D_0 = \frac{1}{2(1+\rho)(\tau - 1)}. \tag{SR1}$$

Here $\mu = m/a^2$, while n, A are unbounded, independently. This section supplies the complete remote join; no remote theorem requiring bounded n is imported as a domain admission.

Two normalized hard state coordinates. Use $\widehat{X} = \sqrt{n}X$, $\widehat{Y} = \sqrt{A}Y$, with $e_x = n^{-1/2}$ and $e_y = A^{-1/2}$ positive infinitesimal. Substitute $X = e_x\widehat{X}$, $Y = e_y\widehat{Y}$ into the original full parent. Its nX^2 and AY^2 packets become the exact hard squares, and every remaining coefficient is bounded. Every cross term with these two normalized states has zero residue. At bounded test residues u_0, v_0, put $U = u_0^2$, $V = v_0^2$. The entire residue of $F/4$ is

$$(1+U)(1+V)(\widehat{X}^2 + \widehat{Y}^2) + \Psi(W, Z),$$

where

$$\begin{aligned} \Psi = {} & m_0(v_0 W - u_0 Z)^2 + C_0(1+V)UW^2 + (D_0 + \tfrac{1}{2})(u_0 W - v_0 Z)^2 \\ & + \tfrac{1}{2}(1+V)UZ^2 - \tfrac{1}{2}(1+U)VZ^2. \end{aligned} \tag{SR2}$$

This retains the complete signed core. If P_2 is its matrix on (W, Z), direct expansion using (SR1) gives

$$\begin{aligned}\det P_2 = {} & D_0\mu_0(\rho V - \tau U)^2 \\ & + \frac{C_0(2m_0+1)}{2}U^2V + C_0D_0UV^2.\end{aligned} \tag{SR3}$$

All three displayed coefficients are strictly positive. For $U > 0$, this determinant is positive for every $V \geq 0$: if $V = 0$ its first term is positive; if $V > 0$ its second is. The W diagonal $m_0V + [C_0(1+V) + D_0 + 1/2]U$ is positive. Thus the full four-state residue is strictly positive. For $U = 0, V > 0$ the same holds when $\rho > 0$; here the two remaining pivots are m_0V, D_0V. Expanding around the real residues and applying the finite separated lift (WA10) proves every such bounded testing chart. All bounded pairs are already covered by the preceding strengthened lemma. The determinant gives a second direct check away from its vanishing locus.

The joint small-a and small-r corner. Only $\rho = 0$, u infinitesimal or zero, and nonzero or infinite v remains outside those arguments. The complete identities (WE8) and (WE10) apply algebraically at arbitrary n, A. In those exact identities make the same pure-source substitutions $X = e_x\widehat{X}$, $Y = e_y\widehat{Y}$, using the five roots

$$(S, R, \widehat{X}, \widehat{Y}, Z) = (aW, uW, \sqrt{n}X, \sqrt{A}Y, Z).$$

All resulting coefficients are bounded polynomials in the testing coordinates with pure-source coefficients. The factors nX^2 and AY^2 again become the exact normalized squares; no term is removed. For $v_0 \neq 0$ finite, the separate pivots on $(S, \widehat{X}, \widehat{Y})$ are

$$\mu_0 V, \qquad 1+V, \qquad 1+V.$$

The remaining block on (R, Z) is

$$\begin{pmatrix} C_0(1+V) + D_0 + 1/2 & -(D_0+1/2)v_0 \\ -(D_0+1/2)v_0 & D_0V \end{pmatrix}, \qquad \det = C_0D_0V^2 > 0. \tag{SR4}$$

Indeed at $\rho = 0$, $C_0 = \tau/2$, $D_0 = 1/[2(\tau-1)]$, and $C_0D_0 - D_0/2 - 1/4 = 0$. For v unbounded put $q = 1/v$ and use the polynomial parent $q^2F(u, 1/q)$. At $u = q = 0$ the full five-root residue of the normalized (WE10) is diagonal with pivots

$$(\mu_0, C_0, 1, 1, D_0) > 0. \tag{SR5}$$

The same finite separated lift covers both charts, including literal $u = 0$ and literal $q = 0$. The roots may be dependent; a strict Gram identity remains a valid finite SOS on them.

Every other projective endpoint. For u unbounded use $p = 1/u$ and the exact polynomial $p^2F(1/p, v)$. With v bounded its normalized full residue is

$$[C_0(1+V) + D_0 + 1/2]W^2 + (1+V)(\widehat{X}^2 + \widehat{Y}^2) + (m_0 + 1/2)Z^2. \tag{SR6}$$

It is strictly positive for every $V \geq 0$, whether ρ vanishes or not. For v unbounded and u bounded the residue is

$$(m_0 + C_0U)W^2 + (1+U)(\widehat{X}^2 + \widehat{Y}^2) + D_0Z^2. \tag{SR7}$$

This is strictly positive if $\rho > 0$ or $U > 0$; the missing case is precisely (SR5). For two unbounded coordinates use the complete polynomial identity (WR6), replace X, Y by their normalized expressions, and keep $P = pZ$, $Q = qZ$ as separate roots. The full residue on $(W, \widehat{X}, \widehat{Y}, P, Q)$ is diagonal with pivots

$$(C_0, 1, 1, D_0, m_0 + 1/2) > 0. \tag{SR8}$$

All exact coefficients are bounded. The finite lift applies in p, q, also at their literal zero values. At $p = q = 0$ both P, Q vanish, preserving the original exact null state direction.

Every bounded pair has both residues zero, $U > 0$, or $U = 0, V > 0$. These are covered by the local lemma, (SR3), and (SR3) or (SR4). Single infinities use (SR5)–(SR7); double infinity uses (SR8). Restoration away from a literal endpoint multiplies by a pure-testing square; no inverse is taken at the endpoint. Every state coordinate and first normal was retained. Applying the real identity separately to real and imaginary state parts gives the complex-state result. No nonemptiness of a residual intersection is asserted.

Corollary 60.2 (Two source-unit assumptions removed)**.** *In the revision-9 weak-C2 source theorem, both additional requirements that r and z/B be bounded units can be omitted, provided $1 - r$ and the other original gaps and ratios remain positive bounded units.*

Proof. The original $0 < r < 1$ is either a unit or infinitesimal. For unit r, revisions 9 and 10 cover unit and infinitesimal z/B. For infinitesimal r, the present theorem covers both possibilities. Positivity of $G/B = 1 - (z/B)(1 - r - H)$ and the retained unit $1 - r$ exclude unbounded z/B. These source-only alternatives are exhaustive. □

The remaining $1-r$-infinitesimal and other source-gap openings are not admitted by this corollary. The failed simple rescaling on that opposite boundary is recorded separately.

Chapter 61

Weak ii source alternatives

Component p268520f6cc26
Source: R13/advances/weak_source/weak_ii_source_alternatives.tex
SHA256: 268520f6cc26c85a8b1785a7b36b1cac0c0cff2ce59142ad1e76834eecf22e4b
Inventory use: WII1-WII6

61.1 Exact source alternatives on the weak-B II face

This is an exhaustive source reduction, not a new generic all-tests admission. Use the original positive source (GE1), with $z, y, B \in \mathrm{Inf}_{>0}$, $B = H + L$, $0 < H, r < 1$, $L > 0$,

$$a_c = 1 + zr, \quad S = a_c + L, \quad j = a_c - L/y > 0, \quad G = B + z(H + r - 1) > 0,$$

and $B - zyr > 0$, $n \geq 0$. Write $T = B + zr$ and $g = G/B$. The already proved common-II source entry gives α weak on this entire face. All comparisons below are in the pure source field. The notation $f \asymp h$ means that both f/h and h/f are bounded.

Uniform estimates before any shape specialization. The exact positive formula for δ gives

$$\frac{\delta}{B + z} \sim 1.$$

Moreover $T \geq B$ and $T > z(1 - H)$, by $G > 0$, while $T \leq B + z$. Consequently T/δ is a positive bounded unit, uniformly without a bound on z/B. Since $a_c, S \sim 1$, the original formulas imply

$$r(2n + 1) = \frac{T a_c S}{\delta} \asymp 1, \qquad \frac{AH}{y} = \frac{a_c T S}{2\delta} \asymp 1. \tag{WII1}$$

Also $0 < g \leq 1 + z$, because $G \leq B + zH$ and $H \leq B$. The exact identity $z(1 - r) = B + zH - G$ rewrites the original m as

$$m = \frac{(1 + y) a_c S}{2} \frac{z}{\delta} \frac{1 + zH/B - g}{g}. \tag{WII2}$$

If g is a unit, this is bounded. If g is infinitesimal, then $z(1-r)/B \sim 1$, so $z/(B+z)$ is a positive unit and $m \asymp 1/g$. In particular there is no omitted intermediate large-m case caused by an independent z/B rate.

Exactly which original coefficients are unbounded. The following five equivalences hold on the complete weak-B face:

coefficient is unbounded	necessary and sufficient source condition
n	$r \in \mathrm{Inf}$
m	$G/B \in \mathrm{Inf}$
A	$H/y \in \mathrm{Inf}$
C	$L/(H + y) \in \mathrm{Inf}$
D	$y/B \in \mathrm{Inf}$ or $j \in \mathrm{Inf}$.

(WII3)

The first three were proved in (WII1)–(WII2). For C, use

$$M = (1+z)T(H+y) - z(1+y)(1-r)L.$$

It bounds C above by a bounded multiple of $(H+y)/L$. If $L/(H+y)$ is infinitesimal, then z/T is bounded and the subtracted term is infinitesimal relative to $T(H+y)$. Thus in that case $C \asymp (H+y)/L$ is unbounded. This proves both directions.

For D, if B/y is infinitesimal then L/y is infinitesimal and $j \sim 1$. The original formula directly gives $D = O(B/y)$, hence D weak. If B/y is noninfinitesimal, then zyr/B is infinitesimal, and the exact positive numerator gives

$$D \asymp B/(yj).$$

Here j is bounded; j infinitesimal forces $L/y \sim 1$ and hence B/y noninfinitesimal. These alternatives give precisely the last line of (WII3). This proof never replaces $B - zyr$ by B in the region where that replacement need not be valid.

The positive-total-scale intersection is explicit. Put $a = H/B \in (0,1)$ and $\tau = y/B > 0$. Every original coefficient is bounded if and only if all six quantities

$$\tau, \quad a, \quad 1-a, \quad j, \quad r, \quad g \quad \text{are positive bounded units.} \tag{WII4}$$

Indeed if τ is infinitesimal, then C, D are unbounded; if τ is unbounded, then C, A are unbounded. With τ a unit, (WII3) says respectively that boundedness of A, C, D, n, m is equivalent to $a, 1-a, j, r, g$ being units. Thus (WII4) is exact and includes arbitrary z/B. The original positive-total-scale theorem already admits this full six-unit interior. After intersecting with $\delta_{\text{tot}} \in \text{Inf}_{>0}$, at least one condition in (WII4) must fail, equivalently at least one line of (WII3) must occur.

The shape alternatives can also be read as follows:

y/B	C	A	D
Inf	unbounded	weak	unbounded
unit	unbounded iff $1-a \in \text{Inf}$	unbounded iff $a \in \text{Inf}$	unbounded iff $j \in \text{Inf}$
unbounded	unbounded	unbounded	weak.

(WII5)

In every row n, m are governed independently by r, g as in (WII3). The exact source inequalities and $n \geq 0$ are still imposed; the table does not assert independent realizability of arbitrary shapes.

The two-weak intersection and the four-product exit. There is a stronger exact equivalence:

$$\boxed{C \in \text{Inf} \iff r, g, H/y, j \in \text{Inf} \iff n, m, A, D \text{ all unbounded.}} \tag{WII6}$$

To prove the forward direction, the positive identity

$$M = (1+z)TH + yG + yzj + zrL(1+y)$$

shows H/L infinitesimal. Since C is bounded, (WII3) makes $L/(H+y)$ a unit. Hence $L \asymp y$, H/y infinitesimal, and $B \asymp y$. The same four positive summands now imply $G/\delta, zr/\delta, zj/\delta$ infinitesimal. The quantity z/δ cannot be infinitesimal: otherwise $B/\delta \sim 1$ and $G/\delta \sim 1$, a contradiction. Thus r, j are infinitesimal. The inequality $G > 0$ then gives B/z a positive-unit lower bound; z/δ a unit supplies its upper bound. Therefore $g = G/B$ is infinitesimal too.

Conversely $r, g \in \text{Inf}$ imply, by $G = B + z(H + r - 1)$,

$$(1-g)B/z = 1 - H - r, \qquad B/z \sim 1.$$

If also $H/y, j \in \text{Inf}$, then $L/y = a_c - j \sim 1$ and $B/y \sim 1$. All four positive summands in $M/(L\delta)$ are now infinitesimal, so C is weak. Finally (WII3) gives one implication of the last equivalence. For

its converse, A unbounded makes H/y infinitesimal and B/y bounded, so D unbounded must mean j infinitesimal. The n, m statements give $r, g \in \mathrm{Inf}$.

The already admitted both-complementary-events-weak theorem removes the first intersection in (WII6). On the remaining directed face C strong and α weak, at least one of n, m, A, D is therefore bounded. Its product with α is infinitesimal. Consequently the four reciprocal-product exit

$$\alpha n, \quad \alpha m, \quad \alpha A, \quad \alpha D \quad \text{all strong}$$

is impossible throughout this directed weak-α/strong-C II face. This is a uniform statement, not a failed-product example.

What this does and does not remove. The exact all-unit region (WII4) is already admitted by the positive-total-scale theorem, and the exact four-large intersection (WII6) is already admitted by the two-weak theorem. Individual real rational source curves remain covered by the separate curve theorem. The latter does not automatically admit these generic multiparameter opening faces. Adjacent dominant-event results require their actual normalized coefficient support and the stated testing-angle hypotheses; two unbounded adjacent coefficients alone do not supply those premises. Thus this reduction identifies the genuine source alternatives still requiring a full-parent argument. It does not establish that their intersection with the entire global complement is nonempty or empty.

Chapter 62

Shared event mechanisms for the remaining C1 sources

Component `p89fd3edb3edd`
Source: `R13/baseline/advances/c1_common/common_remote.tex`
SHA256: `89fd3edb3edd2350eb561f1964d205c81ae33223d97b1f35ac7ab16e710855af`
Inventory use: Original remote/projective testing exits | CE1–CE21 and exact relabelings | CE adjacent C,A noninfinitesimal Alice all-state exit

62.1 Shared event mechanisms for the remaining C1 sources

Normalize the original parent by its positive pure-source coefficient d. Write $D_z = 1 + z^2$, and write the signed core as $2K$, where

$$K = (uW + uvX - Y - vZ)^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_u D_v Z^2. \tag{CE1}$$

The event packets used below are

$$4m(vW - X + uvY - uZ)^2, \quad 4CD_v(uW - Y)^2, \quad 4AD_u D_v Y^2, \quad 4\alpha D_u(vY - Z)^2.$$

Every omitted event and every nonnegative excess is retained as its exact original square. Source and testing fields remain independent. "Strong" means bounded below by a positive real number; "unbounded" means larger than every real number. Neither assertion constrains upper scales.

Theorem 62.1 (Two adjacent unbounded events, with complete raw frames)**.** *If C, A are unbounded, the full original parent is tensor SOS at every projective test for which u is not infinitesimal, and every state. No physicality assumption beyond nonnegative event weights is needed for this particular assertion. By the exact cyclic relabeling below, if A, α are unbounded, the conclusion holds whenever v is not projectively unbounded.*

Proof. Select $C = A = \mu$, with a positive real μ chosen after the real testing residues; unboundedness permits every such choice. The following are exact full quadratic forms, before any restriction to a state ray or contact. They therefore retain every first state normal and hard-normal coupling.

For u a bounded nonzero unit and v bounded, use the four raw roots

$$(W, Y, B, V), \qquad B = X + uZ, \quad V = vZ.$$

The entire core equals

$$K = (uW - Y + uvB)^2 + D_v B^2 + D_u v^2 Y^2 + D_u u^2 V^2 - 2D_u uV(W + vB). \tag{CE2}$$

The two selected event forms remain $4\mu D_v(uW - Y)^2$ and $4\mu D_u D_v Y^2$. In their common real kernel $W = Y = 0$, the core has matrix on (B, V)

$$\begin{pmatrix} 1 + v^2 D_u & -D_u uv \\ -D_u uv & D_u u^2 \end{pmatrix}, \qquad \det = D_u u^2 > 0. \tag{CE3}$$

Here and in residue matrices, bounded testing letters mean their real residues. Thus a sufficiently large finite μ gives a strictly positive residue on the entire four-root space, even when $v = 0$ literally.

For $p = 1/u$ infinitesimal and v bounded, homogenize by p^2 and take

$$(W, Y, B, R), \qquad B = pX + Z, \quad R = vX.$$

The exact full core is

$$K_p = B^2 - 2pvBW - 2vBY + D_pR^2 + 2D_pRW + (W - pY)^2 + D_pv^2Y^2. \tag{CE4}$$

The selected packets are $4\mu D_v(W - pY)^2$ and $4\mu D_pD_vY^2$. At $p = 0$ their sum with $2K_p$ is

$$4\mu D_v(W^2 + Y^2) + 2(W + R)^2 + 2(B - vY)^2, \tag{CE5}$$

strict for every $\mu > 0$. The formula allows literal $p = 0$ and $v = 0$; the raw roots need not form independent state coordinates.

For projectively unbounded v, put $w = 1/v$, homogenize by w^2, and use (W, X, Y, H) with $H = wZ$. The exact full core is

$$\begin{aligned} K_w = {} & u^2H^2 - 2u^2HY - 2uHW + 2uwHX + u^2w^2W^2 \\ & + 2u^2wWX - 2uw^2WY + (D_u + w^2)(X^2 + Y^2) - 2uwXY. \end{aligned} \tag{CE6}$$

The selected packets are $4\mu D_w(uW - Y)^2$ and $4\mu D_uD_wY^2$. If u is a bounded nonzero unit, at $w = 0$ their common kernel has $W = Y = 0$ and $K_w = D_uX^2 + u^2H^2$ strictly positive. A sufficiently large real μ again works. If also $p = 1/u$ is infinitesimal, homogenize (CE6) by p^2. The exact full core is

$$\begin{aligned} K_{pw} = {} & H^2 - 2pHW + 2pwHX - 2HY + w^2W^2 + 2wWX - 2pw^2WY \\ & + (D_p + p^2w^2)(X^2 + Y^2) - 2pwXY. \end{aligned} \tag{CE7}$$

At $p = w = 0$ the full selected residue is

$$4\mu(W^2 + Y^2) + 2X^2 + 2(H - Y)^2 > 0. \tag{CE8}$$

All these exact Gram coefficients are bounded. Strictness in (CE3) and its counterpart following (CE6) implies a finite cap by the elementary real Schur-complement argument: the event Gram is positive definite on the complement of its kernel, and the core is positive definite on the kernel. No source/testing rate comparison is involved. Apply the bounded positive-residue Gram lemma, and restore every retained original square.

For the second assertion, compose the inherited Alice reflection with the swap/reflection. The resulting one-step inverse cycle rotation is

$$(u', v') = (1/v, u), \quad (W', X', Y', Z') = (Y + vZ, -W - vX, Z - vY, vW - X), \tag{CE9}$$

and sends event index i to $i - 1$ in $(\lambda, D, C, A, \alpha, m, \beta, n)$. In particular original A, α become new C, A. Its full-parent identity is

$$v^2F(\mathrm{rot}^{-1}c; 1/v, u; \psi') = D_vF(c; u, v; \psi). \tag{CE10}$$

The state map has determinant of absolute value D_v^2, and its homogeneous version at $v = \infty$ is invertible as well. Thus $u' = 1/v$ not infinitesimal is exactly v not projectively unbounded. Only pure testing factors are restored. Both signs, all literal endpoints, and complex states (by real/imaginary splitting) are included. □

Theorem 62.2 (Four consecutive strong events)**.** *For an original physical parent with C, A, α, m strong, the full parent is tensor SOS at every test except possibly u a bounded nonzero unit and v projectively unbounded. All original upper coefficient scales and all nonnegative unselected coefficients are allowed.*

Proof. Cap these four events at any positive real μ smaller than all four weights, and retain every excess. Noncommuting-residue tests use the inherited physical-source theorem, applied to the original full parent. For commuting-residue tests use the capped parent.

First suppose v is bounded. On the three open commuting axes, the core and the four caps have zero common kernel: at $u = 0$, the core forces $X = Y = 0$, then m and α force $W = Z = 0$ when v is a unit; at $u = \infty$, C, A force $W = Y = 0$, then the core forces $X = Z = 0$; at $v = 0$, A, α force $Y = Z = 0$, then the core forces $W = X = 0$ when u is a unit. These are strictly positive bounded state Grams.

At $(u, v) = (0, 0)$ the residue is

$$(2 + 4\mu)X^2 + (2 + 8\mu)Y^2 + 4\mu Z^2.$$

Only the W ray remains. Normalize $W = 1$ and take the five infinitesimal roots (u, v, X, Y, Z). The full polynomial has no constant or linear term; its quadratic part is

$$2(u - Y)^2 + 2X^2 + 4\mu\{(v - X)^2 + (u - Y)^2 + Y^2 + Z^2\}, \tag{CE11}$$

which is strictly positive. At $(u, v) = (\infty, 0)$ only the X ray remains. Normalize $X = 1$ and use (p, v, W, Y, Z). Its quadratic part is

$$2(W + v)^2 + 2(p + Z)^2 + 4\mu\{(p + Z)^2 + W^2 + Y^2 + Z^2\}, \tag{CE12}$$

also strictly positive. All higher monomials in each chart allocate to products of two displayed roots with infinitesimal bounded coefficients. This is an exact finite Gram, not a truncation certificate.

For $w = 1/v$ infinitesimal use the full raw frame of (CE6). All four selected packets, with every normal present, are exactly

$$4\mu\{(W - wX + uY - uH)^2 + D_w(uW - Y)^2 + D_u D_w Y^2 + D_u(Y - H)^2\}. \tag{CE13}$$

At $u = w = 0$ their sum with $2K_w$ has residue

$$4\mu W^2 + 2X^2 + (8\mu + 2)Y^2 + 4\mu(Y - H)^2, \tag{CE14}$$

strict on all four roots. If $p = 1/u$ is infinitesimal, homogenize again; the exact packets become

$$4\mu\{(pW - pwX + Y - H)^2 + D_w(W - pY)^2 + D_p D_w Y^2 + D_p(Y - H)^2\}. \tag{CE15}$$

Together with (CE7), at $p = w = 0$ the residue is

$$4\mu W^2 + 2X^2 + 4\mu Y^2 + (8\mu + 2)(Y - H)^2, \tag{CE16}$$

again strict. These frames require no state normalization, and no division by either small testing coordinate. Literal $w = 0$ is included. The only chart not covered is ordinary u and infinitesimal w. Restore the original packets and all pure factors as above. □

Lemma 62.3 (Matched weak outer events at the large-Bob corner)**.** *Suppose $q > 0$ is a pure-source infinitesimal and*

$$C \geq cq^2, \quad m \geq m_* q^2, \quad A \geq a_*/q^2, \quad \alpha \geq b_*/q^2 \tag{CE17}$$

for positive real c, m_*, a_*, b_*. At projectively unbounded v, the complete parent is tensor SOS whenever u is infinitesimal or projectively unbounded. In the latter chart the lower bound on m is unnecessary.

Proof. Keep $w = 1/v$ and $H = wZ$ as in (CE6)–(CE7). If u is infinitesimal, use the five raw roots

$$E = qW, \quad X, \quad F = Y/q, \quad G = H/q, \quad V = uwW.$$

The exact whole core (CE6) becomes

$$K_w = V^2 + (D_u + w^2)X^2 + q^2\{u^2(G - F)^2 + D_w F^2\}$$

$$- 2uGE + 2quwX(G - F) + 2uVX - 2uw^2EF. \tag{CE18}$$

Cap only m, A, α at the indicated weights. Their complete packets are respectively

$$4m_*(E - qwX + uq^2(F - G))^2, \quad 4a_*D_uD_wF^2, \quad 4b_*D_u(F - G)^2.$$

The selected residue is

$$2V^2 + 2X^2 + 4m_*E^2 + 4a_*F^2 + 4b_*(F - G)^2 > 0. \tag{CE19}$$

Every coefficient in (CE18) is bounded independently of all relative source/testing rates.

If instead $p = 1/u$ is infinitesimal, use

$$E = qW, \quad F = Y/q, \quad G = H/q, \quad R = X + wW, \quad S = pX.$$

The exact whole core (CE7) is

$$K_{pw} = R^2 + D_wS^2 + q^2(G - F)^2 + p^2D_wq^2F^2 - 2pE(G + w^2F) + 2qwS(G - F). \tag{CE20}$$

Cap only C, A, α. Their complete packets are

$$4cD_w(E - pq^2F)^2, \quad 4a_*D_pD_wF^2, \quad 4b_*D_p(F - G)^2.$$

Together with $2K_{pw}$ their residue is

$$2R^2 + 2S^2 + 4cE^2 + 4a_*F^2 + 4b_*(F - G)^2 > 0. \tag{CE21}$$

Only the pure-source inverse $1/q$ occurs in a root. These exact raw frames include every state normal and every literal endpoint. The strict-residue lemma and restoration of excess packets prove the claim. □

Exact application to the original C1 source rows. The source premises here are precisely the already proved original formulas (NC1)–(NC7), (NI3)–(NI10); no independent source family is substituted. These common theorems give the following new quantified test reductions, always intersected with prior admissions:

1. C1.1, every e type: (NI3) gives C, A unbounded; only u infinitesimal remains.
2. C1.4, L infinitesimal and η unbounded: (NI6) again gives C, A unbounded; only u infinitesimal remains.
3. C1.4, L unbounded: (NI4) and (CE20) leave only u bounded and v unbounded.
4. C1.5 actual UH, η unbounded and ζ unit: (NI9) gives four strong events; only u unit and v unbounded remains.
5. C1.5 actual UH, ζ unbounded: (NI10) and (CE17) leave the same sole strip, u unit and v unbounded.

For the last line choose $q = \zeta^{-1/2}$. Formula (NI10) gives $C, m \asymp q^2$, $\alpha \asymp 1/(q^2Q)$ and $A \asymp 1/r$; the original $Q = 1 - x\zeta r > 0$, with unit x, bounds Aq^2 below. Thus (CE17) holds with positive real constants. The adjacent-event theorem covers bounded v and (CE18)–(CE21) remove both end types of u at unbounded v. Both actual UH source patterns now have the same sole untreated testing strip, ordinary u and unbounded v.

In the L-unbounded C1.4 line take $q = L^{-1/2}$. Then (NI4) gives $C \asymp q^2$, $\alpha \asymp 1/(q^2Q)$ and $Aq^2 \asymp 1/(Lr) > 1$. The p-chart (CE20), which needs no lower bound on m, removes the projectively unbounded-u tests there.

All bounds are relative to original d. The four-source-event result does not require the additional NI9 total-scale condition that at least one opening be infinitesimal. Every literal permitted $D = 0$ or $\beta = 0$ is retained throughout, since neither coefficient was capped. No claim of nonemptiness or full-row completion is made. In particular, the L unit and the L infinitesimal/η bounded parts of C1.4 are unchanged, and these remote results do not supply the missing critical contact conductors.

Scoped corroboration. The checker reconstructs the exact core, all six raw identities (CE2), (CE4), (CE6), (CE7), (CE18), (CE20), their selected packets and residues, the two corner quadratic forms, and the cyclic core and selected-event transport. It checks whole quadratic polynomials, hence all state normals at once. The source entry, ordered-field bounds, real Schur argument, strict-residue lemma, and the inherited noncommuting theorem remain the analytical proof. No whole-row count changes follow from these checks.

Chapter 63

C1 UI: completion of the ordinary/small strip

Component `pa0b854a04f49`
Source: `R13/baseline/advances/c1_ui/ui_all_tests.tex`
SHA256: `a0b854a04f49c38b64e6b24efb55cafc459729aa6cc777d41b9b5c17041d0e9b`
Inventory use: CU1–CU13 and entire full-normal/finite-cap proof | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

63.1 C1 UI: completion of the ordinary/small strip

Use the unchanged native C1 source in the representative of (NI7): k, η are positive bounded units and $\zeta = g^2$ is positive infinitesimal. All original openings and $D = 0$ are retained. The exact stabilizer recovers the other representative; no condition $k \le 1$ is imposed after that change. This section proves the missing ordinary/small testing strip, and joins the remote theorem (CR1)–(CR13).

Exact bounded remainder. Normalize by the positive pure-source a. Write $x = k^2$, $r = 1 - a\eta$, $K = 1 + x$, $R = 1 + xr$, $M = R - t$, and

$$J_s = K - xtg^2 - a, \quad E = 1 - x\eta + tx\eta g^2, \quad Q_c = 1 - xg^2 r, \quad y = x\eta g^2.$$

Here $a, t, r, 1 - r, M, J_s$ are positive bounded openings; $E, Q_c, R, K, r + \eta$ are positive units. The exact polynomials P_4, P_d, P_α are (NC2). The coefficients after division by a are

$$\begin{aligned} c &= 2/a, & A_0 &= \frac{2\eta^2 P_4}{Kr(r+\eta)E}, & d_0 &= \frac{4\eta P_d}{K(r+\eta)E}, \\ D_0 &= \frac{2(1+y)}{J_s E} - d_0/2, & \alpha_0 &= \frac{2x^2\eta P_\alpha}{KQ_c E}, & \alpha/a &= g^2\alpha_0. \end{aligned} \tag{CU1}$$

In particular c, A_0 are strong; d_0, α_0 are units; and $ca, A_0 r, D_0 J_s$ are bounded with continuous rational expressions through every opening. Put

$$h_2 = \frac{2x\eta(1 - tg^2)(1 + y)}{E}, \qquad h_b = \frac{2x\eta[1 + (1 - t)g^2]}{E}, \qquad h_b - h_2 = 2x\eta g^2.$$

Both h_2, h_b are positive units. The exact coefficients are $m/a = h_2/(g^2 M)$ and $\beta/a + d_0/2 = h_b/(g^2 t)$.

For the test put $D_u = 1 + u^2, D_v = 1 + v^2$ and

$$Q = W + uY, \quad B = X + uZ, \quad p = t/R, \quad N = (B - pvQ)/g, \quad L = vQ/g.$$

The inverse $W = Q - uY$, $X = gN + pvQ - uZ$ is polynomial. Retain the two positive squares

$$\frac{4h_2 t}{MR} N^2, \qquad \frac{4h_b M}{tR} D_v N^2. \tag{CU2}$$

Their remainder is exactly (CL7). Choose bounded caps $c_* \le c$, $A_* \le A_0$, $D_* \le D_0$, with $c_*, A_* > 0$, $D_* \ge 0$; bounded original weights can be kept unchanged. Retain the three positive centered excesses

$$\begin{aligned} &4(c-c_*)D_v(r_t-avZ)^2, \quad 4(A_0-A_*)D_uD_v(Y-rvZ/\eta)^2, \\ &4(D_0-D_*)(J_t+J_svZ)^2, \end{aligned} \tag{CU3}$$

where

$$r_t = uQ - D_uY, \quad J_t = r_t - D_uvZ + ugvN + upv^2Q.$$

All five retained rows vanish at the actual contact

$$(u,v) = (k, k\sqrt{\eta}g), \quad (W,X,Y,Z) = (g/\sqrt{\eta}, k(tg^2-1), krg/\sqrt{\eta}, 1). \tag{CU4}$$

Set $c_a = 2 - c_*a$, $A_r = A_0r - A_*r$, $D_J = D_0J_s - D_*J_s$. After removing (CU2)–(CU3), the entire parent divided by a is

$$\begin{aligned} G = \frac{4}{R}&\{(h_2+h_bD_v)N^2 + 2(h_bD_v-h_2)LN + [h_2(1-p)+h_bpD_v]L^2\} \\ &+ 4c_*D_vr_t^2 + 8c_aD_vvZr_t - 4c_aaD_vv^2Z^2 \\ &+ 4A_*D_uD_vY^2 + \frac{8A_r}{\eta}D_uD_vvZY - \frac{4A_rr}{\eta^2}D_uD_vv^2Z^2 \\ &+ 4D_*J_t^2 - 8D_JvZJ_t - 4D_JJ_sv^2Z^2 \\ &+ 4\alpha_0D_u(gvY-gZ)^2 + 2d_0\{J_t^2 + D_u(v^2Y^2 - 2vYZ - v^2Z^2)\}. \end{aligned} \tag{CU5}$$

Thus G is an exact bounded Gram on (N, Q, Y, vZ, gZ, L) whenever u is a bounded unit and v infinitesimal. Every source denominator in its coefficients is a unit. Formula (CU5), rather than a scalar restriction, supplies all normal terms below.

Local contact lifting with all first normals. Put $\kappa = k\sqrt{\eta}$, $\varepsilon = \kappa g$, $s = R/(k\eta)$, $\delta = u - k$, and define

$$E_Q = Q - svZ, \quad E_Y = Y - rvZ/\eta, \quad R_h = (vQ - s\varepsilon^2Z)/\varepsilon, \quad S = v\delta Z, \quad T = \varepsilon\delta Z.$$

Let $c_L = 4[h_2(1-p) + h_bpD_v]/R$, a positive bounded unit. The sole potentially unbounded L row can be replaced exactly:

$$L = \kappa R_h + kRgZ, \qquad L^2 - \kappa^2R_h^2 = 2kRvQZ - k^2R^2g^2Z^2. \tag{CU6}$$

Consequently (CU5) equals $c_L\kappa^2R_h^2$ plus $8\kappa(h_bD_v - h_2)NR_h/R$ and a polynomial G_0. Explicitly, in (CU5) replace its first line by

$$\frac{4}{R}(h_2+h_bD_v)N^2 + 8k(h_bD_v-h_2)gNZ + c_L(2kRvQZ - k^2R^2g^2Z^2). \tag{CU7}$$

All coefficients of G_0 are bounded polynomials in u, v, g with pure-source unit denominators. The coefficient of NR_h is infinitesimal, since $h_bD_v - h_2 = 2x\eta g^2 + h_bv^2$.

Here are exact prescriptions for every normal coefficient, avoiding any implicit omission. Substitute $Q = E_Q + svZ$ and $Y = E_Y + rvZ/\eta$ in G_0 and collect the quadratic polynomial in (N, E_Q, E_Y). The constant coefficient is $\phi(u,\xi)Z^2$, where $\xi = v^2$ and $\phi \in (g^2, \xi)$. The linear coefficients of E_Q, E_Y are $vL_Q(u,\xi)Z, vL_Y(u,\xi)Z$; that of N is $gL_N(u,\xi)Z$, where $L_N \in (g^2, \xi)$. These parity and ideal assertions follow term by term in (CU5),(CU7). The remaining quadratic normal block is the full coefficient collection of the same expression. For explicit pen-and-paper formulas, set

$$b = r/\eta, \quad F_0 = us - D_ub, \quad G_0^{\mathrm{sc}} = F_0 - D_u + up\xi s, \quad H_0 = (8D_* + 4d_0)G_0^{\mathrm{sc}} - 8D_J.$$

Then, writing $D_v = 1 + \xi$ in these polynomials,

$$L_Q = 2c_LkR + 8D_vu(c_*F_0 + c_a) + u(1+p\xi)H_0,$$

$$
\begin{aligned}
L_Y = & -8D_vD_u(c_*F_0+c_a)+8D_uD_v(A_*b+A_r/\eta)-D_uH_0\\
& +8\alpha_0D_ug^2(b\xi-1)+4d_0D_u(b\xi-1),\\
L_N = & \,8k(h_bD_v-h_2)+u\xi H_0.
\end{aligned}
$$

The complete scalar coefficient is

$$
\begin{aligned}
\phi = & \,c_L(2kRs\xi-k^2R^2g^2)+4c_*D_vF_0^2\xi+8c_aD_vF_0\xi-4c_aaD_v\xi\\
& +4A_*D_uD_vb^2\xi+8(A_r/\eta)D_uD_vb\xi-4A_rrD_uD_v\xi/\eta^2\\
& +4D_*(G_0^{\mathrm{sc}})^2\xi-8D_JG_0^{\mathrm{sc}}\xi-4D_JJ_s\xi\\
& +4\alpha_0D_ug^2(b\xi-1)^2+2d_0\{(G_0^{\mathrm{sc}})^2\xi+D_u(b^2\xi^2-2b\xi-\xi)\}.
\end{aligned}
$$

These formulas display the two asserted ideals directly. They also give a short explicit input for the monic divisions, without an unprinted state-normal computation.

The source is an exact stationary zero and every removed square vanishes there. Therefore

$$
\phi=\phi_u=\phi_\xi=0,\qquad L_Q=L_Y=L_N=0\quad\text{at }(u,\xi)=(k,\varepsilon^2).
$$

Successive monic polynomial division gives, with bounded coefficients,

$$
\phi=\delta^2(\varepsilon^2B_0+\xi B_1)+\delta(\xi-\varepsilon^2)B_2+(\xi-\varepsilon^2)^2B_3, \tag{CU8}
$$

$$
L_i=\delta L_{i0}+(\xi-\varepsilon^2)L_{i1}(i=Q,Y),\quad L_N=\delta(g^2L_{N0}+\xi L_{N1})+(\xi-\varepsilon^2)L_{N2}.
$$

For clarity, each quotient is obtained by subtracting the value at $u=k$ and dividing by the monic $u-k$, then by the monic $\xi-\varepsilon^2$. The second u quotient of ϕ stays in (g^2,ξ); its constant and first u terms have the stated double and simple zeros in ξ. The same argument gives the L_N ideal. No small source number is inverted in these divisions.

Put $R_s=(v^2-\varepsilon^2)Z/\varepsilon$. The exact root identities

$$
\varepsilon R_s=(\varepsilon R_h-vE_Q)/s,\quad R_sT=(R_hT-SE_Q)/s,\quad S^2-T^2=\delta(R_hT-SE_Q)/s \tag{CU9}
$$

allocate (CU8) and every first normal on the six roots (N,E_Q,E_Y,R_h,S,T). For example $v\delta Z=S$, $\delta(\xi-\varepsilon^2)Z^2=(R_hT-SE_Q)/s$, and $(\xi-\varepsilon^2)^2Z^2=(\varepsilon R_h-vE_Q)^2/s^2$. The N normal has only infinitesimal couplings. All quadratic normal coefficients are already bounded. This is the required exact full lift.

Strict matched Hessian and source boundary caps. First take finite interior source residues. The leading form is (CL2). Put $V=k\sqrt{\eta}V_1$, $W=W_1/\sqrt{\eta}$, $Y=kY_1/\sqrt{\eta}$, $Z=1$. At the contact $(W_1,Y_1,V_1,u)=(1,r,1,k)$, the Schur complement of the two state coordinates in half the Hessian, in coordinates (V_1,u), is

$$
\begin{aligned}
G_{11} &= \frac{32\eta x^2P_4}{KE_0}, & G_{12} &= \frac{16\eta k^3(R^2-2KE_0)}{KE_0R},\\
G_{22} &= \frac{32\eta x(P_4+xrP_d)}{R^2P_4}, & \det G &= \frac{256\eta^2x^3(4K^2E_0-R^2)}{K^2E_0^2}.
\end{aligned} \tag{CU10}
$$

Here $E_0=1-x\eta$, $P_4=KE_0+x(1-r)$, $P_d=1+2x+rx^2$. There are no opening denominators. The leading $D\ge0$ condition is

$$
K(2+xr)E_0-R^2\ge0,
$$

so $4K^2E_0-R^2\ge K(4K-2-xr)E_0>0$. The state block is uniformly bounded below since c,A_0 are strong. Thus the matched Hessian is strict, uniformly through every permitted source boundary after the appropriate hard directions are retained.

We give the boundary justification explicitly. Divergences can occur only at $a=0$, $r=0$, or $J_s=0$, for c,A_0,D_0, respectively. Since $a+J_s\sim K$, c and D_0 cannot both diverge. If $a=0$, then

$r = 1$, so c is the only divergent weight. The only possible divergent pair is A_0, D_0. Their hard-state rows are E_Y and $kE_Q - KE_Y$, with nonzero determinant k. All angular parts of these centered rows are bounded. Solving these at most two real linear equations therefore supplies a bounded competitor for the minimizing state vector. The bounded finite part and (CU10) imply that the state minimizer is bounded as well, by completing its positive state square. The full matched Hessian hence has a uniform positive lower bound on the Euclidean norm in a nearby finite-source family. On a vector in the limiting hard kernel, adjust only the two state coordinates to annihilate the nearby exact hard rows. This adjustment tends to zero, and the divergent penalties vanish exactly. The bounded finite part is consequently strictly positive on that kernel. The elementary real finite-penalty lemma supplies finite caps.

Every allowed residue is approachable by such finite sources. For fixed $x > 0$, the real leading inequalities are

$$0 \le r \le 1, \qquad \eta_{\min}(r) = \frac{1-r}{K} \le \eta \le \eta_{\max}(r) = \frac{1 - R^2/[K(2+xr)]}{x},$$

with positive interval width $R/[xK(2+xr)]$. Perturb r into $(0,1)$ and its interval coordinate into $(0,1)$. This makes a, r, J and the D numerator strictly positive while converging to the given residue. Taking $t = R/2$ and then sufficiently small real $g > 0$ restores all original source signs, including $\beta > 0$. This construction, rather than substitution into a singular leading denominator, justifies the boundary limit.

In the six-root residue Gram, N is an isolated positive pivot. The (E_Q, E_Y) block is strict, R_h has positive pivot $c_L\kappa^2$, S couples only to E_Q, E_Y, and T only to R_h. Setting $S = T$ is the matched Hessian under an invertible real change of coordinates: R_h/g has relative-angle derivative $2s\kappa \neq 0$, and $T/g = \kappa\delta$ supplies the ordinary-angle derivative. After the hard Schur complements, the sum of the S, T diagonals is therefore strictly positive. Add a real multiple of the last zero in (CU9) to redistribute this sum into two positive diagonals. The extra coefficients contain δ and vanish in the residue. The bounded strict-residue Gram lemma proves the full parent at $u - k$ and v infinitesimal, including every source opening.

Ordinary angles away from contact. Let now $\bar{u}^2 \neq \bar{k}^2$, with u a unit and v infinitesimal. Use the bounded frame (N, r_t, Y, vZ, gZ, L) of (CU5), and the exact zero

$$L(gZ) - Q(vZ) = 0, \qquad Q = (r_t + D_u Y)/u. \tag{CU11}$$

For finite leading source weights, N is an isolated positive pivot. The remaining residue is

$$q^T P q + 2Ub^T q + c_2 U^2 + 4hL^2 + c_0 T^2, \quad q = (r_t, Y)^T,\ U = vZ,\ T = gZ,$$

where the source coefficients are (CL1) and

$$\begin{gathered} s_D = 4D + 2d, \quad s_C = 4C + s_D, \quad P = \operatorname{diag}(s_C, 4AD_u), \\ \ell = (1/u, D_u/u)^T, \quad b = (-s_D D_u, -2dD_u)^T, \\ c_0 = 4\alpha D_u, \qquad c_2 = s_D D_u^2 - 2dD_u. \end{gathered} \tag{CU12}$$

Apply the real rank-one allocation lemma (UA1)–(UA3), using (CU11). We verify its hypotheses uniformly through all source boundaries. Writing $d_a = \ell^T P^{-1}\ell$, $e_a = \ell^T P^{-1} b$, $h_a = c_2 - b^T P^{-1} b$, direct inversion yields

$$\begin{aligned} d_a &= \frac{aJE_0}{8u^2(JE_0 + a)} + \frac{D_u}{4Au^2}, \\ e_a &= -\frac{aD_u}{u(JE_0 + a)} - \frac{dD_u}{2Au}, \\ h_a &= \frac{8D_u^2}{JE_0 + a} - 2dD_u - \frac{d^2 D_u}{A}, \\ P^{-1}b &= (-aD_u/(JE_0 + a), -d/(2A))^T. \end{aligned} \tag{CU13}$$

All are bounded continuous, since $JE_0 + a$ is a positive unit and $1/A$ has a continuous bounded formula proportional to r. The same holds for $P^{-1}\ell$.

Normalize the Laurent polynomial (CL4) by As_C. Its U and T^{-1} coefficients are, respectively, $1/[2(JE_0 + a)]$ and $\alpha/4$, positive units. Its T coefficient is

$$h\{(D + d/2)/s_C + 2CD/(As_C) + d/(8A)\},$$

also a positive unit: when $1/A = 0$, $r = 0$ forces a a unit, so $(D + d/2)/s_C = a/[4(JE_0 + a)]$ is a unit; otherwise the last term is positive. The U^{-1} coefficient is $h\alpha(1/s_C + 1/(4A))$. If it vanishes, $r = 0$ and $J = 0$, and the T/U coefficient has the positive term $hD/s_C = h/4$. All remaining normalized nonconstant coefficients are bounded and nonnegative. The constant is bounded by the exact contact equation. Thus strict convexity and coercivity in logarithmic coordinates persist at every boundary. The normalized Laurent function retains its unique stationary zero $(U, T) = (x, x\eta)$.

For fixed $u^2 \neq x$, the rank-one determinant polynomial is

$$p(T) = c_0 + (h_a + 4hd_ac_0)T + 4h(d_ah_a + e_a^2)T^2 = \frac{16D_u}{As_C}T\mathcal{L}(u^2, T) > 0.$$

Its constant and leading coefficients are positive units by the preceding coefficient bounds. The strict allocation interval therefore survives at every source boundary. The two real Gram blocks have a uniform positive lower bound; (CU13) controls their state minimizers even if P is unbounded. The centered divergent rows have hard parts r_t, Y, r_t; the only simultaneous pair is Y, r_t, whose determinant is one. Exact small state corrections annihilate the large rows in the nearby finite family. The same finite-penalty argument as above supplies caps in (CU3). The exact bounded remainder (CU5), after (CU11), has strict residue. This proves every ordinary/small test away from contact.

Exhaustive admission. The two arguments cover the whole ordinary/small testing strip; the exact gauges cover negative contact signs. The remote theorem (CR1)–(CR13), including CR5a, covers its complement. All transformations are homogeneous in the state; no state coordinate was divided out in the tensor proof. Real and imaginary state parts are treated separately. Hence the entire actual UI subdomain of C1.5, and its exact stabilizer images, has an all-tests tensor-SOS proof. The actual UH subdomain of C1.5 remains a separate obligation.

Chapter 64

A common exact II reduction for the two C2 double corners

Component `pfb3a83f3849a`
Source: `R13/baseline/advances/c2_double/common_ii_reduction.tex`
SHA256: `fb3a83f3849aa5acd6ee74e9116123f4973ae9acfc67b2979d39199f16fcc15e`
Inventory use: source entry and exact C2 reflection; DI1-DI6 as full-parent identities

64.1 A common exact II reduction for the two C2 double corners

This section proves source reduction and an exact full-parent identity. It does not claim a new all-tests admission. In particular, the scalar identity below is not substituted for the missing allocation of its three state normals. Source and testing fields remain independent.

One source chart, with the orientation recorded. Use the positive C2 chart and put

$$z = u_c^2, \quad y = v_c^2, \quad H = v_c Y_c, \quad L = v_c(u_c W_c - Y_c), \quad B = H + L, \quad r = -X_c/u_c.$$

Here the contact is normalized by $Z_c = 1$, $0 < H < 1$, $0 < r < 1$, $L > 0$, and

$$A_c = 1 + zr, \quad j = A_c - L/y > 0, \quad G = B + z(H + r - 1) > 0, \quad B - zyr > 0.$$

Suppose both actual angles are infinitesimal. Then $L < yA_c$ is infinitesimal, and B is bounded. The exact reflection

$$(u_c, v_c; W_c, X_c, Y_c, Z_c; n, m, C, A, D, \alpha) \mapsto (1/v_c, 1/u_c; Z_c, X_c, Y_c, W_c; n, D, \alpha, A, m, C)$$

takes directed weak-C HH to II with C strong and α weak. It takes the strong II/HH orbit to II with both C, α strong. The weak orientation has therefore changed under reflection and must not be relabelled as weak-C II.

Proposition 64.1 (The exact common source face)**.** *For every positive II source, α is weak if and only if B is infinitesimal; α is strong if and only if B is a positive unit. In the latter case C is automatically unbounded.*

Proof. Use the exact δ, M of (GE1), equivalently

$$\begin{aligned} \delta &= z + (1+z)B + (1+z)B^2 + 2z(1+z)Br + z^3r^2 + 2z^2r \\ &\quad + y\{(1+z)B + z + z^2r(2-r)\}, \\ M &= (1+z)(B+zr)(H+y) - z(1+y)(1-r)L. \end{aligned}$$

All terms in δ are nonnegative and $\delta \geq B + z$. The lower bound used below follows from the exact positive identity

$$M = (1+z)(B+zr)H + yG + yzj + zrL(1+y).$$

If B is infinitesimal, then H is infinitesimal, $1-H$ is a unit, and

$$0<\alpha=\frac{(B+zr)M}{2(1-H)\delta}\le\frac{(B+zr)(1+z)(H+y)}{2(1-H)}$$

is infinitesimal. If B is a unit, then $H=B-L$ is a unit, δ is a unit, and the positive decomposition of M gives $M\ge(1+z)(B+zr)H$. Thus M is a unit and α is strong. Moreover $CL=A_cM/(2\delta)$ is a unit and L is infinitesimal, so C is unbounded. These cases exhaust bounded positive B. No lower bound on r, j, H, or $1-H$ was imposed in the weak case, and $n=0$ was not removed. □

Thus the two residual leaves are, respectively, the $B\to0$ face and the B-unit face of one actual II source chart. This is a reduction, not an assertion that the weak face has the same bounded roots as the unit face.

A parity-preserving graph with every state normal. For independent testing angles u,v set $U=u^2$, $V=v^2$ and write

$$X=-\frac{rUv}{B}W+\xi,\qquad Y=\frac{Hu}{B}W+\eta,\qquad Z=\frac{uv}{B}W+\zeta. \tag{DI1}$$

This is an exact polynomial change of all four homogeneous state coordinates, with determinant one. Its only inverse is the positive pure-source inverse $1/B$. It is legal for arbitrary testing signs, at $u=0$ or $v=0$, and for $W=0$. At $(U,V)=(z,y)$ its graph is the exact actual kernel normalized by $W=1$.

Put $q=H+r-1$. The graph amplitudes, with W suppressed, are

$$\begin{aligned}
&x_0=-rUv/B, && y_0=Hu/B, && z_0=uv/B,\\
&M_0=v(B+Uq)/B, && c_0=Lu/B, && J_0=u\{L-V(1+rU)\}/B,\\
&p_0=-(1-H)uv/B, && b_0=(1-r)Uv/B.
\end{aligned} \tag{DI2}$$

The complete state-normal quadratic of the original parent is

$$\begin{aligned}
Q=&(4m+2)(\xi+u\zeta-uv\eta)^2+(4nD_uD_v+2D_uv^2)\xi^2\\
&+(4CD_v+4AD_uD_v+2D_v)\eta^2+4D(uv\xi-\eta-v\zeta)^2\\
&+4\alpha D_u(v\eta-\zeta)^2.
\end{aligned} \tag{DI3}$$

In particular the signed core restricted to $W=0$ is exactly

$$2(\xi+u\zeta-uv\eta)^2+2D_uv^2\xi^2+2D_v\eta^2.$$

This SOS identity uses neither a matrix inverse nor a mixed order.

Here are every first-normal polynomial, without suppressed terms. Define

$$\begin{aligned}
A_x(U)=&-4nr(1+y)(1+U+z)-4mq\\
&+(4D+2)\{L-y[1+r(U+z)]\}+2(1+y)(1-r),\\
B_x(U)=&-4nrU(1+U)-(4D+2)U(1+rU)+2(1-r)U,\\
A_\eta=&\,4myq+4AH(1+y)+(4D+2)ry-(4\alpha+2)y(1-H),\\
B_\eta(U)=&\,4m(B+Uq)-4CL+4AH(1+U)\\
&+(4D+2)(1+rU)-(4\alpha+2)(1+U)(1-H),\\
A_\zeta=&-4mq+(4D+2)ry+(4\alpha+2)(1-H)-2r(1+y),\\
B_\zeta(U)=&\,4D(1+rU).
\end{aligned} \tag{DI4}$$

Then the exact full parent is

$$\begin{aligned}
F=&\,Q+F_{\mathrm{graph}}\\
&+\frac{2W}{B}\big[v\xi\{(U-z)A_x+(V-y)B_x\}+u\eta\{(U-z)A_\eta+(V-y)B_\eta\}\\
&\qquad+uv\zeta\{(U-z)A_\zeta+(V-y)B_\zeta\}\big].
\end{aligned} \tag{DI5}$$

To see the ideal membership without an approximation, the respective linear coefficients divided by $v/B, u/B, uv/B$ are polynomials affine in V and at most quadratic in U. Their values at (z, y) vanish by the exact original state-kernel equations. Subtract their value first at $V = y$ and then at $U = z$; this gives exactly (DI4). The divisions by u and v in this explanatory derivation are polynomial factorizations; (DI4)–(DI5) themselves have no such divisions.

The angular scalar is an exact contact-ideal polynomial. Define positive source coefficients

$$\begin{aligned} p_2 &= r^2(D + n + \tfrac{1}{2}), \quad p_1 = 2Dr + nr^2 + \tfrac{1}{2}r^2, \quad P(U) = p_2U^2 + p_1U + D, \\ q_1 &= AH^2, \quad q_0 = AH^2 + (C + D + \tfrac{1}{2})L^2, \quad d_0 = mB^2, \quad a_y = nr^2 + yp_2, \\ K(U) &= yp_2(U + z) + yp_1 - q_1/y. \end{aligned}$$

The following identity is exact for the entire source chart, including the weak-α face and literal $n = 0$:

$$\boxed{\begin{aligned} F_{\mathrm{graph}} = \frac{4W^2}{B^2}\Big[& V(U - z)^2\Big(a_yU + \frac{d_0}{z^2}\Big) \\ & + U(V - y)^2P(U) + U(U - z)(V - y)K(U)\Big]. \end{aligned}} \tag{DI6}$$

For a short derivation, put $f = B^2F_{\mathrm{graph}}/(4UVW^2)$ formally. Its nine Laurent supports are

$$(2, 1), (2, 0), (1, 1), (1, 0), (1, -1), (0, 1), (0, 0), (0, -1), (-1, 0).$$

All coefficients other than $(1, 0)$ and the constant coefficient are nonnegative; their positive values are respectively $p_2, nr^2, p_1, q_1, D, q_0, d_0$. Thus

$$f = P(U)V + (q_1U + q_0)/V + nr^2U^2 + b_0'U + c_0' + d_0/U.$$

Exact contact and its two angular derivatives give $q_1z + q_0 = y^2P(z)$ and

$$f(U, y) = (U - z)^2\{a_y + d_0/(z^2U)\}.$$

Subtracting $f(U, y)$ proves (DI6). Multiplication by UV removes all testing inverses, so the boxed polynomial identity includes the literal testing axes. In particular its formal $V = y$ slice is SOS, but the mixed term in (DI6) and all three lines of (DI4) remain part of the independent-testing problem.

A full-parent bounded remainder on the strong face. On B-unit II, the source entry gives positive units

$$m/z^2, \quad A/y, \quad CL, \quad Dyj, \quad \alpha(1 - H), \qquad nr \text{ bounded}. \tag{DI7}$$

Here $H = B - L$ and $1 - r$ are units. The latter assertion follows from the exact numerator identity $E(1) = -(H + y)(B + z)(1 + z)$ for $E(r) = (B + zr)(1 + zr)(1 + zr + L) - r\delta$ and $E(r) \geq 0$. All claims in (DI7) retain arbitrary openings $j, 1 - H, r$.

Choose any bounded cap $0 \leq n_* \leq n$ and any positive real cap $0 < \alpha_* \leq \alpha$. Set

$$\begin{aligned} P_C &= 4CD_v\eta^2, \\ P_D &= 4D\{J + (yj/B)uW\}^2, \\ P_n &= 4(n - n_*)D_uD_v\xi^2, \\ P_\alpha &= 4(\alpha - \alpha_*)D_u(v\eta - \zeta)^2. \end{aligned} \tag{DI8}$$

Each displayed root vanishes at the actual contact. The exact identity $F = P_C + P_D + P_n + P_\alpha + G$ has a polynomial remainder G with bounded source coefficients; for bounded affine tests this is a bounded tensor polynomial. Indeed its four changed packets are exactly

$$\begin{aligned} G_C &= 4CD_v\{c_0^2W^2 - 2c_0W\eta\}, \\ G_D &= -8D(yj/B)uWJ - 4D(yj/B)^2u^2W^2, \\ G_n &= 4n_*D_uD_vX^2 + 4(n - n_*)D_uD_v\{x_0^2W^2 + 2x_0W\xi\}, \\ G_\alpha &= 4\alpha_*D_u(vY - Z)^2 + 4(\alpha - \alpha_*)D_u\{p_0^2W^2 + 2p_0W(v\eta - \zeta)\}. \end{aligned} \tag{DI9}$$

Add the unchanged $4mM^2 + 4AD_uD_vY^2$ and the entire signed core. Boundedness follows from (DI7), since the only potentially large products are CL, Dyj, nr, nr^2, $\alpha(1-H)$ and $\alpha(1-H)^2$. No coefficient inverse involving a test is taken. The graph raises the testing degree, so one must not identify its common higher-degree homogenization at a literal projective endpoint with the original parent's endpoint form. Endpoint joins remain part of any future all-tests admission. All removed packets have zero first jet, so G retains the exact contact and every first derivative. This proposition asserts boundedness, not positivity, of G.

The precise remaining obligation. Equations (DI3)–(DI6) expose the same contact ideal on both source faces and print every state normal. Equation (DI8) additionally removes all unbounded source coefficients on the strong face. A completion must allocate the mixed angular term in (DI6) together with all of (DI4), using retained packet roots or further admissible roots, and prove a strict bounded Gram or exact SOS for every independent testing rate. The factors $1/B$ and d_0/z^2 cannot be declared uniformly bounded on the weak face; its source normalization remains a separate task. Neither remaining checklist row is closed by the present reduction.

Chapter 65

C1 UI: a positive Laurent model and the remaining lift

Component `p050c7ae6d9b8`
Source: `R13/baseline/baseline/advances/c1_ui/leading_contact_frontier.tex`
SHA256: `050c7ae6d9b8b1ddb00388ac0efd965dba9855d1aab36301dc8ec51215531a11`
Inventory use: CL1–CL7 only, as exact formulae and real-residue calculations | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

65.1 C1 UI: a positive Laurent model and the remaining lift

This section concerns the leading three-state model. It is not yet a certificate for the exact parent at independent source and test rates. Write $x = k^2$, $r = 1 - a\eta$, $K = 1 + x$, $R = 1 + xr$ and

$$E_0 = 1 - x\eta, \quad J = K - a, \quad P_4 = KE_0 + x(1 - r), \quad P_d = 1 + 2x + rx^2.$$

At a real interior source point the leading coefficients of the parent divided by a are

$$\begin{aligned} h &= \frac{2x\eta}{RE_0}, & C &= \frac{2}{a}, & A &= \frac{2\eta^2 P_4}{Kr(r+\eta)E_0}, \\ d &= \frac{4\eta P_d}{K(r+\eta)E_0}, & D &= \frac{2}{JE_0} - \frac{d}{2}, & \alpha &= \frac{2x^2\eta R}{KE_0}. \end{aligned} \tag{CL1}$$

All are positive except $D \geq 0$, including its literal zero. Here α is the coefficient after extracting the weak g^2 scale. For a test (u, V) put $D_u = 1 + u^2$ and define

$$\begin{aligned} f = {} & 4hV^2(W + uY)^2 + 4C(uW - Y)^2 + 4AD_uY^2 \\ & + (4D + 2d)(uW - Y - VD_uZ)^2 + 4\alpha D_uZ^2 \\ & - 4dD_uVYZ - 2dD_uV^2Z^2. \end{aligned} \tag{CL2}$$

Let M_f be its symmetric three-state matrix and set $U = u^2$, $T = V^2$. Direct collection gives

$$\det M_f = 64D_u^2 UT\,\mathcal{L}(U, T), \tag{CL3}$$

where

$$\begin{aligned} \mathcal{L} = {} & AC(D + d/2)U + \frac{h\alpha(A + C + D + d/2)}{U} + \frac{A\alpha(C + D + d/2)}{T} \\ & + hD(A + C + d/2)\frac{T}{U} + hC(D + d/2)UT \\ & + h(AD + 2CD + Ad/2 + Cd/2 + Dd/2 + d^2/4)T + L_0, \\ L_0 = {} & ACD - ADd/2 + h\alpha(C + D + d/2) - d^2(A + C + D)/4 - d^3/8. \end{aligned} \tag{CL4}$$

Every nonconstant coefficient is nonnegative. The coefficients of U, U^{-1}, T, T^{-1} are strictly positive, also when $D = 0$. Thus in coordinates $(\log U, \log T)$ the Hessian is strictly positive and the function is coercive. Substitution of (CL1) gives

$$\mathcal{L}(x, x\eta) = \partial_U \mathcal{L}(x, x\eta) = \partial_T \mathcal{L}(x, x\eta) = 0. \tag{CL5}$$

Strict convexity proves that $\mathcal{L}$ is positive away from this single contact. The (W, Y) principal block of (CL2) is strictly positive for $U, T > 0$, because its C and A terms already have trivial common kernel. Hence f is positive semidefinite everywhere and positive definite off the contact locus. Literal endpoints follow by continuity. This proof uses no numerical determinant table.

There is a useful corrected formula at the large-D boundary. Impose its hard equation $uW - Y - VD_uZ = 0$, eliminating W. The resulting two-state determinant is $16D_u^2 T\mathcal{L}_D$, where

$$\begin{aligned}\mathcal{L}_D = {} & hT\{(C + A + d/2)/U + (2C + A + d/2) + CU\} \\ & + h\alpha(1 + 1/U) + AC(U + 1) + A\alpha/T - Ad/2 - d^2/4.\end{aligned} \tag{CL6}$$

In particular the coefficients of T/U and UT are $h(C + A + d/2)$ and hC, respectively. Interchanging these coefficients is incorrect. The accompanying checker verifies (CL3), (CL5) and (CL6) by exact rational identities.

The remaining exact lift. For the original source let $\zeta = g^2$, $p = t/R$, and $Q = W + uY$, $B = X + uZ$. The two strong-event normals may be written

$$N = (B - pvQ)/g, \qquad L = vQ/g.$$

Put

$$h_2 = \frac{2x\eta(1 - tg^2)(1 + x\eta g^2)}{E}, \qquad h_\beta = \frac{2x\eta[1 + (1 - t)g^2]}{E}, \qquad h_\beta - h_2 = 2x\eta g^2.$$

After retaining the exact positive packets $4h_2 t\, N^2/(MR)$ and $4h_\beta M D_v N^2/(tR)$, their remaining quadratic is

$$\frac{4}{R}\{(h_2 + h_\beta D_v)N^2 + 2(h_\beta D_v - h_2)LN + [h_2(1 - p) + h_\beta p D_v]L^2\}. \tag{CL7}$$

This is a promising bounded full-state starting point. To complete the ordinary/small testing strip one still must do all of the following: retain the centered excess C, A, D packets with their actual centers; print every first-normal coefficient after that subtraction; lift all contact-vanishing scalar and normal terms to admissible roots; and prove strictness uniformly when a, r or J tends to zero. The leading determinant and contact Hessian alone do not discharge these obligations. No all-tests C1 UI admission is recorded here.

Chapter 66

C1 UI: a full-parent remote complement

Component p013c226a79b3
Source: R13/baseline/baseline/advances/c1_ui/remote_complement.tex
SHA256: 013c226a79b3a1076d4bf4810beef1da98031c61b23c61804687ce57d17a3d8c
Inventory use: CR1–CR13, including CR5a | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

66.1 C1 UI: a full-parent remote complement

The source has actual ordinary Alice angle and infinitesimal Bob angle. Apply the exact native stabilizer (AS5) as in (NI7) and take η a positive bounded unit, $\zeta = g^2$ infinitesimal and k a bounded unit. The reflected orientation need not satisfy $k \le 1$. Its E remains a positive unit by the exact source entry in (NI7). Normalize by the positive pure-source d. Equations (NC2)–(NC4) give

$$C/d,\ A/d,\quad g^2m/d,\quad g^2\beta/d,\quad \alpha/(dg^2)\quad \text{strong},\qquad D/d \ge 0. \tag{CR1}$$

For example $P_d, P_\alpha, Q, r+\eta$ are units, whereas the positive openings $M, t, r, 1-r$ are bounded and may be infinitesimal. The $-1/2$ in β/d disappears after multiplication by g^2. In this particular source subdomain $\beta = 0$ is impossible; $D = 0$ is allowed throughout. Choose positive real constants μ, ν, c, a, b below the five quantities in (CR1), respectively in the order $g^2m/d, g^2\beta/d, C/d, A/d, \alpha/(dg^2)$. Whenever needed we cap these coefficients at

$$m = \mu/g^2,\quad \beta = \nu/g^2,\quad C = c,\quad A = a,\quad \alpha = bg^2. \tag{CR2}$$

Every unused packet, every positive excess, and the complete D packet is retained as a square. The signed core, divided by two, is

$$K = J^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_uD_vZ^2,\quad J = uW + uvX - Y - vZ. \tag{CR3}$$

Only the core and the explicitly selected caps need a bounded Gram. This observation keeps all upper coefficient scales out of the proof.

Theorem 66.1 (C1 UI remote tests)**.** *For every source satisfying (CR1), the complete original C1 parent is tensor SOS at every independent testing pair except possibly u a bounded nonzero unit and v infinitesimal. The conclusion includes literal projective endpoints, both signs, all states, and $D = 0$.*

Proof. All constants in residue forms below are real residues. Roots may be unbounded, but all coefficients in their exact Gram are bounded. First put $Q = W + uY$, $B = X + uZ$; then

$$M = vQ - B,\quad uW - Y = uQ - D_uY,\quad J = uQ - D_uY + uvB - vD_uZ. \tag{CR4}$$

These identities account for every state normal.

If u is a bounded unit and v projectively infinite, put $w = 1/v$ and homogenize by w^2. Use roots $(E_Q, E_B, Y, Z) = (Q/g, B/g, Y, Z)$ and cap only m, β. The exact homogenized core is

$$[w(ugE_Q - D_uY) + ugE_B - D_uZ]^2 + D_u(Y - wZ)^2 + D_w g^2 E_B^2 - D_u D_w Z^2.$$

Its residue is $D_u(Y^2 + u^2Z^2)$. Including the two caps gives

$$4\mu E_Q^2 + 4\nu E_B^2 + 2D_u(Y^2 + u^2Z^2) > 0. \tag{CR5}$$

If u is infinitesimal and v projectively infinite, the preceding four-root frame loses the Z pivot. Keep $w = 1/v$ and instead use (E_Q, E_B, Y, U, T), where $U = uZ$ and $T = gwZ$. Put $A_0 = w(ugE_Q - D_uY) + ugE_B$. The exact homogenized core is

$$K_w = A_0^2 + D_uY^2 + D_w g^2 E_B^2 + D_uU^2 - 2D_u g(wE_Q + E_B)U + 2wD_u uYU.$$

All its coefficients are bounded, with residue $Y^2 + U^2$. Cap m, β, α; their amplitudes after normalization are $E_Q - wE_B$, E_B, and $gY - T$, with the positive factors D_w, D_u in the last two packets. The full selected residue is

$$4\mu E_Q^2 + 4\nu E_B^2 + 2Y^2 + 2U^2 + 4bT^2 > 0. \tag{CR5a}$$

This includes literal $u = 0$ or $w = 0$; no division by either was used.

If u is infinitesimal and v a bounded unit, use $E_Q = Q/g, E_B = B/g, Y, U = uZ, Z_g = gZ$, and cap m, β, α. Substitute $W = gE_Q - uY$, $X = gE_B - uZ$ in (CR3). All terms involving Z collect into uZ; the resulting exact core is a polynomial in $(u, v, g; E_Q, E_B, Y, U)$. Its residue is $D_vY^2 + v^2U^2$. The full selected residue is

$$4\mu(vE_Q - E_B)^2 + 4\nu D_v E_B^2 + 2D_vY^2 + 2v^2U^2 + 4bZ_g^2 > 0. \tag{CR6}$$

The α amplitude is $gvY - Z_g$, so no inverse of u was used in constructing this Gram.

If both u, v are infinitesimal, set

$$U = uQ, \quad V = vQ/g, \quad E_B = B/g, \quad Z_g = gZ, \quad S = uvZ,$$

and include Y as the sixth root. The exact core is

$$\begin{aligned} K = {} & (U - D_uY + uvgE_B)^2 + D_u v^2 Y^2 + D_v g^2 E_B^2 + D_u S^2 \\ & + 2D_u uYS - 2D_u uVZ_g - 2D_u vgE_B S. \end{aligned} \tag{CR7}$$

In particular the apparently problematic term UvZ is exactly uVZ_g, with no mixed inverse. Cap all five weights in (CR2). The entire selected residue is

$$4\mu(V - E_B)^2 + 4\nu E_B^2 + (4c + 2)(U - Y)^2 + 4aY^2 + 4bZ_g^2 + 2S^2 > 0. \tag{CR8}$$

It remains to cover u projectively infinite. Put $p = 1/u$, homogenize by p^2, and use $D_p = 1 + p^2$, $Q = pW + Y$, $B = pX + Z$. Then

$$M = vQ - B, \quad r = W - pY, \quad J = D_p(W + vX) - p(Q + vB). \tag{CR9}$$

For v a bounded unit, use $(Q/g, B/g, W, X)$ and cap m, β, C. The core after $Y = gE_Q - pW$, $Z = gE_B - pX$ is a bounded polynomial with residue $(W + vX)^2$. The selected residue is

$$4\mu(vE_Q - E_B)^2 + 4\nu D_v E_B^2 + 4cD_v W^2 + 2(W + vX)^2 > 0. \tag{CR10}$$

For infinitesimal v, use (W, R, Y, V, E_B) with $R = vX$, $V = vQ/g$, $E_B = B/g$. Cap m, β, C, A, retaining α entirely. The exact core is

$$K_p = (W + D_pR - pY - pvgE_B)^2 + D_v g^2 E_B^2$$

$$+ D_p\{v^2Y^2 - 2vgYE_B + 2pYR - (vgE_B - pR)^2\}. \tag{CR11}$$

Every coefficient is bounded. The selected residue is

$$4\mu(V - E_B)^2 + 4\nu E_B^2 + 4cW^2 + 4aY^2 + 2(W + R)^2 > 0. \tag{CR12}$$

Finally, when both testing angles are infinite, also put $w = 1/v$ and homogenize by p^2w^2. Use $(Q/g, B/g, W, X)$ and cap m, β, C. Substitution $Y = gE_Q - pW$, $Z = gE_B - pX$ in the homogenized core is polynomial and has residue X^2. The selected residue is

$$4\mu E_Q^2 + 4\nu E_B^2 + 4cW^2 + 2X^2 > 0. \tag{CR13}$$

Each displayed positive form is strict on the complete root space. The bounded positive-residue Gram lemma therefore supplies tensor SOS. Restoring the retained packets proves the original parent. The bounded unit/unit testing chart is already covered by the inherited noncommuting-residue theorem. This leaves exactly the ordinary/small testing strip stated in the theorem. The exact gauges $(u, W, X) \mapsto (-u, -W, -X)$ and $(v, W, Y) \mapsto (-v, -W, -Y)$ preserve every event square and the core. All homogeneous formulas allow $p = 0$ or $w = 0$ literally. Real and imaginary state components are treated separately. □

The checker verifies the exact core reductions and their residues; it does not infer positivity from samples. This theorem narrows the actual UI subcase of C1.5. It does not close C1.5, whose actual UH source subcase remains, and it does not yet close UI's ordinary/small strip. The leading-contact calculations stored alongside this proof remain candidate material until their complete normal lift is printed.

Chapter 67

Exact source entry for the strong HI corner

Component `p54b8f20ad2dd`
Source: `R13/baseline/baseline/advances/hi/exact_source_entry.tex`
SHA256: `54b8f20ad2dd194d0e2ed7f5124fa1cc688d91a27ff328d503dc53b0eb45bd9b`
Inventory use: Exact HI entry from n>=0, q,yInf and actual gaps; all four event weights unbounded

67.1 Exact source entry for the strong HI corner

Use the original positive rational C2 source with $d = 1$, actual u_c unbounded and v_c infinitesimal. Put

$$q = u_c^{-2}, \quad y = v_c^2, \quad H = u_c^2 h \in (0,1), \quad R = r/q, \quad B = b/q, \quad L = yt = B - H, \quad T = B - yR.$$

The already proved source identities (WE1)–(WE3) show that R is bounded using only $n \geq 0$, q infinitesimal and y bounded. Here is their short contradiction argument, to make this use explicit. The positive source gaps imply

$$0 < T < H + y, \quad g = H + (1+y)r - 1 + qT > 0,$$

and $n = -qP/(2rQ)$ with $Q > 0$ and

$$P \geq r^2(1+y)(H+yr) - Tqr(y+2) - q^2(T^2+T).$$

If q/r were infinitesimal, then $T/(H+yr)$ and $(T^2+T)/(H+yr)$ would be bounded. For unit r this follows from $T < H + y \leq (H+yr)/r$; for infinitesimal r, the inequality $g > 0$ forces H to have residue one. Division by $r^2(H+yr)$ would give $P > 0$, contrary to $n \geq 0$. Thus R is bounded. (The exact positive remainder in this lower bound is printed in (WE3), so this argument uses no asymptotic sign substitution.)

Now $r = qR$ is infinitesimal and $g > 0$ gives $H_0 = 1$. Define the positive opening

$$E = (1 - H)/q.$$

The same gap implies $E < (1+y)R + T$, so E is bounded. Since $j = 1 + R - t > 0$, t is bounded; consequently

$$H = 1 - qE, \quad B = H + yt \sim 1, \quad j = 1 + R - t, \quad K = B + R - E = g/q > 0.$$

All of E, t, j, K are positive bounded, while $t + j = 1 + R$ and $E + K = B + R$ are positive units.

Let Δ and N denote the unchanged original source polynomials, and set $\mathcal{N} = N/q$. Their exact expressions are

$$\Delta = (1+q)\{B^2 + 2BR + (1+y)B\} + (1-yq)R^2 + 2(1+y)R + 1 + y,$$

$$\mathcal{N} = (1+q)(B+R)(H+y) - (1+y)(1-qR)yt.$$

Thus Δ is a positive bounded unit and

$$\bar{\Delta} = (1+\bar{R})(3+\bar{R}), \qquad \overline{\mathcal{N}} = 1+\bar{R} > 0.$$

Set $S = 1+R+yt$, also a positive unit. Direct substitution into the original coefficients gives the following exact equalities:

$$\begin{aligned} mqK &= \frac{(1+y)(1-qR)(1+R)S}{2\Delta}, & Cyt &= \frac{(1+R)\mathcal{N}}{2\Delta}, \\ \frac{A}{qy} &= \frac{(1+R)(B+R)S}{2H\Delta}, & Dyj &= \frac{(1+q)(B+R)(B-yR)S}{2\Delta}, \\ \alpha qE &= \frac{(B+R)\mathcal{N}}{2\Delta}, & n &= \frac{(B+R)(1+R)S - R\Delta}{2R\Delta}. \end{aligned} \tag{HI-entry}$$

Every one of the first five right-hand sides is a positive bounded unit; in particular $B-yR$ has residue one. The last numerator has residue $(1+\bar{R})(1-\bar{R})$, so $n \geq 0$ implies $0 \leq \bar{R} \leq 1$. This does not imply $R \leq 1$ as a source inequality, and no such stronger claim is used.

Since q, y are infinitesimal and E, t, j, K bounded, the positive numbers qK, yt, yj, qE are infinitesimal. Dividing their unit products in (HI-entry) proves

$$m, C, D, \alpha \quad \text{are all unbounded}, \qquad A \asymp qy.$$

This conclusion follows from the original source formulas and $n \geq 0$; no additional strong-weight assumption is needed for an actual HI source.

Chapter 68

A full-parent remote theorem for strong HI sources

Component `pcee1372511ed`
Source: `R13/baseline/baseline/advances/hi/remote_all_parent.tex`
SHA256: `cee1372511ed2fb356229e5f15bf95a827ce2829e9f8a51a88fe07db4fe72c28`
Inventory use: Every remote original parent state/angle chart and literal critical axis; only nonzero reciprocal-u and vInf corner remains

68.1 A full-parent remote theorem for strong HI sources

Theorem 68.1 (All tests outside the HI critical corner)**.** *Let K_c, K_t be independent real closed extensions of $\mathbb{R}$. Consider the original C2 parent with $d = 1$, $\lambda = \beta = 0$, $n, A \geq 0$, and m, C, D, α positive unbounded source numbers. Then at every projective testing pair except possibly $1/u$ and v both nonzero infinitesimal, and every real or complex testing state, the original parent is a sum of squares in $K_c \otimes_{\mathbb{R}} K_t$. No comparison of source and testing rates is made.*

Proof. For any positive real μ, retain the four nonnegative event packets with coefficients $m-\mu, C-\mu, D-\mu, \alpha-\mu$, and retain the n and A packets in their entirety. It suffices to prove the capped parent obtained by setting $m = C = D = \alpha = \mu$ and $n = A = 0$ is tensor SOS at the given test. The real constant μ may be chosen after the real testing residue is fixed, since each of the original four source coefficients is larger than every real number.

Write $D_u = 1 + u^2$, $D_v = 1 + v^2$. First suppose v is infinitesimal, including zero. Use the roots

$$S = vW, \qquad E = Y - uW, \qquad X, \quad Z.$$

All four capped event amplitudes are bounded linear forms in these roots:

$$M = D_u S - X - uZ + uvE, \quad uW - Y = -E, \quad J = -E + uvX - vZ, \quad vY - Z = uS + vE - Z.$$

The remaining core depends only on the same four roots, so the exact capped parent has a bounded four-root Gram. Let $a = \bar{u}$ and $d_a = 1 + a^2$. Its residue is

$$(8\mu + 2)E^2 + 4\mu(d_a S - X - aZ)^2 + (4\mu + 2)d_a(aS - Z)^2 + 2(X + aZ)^2 - 2d_a Z^2.$$

Put $T = X + aZ$ and $R = aS - Z$; this is an invertible real change of coordinates. The last three-variable form is

$$4\mu(d_a S - T)^2 + 4\mu d_a R^2 + 2T^2 - 2d_a a^2 S^2 + 4d_a aSR.$$

The T and R pivots are $4\mu + 2$ and $4\mu d_a$. The remaining Schur pivot is exactly

$$\frac{d_a[4\mu^2 - a^2(4\mu + 1)]}{\mu(2\mu + 1)}.$$

Taking $\mu = 1 + a^2$ makes its numerator positive, because $4\mu^2 - a^2(4\mu + 1) = 3a^2 + 4$. Thus this bounded Gram has strictly positive real residue and is tensor SOS by the inherited strict-residue Gram lemma.

Next suppose v is a bounded unit. At the real residues $(a, b) = (\bar{u}, \bar{v})$ the four event forms have common kernel

$$(W, X, Y, Z) = W(1, b, a, ab).$$

The core restricted to this line is $2(1 + a^2)(1 + b^2)b^2W^2$, which is strictly positive away from zero. Consequently a sufficiently large positive real μ makes the whole capped residue form positive definite. For completeness, this elementary assertion has a direct quantitative proof. Let $E_0 \succeq 0$ be the event Gram, split the state space into $\ker E_0$ and its orthogonal complement, and let $\sigma > 0$ be its smallest eigenvalue on that complement. Let $\kappa > 0$ be the core's eigenvalue on the one-dimensional kernel and bound the norms of the two remaining core blocks by M. Then $4\mu\sigma > M + M^2/\kappa$ gives a positive Schur complement. Every constant here belongs to $\mathbb{R}$. The exact state Gram is bounded and has this positive residue.

Finally suppose v is unbounded, including the projective endpoint. Set $w = 1/v$, homogenizing before setting $w = 0$. The four event forms become

$$W - wX + uY - uwZ, \quad uW - Y, \quad uwW + uX - wY - Z, \quad Y - wZ,$$

with bounded positive factors D_w or D_u where appropriate. At $w = 0$ their common kernel is $(W, X, Y, Z) = (0, X, 0, aX)$, on which the homogenized core is $2(1 + a^2)X^2$. The same real Schur argument gives a positive residue Gram for a sufficiently large μ. For finite unbounded v, undo homogenization by the pure positive testing square v^2; for the projective endpoint the homogeneous expression is already the required parent. These three cases exhaust every projective Bob test when u is bounded.

It remains to consider $p = 1/u$ infinitesimal, after homogenizing by p^2. If v is a bounded unit, use the four roots

$$S = pY, \qquad T = vY - Z, \qquad W, \quad X.$$

The exact event amplitudes are

$$W - S, \quad J = W + vX - D_vS + pvT, \quad M = pvW - pX + T, \quad T.$$

The signed core also has bounded coefficients in these roots:

$$2J^2 + 2D_pT^2 + 2D_v[p^2X^2 + 2vSX - 2pXT - v^2S^2 + 2pvST - p^2T^2].$$

At $p = 0$, set $b = \bar{v} \neq 0$, $A_0 = W - S$, and $B_0 = X - bS$. The full capped residue is

$$(8\mu + 2)T^2 + 4\mu(1 + b^2)A_0^2 + (4\mu + 2)(A_0 + bB_0)^2 + 4(1 + b^2)bSB_0 + 2(1 + b^2)b^2S^2.$$

As μ grows, the first three positive penalties constrain $T = A_0 = B_0 = 0$, and the residual kernel has positive coefficient $2(1 + b^2)b^2S^2$. The quantitative real Schur argument above therefore gives a finite μ with a strictly positive residue.

If u, v are both projectively unbounded, put $w = 1/v$ and homogenize by p^2w^2. Use $P = Y - wZ$, $S = pZ$, W, X. Now the exact event amplitudes are

$$M = pW - pwX + P, \quad W - pP - wS, \quad J = wW + X - pwP - D_wS, \quad P.$$

The exact core is

$$2J^2 + 2D_pP^2 + 2D_w[p^2X^2 + 2SX - S^2].$$

At $p = w = 0$ the capped residue is exactly

$$4\mu W^2 + (8\mu + 2)P^2 + 4\mu(X - S)^2 + 2X^2,$$

which is strict for every $\mu > 0$. The literal projective endpoints cause no division. The literal critical axes also have direct exact certificates:

$$F|_{p=0} = 4nD_vX^2 + (4m + 4\alpha + 2)(vY - Z)^2 + 4CD_vW^2 + 4AD_vY^2 + (4D + 2)(W + vX)^2,$$
$$F|_{v=0} = 4nD_pX^2 + (4m + 2)(pX + Z)^2 + (4C + 4D + 2)(W - pY)^2 + 4AD_pY^2 + 4\alpha D_pZ^2.$$

Thus only p, v both nonzero infinitesimal remain untreated. Applying the real-state result separately to real and imaginary state components proves the complex-state claim. □

For a strong C2 HI source, the exact source-entry lemma below proves that all four coefficients in the theorem are unbounded. Hence the theorem applies without any additional source-shape assumption. Its precise uncovered complement is the critical testing corner $1/u$ and v both nonzero infinitesimal. Exact sign gauges may make both positive. The original HI source leaf is therefore narrowed but not closed.

Chapter 69

Weak-C IH: the open testing axis

Component `pf95b1c5fc32f`
Source: `R13/baseline/baseline/advances/ih/root/weak_ih_open_axis.tex`
SHA256: `f95b1c5fc32fc0fe354c7c23bcc642ba08597a5178297f8cdcf2898f01ae9e0d`
Inventory use: OA1–OA9 missing open-axis full product-pivot construction, including both rho endpoints

69.1 Weak-C IH: the open testing axis

This argument is reconstructed on 19 October 2026. Its algebraic identities are replayed by `check_weak_ih_open_axis.py`; it does not rely on an unrecovered historical verification receipt.

Domain and source input. The source and testing real closed fields are independent. Let B_c, B_t be their rings of bounded elements and use $B = B_c \otimes_{\mathbb{R}} B_t$ for Gram coefficients. Source square roots and testing square roots are always taken in their respective fields. In the weak-C IH source chart, write $z = u_c^2$ infinitesimal, $y = v_c^2$ unbounded, $H = zh \in (0,1)$, $B_* = zb$ unbounded and $\rho = B_*/y$ bounded. The source entry calculation gives

$$0 \le \rho_0 \le 1, \qquad (B_*C)_0 = \left(\frac{AH}{B_*}\right)_0 = \left(\frac{\alpha(1-H)}{B_*}\right)_0 = c := \frac{1}{2(1+\rho_0)}. \tag{OA1}$$

In particular C is infinitesimal and A, α are unbounded. For $0 < \rho_0 < 1$,

$$D_0 = \frac{\rho_0^2}{2(1-\rho_0^2)}; \tag{OA2}$$

for $\rho_0 = 0$, D is positive infinitesimal; for $\rho_0 = 1$, D is unbounded. These statements follow from $D \sim \rho^2/[2e(1+\rho)]$ and $e_0 = 1 - \rho_0$. No lower bound on $n \ge 0$ is imposed.

Statement. For every testing u with nonzero finite residue and every infinitesimal $w = 1/v$, including $w = 0$, the complete reciprocal-v canonical C2 parent is a tensor sum of squares for every state. The same assertion holds for complex states by applying the real quadratic form to real and imaginary parts separately.

Exact parent and retained squares. Put $D_u = 1 + u^2$, $D_w = 1 + w^2$ and

$$M = W - wX + uY - uwZ, \qquad J = uwW + uX - wY - Z.$$

The parent is

$$\begin{aligned}\widehat{F} = {} & 4nD_uD_wX^2 + 4mM^2 + 4CD_w(uW - Y)^2 + 4AD_uD_wY^2 + 4DJ^2 + 4\alpha D_u(Y - wZ)^2 \\ & + 2\{J^2 + D_u(Y - wZ)^2 + D_w(X + uZ)^2 - D_uD_wZ^2\}.\end{aligned}$$

Retain the entire n, m packets. When $\rho_0 = 0$, retain the entire D packet as well and set $d = 0$. When $0 < \rho_0 < 1$, set $d = D$. When $\rho_0 = 1$, choose a sufficiently large positive real d below D and retain $4(D-d)J^2$. Denote the remainder by F_*. All retained terms are tensor squares or sums of tensor squares.

Use the graph $X = 0$, $Y = HwZ$, and define

$$P = \frac{uW}{\sqrt{B_*}}, \quad E = \sqrt{B_*}\, wZ, \quad V = uwW, \quad E_Y = \sqrt{\frac{B_*}{H(1-H)}}(Y - HwZ). \tag{OA3}$$

Crucially, the exact root relation is

$$PE = VZ. \tag{OA4}$$

On the graph, the C, A, α, d packets are respectively

$$\begin{aligned} &4B_*CD_w(P - HE/B_*)^2, \\ &\frac{4D_u}{B_*}\{AH^2D_w + \alpha(1-H)^2\}E^2, \\ &4d\{V - (1+Hw^2)Z\}^2. \end{aligned}$$

The remaining core is exactly

$$2V^2 - 4(1+Hw^2)PE + \frac{2}{B_*}\{H^2 + u^2(1-H)^2 + H^2w^2\}E^2. \tag{OA5}$$

Thus every graph coefficient is bounded. The last E^2 coefficient is infinitesimal, uniformly for all $0 < H < 1$.

Restoring the Y normal. Let $L = B_*/[H(1-H)]$ and $Y_* = HwZ$. The first normal, divided by four, is

$$\begin{aligned} L_Y := \frac{1}{4}\partial_Y F_*|_{X=0,Y=Y_*} = {} & 2CD_w(Y_* - uW) + 2D_u\{AHD_w - \alpha(1-H)\}wZ \\ & - 2dw\{V - (1+Hw^2)Z\} - w\{V - (1+Hw^2)Z\} + D_u(H-1)wZ. \end{aligned} \tag{OA6}$$

Every coefficient of $L_Y/\sqrt{L}$ in (P, E, V, Z) is infinitesimal when d is bounded, including $d = 0$. Indeed the only potentially finite contribution is

$$2D_u\frac{AH - \alpha(1-H)}{B_*}\sqrt{H(1-H)}E,$$

and it has zero residue by (OA1); the additional w^2 contribution is infinitesimal. The other terms contain C, $w/\sqrt{L}$ or $1/B_*$. The quadratic E_Y coefficient has residue $4cD_{u,0} > 0$, and the XE_Y coefficient has zero residue. These statements remain valid when H or $1-H$ is infinitesimal. No inverse of d, H or $1-H$ enters a Gram coefficient unaccompanied by its compensating factor.

The strict residue when $0 < \rho_0 < 1$. Write $s = \sqrt{1+u_0^2} > 1$ and $r = \rho_0$. Using (OA4), split the core cross term as $-4PE = -4aPE - 4(1-a)VZ$, where a is any real number satisfying

$$\frac{s}{s+r} < a < \min\left\{\frac{s}{1+r}, \frac{s}{s-r}\right\}. \tag{OA7}$$

The interval is nonempty: both upper endpoints exceed $s/(s+r)$. The residue Gram splits into the positive E_Y pivot, a (P, E) block

$$G_{PE} = \begin{pmatrix} 2/(1+r) & -2a \\ -2a & 2s^2/(1+r) \end{pmatrix},$$

and a (V, Z, X) block

$$G_{VZX} = \begin{pmatrix} 2 + 4d_0 & -4d_0 - 2(1-a) & 2u_0(1+2d_0) \\ -4d_0 - 2(1-a) & 4d_0 & -4d_0u_0 \\ 2u_0(1+2d_0) & -4d_0u_0 & 2s^2 + 4d_0u_0^2 \end{pmatrix}. \tag{OA8}$$

The (V, X) principal block is positive definite. Its Schur complement is positive exactly when

$$\frac{s}{s+r} < a < \frac{s}{s-r},$$

as direct substitution of $d_0 = r^2/[2(1-r^2)]$ shows. Hence (OA7) makes both blocks strictly positive. This checks the entire X normal, not merely the scalar graph.

The endpoint $r = 1$. For finite $d > 0$, positivity of the (V, Z, X) block is equivalent to $a_- < a < a_+$, where

$$a_\pm(d) = \frac{(2d+1)s^2 \pm s\sqrt{2d(2d+1)}}{(2d+1)u_0^2 + 1}.$$

As $d \to \infty$, the lower endpoint tends to $s/(s+1)$ and the upper endpoint to $s/(s-1)$. Choose first a real a strictly between $s/(s+1)$ and $\min\{s/2, s/(s-1)\}$, then a sufficiently large finite d. The interval is nonempty because $s > 1$. Since source D is unbounded this cap is admissible, and the full residue is strict.

The endpoint $r = 0$. Take $a = 1$, $d = 0$, having retained the whole D packet, and omit Z from the root list. The residue is the direct sum of

$$2(P-E)^2 + 2u_0^2E^2, \qquad 2(V + u_0X)^2 + 2X^2, \qquad 2D_{u,0}E_Y^2. \tag{OA9}$$

It is strictly positive. The exact rewriting has bounded coefficients: (OA5) uses only P, E, V, and the first X normal from the core is $uV + u(1-H)w^2Z = uV + u(1-H)wE/\sqrt{B_*}$. Thus no unscaled Z or inverse of D remains.

Conclusion. In all three cases the original remainder has a bounded Gram matrix with strictly positive real residue. The bounded-Gram lifting lemma therefore makes it a tensor sum of squares. Adding the retained source-positive packets proves the assertion. The argument is valid for arbitrary $n \geq 0$, including $n = 0$, and literal projective $w = 0$; no comparison between source and testing infinitesimals was used.

Chapter 70

Strong IH: the complete local corner

Component `pf3e0ca8bf054`
Source: `R13/baseline/baseline/advances/ih/strong/ih_local_corner.tex`
SHA256: `f3e0ca8bf0547150036dcd1c624a329da3804b1a781c24835343af7acd1ba115`
Inventory use: SL1–SL5 strong-IH full scalar/current/normal construction on its own source bounds

70.1 Strong IH: the complete local corner

Assume the strong IH source entry of §72.1: z is infinitesimal, y unbounded, $B = H + L$ a bounded positive unit, $0 < H < 1$, $L > 0$, and C, α strong. The source parameter $\rho = B/y$ is infinitesimal, $s = r/\rho$ bounded, and $c_s = (1 + L)/B$ is a positive unit. Testing $u, w = 1/v$ are infinitesimal or zero. We use the exact graph, five roots and both full normal identities (IC3)–(IC12) of the preserved corner calculation; none of those algebraic identities assumes C weak.

The source quantities obey (GE3), and direct rational simplification gives additionally

$$\frac{k}{B\rho} \longrightarrow \frac{2(B^2 s + BHs + BH - Bs - H^2 + H)}{B^2}. \tag{SL1}$$

No opening denominator occurs in this normalized expression.

Set $T_\rho = \rho T$ and $\eta = \sqrt{H(1-H)/B}$, so $Y = HwZ + \eta E_Y$. The full Gram's entire explicit dependence on C is exactly

$$4C(U_1^2 + U_2^2), \quad U_1 = uW - Y - LwZ = \sqrt{B}R_0 - \eta E_Y,$$

$$U_2 = w(uW - Y) - (L/y)Z = R_2 - \sqrt{H/B}E - w\eta E_Y. \tag{SL2}$$

The scalar and normal identities verifying this assertion use

$$\sqrt{H}R_2 - \sqrt{BH}\, wR_0 - \sqrt{B}E = 0. \tag{SL3}$$

Choose a positive real cap $c_* \le C$ and retain $4(C - c_*)(U_1^2 + U_2^2)$. In the remaining exact Gram replace explicit C by c_*, leaving k and all other original source quantities unchanged. The distinction matters: this is an exact square subtraction from the complete Gram, not a perturbation of the source. Add λ times the exact zero (WIE2), where $\lambda > 0$ is a real number to be chosen. All resulting coefficients are bounded, using (GE3), (SL1), and the opening-normalized roots.

We display the entire residue. The roots R_1 and E_X have isolated positive pivots $2c_s$ and 1. The (R_0, E_Y) block is

$$\begin{pmatrix} 4Bc_* & -4c_*\sqrt{H(1-H)} \\ -4c_*\sqrt{H(1-H)} & (4c_* + 4A + 4\alpha + 2)H(1-H)/B \end{pmatrix}. \tag{SL4}$$

Here and below all expressions in a residue matrix mean their bounded residues; products such as AH are evaluated by (GE3), even when A itself is unbounded. Its determinant is exactly $8c_*(1+L) > 0$. Put $a_* = 2(2c_* + 1)$. The (R_2, E) block is

$$\begin{pmatrix} a_* & -a_*\sqrt{H/B} \\ -a_*\sqrt{H/B} & H(4A + a_*)/B \end{pmatrix}, \tag{SL5}$$

with determinant $2a_*c_s > 0$. The only remaining coupling is between R_2 and T_ρ, whose Gram entry is $c_s(H/B - s)$. The T_ρ diagonal is $\lambda + 2c_s s$. Choose λ larger than the finite Schur-complement threshold of the positive block (SL5). This produces a strictly positive residue matrix. Since this choice is made after taking a real residue, it is a real scalar and introduces no mixed inverse.

We spell out why no normal term is hidden in this block description. In (IC11), f_X/ρ is bounded by (GE3) and every scalar-root multiplier contains u or w. In (IC12), the only nonzero limiting term is the explicit C contribution to R_0E_Y, already included in (SL2)–(SL4). After the cap, all other normal multipliers contain z, u, or w. The unbounded factors A and α always occur as AH and $\alpha(1 - H)$ after normalization, hence remain bounded also at $H_0 = 0, 1$. The normal X coefficient divided by $N = 4n + 2$ has residue one, and its coupling to E_Y vanishes. The remaining Y coefficient is exactly the lower-right entry of (SL4). These facts follow directly from the displayed exact normal factorizations, not from scalar-graph positivity.

The bounded positive-residue Gram lemma now gives tensor SOS for the complete original parent, after restoring the retained positive $C - c_*$ squares. This holds for all states and all independent local tests, for both opening endpoints, for arbitrary $n \geq 0$, and at literal $u = 0$ or $w = 0$.

Chapter 71

Strong IH: remote charts and the all-tests join

Component `p7405dbf89341`
Source: `R13/baseline/baseline/advances/ih/strong/ih_remote_charts.tex`
SHA256: `7405dbf893412ea0c95351f3486db35723984e7e5d65073868a21718b179af48`
Inventory use: SR1–SR8 all remaining testing/state charts, with detailed A6 final noncommuting join

71.1 Strong IH: remote charts and the all-tests join

Use the source entry of §72.1. In particular $B = H + L$ is a bounded positive unit, m, D are infinitesimal, C, A, α are strong, and the bounded products satisfy

$$CL \to \tfrac{1}{2}, \qquad AH \to (1+L)/2, \qquad \alpha(1-H) \to B/2. \tag{SR1}$$

Whenever an event coefficient is unbounded, a positive finite real cap may be used and its excess retained in its original positive square. When a coefficient is bounded it may be used unchanged.

Finite-penalty principle. Suppose a real symmetric quadratic form is strictly positive on the common kernel of finitely many real linear forms. Adding sufficiently large positive multiples of their squares makes the complete form strictly positive. To verify this without a limiting assertion, choose an orthogonal splitting into their common kernel and its complement. The upper block has a positive lower eigenvalue c, and the penalty matrix on the complementary block has a positive lower eigenvalue e. If the remaining lower block has norm at most b and the cross block norm is at most a, any penalty M with $Me > b + a^2/c$ makes the Schur complement strictly positive. All choices are real scalars. This principle makes the opening cases below uniform; no unbounded coefficient is inserted into a bounded Gram.

Testing u infinitesimal, v finite. Retain the n, D packets and excess caps of C, A, α. Write their finite selected coefficients as $c, a, b > 0$. Use roots $U = uW, Y, Z, X, V = \sqrt{m}\, vW$. The m packet is exactly

$$4(V - \sqrt{m}X + \sqrt{m}\, uvY - \sqrt{m}\, uZ)^2.$$

Together with the selected packets and the original core this is an exact bounded Gram expression. Its residue has isolated X, V pivots $2(1+v^2), 4$ and the following (U, Y, Z) block (residue notation suppressed):

$$\begin{pmatrix} 2+4cD_v & -2-4cD_v & -2v \\ -2-4cD_v & 2D_v+4cD_v+4aD_v+4bv^2 & -4bv \\ -2v & -4bv & 4b \end{pmatrix}. \tag{SR2}$$

In the interior, use $c = 1/(2L)$, $a = (1+L)/(2H)$ and $b = B/[2(1-H)]$. Its leading principal minors are

$$\frac{2(L+D_v)}{L}, \quad \frac{4(L+1)(D_v-H)(L+D_v)}{HL(1-H)}, \quad \frac{8B(L+1)^2D_v}{HL(1-H)}, \tag{SR3}$$

all positive. At an opening, apply the finite-penalty principle:

- $H_0 = 0$: the large A packet forces $Y = 0$; the restricted two-dimensional determinant is $4(L+1) > 0$.
- $H_0 = 1$: the large α packet forces $Z = vY$; the restricted determinant is $4(L+1)^2D_v/L > 0$.
- $L_0 = 0$: the large C packet forces $U = Y$; the restricted determinant is $4/(1-H) > 0$.
- $L_0 = 0$, $H_0 = 1$: both constraints leave the strictly positive form $2Y^2$.

The case $L_0 = H_0 = 0$ is excluded by $B_0 > 0$. Thus (SR2) can always be made strictly positive with bounded selected coefficients.

Testing u a unit, $w = 1/v$ infinitesimal. Retain n, m and the excess caps of C, A, α; retain the full D packet using its natural root. Put $U = uW$, $E = wZ$, $P_Z = \sqrt{D}Z$. In roots (U, Y, E, X, P_Z) the exact selected expression is

$$\begin{aligned} &4cD_w(U-Y)^2 + 4aD_uD_wY^2 + 4bD_u(Y-E)^2 \\ &\quad + 4\{\sqrt{D}(wU + uX - wY) - P_Z\}^2 \\ &\quad + 2\{(wU + uX - wY)^2 + D_u(Y-E)^2 + D_wX^2 - E^2 - 2UE + 2YE + 2uwXE\}. \end{aligned} \tag{SR4}$$

Its isolated X, P_Z residue pivots are $2D_u, 4$. The (U, Y, E) block is

$$\begin{pmatrix} 4c & -4c & -2 \\ -4c & 4c + (4a+4b+2)D_u & -4bD_u - 2u^2 \\ -2 & -4bD_u - 2u^2 & 4bD_u + 2u^2 \end{pmatrix}. \tag{SR5}$$

For interior openings the leading minors are

$$2/L, \qquad 4(1+L)D_u/[HL(1-H)], \qquad 8u^2(1+L)^2D_u/[HL(1-H)], \tag{SR6}$$

strict because $u_0 \neq 0$. The opening constraints and restricted positive determinants are respectively

$$\begin{array}{c|c|c} H_0 = 0 & Y = 0 & 4u^2(1+1/L) \\ H_0 = 1 & Y = E & 4u^2(1+L)/L \\ L_0 = 0 & U = Y & 4u^2D_u/[H(1-H)] \end{array}$$

and at $L_0 = 0, H_0 = 1$ the remaining form is $2u^2Y^2$. Finite penalties therefore give a strict full Gram at every opening.

Both testing angles unbounded. Use (IR4)–(IR7), replacing C by a finite positive cap c_* and setting $d = D$ (which is infinitesimal in the strong IH source). Retain the excess C packet and the entire n, m packets. The roots are

$$\begin{aligned} &S = \sqrt{c_*}W, \quad E = \sqrt{B}\,wZ, \quad U = pwW, \quad P = \sqrt{D}\,pZ, \\ &E_X = X + wW, \qquad E_Y = \sqrt{B/[H(1-H)]}(Y - HwZ). \end{aligned}$$

The exact graph identity (IR5) remains valid. The E^2 coefficient has residue $2c_s$, $c_s = (1+L)/B$, because its core contribution $2(1-H)^2/B$ must be retained when B is a unit. In (IR7), the potentially nonzero first-normal coefficient is now

$$2\{AH - \alpha(1-H)\} - (1-H),$$

whose residue is zero by (SR1). The normal E_Y pivot is also $2c_s$. All scalar-normal cross coefficients are infinitesimal, since $\sqrt{D}$ is infinitesimal. Thus the full diagonal residue is

$$\operatorname{diag}(4, 2c_s, 2, 4, 2, 2c_s) > 0. \tag{SR7}$$

The remaining remote charts. When u is unbounded and v is a unit, the homogeneous core has residue $2\{(W+vX)^2+(vY-Z)^2\}$. Positive finite caps on the C and A packets supply W^2 and Y^2, respectively, so the residue is strict on (W,X,Y,Z). When u is unbounded and v infinitesimal, put $p=1/u$ and use the five roots $(P,Q,W,Y,Z)=(pX,W+vX,W,Y,Z)$. Positive caps c,a,b give the residue

$$2Q^2+2(P+Z)^2+4cW^2+4aY^2+4bZ^2>0. \tag{SR8}$$

When u is a unit and v infinitesimal, the core has residue $2\{(uW-Y)^2+(X+uZ)^2\}$; the positive A and α caps give Y^2 and Z^2, making the residue strict. In these expressions all excess event coefficients are retained in their positive squares. When both testing angles are units, the inherited noncommuting-test theorem applies independently of source scales.

Exhaustive join. Classify each testing angle in the projective line as infinitesimal, a bounded unit, or unbounded, with literal zero and infinity included in the corresponding homogeneous chart. The infinitesimal/unbounded cell is the local theorem of §70.1. The infinitesimal/finite cells follow from (SR2); the unit/unbounded cell from (SR4); the unbounded/unbounded cell from (SR7); the unbounded/finite and unit/infinitesimal cells from (SR8) and its preceding argument; and the unit/unit cell from the inherited theorem. These nine cells are exhaustive. Both exact sign gauges $(u,W,X)\mapsto(-u,-W,-X)$ and $(v,W,Y)\mapsto(-v,-W,-Y)$ preserve the full parent, so every signed testing chart is included. Every certificate retains the full state normals and the omitted original λ,β remain zero. Therefore the entire strong IH source leaf has all-tests tensor-SOS admission, including $n=0$ and every projective endpoint.

Chapter 72

Strong C2 source entry at an infinitesimal first angle

Component `pf5726218aec8`
Source: `R13/baseline/baseline/advances/ih/strong/strong_angle_entry.tex`
SHA256: `f5726218aec8d9e53923297c5d388116c09d3ea98fc0a916537bc11a9a771245`
Inventory use: GE1–GE3 actual source coquotients, Bunit and all retained opening bounds

72.1 Strong C2 source entry at an infinitesimal first angle

Let $z = u_c^2$ be infinitesimal, $y = v_c^2 > 0$ arbitrary, and use the exact positive C2 source. Set

$$H = zh \in (0,1), \quad L = yt > 0, \quad B = zb = H + L,$$
$$A_c = 1 + zr, \quad S = A_c + L, \quad j = A_c - L/y > 0.$$

Here $0 < r < 1$. Define

$$\begin{aligned}
\delta &= z + (1+z)B + (1+z)B^2 + 2z(1+z)Br + z^3r^2 + 2z^2r \\
&\quad + y\{(1+z)B + z + z^2r(2-r)\}, \\
G &= B + z(H + r - 1), \\
M &= (1+z)(B+zr)(H+y) - z(1+y)(1-r)L \\
&= (1+z)(B+zr)H + yG + yzj + zrL(1+y) > 0.
\end{aligned}$$

The unchanged original coefficient formulas are

$$\begin{aligned}
n &= \frac{(B+zr)A_cS - r\delta}{2r\delta}, & m &= \frac{z^2(1+y)(1-r)A_cS}{2G\delta}, \\
C &= \frac{A_cM}{2L\delta}, & A &= \frac{yA_c(B+zr)S}{2H\delta}, \\
D &= \frac{(1+z)(B+zr)(B-zyr)S}{2yj\delta}, & \alpha &= \frac{(B+zr)M}{2(1-H)\delta}.
\end{aligned} \tag{GE1}$$

Assume C and α are strong, meaning each has a positive real lower bound in the coefficient field. Then B is a bounded positive unit. Indeed

$$\delta \geq (1+y)(B+z), \qquad M \leq (1+z)(B+z)(1+y)$$

give $CL \leq A_c(1+z)/2$, so L and B are bounded. If B were infinitesimal then so would H, while $\alpha(1-H) \leq (B+zr)(1+z)/2$ would force α infinitesimal, a contradiction.

Next, $n \geq 0$ gives $r\delta \leq (B+zr)A_cS$ and therefore $yr \leq A_cS$, since $\delta \geq y(B+zr)$. Thus G, $B - zyr$ and $\delta/(1+y)$ are positive units. Since $L < yA_c$, the ratio $(H+y)/(1+y)$ is a positive unit:

if y is infinitesimal then $H = B - L$ is a unit, and otherwise $y/(1+y)$ is a unit. The positive second expression for M now shows that $M/(1+y)$ is a positive unit.

Finally $1 - r$ is a unit. To see this, let $E(r) = (B + zr)A_c S - r\delta$, holding B, H, y, z fixed. Its coefficients divided by $1 + y$ are bounded, while direct expansion at $r = 1$ gives

$$E(1) = -(H + y)(B + z)(1 + z).$$

The right side divided by $1+y$ is negative with nonzero residue. If $1-r$ were infinitesimal, $E(r)/(1+y)$ would have the same negative residue, contradicting $n \geq 0$.

Consequently the following exact scaled quantities are positive units:

$$\frac{m}{z^2}, \quad CL, \quad \alpha(1 - H), \quad \frac{AH(1+y)}{y}, \quad Dyj(1+y). \tag{GE2}$$

For the IH source subcase y is unbounded, hence $r = O(1/y)$ is infinitesimal, j is a unit, and Dy^2 is a positive unit. Put $\rho = B/y$, $s = r/\rho$, and $c_s = (1 + L)/B$. Then s is bounded and, with $N = 4n + 2$ and $\nu = Nr$,

$$\frac{\nu}{\rho} \longrightarrow 2c_s, \quad \frac{nr}{\rho} \longrightarrow \frac{c_s - s}{2}, \quad \frac{D}{\rho^2} \longrightarrow \frac{c_s}{2}, \quad CL \longrightarrow \frac{1}{2}, \quad \frac{Bm}{z^2} \longrightarrow \frac{c_s}{2},$$

$$AH \longrightarrow \frac{1+L}{2}, \qquad \alpha(1 - H) \longrightarrow \frac{B}{2}. \tag{GE3}$$

Here arrows mean equality of residues of bounded expressions, with B, H, L, s replaced on the right by their residues. The expressions remain valid at $H_0 = 0$, $H_0 = 1$, or $L_0 = 0$ whenever allowed by $B_0 > 0$. The reason is that the scaled rational functions have denominators bounded away from zero after the displayed opening factors have been absorbed; no unscaled $1/H$, $1/(1 - H)$ or $1/L$ is asserted bounded.

Chapter 73

Weak IH: exhaustive all-tests admission

Component p03eb42346077
Source: R13/baseline/baseline/advances/ih/weak/ih_all_tests_join.tex
SHA256: 03eb423460777d8feb007e9f38bd71ebe53220bc4bb561bb30090d2737fbccf8
Inventory use: Exact weak-IH all-testing and source-endpoint union; source frontier no longer used as an unsupported all-tests premise

73.1 Weak IH: exhaustive all-tests admission

The weak-C IH source leaf is admitted at every independent testing pair. The source chart is the unchanged positive C2 source with $d = 1$, $\lambda = \beta = 0$, $n \geq 0$, $z = u_c^2$ infinitesimal, $y = v_c^2$ unbounded, and C weak. The source reduction proves that $B = zb$ is unbounded and $\rho = B/y$ has residue in $[0, 1]$.

For testing $u, w = 1/v$ both infinitesimal, there are exactly three source cases: $\rho_0 = 0$, $0 < \rho_0 < 1$, or $\rho_0 = 1$. They are proved respectively by the first endpoint theorem, the preserved interior-corner theorem, and the second endpoint theorem. For testing u a unit and w infinitesimal, the open-axis theorem covers all three source cases. For testing u unbounded and w infinitesimal, (IR4)–(IR8) cover all source cases. Finally (IR1)–(IR3) cover every finite testing v and every testing u. This partition is exhaustive on the two projective testing lines.

Both sign choices are included. In particular, the exact gauges

$$(u, W, X) \mapsto (-u, -W, -X), \qquad (v, W, Y) \mapsto (-v, -W, -Y)$$

leave the entire original parent unchanged, all unlisted coordinates being fixed. The homogeneous formulas include literal zero and infinity without dividing by a zero testing coordinate. All source scales are independent of all testing scales, and all normal state coordinates are retained. Complex states split into their real and imaginary parts. No lower bound on n is imposed.

The old scope statements in the preserved revision-4 interior proof remain a faithful record of that theorem's original local scope. They are superseded for all-tests coverage by the present join; they are not assumptions or obstructions in this completed argument.

Chapter 74

Weak IH: all remote testing charts

Component `p0fb32cd2f1bc`
Source: `R13/baseline/baseline/advances/ih/weak/ih_remote_charts.tex`
SHA256: `0fb32cd2f1bcee14751012c11d486df66dc64ab47db7590ac916f168ab47a39a`
Inventory use: IR1–IR8 complete remote parent and infinity-infinity state joins

74.1 Weak IH: all remote testing charts

Assume the weak-C IH source entry, with no restriction on $\rho_0 \in [0,1]$, on positive $r, H, 1-H$, or on $n \geq 0$. The uniform source inputs are B unbounded, C, m infinitesimal, $BC, AH/B, \alpha(1-H)/B$ positive units with common residue $c = 1/[2(1+\rho_0)]$, and A, α unbounded. Every square retained below has nonnegative pure-source coefficient; products $1+u^2$ and $1+v^2$ are themselves testing sums of squares.

Finite v, infinitesimal u. Write $Y_A = \sqrt{A}Y$ and $Z_\alpha = \sqrt{\alpha}(vY - Z)$, so $Y = Y_A/\sqrt{A}$ and $Z = vY_A/\sqrt{A} - Z_\alpha/\sqrt{\alpha}$. Retain the complete n, C, D packets. Use the five roots

$$U = uW, \quad X, \quad V = \sqrt{m}\,vW, \quad Y_A, \quad Z_\alpha.$$

The m packet is exactly $4(V - \sqrt{m}X + u\sqrt{m/\alpha}Z_\alpha)^2$. All coefficients in the remaining core are bounded after substituting Y, Z, and all coefficients involving Y or Z are infinitesimal. The full residue Gram is diagonal:

$$2U^2 + 2(1+v_0^2)X^2 + 4V^2 + 4(1+v_0^2)Y_A^2 + 4Z_\alpha^2. \tag{IR1}$$

This includes literal $u = 0$ or $v = 0$; redundant zero roots cause no problem for a positive Gram representation.

Finite v, unit u. Retain n, m, C, D. Use (W, X, Y_A, Z_α). The residue of the core and selected A, α packets is

$$2\{u_0^2(W + v_0X)^2 + (1+v_0^2)X^2\} + 4(1+u_0^2)(1+v_0^2)Y_A^2 + 4(1+u_0^2)Z_\alpha^2, \tag{IR2}$$

which is strictly positive because $u_0 \neq 0$.

Finite v, unbounded u. Use the homogeneous coordinate $p = 1/u$, infinitesimal or zero. Retain n, m, D and use

$$P = pX, \quad Q = W + vX, \quad S = \sqrt{C}W, \quad Y_A, \quad Z_\alpha.$$

The C packet becomes $4(1+v^2)(S - p\sqrt{C/A}Y_A)^2$. The entire selected expression has bounded coefficients, and residue

$$2Q^2 + 2(1+v_0^2)P^2 + 4(1+v_0^2)S^2 + 4(1+v_0^2)Y_A^2 + 4Z_\alpha^2. \tag{IR3}$$

Thus every finite-v chart is complete, independently of n.

Both testing angles unbounded. Let $p = 1/u$, $w = 1/v$ be infinitesimal or zero and put $D_p = 1 + p^2$, $D_w = 1 + w^2$. Retain the entire n, m packets. Set $d = \min(1, D) > 0$ in the source field and retain $4(D - d)J^2$. The remainder is exactly

$$\begin{aligned} F_* &= 4CD_w(W - pY)^2 + 4AD_pD_wY^2 + 4dJ^2 + 4\alpha D_p(Y - wZ)^2 \\ &\quad + 2\{J^2 + D_p(Y - wZ)^2 + D_w(pX + Z)^2 - D_pD_wZ^2\}, \\ J &= wW + X - pwY - pZ. \end{aligned}$$

Use the graph $X = -wW$, $Y = HwZ$, the normals $E_X = X + wW$, $E_Y = \sqrt{L}(Y - HwZ)$ with $L = B/[H(1 - H)]$, and scalar roots

$$S = \sqrt{C}W, \quad E = \sqrt{B}\, wZ, \quad U = pwW, \quad P = \sqrt{d}\, pZ. \tag{IR4}$$

On the graph the exact expression is

$$\begin{aligned} F_*|_{\text{graph}} &= 4D_w(S - pH\sqrt{C/B}E)^2 + \frac{4D_p}{B}\{AH^2D_w + \alpha(1 - H)^2\}E^2 \\ &\quad + 4(1 + Hw^2)^2P^2 + 2D_wU^2 - \frac{4pD_w}{\sqrt{BC}}SE + \frac{2}{B}\{(1 - H)^2 + H^2p^2D_w\}E^2. \end{aligned} \tag{IR5}$$

All coefficients are bounded and its residue is $4S^2 + 4cE^2 + 2U^2 + 4P^2$.

The first X normal, divided by four, is exactly

$$L_X = -pD_wU + \frac{p(1 - H)w}{\sqrt{B}}E - 2\sqrt{d}(1 + Hw^2)P. \tag{IR6}$$

The first Y normal is

$$\begin{aligned} L_Y = &-2p\sqrt{C}D_wS + \frac{E}{\sqrt{B}}\{2Cp^2HD_w + 2D_p(AHD_w - \alpha(1 - H)) \\ &\quad + (2d + 1)p^2(1 + Hw^2) - D_p(1 - H)\}. \end{aligned} \tag{IR7}$$

After dividing by $\sqrt{L}$, all its coefficients are infinitesimal; the only possible nonzero term is proportional to $(AH - \alpha(1 - H))/B$, whose residue is zero. The quadratic normal coefficients have residues $2 + 4d_0$ and $4c$, and their mixed coefficient has zero residue. The only nonzero scalar-normal coupling is the (P, E_X) block

$$\begin{pmatrix} 4 & -4\sqrt{d_0} \\ -4\sqrt{d_0} & 2 + 4d_0 \end{pmatrix}, \tag{IR8}$$

whose determinant is exactly 8. Thus the full Gram is strict even when D is infinitesimal and even when $n = 0$. The normalization uses $\sqrt{d}$ but never $1/\sqrt{d}$ in a Gram coefficient.

Each displayed Gram is a representation of the complete selected state quadratic form, not just its graph restriction. The bounded positive-residue lemma and the retained squares prove tensor SOS. These charts, the open-axis theorem, and the three local source-shape proofs cover every independent testing pair.

Chapter 75

The two missing weak-IH source endpoints

Component `p18dbedfff317`
Source: `R13/baseline/baseline/advances/ih/weak/ih_source_endpoints.tex`
SHA256: `18dbedfff317c23862d5254744a5e0dc33f7149ead8f6cb87f55d9a7f28d610d`
Inventory use: WIE1–WIE6 both rho=0,1 source endpoints with full D packet and complete normals

75.1 The two missing weak-IH source endpoints

We use the exact parent, graph, roots, coefficients K, k, and both normal factorizations (IC3)–(IC12) in the preserved interior-corner proof. Those identities are exact for every positive canonical C2 source; their validity does not require the interior-shape assumption. Here z and testing u, w are infinitesimal (zero testing coordinates are allowed), B is unbounded, $H \in (0,1)$, $\rho = B/y$, and $N = 4n+2$, $\nu = Nr$. All source and testing rates are independent.

The endpoint $\rho_0 = 0$. Put $s = r/\rho$ and $\tau = 1/B$. Nonnegativity of n and the exact source formulas imply that s is bounded. The following normalized source quantities have the indicated residues:

$$\nu/\rho \to 2, \quad nr/\rho \to (1-s)/2, \quad D/\rho^2 \to 1/2, \quad BC, Bm/z^2, AH/B, \alpha(1-H)/B \to 1/2,$$

$$k/(B\rho) \to 2s, \qquad 0 \le s_0 \le 1. \tag{WIE1}$$

These are uniform opening identities: after substituting $B = 1/\tau$, $y = 1/(\tau\rho)$, $r = s\rho$, their normalized rational functions have a nonzero denominator at $z = \tau = \rho = 0$, including $H = 0, 1$. The fresh checker records these denominator residues. Thus they concern simultaneous, not iterated, infinitesimal limits.

Replace T by $T_\rho = \rho T$ in (IC7). Its coefficients are bounded; for example $r^2 n/\rho^2 = s(nr/\rho)$ is bounded. Add the exact zero

$$T_\rho^2 + (u^2 - z)T_\rho R_2 - \rho u(zR_0 - uR_1)(uR_0 - R_1) = 0. \tag{WIE2}$$

This is (IC8a) multiplied by ρ, and uses no mixed inverse. The residue Gram has R_0, R_1, E diagonal entries $2, 2, 2$, and

$$(R_2, T_\rho): \quad \begin{pmatrix} 2 & -s_0 \\ -s_0 & 1+2s_0 \end{pmatrix}.$$

Its determinant $2 + 4s_0 - s_0^2 \ge 2$ is strictly positive on $[0,1]$.

The entire normal restoration survives this rescaling. In (IC11), replace $f_X T$ by $(f_X/\rho)T_\rho$, where

$$f_X/\rho = -2(nr/\rho)(1+w^2) - s(2D+1)$$

is bounded. Every multiplier still contains testing u or w, and $1/\sqrt{N}$ is bounded. The remaining coefficients in (IC11) are unchanged and infinitesimal. Formula (IC12) uses the root R_1 rather than T and remains bounded and infinitesimal unchanged. The normal quadratic residue is $\mathrm{diag}(1,2)$. Hence the full seven-root Gram is strict. This includes arbitrary positive $r, H, 1-H$ and literal $n=0$.

The endpoint $\rho_0 = 1$. Set $e = 1 + zr - \rho + H/y = j > 0$, an infinitesimal, and write

$$\rho = \frac{1 + zr - e}{1 - H\tau}, \qquad \tau = 1/B.$$

The exact source formulas give the simultaneous normalized residues

$$De \to 1/4, \quad \nu \to 1, \quad nr \to (1-2r)/4, \quad BC \to 1/4, \quad Bm/z^2 \to (1-r)/4,$$
$$AH/B, \alpha(1-H)/B \to 1/4, \quad 0 \le r_0 \le 1/2. \tag{WIE3}$$

Put $\kappa = k - 8DHr$; then $\kappa/B \to r$. The scaled rational functions have regular nonzero denominators at $z = \tau = e = 0$ after the displayed powers of e are absorbed.

The large coefficient D must be retained in a square, not treated as a bounded Gram coefficient. Introduce the full-state root

$$F_D = \sqrt{D}\,(J + eZ), \qquad a = \sqrt{H/B}, \qquad Q = R_2 - rT - aE.$$

On the graph, $Q = J + eZ$. Subtracting its square from (IC7) gives the exact identity

$$\begin{aligned} G - 4DQ^2 = {} & (4BC - \kappa z/B)R_0^2 + 4BmR_1^2/z^2 + 2(2C+1)R_2^2 \\ & + r^2(4nw^2 + 4n + 2)T^2 + \frac{H}{B}\{4A(1+u^2) + 4C + 2\}E^2 \\ & + \kappa u R_0 R_1/B + (-4r + \kappa/B)R_2 T - 4a(2C+1)R_2 E. \end{aligned} \tag{WIE4}$$

Add the exact zero (IC8a). Restore the full normals

$$E_X = \sqrt{N}(X + ruZ), \qquad E_Y = \sqrt{B/[H(1-H)]}(Y - HwZ).$$

Use the exact relation

$$R_2 = F_D/\sqrt{D} + rT + aE - uE_X/\sqrt{N} + wE_Y/\sqrt{B/[H(1-H)]}. \tag{WIE5}$$

The inverse $1/\sqrt{D}$ is purely in the source field and infinitesimal. All coefficients after this substitution are bounded.

For completeness, the first-normal quotients of $\widehat{F} - 4F_D^2$ are precisely the D-removed versions of (IC11)–(IC12). Explicitly,

$$\begin{aligned} \bar{f}_X &= -2nr(1+V) - r, \\ \bar{g}_X &= -2mB/z - 2nr(1+z) + 1 - H - r - 2m(H + r - 1), \\ \bar{f} &= 2m - 2C(1+V) - V, \\ \bar{f}_Y &= 2AH(1+V) - 2\alpha(1-H) + (2m+1)(H + r - 1), \\ \bar{g}_Y &= -B(2C+1) + 2AH(1+z) + 2CH + H. \end{aligned}$$

The constant term in the Y monic polynomial acquires $2De$, but its monic quotient coefficients are unchanged; the X polynomial has the constant correction $-2De$. Exact source stationarity still sets the constant remainders to zero because $J + eZ$ vanishes at the contact. Thus the displayed quotient formulas reconstruct both linear normals. Their normalized coefficients are infinitesimal by (WIE3); in particular De is bounded and no bare D survives. The normal quadratic block of the D-removed expression has residue $\mathrm{diag}(1,1)$.

In the order $(R_0, R_1, T, E, F_D, E_X, E_Y)$, the complete residue Gram is diagonal:

$$\operatorname{diag}\{1, 1 - r_0, 1 + r_0 - r_0^2, 1, 4, 1, 1\}. \tag{WIE6}$$

For instance the T^2 entry comes from substituting $R_2 = r_0 T$ in (WIE4): it is $r_0\nu_0 - r_0^2$, plus the unit contribution from (IC8a). Every entry is positive for $0 \le r_0 \le 1/2$. This proves full-parent tensor SOS at the corner also when $1 - \rho$ is infinitesimal. Along with the preserved interior proof and the $\rho_0 = 0$ argument, every allowed source shape is covered at the independent testing corner u, w infinitesimal.

Chapter 76

An exact full-parent Gram at the interior IH double corner

Component `pc00b84929b12`
Source: `R13/baseline/baseline/advances/ih/weak/inherited_ih_interior_corner.tex`
SHA256: `c00b84929b1271fdda012cc24a981ac19c0196ecda9c619805b3adb12932c377`
Inventory use: IC1–IC12 full scalar and both complete normal identities; interior weak source Gram and polynomial identities separately reused by strong source

76.1 An exact full-parent Gram at the interior IH double corner

This is a local testing-chart theorem. It does not admit its source subdomain at all independent tests. In particular, no conclusion for remote commuting testing corners is included below.

Theorem 76.1 (Full-parent IH corner, interior shape)**.** *Use the exact positive C2 source with $d = 1$, the omitted original λ, β coefficients zero, and $n \geq 0$. Suppose its actual angles have type IH, and C is weak. Use the source notation of §77.1; thus z is infinitesimal and B unbounded. Write $\rho = B/y$ (a shape parameter, not an event coefficient). Assume*

$$\rho, \quad 1-\rho \quad \text{are positive bounded units.} \tag{IC1}$$

At every independent testing pair for which u and the homogeneous reciprocal coordinate $w = 1/v$ are infinitesimal or zero, the entire original parent has a separated-field tensor-SOS certificate, for all real or complex state coordinates. The source openings $H, 1-H$ and the positive radial parameter r are arbitrary; n may be unbounded. Literal testing $u = 0$ and $v = \infty$, either testing sign, and literal source $n = 0$ are included. No state coordinate or certificate root is required to be bounded.

Proof. Put $N = 4n + 2$ and $\nu = Nr$. Source reduction (WI1)–(WI3) shows that ν and $1 - r$ are bounded positive units, even when N is unbounded. In residue,

$$\nu_0 = \frac{2\rho_0}{1+\rho_0}, \qquad 2r_0 \leq \nu_0, \qquad D_0 = \frac{\rho_0^2}{2(1-\rho_0^2)}. \tag{IC2}$$

All assumptions and divisions in this sentence concern the coefficient field alone. The uniform coefficient estimates needed below are

$$BC \asymp 1, \quad \frac{Bm}{z^2} \asymp 1, \quad \frac{AH}{B} \asymp 1, \quad \frac{\alpha(1-H)}{B} \asymp 1, \quad D \asymp 1, \quad nr \leq \nu/4.$$

In particular C, m are infinitesimal, nr, nr^2 are bounded, $mB/z \asymp z$, and $m\sqrt{B}/z \asymp z/\sqrt{B}$. These estimates are uniform in $r, H, 1-H$ and do not assert that A/B or α/B is bounded at an opening.

The homogeneous testing parent is exactly

$$\widehat{F} = 4nD_uD_wX^2 + 4mM^2 + 4CD_w(uW - Y)^2 + 4AD_uD_wY^2 + 4DJ^2 + 4\alpha D_u(Y - wZ)^2$$

$$+2\{J^2+D_u(Y-wZ)^2+D_w(X+uZ)^2-D_uD_wZ^2\},$$
$$M=W-wX+uY-uwZ,\qquad J=uwW+uX-wY-Z,\qquad D_u=1+u^2,\quad D_w=1+w^2. \tag{IC3}$$

For $w\neq 0$ it is $w^2F(u,1/w)$; (IC3) defines the literal projective endpoint as well. Let G be (IC3) restricted to the graph

$$X=-ruZ,\qquad Y=HwZ. \tag{IC4}$$

Use the following five scalar roots, all defined using only pure coefficient radicals and inverses:

$$R_0=(uW-BwZ)/\sqrt{B},\qquad R_1=(zW-BuwZ)/\sqrt{B},$$
$$R_2=uwW-\rho Z,\qquad T=(u^2-z)Z,\qquad E=\sqrt{BH}\,(w^2-1/y)Z. \tag{IC5}$$

They vanish at every signed exact source contact. Define the pure coefficient quantities

$$K=4AH^2+8DHr+4\alpha(1-H)^2+4m(H+r-1)^2+2(H+r-1)^2+4nr^2,$$
$$k=K-4B^2m/z^2+8znr^2+8AH^2/y. \tag{IC6}$$

The exact source formulas and state/angular stationarity give the following finite polynomial Gram identity:

$$\begin{aligned}G=&\left(4BC-\frac{kz}{B}\right)R_0^2+\frac{4Bm}{z^2}R_1^2+2(2C+2D+1)R_2^2\\&+r^2\{4nw^2+2(2D+2n+1)\}T^2+\frac{H}{B}\{4Au^2+2(2A+2C+2D+1)\}E^2\\&+\frac{ku}{B}R_0R_1+\left\{-4r(2D+1)+\frac{k}{B}\right\}R_2T-\frac{4\sqrt{H}(2C+2D+1)}{\sqrt{B}}R_2E.\end{aligned} \tag{IC7}$$

For clarity, the only non-obvious algebraic step is the exact toric conductor

$$B(u^2-z)(w^2-1/y)Z^2=(u^2-z)ZR_2-zR_0^2+uR_0R_1. \tag{IC8}$$

Before (IC8), the last mixed radial term is $k(u^2-z)(w^2-1/y)Z^2$. Expanding (IC7), its difference from G has only the four monomials $uwWZ, u^2Z^2, w^2Z^2, Z^2$; the unchanged original source sets all four coefficients exactly to zero. The supplied checker verifies these four rational identities, not just a specialization or a leading-order equality.

All coefficients of (IC7) are bounded: BC, Bm/z^2, AH/B, $\alpha(1-H)/B$, nr, D and k/B are bounded. No inverse of r is used, and $1/\sqrt{N}$ is bounded because $N\geq 2$. Moreover,

$$(k/B)_0=\frac{2r_0}{1+\rho_0}.$$

Add to the right side of (IC7) the following exact zero, obtained by one more toric conductor:

$$\rho T^2+(u^2-z)TR_2-u(zR_0-uR_1)(uR_0-R_1)=0. \tag{IC8a}$$

The new T^2 term has positive unit residue ρ_0. Every other coefficient introduced by (IC8a) is infinitesimal. This step is what makes the proof uniform when r is infinitesimal and n unbounded.

Suppress residue bars temporarily. The R_0, R_1, E diagonal residues are respectively

$$\frac{2}{1+\rho},\qquad \frac{2(1-r)}{1+\rho},\qquad \frac{2}{1+\rho},$$

which are positive. The only surviving off-diagonal block is

$$(R_2,T):\qquad \begin{pmatrix}\dfrac{2}{1-\rho^2} & -\dfrac{r}{1-\rho}\\[2ex] -\dfrac{r}{1-\rho} & \rho+r\nu+4Dr^2\end{pmatrix}. \tag{IC9}$$

Using (IC2), its determinant is exactly

$$\frac{2\rho}{1-\rho^2}+\frac{r\{4\rho-r(3\rho+1)\}}{(1-\rho)(1+\rho)^2}>0.$$

Indeed $0\le r\le \rho/(1+\rho)$, and consequently $4\rho-r(3\rho+1)\ge \rho(3+\rho)/(1+\rho)>0$. The first determinant summand is already strictly positive, including $r_0=0$. Thus (IC7) after the exact zero (IC8a) has strictly positive definite residue, including literal $n=0$ and unbounded n.

It remains to restore the full state normals; the graph certificate alone is not the claimed conclusion. Put

$$E_X=\sqrt{N}(X+ruZ),\qquad E_Y=\sqrt{B/[H(1-H)]}(Y-HwZ).$$

We give exact factorizations of both linear normal terms. Let $U=u^2, V=w^2, s=1/y$ and define L_X, L_Y by $4L_X=\partial_X\widehat{F}|_{\text{graph}}$ and $4L_Y=\partial_Y\widehat{F}|_{\text{graph}}$. Direct expansion gives

$$\begin{aligned}L_X&=(2D+1)uR_2-\frac{2m\sqrt{B}}{z}wR_1+uZP_X(U,V),\\ P_X&=(2D+1)\rho-\frac{2mB}{z}V-2nr(1+U)(1+V)-r(2D+1)U-2D-r\\&\quad+V\{1-H-r-2DH-2m(H+r-1)\}.\end{aligned}\tag{IC10}$$

Every coefficient of P_X is bounded, and source stationarity gives $P_X(z,s)=0$. Write its exact monic division as

$$P_X=(U-z)f_X(V)+(V-s)g_X.$$

Both f_X, g_X have bounded coefficients. Explicitly,

$$\begin{aligned}f_X(V)&=-2nr(1+V)-r(2D+1),\\ g_X&=-2mB/z-2nr(1+z)+1-H-r-2DH-2m(H+r-1).\end{aligned}$$

Thus no inverse of r or of either opening is introduced. We obtain

$$L_X=\left(2D+1+\frac{g_X}{B}\right)uR_2-\frac{2m\sqrt{B}}{z}wR_1+uf_X(V)T-\frac{uwg_X}{\sqrt{B}}R_0.\tag{IC11}$$

Here we used the exact identity $B(V-s)Z=R_2-\sqrt{B}\,wR_0$; this avoids dividing by $\sqrt{H}$ when H is infinitesimal. Every displayed multiplier is infinitesimal in the bounded tensor ring; in particular $m\sqrt{B}/z\asymp z/\sqrt{B}$.

For the other normal, put

$$\begin{aligned}f(V)&=2m-2C(1+V)-(2D+1)V,\\ P_Y(U,V)&=Bf(V)+2AH(1+U)(1+V)+2CH(1+V)+2DHV+2DrU+2D\\&\quad-2\alpha(1-H)(1+U)+2m(H+r-1)U+(H+r-1)U+H(V+1).\end{aligned}$$

Then

$$L_Y=\sqrt{B}f(V)R_0+wZP_Y(U,V),\qquad P_Y(z,s)=0.$$

The coefficients of P_Y/B are bounded. Write $P_Y=(U-z)f_Y(V)+(V-s)g_Y$. The two quotient coefficients are explicitly

$$\begin{aligned}f_Y(V)&=2AH(1+V)+2Dr-2\alpha(1-H)+(2m+1)(H+r-1),\\ g_Y&=-B(2C+2D+1)+2AH(1+z)+2CH+2DH+H.\end{aligned}$$

The displayed source estimates show directly that f_Y/B and g_Y/B have bounded coefficients, including both opening limits. The elementary exact identity

$$(U-z)\sqrt{B}\,wZ=zR_0-uR_1$$

therefore gives

$$\frac{L_Y}{\sqrt{B/[H(1-H)]}} = \sqrt{H(1-H)}\left\{f(V) + \frac{zf_Y(V)}{B}\right\}R_0 - \frac{\sqrt{H(1-H)}uf_Y(V)}{B}R_1 + \frac{\sqrt{1-H}wg_Y}{B}E. \tag{IC12}$$

Again every displayed multiplier is infinitesimal. This argument retains the full X, Y linear terms; none is estimated on the graph and then discarded.

Finally the normal quadratic block, in (E_X, E_Y), is bounded. Its residue is diagonal with entries

$$1, \qquad \frac{2}{1+\rho_0} > 0.$$

Indeed the X^2 coefficient before scaling is $4nD_uD_w + 4mw^2 + (4D+2)u^2 + 2D_w$; the Y^2 coefficient is $4CD_w + 4AD_uD_w + (4D+2)w^2 + (4\alpha+2)D_u + 4mu^2$. Their mixed coefficient is $-4uw(2m+2D+1)$. For the Y normalization, if $L = B/[H(1-H)]$, then A/L and α/L are bounded and

$$\frac{4(A+\alpha)}{L} = 4\left\{\frac{AH}{B}(1-H) + \frac{\alpha(1-H)}{B}H\right\} \longrightarrow \frac{2}{1+\rho_0}.$$

Thus the stated normal residue remains strict even when $H_0 = 0$ or $H_0 = 1$; no denominator involving a vanishing opening was dropped. Equations (IC11)–(IC12) show that all cross terms from these normals to the five scalar roots have zero residue. Thus the exact seven-root Gram for the entire (IC3) is bounded and has positive definite residue. The bounded positive-residue Gram lemma yields tensor SOS. Its roots may be unbounded; it uses no mixed inverse or radical. The homogeneous chart and real/imaginary splitting give all endpoints and states stated in the theorem. □

Precise boundary of this result. The theorem covers only the independent testing corner $(u, v) = (0, \infty)$ in residue. It neither assumes nor concludes all-tests admission. The proof does not cover source ρ infinitesimal or $1-\rho$ infinitesimal. The positive radial parameter r and both positive openings $H, 1-H$ may be arbitrarily infinitesimal; n may be unbounded. The local proof does include literal $n = 0$, but a remote infinity-infinity joining Gram has a pivot proportional to n; that join has not been made uniform as $n \to 0$. No source leaf is removed from the all-tests frontier on the basis of this theorem.

Chapter 77

The genuine directed IH weak-C frontier

Component `p93b05ea6ad1e`
Source: `R13/baseline/baseline/advances/ih/weak/inherited_ih_source_frontier.tex`
SHA256: `93b05ea6ad1e767dea93fe62bf27c28f839db7f81157a435eb35dbf353487678`
Inventory use: Only WI1–WI3 actual weak-IH source entry and opening-normalized coefficient formulas; optional special-curve claims and old open-status statements not imported

77.1 The genuine directed IH weak-C frontier

Scope of retained extract. The current proof imports the complete exact source and opening/coquotient argument (WI1)–(WI3) below. Optional rational-curve examples and historical frontier prose are outside the mathematical scope used in this rederivation and are omitted from this presentation. The complete original file and the exact omitted spans remain in the source archive and presentation manifest.

Write $z = u_c^2$, $y = v_c^2$, and use the exact contact quantities

$$H = v_c Y_c = zh, \quad L = v_c(u_c W_c - Y_c) = yt, \quad B = H + L = u_c v_c W_c = zb, \quad R = -u_c X_c = zr.$$

Thus $0 < H < 1$, $0 < R < z$, $L > 0$, and

$$L < y(1+R), \quad B > yR, \quad G = B + R - z(1-H) > 0.$$

Set $S = 1 + R + L$ and

$$\begin{aligned}\widetilde{\Delta} &= (1+z)B^2 + 2(1+z)BR + (1+z)(1+y)B \\ &\quad + (z-y)R^2 + 2z(1+y)R + z(1+y), \\ \widehat{N} &= (1+z)(B+R)(H+y) - (1+y)(z-R)L.\end{aligned}$$

These are exactly $z\Delta, z^2 N_{\mathrm{src}}$. In particular,

$$\widehat{N} = (1+z)(B+R)H + yG + z\{y(1+R) - L\} + RL(1+y) > 0.$$

The original coefficients, without an approximation, are

$$\begin{aligned} C &= \frac{(1+R)\widehat{N}}{2L\widetilde{\Delta}}, & \alpha &= \frac{(B+R)\widehat{N}}{2(1-H)\widetilde{\Delta}}, \\ A &= \frac{y(1+R)(B+R)S}{2H\widetilde{\Delta}}, & D &= \frac{(1+z)(B+R)(B-yR)S}{2\{y(1+R)-L\}\widetilde{\Delta}}, \\ m &= \frac{z(1+y)(z-R)(1+R)S}{2G\widetilde{\Delta}}, & n + \frac{1}{2} &= \frac{z(B+R)(1+R)S}{2R\widetilde{\Delta}}. \end{aligned} \tag{WI1}$$

The checker verifies all six against the unchanged original source.

Uniform source reduction for IH. Suppose z is positive infinitesimal, y is positive unbounded, and $n \geq 0$. Then C is weak if and only if B is unbounded. To prove the nontrivial direction, suppose B is bounded. Since $R < z$ and $G > 0$, one has $z \leq 4(B + R)$: use $B \geq 1/2$ or, otherwise, $B + R > z(1 - H) > z/2$. Direct division in (WI1) gives bounded $\widetilde{\Delta}/[y(B + R)]$. Moreover,

$$\frac{\widehat{N}}{y(B+R)} = (1+z)(1+H/y) - (1+1/y)\frac{(z-R)L}{B+R}$$

has residue 1, since $L \leq B$ and z is infinitesimal. Consequently $C \geq c/L$ for a positive real c, which is strong because L is bounded.

Now let B be unbounded. Put

$$\rho = B/y, \qquad e = 1 + R - L/y > 0.$$

Here $0 < \rho < 1 + R + H/y$, so ρ is bounded and its residue lies in $[0, 1]$. The following exact normalized quantities have residue 1:

$$\frac{\widetilde{\Delta}}{B(B+y)}, \qquad \frac{\widehat{N}}{yB}, \qquad \frac{L}{B}, \qquad \frac{S}{B}, \qquad \frac{G}{B}. \tag{WI2}$$

They give

$$n + \tfrac{1}{2} = \frac{\rho}{2r(1+\rho)}(1 + o(1)).$$

The inequality $n \geq 0$ therefore gives r/ρ bounded and $r \leq \rho/[1 + \rho](1 + o(1)) \leq 1/2 + o(1)$. In particular $1 - r$ is a positive unit, and $R/\rho = z(r/\rho)$ is infinitesimal. The normalized leading relations, uniform at arbitrary openings $H, 1 - H, e$, are

$$\begin{aligned} C &= \frac{1+o(1)}{2B(1+\rho)}, & A &= \frac{B(1+o(1))}{2H(1+\rho)}, \\ \alpha &= \frac{B(1+o(1))}{2(1-H)(1+\rho)}, & m &= \frac{z^2(1-r)(1+o(1))}{2B(1+\rho)}, \\ D &= \frac{\rho^2(1+o(1))}{2e(1+\rho)}. \end{aligned} \tag{WI3}$$

Here each $1 + o(1)$ means a positive coefficient unit of residue one; it never denotes a comparison with a testing infinitesimal. In particular A, α are unbounded, m, C are infinitesimal, and Cm is infinitesimal for every weak-C IH source. Thus the four reciprocal-product exit can never admit this entire directed leaf. If ρ is infinitesimal then e is a unit and D is weak; if ρ is a unit then D is strong. Literal $n = 0$ remains allowed throughout the reduction. No lower bound on r was used.

Chapter 78

Completion of the strong C2 ordinary–small source chart

Component `p6b22e944bd7c`
Source: `R13/baseline/baseline/advances/ui/ui_all_tests_recovered.tex`
SHA256: `6b22e944bd7c51d3c7e80d46a5a2db0b3ae5a6f6171dfe98a55787befe99a6e4`
Inventory use: UA1–UA3 only: elementary real rank-one allocation lemma and proof | UA1–UA16 complete source openings, finite-cap approximation, remote/testing/projective endpoints and all state ratios | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

78.1 Completion of the strong C2 ordinary–small source chart

Theorem 78.1 (Strong C2 UI, all independent tests)**.** *Use the original C2 source (UR1)–(UR3), with $d = 1$, actual source angles $u_c = a > 0$ a bounded unit and $v_c = k > 0$ infinitesimal, and C, α strong. Then the entire parent (UR6) is a sum of squares in the tensor product of the independent source and testing fields, for every testing angle, including projective endpoints, and every homogeneous state. The statement retains the literal boundary $n = 0$ and arbitrary positive rates of every source opening.*

The local contact certificate in Section 84.1 already covers v infinitesimal and u with the same residue as a. We supply its complete complement. No comparison between v and k is made. The argument uses one elementary rank-one allocation, followed by short positive frames at the testing endpoints.

A rank-one allocation lemma. Let P be a positive definite real symmetric matrix, let ℓ, b be real columns, and let $H, c_0 > 0$. Set

$$d = \ell^T P^{-1} \ell, \qquad e = \ell^T P^{-1} b, \qquad h = c_2 - b^T P^{-1} b.$$

Suppose

$$p(T) = c_0 + (h + 4Hdc_0)T + 4H(dh + e^2)T^2 > 0 \quad (T \geq 0), \qquad dh + e^2 > 0. \tag{UA1}$$

There exists a real λ such that

$$|\lambda| < \sqrt{4Hc_0}, \qquad h + 2e\lambda - d\lambda^2 > 0. \tag{UA2}$$

Indeed put $s = \sqrt{4Hc_0}$. If $d > 0$ and $|e| < ds$, take $\lambda = e/d$. If $|e| \geq ds$, the value at the appropriate endpoint is $h + 2|e|s - ds^2$. Were this nonpositive, writing $x = |e| - ds \geq 0$ would give $h + ds^2 \leq -2sx$ and $dh + e^2 \leq x^2$; the quadratic in (UA1) would then have a nonpositive value at a positive argument. Thus a nearby interior λ works. This includes equality $|e| = ds$. The scalar interval argument also holds in a limiting case $d = 0$; then $e \neq 0$ follows from $dh + e^2 > 0$. (For a finite positive P itself, $d = 0$ would force $\ell = e = 0$, so this case occurs only when using the continuous scalar extension at a boundary.)

Consequently the two real matrices

$$\begin{pmatrix} P & b-\lambda\ell \\ (b-\lambda\ell)^T & c_2 \end{pmatrix}, \qquad \begin{pmatrix} 4H & \lambda \\ \lambda & c_0 \end{pmatrix} \tag{UA3}$$

are positive definite. This is an elementary real lemma. Its inverse and square root are never taken in the tensor ring.

The finite ordinary angle away from contact. Assume v infinitesimal and u a bounded unit whose real residue has square different from a_0^2. Keep the centered packets $P_L, P_C, P_n, P_m, P_\alpha$ of (UR12), (UR14) outside the Gram, and combine P_H with the exact remainder. Use the six separated roots

$$r_0 = uW - Y, \quad X, \quad Z, \quad V_0 = vW, \quad K_0 = kW, \quad R_0 = \frac{v(uX-Z)}{k}. \tag{UA4}$$

They satisfy the exact zero relation

$$R_0K_0 - (uX-Z)V_0 = 0. \tag{UA5}$$

All divisions in (UA4) are pure source divisions.

For completeness the boundedness of the C, D remainder in this frame is explicit. Write $\ell_0 = L\tau$, $c_R = C\rho k^2$, and $Q = uX - Z$. Then (UR13) gives exactly

$$\begin{aligned} G_{CD} + P_H = {} & 4\gamma R_0^2 + 8\ell_0 kK_0 r_0 + 8\ell_0 qk^2 V_0 Q - 4\ell_0\tau k^2 K_0^2 \\ & + 8c_R v V_0 r_0 - 4c_R\rho k^2 V_0^2. \end{aligned} \tag{UA6}$$

All displayed coefficients are bounded by (UR9)–(UR11); all terms except the first have zero residue. The A packet is exactly $4(A/k^2)D_uD_v(uK_0 - kr_0)^2$. Substituting $Y = uW - r_0$ in (UR15) and the signed core gives a bounded Gram on (UA4), with no discarded normal term. Its residue has an isolated $2r_0^2$ term. First suppose the leading source weights are finite. The remaining residue is

$$q_s^T P q_s + 2V_0 b_*^T q_s + c_2 V_0^2 + 4HR_0^2 + c_0 K_0^2, \quad q_s = (X, Z)^T, \tag{UA7}$$

where $D_u = 1 + u^2$, $H = H_0$, $A_w = A_0^{\text{weak}}$ are (RC3), and

$$\begin{aligned} P &= \begin{pmatrix} 4nD_u + 4m + 2 & (4m+2)u \\ (4m+2)u & (4m+2)u^2 + 4\alpha D_u \end{pmatrix}, \quad \ell = (u, -1)^T, \\ b_* &= \begin{pmatrix} -4mD_u \\ -D_u u(4m + 4\alpha + 2) \end{pmatrix}, \quad c_0 = 4A_w D_u u^2, \quad c_2 = 4mD_u^2 + (4\alpha+2)D_u u^2. \end{aligned} \tag{UA8}$$

The matrix P is strictly positive because m, α are strong. The last sign-changing part of (UA7) is handled by adding 2λ times (UA5). This gives precisely the two blocks (UA3).

Here is a direct proof of (UA1) for every ordinary angle away from contact. Put $N_* = 2n+1$, $T_* = 2m + 2\alpha + 1$, $U = u^2$, and $T = V^2$. The leading state polynomial (RC4), with its testing a replaced by u but its source unchanged, has determinant

$$\det f = 8D_u^2 u^2 V^2 \, \mathcal{L}(U, T), \tag{UA9}$$

where

$$\begin{aligned} \mathcal{L} = {} & \frac{4\alpha m N_*}{U} + 4A_w H T_* U + \frac{4A_w\alpha(2m+N_*)}{T} + \frac{4A_w n T_* U}{T} \\ & + 2HN_*T_*T + \frac{4HmN_*T}{U} + 4A_wH(2m+N_*) - N_*T_*. \end{aligned}$$

After division by N_*T_* all possible opening poles cancel. Define

$$i_N = N_*^{-1}, \quad t_m = m/T_*, \quad t_{am} = \alpha m/T_*, \quad q_\alpha = \frac{\alpha(2m+N_*)}{N_*T_*}, \quad i_\alpha = 1/\alpha.$$

Writing $z = a^2$, $A_c = 1 + zr$, and using the leading Δ of (RC3), these quantities are exactly

$$\begin{aligned} i_N &= \frac{r\Delta}{(b+r)A_c^2}, & t_m &= \frac{(1-r)(1-zb)}{2b(b+r)(z+1)}, \\ t_{am} &= \frac{z^2(b+r)(1-r)}{4\Delta}, & q_\alpha &= \frac{bz^2\{(z+1)b+(z+2)r-1\}}{2A_c^2}, \\ i_\alpha &= \frac{2(1-zb)\Delta}{z^2(z+1)b(b+r)^2}. \end{aligned} \tag{UA10}$$

All denominators here are units. The normalized Laurent polynomial is

$$\frac{\mathcal{L}}{N_*T_*} = \frac{4t_{am}}{U} + 4A_wHi_NU + \frac{4A_wq_\alpha}{T} + 2A_w(1-i_N)\frac{U}{T} + 2HT + 4Ht_m\frac{T}{U} + 4A_wHq_\alpha i_\alpha - 1. \tag{UA11}$$

Its value and its two first derivatives vanish at $(U,T) = (z,1)$, as direct substitution verifies. In logarithmic coordinates the nonconstant terms have nonnegative coefficients. The coefficients of U^{-1} and T are positive units. If $i_N > 0$, the coefficients of U and T^{-1} are positive; if $i_N = 0$, that of U/T is positive. The function is therefore strictly convex in logarithmic coordinates and has its unique minimum at $(z,1)$. This proves strict positivity at fixed $U \neq z$ for every $T > 0$. The constant and leading coefficients of the equivalent quadratic (UA1) are strictly positive: $c_0 > 0$, and the positive T coefficient in (UA11) gives the second assertion. Thus (UA2)–(UA3) prove the required strict residue Gram.

Opening boundaries and divergent source weights. The preceding proof extends to all allowed opening residues, rather than assuming n, m, α bounded. The following formulas make the needed uniformity checkable. Put

$$J_0 = q_\alpha + \frac{u^2(1-i_N)}{2}.$$

Then J_0 is a positive unit, and direct two-by-two inversion gives

$$\begin{aligned} \frac{\det P}{N_*T_*} &= 8D_uJ_0, \\ d &= \frac{u^2i_N + q_\alpha i_\alpha}{4J_0}, & e &= \frac{uD_u}{2J_0}, & h &= \frac{D_u(4t_{am}-u^2)}{J_0}, \\ (P^{-1}b_*)_1 &= \frac{i_N(u^2-4t_{am})}{2J_0}, & (P^{-1}b_*)_2 &= -\frac{u\{4t_{am}i_N + (1-i_N)u^2 + 1\}}{2J_0}. \end{aligned} \tag{UA12}$$

To check the unit assertion, if $i_N = 0$ then $J_0 = u^2/2 > 0$. If $i_N > 0$, the weight N_* is finite and the strong lower bounds on m, α give $q_\alpha > 0$: this follows directly from $\alpha(2m+N_*)/[N_*(2m+2\alpha+1)]$, whether both weights are finite or one diverges. Thus (UA12) has only unit denominators on the entire source residue domain.

The strictly positive leading coefficient is also explicit:

$$dh + e^2 = \frac{D_u(u^2+2t_m)}{2J_0} > 0.$$

Equations (UA10)–(UA12) prove continuity of d, e, h and of the minimizing hard vector. The convexity proof for (UA11) remains strict at every allowed boundary. Choose λ from the strict interval (UA2) at that boundary; the same strict inequalities hold nearby. Moreover P is bounded below by a positive real matrix, while $P^{-1}b_*$ and $P^{-1}\ell$ are bounded. Its Schur factorization therefore bounds the first block in (UA3) below by a positive real multiple of the identity, uniformly in the nearby source. The second block is a strictly positive real matrix as well.

The finite leading sources used in this limiting argument are valid real interior sources. This point must be established separately: the actual source has $b = h + (k^2/z)t$, and it is not legitimate

to substitute that actual b into a leading formula with denominator $1 - zb$ when this denominator has an infinitesimal sign change. For the limiting real parameters, put

$$f(b, r) = 1 + r(z - 1) - brz(z + 1) - r^2 z(z + 2).$$

Their domain is $b > 0$, $0 \le r < 1$, $g = (1+z)b-1+r \ge 0$, $1 - zb \ge 0$, and $f(b, r) \ge 0$. For $0 < s < 1$, let

$$b_s = sb + (1 - s)/(1 + z), \qquad r_s = sr.$$

Direct expansion gives

$$g_s = sg, \qquad 1 - zb_s > 0, \qquad f(b_s, r_s) = (1 - s)(1 + s - sr) + s^2 f(b, r) > 0.$$

Increase b_s by a sufficiently small $\epsilon > 0$. This makes g_s strict, decreases the last two strict margins by respectively $z\epsilon$ and $r_s z(z + 1)\epsilon$, and therefore preserves them. If $r = 0$, then increase r_s by a sufficiently small positive amount; g increases and f remains positive by continuity. Letting these perturbations tend to zero as $s \to 1$ supplies valid finite real leading sources approaching every allowed residue. Their t may be chosen as $(1 + zr)/2$, since the leading harmonic weight is independent of t. The normalized leading identity $2rn\Delta = bf(b, r)$ verifies the nonnegative-n requirement. The same construction supplies the real-interior approximation used in the finite-cap step of the local UI certificate.

At most two of n, m, α diverge: m can diverge only at $g = 0$, and α only at $1 - zb = 0$, which cannot occur together. Their centered excess rows have hard-state parts proportional to

$$X, \qquad X + uZ, \qquad Z.$$

Any possible pair is independent because u is a unit. For a vector in their common limiting kernel, change only (X, Z) by a vanishing amount to annihilate the corresponding nearby exact rows. Their large penalties then vanish exactly. Taking the limit in the uniform strict lower bound proves that the bounded finite part is strictly positive on that limiting kernel. The elementary finite-dimensional penalty lemma used in the local theorem now supplies finite real caps n_*, m_*, α_*. Their centered excesses are precisely the retained packets (UR14). The exact bounded remainder with those caps, after adding (UA5), has a strictly positive residue Gram on (UA4). This proves the finite ordinary-angle complement for all source openings and literal $n = 0$.

Endpoint frames. Choose positive real μ, m_*, α_* below, respectively, k^2C, k^2D, m, and α. The products k^2C, k^2D are strong by (UR5), and $A_w = A/k^2$ is a positive unit. Retain n and all positive excesses in full. The following formulas concern the parent with $C = D = \mu/k^2$, $m = m_*$, $\alpha = \alpha_*$ and $A = A_w k^2$. They therefore suffice. The root coordinates need not themselves be bounded; only their displayed Gram coefficients must be bounded.

At u, v infinitesimal use

$$E = (uW - Y)/k, \quad R = v(uX - Z)/k, \quad V = vW, \quad K = kuW, \quad X, Z.$$

The exact amplitudes are

$$J = k(E + R), \quad M = D_u V - X - uZ - uvkE, \quad vY - Z = uV - Z - vkE, \quad kY = K - k^2 E.$$

Their full residue form is

$$4\mu\{E^2 + (E + R)^2\} + 4m_*(V - X)^2 + 4\alpha_* Z^2 + 4\overline{A_w}K^2 + 2X^2, \tag{UA13}$$

which is strict. Literal zero testing angles are included without any division by a testing angle.

For bounded u and finite unit v, put $E = (uW - Y)/k$, $F = J/k$. Then

$$Y = uW - kE, \quad Z = uX + k(E - F)/v.$$

The roots (E, F, W, X) have strict residue

$$4\mu D_v E^2 + 4\mu F^2 + \{4m_* D_u^2 + (4\alpha_* + 2)D_u u^2\}(vW - X)^2 + 2D_v D_u X^2. \tag{UA14}$$

For bounded u and $v = \infty$ write $q = 1/v$ and homogenize by q^2. The same roots are defined using the homogeneous J, and

$$Y = uW - kE, \qquad Z = uX + kqE - kF.$$

The strict residue is (UA14) with $D_v = 1$ and $(vW - X)^2$ replaced by W^2. This includes the literal endpoint $q = 0$.

For $u = \infty$, put $p = 1/u$, $D_p = 1 + p^2$, and homogenize by p^2. Write $r_p = W - pY$, $Q_p = X - pZ$, and $P = vY - Z$. Then

$$J = r_p + vQ_p, \qquad M = D_p P - pQ_p + pvr_p.$$

For v infinitesimal use the six roots

$$E = r_p/k, \quad F = vQ_p/k, \quad P, \quad R = pQ_p, \quad K = kY, \quad S = pZ.$$

The signed core is exactly

$$2k^2(E + F)^2 + 2D_p P^2 + 2D_v R^2 - 4D_p D_v RP + 4D_p D_v pFK + 2D_p D_v S^2.$$

Its cross-term reduction uses $RvY = pFK$, an exact identity. Adding every capped packet gives the strict residue

$$4\mu\{E^2 + (E + F)^2\} + (4m_* + 2)(P - R)^2 + 4\alpha_* P^2 + 4\overline{A_w}K^2 + 2S^2. \tag{UA15}$$

For finite unit v, instead use $E = r_p/k$, $F = J/k$, P, K, S. Since $Q_p = k(F - E)/v$, the strict residue is

$$4\mu D_v E^2 + 4\mu F^2 + (4m_* + 4\alpha_* + 2)P^2 + 4\overline{A_w}D_v K^2 + 2D_v S^2. \tag{UA16}$$

Finally, at both infinite endpoints put $q = 1/v$ and homogenize by $p^2 q^2$. Use

$$E = (W - pY)/k, \quad F = (qW + X - pqY - pZ)/k, \quad P = Y - qZ, \quad K = kY, \quad S = pZ.$$

Here $M = D_p P + pkD_q E - pqkF$ and the entire signed core is

$$2k^2 F^2 + 2D_p P^2 + 2D_q p^2 k^2 (F - qE)^2 + 4D_q D_p k(F - qE)S + 2D_q D_p S^2.$$

The strict residue is (UA16) with $D_v = 1$.

Exhaustion and restoration. A testing angle is infinitesimal, a bounded unit, or projectively infinite. Equations (UA13)–(UA16) cover every pair except ordinary u with infinitesimal v. The local theorem covers the equal-residue contact, and (UA4)–(UA12) cover its complement. Exact sign gauges are

$$(u, W, X) \mapsto (-u, -W, -X), \qquad (v, W, Y) \mapsto (-v, -W, -Y);$$

all unspecified variables are fixed. Both preserve the full parent, so negative ordinary contact residues are included. All endpoint frames have been written after homogenization, including literal projective endpoints. Each proof has an exact bounded Gram with strict real residue; the strict-residue Gram lemma gives tensor SOS. Restoring all retained packets completes the original parent. Real and imaginary parts treat complex states. This closes the entire strong C2 UI admission and its exact symmetry image HU.

Reproducibility. The recovered endpoint and allocation identities are replayed by two scripts:

- `check_ui_projective_recovery.py`: seven exact checks.
- `check_ui_all_tests_recovery.py`: thirty-one exact checks.

Those scripts verify symbolic equalities, sign gauges, all five pole-free formulas, determinant factorization, contact derivatives, and displayed endpoint residue forms. The explicit real interior approximation also incorporates the same-system mathematical audit in `ui_boundary_audit.tex`; it is not an independent external verification. Positivity, boundedness and exhaustion are supplied by the preceding analytical arguments; passing algebraic checks alone is not used as their substitute.

Chapter 79

The centered-A certificate for every bounded positive δ

Component `pccaa793fd0bf`
Source: `R13/baseline/baseline/baseline/advances/c1_scales/centered_a_bounded_delta.tex`
SHA256: `ccaa793fd0bf7269744455946c8b647a4d93402c3779bb9d93461724797a5985`
Inventory use: entire theorem; CS1–CS2 and normal/restoration proof | CS1-CS4 and exhaustive outer restoration | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

79.1 The centered-A certificate for every bounded positive δ

Theorem 79.1. *Use the unchanged physical native C1 source with $k, \zeta, r = 1 - a\eta$ positive infinitesimals and η a positive bounded unit. Suppose $e = k\sqrt{\eta/\zeta}$ is bounded below by a positive real number; it may be unbounded. Then the entire original parent is tensor SOS at every independent projective test and every real or complex state. Every positive opening rate, literal $\beta = 0$, and every rate of k^2/r is included.*

Use all exact definitions and the centered positive packet (CA1) in `centered_a_unit_e.tex`. The sole change is that $\delta = \Delta R/K$ is now any positive bounded coefficient, so it must not be inverted in a bounded Gram. The source chart (CA3), $\ell = Qr = K - aE$, and the exact cancelled coquotients remain unchanged. Their only denominators are positive units, independent of the rate of ℓ relative to k.

Scale the entire remainder and its full normal. Let $\mathscr{G}$ be the exact full remainder (CA5). Put

$$\widehat{G} = \delta^2 \mathscr{G}, \qquad \widehat{M} = \delta M_B, \quad \widehat{Y} = \delta Y, \quad \widehat{W} = \delta W.$$

The original testing variables have not been rescaled against a coefficient rate; these displayed quantities are explicit separated tensor roots. Their complete formulas are

$$q_A = vD_u w, \quad q_B = v(D_u - uN - puD_u vw),$$

$$\widehat{W} = \delta w + u(\widehat{M} + \delta h q_B)/D_u, \quad \widehat{Y} = \delta uw - (\widehat{M} + \delta h q_B)/D_u.$$

Set

$$\widehat{B}_2 = \frac{8a\widetilde{c}_2 K^2}{R\widetilde{c}_3 Q^2}, \qquad \widehat{B}_b = \frac{8a\widetilde{c}_b K^2}{R\widetilde{c}_3 Q^2}.$$

The entire exact scaled remainder is

$$\begin{aligned}\widehat{G} = {} & \widehat{B}_2\{N^2 - 2q_A N + (1-p)q_A^2\} + \widehat{B}_b D_v\{N^2 + 2q_A N + pq_A^2\} \\ & + B_3 D_v\{\widehat{M}^2 + 2\delta q_B \widehat{M} + \delta^2 h q_B^2\} + B_5\{\widehat{M}^2 - 2\delta q_B \widehat{M} + \delta^2(1-h)q_B^2\} \\ & + 4A_* D_u D_v(\widehat{Y} - ru\widehat{W})^2 + 4LD_u D_v(2u\widehat{W}\widehat{Y} - ru^2\widehat{W}^2)\end{aligned}$$

$$+ 4\alpha D_u(v\widehat{Y} - \delta)^2 + 2dD_u\{(v\widehat{Y} - \delta)^2 - \delta^2 D_v\}. \tag{CS1}$$

All coefficients are bounded; every occurrence of δ has nonnegative exponent. There is no inverse of δ, ℓ, k, M, J or t. The inverse $1/a$ in A_* is a positive unit. Every first-normal and hard-normal term is retained in (CS1).

Uniform contact grading. Keep the exact roots $A_0, B_0, C_0, D_0, x, y, z, h_0, q_1, q_2$ and polynomial P in (CA6). Use the same weights $\mathrm{wt}(\delta, k, E, \ell, t, u, w, v) = (0, 1, 0, 1, 0, 1, 1, 2)$. Weight zero means bounded here, not bounded away from zero. The exact finite coefficient collection from (CS1) is

$$\begin{aligned}
(\widehat{G}|_{N=\widehat{M}=0})_{\le 6} &= \frac{4P}{E^4}x^2 - \frac{16(k+u)}{E^2}xz + \frac{8}{E}T_4,\\
(\widehat{L}_N)_{\le 3} &= 0,\\
(\widehat{L}_M)_{\le 3} &= \frac{8\{E^2 + 2E - 2\ell(E+1)\}}{E^2}x.
\end{aligned} \tag{CS2}$$

The full hard block on $(N, \widehat{M})$ has residue

$$\mathrm{diag}(16/E, 4(E^2+4)/E). \tag{CS3}$$

These are precisely the old identities after the indicated scaling, but their validity for an infinitesimal δ requires the nonnegative-exponent assertion in (CS1), not cancellation of a purported unit δ.

The scalar is in $(A_0, B_0, C_0)^2 + (D_0^2)$ over the ring with E and the displayed positive units inverted. Both complete first normals are in (A_0, B_0, C_0). Direct monomial allocation in (CS1) proves these memberships with nonnegative δ exponents. At zero normals, $q_A = A_0 D_u$, $\delta q_B = B_0 D_u(1 - puvw)$, and both $\widehat{Y}, u\widehat{W}$ belong to (B_0, C_0) with bounded-unit coefficients. The sole constant from the α packet satisfies $\alpha\delta^2/D_0^2 = 2a\tilde{c}_\alpha/\tilde{c}_3$ exactly. Thus these are memberships of the entire rational expression, not merely its finite collection. The unchanged monic contact conductor lifts them without a δ inverse. Its contact substitutions preserve the weights, even if δ is infinitesimal. After (CS2), all remaining scalar weights are at least seven and all remaining normal weights at least four, hence the lifted correction coefficients have zero residue. The finite checker verifies complete lifts of the displayed low-weight collection and explicitly rejects every negative exponent other than the allowed positive-unit E inverse.

The leading soft form is

$$\frac{4(E+4)}{E^2}x^2 + \frac{8}{E}T_4.$$

Use the strict two-pairing circuit on T_4. Its only hard coupling is the full first normal $8(E+2)x/E$ on $\widehat{M}$, and

$$\det\begin{pmatrix} 4(E^2+4)/E & 4(E+2)/E \\ 4(E+2)/E & 4(E+4)/E^2 \end{pmatrix} = 256/E^3 > 0. \tag{CS4}$$

Together with $16/E$ on N, this gives the strict eight-root Gram.

Exhaustion and restoration. The independent outer/state lemma `centered_a_outer.tex` uses the unit-e hypothesis only to infer a positive lower bound on m/d in (CAO2). When e is unbounded, m/d is instead unbounded, so that same lower bound and every clipping argument hold. Its $c/d, A/d, D/d$ bounds, exact centered root identity, ordinary-u certificate, unit-w certificate, and literal projective endpoint arguments do not contain e or δ. Thus it reduces every remaining independent test to exactly the contact chart settled above. Restore δ^2 times every retained original packet, including (CA1), and undo the positive pure coefficient factor δ^2. All factors and radicals remain separated between the two fields. Complex states split into real and imaginary evaluations.

This closes the formerly missing infinitesimal-r, unbounded-e slice, and incorporates the previous unit-e theorem. Together with the inherited unit-r results, it covers all positive native r when η is a unit and e is bounded below. It does not yet claim the infinitesimal-e slice.

Chapter 80

The outer and unit-state join for infinitesimal e

Component p05a86be0046c
Source: R13/baseline/baseline/baseline/advances/c1_scales/outer_small_e/outer_small_e.tex
SHA256: 05a86be0046c9b9ee011ff502597dbc9f6c3153d2ee20ef81aa92d23d11a005b
Inventory use: SEO1–SEO20 and all exact endpoint/critical-current proofs | SEO1-SEO20, full critical-wall/current identities and endpoints | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

80.1 The outer and unit-state join for infinitesimal e

Lemma 80.1. *Use the unchanged physical native C1 source with $c = 2$, $\lambda = n = 0$, positive original openings, and nonnegative original event coefficients. Suppose $k, \zeta, e = k\sqrt{\eta/\zeta}$ are positive infinitesimals, η is a positive bounded unit, and $r = 1 - a\eta > 0$. Then every independent projective test and state is tensor SOS except possibly the chart*

$$u, v, w > 0 \text{ infinitesimal}, \qquad Z = 1, \qquad w = (W + uY)/(1 + u^2). \tag{SEO1}$$

There is no restriction on the positive relative scales of a, r, J or on k^2/r. This lemma supplies a join, not a certificate for the remaining chart (SEO1).

The small weak-event scale. Put $\kappa = k^2$, $K = 1 + \kappa$, $H = r + \eta$, $V_c = \kappa\eta\zeta$, $S = 1 - \kappa\eta + tV_c$, $R = 1 + \kappa r$, $T = K - \kappa\zeta t$, $M = R - t$, and $J = T - a$. Let P_4, P_d be the short native polynomials printed in the centered-A outer lemma. All of K, H, S, P_4, P_d are positive bounded units. Set

$$L = KHS, \quad N = KH(1 + V_c) - J\eta P_d, \quad d_0 = 4\eta P_d, \quad A_0 = 2\eta^2 P_4, \quad D_0 = 2N.$$

After exact cancellation of the native r factors, the source is

$$d = \frac{ad_0}{L}, \qquad A = \frac{aA_0}{rL}, \qquad D = \frac{aD_0}{JL}, \qquad \frac{m}{ae^2} = \frac{2(1 - t\zeta)(1 + V_c)}{MS}. \tag{SEO2}$$

Here $\bar{N} = 1$. Native $\beta \geq 0$ gives the exact bound

$$\frac{t}{e^2} \leq \frac{2c_b}{\eta c_d} = \frac{K[1 + (1 - t)\zeta]H}{\eta P_d}.$$

Thus t is infinitesimal, $M \sim 1$, and $m/(ae^2) \sim 2$. In particular

$$\frac{k^2}{e^2} = \frac{\zeta}{\eta} \quad \text{is infinitesimal.} \tag{SEO3}$$

The ratios $d/a, A/a, D/a$ have positive real lower bounds, while d/a is a bounded unit. These statements allow arbitrary positive opening rates and literal $\beta = 0$.

Choose the critical parameter without a small denominator. There are two source cases. If a is a bounded unit, keep A, D, d and replace $c = 2$ by the critical number

$$c_* = \frac{d(d + 2A)(d + 2D)}{2(4AD - d^2)}. \tag{SEO4}$$

This is positive and bounded, and $c_* - 2 = O(k^2)$. Here is an exact short verification of the potentially delicate bound. Define

$$Q = 4A_0D_0 - rJd_0^2,$$

so $\bar{Q} = 16\bar{\eta}^2\bar{a}\bar{H} > 0$. Then

$$c_* - 2 = \frac{ad_0(d_0r + 2A_0)(d_0J + 2D_0) - 4QL}{2QL}. \tag{SEO5}$$

After $r = 1 - a\eta$ is inserted, its polynomial numerator vanishes identically at $\kappa = 0$, with ζ still free. It is therefore κ times a polynomial in bounded native variables; its denominator is a positive unit. No inverse of r or J occurs.

If a is infinitesimal, then $r, J \sim 1$. Divide the whole parent by a and replace the original A by

$$A_* = \frac{d^2(4 + d + 2D)}{2(8D - 2dD - d^2)}. \tag{SEO6}$$

The denominator is positive. In this case $A_*/a, D/a, d/a$ are positive bounded units, $c/a = 2/a$ is unbounded, and $(A_* - A)/a = O(k^2)$. To verify this without a rate comparison, put

$$B_0 = 8D_0L - a(2d_0D_0 + d_0^2J).$$

Its residue is $16\bar{H} > 0$, and

$$\frac{A_* - A}{a} = \frac{rd_0^2\{4JL + a(d_0J + 2D_0)\} - 2A_0B_0}{2rLB_0}. \tag{SEO7}$$

Again the numerator vanishes identically at $\kappa = 0$, so it has an exact polynomial factor κ, with a unit denominator.

Use $\sigma = 1$ in the first case and $\sigma = a$ in the second, and henceforth normalize all parent coefficients by σ. Let ϵ be the positive part of the critical increase in the chosen normalized coefficient. From (SEO2)–(SEO7),

$$\epsilon \text{ is infinitesimal}, \qquad \frac{\epsilon}{m/\sigma} = O(\zeta) \text{ is infinitesimal}. \tag{SEO8}$$

If the chosen critical value is smaller than the actual coefficient, retain the positive excess packet instead. In either source case the chosen critical parent obeys

$$2c(4AD - d^2) = d(d + 2A)(d + 2D), \tag{SEO9}$$

with the letters in (SEO9) now denoting the normalized critical coefficients. In the first case c, d are units and A, D are bounded below by positive units; in the second A, D, d are units and c may be unbounded.

A critical-wall frame permitting the large coefficients. Write $D_u = 1 + u^2$, $D_v = 1 + v^2$ and

$$B = X + uZ, \quad U = uW, \quad V = vZ, \quad J_t = U - Y, \quad C = 4D + 2d,$$

and define the pure-coefficient numbers

$$y = \frac{d}{2A}, \quad b = \frac{4AD - d^2}{AC}, \quad s = 1 - b, \quad q = y + b.$$

The critical identity gives the useful exact relations

$$b = \frac{C}{4c + C}, \quad Cs = 4cb, \quad cb = \frac{d(2A + d)}{4A}. \tag{SEO10}$$

In both source cases, y, b, s, q are bounded and $b, s > 0$. The entire signed core plus the three critical packets is exactly

$$\begin{aligned} F_* = {} & 4cD_v J_t^2 + 4AD_u D_v Y^2 + C(J_t + uvB - D_u V)^2 + 2dD_v B^2 \\ & + 2dD_u\{v^2 Y^2 - 2YV - V^2\}. \end{aligned} \tag{SEO11}$$

In the first source case use

$$H_A = \sqrt{A}(Y - yV), \qquad H_D = \sqrt{C}(J_t + uvB - (b + u^2)V). \tag{SEO12}$$

The inverse is $Y = yV + H_A/\sqrt{A}$ and $J_t = (b + u^2)V - uvB + H_D/\sqrt{C}$. All coefficients in (SEO11) after this substitution are bounded. Indeed the large A, C occur only in the normalized squares and in the bounded combinations Ay^2, $\sqrt{A}\, y$, Cs^2, $\sqrt{C}\, s$; use $Cs = 4cb$ and bounded c for the last two. At $u = v = 0$, the hard block on (B, H_A, H_D) is

$$\operatorname{diag}(2d, 4, 1 + 4c/C) \succ 0.$$

On its exact zero section the scalar coefficient of V^2 is

$$gu^2 + \gamma v^2 + 4cu^4 + g(y + 2)u^2 v^2 + 4cu^4 v^2, \quad g = \frac{d(2A + d)}{A}, \quad \gamma = \frac{Dd(2A + d)^2}{A^2(2D + d)}. \tag{SEO13}$$

Both g, γ are positive bounded units. Every hard-V entry has angular degree at least two, with bounded coefficients; this is an exact full-parent polynomial statement. Allocate it on uV, vV, leaving a factor u or v in its coefficient. The resulting exact five-root Gram on (B, H_A, H_D, uV, vV) therefore has strict residue.

In the second source case set $L_c = 4c + C$ and instead use

$$H_A = \sqrt{A}(Y - yV), \qquad H_c = \sqrt{L_c}(J_t - bV). \tag{SEO14}$$

Its coefficients are bounded because A, D, d are units and c/L_c, $cb/\sqrt{L_c}$ and cb^2 are bounded by (SEO10). Its hard block is $\operatorname{diag}(2d, 4, 1)$. The exact zero-section scalar is

$$gu^2 + \gamma v^2 + Cu^4 + gy\, u^2 v^2. \tag{SEO15}$$

The same complete hard-V allocation gives a strict five-root Gram on (B, H_A, H_c, uV, vV). Thus this is a full critical-wall certificate in both source cases, even with the stated unbounded coefficients. Every radical and denominator used in the roots is pure coefficient; no mixed inverse is used.

All unit-W states at the small-angle corner. Normalize $W = 1$, with Z, Y, X bounded, and take u, v infinitesimal. Retain the complete original α, β packets. Put

$$R_m = v - X + uvY - uZ = v(1 + uY) - B, \qquad V = \frac{Z}{1 + uY}(R_m + B). \tag{SEO16}$$

The displayed denominator is a positive pure-testing unit. In the first source case the only negative correction to the critical parent is $-4\epsilon D_v J_t^2$; (SEO12) and (SEO16) express $\sqrt{\epsilon} J_t$ as a bounded combination of the five critical roots and $\sqrt{m/\sigma}\, R_m$, with every coefficient having zero residue. In the second case the only negative correction is $-4\epsilon D_u D_v Y^2$; use $Y = yV + H_A/\sqrt{A}$ and (SEO16) for the identical conclusion. The only possibly delicate coefficient is a bounded testing multiple of $\sqrt{\epsilon/(m/\sigma)}$, and it is infinitesimal by (SEO8). The extra original m square contributes the strict diagonal 4. Hence the complete perturbed Gram remains strict. This includes unit or infinitesimal Z, literal $Z = 0$, and both signs of the bounded testing state. It uses the entire original weak m event, without a false positive-unit lower bound on m.

Finite outer angles without the weak event. The same critical parent also gives the two finite outer charts. In (SEO11) regard B, Y, J_t, V as independent roots. Exactly,

$$F_*(0,v) = F_*(0,0) + v^2\{2dB^2 + (4A+2d)Y^2 + 4cJ_t^2\}. \tag{SEO17}$$

The zero-angle form has three positive hard pivots and null ray $(B,Y,J_t,V) = (0,y,b,1)V$. On this null ray the additional coefficient is $\gamma v^2 > 0$. The normalized hard frames (SEO12) or (SEO14) keep every coefficient bounded. Thus when u is infinitesimal and v a bounded unit the full four-root residue is strict. The infinitesimal critical-parameter deficit is absorbed directly on these four roots; retain the entire m, α, β packets.

For bounded-unit u and infinitesimal v, set $V' = D_u V$. The other exact identity is

$$F_*(u,0) = F_*(0,0;V') + u^2\{4AY^2 + 2d(V')^2/D_u\}. \tag{SEO18}$$

Use the same normalized critical roots with V' in place of V. In the first source case include uvB in the H_D root so that the unbounded D event remains isolated for nonzero small v. The extra null mass in (SEO18) is positive in residue because d and u are units. The bounded full four-root Gram is strict, and the deficit is again a zero-residue correction. These two identities include every state, not merely the normal restriction.

For completeness the same argument covers both finite angles of nonzero residue, without a separate physical-source admission. The exact determinant of (SEO11) on (B,Y,J_t,V) is

$$64cdD_u^2D_v^2\{A(2D+d)u^2 + D(2A+d)v^2\}. \tag{SEO20}$$

The determinant in the normalized frame (SEO12) is this number divided by AC, namely

$$32cdD_u^2D_v^2\left\{u^2 + \frac{D(2A+d)}{A(2D+d)}v^2\right\}.$$

It is a positive unit whenever at least one finite angle has nonzero residue. The hard restriction $V = 0$ is explicitly

$$\left(4D_uD_v + \frac{2dD_uv^2}{A}\right)H_A^2 + H_D^2 + 4cD_v(H_D/\sqrt{C} - uvB)^2 + 2dD_vB^2,$$

which has a strict bounded residue. Hence its full residue Gram is positive definite by (SEO20). In frame (SEO14), the determinant is (SEO20) divided by $A(4c+C)$. Here A, D, d are units and $c/(4c+C) \sim 1/4$, so it too is a positive unit. Its hard restriction is

$$\left(4D_uD_v + \frac{2dD_uv^2}{A}\right)H_A^2 + \frac{4cD_v}{4c+C}H_c^2 + C(H_c/\sqrt{4c+C} + uvB)^2 + 2dD_vB^2,$$

again strict. All entries of both full transformed Grams are bounded as already proved. The infinitesimal deficit is a zero-residue correction on these four roots. This settles every finite angle pair outside the double-infinitesimal corner.

Projective infinity charts. For these charts normalize by a, clip the positive coefficients $c/a, A/a, D/a$ at suitable positive real lower bounds, and retain their positive original excess squares. Retain the whole original m, α, β packets. The normalized d/a is a unit. All the infinity-infinity and infinity-zero four-root identities in (BE2) and the subsequent outer routing apply without m. For $u = \infty$ in residue and v a bounded unit, the signed-core residue itself retains $2d\{(W+vX)^2 + (vY-Z)^2\}$; together with cW^2, AY^2 it is strict, so no $m(vY-Z)^2$ lower bound is needed. For $v = \infty$ and u a bounded unit the old signed-core and c, D root argument is unchanged. At the remaining $(u, 1/v) = (0,0)$ corner, replace the obsolete state-W mass argument by the exact root ideal $(X, Y, U = uW, Z)^2$ after retaining m. In $q^2F(u,1/q)$ its complete selected residue is

$$2dX^2 + 4c(U-Y)^2 + (4A+2d)Y^2 + 4DZ^2 \succ 0. \tag{SEO19}$$

Every remaining occurrence of W carries u, so this is a full bounded Gram including all states. Literal infinity is included by these homogeneous identities. Literal zero axes have their unchanged original-event SOS. Finite tests with both noncommuting residues were included directly in (SEO20).

The residual state chart. Only u, v infinitesimal remain. Normalize the state by a maximal coordinate. On a negative pure-testing-core branch, X, Y have zero residue. The unit-W chart was just proved for every bounded Z. Otherwise Z is a unit, so normalize $Z = 1$ and W is infinitesimal. Then $w = (W + uY)/D_u$ is infinitesimal. The inherited exact testing Schur identity and sign gauges give (SEO1). Complex states split into their real and imaginary evaluations. The exact checker accompanying this lemma verifies its source cancellations, critical roots, full hard-normal orders, both finite outer identities, and the repaired projective-corner identity.

Chapter 81

Small e: a bounded full-parent contact certificate for every r

Component `pa7e744614f49`
Source: `R13/baseline/baseline/baseline/advances/c1_scales/small_e_all_r_contact.tex`
SHA256: `a7e744614f496e9934ed7f03c4655c5f509f389faa5dd2056eb1dd4eb8875856`
Inventory use: SE1–SE8 and complete full-normal residue/restoration proof | SE1-SE11 | C1_BARE_ OTHER_TERMINALS; precise subordinate uses retained below

81.1 Small e: a bounded full-parent contact certificate for every r

Proposition 81.1 (Local part of the small-e theorem)**.** *Use the unchanged physical native C1 source, with k, ζ, e positive infinitesimals and η a positive bounded unit. Every native positive $r = 1 - a\eta$, and every positive opening rate, is allowed. At every independent test with u, v, w infinitesimal, $Z = 1$, and $w > 0$, the entire original parent is tensor SOS. Literal $\beta = 0$ is included.*

This proposition is the contact part only. The independent outer/state join is supplied separately; no assertion here identifies a testing rate with a source rate.

The source entry forced by the original sign. Keep K, Q, E, ℓ, δ from (CA3), with $E = b^2$ and $\ell = Qr = K - aE$. Here E is a positive unit, $0 < \ell < K$, and no lower bound on a or ℓ is assumed. Put

$$\varepsilon = 1/\delta, \qquad j = \delta k, \qquad t = \varepsilon^2 \tau.$$

Both ε, j are infinitesimal: $e = \varepsilon K/Q$ and $\zeta = j^2 QE/K$. The original condition $\beta \geq 0$ gives

$$\frac{t}{e^2} \leq \frac{2c_b}{\eta c_d},$$

whose right side is bounded after cancelling the common native r factor. Thus τ is positive bounded. The exact chart is

$$k = \varepsilon j, \quad K = 1 + \varepsilon^2 j^2, \quad Q = K^2 - \ell \varepsilon^2 j^2, \quad a = (K - \ell)/E,$$

$$\begin{aligned} &\eta = KE/Q, \quad r = \ell/Q, \quad H = r + \eta, \quad e^2 = \varepsilon^2 K^2/Q^2, \\ &\zeta = j^2 QE/K, \quad R = K^2/Q, \quad T = K - k^2 t\zeta. \end{aligned} \tag{SE1}$$

All denominators displayed here are positive units. The cancelled coquotients $\widetilde{c}_i = c_i/\ell$ are the same short exact formulas as in (CA3); the cancellation introduces no inverse of ℓ. Their residues are $2(E + \ell)$ for $i = 2, b, 3, 5, \alpha$, and 4 for $i = d$; c_4 has residue 2. These formulas hold uniformly as ℓ tends to zero or to one.

One bounded centered remainder, including both normals. Set

$$A_* = \frac{ArH}{1+rH}, \qquad A - A_* = \frac{A}{1+rH} > 0. \tag{SE2}$$

Retain the entire square $4(A-A_*)D_uD_v(Y-ruW)^2$. It and its full first derivative vanish at exact native contact. Divide the remaining parent by the positive pure coefficient a; every factor a cancels before coefficients are used. Write

$$\begin{aligned} &b_2 = 8\widetilde{c}_2e^2/(R\widetilde{c}_3), && b_b = 8\widetilde{c}_be^2/(R\widetilde{c}_3),\\ &b_3 = 8/T, && b_5 = 8\widetilde{c}_5/(T\widetilde{c}_3),\\ &L = Ar/a = 2\eta^2c_4/(Q\widetilde{c}_3), && d_0 = d/a = 2\eta\widetilde{c}_d/\widetilde{c}_3,\\ &A_0 = A_*/a = LH/(1+rH), && \alpha_0 = \alpha/a = 2\varepsilon^4j^6Q^2\eta^2\widetilde{c}_\alpha/(K^2\widetilde{c}_3). \end{aligned} \tag{SE3}$$

Every quantity in (SE3) is bounded with unit denominators, and b_2, b_b contain ε^2. Use the original normals N, M_B, put $p = t/R$, $h = a/T$, and write

$$\begin{aligned} &q_A = vD_uw, \qquad q_B = v(D_u - uN - puD_uvw),\\ &W = w + u(M_B + hq_B)/D_u, \qquad Y = uw - (M_B + hq_B)/D_u. \end{aligned}$$

The entire bounded remainder is

$$\begin{aligned} G_s = {}& b_2\{N^2 - 2q_AN + (1-p)q_A^2\} + b_bD_v\{N^2 + 2q_AN + pq_A^2\}\\ &+ b_3D_v\{M_B^2 + 2q_BM_B + hq_B^2\} + b_5\{M_B^2 - 2q_BM_B + (1-h)q_B^2\}\\ &+ 4A_0D_uD_v(Y-ruW)^2 + 4LD_uD_v(2uWY - ru^2W^2)\\ &+ 4\alpha_0D_u(vY-1)^2 + 2d_0D_u\{(vY-1)^2 - D_v\} + \nu D_vN^2, \qquad \nu = d_0(R-t)/R. \end{aligned} \tag{SE4}$$

This is (SD4), after the indicated exact centered subtraction and normalization, except that a positive part of the original Π_2 is kept inside the remainder. Indeed the original coefficient of D_vN^2 in Π_2/a is

$$\frac{4\beta + 2d}{a}\frac{R-t}{R} \;\geq\; 2\nu.$$

Thus subtracting only $\Pi_2/a - \nu D_vN^2$ remains legitimate, even when $\beta = 0$. The other three original packets are retained unchanged. All full first normals and every hard quadratic term are present in (SE4); there are no terms of higher normal degree.

A conductor that never divides by ε. Define the separated monomials and differences

$$A = \varepsilon wv, \quad B = v, \quad C = Euw, \quad D_0 = \varepsilon^2j^3E, \quad k = \varepsilon j,$$

$$x = B - C, \quad y = A - kC, \quad z = uB - kC, \quad h_0 = kC - D_0.$$

The two standard radical pairings are obtained by dividing the original four squared circuit monomials by δ: they are A, uB, kC, D_0, and their product is $(k^2BC)^2$. Their strict circuit is $T_4 = y^2 + z^2 + h_0^2 - 2yz + 2q_1^2$. The exact contact is $(u, w, v) = (k, j, \varepsilon j^2E)$. Besides the three basic conductor identities,

$$C(u-k) = z - ux, \quad \varepsilon C(w-j) = y - \varepsilon wx, \quad C(v - \varepsilon j^2E) = E(wz + jy - kwx), \tag{SE5}$$

use the additional, division-free identity

$$Cv(w-j) = Ew(w-j)z + Ejwy - Ekw^2x. \tag{SE6}$$

For a monic contact division $S = (u-k)T_u + (w-j)T_w + (v - \varepsilon j^2E)T_v$, whenever $T_w = \varepsilon U + vV$, the bounded conductor is

$$\begin{aligned} CS = {}& x[-uT_u - \varepsilon wU - Ekw^2V - EkwT_v]\\ &+ y[U + EjwV + EjT_v] + z[T_u + Ew(w-j)V + EwT_v]. \end{aligned} \tag{SE7}$$

No division by $\varepsilon, j, k, r, a, t$, or a contact difference occurs.

Here is the full rational membership argument required for applying (SE7), not only a finite truncation. At zero normals let $V_0 = 1 - puvw$. Then

$$q_A = D_u wv, \quad q_B = D_u vV_0, \quad uW = C/E + u^2 hvV_0, \quad Y = C/E - hvV_0.$$

The scalar of (SE4) therefore has an exact raw allocation

$$f_A A^2 + f_{BB} B^2 + f_{BC} BC + f_{CC} C^2 + \rho D_0^2. \tag{SE8}$$

Expand only the nonconstant part of $(vY - 1)^2$ into B, C; its constant has $\alpha_0/D_0^2 = 2\widetilde{c}_\alpha/\widetilde{c}_3$. The coefficients in (SE8) have unit denominators independent of w. Every dependence on w in them comes through V_0, and hence has a factor ε^2; modulo ε they are independent of w and even in u, v. Both complete first normals are in (A, B, C) with bounded coefficients by the same exact formulas.

Clear the finitely many unit denominators, chosen even in u, v and independent of w, and use the unchanged scalar square-lift algorithm. For its first scalar conductor, the only w-dependent coefficient T_w is divisible by ε (in fact by ε^2). At $\varepsilon = u = 0$, its residual polynomial S is even in v, and contact makes its constant zero; thus $S \in (v^2)$ there. More explicitly, modulo ε the first divided differences satisfy $T_u \in (u)$, $T_w = 0$, and $T_v \in (v)$ after the preceding monic substitutions. It follows that after this first conductor, the sole possible w dependence modulo ε, u in the four second-stage remainders is of the form wv times a polynomial in v. Hence their T_w all belong to (ε, v), exactly the hypothesis of (SE7). The complete normal raw coefficients are independent of w modulo ε, so their one conductor has $T_w \in (\varepsilon)$. The retained contact squares and the original native balances supply the exact contact value and first-jet identities throughout. This proves the full scalar-square and complete first-normal lifts over the bounded ring.

Finite residues of that full lift. Give $(\varepsilon, j, u, w, v, N, M_B)$ weights $(1, 1, 2, 1, 3, 5, 3)$; E, ℓ, τ have weight zero. Every positive-weight variable is infinitesimal, while all weight-zero coefficients are bounded. The contact substitutions preserve the grading; x has weight three and the other circuit roots weight five. Both (SE5) and (SE6) preserve weights, so after the scalar collection through weight ten and full first normals through weight five, every lifted remainder coefficient has positive weight. Restore the unit denominators afterward.

Set $H_0 = E + \ell$ and $V_* = 1 + \ell H_0$. Direct exact multiplication of (SE1)–(SE4), followed by (SE7), gives scalar residue

$$\frac{8(E+2)}{H_0 V_*} x^2 + 8T_4, \tag{SE9}$$

and complete first-normal residues

$$L_N = 0, \qquad L_M = \frac{16E(H_0+1)}{H_0 V_*}\, x.$$

The hard block has residue

$$\operatorname{diag}\left(\frac{4E}{H_0}, \frac{8(E^2 + 2E\ell + 2\ell^2 + 2)}{V_*}\right). \tag{SE10}$$

These statements include coefficient corrections of positive weight; they are not asserted as literal equalities of the full rational scalar. The exact checker reconstructs (SE4), verifies the full finite lifts, and compares their residues with (SE9)–(SE10). The only hard-soft coupled determinant is

$$\det G_{M,x} = \frac{64(E^2 + 2E\ell + 4E + 4\ell)}{H_0^2 V_*} > 0. \tag{SE11}$$

Together with $4E/H_0 > 0$ and the strict two-pairing circuit this gives the full bounded strict eight-root Gram. All original positive packets, the centered square, and the positive pure-coefficient factor a are then restored. All inverses and radicals remain separated between the two independent real closed fields.

Chapter 82

Completion of the unit-η, infinitesimal-e source domain

Component p29aeda9b68a4
Source: R13/baseline/baseline/baseline/advances/c1_scales/small_e_all_tests.tex
SHA256: 29aeda9b68a4eaee13258b11d90f4008a3f18acf19dfb3778602ced74ef5ad3f
Inventory use: entire all-tests join | entire theorem and proof | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

82.1 Completion of the unit-η, infinitesimal-e source domain

Theorem 82.1 (Small e, every positive native r, all independent tests)**.** *Use the unchanged physical native C1 source with $c = 2$, $\lambda = n = 0$, all original event coefficients nonnegative, and all original openings a, t, M, J positive. Suppose k, ζ are positive infinitesimals, η is a positive bounded unit, $e = k\sqrt{\eta/\zeta}$ is a positive infinitesimal, and $r = 1 - a\eta > 0$. Then the entire original parent is tensor SOS at every independent projective test and every real or complex state. This includes every relative rate of k^2/r, every positive opening rate, and literal $\beta = 0$.*

Proof. The exact original β sign supplies the bounded opening scale t/e^2 in (SE1) and (SEO2), without a lower bound on t. The outer/state lemma `outer_small_e.tex` proves every independent test outside the chart $Z = 1$ with $u, v, w > 0$ infinitesimal. Its critical-source deficits have bounded exact coquotients after a single factor k^2 is removed. The weak original m event absorbs those deficits because $k^2/e^2 = \zeta/\eta$ is infinitesimal. The two source cases a unit and a infinitesimal exhaust all positive native a, and the homogeneous endpoint identities include every projective corner and every state normalization.

On the remaining chart, the proposition in `small_e_all_r_contact.tex` gives the bounded full-parent remainder (SE4). Its centered cap is uniform in r; the reserved positive part of Π_2 keeps the full N pivot positive even at $\beta = 0$. The division-free conductor (SE5)–(SE8) applies to the entire rational scalar and both complete first normals. Its finite residues (SE9)–(SE11) give the strict eight-root Gram. Restore all retained original packets, the centered square, and the positive pure coefficient normalization. All testing and source inverses and radicals remain separated. The real and imaginary evaluations add to give the complex-state assertion. □

Together with the bounded-positive-δ theorem and the inherited unit-r results, this completes every positive e and every positive native r when k, ζ are infinitesimal and η is a positive bounded unit. It makes no assertion about infinitesimal or unbounded η, or about another native C1 source branch.

Chapter 83

The exact strong-C2 ordinary–small source reduction

Component `pc3daa35e9a9b`
Source: `R13/baseline/baseline/baseline/advances/c2_ui/full_parent_reduction.tex`
SHA256: `c3daa35e9a9bf4f24e2dd222610b0b42d4288eef948dd408990bf1f03a444b47`
Inventory use: UR1–UR17 entire original parent, exact centered packet retention and all normalized source differences including Ct-Dj

83.1 The exact strong-C2 ordinary–small source reduction

This section records the original source, every parent packet, and the bounded products used in the following reciprocal and local certificates. It is an exact algebraic reduction, not an all-tests admission. The coefficient and testing fields remain independent real closed extensions of $\mathbb{R}$.

Source definitions and physical domain. Take the actual source angles $u_c = a$ and $v_c = k$, where $a > 0$ is a bounded unit and $k > 0$ is infinitesimal. Set $z = a^2$ and $\varepsilon = k^2/z$. In the original canonical C2 chart put

$$h = b - \varepsilon t, \quad A_c = 1 + zr, \quad j = A_c - t, \quad g = (1+z)b - 1 + r - \varepsilon zt, \quad S = A_c + \varepsilon zt. \tag{UR1}$$

The native positive gaps are $r, 1-r, t, h, g, j, 1-zh$; assume all original event signs, including $n \geq 0$, and assume C, α are bounded below by positive real numbers. No positive lower bound on a native opening is imposed. Define the two complete source polynomials

$$\begin{aligned} \Delta &= 1 + (1+z)b + z(1+z)b^2 + 2z(1+z)br + z^2r^2 + 2zr \\ &\quad + \varepsilon z\{(1+z)b + 1 + zr(2-r)\}, \\ N &= (1+z)(b+r)\{b + \varepsilon(1-t)\} - \varepsilon(1+\varepsilon z)(1-r)t. \end{aligned} \tag{UR2}$$

With $d = 1$ and $\lambda = \beta = 0$, the unchanged original coefficients are

$$\begin{aligned} n &= \frac{(b+r)A_cS - r\Delta}{2r\Delta}, & m &= \frac{(1+k^2)(1-r)A_cS}{2g\Delta}, \\ C &= \frac{A_cN}{2\varepsilon t\Delta}, & A &= \frac{\varepsilon A_c(b+r)S}{2h\Delta}, \\ D &= \frac{(1+z)(b+r)(b-k^2r)S}{2\varepsilon j\Delta}, & \alpha &= \frac{z^2(b+r)N}{2(1-zh)\Delta}. \end{aligned} \tag{UR3}$$

These are the original rational C2 source formulas, with its angle-square parameter replaced by z and its squared angle-ratio parameter by ε; all quantities formerly supplied by a source-file reference are displayed in (UR1)–(UR3).

The source bounds used below follow directly from these definitions. First $t < A_c$, $h < 1/z$, and $b = h + \varepsilon t$ give bounded t, h, b. The inequality $(1+z)b + r > 1$ gives a positive lower bound on $b + r$; (UR2) makes Δ a positive bounded unit. The exact positive identity

$$N = (1+z)(b+r)h + \varepsilon\{g + j + rt(1+\varepsilon z)\} \tag{UR4}$$

shows that infinitesimal h would make b, N infinitesimal and $1 - zh$ a unit, contradicting strong α. Thus h, b, N are units. The numerator of $2rn\Delta$ evaluated at $r = 1$ is exactly

$$-z(\varepsilon + h)(b+1)(z+1).$$

It has negative unit residue, so $n \geq 0$ forces $1 - r$ to be a unit. Consequently

$$mg, \quad A/\varepsilon, \quad \varepsilon Ct, \quad \varepsilon Dj, \quad \alpha(1 - zh) \quad \text{are positive bounded units,} \qquad nr \text{ is bounded.} \tag{UR5}$$

In particular m is strong, A is infinitesimal, and C, D may have arbitrarily large positive sizes. Literal $n = 0$ is retained.

The entire parent and exact contact. For independent testing angles and a homogeneous state define

$$D_u = 1 + u^2, \quad D_v = 1 + v^2, \quad r_{\mathrm{amp}} = uW - Y, \quad Q = uX - Z,$$
$$J = r_{\mathrm{amp}} + vQ = uW + uvX - Y - vZ,$$
$$M = vW - X + uvY - uZ, \quad p = vY - Z, \quad B = X + uZ.$$

The signed core and the full original parent are

$$\mathcal{K} = 2\{J^2 + D_u p^2 + D_v B^2 - D_u D_v Z^2\},$$

$$F = 4nD_uD_vX^2 + 4mM^2 + 4CD_v r_{\mathrm{amp}}^2 + 4AD_uD_vY^2 + 4DJ^2 + 4\alpha D_u p^2 + \mathcal{K}. \tag{UR6}$$

Normalize the actual contact by $W_c = 1$. Introduce the bounded pure-source quantities

$$x = -r/b, \quad \rho = t/(ab), \quad \sigma = -j/(ab), \quad s = \sigma - \rho = -A_c/(ab), \quad \mu = g/b, \quad \pi = (zh-1)/(ab). \tag{UR7}$$

Here s is a nonzero bounded unit. The exact stationary contact is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(a, k; 1, kx, a - \rho k^2, \frac{k}{ab}\right), \tag{UR8}$$

and its amplitudes are

$$r_{\mathrm{amp},c} = \rho k^2, \quad J_c = \sigma k^2, \quad M_c = \mu k, \quad p_c = \pi k, \quad Q_c = sk.$$

Substitution of (UR3) gives $F = 0$ and all four state and both angle first derivatives zero at (UR8). These are the unchanged native contact identities; no equality of independent testing rates is used.

Products with all possible opening poles cancelled. Define two positive bounded units

$$c_1 = \frac{zA_cN}{2\Delta}, \qquad c_2 = \frac{z(1+z)(b+r)(b-k^2r)S}{2\Delta}.$$

The exact products required by the local cap argument are

$$\begin{aligned} nx &= -\frac{(b+r)A_cS - r\Delta}{2b\Delta}, & m\mu &= \frac{(1+k^2)(1-r)A_cS}{2b\Delta}, \\ \alpha\pi &= -\frac{z^2(b+r)N}{2ab\Delta}, & C\rho k^2 &= \frac{c_1}{ab}, \qquad D\sigma k^2 = -\frac{c_2}{ab}. \end{aligned} \tag{UR9}$$

Their products with the bounded factors x, μ, π also show $nx^2, m\mu^2, \alpha\pi^2$ bounded. Crucially, cancellation before estimating gives the further exact identity

$$Ct - Dj = \frac{(1+z)(b+r)\{A_c j - zt(b-\varepsilon z r)\} - A_c(1+\varepsilon z)(1-r)t}{2\Delta}. \tag{UR10}$$

Thus $C\rho + D\sigma = (Ct - Dj)/(ab)$ is bounded for every positive rate of t, j, including infinitesimal openings.

Set

$$L = C + D, \qquad q = D/L, \qquad H = CD/L, \qquad \tau = (C\rho + D\sigma)/L, \qquad \gamma = k^2 H.$$

Here $q, 1/L, \tau, L\tau$ are bounded, and

$$\gamma = \frac{c_1 c_2}{c_1 j + c_2 t} \asymp 1, \qquad t + j = A_c \asymp 1. \tag{UR11}$$

The denominator in (UR11) is a positive unit even if either opening is infinitesimal. At finite source residues it gives precisely H_0 in (RC3). Equations (UR9)–(UR11) also show that the bounded finite part extends continuously to the allowed boundary residues: no inverse of $r, g, 1 - zh, t, j$ remains in these products.

Proposition 83.1 (Exact bounded UI remainder)**.** *For bounded source caps $0 \le n_* \le n$ and positive bounded-unit caps $m_* \le m$, $\alpha_* \le \alpha$, the parent (UR6) is the sum of explicit positive centered packets and a bounded polynomial remainder G. Every centered packet vanishes with its full first jet at (UR8), so G has exact zero value and all first derivatives there.*

Proof. The exact original C, D combination is

$$CD_v r_{\mathrm{amp}}^2 + DJ^2 = L(r_{\mathrm{amp}} + qvQ)^2 + Hv^2Q^2 + Cv^2 r_{\mathrm{amp}}^2.$$

Retain the three nonnegative packets

$$\begin{aligned} P_L &= 4L(r_{\mathrm{amp}} + qvQ - \tau k^2 W)^2, \\ P_H &= 4H(vQ - sk^2 W)^2, \\ P_C &= 4Cv^2(r_{\mathrm{amp}} - \rho k^2 W)^2. \end{aligned} \tag{UR12}$$

Their subtraction leaves exactly

$$\begin{aligned} G_{CD} = {} & 8L\tau k^2 W(r_{\mathrm{amp}} + qvQ) - 4L\tau^2 k^4 W^2 \\ & + 8Hsk^2 vWQ - 4Hs^2 k^4 W^2 \\ & + 8C\rho k^2 v^2 W r_{\mathrm{amp}} - 4C\rho^2 k^4 v^2 W^2. \end{aligned} \tag{UR13}$$

All its coefficients are bounded by (UR9)–(UR11), including $L\tau^2 = (L\tau)^2/L$. Centering at $k^2 W$ is essential to this boundedness assertion.

Also retain

$$\begin{aligned} P_n &= 4(n - n_*) D_u D_v (X - xvW)^2, \\ P_m &= 4(m - m_*)(M - \mu v W)^2, \\ P_\alpha &= 4(\alpha - \alpha_*) D_u (p - \pi v W)^2. \end{aligned} \tag{UR14}$$

The exact corresponding remainders are

$$\begin{aligned} G_n &= 4n_* D_u D_v X^2 + 8(n - n_*) xvW D_u D_v X - 4(n - n_*) x^2 v^2 W^2 D_u D_v, \\ G_m &= 4m_* M^2 + 8(m - m_*)\mu v W M - 4(m - m_*)\mu^2 v^2 W^2, \\ G_\alpha &= 4\alpha_* D_u p^2 + 8(\alpha - \alpha_*)\pi v W D_u p - 4(\alpha - \alpha_*)\pi^2 v^2 W^2 D_u. \end{aligned} \tag{UR15}$$

These coefficients are bounded by (UR9) and the bounded caps. With

$$G = G_{CD} + G_n + G_m + G_\alpha + 4AD_uD_vY^2 + \mathcal{K},$$

the entire exact identity is

$$F = P_L + P_H + P_C + P_n + P_m + P_\alpha + G. \tag{UR16}$$

It includes the signed core and every original state-normal term. All six retained roots vanish at (UR8), proving the contact-jet assertion. This proposition does not claim positivity of G. □

For the local theorem retain $P_L, P_C, P_n, P_m, P_\alpha$ outside the bounded Gram and combine G with

$$P_H = 4\gamma R_h^2, \qquad R_h = (vQ - sk^2W)/k. \tag{UR17}$$

The factor $1/k$ is pure coefficient. The exact graph normals are

$$E_X = X - xvW, \quad E_Z = Z - (ux - s)vW, \quad E_r = r_{\text{amp}} - \rho k^2 W.$$

Their inverse is polynomial:

$$X = E_X + xvW, \quad Z = E_Z + (ux - s)vW, \quad Y = uW - \rho k^2 W - E_r, \quad Q = svW + uE_X - E_Z.$$

Thus no state is discarded, including $W = 0$. The following reciprocal lemma identifies the leading relative-angle completion; the local certificate supplies the complete normal lift, cap choice, and strict Gram. The outer-angle and other commuting-axis join remains a separate obligation, so this reduction adds no all-tests source admission.

Chapter 84

A local strong-C2 certificate at the ordinary–small contact

Component p76545700d90b
Source: R13/baseline/baseline/baseline/advances/c2_ui/local_ui_certificate.tex
SHA256: 76545700d90b3f78362588444e472f5d9f96dd8938ed8d5aee2daf41a711f393
Inventory use: Complete strong-UI local scalar and all three state normals with actual-source Hessian

84.1 A local strong-C2 certificate at the ordinary–small contact

This section uses the exact full-parent reduction and source notation in Section 83.1. It proves a local independent-test statement. The other commuting axis and the outer projective join are not included.

Theorem 84.1 (Local ordinary–small C2 certificate)**.** *Use the unchanged physical C2 source with $d = 1$, actual contact $u_c = a$ a positive bounded unit and $v_c = k$ a positive infinitesimal, and C, α bounded below by positive real numbers. Then the full parent is tensor SOS for every independent test with v infinitesimal and $u - a$ infinitesimal, and every projective state. The original $n = 0$ boundary and all positive source opening rates are included. No upper bound on n, m, C, D, α is imposed.*

Here $u - a$ is the difference of the corresponding tensor factors. The residue condition means that their real residues agree; it does not identify the source and testing fields or compare their rates. The exact sign gauges give the corresponding negative-angle charts.

Bounded full remainder and its exact graph. The packet reduction leaves a bounded polynomial remainder G and the positive relative square

$$4\gamma R_h^2, \qquad \gamma = k^2 CD/(C + D) \asymp 1, \qquad R_h = \frac{v(uX - Z) - sk^2 W}{k}.$$

Every other retained packet is an exact positive square vanishing at the full actual contact. For positive bounded caps m_*, α_* and $0 \le n_* \le n$, set

$$E_X = X - xvW, \quad E_Z = Z - (ux - s)vW, \quad E_r = uW - Y - \rho k^2 W, \qquad Q_N = uE_X - E_Z.$$

The full inverse is polynomial. All terms of G, including its complete first-normal rows and quadratic normal block, are retained. There are no higher state-normal terms.

Put $w = u - a$, $\xi = v^2$, and use the additional separated roots

$$S = vwW, \qquad T = kwW, \qquad R = (v^2 - k^2)W/k.$$

The exact identities

$$kR = \frac{kR_h - vQ_N}{s}, \qquad RT = \frac{R_h T - SQ_N}{s}, \qquad S^2 - T^2 = \frac{w}{s}(R_h T - SQ_N) \tag{UI1}$$

use only the bounded pure-source inverse $1/s$.

Complete scalar and normal lifting. On the graph $E_X = E_Z = E_r = 0$, write $G = \phi(u,\xi)W^2$. Its bounded polynomial coefficients and the original parity imply $\phi \in (k^2, \xi)$. The exact contact identities give $\phi = \phi_u = \phi_\xi = 0$ at $(u,\xi) = (a, k^2)$. Successive monic Taylor division gives

$$\phi = w^2(k^2 A_0 + \xi A_1) + w(\xi - k^2)B + (\xi - k^2)^2 C_0, \tag{UI2}$$

with bounded coefficients. To see the asserted first factor directly, divide first in $u - a$; the second u coquotient stays in (k^2, ξ). The constant and first u terms are then divided in $\xi - k^2$. No source infinitesimal is inverted by these operations. Equations (UI1) allocate the three terms of (UI2) on (S, T, R_h, E_X, E_Z).

The full first E_X, E_Z coefficients are $vL_X(u,\xi)$ and $vL_Z(u,\xi)$. Exact contact gives $L_i(a, k^2) = 0$, so they are

$$v\{wL_{i0} + (\xi - k^2)L_{i1}\}.$$

The first summand is allocated on S, and the second uses $vkR = v(kR_h - vQ_N)/s$. The entire E_r coefficient belongs to (k^2, ξ) and vanishes at contact. The same division writes it as

$$w(k^2 L_{r0} + \xi L_{r1}) + (\xi - k^2)L_{r2}.$$

It consequently uses kT, vS, kR. All these E_r couplings have zero residue. Thus (UI1)–(UI2) produce an exact bounded Gram on

$$(E_r, E_X, E_Z, R_h, S, T). \tag{UI3}$$

This is a full-parent lift, not a restriction to a state kernel.

The strict matched Hessian. First take finite positive residue source weights. The leading model before setting the ordinary test angle equal to a is (RC4), with every testing occurrence of a replaced by u; its source coefficients remain those of (RC3). Let G denote one half of its angular Hessian after eliminating its two state directions, in variables (V, u) at the exact contact. Direct differentiation and two-by-two Schur elimination give

$$\mathsf{G}_{uu} = \frac{8(1-r)(1+zr)^2}{zb\Delta} > 0, \tag{UI4}$$

$$\det \mathsf{G} = \frac{16(b+r)^2(1+zr)^3\{3z+4-(z+2)^2 r\}}{b^2\Delta^2}. \tag{UI5}$$

The other two entries, making boundedness explicit, are

$$\mathsf{G}_{Vu} = \frac{4(1+zr)^2\{zb+zr+2r-2\}}{ab\Delta},$$
$$\mathsf{G}_{VV} = \frac{8(1+zr)\{z^2b^2+z^2br+zb^2+2zbr-zb-r+1\}}{b\Delta}.$$

For a short check of strictness, the exact leading n numerator is bQ_n, where

$$Q_n = 1 + r(z-1) - z(z+2)r^2 - z(z+1)br = (1-(z+1)r)(1+zr) - zrg.$$

Since $Q_n \geq 0$ and $g \geq 0$, one has $r \leq 1/(z+1)$. Consequently

$$3z + 4 - (z+2)^2 r \;\geq\; \frac{z(2z+3)}{z+1} > 0. \tag{UI6}$$

All denominators in (UI4)–(UI6) are positive bounded units. These formulas therefore give a strict, uniformly bounded-away-from-zero angular Schur form on the source residue domain, including $n = 0$. They contain no inverse of r, g, or $1 - zb$.

Large weights and finite real caps. We use the elementary finite-dimensional penalty lemma: if a real symmetric form B is strictly positive on the common kernel of the rows of a matrix V, then $B + MV^TV$ is positive definite for some finite real $M > 0$. A proof is to split into $\ker V$ and its orthogonal complement. The first diagonal block is strict; taking M sufficiently large makes its Schur complement strict too.

The bounded finite part obtained by the centered packets is continuous at all source boundary residues. This assertion is algebraic: its only large-weight products are

$$nx,\ nx^2,\ m\mu,\ m\mu^2,\ \alpha\pi,\ \alpha\pi^2,\quad C\rho k^2,\ D\sigma k^2,\ k^2H,$$

and their bounded cap analogues. Substitution of the original source formulas cancels the possible $r, g, 1 - zh, t, j$ poles in those products; their denominators are the displayed positive units. For example,

$$m\mu = \frac{(1+k^2)(1-r)A_cS}{2b\Delta}, \qquad \alpha\pi = -\frac{z^2(b+r)N}{2ab\Delta}.$$

The other cancellations are

$$nx = -\frac{(b+r)A_cS - r\Delta}{2b\Delta}, \quad nx^2 = x(nx), \quad m\mu^2 = \mu(m\mu), \quad \alpha\pi^2 = \pi(\alpha\pi),$$

$$C\rho k^2 = \frac{c_1}{ab}, \qquad D\sigma k^2 = -\frac{c_2}{ab}, \qquad c_1 = \frac{zA_cN}{2\Delta}, \quad c_2 = \frac{z(1+z)(b+r)(b-k^2r)S}{2\Delta}.$$

Finally $C\rho + D\sigma = (Ct - Dj)/(ab)$, using the exact canceled formula in the full-parent reduction. Thus even this difference has no hidden small-opening denominator. For k^2H, write $k^2C = c_1/t$, $k^2D = c_2/j$, with c_1, c_2 positive bounded units. Then $k^2H = c_1c_2/(c_1j + c_2t)$, whose denominator is a unit because $t + j = A_c$ is a unit. Its residue is exactly H_0 in (RC3).

At most two of n, m, α can diverge. Indeed m can diverge only at $g = 0$, while α can diverge only at $1 - zb = 0$; these two equalities would give $g = 1/z + r > 0$. The hard-state parts of the three possible diverging centered rows are, up to nonzero unit factors,

$$E_X, \qquad E_X + aE_Z, \qquad E_Z.$$

Every possible pair is independent with a unit determinant.

Here is why their constrained limiting finite form is strict. Approach the boundary by the positive leading-source domain. The state block is bounded below by a positive real form because m and α are strong. Its minimizing hard vector for a fixed angular vector is bounded: solve the at most two diverging packet equations in the hard variables, using the unit determinants just displayed; this supplies a bounded competitor, and all remaining coefficients stay bounded. If h_0 is that competitor and $h_{\min}$ the state minimizer for angular vector q, the exact identity

$$(h_0 - h_{\min})^T P(h_0 - h_{\min}) = F(h_0, q) - q^T \mathsf{G} q$$

and $P \succeq cI$ make boundedness of $h_{\min}$ explicit. Together with (UI4)–(UI6), the state Schur factorization then bounds the full matched Hessian below by a positive real multiple of the Euclidean norm. For a vector in the common limiting packet kernel, adjust only its hard coordinates to lie in the exact nearby packet kernels. The adjustment tends to zero and annihilates the large weights exactly. Taking the limit proves strict positivity of the bounded finite part on that common kernel. The penalty lemma now chooses finite real caps for every diverging weight. Bounded source weights can be retained in full. All caps are below their original weights, so the remaining centered excesses are positive squares.

This limiting argument uses only real finite-dimensional forms to choose caps. No inverse of a tensor matrix and no mixed root are used in the full-parent identity. The boundary residues are limits of the positive leading-source domain: g and $1 - zb$ specify the interval for b, and decreasing a positive r if necessary makes $Q_n \geq 0$ strict. Thus the argument also covers a literal original $n = 0$ and all combinations of the allowed boundary residues.

Separate angular roots and full strictness. The residue of the hard block in (UI3) is

$$2E_r^2 + 4n_*(1+a^2)E_X^2 + (4m_*+2)(E_X + aE_Z)^2 + 4\alpha_*(1+a^2)E_Z^2 + 4\gamma R_h^2,$$

which is strict. In the residue Gram, S couples only to the two ordinary hard normals, and T only to R_h; the E_r couplings vanish. Setting $S = T$ gives the matched leading Hessian under an invertible bounded change of coordinates. Indeed at $V = 1$ the ordinary normals are its two state deviations, R_h supplies the relative-angle deviation with nonzero coefficient $2s$, and $S = T$ supplies the ordinary-angle deviation. The chosen caps therefore make the sum of the two angular Schur diagonals strictly positive.

Add an appropriate real multiple of the last zero identity in (UI1). It redistributes mass between S^2 and T^2; all accompanying coefficients contain w and have zero residue. Since their Schur diagonal sum is positive, choose this multiple so that both Schur diagonals are positive. The entire six-root residue Gram is now positive definite. The strict-residue Gram lemma proves the exact bounded remainder plus relative packet is tensor SOS.

Finally restore every centered excess and both other C/D packets. The construction was homogeneous in W and its state inverse was polynomial, so it includes $W = 0$ and every projective state. Real and imaginary parts give complex states. This completes only the stated local independent-test domain; the outer-angle and other commuting-axis certificates are still required before admitting the full UI source leaf.

Chapter 85

A reciprocal contact completion for a singular angle

Component `pa142a027a282`
Source: `R13/baseline/baseline/baseline/advances/c2_ui/reciprocal_contact_lemma.tex`
SHA256: `a142a027a282606d8d7a73067aa75177207f544d3663e7826cf196d8c1834605`
Inventory use: RC1–RC5 reciprocal contact ideal and complete radical allocation

85.1 A reciprocal contact completion for a singular angle

The following exact algebraic lemma isolates a useful mechanism. It does not assert a complete C2 all-tests theorem.

Lemma 85.1 (Reciprocal contact completion)**.** *Let A_0, A_2 be symmetric matrices over a real closed field, let q_0 be a column vector, and consider*

$$f(q, V, W) = q^T(A_0 + V^2 A_2)q + 2VWb^T q + (c_0 + c_2 V^2)W^2.$$

Suppose $(q, V, W) = (q_0, 1, 1)$ is a zero with all first derivatives zero. Then

$$b = -(A_0 + A_2)q_0, \qquad c_0 = q_0^T A_2 q_0, \qquad c_2 = q_0^T A_0 q_0,$$

and the entire polynomial has the exact decomposition

$$\boxed{f = (q - Vq_0W)^T A_0(q - Vq_0W) + (Vq - q_0W)^T A_2(Vq - q_0W).} \tag{RC1}$$

Consequently $A_0, A_2 \succeq 0$ imply a sum of squares. If $A_0 \succ 0$ and $q_0^T A_2 q_0 > 0$, its nonzero zero states have $V = \pm 1$ and $q = Vq_0W$.

Proof. The q equation gives the displayed value of b. The V equation then gives $c_2 = q_0^T A_0 q_0$, and the W equation gives $c_0 = q_0^T A_2 q_0$. Expanding the right side of (RC1) proves the identity. The zero statement follows first from the strictly positive A_0 term and then from $(V^2 - 1)^2 W^2 q_0^T A_2 q_0$. □

Homogeneous form without a mixed inverse. For a positive pure-coefficient k, substitute $q = Q/k$ and $V = v/k$, and multiply (RC1) by k^2. It becomes

$$(Q - vq_0W)^T A_0(Q - vq_0W) + \frac{1}{k^2}(vQ - k^2 q_0W)^T A_2(vQ - k^2 q_0W). \tag{RC2}$$

Thus its roots use only the pure-coefficient factor $1/k$. There is no inverse or radical of a mixed tensor element and no comparison between a testing rate and k.

The bounded-source C2 UI leading model. Here $u_c = a$ is a positive bounded unit and $v_c = k$ is infinitesimal, with $d = 1$. To state a residue model, first restrict to the subdomain where the original n, m, α are bounded and the source parameters $r, g, 1 - a^2 b$ have nonzero residues. In the following formulas a, b, r denote those residues, $z = a^2$, and

$$A_c = 1 + zr, \quad g = (1+z)b - 1 + r, \quad \Delta = 1 + (1+z)b + z(1+z)b^2 + 2z(1+z)br + z^2 r^2 + 2zr.$$

The source formulas give

$$\begin{aligned} n &= \frac{(b+r)A_c^2 - r\Delta}{2r\Delta}, \qquad m = \frac{(1-r)A_c^2}{2g\Delta}, \qquad \alpha = \frac{z^2(1+z)b(b+r)^2}{2(1-zb)\Delta}, \\ H_0 &= \overline{k^2 CD/(C+D)} = \frac{z(1+z)(b+r)b}{2\Delta}, \qquad A_0^{\mathrm{weak}} = \overline{A/k^2} = \frac{A_c^2(b+r)}{2zb\Delta}. \end{aligned} \tag{RC3}$$

Normalize the actual contact by $W = 1$. With $v = kV$, $X = k\xi$, $Z = k\zeta$ and the leading constraint $Y = aW$, the leading polynomial of F/k^2 is

$$\begin{aligned} f = {} & 4n(1+a^2)\xi^2 + 4m\{V(1+a^2)W - \xi - a\zeta\}^2 + 4H_0 V^2(a\xi - \zeta)^2 \\ & + 4A_0^{\mathrm{weak}}(1+a^2)a^2 W^2 + (4\alpha+2)(1+a^2)(VaW - \zeta)^2 \\ & + 2(\xi + a\zeta)^2 - 2(1+a^2)\zeta^2. \end{aligned} \tag{RC4}$$

It has the form of the lemma, with $q = (\xi, \zeta)^T$,

$$q_0 = (-r/b, 1/(ab))^T,$$

$$q^T A_0 q = 4n(1+a^2)\xi^2 + (4m+2)(\xi + a\zeta)^2 + 4\alpha(1+a^2)\zeta^2, \qquad q^T A_2 q = 4H_0(a\xi - \zeta)^2.$$

Here $A_0 \succ 0$ and $H_0 > 0$. Put $s_0 = aq_{0,1} - q_{0,2} = -(1 + a^2 r)/(ab) \neq 0$. The exact source identities (RC3) verify all three coefficient relations in the lemma. Hence the apparently problematic relative normal and scale difference combine into the single root

$$\frac{v(aX - Z) - s_0 k^2 W}{k}. \tag{RC5}$$

The ordinary centered normals $(X, Z) - vq_0 W$ supply the remaining positive directions. Requiring separate positive pivots for the relative normal and the scale difference would obscure this completion.

This proves an exact leading-model identity on its stated bounded subdomain. It does not discard the additional large source scales: those are handled algebraically by the centered-packet reduction in Section 83.1. The complete local lifted Gram is proved in Section 84.1; the outer projective join remains a separate obligation.

Chapter 86

Five consecutive lower-bounded event packets

Component p80253fe9bc1b
Source: R13/baseline/baseline/baseline/advances/dependencies/five_retained_events.tex
SHA256: 80253fe9bc1be97cd3abb6df3c3c05b7b6e86541180b47263f28f69eb8b5605b
Inventory use: entire theorem and NW1–NW5 | Entire selected five-event commuting-axis/corner proof; noncommuting testing uses the original physical source and the detailed A6 bodies | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

86.1 Five consecutive lower-bounded event packets

The following argument permits every retained event coefficient to be unbounded. It requires no canonical omission, actual-contact coordinates, or source-shape entry. Five consecutive events in the eight-cycle suffice.

Theorem 86.1 (Five consecutive lower bounds)**.** *Let the original canonical source be physical, with $d > 0$ and all eight event coefficients nonnegative. Suppose*

$$d/m,\ d/c,\ d/A,\ d/D,\ d/\alpha \quad \textit{are positive bounded coefficient elements.}$$

Then the entire original parent is tensor SOS at every independent projective measurement pair and every real or complex projective state. There is no upper bound on any event coefficient divided by d. By exact CHSH relabeling the same conclusion holds if any five consecutive event coefficients in the cycle $(\lambda, D, c, A, \alpha, m, \beta, n)$ have these lower bounds.

Proof. Divide by the positive pure coefficient d, so $d = 1$. Let $\mu = \min(1, m, c, A, D, \alpha)$. It is a positive bounded unit. Select the parent F_μ with $m = c = A = D = \alpha = \mu$ and $\lambda = n = \beta = 0$. Keep every nonnegative excess and all three unselected coefficients as their original event squares. This uses only pure coefficient square roots. It makes no assertion that F_μ is globally physical.

At noncommuting residue tests apply the inherited noncommuting-residue theorem to the full physical source. It performs its own positive coefficient normalization if the event weights are unbounded. It remains to certify F_μ near every commuting residue test. Its coefficients are bounded, so a positive-definite real residue Gram is sufficient by the bounded positive-residue Gram lemma.

Write $\mathcal{H}$ for the signed core with $d = 1$ and $D_u = 1 + u^2, D_v = 1 + v^2$. On the four open commuting axes its exact homogeneous restrictions are

$$\begin{aligned} \mathcal{H}|_{u=0} &= 2D_v(X^2 + Y^2), \\ \mathcal{H}|_{v=0} &= 2\{(uW - Y)^2 + (X + uZ)^2\}, \\ \mathcal{H}|_{u=\infty} &= 2\{(W + vX)^2 + (vY - Z)^2\}, \\ \mathcal{H}|_{v=\infty} &= 2D_u(X^2 + Y^2). \end{aligned} \tag{NW1}$$

The remaining finite angle has nonzero residue. At $u = 0$, the retained m, D packets force $W = Z = 0$ on the core kernel. At $v = 0$, the retained A, α packets force $Y = Z = 0$, and then $W = X = 0$. At $u = \infty$, the retained c, A packets force $W = Y = 0$, and then $X = Z = 0$. At $v = \infty$, the retained c, D packets force $W = Z = 0$ after the core has forced $X = Y = 0$. Thus the selected residue state matrix is positive definite on every open axis, for either sign of the other finite angle.

Normalize a nonzero testing state by a coordinate of largest absolute value. Its coordinates are then bounded and one is a unit. At the four double corners, write F_{ab} for the homogeneous restriction of F_μ to $(u, v) = (a, b)$. Directly,

$$\begin{aligned} F_{00} &= (2 + 4\mu)X^2 + (2 + 12\mu)Y^2 + 4\mu Z^2, \\ F_{\infty 0} &= (2 + 8\mu)(W^2 + Z^2) + 4\mu Y^2, \\ F_{0\infty} &= 4\mu(W^2 + Z^2) + 2X^2 + (2 + 12\mu)Y^2, \\ F_{\infty\infty} &= 4\mu W^2 + (2 + 4\mu)X^2 + (2 + 12\mu)Y^2. \end{aligned} \tag{NW2}$$

Here infinity means the polynomial homogeneous chart, obtained by dividing the finite parent by the appropriate pure-testing squared angle. Every form is strict except its explicitly missing state coordinate. Consequently the only nontrivial residue states are the W ray at $(0, 0)$, the X ray at $(\infty, 0)$, and the Z ray at (∞, ∞).

At $(0, 0)$ normalize $W = 1$. The five local coordinates (u, v, X, Y, Z) are infinitesimal. The complete polynomial F_μ has no constant or linear term, and its degree-two part is

$$(2 + 8\mu)(u - Y)^2 + 2X^2 + 4\mu\{(v - X)^2 + Y^2 + Z^2\}. \tag{NW3}$$

Its residue is positive definite: vanishing successively gives $X = Y = Z = v = u = 0$.

At the corner $(\infty, 0)$ normalize $X = 1$ and put $p = 1/u$. The five infinitesimal coordinates are (p, v, W, Y, Z). The complete homogeneous polynomial again has no terms below degree two, and its quadratic part is

$$(2 + 4\mu)(W + v)^2 + (2 + 4\mu)(p + Z)^2 + 4\mu(W^2 + Y^2 + Z^2). \tag{NW4}$$

This is positive definite. At the corner (∞, ∞), normalize $Z = 1$ and put $p = 1/u, q = 1/v$. On (p, q, W, X, Y) the exact quadratic part is

$$(2 + 8\mu)(q - Y)^2 + 2X^2 + 4\mu\{(p - X)^2 + Y^2 + W^2\}. \tag{NW5}$$

It is positive definite as well. One can obtain (NW5) directly from (NW3) by the exact symmetry $(u, v; W, X, Y, Z) \mapsto (1/v, 1/u; Z, X, Y, W)$, which permutes the five equal retained weights.

For completeness, these quadratic calculations control the entire polynomial. In each local chart, allocate every higher monomial to a product of two of the five coordinate roots. The remaining monomial has positive degree, hence is infinitesimal; its coefficient is bounded because it is a real polynomial in the bounded unit μ. Symmetrizing gives an exact finite five-root Gram whose residue is (NW3), (NW4), or (NW5). No inversion, root, or comparison mixing the two fields occurs. The bounded positive-residue Gram lemma therefore certifies the full selected polynomial, including all higher terms.

Restore the excess event squares, the positive pure-testing homogeneous and projective state factors, and the positive coefficient d. The zero state is immediate. Splitting a complex state into real and imaginary parts proves the complex assertion.

Finally the exact full-parent Alice reflection acts on the displayed cycle indices as $i \mapsto 7 - i$, and the swap/reflection $(u, v; W, X, Y, Z) \mapsto (1/v, 1/u; Z, X, Y, W)$ acts as $i \mapsto 6 - i$. Their composition gives a one-step cycle rotation. Their full-parent identities have only positive pure-testing homogeneous factors, so they transport the five-packet conclusion to every cyclic translate. □

This theorem is a sufficient source domain, not an exhaustive zero-d source cover. It requires five selected events to be positive and noninfinitesimal relative to d. Its all-tests conclusion is independent of actual contact angles and of all original event hierarchies. In particular, it contains both C1 and C2 sources with all six retained events bounded below relative to d.

Chapter 87

An unbounded first actual angle forces a strong C packet

Component `pb7189b9c1f6e`
Source: `R13/baseline/baseline/baseline/advances/source/bounded_second_angle_exclusion.tex`
SHA256: `b7189b9c1f6e2bcb933cf7d128ea76d1ba7f7adeb5857773f5054f2a726e0bc3`
Inventory use: WE1–WE6 exact n numerator contradiction for weak-C actual HI/HU

87.1 An unbounded first actual angle forces a strong C packet

Theorem 87.1 (The directed weak-C types HI and HU are empty)**.** *Use the exact positive rational C2 source, with $d = 1$ and $n \geq 0$. If the first actual half-angle u_c is unbounded and the second actual half-angle v_c is bounded, then $1/C$ is bounded. Consequently no exactly-one-weak source oriented with C weak can have directed actual type HI or HU. Literal $n = 0$ is included.*

This is an original-source impossibility theorem. It makes no claim that a scalar contact identity by itself is a tensor-SOS certificate. All independent testing variables and state coordinates remain unrestricted; the two source leaves are empty before any testing restriction is imposed.

Proof. Write

$$q = u_c^{-2} > 0, \qquad y = v_c^2 > 0, \qquad H = u_c^2 h = v_c Y_c \in (0, 1),$$

and retain the original $r \in (0, 1)$. Here q is infinitesimal and y is bounded; y may itself be infinitesimal. Put

$$R = r/q = -u_c X_c, \quad B = u_c v_c W_c, \quad L = v_c(u_c W_c - Y_c), \quad T = B - yR.$$

The exact positive gaps give

$$B = H + L = yR + T, \qquad 0 < T < H + y, \qquad L > 0.$$

In particular H, T, y, r are bounded. The original source coordinates become exactly

$$z = 1/q, \quad a^2 = yq, \quad b = yr + qT, \quad h = qH, \quad t = r/q + (T - H)/y,$$

and the two gaps satisfy

$$g = H + (1 + y)r - 1 + qT > 0, \qquad j = (H + y - T)/y > 0. \tag{WE1}$$

The displayed expressions are pure coefficient identities; no inverse of a mixed source/testing element is used.

We first show that R is bounded. Set $Q = q^2\Delta > 0$. Direct expansion of the unchanged original formula for n gives

$$n = -\frac{qP}{2rQ}, \tag{WE2}$$

where the following exact decomposition of P is sufficient:

$$\begin{aligned} P &= r^2(1+y)(H+yr) - Tqr(y+2) - q^2(T^2+T) + P_+, \\ P_+ &= HTq^2 + HTqr + Hqr(y+1) + T^2q^2r \\ &\quad + Tq^2r(y+1) + 2Tqr^2(y+1) + qr^2y(y+1) > 0. \end{aligned} \tag{WE3}$$

Suppose for contradiction that R is unbounded, and put $\kappa = q/r = 1/R$, which is infinitesimal. Let $S = H + yr > 0$. The ratio T/S is bounded. Indeed, if r is a unit, then

$$T < H + y \le S/r.$$

If r is infinitesimal, (WE1) and boundedness of y, T imply $H_0 = 1$, so S is a unit and T/S is again bounded. Since T is bounded, $(T^2+T)/S$ is bounded as well. Dividing the lower bound in (WE3) by the positive r^2S gives

$$\frac{P}{r^2S} \ge (1+y) - \kappa\frac{T(y+2)}{S} - \kappa^2\frac{T^2+T}{S}.$$

The last two terms are infinitesimal, while $1 + y \ge 1$. Thus $P > 0$. Equation (WE2) contradicts $n \ge 0$, including the equality case $n = 0$. Therefore R is bounded.

Now $r = qR$ is infinitesimal. Equation (WE1) becomes

$$g = H - 1 + q\{(1+y)R + T\} > 0,$$

so $H_0 = 1$. Both $B = yR + T$ and $L = B - H$ are bounded. The original denominator has the exact form

$$\Delta = (1+q)\{B^2 + 2BR + (1+y)B\} + (1-yq)R^2 + 2(1+y)R + 1 + y. \tag{WE4}$$

It is a positive bounded unit: all quantities are bounded, and $1 - yq$ is a positive unit, with constant contribution $1 + y \ge 1$. The unchanged numerator N_{src} satisfies

$$\frac{N_{\mathrm{src}}}{q} = (1+q)(B+R)(H+y) - (1+y)(1-qR)L. \tag{WE5}$$

Since $H_0 = 1$ and $L = B - H$, its residue is

$$(N_{\mathrm{src}}/q)_0 = (1+y_0)(1+R_0) > 0.$$

Thus N_{src}/q is a positive bounded unit. Finally, $A_c = 1 + R$ and $a^2t = qL$, so the original coefficient is exactly

$$C = \frac{(1+R)(N_{\mathrm{src}}/q)}{2L\Delta}. \tag{WE6}$$

The numerator has bounded positive inverse, and $L\Delta$ is bounded. Therefore $1/C$ is bounded, as asserted. No lower bound on L was required; if L is infinitesimal the conclusion is stronger. □

The exact C2 reflection preserves directed type HI and exchanges C, α. Applying the theorem in both orientations shows that every physical HI source has both complementary packets strong. For type HU the reflected type is UI; the reflected consequence is that α is automatically strong at every UI actual contact. These observations respect the directed weak-C orientation; they do not identify IH with HI.

Chapter 88

Further exact source information on the three broad C1 rows

Component `p51451360ff4a`
Source: `R13/baseline/baseline/baseline/advances/source/further_source_implications.tex`
SHA256: `51451360ff4aede0a2323ffc6f17ef68f11b7cf4dd457fc8207c3a5e3c365ac5`
Inventory use: NI3–NI10, especially NI4–NI6 | NI1–NI10 and the accompanying boundedness arguments | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

88.1 Further exact source information on the three broad C1 rows

This section records source implications of (NC1)–(NC7), not further all-tests certificates. All quantities are in the coefficient field. Write $f \asymp g$ for two strictly positive quantities whose quotient and reciprocal quotient are bounded. Set

$$L = x\zeta, \qquad S = 1 + L + y, \qquad \rho = r + \eta.$$

The native signs and (NC5) imply the uniform estimates

$$S \le P_d \le 6S, \qquad \frac{1+L}{54} \le P_4 \le 3(1+L), \qquad P_\alpha \asymp h(1+y) + \zeta M. \tag{NI1}$$

Indeed $rx^2\zeta < x$ follows from $Q > 0$, and $\eta x^2 h\theta = x\theta(1-E) \le 1$, giving the P_d bounds from (NC2). For P_4, if $L \le 2$ use $E \ge 1/18$; if $L \ge 2$, then $r < 1/L \le 1/2$ and $P_4 \ge L/2$. Its upper bound follows immediately from (NC2). The last comparison uses $1 \le R \le 2$ and $1/18 \le E < 1$. Consequently the unchanged original source satisfies

$$\begin{aligned} C/d &\asymp \frac{\rho}{(1-r)S}, & A/d &\asymp \frac{\eta(1+L)}{rS}, \\ m/d &\asymp \frac{x\rho h(1+y)}{M\zeta S}, & \alpha/d &\asymp \frac{x^2\zeta\rho\{h(1+y)+\zeta M\}}{QS}. \end{aligned} \tag{NI2}$$

No opening has been bounded away from zero in these statements. In particular, E and P_4 do not supply additional hidden small source denominators on C1.1, C1.4 or C1.5.

C1.1: two unavoidable large weights and its actual angle. Here x, ζ are infinitesimal and η is unbounded. The inherited original-D cap gives $q = x\eta$ bounded with residue at most $1/2$. Hence $y = q\zeta$ is infinitesimal: the actual angle type is necessarily (I, I), independently of e. The exact formulas give

$$C/d \asymp \frac{\eta}{1-r}, \qquad A/d \asymp \frac{\eta}{r}, \qquad m/d \asymp \frac{e^2}{M}, \qquad \alpha/d \asymp xq\zeta, \qquad \beta/d + \frac{1}{2} \asymp \frac{e^2}{t}. \tag{NI3}$$

Thus C, A are unbounded relative to d, and α is weak. The inherited bounded polynomial formula for B_5 gives $D/d = O(1/\eta)$, so D is weak too, including its literal zero. In (NI3), $\beta \geq 0$ forces $t = O(e^2)$. Therefore when e is infinitesimal, t is infinitesimal, M is a unit and m is weak. When e is a unit, m is strong; when e is unbounded, both m, β are unbounded. These are restrictions on the original C1.1 row, not new admissions. Its positive infinitesimal δ_{tot} condition is automatic because $C/d, A/d$ are unbounded.

C1.4: the natural scale is $L = x\zeta$. Now x is infinitesimal and ζ is noninfinitesimal. The following implications follow uniformly through all retained coefficient boundaries.

1. If L is unbounded, $Q > 0$ forces r infinitesimal, and $1 - r = a\eta < 2\eta$ makes η noninfinitesimal. Hence $y = L\eta$ is unbounded: the actual type is (I, H). Here ζ is unbounded, $t = \theta/\zeta$ is infinitesimal, and M is a unit. Since $hy/\zeta = x\eta h < 1$, (NC2) gives $P_\alpha \asymp \zeta$. Equations (NI1)–(NI2) then yield

$$C/d \asymp L^{-1}, \qquad A/d \asymp r^{-1}, \qquad \alpha/d \asymp L/Q, \qquad m/d \asymp (x\eta h)/\zeta \leq O(\zeta^{-1}). \tag{NI4}$$

   In particular C, m are weak, A, α unbounded, and $(m/d)/(C/d) = O(x)$ is infinitesimal. The original total-scale intersection is automatic.

2. If L is a bounded unit, ζ is unbounded, M is a unit, $P_\alpha \asymp \zeta$, and $S \asymp 1 + \eta$. The identity $1 = r + a\eta$ gives $\rho/(1 + \eta)$ a positive bounded unit. Thus

$$C/d \asymp (1 - r)^{-1}, \qquad \alpha/d \asymp Q^{-1}, \qquad A/d \asymp \frac{\eta}{r(1 + \eta)}, \qquad m/d = O(x). \tag{NI5}$$

   So C, α are strong and m weak. Moreover A is weak exactly when η is infinitesimal; that case has r a unit and D/d unbounded, by $B_5 \sim 1$ in (NC3). The actual Bob angle is infinitesimal, a unit, or unbounded exactly as η is, because $y = L\eta$. No restriction on D, β is added in the other cases.

3. If L is infinitesimal and η is bounded, then y is infinitesimal. The $B_\beta \geq 0$ equation bounds $\theta \leq Kx\rho G/P_d = O(x + L)$, so h, M are units. Now (NI2) gives C strong, m, α weak, and A weak exactly when η is infinitesimal. In that last case D/d is unbounded, as in the preceding item. The actual type is (I, I).

4. If L is infinitesimal and η is unbounded, then

$$C/d \asymp \frac{\eta}{(1 - r)(1 + y)}, \qquad A/d \asymp \frac{\eta}{r(1 + y)}. \tag{NI6}$$

   Both are unbounded since $\eta/(1 + y) = (\eta^{-1} + L)^{-1}$. Also Q is a unit and (NI2) bounds

$$\alpha/d \leq O\left(x^2\zeta\eta h + \frac{x^2\zeta^2\eta}{1 + y}\right) = O(L),$$

   so α is weak. Finally $J \sim 1$ and $P_d/(1+y) \sim 1$ directly in (NC2); (NC4) gives $D/d + 1/2 \sim 1/2$, hence D weak. This extends the C, A-large/D, α-weak source pattern of C1.1 into C1.4, without claiming that its existing conductors apply. Here y may be infinitesimal, a unit, or unbounded. The total-scale intersection is again automatic.

These four alternatives exhaust C1.4, but they are necessary source reductions only. In particular no entire C1.4 leaf is closed.

C1.5 after the actual unit/unit admission. Here k is a unit. The new theorem removes actual (U,U); the inherited both-η,ζ-infinitesimal theorem is also subtracted. Two useful remaining source types can be specified more tightly.

If v_c is infinitesimal, $\eta\zeta$ is infinitesimal. Neither η nor ζ can be unbounded: ζ infinitesimal forces θ infinitesimal and excludes unbounded η by $E>0$; η infinitesimal gives $r\sim 1$ and excludes unbounded ζ by $Q>0$. Thus exactly one of η,ζ is a unit in this residual subcase. The exact stabilizer (AS5) exchanges these two alternatives. It keeps k a bounded unit, although a second orientation need not keep $k\le 1$. Choose the representative η a unit and ζ infinitesimal. Then

$$m/d,\beta/d \text{ are unbounded}, \qquad C/d, A/d \text{ are strong}, \qquad \alpha/d \text{ is weak}. \tag{NI7}$$

Indeed $\theta=t\zeta$ is infinitesimal, E is a unit, $P_d,P_4,P_\alpha,r+\eta$ are units, and (NC4) gives $m/d\asymp 1/(M\zeta)$ and $\beta/d+1/2\asymp 1/\theta$. The reflected representative also has a unit E: its exact value is $E'=h(1+y)/G$ in the original coordinates, with h a unit by the same $\theta=1$ exclusion used in the ordinary-angle proof. There is no conclusion about the size of D/d in (NI7).

If v_c is unbounded, both η,ζ are noninfinitesimal and at least one is unbounded. To see the first assertion, an infinitesimal ζ would force $h\sim 1$ and η bounded by $E>0$; an infinitesimal η would force $r\sim 1$ and ζ bounded by $Q>0$. Two rate regimes are then useful.

- If η is unbounded and ζ a unit, then h is infinitesimal. The exact alternative numerator identity

  $$B_\beta=(x\eta h)\{K(1+\zeta)-x\theta^2\}+xrGT-\theta(1+2xh+x\zeta) \tag{NI8}$$

  shows that $x\eta h$ is a unit: otherwise its residue at $\theta=1$ would be $-1-x\zeta(1-r)<0$. Thus P_α is a unit and $P_d\asymp\eta$. Equations (NC4) give

  $$C/d\asymp(1-r)^{-1}, \quad A/d\asymp r^{-1}, \quad m/d\asymp M^{-1}, \quad \alpha/d\asymp Q^{-1}, \quad D/d,\beta/d \text{ weak}. \tag{NI9}$$

  For the last assertion use $J\sim 1$, $\theta\sim 1$ and $P_d/(Ky)\sim 1$ in the original $D/d+1/2,\beta/d+1/2$ formulas. The total-scale condition δ_{tot} infinitesimal is therefore equivalent on this typed branch to at least one of $r,1-r,M,Q$ being infinitesimal. If all four are units, the inherited positive-total-scale theorem already applies.

- If ζ is unbounded, r is infinitesimal and M a unit. The noninfinitesimal η obeys $\eta h\asymp 1$. For if θ is infinitesimal this follows from $E>0$ and $h\sim 1$; otherwise divide (NI8) by ζ, use $xrG/\zeta=O(r)$, and obtain that $x\eta h$ has a positive-unit lower bound. Therefore (NC2) and (NC4) give

  $$C/d\asymp\zeta^{-1}, \quad m/d\asymp\zeta^{-1}, \quad A/d\asymp r^{-1}, \quad \alpha/d\asymp\zeta/Q. \tag{NI10}$$

  Thus C,m are weak and A,α unbounded. No size restriction on D,β is inferred. This branch already has infinitesimal δ_{tot}.

All implications retain the original physicality and exact source openings. They identify the missing source scales for further full-parent proofs; only the actual (U,U) subcase has received a new all-tests admission in this section and its companion.

Chapter 89

Bounded unit native state scale and bounded unit e

Component p69995d829bb2
Source: R13/baseline/baseline/baseline/baseline/advances/c1/bounded_e_all_tests.tex
SHA256: 69995d829bb284070279e022fc25cb48a65144979349551aa30b0b267df7a7f4
Inventory use: BE1–BE4; exact projective selected-parent identities and state entry | BE2-BE4, state entry, and exact infinity/zero selected-parent identity | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

89.1 Bounded unit native state scale and bounded unit e

Theorem 89.1. *Use the unchanged physical native C1 source, $c = 2$, $\lambda = n = 0$, with all original event signs and positive openings a, t, M, J. Suppose k, ζ are positive infinitesimals, and*

$$\eta, \quad r = 1 - a\eta, \quad e = k\sqrt{\eta/\zeta}$$

are positive bounded units. Then the complete original parent is tensor SOS at every independent projective test and every real or complex state. The openings may be arbitrarily small, and $\beta = 0$ is included.

Correction of the previously recorded outer obstruction. The claimed critical determinant in BOUNDED_E_LOCAL_FRONTIER.md omitted the term $2dq^2$. There is no equality corner of the stated kind. Indeed, with $p = 1/u$, $q = 1/v$, the exact homogeneous parent is

$$\begin{aligned} F_{\infty\infty} = {} & 4m(pW - pqX + Y - qZ)^2 + 4c(1+q^2)(W - pY)^2 + 4A(1+p^2)(1+q^2)Y^2 \\ & + (4D + 2d)(qW + X - pqY - pZ)^2 + (4\alpha + 2d)(1+p^2)(Y - qZ)^2 \\ & + (4\beta + 2d)(1+q^2)(pX + Z)^2 - 2d(1+p^2)(1+q^2)Z^2. \end{aligned}$$

At $Z = 1$, after retaining α, β, its complete quadratic part is

$$4cW^2 + 2dX^2 + 4D(X - p)^2 + 4AY^2 + (4m + 2d)(Y - q)^2. \tag{BE1}$$

In particular the (Y, q) determinant is $4A(4m + 2d) > 0$. This is an exact full-parent expansion; no physicality argument or opening assumption repairs the erroneous smaller quadratic.

An even shorter full-parent certificate retains every event except c, A. The remaining polynomial belongs to $(W, X, Y, qZ)^2$, with bounded coefficients at $p, q = 0$. Its residue Gram is the quadratic form

$$4cW^2 + 2dX^2 + 4AY^2 + 2d(Y - qZ)^2. \tag{BE2}$$

It is strictly positive when $c/d, A/d$ are bounded below by positive units. All cross terms of the complete polynomial are allocated on these four roots, so (BE2) handles arbitrary states and both literal projective endpoints, without a normal restriction.

Uniform lower bounds and outer tests. Write $H = r + \eta$. The short coquotients have residues $2rH$ for $c_2, c_b, c_3, c_5, c_\alpha$, 2 for c_4, and $4r$ for c_d. The source identities imply

$$\frac{m}{d} = \frac{e^2 c_2}{M\eta c_d}, \quad \frac{c}{d} = \frac{c_3}{a\eta c_d}, \quad \frac{A}{d} = \frac{\eta c_4}{c_d}, \quad \frac{D}{d} = \frac{2c_5 - J\eta c_d}{2J\eta c_d}. \tag{BE3}$$

Every ratio is bounded below by a positive real number. For the last, $J + a = T \sim 1$ and

$$\overline{2c_5 - J\eta c_d} = 4\bar{r}(\bar{H} - \bar{J}\bar{\eta}) = 4\bar{r} > 0.$$

No reciprocal of an opening is required to be bounded. Choose positive real lower bounds for the four ratios, cap these four coefficients at the corresponding multiples of d, and retain their nonnegative excess event squares. All selected-parent Gram coefficients divided by d are then bounded.

The physical-source theorem covers every noncommuting residue test. For u infinitesimal and v a bounded unit, the selected residue form is

$$4m(vW - X)^2 + 2d(1 + v^2)(X^2 + Y^2) + 4(c + A)(1 + v^2)Y^2 + 4D(Y + vZ)^2 \succ 0.$$

For $u = \infty$ in residue and v a bounded unit, retain the signed core at that commuting residue and use the four positive squares cW^2, AY^2, $m(vY - Z)^2$, $D(W + vX)^2$; their common kernel is zero. For $v = \infty$ in residue and u a bounded unit, use the core $2d(1 + u^2)(X^2 + Y^2)$ and the $c(uW - Y)^2, D(uX - Z)^2$ packets. The $(0, \infty)$ corner has the strictly positive state residue

$$4mW^2 + 2dX^2 + \{4(c + A) + 2d\}Y^2 + 4DZ^2.$$

The (∞, ∞) corner is (BE2). At $(\infty, 0)$ retain all events except c, A and use the exact four-root allocation on

$$(pX + Z,\ W,\ Y,\ vX), \qquad p = 1/u.$$

Its residue is

$$2d(pX + Z)^2 + 2d(W + vX)^2 + 4cW^2 + 4AY^2 \succ 0.$$

The full selected polynomial lies in the square of this displayed root ideal; the pure-testing change $Z = (pX + Z) - pX$ makes this immediate. The other coefficients have zero residue. These homogeneous identities include literal infinities. Literal zero axes have a nonnegative pure-testing core and are immediate original-event SOS. Only bounded u and infinitesimal v remain.

State entry and the W endpoint. Normalize the state by a maximal coordinate. Set $D_u = 1 + u^2$ and $w = (W + uY)/D_u$. On the negative pure-core branch, if w is infinitesimal then Z is a unit: otherwise the residue core $2\{(X + uZ)^2 + (uW - Y)^2\}$ forces every normalized state coordinate to have zero residue. Normalize $Z = 1$; the exact pure-testing Schur identity in (SD5) gives $w > 0$ after the standard angle gauges.

If w is a unit and u is a unit, a zero of that core residue has $Y = uw \neq 0$. The positive lower bound for A/d gives a strict state residue after clipping the positive events. Thus only u, v infinitesimal and $W \sim w$ a unit need an additional state check. When Z is infinitesimal, retain all events except clipped m, A and use the roots (X, Y, uW, vW). Since Z/W is a pure-testing infinitesimal, the complete selected polynomial has residue

$$2d\{X^2 + (uW - Y)^2\} + 4m(vW - X)^2 + 4AY^2 \succ 0. \tag{BE4}$$

Every remaining coefficient contains u, v or Z/W; there is no constant or first-normal term outside this frame. This includes literal $Z = 0$. If Z is a unit, normalize $Z = 1$ and use the next raw scalar frame with w a unit.

The exact remainder and its complete normals. Use exactly (SD3)–(SD8), their four positive packets, and the positive clearing factor $\mathcal{C}$. Put

$$\delta = \Delta R/K, \quad E = b^2 = \eta K/R, \quad A_0 = wv,\ B_0 = \delta v,\ C_0 = \delta E u w,\ D_0 = \delta^2 k^3 E, \quad x = B_0 - C_0.$$

Here δ, E are positive bounded units, and a, t are bounded. The factored five-term scalar is in $(A_0, B_0, C_0)^2 + (D_0^2)$, and the two complete first normals are in (A_0, B_0, C_0). These are the full formulas, without restricting normals or replacing native coefficients by leading surrogates.

At $k = v = 0$ and at the relevant small-w or small-u endpoint, put $r_0 = 1 - aE$ and $H_0 = r_0 + E$. Direct collection of the short formulas gives the raw scalar residue

$$8r_0H_0D_u^2A_0^2 + 8D_u\{x^2 + u^2r_0H_0B_0^2\} + 8r_0H_0D_uD_0^2. \tag{BE5}$$

The hard block on $(N, \delta M_B)$ is

$$\operatorname{diag}(16r_0H_0, 16r_0H_0 + 8E^2/D_u),$$

and the complete first normals have residues $0, 16Ex$. The coupled $(\delta M_B, x)$ determinant is $128D_ur_0H_0 > 0$.

If u is a unit and w infinitesimal, (BE5) is a strict Gram on $(A_0, x, B_0, D_0, N, \delta M_B)$. If u is infinitesimal and w a unit, use the exact relations

$$B_0 = (\delta/w)A_0, \qquad C_0 = (\delta/w)A_0 - x.$$

The factor δ/w is a product of a pure-coefficient unit and a pure-testing unit; it is not a mixed inverse. The resulting residue is the strict form

$$8r_0H_0A_0^2 + 8x^2 + 8r_0H_0D_0^2 + 16r_0H_0N^2 + (16r_0H_0 + 8E^2)(\delta M_B)^2 + 16E(\delta M_B)x. \tag{BE6}$$

This supplies precisely the unit-w chart for which the older Δ-small retained-current argument is not applicable.

For transparency, the only nontrivial raw coefficient collection in (BE5) is

$$r_0H_0D_u + a^2E^2 + r_0E(2a-1) = 1 + u^2r_0H_0, \qquad aE + r_0 = 1.$$

The cross coefficient is $-16D_u(aE + r_0) = -16D_u$. In the full normal formula S_B has residue $-16E$ and S_C has residue $16E$. These prove the claimed full normal residues directly.

The sole contact chart. For u, w, v infinitesimal and $Z = 1$, use the unchanged monic contact–conductor identities and the two-pairing circuit. Use weights

$$(\delta, k, b, a, t, u, w, v): \quad (0, 1, 0, 0, 0, 1, 1, 2).$$

The exact source collection is

$$(E_0)_{\le 6} = 8\rho T_4 - 8(\rho + E)k^2x^2, \quad (L_1)_{\le 3} = 0, \quad (L_2)_{\le 3} = 16Ex, \qquad \rho = r_0H_0 - 1.$$

The source/contact substitutions preserve this grading. After the displayed terms are removed, the scalar and normal lift coefficients have positive weights, hence zero residues. The term k^2x^2 also has a zero-residue coefficient. Thus the soft leading form is $8x^2 + 8r_0H_0T_4$, with the same strict two-pairing circuit frame. The hard block and its only first-normal coupling are those of (BE6), and their determinant is $128r_0H_0 > 0$. Retain the old positive g_4^2 square. The full eight-root remainder Gram is strict. This invokes only the polynomial monic conductor already proved; no opening or contact difference is inverted.

Finally restore the four complete positive packets, the retained positive square, and all pure-coefficient and pure-testing factors. The cases above exhaust both projective angle lines and every projective state. Complex states split into their real and imaginary evaluations. Neither this theorem nor the correction of (BE1) asserts coverage when r is infinitesimal, η unbounded, or e infinitesimal; those are separate source domains.

Chapter 90

Outer tests and state entry when the native r is infinitesimal

Component p364e14238053
Source: R13/baseline/baseline/baseline/baseline/advances/c1/centered_a_outer.tex
SHA256: 364e1423805318f1c173683bd3eb63c27eba6442d586b66998103a7db65671ec
Inventory use: CAO1–CAO8 and all state-entry paragraphs | CAO1-CAO8 and full outer/state lemma | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

90.1 Outer tests and state entry when the native r is infinitesimal

Lemma 90.1. *Use the unchanged physical native C1 source, normalized by $c = 2$ and $\lambda = n = 0$, with all original event signs and positive openings a, t, M, J. Suppose $k, \zeta, r = 1 - a\eta$ are positive infinitesimals and $\eta, e = k\sqrt{\eta/\zeta}$ are positive bounded units. Then the outer-test and state-entry arguments reduce the all-tests problem to precisely*

$$u, v, w > 0 \text{ infinitesimal}, \qquad Z = 1, \qquad w = (W + uY)/(1 + u^2). \tag{CAO1}$$

All other independent projective tests and states admit a full-parent tensor SOS. No comparison of k^2 with r is made, and arbitrarily small positive openings and literal $\beta = 0$ are included.

Source ratios with the common factor cancelled. Here r always denotes the native source parameter, not the event root $uW - Y$. Put

$$K = 1 + k^2, \quad H_c = r + \eta, \quad V_c = k^2\eta\zeta, \quad S_3 = 1 - k^2\eta + tV_c,$$

The two required short native polynomials are

$$\begin{aligned} P_4 &= 1 + k^2 + \eta k^2(a - k^2 - 1) - \zeta k^2 t \\ &\quad + V_c[-\zeta k^2 t^2 + 2k^2 t + a(1 - t) + t], \\ P_d &= (1 + k^2)^2 - a\eta k^4 + \zeta k^2[(k^2 + 1) - (k^2 + 2)t] \\ &\quad + V_c[-\zeta k^2 t^2 + k^2 a(t - 1) + k^2 t + k^2 + 1]. \end{aligned}$$

They are positive bounded units with residue one. The exact source identities, after cancelling the common factor r in the coquotients, are

$$d = \frac{4a\eta P_d}{KH_cS_3}, \qquad Ar = \frac{2a\eta^2 P_4}{KH_cS_3},$$
$$\frac{m}{d} = \frac{e^2 K(1 - t\zeta)(1 + V_c)H_c}{2M\eta P_d}, \qquad \frac{c}{d} = \frac{KH_cS_3}{2a\eta P_d},$$

$$\frac{A}{d}=\frac{\eta P_4}{2rP_d},\qquad\qquad \frac{D}{d}=\frac{KH_c(1+V_c)-J\eta P_d}{2J\eta P_d}. \tag{CAO2}$$

The positive openings obey $M+t=1+k^2r$ and $J+a=1+k^2-k^2\zeta t$; hence a,t,M,J are bounded. Since $a\eta=1-r$, one has

$$\bar{\eta}\geq 1,\qquad \bar{d}=4/\bar{\eta},\qquad Ar\sim 2,\qquad A \text{ positive unbounded.} \tag{CAO3}$$

The numerator of D/d in (CAO2) has residue $\bar{H}_c-\bar{J}\bar{\eta}=1$. Therefore $m/d, c/d, D/d, A/d$ all have positive real lower bounds. This conclusion uses no inverse bound on an opening. Moreover,

$$\begin{cases}\bar{D}=\dfrac{2}{\bar{\eta}(\bar{\eta}-1)}, & \bar{\eta}>1,\\ D \text{ positive unbounded}, & \bar{\eta}=1.\end{cases} \tag{CAO4}$$

The previously proved outer routing still applies. Clip m, c, A, D at positive real multiples of d, retaining all their nonnegative excess original event squares. The resulting coefficients divided by d are bounded. The outer-test argument in `bounded_e_all_tests.tex`, through (BE2)–(BE4), uses only these lower bounds and the independent physical-source admission at noncommuting residue tests. Its exact homogeneous identities and four-root ideal memberships are unchanged. It therefore includes all projective infinity charts, the $(0,\infty)$, $(\infty,0)$ and (∞,∞) corners, and the unit-W, infinitesimal-Z state chart. The original-event SOS on literal zero axes is unchanged as well. This invokes none of the bounded-r scalar or normal conclusions (BE5)–(BE6). Only bounded u and infinitesimal v remain.

An exact centered root identity. Write $D_u=1+u^2$, $D_v=1+v^2$ and introduce

$$h=A^{-1/2},\quad L=Ar,\qquad H=\sqrt{A}(Y-ruW),\quad B=X+uZ,\quad U=uW,\quad V=vZ. \tag{CAO5}$$

Thus $Y=rU+hH$, while h,r are infinitesimal and $L\sim 2$. All coefficient radicals in (CAO5) lie in the coefficient field; B,U,V lie in the testing field. The root H is their finite tensor combination, with no mixed inverse or mixed radical.

For clarity, the complete original finite parent used here is

$$\begin{aligned}F={}&4m(vW+uvY-X-uZ)^2+8D_v(uW-Y)^2+4AD_uD_vY^2\\&+(4D+2d)(uW+uvX-Y-vZ)^2+(4\alpha+2d)D_u(vY-Z)^2\\&+(4\beta+2d)D_v(X+uZ)^2-2dD_uD_vZ^2.\end{aligned}$$

Choose $0<D_*\leq D$ bounded, and let $m_*=0$ or choose a positive real $m_*\leq m$, as specified below. Retain the four nonnegative original event contributions with weights $m-m_*, D-D_*, \alpha, \beta$. For $C=4D_*+2d$, the exact remaining parent is

$$\begin{aligned}F_*={}&8D_v[(1-r)U-hH]^2+4D_uD_vH^2+4LrD_uD_vU^2+8LhD_uD_vUH\\&+C[(1-r)U-hH+uvB-D_uV]^2+2dD_vB^2\\&+2dD_u\{v^2(rU+hH)^2-2(rU+hH)V-V^2\}+4m_*R_m^2,\\R_m={}&vW+uv(rU+hH)-B.\end{aligned} \tag{CAO6}$$

In particular, the centered A contribution is the exact identity

$$4AD_uD_vY^2=4D_uD_vH^2+4LrD_uD_vU^2+8LhD_uD_vUH.$$

This retains its complete hard root. The last two coefficients have zero residue. Formula (CAO6) follows directly from the original parent and includes the complete signed core; it is not an eliminated scalar or a restriction to state normals. Every displayed coefficient is bounded. The term involving Z in that core has already cancelled to the displayed expression in $V=vZ$.

Every state when u is a bounded unit. Set $m_* = 0$, so the entire original m event is retained. Formula (CAO6) is an exact Gram allocation on (H, B, U, V). At infinitesimal v its residue is

$$4D_uH^2 + 2dB^2 + 8U^2 + C(U - D_uV)^2 - 2dD_uV^2. \tag{CAO7}$$

Residues are understood in this and the following determinant calculations. The U, V determinant is

$$D_u\{8CD_u - 2d(8 + C)\}.$$

If $\bar{\eta} > 1$, take $D_* = D$, which is bounded by (CAO4). Then $C = 8/(\bar{\eta} - 1)$ in residue and this determinant is

$$\frac{64u^2(1 + u^2)}{\bar{\eta} - 1} > 0.$$

If $\bar{\eta} = 1$, choose a sufficiently large positive real D_*; this is permitted because D is unbounded. The determinant is then

$$32(1 + u^2)(D_*u^2 + 2u^2 - 2) > 0.$$

The first diagonal is $8 + C > 0$, so the complete four-root residue Gram is positive definite in both cases. The cap is a real constant chosen from the nonzero testing residue of u; no division of tensor elements is introduced. The bounded full Gram is therefore tensor SOS, and restoring the retained events proves the claim for every state in this chart.

The remaining unit-w state chart. Suppose u, v are infinitesimal and w is a unit. Then W is a unit on the negative-core branch. When Z is infinitesimal the inherited (BE4) argument already applies. Otherwise normalize the state so that Z is a unit and put $q = W/Z$, a pure-testing unit. Choose a positive real lower bound $m_* \le m$. Now

$$R_m = qV + uv(rU + hH) - B,$$

so (CAO6) is again an exact bounded Gram on (H, B, U, V), with residue

$$4H^2 + 2dB^2 + 4m_*(qV - B)^2 + 8U^2 + C(U - V)^2 - 2dV^2. \tag{CAO8}$$

The B pivot is positive. Its Schur complement adds

$$\mu V^2, \qquad \mu = \frac{4dm_*q^2}{d + 2m_*} > 0$$

to the U, V block. This is just a residue determinant computation; no mixed inverse is used to define a root or a full-parent coefficient. If $\bar{\eta} > 1$, take $D_* = D$ as before. The determinant becomes

$$\frac{8\bar{\eta}\mu}{\bar{\eta} - 1} > 0.$$

If $\bar{\eta} = 1$, choose a sufficiently large positive real cap D_* so that $\mu(C+8) - 64 > 0$. The unboundedness of D permits this choice. Thus the complete residue Gram in (CAO8) is strict, including all hard-soft terms, and restoring the retained events gives the full-parent tensor SOS.

Completion of state entry. Normalize the state by a maximal coordinate and first remove the nonnegative pure-testing-core branch, which has its direct original SOS. On the negative-core branch with v infinitesimal, the residue core is

$$2\{(X + uZ)^2 + (uW - Y)^2\}.$$

Consequently these two roots have zero residue. If w and Z were both infinitesimal, the exact inverse relations

$$W = w + \frac{u(uW - Y)}{D_u}, \quad Y = uw - \frac{uW - Y}{D_u}, \quad X = (X + uZ) - uZ$$

would make every normalized state coordinate infinitesimal. Hence small w forces Z to be a unit. The bounded-unit u chart was settled by (CAO7), and all unit-w states by (CAO8) and (BE4). Normalize the only remaining states by $Z = 1$. The same exact pure-testing Schur identity (SD5) and testing sign gauges give $u, v, w > 0$. This is exactly (CAO1). Literal axes and literal state zeros have already been included. Complex states split into their real and imaginary evaluations because the parent is real.

The accompanying `check_centered_a_outer.py` independently checks the native source cancellations, the exact selected-parent identity, both complete root allocations, their residues and all displayed determinants. It does not import the contact scalar calculations used for the remaining chart.

Chapter 91

An infinitesimal native r with unit η and unit e

Component p3763434080c2
Source: R13/baseline/baseline/baseline/baseline/advances/c1/centered_a_unit_e.tex
SHA256: 3763434080c27994309135baaeef0216f6916b0507db82e8a646169a209c2016
Inventory use: CA1–CA8, complete raw scalar/normal dictionary | CA1-CA8 | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

91.1 An infinitesimal native r with unit η and unit e

Theorem 91.1 (The centered-A unit-e slice)**.** *Use the physical native C1 source with its unchanged original events, $c = 2$, $\lambda = n = 0$, all positive openings a, t, M, J, and all original nonnegative signs, including literal $\beta = 0$. Suppose $k, \zeta, r = 1 - a\eta$ are positive infinitesimals, while η and $e = k\sqrt{\eta/\zeta}$ are positive bounded units. Then the full parent is tensor SOS at every independent projective test and every real or complex state. No bound on k^2/r is imposed.*

The theorem is restricted to unit e. Its proof uses δ^{-1} as a bounded pure coefficient number; it does not assert the cases of infinitesimal or unbounded e.

Centering the sole large event. The exact native contact has $Y_c = ru_cW_c$. Consequently

$$L_A = Y - ruW$$

vanishes at contact with no asymptotic replacement. The native formulas give a a positive unit, A unbounded and Ar a positive unit of residue 2. Put $A_* = 1/a$, a positive bounded pure coefficient unit. The complete packet

$$4(A - A_*)D_uD_vL_A^2 \tag{CA1}$$

is positive and has zero value and full first derivative at the exact native contact. Retain it as a square.

It is essential to center this packet before taking residues. The remaining A channel is exactly

$$4A_*D_uD_vL_A^2 + 4(Ar)D_uD_v\{2uWY - ru^2W^2\}. \tag{CA2}$$

Only Ar and r occur in its coefficients. In particular this is a bounded expression for every positive rate of r relative to k.

A source chart with no hidden k^2/r quotient. Keep the exact contact variables $\delta, k, E = b^2$ from (SD6), and put

$$K = 1 + k^2, \quad \ell = K - aE = Qr > 0, \quad Q = K^2 - \ell k^2, \quad a = (K - \ell)/E.$$

Here δ, E, a are positive bounded units and k, ℓ are positive infinitesimals. The exact source substitution is

$$\eta = KE/Q, \quad r = \ell/Q, \quad \Delta = \delta Q/K, \quad \zeta = \delta^2 k^2 QE/K, \quad R = K^2/Q, \quad T = K - k^2 t\zeta. \tag{CA3}$$

All denominators in (CA3) are positive pure coefficient units. For $i = 2, b, 3, 5, \alpha, d$ let $\widetilde{c}_i = c_i/\ell$, defined by cancelling the explicit $L_c = \ell/Q$ factor in the short native formulas. This cancellation gives rational expressions with unit denominators; it does not leave an inverse of ℓ. For example, writing $H_c = r + \eta$ and $V_c = k^2\eta\zeta$,

$$\widetilde{c}_3 = \frac{2KH_cQ_c(1 - k^2\eta + tV_c)}{Q}, \qquad \widetilde{c}_d = \frac{4Q_cP_d}{Q}.$$

Their residues are $2E$ for $i = 2, b, 3, 5, \alpha$, and 4 for $i = d$; c_4 has residue 2.

Define the following bounded pure coefficient quantities:

$$\begin{aligned} B_2 &= \frac{8a\widetilde{c}_2}{R\widetilde{c}_3\Delta^2}, & B_b &= \frac{8a\widetilde{c}_b}{R\widetilde{c}_3\Delta^2}, \\ B_3 &= 8a/T, & B_5 &= \frac{8a\widetilde{c}_5}{T\widetilde{c}_3}, \\ L &= Ar = \frac{2a\eta^2 c_4}{Q\widetilde{c}_3}, & d &= \frac{2a\eta\widetilde{c}_d}{\widetilde{c}_3}, \\ \alpha &= \frac{2a\Delta^2 k^6\eta^2\widetilde{c}_\alpha}{\widetilde{c}_3}. \end{aligned} \tag{CA4}$$

These formulas explicitly account for the cancellation needed below.

The exact full remainder, before any collection. Retain the four original (SD3) packets, divided by the positive source factor $\Delta^2\Lambda$, and retain (CA1). Use the unchanged pure testing coordinate $w = (W + uY)/D_u$ and the hard normals N, M_B. For $Z = 1$ put $p = t/R$, $h = a/T$ and

$$q_A = vD_uw, \quad q_B = v(D_u - uN - puD_uvw),$$

$$W = w + u(M_B + hq_B)/D_u, \qquad Y = uw - (M_B + hq_B)/D_u.$$

The complete remainder is exactly

$$\begin{aligned} \mathscr{G} = {} & B_2\{N^2 - 2q_AN + (1-p)q_A^2\} + B_bD_v\{N^2 + 2q_AN + pq_A^2\} \\ & + B_3D_v\{M_B^2 + 2q_BM_B + hq_B^2\} + B_5\{M_B^2 - 2q_BM_B + (1-h)q_B^2\} \\ & + 4A_*D_uD_v(Y - ruW)^2 + 4LD_uD_v(2uWY - ru^2W^2) \\ & + 4\alpha D_u(vY - 1)^2 + 2dD_u\{(vY - 1)^2 - D_v\}. \end{aligned} \tag{CA5}$$

There are no normal terms above degree two. Every coefficient of (CA5) is bounded. The only testing denominator is a power of the positive unit D_u; every source denominator in (CA3)–(CA5) is a positive unit. The possibly unbounded opening ratios remain exclusively in the four retained positive packets.

The exact low-weight collection. The outer/state lemma in `centered_a_outer.tex` reduces the independent tests to $Z = 1$ and u, w, v infinitesimal, with $w > 0$ on the negative pure-core branch. Use the unchanged circuit monomials

$$A_0 = wv, \quad B_0 = \delta v, \quad C_0 = \delta Euw, \quad D_0 = \delta^2 k^3 E,$$

$$x = B_0 - C_0, \quad y = A_0 - kC_0, \quad z = uB_0 - kC_0, \quad h_0 = kC_0 - D_0.$$

Let q_1, q_2 be the same two separated radical pairings as in the contact–conductor proof, and let

$$T_4 = y^2 + z^2 + h_0^2 - 2yz + 2q_1^2.$$

Give $(\delta, k, E, \ell, t, u, w, v)$ weights $(0,1,0,1,0,1,1,2)$. Then x has weight two and y, z, h_0, q_1, q_2 have weight three. Denote the scalar of (CA5) by S, and its two complete first normals by L_N, L_M. Direct multiplication of (CA3)–(CA5) through the indicated weights gives

$$\begin{aligned} S_{\le 6} &= \frac{4P}{E^4\delta^2}x^2 - \frac{16(k+u)}{E^2\delta^2}xz + \frac{8}{E\delta^2}T_4, \\ (L_N)_{\le 3} &= 0, \\ (L_M)_{\le 3} &= \frac{8\{E^2+2E-2\ell(E+1)\}}{E^2\delta}\,x, \end{aligned} \tag{CA6}$$

where the short coefficient polynomial is

$$\begin{aligned} P = {} & E^2(E+4) - E(E^2+6E+4)\ell + 2(E+1)(E+2)\ell^2 \\ & + E^2(8-3E)k^2 + 4E^2ku + E^2(E+4)u^2. \end{aligned}$$

The full hard-normal block, from the quadratic part of (CA5), has residue

$$\operatorname{diag}\left(\frac{16}{E\delta^2}, \frac{4(E^2+4)}{E}\right). \tag{CA7}$$

Here and in the residue Grams, E, δ denote their positive residues. Formula (CA6), rather than a large expansion, is the finite coefficient identity used by the proof. Its 37 raw scalar monomials collect to the displayed three terms. The exact checker reconstructs the six cancelled native coquotients, the entire expression (CA5), and both complete normals before making this collection.

Why all higher terms have harmless coefficients. The scalar of (CA5), after clearing the stated units, belongs to $(A_0, B_0, C_0)^2 + (D_0^2)$. This follows directly from $q_A = wvD_u$, $q_B|_{N=M_B=0} \in (v)$, and Y, W in (CA5): the centered scalar $Y - ruW$ lies in (v, uw); its square is in the same raw monomial square ideal since δ, E are units. The sole constant state term comes from α, which has the factor k^6, and is a bounded unit multiple of D_0^2. Both full first normals belong to (A_0, B_0, C_0) by the same displayed expression.

All retained packets vanish to first order at the exact contact

$$(u, w, v, N, M_B) = (k, \delta k, \delta k^2 E, 0, 0).$$

Thus the scalar belongs to the square of its contact ideal and the first normals belong to that ideal. The unchanged monic contact–conductor identities therefore apply, over the coefficient ring in which only the displayed positive units have been inverted. They never divide by k, ℓ, M, J, t, or a contact difference; the displayed $1/a$ is the inverse of a known positive unit. The contact substitutions preserve the new weights. After (CA6) is subtracted, scalar weights are at least seven and normal weights at least four. Their lifted coefficients consequently have positive weights, hence zero residues. Clearing the finitely many unit denominators before this polynomial lift and restoring them afterward changes no residue sign and introduces no mixed inverse.

Strictness, full normals, and restoration. The leading scalar is

$$\frac{4(E+4)}{E^2\delta^2}x^2 + \frac{8}{E\delta^2}T_4.$$

The standard two-pairing circuit supplies a strict Gram on (y, z, h_0, q_1, q_2). The only nonzero first-normal residue is $8(E+2)x/(E\delta)$, the coefficient of the full normal M_B. Accordingly the coupled (M_B, x) Gram is

$$\begin{pmatrix} 4(E^2+4)/E & 4(E+2)/(E\delta) \\ 4(E+2)/(E\delta) & 4(E+4)/(E^2\delta^2) \end{pmatrix}, \qquad \det = \frac{256}{E^3\delta^2} > 0. \tag{CA8}$$

Together with the independent N pivot in (CA7), this gives a bounded strict eight-root Gram for the entire remainder (CA5). Restore (CA1), all four original positive packets, and all separated normalization factors. The outer/state lemma exhausts the remaining tests; real and imaginary state evaluations give the complex case.

This proves precisely the stated unit-e, unit-η, infinitesimal-r slice. Combined with the bounded-unit-r theorem, it covers every positive r when η, e are bounded units and k, ζ infinitesimal. It makes no assertion about unbounded η or either nonunit alternative for e.

Chapter 92

The joint rational C2 opening: a full-parent certificate

Component p8f912583feaa
Source: R13/baseline/baseline/baseline/baseline/advances/c2_joint/joint_centered_packets.tex
SHA256: 8f912583feaa7f930694a5996b34ad984e1af7b157dbfae029f41c58a4a2ef70
Inventory use: JC1–JC13 complete simultaneous r,gInf source with both centered excess packets

92.1 The joint rational C2 opening: a full-parent certificate

Theorem 92.1 (Both event openings may close)**.** *Use the exact rational C2 source and original parent of general_single_angle.tex. Suppose $x > 0$ is infinitesimal, k, b are positive bounded units, $0 < r < 1$, $0 < h < b$, and $g = b - 1 + r + xh > 0$. Suppose r, g are infinitesimal, $n \geq 0$, and the source is physical. Then the entire original parent is tensor SOS at every independent real or complex projective testing state and every independent projective pair of testing angles. No lower bound on either h or $b - h$ is required.*

Together with the three-opening and last-opening theorems this removes the condition that at least one of r, g be a unit in the bounded-k, b rational C2 source family. It does not classify all physical one-weak C2 sources, nor admit a nonunit k or unbounded b. All tensor-ring conventions and the strict bounded tensor Gram lemma are those of the authoritative package. In particular, a root vector need not be linearly independent, and its entries may be unbounded.

Source bounds, including both complementary openings. Write $\rho = \sqrt{x}$, $\eta = h/b$, $\kappa = b/k$, $K = k_0 > 0$, and $D_K = 1 + K^2$. Since r, g are infinitesimal, $b_0 = 1$. The source formulas give

$$(nr)_0 = \frac{1}{4D_K}, \qquad (mg)_0 = \frac{1}{4}, \qquad \Delta_0 = 2D_K.$$

Thus n, m are positive unbounded pure coefficient quantities, nr, mg are bounded units, and nr^2, mg^2 are infinitesimal. In particular $m - 1 > 0$. The existing opening cancellation identities, with $T_C = xC(b - h)$ and $T_A = xAh$, give bounded positive units

$$(T_C)_0 = (T_A)_0 = \frac{K^2}{4D_K}, \qquad (D/x)_0 = \frac{1}{4K^2 D_K}, \quad (\alpha/x)_0 = \frac{K^2}{4D_K}.$$

They hold even if h or $b - h$ is infinitesimal. Specifically,

$$xC(1-\eta) = T_C/b, \quad xA\eta = T_A/b, \quad xC(1-\eta)^2 = T_C(b-h)/b^2, \quad xA\eta^2 = T_A h/b^2. \tag{JC1}$$

Put $H_c = x(C + A)$. This quantity is bounded below by a positive real, although it can be unbounded. Its reciprocal and reciprocal square root are bounded pure coefficient elements.

Center the excess positive packets before taking normals. Retain exactly the original amplitudes

$$M = vW - X + uvY - uZ, \qquad J = uW + uvX - Y - vZ, \qquad D_u = 1 + u^2, \quad D_v = 1 + v^2.$$

Let F denote the full original parent, including its signed core H. Define G by the following exact polynomial identity:

$$F = G + 4nD_uD_v(X + ruZ)^2 + 4(m-1)(M - guZ)^2. \tag{JC2}$$

An explicit expression, requiring no elimination or mixed inverse, is

$$\begin{aligned} G = {} & 4M^2 + 4CD_v(uW - Y)^2 + 4AD_uD_vY^2 + 4DJ^2 + 4\alpha D_u(vY - Z)^2 + H \\ & - 8nrD_uD_vuZX - 4nr^2D_uD_vu^2Z^2 \\ & + 8(m-1)g\,uZM - 4(m-1)g^2u^2Z^2. \end{aligned} \tag{JC3}$$

Every coefficient in (JC3) except C, A is bounded. At the exact original contact

$$(u, v; W, X, Y, Z) = (\rho, k; \kappa\rho, -r\rho, xh/k, 1)$$

both subtracted squares and all of their first derivatives vanish. Consequently G has exactly the same zero and full stationary contact as F. This statement is essential: merely capping the uncentered original packets would not preserve the contact.

On the finite $u = 0$ residue axis use

$$\begin{aligned} X_g = -ruZ, \qquad Y_g = \eta uW, \qquad E_X = \sqrt{n}(X + ruZ), \\ E_Y = \sqrt{C + A}(Y - \eta uW), \qquad E_M = \sqrt{m-1}(M - guZ). \end{aligned} \tag{JC4}$$

Expand only G in E_X, E_Y and retain $4E_M^2$ as written. The root E_M is allowed to be redundant. No substitution for W is made. In particular, the positive C packet introduces no coefficient of the form Cu^2/n or Cu^2/m.

The complete graph scalar and its global leading inequality. Write $f = G|_g = f_{ww}W^2 + f_{wz}WZ + f_{zz}Z^2$. The same exact monic divisions as in the prior single-angle proof give

$$f = aP^2 + b_sB^2 + c_sV^2 + eE^2 + g_sBV, \qquad (P, B, V, E) = (uW/\rho, uZ, vW, \rho Z), \tag{JC5}$$

where

$$a = x\frac{f_{ww} - f_{ww}|_{u=0}}{u^2}, \quad b_s = \frac{f_{zz} - f_{zz}|_{u=0}}{u^2}, \quad c_s = \frac{f_{ww}|_{u=0}}{v^2}, \quad e = \frac{f_{zz}|_{u=0}}{x}, \quad g_s = \frac{f_{wz}}{uv}.$$

These are polynomial coquotients, not inverses of testing infinitesimals. Their coefficients are bounded by (JC1)–(JC3). The exact constant $f_{zz}|_{u=0} = 4Dv^2 + 4\alpha$ supplies the x factor. For finite v their residues are

$$q(v) = \frac{K^2(1 + v^2)}{D_K}, \qquad b_s = 4, \quad c_s = 4, \quad g_s = -10, \qquad e(v) = \frac{v^2/K^2 + K^2}{D_K}. \tag{JC6}$$

Thus the leading scalar, using the exact identity $BV = vPE$, is

$$qP^2 - 2vPE + eE^2 + 4(V - B)^2. \tag{JC7}$$

The key identity is now an identity for the graph of a full bounded remainder, rather than for the overconstrained graph from the earlier frontier note:

$$q(v)e(v) - v^2 = \frac{(v^2 - K^2)^2}{D_K^2}. \tag{JC8}$$

If v is a finite nonzero residue with $v^2 \neq K^2$, then $10|v| < 8|v| + 2\sqrt{qe}$. Splitting the exact $BV = vPE$ cross term with the pure testing function

$$\theta = \frac{4|v|}{4|v| + \sqrt{q(v)e(v)}}$$

gives strictly positive residue blocks on (B, V) and (P, E). At $v_0 = 0$, allocate the entire cross term using $BV = vPE$; (JC5) then has strictly positive diagonal residue $(q(0), 4, 4, e(0))$ and zero cross residues. There is no exceptional $n_0 = 0$ corner in the joint regime.

Here is also a global equality and Hessian check needed at the matched angles. For $z = v^2 > 0$ and $s > 0$, set

$$\mathcal{D}(s, z) = (qs + 4z)(4s + e) - 25zs.$$

Exactly,

$$\begin{aligned} \frac{\mathcal{D}(s,z)}{s} &= 4qs + 4ez/s + qe - 9z \\ &= 4\big(\sqrt{qs} - \sqrt{ez/s}\big)^2 + (\sqrt{qe} - \sqrt{z})(\sqrt{qe} + 9\sqrt{z}). \end{aligned} \tag{JC9}$$

The second term is nonnegative by (JC8); its equality forces $z = K^2$. The first then forces $s = 1$. Moreover the second term is

$$\frac{(z - K^2)^2}{D_K^2} \frac{\sqrt{qe} + 9\sqrt{z}}{\sqrt{qe} + \sqrt{z}},$$

whose quadratic coefficient at contact is strictly positive. The first square supplies a nonzero s derivative there. Hence $\mathcal{D}$ has a positive definite Hessian at $(1, K^2)$. These radicals are used only in the residue-field inequality; no radical of an element mixing coefficient and testing fields is used.

Restore every normal, with bounded coefficients. From (JC2)–(JC4) the entire parent is exactly

$$F = f + E_X L_X + E_Y L_Y + H_{XX} E_X^2 + 2H_{XY} E_X E_Y + H_{YY} E_Y^2 + 4E_M^2. \tag{JC10}$$

The first normals are $L_X = G_X|_g/\sqrt{n}$ and $L_Y = G_Y|_g/\sqrt{C + A}$. The only unbounded weights C, A do not enter the X row. Thus

$$L_X = V\ell_w + B\ell_z$$

with bounded coefficient functions carrying the infinitesimal pure coefficient factor $1/\sqrt{n}$. Both have identically zero residue on the entire finite $u = 0$ axis, including $v = 0$.

For the second normal the exact form is

$$L_Y = \frac{1}{\sqrt{H_c}} \{P(B_2 + xe_w) + B(\rho u e_z) + Ee_*\}, \tag{JC11}$$

where

$$B_2 = 8D_v\{(xA)\eta D_u - (xC)(1 - \eta)\}, \qquad e_* = 8v(D - \alpha).$$

All e_w, e_z are bounded: all new terms in (JC3) have bounded coefficients, and their Y derivative has an additional $u^2 Z$ factor. Equation (JC1) makes B_2 bounded with identically zero residue, even at either closing complementary opening. Thus every coefficient in (JC11) has zero residue.

The entire hard block is bounded and has residue

$$\operatorname{diag}(4D_{v,0}, 4D_{v,0}, 4) \quad \text{on } (E_X, E_Y, E_M). \tag{JC12}$$

The first entry comes from the retained centered n packet; the bounded G contribution is divided by n. The second is

$$4D_v \frac{C + AD_u}{C + A} = 4D_v \left(1 + \frac{A}{C + A} u^2\right)$$

plus bounded terms divided by $C + A$. The mixed entry contains $1/(\sqrt{n}\sqrt{C + A})$ times a bounded function. The third entry is the unexpanded centered m packet. This proves the full-parent assertion at all remote finite-axis tests, by the strict Gram in (JC5) and the zero-residue first normals.

The matched finite-angle residues. At $v_0 = K$, normalize $Z = 1$ and suppose W is infinitesimal. Use exactly the prior contact conductors

$$R = (uW - \kappa x)/\rho, \quad S = u - \kappa^{-1}W, \quad U = \rho(v - k), \quad T = u(v - k),$$

with exact identities

$$u^2 - x = uS + \kappa^{-1}\rho R, \quad W^2 - \kappa^2 x = -\kappa WS + \kappa\rho R, \quad uW - \kappa x = \rho R. \tag{JC13}$$

Here $f = p^2 Q_b/x + B_b$, with bounded B_b and

$$Q_b = 4D_v\{(xC)(1 - \eta)^2 + (xA)\eta^2 D_u\}.$$

Boundedness follows from (JC1); the remaining terms use only the bounded coefficients of (JC3). The full exact stationary contact of G proved after (JC3) supplies every vanishing required by the prior monic-contact-division argument. Successive monic differences in $u^2 - x, W^2 - \kappa^2 x, uW - \kappa x, v - k$ therefore give a bounded Gram on R, S, U, T.

For precision, its strict residue follows from the same explicit Schur-complement step in that argument, now with

$$\Phi = q(v)s^2w^2 + 4s^2 + 4v^2w^2 - 10vsw + e(v).$$

Completing the w square gives

$$\Phi = (qs^2 + 4v^2)\left(w - \frac{5vs}{qs^2 + 4v^2}\right)^2 + \frac{\mathcal{D}(s^2, v^2)}{qs^2 + 4v^2}.$$

Equation (JC9) proves a positive definite three-variable Hessian at $(s, w, v) = (1, 1/K, K)$. Consequently the residue before the conductor-zero correction has form

$$q(K)R^2 + 4S^2 + A_1RU + B_1ST + E_1U^2, \qquad E_1 > A_1^2/(4q(K)) + B_1^2/16.$$

Choose a real δ strictly between $B_1^2/16$ and $E_1 - A_1^2/(4q(K))$, and add δ times the exact zero

$$T^2 - U^2 - ST(v - k) - \kappa^{-1}RU(v - k) = 0.$$

The two residue blocks (R, U) and (S, T) are strictly positive.

An explicit equivalent residue check can be used in place of this Schur choice. The capped current has the exact conductor lift

$$(M - gu)|_g = -v\kappa S + \kappa T + \eta uv\rho R + \eta u\kappa\rho U.$$

Retaining its T^2 term, rather than first moving that term to U^2, gives the scalar residue

$$K^2R^2 + \frac{2(K^2 - 1)}{D_K}RU + \frac{U^2}{K^2} + 4(S - T/K)^2.$$

Adding $2/D_K^2$ times the same conductor zero leaves the (R, U) Schur margin $2/D_K^2 > 0$, and leaves $4(S - T/K)^2 + (2/D_K^2)T^2$ on (S, T), also strictly positive.

The full-normal lifts are also the ones actually required for (JC10), rather than scalar-only lifts. Write $L_X = uA_1' + WB_1' = -\kappa B_1'S + u(A_1' + \kappa B_1')$. Both functions and all their bounded monic coquotients retain $1/\sqrt{n}$. Exact $G_X = 0$ at contact supplies vanishing of $A_1' + \kappa B_1'$. Using (JC13) gives zero-residue coefficients on R, S, U, T. For (JC11), write its numerator as $RB_2 + \rho E_2$. Both B_2, E_2 have identically zero leading functions and E_2 vanishes at contact. The same monic divisions give factors $x, \rho u, \rho W$, or a zero-leading-function $(v - k)$ coquotient; division by $\sqrt{H_c}$ preserves boundedness and zero residue. Thus the complete seven-root Gram on $(R, S, U, T, E_X, E_Y, E_M)$ is strict in residue.

If W/Z is not infinitesimal, normalize $W = 1$ and keep Z bounded. Then the scalar has a strict residue on $(P, 1)$: its positive constant contribution is $4K^2$. All normalized first normals have zero residue as above. For $Z = 0$, the original parent is already the literal sum of its event packets and three positive core squares. The simultaneous sign copy gives $v_0 = -K$.

Every other independent test. The outer-chart proof need not use the centered decomposition. Use the original parent, cap C, A at σ/x, where $\sigma = \min(1, xC, xA)$ is a positive bounded unit, and cap m at 1. Retain the original nonnegative excess packets and the entire original n packet. The four exact outer-core identities in the prior single-angle theorem apply without change: their strict residues require only $\sigma_0 > 0$, the selected $m = 1$, and $(D/x)_0 > 0$. They never bound n. This covers the finite unit u axis, both reciprocal axes, all their intersections, and literal projective ends. The inherited physical noncommuting-residue theorem covers the remaining tests.

These charts and the entire finite $u = 0$ axis exhaust all independent projective measurement pairs. All state charts were restored above; complex states split into their real and imaginary parts as in the prior proof. Every inverse or radical used in a tensor coefficient is pure coefficient or pure testing. No comparison of a source infinitesimal with a testing infinitesimal, and no mixed inverse or radical, occurs. Restoring the retained packets proves the theorem.

Chapter 93

Strong C2 pitchfork with arbitrary complementary scales

Component `pf7943b34a7dc`
Source: `R13/baseline/baseline/baseline/baseline/advances/c2_strong/strong_complement_pitchfork.tex`
SHA256: `f7943b34a7dce78326e3e9ae90456999195c03136f243662c3381e74e9e416bb`
Inventory use: SC1–SC23 unbounded C or alpha and all upper event scales, full three normals, strict real boundary Schur, complete cap/corner outer proof

93.1 Strong C2 pitchfork with arbitrary complementary scales

Theorem 93.1 (No upper bound on the two complementary events)**.** *Let $K_c, K_t \supset \mathbb{R}$ be independent real closed fields. Let the original canonical C2 parent be physical over K_c, with $\lambda = \beta = 0$, d a positive bounded unit, $C, \alpha > 0$ having bounded reciprocals, and $n, m, A, D \geq 0$ arbitrary. Suppose it has an actual finite noncommuting stationary zero with $u_c = k > 0$ infinitesimal and $v_c = a > 0$ a bounded unit. Then the entire original parent is tensor SOS at every independent projective measurement pair and every real or complex projective state. There is no upper bound or relative-scale condition on any event weight.*

The case of bounded C, α is the existing three-large-event theorem. We prove both remaining cases, without modifying the parent or identifying the source and testing fields. We use

$$\begin{gathered} D_u = 1 + u^2, \quad D_v = 1 + v^2, \quad r = uW - Y, \quad p = vY - Z, \\ J = uW + uvX - Y - vZ, \quad R_2 = vW - X + up, \quad B = X + uZ, \\ H = 2\{J^2 + D_u p^2 + D_v B^2 - D_u D_v Z^2\}, \\ F = dH + 4nD_u D_v X^2 + 4mR_2^2 + 4CD_v r^2 + 4AD_u D_v Y^2 + 4DJ^2 + 4\alpha D_u p^2. \end{gathered} \tag{SC1}$$

Every square in this formula retains its original coefficient.

Necessary reductions from the full parent. Cap only C, α at positive bounded units in the initial hard state estimate. At the actual angles the bounded selected form is bounded below by $b(X^2 + Y^2 + Z^2) - B_0 k^2 W^2$, with positive real b, B_0. Every retained packet is nonnegative. Hence the contact has $W_c \neq 0$; on setting $W_c = 1$,

$$X_c, Y_c, Z_c = O(k), \qquad m = O(k^2).$$

Each positive event packet is $O(k^2)$ as well. Thus, writing $y = Y_c/k$, $z = Z_c/k$, $x_1 = X_c/k$, $j = J_c/k$,

$$Ay^2,\ Dj^2,\ C(1-y)^2,\ \alpha(ay - z)^2 = O(1), \qquad j = 1 - y - az + aX_c. \tag{SC2}$$

If C is unbounded then $\bar{y}=1$, so A is bounded. If D were also unbounded, j,z would be infinitesimal, and the residue of

$$H_c/(2k^2)=j^2+(1+k^2)(ay-z)^2+(1+a^2)(x_1+Z_c)^2-(1+k^2)(1+a^2)z^2$$

would be $\bar{a}^2+(1+\bar{a}^2)\bar{x}_1^2>0$. This contradicts $H_c\le 0$ at the zero of (SC1). Therefore C unbounded forces both A,D bounded. If α is unbounded and A were unbounded, (SC2) would make y,z infinitesimal and $\bar{j}=1$; the same residue would instead be $1+(1+\bar{a}^2)\bar{x}_1^2>0$. Thus α unbounded forces A bounded. It remains to prove precisely

$$\text{(I) } C \text{ unbounded, } A,D \text{ bounded;} \qquad \text{(II) } C \text{ bounded, } \alpha \text{ unbounded, } A \text{ bounded.} \tag{SC3}$$

Uniform stationary scale entry. Set $U=k^2$, $V=a^2$, $N=d+2n$, $\theta=n/N$, and

$$t_2=d+2(\alpha+m+A),\quad T=t_3=d+2(C+D),\quad t_4=d+2(\alpha+m),$$
$$t_6=d+2(\alpha+C+A),\quad t_8=d+2(C+A+D).$$

The exact original Laurent stationary equations are

$$\begin{aligned} U^2At_4\{\theta T+C(1+2D/N)V\}&=mV\{\alpha t_8+Dt_6V\},\\ \alpha A(1+2m/N)T+\theta ATt_4U&=\{mDt_6/U+CDt_2\}V^2.\end{aligned} \tag{SC4}$$

Define only pure coefficient quantities

$$\delta=CD/T,\quad e=(D/T)(t_6/\alpha),\quad b=t_2/\alpha,\quad \mathcal{B}=\theta+(C/T)(1+2D/N)V,\quad \mathcal{G}=t_8/T+eV.$$

Because one of C,D is bounded in either case (SC3), $\delta\le\min(C,D)/2$ is bounded. The identities

$$e=(D/T)\{2+(d+2A)/\alpha\}+2\delta/\alpha,\quad \mathcal{G}=1+2A/T+eV,\quad \mathcal{B}=\theta+(C/T)V+(2\delta/N)V$$

show boundedness of all these quantities. The quantities $b,\mathcal{G},t_4/\alpha$ are positive bounded units. So is $\mathcal{B}$: since C,N have bounded reciprocals and d is bounded,

$$\frac{T}{C(N+2D)}\le\frac{d}{CN}+\frac{2}{N}+\frac{1}{C}=O(1),$$

and consequently

$$\theta+(C/T)(1+2D/N)=\frac{n+C(N+2D)/T}{d+2n}\ge\frac{n+c_0}{d+2n}$$

for a positive real c_0. The bounded-unit factor V preserves this lower bound. Dividing (SC4) by αT gives exactly

$$\frac{m}{U^2}=A\rho,\qquad A\mathcal{H}=\delta bV^2,\qquad \rho=\frac{(t_4/\alpha)\mathcal{B}}{V\mathcal{G}},\quad \mathcal{H}=1+2m/N+\theta(t_4/\alpha)U-U\rho eV^2. \tag{SC5}$$

Here ρ is a positive bounded unit and $\mathcal{H}=1+O(U)$. These identities were obtained before inverting A or D. They show $A\asymp\delta$. Since C has a bounded reciprocal, δ infinitesimal implies D infinitesimal. If A were infinitesimal, therefore, so would D,m be. But the exact physical test

$$u=v=1,\qquad (W,X,Y,Z)=(1,0,1,1)$$

has $X=r=p=0$, $R_2=1$, $J=-1$, $H=-2$, and hence $F=-2d+4m+16A+4D$. Its residue would be negative. We conclude

$$A\asymp 1,\qquad D^{-1}=O(1),\qquad m/k^4\asymp 1,\qquad \gamma=4ma^2/k^4\asymp 1. \tag{SC6}$$

In case (I), D is a bounded unit; in case (II), it may be arbitrarily large. In both cases δ is a positive bounded unit.

Full state equations and bounded graph data. For later use, the three entire state derivatives of (SC1) are

$$\begin{aligned} F_X/4 &= D_v[d+2nD_u]X + (d+2D)uvJ + duD_vZ - 2mR_2,\\ F_Y/4 &= -(d+2D)J + (d+2\alpha)vD_up - 2CD_vr + 2AD_uD_vY + 2muvR_2,\\ F_Z/4 &= -(d+2D)vJ - (d+2\alpha)D_up + duD_vB - dD_uD_vZ - 2muR_2. \end{aligned} \tag{SC7}$$

In case (I), the Z equation first yields $\alpha p_c = O(k)$, because D is bounded and every core coordinate is $O(k)$. Then the Y equation yields $Cr_c = O(k)$. In case (II), C, A are bounded, so the Y, Z equations are a system with bounded right side $O(k)$ and unit coefficient matrix

$$\begin{pmatrix} -1 & a \\ -a & -1 \end{pmatrix}$$

on $((d+2D)J_c, (d+2\alpha)(1+k^2)p_c)$. They yield $DJ_c, \alpha p_c = O(k)$. Finally the X equation in either case yields $NX_c = O(k^2)$. Thus put $x = X_c/k^2$ and

$$\begin{aligned} &\text{(I)} \quad r_0 = r_c/k,\ p_0 = p_c/k, \qquad Nx, Cr_0, \alpha p_0 = O(1),\\ &\text{(II)} \quad j_0 = J_c/k,\ p_0 = p_c/k, \qquad Nx, Dj_0, \alpha p_0 = O(1). \end{aligned} \tag{SC8}$$

All unweighted graph constants are bounded as well. The reciprocals of N, C, D, α used below are bounded pure coefficient elements.

Two exact graphs retaining the large packets. For independent finite testing angles, case (I) uses

$$X_g = xu^2W, \quad r_g = r_0uW, \quad p_g = p_0uW, \quad Y_g = (1-r_0)uW, \quad Z_g = \{v(1-r_0) - p_0\}uW. \tag{SC9}$$

Its exact normals and inverse are

$$\begin{aligned} E_X = \sqrt{N}(X - xu^2W), \quad E_r = \sqrt{C}(r - r_0uW), \quad E_p = \sqrt{\alpha}(p - p_0uW),\\ X = xu^2W + E_X/\sqrt{N}, \quad Y = (1-r_0)uW - E_r/\sqrt{C}, \quad Z = vY - p_0uW - E_p/\sqrt{\alpha}. \end{aligned} \tag{SC10}$$

Case (II) uses

$$X_g = xu^2W, \quad J_g = j_0uW, \quad p_g = p_0uW, \quad Y_g = \frac{u(1 - j_0 + vp_0 + xu^2v)}{D_v}W, \quad Z_g = vY_g - p_0uW. \tag{SC11}$$

Here the exact normals and inverse are

$$\begin{aligned} E_X = \sqrt{N}(X - xu^2W), \quad E_J = \sqrt{D}(J - j_0uW), \quad E_p = \sqrt{\alpha}(p - p_0uW),\\ X = xu^2W + E_X/\sqrt{N}, \quad J = j_0uW + E_J/\sqrt{D}, \quad p = p_0uW + E_p/\sqrt{\alpha},\\ Y = (uW + uvX + vp - J)/D_v, \qquad Z = vY - p. \end{aligned} \tag{SC12}$$

Both graphs match the entire actual contact. All denominators in these formulas are pure coefficient factors or the pure testing unit D_v. No inverse of a mixed tensor element occurs. The possibly unbounded packets become respectively

$$\begin{aligned} &\text{(I)} \quad 4(n/N)D_uD_vE_X^2, quad4D_vE_r^2, quad4D_uE_p^2,\\ &\text{(II)} \quad 4(n/N)D_uD_vE_X^2, quad4E_J^2, quad4D_uE_p^2 \end{aligned}$$

in the quadratic normal remainder. Their graph and first-normal coefficients are bounded by (SC8), for example $nx, nx^2, Cr_0^2, \sqrt{C}r_0, \alpha p_0^2, \sqrt{\alpha}p_0$ in case (I), and $Dj_0^2, \sqrt{D}j_0$ in case (II). Every remaining source coefficient is bounded.

Entire scalar and entire first-normal division. Set $W = 1$, $K = k^2$, and $R = u^2 - (K/a)v$, $T_s = u(v - a)$. Since $R_2|_g = v + (p_0 - x)u^2$ for both graphs, the entire scalar has the exact shape

$$\Phi = F|_g = 4mv^2 + u^2 H_0(u^2, v), \tag{SC13}$$

with bounded polynomial coefficients in case (I), and bounded coefficients in $B_c[v, D_v^{-1}]$ in case (II). The three scalar contact jets give $H_0(K, a) = -\gamma K$, $H_{0,U}(K, a) = \gamma$, and $H_{0,v}(K, a) = -2\gamma K/a$. In particular the explicit expression

$$\Psi = H_0 - \gamma u^2 + 2\gamma(K/a)v$$

satisfies $\Phi - \gamma R^2 = u^2\Psi$, and its value and both first derivatives vanish at $(u^2, v) = (K, a)$. Finite Taylor division of Ψ therefore gives

$$\Phi = \gamma R^2 + u^2\{(u^2 - K)^2 H_{20} + (u^2 - K)(v - a)H_{11} + (v - a)^2 H_{02}\}, \tag{SC14}$$

with bounded coquotients. This is an exact bounded Gram on (R, T_s), with separate residue pivot $\bar{\gamma}$ on R. For the rational case, Taylor division is legitimate because

$$D_v^{-r} - D_a^{-r} = -(v - a)(v + a)\frac{\sum_{i=0}^{r-1} D_v^i D_a^{r-1-i}}{D_a^r D_v^r}.$$

Iterating this identity gives the second-order division; only pure factors have been inverted.

Here are the full first normals, rather than initial forms. In case (I) they are

$$\mathcal{N}_X = F_X|_g, \quad \mathcal{N}_r = -(F_Y + vF_Z)|_g, \quad \mathcal{N}_p = -F_Z|_g, \tag{SC15}$$

and in case (II) they are

$$\mathcal{N}_X = \{F_X + (uv/D_v)(F_Y + vF_Z)\}|_g, \quad \mathcal{N}_J = -(F_Y + vF_Z)|_g/D_v, \quad \mathcal{N}_p = (vF_Y - F_Z)|_g/D_v. \tag{SC16}$$

The derivatives in (SC7) specify every term. In particular the following exact packet expressions expose the cancellations:

$$\begin{aligned} -(F_Y + vF_Z)/(4D_v) &= (d + 2D)J + 2Cr - 2AD_uY - duvB + dvD_uZ, \\ (vF_Y - F_Z)/(4D_v) &= (d + 2\alpha)D_up - 2Cvr + 2AvD_uY + 2muR_2 - duB + dD_uZ. \end{aligned} \tag{SC17}$$

The actual coefficients of the normal roots are obtained by dividing these rows by $\sqrt{N}, \sqrt{C}, \sqrt{\alpha}$ in case (I), and by $\sqrt{N}, \sqrt{D}, \sqrt{\alpha}$ in case (II). Using (SC8), their exact forms after those divisions are

$$\mathcal{N}_X/\sqrt{N} = -8mv/\sqrt{N} + u^2 L_X(u^2, v), \qquad \mathcal{N}_i/\sqrt{w_i} = uL_i(u^2, v), \tag{SC18}$$

where all L_i have bounded coefficients in the indicated ring. The contact equations give $L_X(K, a) = 8ma/(K\sqrt{N})$ and $L_i(K, a) = 0$. Writing $L_X = L_X(K, a) + (u^2 - K)L_{X1} + (v - a)L_{X2}$, and similarly for L_i, gives the full exact row identities

$$\begin{aligned} \mathcal{N}_X/\sqrt{N} &= (8ma/(K\sqrt{N}) + u^2 L_{X1})R + u\{(K/a)L_{X1} + L_{X2}\}T_s, \\ \mathcal{N}_i/\sqrt{w_i} &= uL_{i1}R + \{(K/a)L_{i1} + L_{i2}\}T_s. \end{aligned} \tag{SC19}$$

Thus the X row has infinitesimal couplings to both soft roots, and the other rows have infinitesimal couplings to R. The exact state degree is two, so the bounded quadratic remainder already accounts for every higher-normal term. Equations (SC10)–(SC19) give an exact full-parent bounded five-root Gram.

A common exact real Schur certificate. The X and R residue pivots are $2(1+\bar{a}^2)$ and $\bar{\gamma}$. The unscaled boundary hard form on Y, Z is

$$\mathsf{B}(v)[Y,Z] = sD_vY^2 + 4D(Y+vZ)^2 + 4\alpha(vY-Z)^2, \qquad s = 4(C+A)+2d. \tag{SC20}$$

In case (I), $Y = -E_r/\sqrt{C}$, $Z = -vE_r/\sqrt{C} - E_p/\sqrt{\alpha}$ at $u=0$. The form has the explicit pivots $4D_vE_r^2 + 4E_p^2$ plus a bounded nonnegative form. In case (II), $Y = (vE_p/\sqrt{\alpha} - E_J/\sqrt{D})/D_v$ and $Z = (-E_p/\sqrt{\alpha} - vE_J/\sqrt{D})/D_v$. The explicit pivots are $4E_J^2 + 4E_p^2$, again with a bounded nonnegative remainder. The hard residue is strict in both cases.

The determinant and the exact boundary Schur numerator are

$$\begin{aligned} \det \mathsf{B}(v) &= 4(1+v^2)\{s(D\,v^2+\alpha) + 4D\alpha(1+v^2)\}, \\ Q_{10} &= 4\alpha A[d+2(C+D)], \qquad Q_{12} = 4CD[d+2(\alpha+A)], \\ Q_0 &= 8ACD + 8AC\alpha - 4ADd - 2Ad^2 - 2Cd^2 - 4Cd\alpha - 2Dd^2 - 4Dd\alpha - d^3 - 2d^2\alpha. \end{aligned} \tag{SC21}$$

Put $S = \alpha T$. Every coefficient of $\det \mathsf{B}/S$ and of $Q_{10}/S, Q_0/S, Q_{12}/S$ is bounded. This follows directly from (SC21), bounded A, bounded $C/T, D/T, \delta$, and bounded $1/\alpha$. Moreover

$$Q_{10}/S = 4A \asymp 1, \qquad Q_{12}/S = 4\delta[d+2(\alpha+A)]/\alpha \asymp 1. \tag{SC22}$$

The normalized determinant has a positive lower bound for every bounded unit real v. Indeed it equals C/T times the determinant of the scaled hard form in case (I), and equals $(D/T)D_v^2$ times that determinant in case (II). The factors C/T and D/T, respectively, are positive units in their cases.

We now take residues of these bounded coefficients and use an ordinary real variable v. The exact real hard Schur function is

$$\bar{f}(v) = \frac{8(1+v^2)}{\overline{\det \mathsf{B}/S}(v)} \left\{ \overline{Q_{10}/S} + \overline{Q_0/S}v^2 + \overline{Q_{12}/S}v^4 \right\}. \tag{SC23}$$

This formula is an ordinary real Schur calculation, never an inverse of $\det \mathsf{B}(v)$ in $K_c \otimes K_t$. The contact first-normal identities identify the graph at $\bar{a}$ with the unique minimizer of the scaled real hard form. The scalar contact jets give $\bar{f}(\bar{a}) = \bar{f}'(\bar{a}) = 0$. Because $\bar{a} \neq 0$, the quartic bracket in (SC23) is exactly

$$\overline{Q_{12}/S}(v^2 - \bar{a}^2)^2.$$

Its strict second derivative supplies the soft Schur pivot

$$\tfrac{1}{2}\bar{f}''(\bar{a}) = \frac{32(1+\bar{a}^2)\bar{a}^2\overline{Q_{12}/S}}{\overline{\det \mathsf{B}/S}(\bar{a})} > 0.$$

The entire five-root Gram therefore has strictly positive residue when u is infinitesimal or zero and $\bar{v} = \bar{a}$. The strict-residue Gram lemma proves tensor SOS. The exact identity

$$F(u,-v;W,-X,Y,-Z) = F(u,v;W,X,Y,Z)$$

supplies $\bar{v} = -\bar{a}$.

All other independent tests, including literal endpoints. If u is infinitesimal and v is a bounded unit with $\bar{v} \neq \pm\bar{a}$, retain the full original m, n packets. In case (I) use the roots

$$X, \quad K_C = \sqrt{C}r, \quad K_\alpha = \sqrt{\alpha}p, \quad uW$$

and the exact inverse $Y = uW - K_C/\sqrt{C}$, $Z = vY - K_\alpha/\sqrt{\alpha}$. In case (II) use

$$X, \quad K_D = \sqrt{D}J, \quad K_\alpha = \sqrt{\alpha}p, \quad uW$$

and the inverse in (SC12), with unshifted J, p. The entire remaining parent is a bounded quadratic Gram on these four roots. Its separate X residue pivot is $2\bar{d}(1+\bar{v}^2)$; its hard pair is the strict form (SC20) in the scaled coordinates just displayed, and its soft Schur complement is $\bar{f}(\bar{v}) > 0$ by (SC23).

For the two endpoint neighborhoods $u, v \sim 0$ and $u, q = 1/v \sim 0$, divide by d, choose

$$\mu = \min(1, C/d, A/d, D/d, \alpha/d) > 0,$$

a bounded unit by (SC6), and retain all excesses together with the full m, n packets. The selected bounded parent has signed weight one, $C = A = D = \alpha = \mu$, and $m = n = 0$. In the finite chart it is exactly a Gram on (X, Y, Z, uW). At $(u, v) = (0, 0)$ its hard block is positive and its soft Schur complement is $8Q_{10,\mu}/\det \mathsf{B}_\mu(0) > 0$. For the reciprocal chart use the polynomial homogeneous parent $q^2 F_\mu(u, 1/q)$, on the same four roots. At $(u, q) = (0, 0)$ its hard block is $\mathrm{diag}(2, 4(C + A + \alpha) + 2, 4D)$ in the (X, Y, Z) variables, evaluated at the capped weights, and its soft Schur complement is $8Q_{12,\mu}/\det \mathsf{B}_{\mu,\infty} > 0$. These are bounded polynomial Grams, so they include the full infinitesimal neighborhoods and literal $v = 0$ or $q = 0$.

For completeness, the other open commuting axes use the same cap; here the other angle has a nonzero finite residue. At $v = 0$ the core is $2\{(uW - Y)^2 + (X + uZ)^2\}$ and the A, α packets force $Y = Z = 0$; finite nonzero u then forces $W = X = 0$. At $u = \infty$ the homogeneous core is $2\{(W + vX)^2 + (vY - Z)^2\}$; the C, A packets force $W = Y = 0$, and finite nonzero v forces $X = Z = 0$. At $v = \infty$ the core is $2(1 + u^2)(X^2 + Y^2)$; the C, D packets force $W = Z = 0$ for finite nonzero u. Thus these complete state matrices are strict.

Only the two corners with $u = \infty$ remain. Put $p_u = 1/u$. At $(p_u, v) = (0, 0)$ the selected state form is $(2 + 8\mu)W^2 + 4\mu Y^2 + (2 + 4\mu)Z^2$. Its sole zero ray is X. In the chart $X = 1$, the five small variables (p_u, v, W, Y, Z) have exact quadratic part

$$(2 + 4\mu)(W + v)^2 + 2(Z + p_u)^2 + 4\mu(W^2 + Y^2 + Z^2) \succ 0.$$

At $(p_u, q) = (0, 0)$ the selected state form is $4\mu W^2 + (2 + 4\mu)X^2 + (2 + 8\mu)Y^2$. In its sole zero-ray chart $Z = 1$, the variables (p_u, q, W, X, Y) have exact quadratic part

$$2X^2 + (2 + 4\mu)(Y - q)^2 + 4\mu\{W^2 + Y^2 + (X - p_u)^2\} \succ 0.$$

There are no lower-order terms. Allocating every higher monomial to two coordinate roots leaves an infinitesimal factor, giving a strict-residue Gram for the entire selected polynomial at each corner. Retain the original excess packets throughout. After pure testing projective normalization, states outside each displayed zero ray have positive residue value and are covered by the scalar strict-residue lemma.

Noncommuting-residue tests use the inherited arbitrary-physical-source theorem. This exhausts all independent projective tests. Finally homogenize the local graph identities in W, replacing R, T_s by $RW, T_s W$; they then include $W = 0$. Restore all pure positive coefficient and testing factors, and split complex states into real and imaginary parts. No source infinitesimal was compared with an independent testing infinitesimal anywhere in the argument.

Exact scope. This completes the strong C2 finite pitchfork for $C/d, \alpha/d$ bounded below, with arbitrary upper scales of all six events. The reflected actual-contact domain follows by the established C2 reflection. It does not cover other actual-angle residue types or sources in which either complementary ratio has no positive real lower bound; no exhaustive global C2 join is asserted here.

Chapter 94

A homogeneous scalar lift at the unbounded-state contact

Component `pbb29c73cc124`
Source: `R13/baseline/baseline/baseline/baseline/advances/c2_unbounded/homogeneous_scalar_lemma.tex`
SHA256: `bb29c73cc124d2a0c215bb3baf3e7812d921f83f2684457880bf93b0f1953f2e`
Inventory use: HL1–HL11 complete scalar ideal lift at arbitrary actual state ratio

94.1 A homogeneous scalar lift at the unbounded-state contact

This lemma is a scalar lifting statement. It neither supplies the full state normals nor proves positivity of its coefficient Gram by itself. Its purpose is to avoid any bound on the contact state ratio.

Let $B, \sigma > 0$ be pure coefficient infinitesimals, let $k > 0$ be a pure coefficient bounded unit, and put $x = \sigma B$, $\xi = u^2$. Suppose

$$f = a(\xi, v)W^2 + 2ud(\xi, v)WZ + c(\xi, v)Z^2, \quad a = \frac{\xi}{B}Q + \sigma^2 a_0, \qquad c = Be + \xi b, \tag{HL1}$$

where Q, a_0, d, e, b have bounded polynomial coefficients. Bounded pure testing unit denominators may be cleared beforehand. Assume the value and all first derivatives of f vanish at

$$u = \sqrt{x}, \quad v = k, \quad W = \frac{1}{k}\sqrt{B/\sigma}, \quad Z = 1. \tag{HL2}$$

No bound on B/σ is assumed.

Define the homogeneous roots

$$R = \frac{uW - (B/k)Z}{\sqrt{B}}, \qquad S = uZ - \sigma kW, \qquad U = \sqrt{B}(v - k)Z, \qquad T = u(v - k)Z. \tag{HL3}$$

They satisfy the exact conductors

$$(\xi - x)W = u\sqrt{B}R + (B/k)S, \qquad (\xi - x)Z = uS + \sigma k\sqrt{B}R. \tag{HL4}$$

Then f has an exact symmetric Gram on (R, S, U, T) whose entries are bounded polynomials. In particular no inverse of σ, B, a testing infinitesimal, or a mixed expression is required in these Gram entries. The roots themselves need not be bounded.

First set $v = k$. In this paragraph every scalar coefficient is evaluated at $v = k$. State stationarity and u stationarity give

$$a_c = -\sigma k d_c, \qquad c_c = -Bd_c/k,$$

$$B(a_\xi)_c + kd_c + 2kx(d_\xi)_c + \sigma k^2(c_\xi)_c = 0, \tag{HL5}$$

where c on a subscript means evaluation at $\xi = x$. These are identities in the pure coefficient field; deriving them does not insert an inverse into a tensor Gram.

Set, by monic polynomial division in ξ,

$$s = \frac{c + Bd/k}{\xi - x}, \qquad q = -kd - \sigma k^2 s, \qquad h_2 = \frac{Ba + k\xi d + \sigma k^2 c + \sigma Bkd}{(\xi - x)^2}. \tag{HL6}$$

All three quotients have bounded polynomial coefficients. The first numerator vanishes at x by (HL5), and the last numerator and its ξ derivative vanish there by the same three identities. Moreover $Ba = \xi Q + B\sigma^2 a_0$ is bounded. Division by a monic polynomial preserves bounded coefficients and is not evaluation of an inverse of $\xi - x$.

Direct multiplication gives the exact identity

$$f(\xi, k) = qR^2 + sS^2 + h_2 \left(uR + \frac{\sqrt{B}}{k} S \right)^2. \tag{HL7}$$

For later residue computations, if $E_0 = e + d/k$, then

$$s = b + B\frac{E_0(\xi) - E_0(x)}{\xi - x} + x\frac{b(\xi) - b(x)}{\xi - x}. \tag{HL8}$$

Thus $s_0 = b_0$, $q_0 = -k_0 d_0$, and the h_2 correction has zero residue in every Gram entry. Here subscripts 0 denote residues, not evaluation at the contact polynomial variable.

The complete angular coquotient. Write $\tau = v - k$ and, for each of Q, a_0, d, e, b, form its bounded monic difference in v, denoted respectively by $Q_1, a_{01}, d_1, e_1, b_1$:

$$Q(\xi, v) - Q(\xi, k) = \tau Q_1(\xi, v),$$

and similarly for the other four functions. Put

$$H_1 = Q_1/k^2 + 2d_1/k + e_1, \qquad L_1 = a_{01}/k^2 + b_1, \qquad J_1 = H_1 + \sigma L_1.$$

Angular stationarity at (HL2) gives exactly $J_1(x, k) = 0$. Consequently the successive monic differences

$$J_1 = (\xi - x)J_\xi + \tau J_v, \quad J_\xi = \frac{J_1(\xi, v) - J_1(x, v)}{\xi - x}, \qquad J_v = \frac{J_1(x, v)}{v - k} \tag{HL9}$$

are bounded polynomials. Let $D_1 = L_1 + BJ_\xi$. The full exact angular correction is

$$\begin{aligned} f(\xi, v) - f(\xi, k) = \tau Q_1 R^2 + \frac{\tau a_{01}}{k^2} S^2 \\ + \{2(Q_1/k + d_1) + \sigma k D_1\} RU \\ + \left\{ D_1 - 2a_{01}/k^2 \right\} ST + J_v U^2. \end{aligned} \tag{HL10}$$

To check this without any division, use $uW/\sqrt{B} = R + \sqrt{B}Z/k$ and $\sigma W = (uZ - S)/k$. The only term not already among the displayed root products becomes

$$\tau Z^2 \{BH_1 + \xi L_1\} = D_1 \tau(\xi - x)Z^2 + BJ_v \tau^2 Z^2 = D_1(ST + \sigma kRU) + J_v U^2,$$

by (HL4). Equations (HL7) and (HL10) prove the asserted homogeneous Gram lift for every state W, Z simultaneously. There is no state normalization and no discarded endpoint $Z = 0$.

The usual exact zero, now with the new source scale, is

$$T^2 - \sigma U^2 - ST(v - k) - \sigma kRU(v - k) = 0. \tag{HL11}$$

This is the appropriate identity for redistributing the angular diagonal. Its U^2 coefficient is σ, which is infinitesimal; replacing it by 1 would erase the smaller radial scale. Any positivity argument must use the actual residue coefficients of (HL7)–(HL10), rather than importing the bounded-state Schur test without this scale correction.

Strict scalar corollary for the small-B source. Suppose additionally that the coefficientwise leading functions on the finite $u = 0$ axis are

$$Q_0(v) = \frac{2K^2(1+v^2)}{1+K^2}, \qquad d_0(v) = -2v, \qquad e_0(v) = \frac{2(v^2/K^2+K^2)}{1+K^2}, \qquad b_0(K) > 0,$$

where $K = k_0$. Then the (R, U) residue block produced above is

$$2K^2R^2 + \frac{4(K^2-1)}{1+K^2}RU + \frac{2}{K^2}U^2,$$

with Schur margin $8/(1+K^2)^2 > 0$. The S^2 coefficient has residue $b_0(K) > 0$, the RS residue is zero, and the only other possible residue entry is a bounded ST coefficient, say ℓ. Add a positive real multiple δ of (HL11), with $\delta > \ell^2/(4b_0(K))$. Its correction to the U^2 entry has zero residue because σ is infinitesimal, whereas it supplies the positive T^2 diagonal δ. Both residue blocks are then strictly positive. This proves the scalar strict bounded Gram for all homogeneous states in the matched-angle chart. Full normals and the other testing charts remain separate requirements.

Chapter 95

The unbounded-b source with xb infinitesimal

Component `pab6222b899ef`
Source: `R13/baseline/baseline/baseline/baseline/advances/c2_unbounded/unbounded_b_small_B.tex`
SHA256: `ab6222b899ef922dd4814c71231c2778c17c53fdee166bae7e3618224b5641ec`
Inventory use: UB1–UB17 actual full homogeneous scalar, both full normals, hard block, literal n=0, every state ratio and every testing chart

95.1 The unbounded-b source with xb infinitesimal

Theorem 95.1 (The remaining unbounded-state rational C2 source)**.** *Use the exact original rational C2 source. Suppose $x > 0$ is infinitesimal, k is a positive bounded unit, b is positive unbounded, $B = xb$ is infinitesimal, $0 < r < 1$, $0 < h < b$, and all original positive-chart inequalities hold. Assume $n \geq 0$ and physicality of the source. Then the full original parent has tensor SOS for every independent real or complex projective state and every independent projective pair of measurement angles.*

The conclusion includes $n = 0$, arbitrarily large n, arbitrary complementary openings, and every value of the pure coefficient ratio B/σ, where $\sigma = 1/b$. In particular, the actual contact ratio $W/Z = k^{-1}\sqrt{B/\sigma}$ need not be bounded. This proof uses the homogeneous scalar lemma in Section 94.1; it restores both full state normals and exhausts the other testing charts.

Exact source scales. Put

$$\sigma = 1/b, \quad x = \sigma B, \quad \eta = h/b, \quad y = B\eta, \quad z = B(1-\eta), \quad A_c = 1 + \sigma B r, \quad S_c = A_c + z.$$

The positive chart gives $0 < \eta < 1$; both y, z are infinitesimal. Let $\widehat{\Delta} = \sigma\Delta$, $\widehat{N} = \sigma^2 B N_{\mathrm{src}}$, $\widehat{g} = \sigma g = 1 + \sigma(r - 1 + B\eta)$ and $j = A_c - z/k^2$. The cancelled exact formulas are

$$\begin{aligned}
nr &= \frac{(1+\sigma r)A_c S_c - r\widehat{\Delta}}{2\widehat{\Delta}}, \\
m &= \frac{\sigma^2(1+k^2)(1-r)A_c S_c}{2\widehat{g}\widehat{\Delta}}, \\
C &= \frac{A_c \widehat{N}}{2z\widehat{\Delta}}, \qquad A = \frac{k^2 A_c(1+\sigma r)S_c}{2y\widehat{\Delta}}, \\
D &= \frac{B(1+\sigma B)(1+\sigma r)(1-\sigma k^2 r)S_c}{2k^2 j\widehat{\Delta}},
\end{aligned}$$

$$\alpha = \frac{B(1+\sigma r)\widehat{N}}{2(1-y)\widehat{\Delta}}. \tag{UB1}$$

Here $\widehat{\Delta}, \widehat{N}, \widehat{g}, j, 1-y$ are bounded positive units. Write $K = k_0 > 0$ and $D_K = 1 + K^2$. Their relevant residues imply

$$\begin{gathered}(nr)_0 = \tfrac{1}{2}(D_K^{-1} - r_0), \quad (m/\sigma^2)_0 = (1-r_0)/2, \\ (Cz)_0 = (Ay)_0 = K^2/(2D_K), \quad (D/B)_0 = 1/(2K^2 D_K), \quad (\alpha/B)_0 = K^2/(2D_K). \end{gathered} \tag{UB2}$$

Since $n \geq 0$, $r_0 \leq D_K^{-1} < 1$, so $1 - r$ is a unit. Set $N = 4n + 2 \geq 2$. Exactly,

$$Nr = \frac{2(1+\sigma r)A_c S_c}{\widehat{\Delta}}, \tag{UB3}$$

which is a bounded positive unit. Thus nr, nr^2 are bounded even when n is unbounded, while $1/N$ and $1/\sqrt{N}$ are bounded.

All opening cancellations needed below are uniform:

$$BC(1-\eta) = Cz, \quad BA\eta = Ay, \quad BC(1-\eta)^2 = Cz(1-\eta), \quad BA\eta^2 = Ay\eta. \tag{UB4}$$

The quantity $H_B = B(C + A)$ is bounded below by a positive real; its reciprocal and reciprocal square root are bounded. No assumption that η or $1 - \eta$ is a unit is made.

The whole finite $u = 0$ axis. Use the original parent and the exact-contact graph

$$X_g = -ruZ, \qquad Y_g = \eta u W, \qquad E_X = \sqrt{N}(X + ruZ), \quad E_Y = \sqrt{C + A}(Y - \eta u W). \tag{UB5}$$

The actual contact is

$$u = \sqrt{\sigma B}, \quad v = k, \quad W = k^{-1}\sqrt{B/\sigma}, \quad Z = 1.$$

Let $\xi = u^2$ and $f = F|_g$. Exact polynomial coquotients give

$$f = a(\xi, v)W^2 + 2ud(\xi, v)WZ + c(\xi, v)Z^2, \quad a = \frac{\xi}{B}Q + \sigma^2 a_0, \quad c = Be + \xi b. \tag{UB6}$$

Every coefficient of Q, a_0, d, e, b is bounded by (UB3)–(UB4). For example $a_0 = 4(m/\sigma^2)v^2$, $e = 4(D/B)v^2 + 4\alpha/B$, and $Q = B(a - a|_{u=0})/u^2$. These are monic polynomial divisions. The full leading functions at $u = 0$ are

$$\begin{aligned} q(v) &= 2K^2(1+v^2)/D_K, & d_0(v) &= -2v, \\ e_0(v) &= 2(v^2/K^2 + K^2)/D_K, & (a_0)_0(v) &= 2(1-r_0)v^2, \\ b_0(v) &= 2r_0(1+v^2)/D_K - 4r_0 + 2. \end{aligned} \tag{UB7}$$

For remote tests use

$$P = uW/\sqrt{B}, \quad B_u = uZ, \quad V = \sigma v W, \quad E = \sqrt{B}Z.$$

Then (UB6) has the exact bounded Gram

$$QP^2 + bB_u^2 + (a_0/v^2)V^2 + eE^2 + 2dPE, \tag{UB8}$$

where a_0/v^2 is a polynomial coquotient. The (P, E) residue block is strictly positive whenever $v_0^2 \neq K^2$, since

$$q(v)e_0(v) - 4v^2 = 4(v^2 - K^2)^2/D_K^2. \tag{UB9}$$

This is a global identity for every finite residue v, not a local-contact positivity inference.

Both complete state normals. The exact quadratic normal expansion is

$$F = f + E_X L_X + E_Y L_Y + H_{XX} E_X^2 + 2H_{XY} E_X E_Y + H_{YY} E_Y^2. \tag{UB10}$$

Its entire hard block is bounded with residue $\mathrm{diag}(D_{v,0}, 4D_{v,0})$. Indeed the X pivot is $ND_uD_v + O(1)$ before division by N, with the remaining terms either infinitesimal in u or of order m; the Y pivot from the strong packets is exactly $4D_v(1 + [A/(C + A)]u^2)$. All mixed entries have zero residue and contain no C, A contribution.

The complete unnormalized X derivative has the form

$$F_X|_g = uZA_X + W(\xi B_1 + \sigma^2 B_0), \tag{UB11}$$

with bounded polynomial coefficients, where

$$\begin{aligned} B_0 &= -8(m/\sigma^2)v(1 + \eta\xi), \\ B_1 &= v(8D + 4)(1 - \eta), \\ A_X &= -8nrD_uD_v + 8m(1 - r) - (8D + 4)v^2(1 + r\xi) + 4D_v(1 - r). \end{aligned}$$

Thus in the remote roots its P, V coefficients have zero residue; its only possible nonzero residue coupling is to B_u. The exact residue Schur expression for the (B_u, E_X) block is

$$(Nr^2 D_v - 4r + 2) - \frac{(4 - 2NrD_v)^2}{4ND_v} = 2 - \frac{4}{ND_v}. \tag{UB12}$$

It is strictly positive for every finite nonzero residue v, because $N \geq 2$. At $v_0 = 0$ it is strict if n is unbounded or has positive residue. The sole zero case $n_0 = v_0 = 0$ is treated below.

For the other complete first normal,

$$L_Y = \frac{1}{\sqrt{H_B}}\{P(B_2 + Be_w) + B_u(\sqrt{B}\, ue_z) + Ee_*\}, \tag{UB13}$$

where e_w, e_z are bounded,

$$B_2 = 8D_v\{(BA)\eta D_u - (BC)(1 - \eta)\}, \qquad e_* = 8v(D - \alpha).$$

All coefficients have zero residue, by (UB4), $B \to 0$, and $u \to 0$. This verifies both full normals and proves the full parent at every remote finite-axis test covered by (UB12).

Matched residues without any state-ratio restriction. For $v_0 = K$, apply the homogeneous scalar lemma to (UB6). Its contact and full stationarity hypotheses hold exactly because the graph (UB5) passes through the original stationary contact. Use the homogeneous roots

$$R = \frac{uW - (B/k)Z}{\sqrt{B}}, \quad S = uZ - \sigma kW, \quad U = \sqrt{B}(v - k)Z, \quad T = u(v - k)Z. \tag{UB14}$$

They satisfy

$$(\xi - x)W = u\sqrt{B}R + (B/k)S, \qquad (\xi - x)Z = uS + \sigma k\sqrt{B}R. \tag{UB15}$$

The lemma constructs an exact bounded scalar Gram, valid for every W, Z simultaneously. Its (R, U) residue is

$$2K^2R^2 + \frac{4(K^2 - 1)}{D_K}RU + \frac{2}{K^2}U^2,$$

whose Schur margin is $8/D_K^2 > 0$. Its radial S^2 residue is $s_0 = 2(1 - r_0) > 0$; its only other possible residue coupling is ℓST, with a bounded real residue ℓ.

To lift the full X normal, set

$$L = A_X + (B/k)B_1 + (\sigma/k)B_0.$$

Equations (UB11) and (UB15) give exactly

$$F_X|_g = u\sqrt{B}B_1R - (\sigma/k)B_0S + uZL.$$

At the source contact $L(x,k) = 0$, because $F_X = 0$ there. Write its bounded monic differences as $L = (\xi - x)L_\xi + (v - k)L_v$. Then

$$F_X|_g = u\sqrt{B}(B_1 + \sigma k L_\xi)R + \{\xi L_\xi - (\sigma/k)B_0\}S + L_v T. \tag{UB16}$$

After division by $\sqrt{N}$, all coefficients are bounded. The R, S coefficients have zero residue; the T coefficient may have a finite residue γ (it is $-8K/(D_K\sqrt{N_0})$ when N is bounded). This nonzero full-normal coupling is retained, not silently discarded.

For (UB13), put $A_Y = B_2 + Be_w$ and $L'_Y = A_Y/k + \xi e_z + e_*$. Its numerator is exactly $RA_Y + \sqrt{B}ZL'_Y$. Exact normal contact gives $L'_Y(x,k) = 0$. Monic differences and (UB15) yield bounded coefficients on R, S, U, all of zero residue: the R, S corrections carry factors σB or $u\sqrt{B}$, and the U coefficient is the angular coquotient of an identically zero leading function. Division by $\sqrt{H_B}$ preserves these bounds.

Finally add δ times the exact zero

$$T^2 - \sigma U^2 - ST(v - k) - \sigma k RU(v - k) = 0. \tag{UB17}$$

Choose a real $\delta > \ell^2/(4s_0) + \gamma^2/(4D_K)$. This supplies a T^2 residue δ at zero residue cost to the U^2 coefficient, since σ is infinitesimal. The block on (S, E_X, T) has strictly positive Schur complement; the (R, U) block and E_Y diagonal are already strict. Thus the complete six-root Gram on (R, S, U, T, E_X, E_Y) is strictly positive in residue. The calculation is homogeneous in all four state coordinates and needs no bound on W/Z, no comparison of B with σ, and no state endpoint exclusion. The exact simultaneous sign copy treats $v_0 = -K$.

The double-zero test, including literal $n = 0$. Suppose now $n_0 = v_0 = 0$. Use the pure coefficient quantities

$$H_c = 4n + 4m + 2, \quad \zeta = (4m + 2)/H_c, \quad \xi_c = 4m/H_c, \quad \eta_c = C/(C + A),$$

and graph $X_g = \xi_c vW - \zeta uZ$, $Y_g = \eta_c uW$. Normalize its X normal by $\sqrt{H_c}$ and its Y normal by $\sqrt{C + A}$. Put

$$\epsilon = 4\alpha + 4n(4m + 2)/H_c > 0.$$

Then ϵ is infinitesimal, n/ϵ and B/ϵ are bounded, and the harmonic product $BCA/(C + A)$ is a unit. Use the five scalar roots

$$P = uW/\sqrt{B}, \quad V = \sigma vW, \quad E = \sqrt{B}Z, \quad B_\epsilon = \sqrt{\epsilon}\, uZ, \quad T_0 = uvZ.$$

The exact inherited identity, with its original coefficients, is

$$f_{zz} = 4\alpha + \epsilon u^2 + 4n\zeta^2 u^4 + 4Dv^2 + u^2 v^2 B_b, \qquad (B_b)_0 = 2.$$

The W^2 coefficient at $u = 0$ is exactly

$$\frac{4m(2n + 1)}{2n + 2m + 1} v^2 + (4n + 2)\xi_c^2 v^4.$$

Therefore its coefficient on V^2 has positive residue $4(m/\sigma^2)_0 = 2(1 - r_0)$. The scalar residue on the five roots is strictly diagonal, with entries

$$\left(\frac{2K^2}{D_K},\ 2(1 - r_0),\ \frac{2K^2}{D_K},\ 1,\ 2 \right).$$

The cross term is allocated using $uvWZ = vPE$, so it has zero residue.

Both full normals are restored, including when $n = 0$ exactly. At $v = 0$ the X derivative is $-8n\zeta u^3 Z$, whose coefficient on B_ϵ is infinitesimal since n/ϵ is bounded. Its remaining Z coefficient is uv^2 times a bounded function, so it has zero residue on T_0. Its W coefficient is a sum of $u^2 vW$ times a bounded function and mvW times a bounded function divisible by v^2 at $u = 0$. Allocate the former to P with coefficient $u\sqrt{B}\, v$, and the latter to V with coefficient m/σ; both have zero residue. In particular one must not divide the former by σ. The Y derivative is the inherited harmonic-opening identity: $B_2 = 8D_v(BA)\eta_c u^2$, with bounded residual coquotients. Its coefficients on P, T_0, E all have zero residue. The hard residue is $\mathrm{diag}(1, 4)$. This proves the full-parent corner without deleting a state coordinate.

All remaining projective tests. For the other commuting charts use the four original outer-core identities with ρ replaced by $\sqrt{B}$, and cap C, A at s/B, where $s = \min(1, BC, BA)$ is a positive bounded unit. Retain both original nonnegative excess packets and the entire original n packet. No positive lower bound on m is needed: the first two outer pivots are $4m_0 + 2 = 2$, while the other two charts retain the whole m packet and do not use it for strictness. The reciprocal-v chart uses $(D/B)_0 > 0$ exactly as before. Thus the finite unit u axis, both reciprocal axes, all double corners, and literal projective endpoints have strict bounded Grams. The inherited physical noncommuting-residue theorem covers all remaining testing pairs.

The finite-axis certificates above are homogeneous in the state, including the previously problematic unbounded actual state ratio. Splitting a complex state into its real and imaginary parts proves the complex assertion without an additional chart. All inverses and radicals used in coefficients are pure coefficient or pure testing operations; every displayed mixed quotient is explicit monic polynomial division. The roots may be unbounded. The strict bounded tensor Gram lemma and restoration of the retained packets complete the all-tests proof.

Chapter 96

Further actual-angle reductions from exact source entry

Component `pe56fb394b522`
Source: `R13/baseline/baseline/baseline/baseline/advances/source/angle_reductions.tex`
SHA256: `e56fb394b52263918d985b9437e107edea077b0ea105beebe65fca8b1c306250`
Inventory use: Actual directed weak II/IU/UI/UU exclusions and both-strong UU five-event entry

96.1 Further actual-angle reductions from exact source entry

All angles in this section are actual source contacts. They are strictly positive finite field elements; "infinite" below means unbounded in the coefficient field, not a literal projective endpoint. Write $z = c^2 = u_c^2$, $y = a^2 z = v_c^2$, and $A_c = 1 + zr$, $S = A_c + a^2 zt$. The exact source identities used are

$$n \geq 0 \iff (b+r)A_c S - r\Delta \geq 0, \qquad N = (1+z)(b+r)h + a^2\{g + j + rt(1+y)\} > 0.$$

Theorem 96.1 (The ordinary–ordinary source orbit is already closed)**.** *If both actual C2 half-angles are positive bounded units, then the source belongs either to the two-weak-complementary-event all-tests exit or to the five-consecutive-strong-event all-tests exit. No upper bound on an original event coefficient is required.*

Proof. Here a, z are units. The positive gaps give $0 < h < 1/z$ and $0 < t < 1 + zr$, so $h, t, b = h + a^2 t$ are bounded. From $g = b - 1 + r + zh > 0$ we obtain $1 < b + r + zh \leq (1+z)b + r$; thus $b + r$ is a positive unit. The positive expansion of Δ makes it a bounded unit too. The original A formula has unit numerator and bounded denominator h, so A is strong.

The exact numerator for n, after substituting $t = (b-h)/a^2$, satisfies the polynomial specialization

$$[(b+r)A_c S - r\Delta]_{r=1} = -z(a^2 + h)(b+1)(z+1).$$

If $1 - r$ were infinitesimal, taking residues would contradict $n \geq 0$: the right side is strictly negative. Hence $1 - r$ is a unit. The original m numerator is now a unit and $g\Delta$ is bounded, so m is strong. Apply the exact C2 reflection; it preserves ordinary actual angles and exchanges m, D. Thus D is strong.

If either C or α is weak, its original rational formula implies N infinitesimal: all the multiplying denominator factors are bounded and $z, a, b + r, \Delta$ have bounded inverses. The positive identity for N then forces h, j infinitesimal. Consequently $t = A_c - j$ and $1 - zh$ are units. Both original complementary formulas therefore give C, α infinitesimal. The two-weak theorem applies. Otherwise C, α are both strong; with m, A, D strong, the five-consecutive-event theorem applies. These cases exhaust the source. □

Some directed one-weak angle types are empty. Orient an exactly-one-weak source so that C is weak and α strong. In addition to the ordinary–ordinary exclusion above, its actual angle types cannot be $(\text{unit}, \text{Inf})$, $(\text{Inf}, \text{unit})$ or (Inf, Inf). Here Inf denotes positive infinitesimal.

For $(\text{unit}, \text{Inf})$, z is a unit and a^2 is infinitesimal. Again t, h, b, Δ are bounded, Δ is a unit, and $b+r$ is a unit. Weak C implies N/a^2 infinitesimal. The positive identity forces j and rt infinitesimal. Since $t = A_c - j$ is a unit, r is infinitesimal. Then $g > 0$ makes $b + zh$ a unit, and $b = h + a^2 t$ makes h a unit. The first positive summand of N is consequently a unit, contradicting N/a^2 infinitesimal.

For $(\text{Inf}, \text{unit})$, bounded b is a unit by the inherited source-entry inequality. Explicitly,

$$xC(b-h) = \frac{A_c(xN)}{2\Delta}, \qquad xN = (1+x)(b+r)(xh+k^2) - x(1+k^2)(1-r)(b-h).$$

The last numerator has residue $k_0^2(b_0 + r_0) > 0$, independently of any lower bound on g. Thus $xC(b-h)$ is a unit and C is strong. For unbounded b, (SR1) gives $C \asymp 1/L$ with bounded L, again strong. This exhausts b and excludes weak C in this angle orientation.

For (Inf, Inf), both z, y are infinitesimal and $t < A_c$ is bounded. Put $H = zh$, $L = z(b-h) = yt$ and $B = zb = H + L$. Here $0 < H < 1$, L is infinitesimal, and B is bounded. If H is a unit then b is unbounded; direct division of the original Δ by b gives a positive unit of residue $1 + B_0$. The positive first summand of N consequently gives $C \geq c_0 H/L$ for some positive real c_0, contradicting weak C. Thus weak C forces H, B infinitesimal. The exact ratio is then

$$\frac{\alpha}{C} = \frac{z^2(b+r)(b-h)}{A_c(1-H)} = \frac{(B+zr)L}{A_c(1-H)},$$

which is infinitesimal. Therefore α is weak as well, contrary to the selected one-weak orientation.

After these exclusions, an exactly-one-weak source oriented with weak C has only the following candidate actual types:

$$(\text{Inf}, \text{unbounded}), \quad (\text{unit}, \text{unbounded}), \quad (\text{unbounded}, \text{Inf}),$$
$$(\text{unbounded}, \text{unit}), \quad (\text{unbounded}, \text{unbounded}).$$

The second type is the reflection of the single-angle domain and is now fully covered by the homogeneous unbounded-b theorem and the other single-angle admissions. The four other listed directed types remain conditional source leaves; no nonemptiness is inferred from their listing.

Other strong-event angle types are not all empty. For $x > 0$ infinitesimal, the exact positive chart

$$c = \sqrt{x}, \quad a = 1, \quad b = 1/x, \quad r = 1/4, \quad t = 1/2$$

has $h = 1/x - 1/2$, $g = 1/x + 1/4 - x/2$, $j = 1/2 + x/4$ and all remaining chart gaps positive. Its original coefficients are

$$n = \frac{8-5x^2}{16(1+x)^2}, \qquad m = \frac{3x^2(3x+4)}{16(1+x)(4+x-2x^2)},$$
$$C = \alpha = \frac{4-x}{8x}, \qquad A = \frac{x(x+4)(3x+4)}{32(2-x)(1+x)^2},$$
$$D = \frac{(2-x)(3x+4)}{16x(1+x)}.$$

Every coefficient is positive. The unchanged exact contact is

$$(u_c, v_c; W, X, Y, Z) = (\sqrt{x}, \sqrt{x}; 1/x, -\sqrt{x}/4, 1/\sqrt{x} - \sqrt{x}/2, 1).$$

Original source stationarity, positive transverse state inertia and Laurent convexity prove physicality, including projective limits. Here C, α, D are unbounded, $m \sim 3x^2/16$, $A \sim x/4$, and $n \to 1/2$. Thus a genuine strong-complementary source can have two infinitesimal actual angles and weak m, A. This is an obstruction to deleting that angle orbit, not a counterexample to tensor SOS.

Chapter 97

The exact unbounded-state-scale C2 complement

Component `pb60046a37191`
Source: `R13/baseline/baseline/baseline/baseline/advances/source/unbounded_b_reduction.tex`
SHA256: `b60046a37191c59da6ebca1e87e245f5c8df75d7c802327c1cb821c8629f81a3`
Inventory use: SR1 exact opening-cancelled source scales and exhaustive B=xb weak/unit split

97.1 The exact unbounded-state-scale C2 complement

Use the original positive rational source with $d = 1$, actual $u_c = \sqrt{x}$ positive infinitesimal and $v_c = k$ a positive bounded unit. The global entry theorem supplies its positive chart inequalities. The preceding rational theorems, including the joint-opening theorem, now cover every bounded b: the inherited $n \geq 0$ entry inequality makes bounded b a unit. We classify the remaining b unbounded without changing any original coefficient.

Put

$$B = xb, \qquad H = xh, \qquad L = x(b-h) = B - H, \qquad a^2 = k^2/x, \quad t = L/k^2.$$

The original gaps give

$$0 < H < 1, \qquad 0 < L < k^2(1 + xr), \qquad B = H + L.$$

Thus B, H, L are bounded and $x/B = 1/b$ is infinitesimal. Set $A_c = 1 + xr$ and $S = A_c + L$, both positive bounded units. Also

$$g/b = 1 + (r-1)/b + H/b \longrightarrow 1, \qquad j = 1 + xr - L/k^2 > 0, \quad j \leq 1 + x.$$

The following exact rearrangements of the original short polynomials control every event scale:

$$\begin{aligned} x^2 N &= (1+x)(B+xr)(H+k^2) - x(1+k^2)(1-r)L, \\ \Delta/b &= (1+x)B + 2x(1+x)r + (1+x)(1+k^2) \\ &\quad + (x^2 - k^2 x)r^2/b + 2x(1+k^2)r/b + (1+k^2)/b. \end{aligned}$$

Consequently $xN/b = x^2 N/B$ and Δ/b are positive bounded units, with respective residues

$$H_0 + k_0^2, \qquad B_0 + 1 + k_0^2.$$

Here $H, L \leq B$ controls the divided corrections, and $x/B \to 0$.

The exact physical condition $n \geq 0$ is

$$r(\Delta/b) \leq (1 + r/b)A_c S.$$

It implies

$$r_0 \le \frac{1+L_0}{1+k_0^2+B_0}, \qquad 1-r_0 \ge \frac{k_0^2+H_0}{1+k_0^2+B_0} > 0.$$

Thus $1-r$ is a positive bounded unit without any lower bound on r. The original formulas now give the exact comparabilities

$$m \asymp b^{-2}, \qquad C \asymp L^{-1}, \qquad A \asymp H^{-1}, \qquad D \asymp B/j, \qquad \alpha \asymp B/(1-H). \tag{SR1}$$

Every omitted factor in $\asymp$ is a positive bounded unit with bounded reciprocal. For instance,

$$CL = \frac{A_c(xN/b)}{2\Delta/b}, \qquad AH = \frac{k^2 A_c(1+r/b)S}{2\Delta/b}$$

are positive units; the other identities follow using $g/b \to 1$ and $(b-k^2r)/b \to 1$.

There are exactly two unbounded-b scale leaves:

1. If $B = xb$ is infinitesimal, so are H, L, while $j \to 1$ and $1-H \to 1$. Thus C, A are unbounded and m, D, α are weak, with $m \asymp b^{-2}$ and $D \asymp \alpha \asymp xb$. This is the one-weak-complementary-event frontier, with weak α and strong C in this actual-angle orientation.

2. If B is a positive bounded unit, then C, A, D, α are all strong and m is weak. The new strong-complement pitchfork applies to this entire leaf at all upper event scales. In particular it includes C unbounded when L is infinitesimal and α unbounded when $1-H$ is infinitesimal.

Unbounded B is impossible by the original chart inequalities. The homogeneous all-tests theorem of Section 95.1 closes the remaining infinitesimal-B leaf without assuming a bounded actual state ratio. Thus both unbounded-b leaves, as well as all bounded-b openings, are now covered. Literal $n = 0$ is preserved.

Chapter 98

A unit native state scale: the same circuit with a different mass

Component `p890eea5f538b`
Source: `R13/baseline/baseline/baseline/baseline/baseline/advances/c1/unit_eta_unbounded_e.tex`
SHA256: `890eea5f538b9dae39c62552c1150d873a898f4e2a9e6e261d95f0c6b8b744c2`
Inventory use: entire scoped proof

98.1 A unit native state scale: the same circuit with a different mass

Theorem 98.1 (Unit η and unbounded e)**.** *Use the unchanged physical native C1 source, its original event signs, and the positive shapes and gaps a, t, M, J. Suppose k, ζ are positive infinitesimals, η is a positive bounded unit, $r = 1 - a\eta$ is a positive bounded unit, and $e = k\sqrt{\eta/\zeta}$ is unbounded. Then the full original parent is tensor SOS at every independent projective test and every real or complex state. No positive lower bound on a, t, M, J is imposed.*

Write $H = r + \eta$ and $\Delta = 1/e$. The seven native coquotients have residues $2rH$ for $c_2, c_b, c_3, c_5, c_\alpha$, 2 for c_4, and $4r$ for c_d. All are positive units. In particular the four retained packets of the contact–conductor proof remain positive and the remainder has bounded coefficients. We reuse their exact definitions, not asymptotic substitutes for the original source.

Outer tests. The exact formulas give

$$\frac{m}{d} = \frac{e^2 c_2}{M\eta c_d}, \qquad \frac{\beta}{d} = \frac{e^2 c_b}{t\eta c_d} - \frac{1}{2}.$$

Both ratios are unbounded. Apply the exact full-parent Alice reflection (AS3)–(AS4), which exchanges $m \leftrightarrow c$ and $\beta \leftrightarrow D$ while leaving v unchanged. The already proved two-strong-current identity then covers every test with $1/v$ bounded, for every Alice angle and every state. Its homogeneous form includes both literal Alice endpoints, so no inverse-angle point is lost.

For v infinitesimal and Alice infinity, use the critical-current identity with $g_c = cD/(c + D)$, $\epsilon = d/g_c$ and $\gamma = Ag_c/d^2$. Its actual sufficient conditions on this chart are bounded ϵ and positive-unit γ: in its exact four-root identity the correction L_0 has factors c_u or v, where $c_u = (1 + u^2)^{-1/2}$ is infinitesimal. Thus $\sqrt{\epsilon}$ need only be bounded. The residue Gram is $\mathrm{diag}(1, 2, 2, 4\bar{\gamma})$. Use the exact relation $J + a = T$ with $T - 1$ infinitesimal, and the cancelled original coefficient formula

$$\frac{D}{a} = \frac{2c_5/c_3 - J\eta c_d/c_3}{J}.$$

Its numerator has residue $2/\bar{H}$. The unchanged native formulas give

$$\frac{d}{a} \sim \frac{4\eta}{H}, \quad \frac{A}{a} \sim \frac{2\eta^2}{rH}, \quad \frac{D}{a} \sim \frac{2}{JH}, \qquad \epsilon \sim \frac{2\eta(JH+a)}{H}, \quad \gamma \sim \frac{H}{4r(JH+a)}.$$

The identity $\overline{JH+a} = 1 + \bar{\eta}(1-\bar{a})^2$ shows that the two required conditions hold even when a or J is infinitesimal. The calculation uses the cancelled harmonic coefficient g_c/a, so no lower bound on an opening is inserted. Homogeneous angle normalization includes literal Alice infinity. It remains to treat bounded u and infinitesimal v.

State entry. Normalize the state by a coordinate of maximal absolute value and put $D_u = 1 + u^2$ and $w = (W + uY)/D_u$. If w is a unit, the exact five-root current identity (UW2)–(UW3) applies with $Q = W + uY = D_u w$ a unit. Its first four diagonal residues are $8rH$ and its fifth is $8D_u$; both negative corrections have factors Δ or $\Delta^2\eta$. Thus this chart is strict.

If w is infinitesimal, the nonnegative pure-testing core is a direct exit. On the negative-core branch Z is a unit. Indeed its residue at $v = 0$ is $2\{(X + uZ)^2 + (uW - Y)^2\}$; if Z, w and both currents had zero residue, every normalized state coordinate would have zero residue. Normalize $Z = 1$. The exact core Schur identity gives $w > 0$ after the usual positive-angle sign gauges. All transformations here belong solely to the testing field.

The ordinary-u chart needs no contact division. Use the exact source change

$$\delta = \Delta R/K, \quad b^2 = \eta K/R, \quad K = 1 + k^2, \quad Q = K + ak^2b^2, \quad T_n = K^2 - t\delta^2k^4Qb^2.$$

All source denominators are positive units, δ, k are infinitesimal and $b^2 \sim \eta$. Let

$$A_0 = wv, \quad B_0 = \delta v, \quad C_0 = \delta b^2 uw, \quad D_0 = \delta^2 k^3 b^2, \qquad x = B_0 - C_0.$$

The five-term exact scalar belongs to $(A_0, B_0, C_0)^2 + (D_0^2)$ and both complete first normals belong to (A_0, B_0, C_0). Collecting these displayed factors before taking residues is legitimate over the bounded coefficient ring. At $\delta = k = w = v = 0$, keeping bounded u free, the scalar Gram is

$$8rHD_u^2A_0^2 + 8D_u\{x^2 + u^2rHB_0^2\} + 8rHD_uD_0^2. \tag{UE1}$$

The hard block on $(N, \delta M_B)$ is $\mathrm{diag}(16rH, 16rH + 8\eta^2/D_u)$, and the complete first normals have residues 0 and $16\eta x$. The latter is the coefficient of δM_B in the full polynomial, so the symmetric Gram off-diagonal entry between δM_B and x is 8η. The coupled $(\delta M_B, x)$ block has determinant

$$(16rH + 8\eta^2/D_u)\,8D_u - (8\eta)^2 = 128D_urH > 0. \tag{UE2}$$

If u is a unit, the remaining B_0 pivot in (UE1) is strict, as are the A_0, D_0, N pivots. Every omitted coefficient has zero residue. This proves the chart by a bounded strict Gram on these six roots, without any mixed inverse. Equation (UE1) follows directly from $a\eta + r = 1$: its (B_0, C_0) part before collection is $8D_u\{(1 + u^2rH)B_0^2 - 2B_0C_0 + C_0^2\}$.

The infinitesimal-u chart uses the old conductor. Now δ, k, u, w, v are infinitesimal, while b, a, t are bounded. Use the unchanged roots

$$y = A_0 - kC_0, \quad z = uB_0 - kC_0, \quad h = kC_0 - D_0, \quad q_1 = a_1a_2 - a_3a_4,$$

and the second pairing $q_2 = a_1a_3 - a_2a_4$ of the four separated radicals. Set

$$T_4 = \sum_{i=1}^{4} a_i^4 - 4\prod_{i=1}^{4} a_i = y^2 + z^2 + h^2 - 2yz + 2q_1^2, \qquad \rho = (1 - ab^2)(1 + (1-a)b^2) - 1.$$

Give $(\delta, k, b, a, t, u, w, v)$ weights $(1, 1, 0, 0, 0, 1, 2, 3)$. The root x has weight four and the other soft roots have weight five. Substitution at the exact contact preserves this grading. Writing E_0 for the complete scalar correction relative to the old leading polynomial, the five-term factorization and the two complete normal formulas give

$$\begin{aligned} E_0 &= 8\rho T_4 - 8(\rho + b^2)k^2x^2 + O_{\mathrm{wt}}(12), \\ L_1 &= O_{\mathrm{wt}}(6), \qquad L_2 = 16b^2x + O_{\mathrm{wt}}(6). \end{aligned} \tag{UE3}$$

These identities require only multiplication of the short source polynomials. In particular the residue correction is the single factor ρ multiplying the same circuit, rather than a new polynomial positivity problem.

The unlocalized monic contact–conductor lift applies unchanged. After the terms explicitly displayed in (UE3) are removed, scalar coefficient weights are at least two and normal coefficient weights are at least one. Therefore they all have zero residue. There is no division by k, b, δ, a, t or a contact difference. The leading strict soft Gram is now

$$8x^2 + 8rHT_4.$$

The two-pairing circuit identity supplies a strict Gram on (y, z, h, q_1, q_2), and rH is a positive unit. The complete hard block is $\mathrm{diag}(16rH, 16rH + 8\eta^2)$; its only nonzero first-normal residue is $16\eta x$. Its coupling to the $8x^2$ pivot has determinant $128rH > 0$. Retain the old positive g_4^2 square. The resulting eight-root remainder Gram is strict.

Finally restore all four positive packets, the positive square just retained, and the positive pure-coefficient multiplier. The state and angle cases above exhaust the independent tests. Complex states split into their real and imaginary quadratic evaluations. The result closes the entire stated source domain, rather than only an optimizing or matching testing chart.

Scope. This theorem adds the positive-unit η, r slice with unbounded e. It does not cover infinitesimal r, unbounded η, bounded e, or the other native complement rows. Those domains must not be marked complete by this result.

Chapter 99

Physical entry excludes a small state parameter

Component `p7294aae40ee2`
Source: `R13/baseline/baseline/baseline/baseline/baseline/advances/c2/c2_bounded_state_entry.tex`
SHA256: `7294aae40ee2774cc19176ebd1d92b4079f29a9b74246870d3d965b62bc97f1a`
Inventory use: Exact lower bound for bounded b, before any single-opening theorem

99.1 Physical entry excludes a small state parameter

This short reduction concerns the same exact rational source, before assuming that b is a unit. Suppose $x > 0$ is infinitesimal, k is a positive bounded unit, $b > 0$ is bounded, $0 < r < 1$, $0 < h < b$, $g = b - 1 + r + xh > 0$, and $n \geq 0$. No further source physicality is needed for the following necessary bound.

All quantities in

$$(b + r)A_c S_c \geq r\Delta$$

are bounded, so taking residues gives

$$r_0\{b_0 + k_0^2(b_0 + 1)\} \leq b_0.$$

The brace is strictly positive even if $b_0 = 0$. On the other hand $g > 0$ gives $r_0 \geq 1 - b_0$. If $b_0 \leq 1$, combining the two inequalities yields

$$(1 - b_0)\{b_0 + k_0^2(b_0 + 1)\} \leq b_0, \quad \text{or equivalently} \quad b_0^2 \geq \frac{k_0^2}{1 + k_0^2}.$$

For $b_0 \geq 1$ this lower bound is automatic. Hence in every case

$$b_0 \geq \frac{k_0}{\sqrt{1 + k_0^2}} > 0.$$

Thus bounded b is necessarily a unit. In particular there is no additional b-infinitesimal source domain in this actual-contact scaling. Combining the two new single-angle certificates, the uncaptured part of this bounded-b entry domain has exactly the joint necessary degeneration $r_0 = g_0 = 0$, hence $b_0 = 1$; the other remaining scale is unbounded b. This is a necessary source reduction, not a certificate for the joint degeneration.

Chapter 100

The remaining opening when the opposite event is bounded

Component `pd7bd7f0d7fb6`
Source: `R13/baseline/baseline/baseline/baseline/baseline/advances/c2/c2_last_opening.tex`
SHA256: `d7bd7f0d7fb66e4e45a7fceaac4ff793b5d046b7874bc3879cd8bd20219c3c81`
Inventory use: CG1–CG10 complete gInf,runit opening, triangular m hard normal and literal n=0

100.1 The remaining opening when the opposite event is bounded

Theorem 100.1 (The unbounded-m opening with bounded n)**.** *In Theorem 101.1, replace the assumption that $g = b - 1 + r + xh$ is a positive bounded unit by $g > 0$ and the assumption that at least one of r, g is a bounded unit. The all-tests tensor-SOS conclusion still holds.*

The only new case is infinitesimal g; it is automatically bounded. Our hypothesis now makes r a unit. Since $b_0 + r_0 = 1$ and b is a positive unit, $r_0 < 1$. Thus n is bounded, $0 < b_0 < 1$, and $1 - r$ is a unit. The source formulas imply

$$mg = \frac{(1 + k^2)(1 - r)A_c S_c}{2\Delta}, \tag{CG1}$$

which is a positive bounded unit. Thus m is unbounded. The two-opening bounds (CO1), positivity of $D/x, \alpha/x$, and the consequence that $b - k^2 r$ is a unit remain valid.

Use the original strong amplitude as the hard normal. Put $\eta = h/b$ and use the exact contact graph

$$Y_g = \eta u W, \qquad X_g = v(1 + \eta u^2)W - (1 + g)uZ. \tag{CG2}$$

At contact this gives $X_g = \rho[b + xh - 1 - g] = -\rho r$. The original m amplitude satisfies exactly $M|_g = guZ$. Normalize the full hard directions by

$$E_M = \sqrt{m}\{X - X_g - uv(Y - Y_g)\}, \qquad E_Y = \sqrt{C + A}(Y - Y_g). \tag{CG3}$$

In these variables

$$M = guZ - E_M/\sqrt{m}, \quad X = X_g + E_M/\sqrt{m} + uvE_Y/\sqrt{C + A}.$$

Consequently the unbounded packet has no E_Y contribution at all. This triangular change is essential: scaling X alone would leave the uncontrolled coefficient $\sqrt{m/(C + A)}uv$. The entire scalar contribution of that packet is $4mg^2u^2Z^2$, which is bounded by (CG1). Its full X derivative contributes $-8mg\,uZ$, also bounded, before division by $\sqrt{m}$.

All other source weights entering the X row are n, D and the unit signed weight, so the full X derivative on (CG2) has bounded coefficients on $V = vW, B = uZ$. Hence the E_M first normal has zero-residue bounded coefficients on those roots. The E_Y first normal is

$$\frac{(F_Y + uvF_X)|_g}{\sqrt{C + A}}. \tag{CG4}$$

The m packet cancels identically in $F_Y + uvF_X$, before any graph substitution. The remaining large terms are exactly the C, A terms controlled by (CO1); all other coefficients are bounded. In particular the old (CO5) form on P, B, E holds with bounded coquotients and identically zero leading corrections. The complete quadratic hard block is bounded and has residue

$$\operatorname{diag}(4, 4D_{v,0}) \tag{CG5}$$

on the finite $u = 0$ axis. The m packet gives precisely its first diagonal entry, the C, A packets give the second, and all mixed entries from bounded packets contain $1/(\sqrt{m}\sqrt{C + A})$.

The remaining five Laurent terms already force strictness. The exact scalar divisions on P, B, V, E are bounded for the same reason: C, A enter through (CO1), and m only through mg^2. Their leading weighted scalar is

$$q\,u^2W^2/x + b_s u^2 Z^2 + c_s W^2 - d_s uWZ + exZ^2,$$

where, for $z = v^2$ and $\nu = 2n_0 + 1 \geq 1$,

$$\begin{aligned} &q = Q(1 + z), \quad b_s = 2\nu(1 + z) - 2, \quad c_s = 2z\nu(1 + z), \quad d_s = 4v\nu(1 + z), \\ &e = E_2 z + E_0, \qquad Q = 4T_0/b_0, \quad E_2 = 4(D/x)_0 > 0, \quad E_0 = 4(\alpha/x)_0 > 0. \end{aligned} \tag{CG6}$$

Set $\mathcal{D}(s, z) = (qs + c_s)(b_s s + e) - d_s^2 s/4$. Its exact Laurent identity is

$$\begin{aligned} \frac{\mathcal{D}(s, z)}{(1 + z)sz} = {} & 2Q(\nu - 1)s/z + 2Q\nu s + 2\nu E_2 z/s + 2\nu E_0/s \\ & + QE_0/z + QE_2 - 4\nu. \end{aligned} \tag{CG7}$$

Contact and stationarity give a zero at $(s, z) = (1, k_0^2)$. After $z = k_0^2 t$, write the five coefficients in order as a, b, c, d, e. Here $a \geq 0$ and $b, c, d, e > 0$. Stationarity gives $c = a + e$ and $b = d + e$. The zero value fixes the constant, so exactly

$$\mathcal{L} = a(s/t + t/s - 2) + d(s + 1/s - 2) + e(t/s + s + 1/t - 3). \tag{CG8}$$

Every summand is nonnegative by AM–GM. The last bracket alone forces $s = t = 1$ at equality: its three positive terms have product one, and equality makes all three equal. Its exponent vectors $(-1, 1), (1, 0), (0, -1)$ span two dimensions; equivalently its Hessian in $\log s, \log t$ at $(1, 1)$ is $\left(\begin{smallmatrix} 2 & -1 \\ -1 & 2 \end{smallmatrix}\right)$. Hence it already has a positive definite contact Hessian. This proves global uniqueness and local nondegeneracy even when $n_0 = 0$ and the first pair is absent.

For remote finite nonzero v the same scalar cross-term split as before gives a strict Gram. The case $v_0 = 0, n_0 > 0$ is also strict on P, B, V, E. The full normals are restored by (CG3)–(CG5). At matched finite-angle residues the whole scalar still has the form $p^2 Q_b/x + B_b$ with bounded Q_b, B_b. The four monic conductors R, S, U, T of the prior theorem therefore give a bounded Gram, and (CG8) proves its strict Schur inequality.

The full first normals have the required same conductor lifts. For E_M, their coefficient is $(uA_1 + WB_1)/\sqrt{m}$ with bounded polynomial A_1, B_1. Monic contact division retains the factor $1/\sqrt{m}$ in every coquotient. Exact contact makes the residual coefficient $A_1 + \kappa B_1$ vanish, giving the old four-root lift with zero residue. For E_Y, the cancellation in (CG4) leaves precisely the old large-packet term B_2 plus bounded residual terms; its exact contact and identically zero leading part give the same $RB_2 + \rho E_2'$ division, then the bounded factor $1/\sqrt{H}$. These are full normal identities, not just Hessian checks. Quadratic state degree and (CG5) finish the complete matched-contact certificate.

The double-zero test when $n_0 = 0$. The graph (CG2) is replaced in this remote corner. Put

$$H_c = 4m + 4n + 2, \quad \xi = 4m/H_c, \quad \zeta = (4m+2)/H_c, \quad \eta_c = C/(C+A).$$

Use

$$Y_g = \eta_c u W, \quad X_g = \xi v(1 + \eta_c u^2)W - \zeta u Z,$$

and the triangular normals

$$E_X = \sqrt{H_c}\{X - X_g - uv(Y - Y_g)\}, \qquad E_Y = \sqrt{C+A}(Y - Y_g). \tag{CG9}$$

The entire graph amplitude is

$$M|_g = (1-\xi)v(1+\eta_c u^2)W + (\zeta - 1)uZ.$$

Both $m(1-\xi)$ and $m(\zeta - 1)$ are bounded because n is bounded. Thus the scalar and full first normals have bounded coquotients after the same x scaling as before.

Set $\epsilon = 4\alpha + 4n\zeta > 0$. It is infinitesimal, with $n/\epsilon, x/\epsilon$ bounded. The complete scalar Z^2 coefficient is exactly

$$f_{zz} = 4\alpha + \epsilon u^2 + 4n\zeta^2 u^4 + 4Dv^2 + u^2 v^2 B_b, \tag{CG10}$$

where B_b is bounded with residue 2. This identity follows directly from the packets and the exact relation $4n\zeta^2 + (4m+2)(1-\zeta)^2 = 4n\zeta$. The W^2 coefficients on $P = uW/\rho, V = vW$ have positive residues $Q, 2$. The cross term is $uvWZ$ times a bounded function and can be written as a zero-residue multiple of PE using $uvWZ = vPE$. Thus the five scalar roots

$$P, \quad V, \quad E = \rho Z, \quad B_\epsilon = \sqrt{\epsilon}\, uZ, \quad T = uvZ$$

have strict diagonal residue $(Q, 2, E_0, 1, 2)$.

For the full X derivative, at $v = 0$ the Z coefficient is exactly $-8n\zeta u^3$. After division by $\sqrt{H_c}$ its coefficient on B_ϵ is infinitesimal because $n/\sqrt{\epsilon} = (n/\epsilon)\sqrt{\epsilon}$ is infinitesimal. The remaining Z coefficient has the factor uv^2 and bounded coquotient; its coefficient on T has zero residue. The W coefficient has a factor v and bounded coquotient, multiplied by $1/\sqrt{H_c}$. For E_Y the exact m cancellation in $F_Y + uvF_X$ again applies; (CO6) gives $B_2 = 8D_v(xA)\eta_c u^2$, and the remaining lift is

$$P(B_2 + xe_w) + T(\rho u e_z) + E\, 8v(D - \alpha),$$

divided by $\sqrt{H}$, with bounded e_w, e_z. Every coefficient has zero residue. The hard residue of (CG9) is $\mathrm{diag}(1, 4)$, and all mixed hard entries vanish in residue. This proves the full-parent corner including literal $n = 0$.

All other tests. For the remaining commuting axes use the earlier outer identities, cap C, A at σ/x as before, and also cap m at $\min(m, 1)$. Retain the entire nonnegative excess m packet. The selected m weight is a unit and all previous displayed strict residues apply. The noncommuting-residue theorem covers the other tests. Restore all positive factors and split complex states as before. This proves the theorem for arbitrary positive g, including the new unbounded-m regime.

The remaining assumptions in this source family are now only that k, b are units, x is positive infinitesimal, all original contact openings are positive, at least one of r, g is a unit, $n \geq 0$, and the source is physical. The joint degeneration $r, g \to 0$, where both n, m can be unbounded, is not covered. The result does not yet treat unbounded b or a degenerate residue of the finite actual angle k. Section 99.1 excludes infinitesimal b under these entry hypotheses.

Chapter 101

The one-weak C2 family with three arbitrarily small openings

Component p81893ad08bed
Source: R13/baseline/baseline/baseline/baseline/baseline/advances/c2/c2_openings_and_unbounded_n.tex
SHA256: 81893ad08bed97aa8f133eedf1f5dfa35d77c4a9500ec4add025d102f2c451fb
Inventory use: CO1–CO11 complete h,b-h,r openings with gunit, full large-n normal and all tests

101.1 The one-weak C2 family with three arbitrarily small openings

This theorem removes three unit assumptions from the source of Theorem 105.1. Both positive openings h and $b-h$ may approach zero, and r may approach zero, making n arbitrarily large. The conclusions concern the entire original parent and every independent test. Source physicality remains an explicit hypothesis; no new source classification is claimed.

Theorem 101.1 (Three-opening single-angle source)**.** *Use exactly the rational C2 source and contact of Section 105.1. Suppose $x>0$ is infinitesimal, k,b are positive bounded units,*

$$0<r<1, \qquad 0<h<b,$$

and $g=b-1+r+xh$ is a positive bounded unit. Assume $n\geq 0$ and physicality of the original source. There is no lower bound on $r,h,b-h$. Then the entire original parent has tensor SOS at all independent projective measurement pairs and all real or complex projective states. Literal $n=0$ is included.

We write $\rho=\sqrt{x}$, $\eta=h/b$, and retain the exact source quantities A_c,S_c,Δ from that section. All arguments below take place over the independent coefficient and testing fields. Every coefficient inverse used below is pure coefficient or pure testing, as specified; no mixed denominator is inverted.

Entry and the two complementary openings. First the former assumptions that $1-r$ and $b-k^2r$ are units follow from $n\geq 0$. Indeed all quantities in

$$(b+r)A_cS_c\geq r\Delta$$

are bounded, and $\Delta_0=(1+k_0^2)(b_0+1)$. Taking residues yields

$$r_0\leq \frac{b_0}{b_0+k_0^2(b_0+1)}<1, \qquad b_0-k_0^2r_0\geq \frac{b_0^2(1+k_0^2)}{b_0+k_0^2(b_0+1)}>0.$$

Thus m is a positive bounded unit, and $D/x,\alpha/x$ are positive bounded units, regardless of whether r has zero residue.

Set

$$T_C = xC(b-h), \qquad T_A = xAh, \qquad H = x(C+A).$$

The original source formulas, with their apparent opening denominators cancelled, show that T_C, T_A are bounded positive units and have the same residue

$$T_0 = \frac{k_0^2(b_0+r_0)}{2\Delta_0} > 0.$$

Consequently H is bounded below by a positive real, although it may be unbounded. The four identities

$$\begin{aligned} xC(1-\eta) &= T_C/b, & xA\eta &= T_A/b, \\ xC(1-\eta)^2 &= T_C(b-h)/b^2, & xA\eta^2 &= T_A h/b^2 \end{aligned} \tag{CO1}$$

are the reason no new scalar singularity appears when either opening collapses. In particular their last two terms add to a bounded unit, with residue T_0/b_0. The effective scalar coefficient is

$$Q = 4T_0/b_0 > 0. \tag{CO2}$$

It is independent of the limiting value of h.

A bounded n: restoration with the normalized hard root. If r is a unit, the exact formula for n makes n bounded. Use the contact graph $X_g, Y_g = \eta uW$ of the prior theorem. All of its scalar coefficients remain bounded: the only potentially unbounded weights are C, A, and on the graph they occur solely through the last two products in (CO1), divided by x and multiplied by u^2W^2. Its leading scalar is exactly the scalar in that theorem with Q replaced by (CO2). The quantities n_0, m_0, D_*, E_* have their former values. Hence the entire six-term Laurent identity, its balanced AM–GM decomposition, global equality characterization, and positive contact Hessian apply without change.

Replace the old hard root $N_Y = (Y - Y_g)/\rho$ by

$$\widehat{N}_Y = \sqrt{H}\, N_Y = \sqrt{C+A}(Y - Y_g). \tag{CO3}$$

Its exact hard coefficient from the C, A packets is

$$4D_v \frac{C + AD_u}{C+A} = 4D_v \left(1 + \frac{A}{C+A}u^2\right), \tag{CO4}$$

which is bounded and has residue $4D_{v,0}$ on the finite $u = 0$ residue axis. The other packet contributions to this coefficient and to the mixed hard entry are bounded: $1/(C+A) = x/H$ and $1/\sqrt{C+A} = \rho/\sqrt{H}$ are bounded and infinitesimal.

For clarity the complete first-normal verification is also retained. The previous unnormalized Y derivative has exactly the form

$$L_Y = P(B_2 + xe_w) + B(\rho u e_z) + Ee_*, \qquad B_2 = 8D_v\{(xA)\eta D_u - (xC)(1-\eta)\}, \quad e_* = 8v(D-\alpha). \tag{CO5}$$

Here $P = uW/\rho$, $B = uZ$, $E = \rho Z$. The functions e_w, e_z come only from the bounded n, m, D, α packets and the signed core, so remain bounded. Equation (CO1) makes B_2 bounded with identically zero residue for every finite v on $u = 0$; e_* has the same property. The coefficient of $\widehat{N}_Y$ is exactly $L_Y/\sqrt{H}$, so its bounded zero-residue lifts are preserved. The X derivative contains neither C nor A and is unchanged. Thus all four-root remote certificates restore both full normals.

For matched finite-angle residues the scalar has exactly the old form $f = p^2 Q_b/x + B_b$, with

$$Q_b = 4D_v\{(xC)(1-\eta)^2 + (xA)\eta^2 D_u\}.$$

Both Q_b and B_b have bounded coefficients by (CO1). The monic divisions in $u^2 - x$, $W^2 - \kappa^2 x$, $uW - \kappa x$, and $v - k$, where $\kappa = b/k$, therefore give precisely the same bounded Gram on R, S, U, T. Its residue is unchanged. The old first-normal contact divisions use only bounded B_2, e_w, e_z and exact stationary contact; multiplication by the pure coefficient $1/\sqrt{H}$ preserves their zero-residue lifts. Hence the complete six-root Gram has strictly positive residue.

If $n_0 = 0$ at the double-zero test, use the previous exact graph with $\eta_c = C/(C + A)$. Now

$$xC(1 - \eta_c) = xA\eta_c = \frac{xCA}{C + A}, \qquad \left(\frac{xCA}{C + A}\right)_0 = T_0/b_0. \tag{CO6}$$

These quantities are bounded units even when H is unbounded. Thus the exact ϵ reserve and five scalar roots in that proof remain valid. Its Y normal is again divided by $\sqrt{H}$, and its hard Y residue is now 4 rather than $4(C_* + A_*)$. This proves the literal $n = 0$ case as well as its infinitesimal neighborhood without deleting a state coordinate.

An infinitesimal r: the opposite event can be unbounded. Suppose now r is infinitesimal. Let $N = 4n + 2$. The exact source formula gives

$$Nr = \frac{2(b + r)A_c S_c}{\Delta}. \tag{CO7}$$

Thus Nr is a bounded positive unit, N is unbounded, and both nr and nr^2 are bounded. Use the exact-contact graph

$$X_g = -ruZ, \qquad Y_g = \eta uW, \qquad \widehat{N}_X = \sqrt{N}(X + ruZ), \quad \widehat{N}_Y = \sqrt{C + A}(Y - \eta uW). \tag{CO8}$$

At the original contact $u = \rho, v = k, W = \rho b/k, Z = 1$ this graph has exactly the required X, Y values. The potentially large n packet contributes $4nr^2 D_u D_v u^2 Z^2$ to the graph scalar, and $-8nrD_u D_v uZ$ to its unscaled X derivative. Equation (CO7) therefore controls the entire scalar and every full first normal, not only their values at contact.

The hard residue on $(\widehat{N}_X, \widehat{N}_Y)$ is

$$\operatorname{diag}(D_{v,0}, 4D_{v,0}). \tag{CO9}$$

Indeed $H_{XX}/N = D_u D_v + O(1/N)$, since the remaining source weights that enter H_{XX} are m, D; (CO4) controls H_{YY}. The mixed entry is bounded times $1/(\sqrt{N}\sqrt{C + A})$ and has zero residue.

The five exact scalar monic divisions on P, B, V, E from the previous proof still apply. Their residues, with $z = v^2$, are now

$$q = Q(1 + z), \quad b_s = 2\mu, \quad c_s = 2(\mu - 1)z, \quad d_s = 4\mu v, \quad e = E_2 z + E_0, \tag{CO10}$$

where $\mu = 2m_0 + 1 > 1$, $E_2 = 4(D/x)_0 > 0$, $E_0 = 4(\alpha/x)_0 > 0$, and Q is given by (CO2). Define $\mathcal{D}(s, z) = (qs + c_s)(b_s s + e) - d_s^2 s/4$. Exactly,

$$\begin{aligned}\frac{\mathcal{D}(s, z)}{sz} &= 2Q\mu s/z + 2Q\mu s + 2(\mu - 1)E_2 z/s + 2(\mu - 1)E_0/s \\ &\quad + QE_0/z + QE_2 z + Q(E_0 + E_2) - 4\mu.\end{aligned} \tag{CO11}$$

All six nonconstant Laurent coefficients are positive. Exact stationary contact, applied after the bounded graph substitution $u = \rho s_1, W = \rho w_1$, gives a stationary zero at $(s, z) = (1, k_0^2)$. Alternatively this follows by direct substitution of

$$m_0 = \frac{1}{2(b_0 - 1)(b_0 + 1)}, \quad Q = \frac{2k_0^2}{\Delta_0}, \quad E_2 = \frac{2b_0^2}{k_0^2 \Delta_0}, \quad E_0 = \frac{2k_0^2 b_0^2}{\Delta_0}.$$

Here $b_0 > 1$ follows from $g_0 = b_0 - 1 > 0$. Rescale $z = k_0^2 t$ and use exactly the balanced reciprocal-pair and three-cycle identity of the previous theorem. Its $s + 1/s - 2$ and $t + 1/t - 2$ coefficients are

strictly positive. Thus (CO11) has a unique zero in the positive quadrant and a positive definite Hessian there. This proves remote scalar strictness and the matched-contact Schur inequality, with no appeal to numerical evidence.

For the full X normal, direct differentiation and parity give

$$F_X|_g = VA_X + BB_X,$$

where the complete coefficients are

$$\begin{aligned} A_X =& -8m(1+\eta u^2) + (8D+4)(1-\eta)u^2, \\ B_X =& -8nrD_uD_v + 8m(1-r) - (8D+4)v^2(1+ru^2) + 4D_v(1-r). \end{aligned}$$

These are bounded polynomials in the bounded testing coordinates. Consequently the coefficient of $\widehat{N}_X$ has bounded zero-residue coefficients on V, B because $1/\sqrt{N}$ is infinitesimal. The full Y normal is (CO5), with the bounded graph (CO8); (CO1) gives all of its required bounds and zero residues. Its complete residual coefficients are

$$\begin{aligned} e_w &= 8mv^2(1+\eta u^2) - (8D+4)(1-\eta) + (8\alpha+4)D_uv^2\eta, \\ e_z &= v\{-8m(1-r) + (8D+4)r - (8\alpha+4)\}. \end{aligned}$$

This restores the remote scalar certificates, including $v_0 = 0$, where (CO10) has strictly positive coefficients on P, B, V, E and zero cross residues. No special $n_0 = 0$ corner occurs in this case.

At a matched residue write, exactly,

$$\frac{F_X|_g}{\sqrt{N}} = uA_1 + WB_1 = -\kappa B_1 S + u(A_1 + \kappa B_1).$$

Both A_1, B_1 have identically zero residue for every finite v; their finite monic coquotients also retain the factor $1/\sqrt{N}$. Exact normal contact makes $A_1 + \kappa B_1$ vanish at the source contact. The same four conductors R, S, U, T therefore give bounded zero-residue coefficients, exactly as in the bounded-n argument. For the Y normal the previous division $L_Y = RB_2 + \rho E_2'$ has bounded B_2, E_2' with identically zero leading functions; division by $\sqrt{H}$ preserves those properties. All quadratic normal coefficients are already bounded by (CO4),(CO7),(CO9). Thus the matched-contact Gram for the entire parent is strictly positive in residue. The other state charts and the simultaneous sign copy use precisely the previous proof.

Every other commuting test. Choose the pure coefficient

$$\sigma = \min(1, xC, xA).$$

It is a positive bounded unit, since T_C, T_A are units while $h, b-h$ are bounded. Retain the nonnegative excess packets

$$4(C - \sigma/x)D_v(uW - Y)^2, \qquad 4(A - \sigma/x)D_uD_vY^2,$$

and use the four outer core identities in the previous theorem with $C = A = \sigma/x$. These identities retain the entire n packet; they never use boundedness of n. Their displayed hard entries $4C_*, 4A_*$ are simply replaced by $4\sigma_0$, while $m_0 > 0$ and $(D/x)_0 > 0$ are unchanged. Every residue Gram stays strictly positive. This covers the finite-unit u axis, both reciprocal axes and their double corners, including literal projective ends.

The finite $u = 0$ axis, those outer axes, and the inherited theorem for noncommuting-residue tests exhaust all independent measurements. Restore the retained packets and positive homogeneous factors, and split complex states into real and imaginary parts. This proves Theorem 101.1. In particular the newly admitted source scales permit either A or C to be arbitrarily larger than x^{-1}, independently of an arbitrarily large n. They cannot both have unbounded ratios to this scale because $h + (b - h) = b$ is a unit. There is no prescribed comparison between these coefficient scales and any testing infinitesimal.

Chapter 102

C2 pitchfork with three arbitrary event weights

Component `p6c62a53b1454`
Source: `R13/baseline/baseline/baseline/baseline/baseline/advances/c2/c2_three_large_events_pitchfork.tex`
SHA256: `6c62a53b1454a209db8cb324e8d59bbea44a164545fb7db47f20f972f250ba3f`
Inventory use: TA1–TA11 all n,A,D upper scales with bounded-unit C,alpha, full-parent normalization and real Schur proof; outer cap supplied explicitly by next body

102.1 C2 pitchfork with three arbitrary event weights

Theorem 102.1 (All scales of the three retained events)**.** *Let K_c, K_t be independent real closed extensions of $\mathbb{R}$. Let a physical canonical C2 source over K_c have $\lambda = \beta = 0$, with d, C, α positive bounded units and $n, m, A, D \geq 0$ arbitrary. Suppose an actual finite noncommuting stationary zero has $u_c = k > 0$ infinitesimal and $v_c = a > 0$ a bounded unit. Then the entire original parent is tensor SOS at every independent projective measurement pair and every real or complex projective state. No ratios among n, A, D are restricted.*

The bounded-A result is a special case. The mechanism is to keep all three possibly large event amplitudes as state coordinates. The exact parent and its amplitudes are (TD1); in particular $J = uW + uvX - Y - vZ$.

Source entry with three large weights. The initial coercivity estimate in the preceding proof uses only d, C, α, not A. Thus after $W_c = 1$,

$$X_c, Y_c, Z_c = O(k), \qquad m = O(k^2).$$

Now use $F_Z = 0$ first. The A and n packets do not enter this state equation. Every term other than $-8DvJ$ is $O(k)$, because $v_c = a$ is a unit and the just displayed bounds hold. Therefore $DJ_c = O(k)$. The Y equation then gives $AY_c = O(k)$, and the X equation gives $(d + 2n)X_c = O(k^2)$. Write

$$N = d + 2n, \quad T_A = d + 2A, \quad L = d + 2D, \qquad x = X_c/k^2, \quad y = Y_c/k, \quad j = J_c/k.$$

Consequently

$$x, y, j, Nx, Ay, Dj \text{ are bounded}, \qquad N^{-1} \text{ bounded}. \tag{TA1}$$

These estimates do not yet invert A or D.

Use the exact stationary equations (TD3) and divide both by $T_A L$. Now $t_2/T_A, t_6/T_A, t_3/L, t_4$ are positive bounded units. In the first equation the coefficient multiplying U^2 is (A/T_A) times a positive bounded unit: its remaining bracket is

$$4t_4 \left\{ \theta \frac{t_3}{L} + CV \frac{1 + 2D/N}{L} \right\},$$

whose lower bound follows from (TD4). The coefficient multiplying m is also a positive bounded unit:

$$4V\left\{\alpha\frac{t_8}{T_AL}+\frac{D}{L}\frac{t_6}{T_A}V\right\}. \tag{TA2}$$

Indeed, if D/L is bounded below, the second summand is a unit. Otherwise D is infinitesimal and L is a unit, while $t_8/T_A = 1 + 2(C + D)/T_A$ is a unit. All displayed factors have bounded upper bounds in either case. Hence

$$\frac{m}{U^2} = \frac{A}{T_A}\cdot(\text{positive bounded unit}). \tag{TA3}$$

The second stationary equation becomes

$$\frac{A}{T_A}\cdot(\text{positive unit}) + O\left(\frac{A}{T_A}U\right) = \frac{D}{L}\cdot(\text{positive unit}) + O\left(\frac{A}{T_A}U\right). \tag{TA4}$$

Thus A/T_A and D/L are comparable. If they were infinitesimal, A, D, m would all be infinitesimal, contradicting exactly the physicality test printed after (TD5). Therefore

$$A/T_A,\ D/L,\ m/k^4 \quad \text{are positive bounded units.} \tag{TA5}$$

In particular A, D have bounded reciprocals, but may be arbitrarily large. Set $\gamma = 4ma^2/k^4$, also a positive bounded unit.

The three-packet graph. On the local testing domain $\bar{v} = \bar{a} \neq 0$, put $b = 1 - j - y$ and use

$$X_g = xu^2W, \qquad Y_g = yuW, \qquad J_g = juW, \qquad Z_g = u\left(\frac{b}{v}+xu^2\right)W. \tag{TA6}$$

This graph matches the whole actual contact: the definition of J_c gives $Z_c/k = b/a + xk^2$. Its three hard normals are

$$E_X = \sqrt{N}(X - xu^2W), \quad E_Y = \sqrt{A}(Y - yuW), \quad E_J = \sqrt{D}(J - juW). \tag{TA7}$$

The exact inverse is

$$X = xu^2W + E_X/\sqrt{N}, \quad Y = yuW + E_Y/\sqrt{A}, \quad Z = \frac{uW + uvX - Y - juW - E_J/\sqrt{D}}{v}. \tag{TA8}$$

Every denominator is either pure coefficient or the pure testing unit v. All graph coefficients and all scaled normal coefficients are bounded. For the three large packets this follows explicitly from

$$nx^2,\ nx,\ n/N, \quad Ay^2,\ \sqrt{A}\,y, \quad Dj^2,\ \sqrt{D}\,j \quad \text{bounded}, \tag{TA9}$$

using (TA1),(TA5). Their quadratic normal contributions are exactly $4(n/N)D_uD_vE_X^2$, $4D_uD_vE_Y^2$, and $4E_J^2$. Every other original source coefficient is bounded, including m.

For clarity, the finite Taylor division used below works in the bounded Laurent polynomial ring in v: for positive integer r,

$$v^{-r} - a^{-r} = -(v-a)\frac{\sum_{i=0}^{r-1} v^i a^{r-1-i}}{a^r v^r}.$$

Thus a Laurent polynomial vanishing at $(U, v) = (K, a)$ belongs to $(U - K, v - a)$ with bounded Laurent coquotients; repeating the same identity gives the second-order division. There is no localization at a difference of the two fields.

Set $K = k^2$, $R = u^2 - (K/a)v$ and $T = u(v - a)$. The entire graph scalar is again $4mv^2 + u^2H_0(u^2, v)$, now with bounded Laurent coefficients in v. Its three scalar contact jets give (TD10),

and hence an exact bounded Gram on (R,T) with R residue pivot $\bar{\gamma}$. The full first normals before their pure source scale divisions are

$$\begin{aligned}(F_X+uF_Z)|_g &= -8mv+u^2L_X(u^2,v),\\ (F_Y-F_Z/v)|_g &= uL_Y(u^2,v),\\ (-F_Z/v)|_g &= uL_J(u^2,v).\end{aligned} \tag{TA10}$$

Divide the rows by $\sqrt{N},\sqrt{A},\sqrt{D}$, respectively. The scaled coefficients are bounded by (TA9). At contact they vanish. Laurent Taylor division gives exactly the root membership and infinitesimal R couplings proved in (TD11)–(TD12). The X row has infinitesimal coefficients on both R,T because $m/K=O(K)$; the other rows have infinitesimal coefficients on R. The exact quadratic state remainder is bounded by (TA8)–(TA9), so the entire original parent has a bounded five-root Gram on (E_X,E_Y,E_J,R,T).

Uniform strictness without a largest-event choice. The X and R residue pivots are $2(1+\bar{a}^2)$ and $\bar{\gamma}$. The remaining hard state form is still (TD13). At $u=0$, (TA8) gives

$$Y=E_Y/\sqrt{A},\qquad Z=-\frac{E_Y}{v\sqrt{A}}-\frac{E_J}{v\sqrt{D}}.$$

In these coordinates (TD13) is the sum of the two explicit pivots

$$4(1+v^2)E_Y^2+4E_J^2$$

and the nonnegative bounded form $(4C+2d)(1+v^2)Y^2+4\alpha(vY-Z)^2$. Its residue is therefore positive definite for every combination of bounded or unbounded A,D.

The boundary Schur function is the exact expression

$$f(v)=\frac{8(1+v^2)}{\det B(v)/(T_AL)}\left(\frac{Q_{10}}{T_AL}+\frac{Q_0}{T_AL}v^2+\frac{Q_{12}}{T_AL}v^4\right), \tag{TA11}$$

with Q_{10},Q_{12} as in (TD14). For v a bounded unit all its coefficients and denominator are bounded, and

$$\frac{Q_{10}}{T_AL}=4\alpha\frac{A}{T_A}\frac{t_3}{L},\qquad \frac{Q_{12}}{T_AL}=4C\frac{D}{L}\frac{t_2-2m}{T_A}$$

are positive units. The determinant has the positive term $16AD(1+v^2)v^2$, so its quotient by T_AL has a positive lower bound. Its other terms are bounded after that division. The exact middle coefficient is polynomial of degree at most one in each of A,D, so $Q_0/(T_AL)$ is bounded too.

The contact normal equations and scalar jets give $\bar{f}(\bar{a})=\bar{f}'(\bar{a})=0$ as before. Therefore the real quartic bracket equals $\overline{Q_{12}/(T_AL)}(v^2-\bar{a}^2)^2$. Its strict second derivative gives the positive soft Schur pivot on T. The whole five-root residue Gram is strict. This proves the local all-state certificate; the Bob sign identity gives its negative sign copy. Homogenization in W includes $W=0$.

Outer tests when both A and D are large. For u infinitesimal and v a bounded unit with $\bar{v}\neq\pm\bar{a}$, retain the entire m,n packets and use the four roots

$$X,\quad K_A=\sqrt{A}Y,\quad K_D=\sqrt{D}J,\quad uW.$$

Substitute $Y=K_A/\sqrt{A}$ and $Z=(uW+uvX-K_A/\sqrt{A}-K_D/\sqrt{D})/v$. The exact remainder is a bounded quadratic Gram. Its hard residue has the positive X pivot $2\bar{d}(1+\bar{v}^2)$ and the two explicit pivots just displayed; its soft Schur complement is $\bar{f}(\bar{v})>0$ by (TA11).

At u,v both infinitesimal, and at $u,1/v$ both infinitesimal, one need not divide by v. Cap C,A,D,α at the positive unit $\mu=\min(1,C/d,A/d,D/d,\alpha/d)$ after division by d, and retain all excesses and the full m,n packets. The bounded selected parent has an exact Gram on (X,Y,Z,uW) in the finite chart and on the same four roots in the homogeneous reciprocal chart. At $(u,v)=(0,0)$

its soft Schur complement is $8Q_{10,\mu}/\det B_\mu(0) > 0$; at $(u, 1/v) = (0, 0)$ it is $8Q_{12,\mu}/\det B_{\mu,\infty} > 0$. The hard blocks are positive by (TD13) and its homogeneous limit. Hence both entire endpoint neighborhoods have strict-residue Grams, including literal zeros and infinities. These expressions use only the bounded selected parent; no residue of an unbounded A or D is taken.

All other commuting tests use the same cap and the already printed open-axis and corner forms in the preceding all-tests proof. Their hypotheses require only the four lower bounds supplied by (TA5). Noncommuting residue tests use the retained arbitrary-physical-source theorem. Restore the positive pure factors and split complex states into real and imaginary parts. This completes every independent test for the stated source domain.

Coverage gained and remaining restriction. The theorem removes every relative-scale restriction among n, A, D from the finite pitchfork. Its reflected form covers the corresponding ordinary–infinite actual contacts. It still assumes C/d and α/d are bounded units and this actual-angle residue type. Unbounded C/d or α/d, other actual-angle types, and the exhaustive C2 join remain outside this theorem.

Chapter 103

C2 pitchfork with two arbitrarily large retained events

Component p0c7a15f800d9
Source: R13/baseline/baseline/baseline/baseline/baseline/advances/c2/c2_two_large_events_pitchfork.tex
SHA256: 0c7a15f800d954d9237cb833959cb1c13e6604732b368deb1ccf87b4678a016f
Inventory use: TD1–TD14 exact stationary source balances and whole-normal Taylor construction; outer reference replaced by explicit cap/corner proof in strong-complement body

103.1 C2 pitchfork with two arbitrarily large retained events

Theorem 103.1 (The n, D-unbounded C2 pitchfork)**.** *Let the physical canonical C2 source over a real closed coefficient field have $\lambda = \beta = 0$, with d, C, α positive bounded units, $A \geq 0$ bounded, and $n, m, D \geq 0$ arbitrary. Suppose there is an actual finite noncommuting stationary zero with $u_c = k > 0$ infinitesimal and $v_c = a > 0$ a bounded unit. Then the entire original parent is tensor SOS at every independent projective measurement pair and every real or complex projective state. There is no bound on n, D, or on their ratio. In fact physicality and contact force A to be a positive unit, D to be bounded below by a positive real, and m/k^4 to be a positive bounded unit.*

This extends the preceding pitchfork by removing the upper bound on D. A bound on m need not be assumed: it follows from physicality. The original D packet supplies a state normal, so an unbounded weight is not put into a bounded Gram coefficient.

Write the unchanged parent as

$$\begin{aligned} F &= dH + 4nD_uD_vX^2 + 4mR_2^2 + 4CD_vr^2 + 4AD_uD_vY^2 + 4DJ^2 + 4\alpha D_up^2, \\ H &= 2\{J^2 + D_up^2 + D_vB^2 - D_uD_vZ^2\}, \\ R_2 &= vW - X + uvY - uZ, \quad r = uW - Y, \quad J = uW + uvX - Y - vZ, \\ p &= vY - Z, \quad B = X + uZ, \quad D_u = 1 + u^2, \quad D_v = 1 + v^2. \end{aligned} \tag{TD1}$$

Entry and bounded contact coordinates. At $u = 0, v = a$, the state block on (X, Y, Z) of $dH + 4CD_vr^2 + 4\alpha D_up^2$ has positive definite residue. Its W column is $O(k)$ and its W^2 entry is $O(k^2)$. Completing this ordinary bounded hard block gives, for positive real constants b, B_0, the coefficient-field inequality

$$dH + 4CD_vr^2 + 4\alpha D_up^2 \geq b(X^2 + Y^2 + Z^2) - B_0k^2W^2$$

at the actual angles. All omitted packets are nonnegative. Thus an actual zero has $W \neq 0$ and, on setting $W = 1$, $X, Y, Z = O(k)$. Since $R_2 = a + O(k)$ is a unit, the same inequality and its retained m packet give $m = O(k^2)$.

The state equation $F_Y = 0$ gives $DJ = O(k)$: every other term is $O(k)$, using bounded A, C, α, d, $m = O(k^2)$ and the bounds just obtained. The equation $F_X = 0$ then gives $(d+2n)X = O(k^2)$, since the J contribution is $O(kDJ)$ and the m contribution is $O(m)$, while $Z = O(k)$. Consequently, with

$$N = d + 2n, \qquad x = X_c/k^2, \quad z = Z_c/k, \quad j = J_c/k,$$

we have

$$Nx,\ z,\ j,\ Dj \quad \text{bounded}, \qquad N^{-1} \text{ bounded}. \tag{TD2}$$

The bound on j follows directly from $X, Y, Z = O(k)$, even before a lower bound for D has been obtained. No inverse of D has yet been used.

For the scale entry, put $U = k^2, V = a^2$, $L = d + 2D$, $\theta = n/N$ and use

$$t_2 = d + 2(\alpha + m + A), \quad t_3 = d + 2(C + D), \quad t_4 = d + 2(\alpha + m),$$
$$t_6 = d + 2(\alpha + C + A), \quad t_8 = d + 2(C + A + D).$$

The original Laurent-determinant stationary equations, divided only by N, are

$$\begin{aligned} U^2\{4\theta A t_3 t_4 + 4CA(1 + 2D/N)t_4 V\} &= mV\{4\alpha t_8 + 4Dt_6 V\}, \\ 4\alpha A(1 + 2m/N)t_3 + 4\theta A t_3 t_4 U &= \{4mDt_6/U + 4CDt_2\}V^2. \end{aligned} \tag{TD3}$$

Here $t_2, t_4, t_6, t_3/L, t_8/L$ are positive bounded units. After division by L, the braces in the first equation are A times a positive unit on the left and a positive unit on the right. To check the lower bound uniformly when both n, D diverge, observe

$$\theta + \frac{1 + 2D/N}{L} = \frac{n+1}{N} + \frac{2n}{NL}, \tag{TD4}$$

which is a positive bounded unit; all its neighboring factors in (TD3) have positive upper and lower real bounds. Thus m/U^2 is A times a positive unit. The second equation, again divided by L, has the form

$$A \cdot (\text{positive unit}) + O(AU) = (D/L) \cdot (\text{positive unit}) + O(AU). \tag{TD5}$$

If A were infinitesimal, then D/L and hence D would be infinitesimal, as would m. This is incompatible with physicality: on the test state $(W, X, Y, Z) = (1, 0, u, zu)$ the n packet vanishes exactly, and at the residue source $m = A = D = 0$ choose

$$z = \frac{v(2\bar{\alpha} + \bar{d})}{2\bar{\alpha}}.$$

The coefficient of u^2 in the full polynomial is then $-\bar{d}v^2(2\bar{\alpha} + \bar{d})/\bar{\alpha} < 0$ for real $v \neq 0$. The bounded residue polynomial cannot be nonnegative for all real small u. Therefore A is a unit. Equations (TD3)–(TD5) now prove that D/L and m/k^4 are positive bounded units. In particular $D > 0$ and D^{-1} is bounded. Set $\gamma = 4ma^2/k^4$.

A graph that keeps both large packets bounded. For arbitrary independent testing angles use the exact graph

$$X_g = xu^2W, \qquad Z_g = zuW, \qquad J_g = juW,$$

which fixes

$$Y_g = u(1 - j - vz + xu^2 v)W. \tag{TD6}$$

It matches the entire actual contact, including its Y coordinate. The three exact state normals are

$$E_X = \sqrt{N}(X - xu^2W), \qquad E_J = \sqrt{D}(J - juW), \qquad E_Z = Z - zuW. \tag{TD7}$$

The inverse uses only pure coefficient inverses:

$$X = xu^2W + E_X/\sqrt{N}, \quad Z = zuW + E_Z, \quad Y = u(1 - j)W + uvX - vZ - E_J/\sqrt{D}. \tag{TD8}$$

The n packet on the graph is $4nx^2u^4D_uD_vW^2$, and the D packet is $4Dj^2u^2W^2$. Both have bounded coefficients by (TD2). Their linear coefficients are bounded too, since nx and $\sqrt{D}\,j = (Dj)/\sqrt{D}$ are bounded. The exact quadratic normal block is bounded: its only large packets become $4(n/N)D_uD_vE_X^2$ and $4E_J^2$.

Set $W = 1$, $K = k^2$, and

$$R = u^2 - (K/a)v, \qquad T = u(v-a).$$

The entire graph scalar has the exact shape

$$F|_g = 4mv^2 + u^2H_0(u^2, v), \qquad H_0 \in B_c[u^2, v]. \tag{TD9}$$

The three contact scalar jets give exactly the same monic division as for the preceding pitchfork:

$$F|_g = \gamma R^2 + u^2\{(u^2-K)^2H_{20} + (u^2-K)(v-a)H_{11} + (v-a)^2H_{02}\}, \tag{TD10}$$

with bounded polynomial coquotients. After $u^2 - K = R + (K/a)(v-a)$ this is a bounded Gram on (R, T). Its R residue pivot is $\bar{\gamma} > 0$.

The full first normals, before the indicated scale divisions, have exact shapes

$$(F_X + uvF_Y)|_g = -8mv + u^2L_X(u^2, v), \quad (-F_Y)|_g = uL_J(u^2, v), \quad (F_Z - vF_Y)|_g = uL_Z(u^2, v). \tag{TD11}$$

The coefficient of E_X is the first expression divided by $\sqrt{N}$; that of E_J is the second divided by $\sqrt{D}$. These scaled polynomials have bounded coefficients, as shown above. Contact makes $L_X(K, a) = 8ma/K$, and makes the scaled J, Z functions vanish at (K, a). Monic division by $u^2 - K, v - a$ therefore puts every first normal in (R, T) with bounded coquotients. Every coefficient coupling E_X to R, T is infinitesimal; the R coefficients in the E_J, E_Z rows are infinitesimal. For example, the X row is exactly

$$\frac{1}{\sqrt{N}}\left[(8ma/K + u^2L_{X1})R + u\{(K/a)L_{X1} + L_{X2}\}T\right], \tag{TD12}$$

and the other two rows have form $uL_{i1}R + \{(K/a)L_{i1} + L_{i2}\}T$ after their scale division. State degree is two, so there are no omitted higher-normal terms. Equations (TD7)–(TD12) give an exact bounded full-parent Gram on (E_X, E_J, E_Z, R, T).

The residue Gram is strict. The X pivot has residue $2(1 + \bar{a}^2)$ and the R pivot has residue γ. For the remaining hard pair the unscaled state form is

$$B(v)[Y, Z] = \{4(C + A) + 2d\}(1 + v^2)Y^2 + 4D(Y + vZ)^2 + 4\alpha(vY - Z)^2. \tag{TD13}$$

In the scaled coordinates $Y = -vE_Z - E_J/\sqrt{D}$, $Z = E_Z$ its coefficients are bounded and its residue is positive definite. If D is bounded, this follows from the invertible residue change and the positive form (TD13). If D is unbounded, its residue is

$$4E_J^2 + \big(\{4(\bar{C} + \bar{A}) + 2\bar{d}\}(1 + v^2)v^2 + 4\bar{\alpha}(1 + v^2)^2\big)E_Z^2 \succ 0.$$

The exact minimized boundary coefficient of u^2, before residue, is

$$f(v) = \frac{8(1 + v^2)}{\det B(v)/L}\left(\frac{Q_{10}}{L} + \frac{Q_0}{L}v^2 + \frac{Q_{12}}{L}v^4\right), \tag{TD14}$$

where

$$Q_{10} = 4\alpha A[d + 2(C + D)], \qquad Q_{12} = 4CD[d + 2(\alpha + A)].$$

The middle coefficient Q_0 is defined by the exact Schur numerator; its explicit value is unnecessary. All quantities in (TD14) are bounded, $\det B/L$ is a positive unit for bounded real v, and $Q_{10}/L, Q_{12}/L$ are positive units. One may verify boundedness of Q_0/L directly from the determinant expansion,

which is affine in D. This is only a residue calculation of a real hard Schur complement, not inversion of a mixed tensor element.

The full contact normal and scalar equations imply that the residue of (TD6) at $v = \bar{a}$ minimizes the hard form, and that $\bar{f}(\bar{a}) = \bar{f}'(\bar{a}) = 0$. Thus the real quartic bracket in (TD14) equals

$$\overline{Q_{12}/L}\,(v^2 - \bar{a}^2)^2.$$

Its second derivative at $\bar{a}$ is strictly positive. This is exactly the Schur complement on T of the remaining residue block (E_J, E_Z, T). Hence the whole five-root residue Gram is positive definite. The bounded strict-residue Gram lemma proves the local certificate at $\bar{u} = 0, \bar{v} = \bar{a}$. The Bob sign copy covers $\bar{v} = -\bar{a}$.

All other independent tests. For u infinitesimal and v bounded with $\bar{v} \neq \pm\bar{a}$, retain the original m, n packets. Set $K_J = \sqrt{D}J$ and substitute

$$Y = uW + uvX - vZ - K_J/\sqrt{D}.$$

The exact remainder is a bounded quadratic Gram on (X, Z, K_J, uW). Its hard residue is the block just considered with X pivot $2\bar{d}(1 + \bar{v}^2)$; its soft Schur complement is $\bar{f}(\bar{v}) > 0$. This includes $v = 0$ and all infinitesimal v. For $u, q = 1/v$ infinitesimal use the homogeneous parent $q^2F(u, 1/q)$. Here $J_h = quW + uX - qY - Z$. Retain m, n and substitute $Z = quW + uX - qY - K_J/\sqrt{D}$ with $K_J = \sqrt{D}J_h$. The remainder is a bounded Gram on (X, Y, K_J, uW). Its soft residue is $\lim_{q\to 0} q^2\bar{f}(1/q) > 0$, since $\overline{Q_{12}/L} > 0$ and $\overline{\det B_\infty/L} > 0$. These are polynomial homogeneous expressions, so literal projective endpoints are included.

At every other commuting test the capping proof from the preceding all-tests corollary applies verbatim: choose $\mu = \min(1, C/d, A/d, D/d, \alpha/d)$, a positive unit, retain every excess and the whole m, n packets, and use its displayed open-axis state forms and two corner quadratic forms. That proof used lower bounds, not upper bounds, on these four source weights. Noncommuting residue tests use the retained arbitrary-physical-source theorem. Finally homogenize (TD6)–(TD12) in W, replacing R, T by RW, TW; restore pure testing projective factors and split complex states into real and imaginary parts. This proves the theorem for all states, including $W = 0$, without comparing k, n, D with independent testing infinitesimals.

Exact scope. This is an all-tests theorem for the stated actual-contact domain. It does not bound or remove the residual regimes where C/d or α/d is unbounded, A/d is unbounded, or the actual contact has a different angle-residue type. The C2 reflection $(m, C, D, \alpha; u_c, v_c) \mapsto (D, \alpha, m, C; 1/v_c, 1/u_c)$ supplies its corresponding reflected domain. No exhaustive C2 join follows from these two domains alone.

Chapter 104

Ordinary actual Alice angle, with arbitrary state ratio

Component `p9f00f949fc8c`
Source: `R13/baseline/baseline/baseline/baseline/baseline/baseline/advances/c1/arbitrary_state_ratio.tex`
SHA256: `9f00f949fc8c2e401f57d03bb695867f73c6581778eb8351d4f5dcbdfa388eba`
Inventory use: entire scoped proof | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

104.1 Ordinary actual Alice angle, with arbitrary state ratio

The positive bounded-unit assumption on $s = g/f$ can be removed from the arbitrary-opening C1 theorem. The new ingredient is a different grading of the already proved polynomial contact–conductor identities. No new ideal membership or division is required.

Theorem 104.1. *Use the unchanged native C1 source, normalized by $c = 2$ and $\lambda = n = 0$. Suppose k is a positive bounded unit, $f, g > 0$ are infinitesimal, and*

$$\begin{aligned} &\eta = f^2, \quad \zeta = g^2, \quad K = 1 + k^2, \qquad R = K - ak^2\eta, \quad T = K - k^2 t\zeta, \\ &\qquad a, t > 0, \qquad M = R - t > 0, \qquad J = T - a > 0. \end{aligned} \tag{AS1}$$

Then the full original parent is tensor SOS at every independent projective test and every real or complex state. The ratio $s = g/f$ is arbitrary positive. No positive lower bound on a, t, M, J and no boundedness of original event coefficients is assumed.

The native source and every retained packet are exactly those of (OA2), (AO3)–(AO5), with $s = g/f$ and $\Delta = s/k$. Since $a < T < K$ and $t < R < K$, the shapes are bounded, while R, T are units of residue K. The six coquotients $c_2, c_b, c_3, c_5, c_4, c_\alpha$ have residue $2K$, and c_d has residue $4K^2$. All original events are positive. For the two subtracted expressions, this follows from

$$\frac{D}{a} = \frac{1}{J}\left(\frac{2c_5}{c_3} - J\eta\frac{c_d}{c_3}\right), \qquad \frac{\beta}{a} = \frac{1}{s^2 t}\left(\frac{2k^2 c_b}{c_3} - t\zeta\frac{c_d}{c_3}\right). \tag{AS2}$$

The parentheses have positive unit residues 2 and $2k^2$.

Exact reflection reduces to bounded s. The whole-parent reflection is

$$u' = 1/u, \quad v' = v, \quad (W', X', Y', Z') = (vW - X, -W - vX, Z - vY, Y + vZ), \tag{AS3}$$

with $(\lambda\ n)(m\ c)(A\ \alpha)(D\ \beta)$ on events and

$$u^2 F(\mathrm{ref}(c); 1/u, v; \psi') = (1 + v^2) F(c; u, v; \psi). \tag{AS4}$$

This is an identity of the full parent, including the signed core. At the native contact set $H_f = 1+(1-a)\eta$, $H_g = 1+(1-t)\zeta$, and $V = 1+k^2\eta\zeta$. Both H_f, H_g are positive units. After normalizing the reflected contact by its positive Z' coordinate, the exact reflected native parameters are

$$\begin{aligned} k' &= 1/k, \qquad s' = \frac{H_g}{sH_f}, \qquad \eta' = \frac{k^2\zeta H_f}{H_g}, \qquad \zeta' = \frac{k^2\eta H_g}{H_f}, \\ a' &= \frac{M}{k^2 H_f}, \quad t' = \frac{J}{k^2 H_g}, \qquad M' = \frac{aV}{k^2 H_g}, \quad J' = \frac{tV}{k^2 H_f}. \end{aligned} \tag{AS5}$$

For example, the reflected state has $r' = Q_c/H_g$, $\theta' = \eta J/H_f$, where $Q_c = 1 - k^2\zeta(1-a\eta)$; the differences $H_g - Q_c = \zeta M$ and $H_f - E = \eta J$ prove (AS5). The reflected coefficient vector, after positive normalization to $c' = 2$, is the reflected native source: this follows from the same six contact equations and their nonzero selected minor used for the original native source. All displayed shapes and gaps are positive. Thus unbounded s becomes infinitesimal s', with both η', ζ' infinitesimal and k' a unit. Unit s is the previous arbitrary-opening theorem. It remains to prove the case s infinitesimal.

Outer angles require only a positive current mass. Put $D_1 = 2c_5/c_3 - J\eta c_d/c_3$, a unit of residue 2. For $g_c = cD/(c+D)$ the exact cancellation

$$\frac{g_c}{a} = \frac{2D_1}{2J + aD_1}, \qquad 2J + aD_1 = 2T + a(D_1 - 2) \tag{AS6}$$

gives

$$d/c,\ d/D \text{ infinitesimal}, \qquad \gamma = \frac{Ag_c}{d^2} \text{ a positive bounded unit}, \qquad \bar{\gamma} = \frac{1}{4\bar{K}^3}. \tag{AS7}$$

The two-strong-current identity therefore covers every test with $1/v$ bounded, including Bob infinity, for every Alice angle and state. For v infinitesimal and Alice unbounded, use the exact critical-current identity with residue diagonal

$$\operatorname{diag}(1, 2, 2, 4\bar{\gamma} - \bar{c}_u^2), \qquad c_u = (1+u^2)^{-1/2}. \tag{AS8}$$

Here $\bar{c}_u = 0$ and $\bar{\gamma} > 0$, so it is strict. The former sufficient assumption $\bar{\gamma} \geq 1/4$ is unnecessary on this particular reciprocal-Alice chart. Its same exact identity and homogeneous normalization also include literal Alice infinity. Consequently only bounded u and infinitesimal v remain.

The complementary state chart. Normalize the testing state by a coordinate of maximal absolute value. Put $D_u = 1 + u^2$ and $w = (W + uY)/D_u$, a pure testing coordinate. If w is a unit, the retained-current identity (UW2)–(UW3) applies with $Q = W + uY = D_u w$ a unit. It requires only bounded u, v, Z, not infinitesimal u or unit k. The five roots are

$$r_2,\ B,\ \Delta r,\ \Delta r_5,\ \Delta\eta Y.$$

Their positive diagonal coefficients are $H_2, H_b D_v, H_3 D_v, H_5, 4RTc_4 D_u D_v$, all units. The only two negative corrections have factors $\Delta Z/Q$ and $\Delta^2\eta Z^2/Q^2$, hence zero residue. Restoring the retained packets proves this chart.

Otherwise w is infinitesimal. Literal axes and a nonnegative pure testing core are direct original-event SOS exits. On the negative-core branch, sign gauges give $u, v > 0$. The coordinate Z must be a unit: at v of zero residue the core has residue $2((X+uZ)^2 + (uW - Y)^2)$. If Z, w and these two currents all had zero residue, the normalized state would have every coordinate of zero residue. Thus a negative-core test with small w has unit Z; normalize $Z = 1$ by a pure testing factor. The pure-testing Schur identity now gives $w > 0$; before this normalization its conclusion is $wZ > 0$.

The same conductor, with k of weight zero. In this last chart use exactly (SD3)–(SD8), retaining all four positive packets. Set

$$\delta = \Delta R/K, \qquad b^2 = \eta K/R, \qquad Q = K + ak^2b^2, \quad T_n = K^2 - t\delta^2k^4Qb^2, \quad \mathcal{C} = K^6Q^4T_n.$$

Here δ, b are positive infinitesimals, and k, a, t, u are bounded. The exact source substitution is unchanged:

$$\eta = Kb^2/Q, \quad \Delta = \delta Q/K, \quad \zeta = \delta^2k^2Qb^2/K, \quad R = K^2/Q, \quad T = T_n/K.$$

Let E_0, L_1, L_2 be the scalar correction and two full first normals in the contact–conductor proof. Its polynomial construction places

$$E_0 \in (x, y, z, h, q_1)^2, \qquad L_1, L_2 \in (x, y, z), \tag{AS9}$$

where

$$A_0 = wv, \ B_0 = \delta v, \ C_0 = \delta b^2uw, \ D_0 = \delta^2k^3b^2,$$
$$x = B_0 - C_0, \quad y = A_0 - kC_0, \quad z = uB_0 - kC_0, \quad h = kC_0 - D_0.$$

These are universal polynomial identities; their proof never assumed that k was small or inverted it.

Give $(\delta, k, b, a, t, u, w, v)$ weights $(1, 0, 1, 0, 0, 0, 1, 3)$. All four monomials A_0, B_0, C_0, D_0, and all the soft roots, have weight 4. The substitutions at contact $(u, w, v) = (k, \delta k, \delta k^2b^2)$ preserve this grading. The factored five-term scalar gives

$$\operatorname{wt} E_0 \geq 10, \qquad \operatorname{wt} L_1 \geq 8, \qquad \operatorname{wt} L_2 \geq 6. \tag{AS10}$$

Here is a short way to read these bounds. One has $Q - K = O(2)$, $T_n - K^2 = O(4)$, $V_n - K^2 = O(4)$, $D_v - 1 = O(6)$, $Y_n = O(1)$ and $B_n + KT_n = O(4)$. The six cleared coquotients equal $2K^3Q^3 + O(2)$, while $C_d = 4K^4Q^3 + O(2)$. In the five-term scalar formula, discard terms of weight at least 10; its four weight-eight squares and cross term are precisely $8K^2D_u\mathcal{C}P$. For the normals use

$$K(C_b - C_2) = \delta^2k^2Qb^2C_3, \qquad Q(C_3 - C_5) = -k^2Kb^2C_b,$$

and $S_B, S_C = O(2)$ in their displayed factorizations. This gives (AS10) without an expanded source calculation.

Monic contact division preserves these weights. The conductor lifts therefore have coefficient weights at least $2, 4, 2$, respectively. Every such polynomial coefficient contains a positive power of δ, b, w, or v, and has zero residue. There are no negative exponents, including on a, t; the only source denominators used before this polynomial construction are the displayed positive bounded units.

The leading scalar remains exactly

$$P = D_u(A_0^2 + B_0^2) + K(C_0^2 + D_0^2) - 2K^2B_0C_0 = x^2 + g_4^2 + \sum_{i=1}^{4} a_i^4 - 4\prod_{i=1}^{4} a_i.$$

Here $g_4 = uwv - \delta^2k^4b^2$ and

$$a_1 = \sqrt{wv}, \quad a_2 = \sqrt{\delta uv}, \quad a_3 = b\sqrt{\delta kuw}, \quad a_4 = \delta k^{3/2}b,$$
$$q_1 = a_1a_2 - a_3a_4, \qquad q_2 = a_1a_3 - a_2a_4.$$

Their radicals separate between coefficient and testing fields; $y = a_1^2 - a_3^2$, $z = a_2^2 - a_3^2$, $h = a_3^2 - a_4^2$. The universal two-pairing circuit identity gives a strict constant Gram on (x, y, z, h, q_1, q_2). Retain g_4^2 as a positive square; its coefficients need not have zero residue when k, u are units. The complete hard block on $(N, \delta M_B)$ has positive residue $16\bar{K}^{14}I_2$ after multiplication by $\mathcal{C}$. Indeed $\Delta/\delta = Q/K \sim 1$, $H_i \sim 8K^2$, and the remaining hard-block terms have factors η or v. The full first normals are exactly those in (AS9), not only their restrictions at contact. Thus (AS9)–(AS10) give a bounded strict eight-root remainder Gram. Restore g_4^2 and the four positive packets, then undo the positive pure source multiplier $\mathcal{C}\Delta^2\Lambda$.

Exhaustion and precise advance. The source reflection, unit-s theorem, two outer-angle identities, unit-w current identity, and small-w conductor cover every source and every independent projective test under (AS1). Pure testing normalizations and pure source multipliers are restored separately; no mixed inverse or radical occurs. Complex states split into real and imaginary parts.

Together with the complete infinitesimal-k theorem and the exact orientation $0 < k \leq 1$, this proves the all-tests assertion whenever the actual native entry has both η and ζ infinitesimal, for every positive k in that orientation and every state ratio. The remaining native complement is therefore: k infinitesimal with ζ infinitesimal and η a unit or unbounded; k infinitesimal with ζ non-infinitesimal; and k a unit with at least one of η, ζ non-infinitesimal. Each is still intersected with physicality and the complements of all other inherited admissions and symmetry images. No whole-global completion is asserted.

Chapter 105

A bounded-shape one-weak C2 family, at all tests

Component `pec59340d644f`
Source: `R13/baseline/baseline/baseline/baseline/baseline/baseline/advances/c2/general_single_angle.tex`
SHA256: `ec59340d644fc7d091e3d6cf8f9da2f1a1096c7fb209e9c0a8233c82509a782f`
Inventory use: Complete six-term Laurent identity, full scalar and both normal monic divisions, literal n=0 corner and all four outer-core identities

105.1 A bounded-shape one-weak C2 family, at all tests

This module gives a source-domain extension of the fixed rational family. Its scalar uniqueness proof is an exact six-term Laurent identity and a three-cycle arithmetic–geometric mean inequality. The full-normal and outer-chart arguments include literal $n = 0$.

Put $x > 0$ infinitesimal in the coefficient field and $\rho = \sqrt{x}$. Choose positive bounded coefficient units k, b, r, h such that

$$1 - r, \quad b - h, \quad g = b - 1 + r + xh, \quad b - k^2 r$$

are positive bounded units. Set

$$c = \rho, \quad a = k/\rho, \quad t = x(b-h)/k^2, \quad j = 1 + xr - t, \quad a_2 = k^2/x, \quad A_c = 1 + xr, \quad S_c = A_c + a_2 xt.$$

Thus $h = b - a_2 t$, $g = (1 + x)b - (1 - r + a_2 xt)$, and $j = 1 + xr - t$. Use the exact original rational source

$$\begin{aligned}
\Delta &= (1+x)xb^2 + 2x(1+x)br + (1+x)(1+a_2x)b + x^2(1-a_2)r^2 + 2x(1+a_2x)r + 1 + a_2x,\\
N &= (1+x)(b+r)(b+a_2(1-t)) - a_2(1+a_2x)(1-r)t,\\
n &= \frac{(b+r)A_cS_c - r\Delta}{2r\Delta},\\
m &= \frac{(1+a_2x)(1-r)A_cS_c}{2g\Delta},\\
C &= \frac{A_cN}{2a_2t\Delta},\\
A &= \frac{a_2A_c(b+r)S_c}{2h\Delta},\\
D &= \frac{(1+x)(b+r)(b-a_2xr)S_c}{2a_2j\Delta},\\
\alpha &= \frac{x^2(b+r)N}{2(1-xh)\Delta}.
\end{aligned}$$

Assume $n \geq 0$ and physicality of this canonical source. The latter is the same inherited source-physicality hypothesis used for the general noncommuting-residue exit; no conclusion below is predicated on a new source enumeration. Literal $n = 0$ is allowed.

The parent, with $d = 1$ and $\lambda = \beta = 0$, is

$$\begin{aligned} F &= 4nD_uD_vX^2 + 4mM^2 + 4CD_v(uW - Y)^2 + 4AD_uD_vY^2 + 4DJ^2 + 4\alpha D_u(vY - Z)^2 + H, \\ M &= vW - X + uvY - uZ, \qquad J = uW + uvX - Y - vZ, \\ H/2 &= J^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_uD_vZ^2, \qquad D_u = 1 + u^2, \quad D_v = 1 + v^2. \end{aligned}$$

Its exact stationary contact is

$$(u, v; W, X, Y, Z) = (\rho, k; \rho b/k, -\rho r, xh/k, 1).$$

Writing $\kappa = b/k$ and using a subscript zero for residues gives

$$\begin{aligned} \Delta_0 &= (1 + k_0^2)(b_0 + 1), \qquad n_0 = \frac{b_0 + r_0 - r_0\Delta_0}{2r_0\Delta_0} \geq 0, \\ m_0 &= \frac{1 - r_0}{2(b_0 - 1 + r_0)(b_0 + 1)} > 0, \\ C_* := (xC)_0 &= \frac{k_0^2(b_0 + r_0)}{2(b_0 - h_0)\Delta_0}, \qquad A_* := (xA)_0 = \frac{k_0^2(b_0 + r_0)}{2h_0\Delta_0}, \\ D_* := (D/x)_0 &= \frac{(b_0 + r_0)(b_0 - k_0^2 r_0)}{2k_0^2\Delta_0}, \qquad E_* := (\alpha/x)_0 = \frac{k_0^2(b_0 + r_0)^2}{2\Delta_0}. \end{aligned}$$

All starred quantities are positive. Thus n is bounded, m is a positive unit, C, A have scale x^{-1}, and D, α have scale x.

Theorem 105.1 (General bounded-shape single-angle source)**.** *Every source in the displayed domain has tensor SOS for the full original parent at all independent real or complex projective testing states and all projective testing angles. No comparison between a coefficient infinitesimal and a testing infinitesimal is needed. The conclusion includes $n = 0$. Applying the exact C2 reflection gives the corresponding original orientation with weak C and strong α; this is a sufficient source domain, not a global one-weak C2 classification.*

The finite $u = 0$ axis and its full normals. Set $a_0 = 4m_0$, $b_0' = a_0 + 2$, and

$$H_0(v) = (4n_0 + 2)(1 + v^2) + a_0, \quad \eta = h/b, \quad \gamma = -r - \frac{a_0 b - b_0'}{H_0(k)}.$$

Here γ is infinitesimal and $\eta_0 = C_*/(C_* + A_*)$. Use the exact contact graph

$$X_g = \frac{a_0 vW - b_0' uZ}{H_0(v)} + \gamma uZ, \qquad Y_g = \eta uW.$$

The denominator $H_0(v)$ is a pure testing unit and $H_0(k)$ is a pure coefficient unit. Let $N_X = X - X_g$, $N_Y = (Y - Y_g)/\rho$. Writing $f = F|_g = f_{ww}W^2 + f_{wz}WZ + f_{zz}Z^2$, exact monic divisions give

$$f = aP^2 + b_sB^2 + c_sV^2 + eE^2 + g_sBV, \qquad (P, B, V, E) = (uW/\rho, uZ, vW, \rho Z),$$

where

$$\begin{aligned} a &= x(f_{ww} - f_{ww}|_{u=0})/u^2, \qquad b_s = (f_{zz} - f_{zz}|_{u=0})/u^2, \\ c_s &= f_{ww}|_{u=0}/v^2, \qquad e = f_{zz}|_{u=0}/x, \qquad g_s = f_{wz}/(uv). \end{aligned}$$

These displayed quotients are polynomial divisions, not inverses of testing infinitesimals. Parity gives the divisions by u^2, v^2, uv; $f_{zz}|_{u=0} = 4Dv^2 + 4\alpha$ gives the x factor. Only C, A can produce coefficient poles, and they occur in f_{ww} with a factor u^2; its prefactor x removes them.

Put $Q = 4C_*A_*/(C_* + A_*) > 0$. The bounded coefficient residues are

$$q(v) = a_0^{\mathrm{sc}} = QD_v, \qquad b(v) = b_0' \frac{H_0 - a_0 - 2}{H_0},$$
$$\bar{c}(v) = a_0 \frac{H_0 - a_0}{H_0}, \qquad g(v) = -2b_0' \frac{H_0 - a_0}{H_0}, \qquad e(v) = 4D_* v^2 + 4E_*.$$

Thus $c(v) := v^2 \bar{c}(v)$ and $d(v) := -vg(v)$ give the weighted leading scalar

$$q(v)u^2W^2/x + b(v)u^2Z^2 + c(v)W^2 - d(v)uWZ + e(v)xZ^2.$$

The full expansion is

$$F = f + N_X L_X + N_Y L_Y + H_{XX} N_X^2 + 2H_{XY} N_X N_Y + H_{YY} N_Y^2.$$

Its hard residue is $\mathrm{diag}(H_0(v), 4D_v(C_* + A_*))$. Direct differentiation gives

$$L_X = V\ell_w + B\ell_z, \qquad L_Y = P(B_2 + xe_w) + B(\rho u e_z) + Ee_*,$$

where

$$B_2 = 8D_v\{(xA)\eta D_u - (xC)(1 - \eta)\}, \qquad e_* = 8v(D - \alpha).$$

All ℓ_w, ℓ_z, e_w, e_z are bounded; the residues of ℓ_w, ℓ_z, B_2, e_* vanish identically in finite v. For the second identity, after subtracting uWB_2/x from $F_Y|_g$, the remaining W coefficient is ue_w and the remaining Z coefficient is $u^2 e_z + e_*$. The constant signed-core derivatives cancel. This proves full normal restoration wherever the scalar Gram on these four roots is strict.

Global scalar strictness by an exact cycle identity. Let $\nu = 2n_0 + 1 \geq 1$, $\mu = 2m_0 + 1 > 1$, $z = v^2$, $\mathcal{H} = \nu(1 + z) + \mu - 1$, $e_2 = 4D_*$ and $e_0 = 4E_*$. Then

$$b = \frac{2\mu(\nu(1 + z) - 1)}{\mathcal{H}}, \quad c = \frac{2(\mu - 1)z\nu(1 + z)}{\mathcal{H}}, \quad d = \frac{4\mu v\nu(1 + z)}{\mathcal{H}}, \quad e = e_2 z + e_0.$$

For $s, z > 0$ define $\mathcal{D}(s, z) = (qs + c)(bs + e) - d^2 s/4$. The following exact identity is the key:

$$\begin{aligned}\frac{\mathcal{H}\mathcal{D}(s, z)}{(1 + z)sz} = {} & 2Q\mu(\nu - 1)s/z + 2Q\mu\nu s + 2(\mu - 1)\nu e_2 z/s + 2(\mu - 1)\nu e_0/s \\ & + Qe_0(\nu + \mu - 1)/z + Qe_2\nu z \\ & + Q\{e_2(\nu + \mu - 1) + e_0\nu\} - 4\mu\nu.\end{aligned}$$

Every nonconstant coefficient is positive, except the first may be zero when $n_0 = 0$. Exact contact and stationarity, followed by taking residues, give a stationary zero at $(s, z) = (1, k_0^2)$. Here is an algebraic proof that it is the unique zero on the whole positive quadrant. Rescale $z = k_0^2 t$ and write the Laurent expression as

$$\mathcal{L} = a_\ell s/t + b_\ell s + c_\ell t/s + d_\ell/s + e_\ell/t + f_\ell t - \Sigma, \qquad \Sigma = a_\ell + b_\ell + c_\ell + d_\ell + e_\ell + f_\ell.$$

The constant term has this form by its zero value at $(1, 1)$. Stationarity gives $a_\ell - c_\ell = d_\ell - b_\ell = f_\ell - e_\ell =: \tau$. Put $p_\ell = \min(a_\ell, c_\ell) \geq 0$, $q_\ell = \min(b_\ell, d_\ell) > 0$, $r_\ell = \min(e_\ell, f_\ell) > 0$. Exactly,

$$\begin{aligned}\mathcal{L} = {} & p_\ell(s/t + t/s - 2) + q_\ell(s + 1/s - 2) + r_\ell(t + 1/t - 2) \\ & + \begin{cases}\tau(s/t + 1/s + t - 3), & \tau \geq 0, \\ (-\tau)(t/s + s + 1/t - 3), & \tau \leq 0.\end{cases}\end{aligned}$$

Each bracket is nonnegative by the two- or three-term arithmetic–geometric mean inequality. The two strictly positive pair terms force $s = t = 1$ at equality. They also give a positive definite Hessian there: the other brackets are nonnegative and vanish at that point. Thus both remote uniqueness

and local nondegeneracy are proved by the same finite algebraic identity, over any real closed residue field. No inference from a merely local Hessian is being used.

Consequently, for finite nonzero v with $v^2 \neq k_0^2$,

$$|d(v)| < 2\sqrt{b(v)c(v)} + 2\sqrt{q(v)e(v)}.$$

Indeed the minimum of $\mathcal{D}(s,z)/s$ over $s>0$ is $(\sqrt{bc}+\sqrt{qe})^2 - d^2/4$ and is attained, so positivity is strict. Split the exact cross term using $BV = vPE$ and the pure testing function

$$\theta = \frac{|v|\sqrt{b(v)\bar{c}(v)}}{|v|\sqrt{b(v)\bar{c}(v)} + \sqrt{q(v)e(v)}}.$$

The resulting residue blocks on (B,V) and (P,E) are strictly positive. The denominator is a pure testing unit. At $v_0 = 0$ the same conclusion holds when $n_0 > 0$, with zero cross residues. The case $n_0 = 0, v_0 = 0$ is treated separately below.

The matched finite-angle residue. For $v_0 = k_0$, normalize $Z = 1$ when $Z \neq 0$. If W is infinitesimal, put

$$R = (uW - \kappa x)/\rho, \quad S = u - \kappa^{-1}W, \quad U = \rho(v-k), \quad T = u(v-k).$$

The exact conductors are

$$u^2 - x = uS + \kappa^{-1}\rho R, \qquad W^2 - \kappa^2 x = -\kappa WS + \kappa\rho R, \qquad uW - \kappa x = \rho R.$$

Write $\xi = u^2, \omega = W^2, p = uW$ and

$$f = p^2 Q_b/x + B_b, \qquad Q_b = 4D_v\{(xC)(1-\eta)^2 + (xA)\eta^2 D_u\}.$$

Here B_b is bounded and polynomial in these three invariants with only the separate unit denominators already specified. Put $G = \kappa^2 x Q_b + B_b$ and take successive monic differences

$$G = G_0(v) + (\xi - x)G_\xi + (\omega - \kappa^2 x)G_\omega + (p - \kappa x)G_p.$$

Exactly,

$$f = R^2 Q_b + \kappa^2 S^2 G_\omega + \rho R\mathcal{A} + uS\mathcal{B} + G_0,$$

where $\mathcal{A} = 2\kappa Q_b + \kappa^{-1}G_\xi + \kappa G_\omega + G_p$ and $\mathcal{B} = G_\xi - \kappa^2 G_\omega$. The contact Jacobian of (R,S) with respect to (u,W) has determinant -2. Exact stationarity therefore gives $\mathcal{A}_c = \mathcal{B}_c = 0$. Cancelling ρ here proves vanishing and does not introduce a Gram coefficient inverse. Monic contact division of these two functions and the conductors above give a bounded Gram on (R,S,U,T). Also $G_0 = xg_0(v)$ with bounded g_0, and exact contact gives a double factor $(v-k)^2$, producing its bounded U^2 coefficient.

The full normals have zero-residue lifts on these roots. For L_X, write $L_X = uA_1 + WB_1 = -\kappa B_1 S + u(A_1 + \kappa B_1)$. Both coefficient functions have identically zero leading part for every v; exact normal contact makes $A_1 + \kappa B_1$ vanish at contact. Monic division gives coefficients with factors $u\rho$, u^2, or uW, and a $(v-k)$ quotient of an identically zero leading function. All have zero residue. For L_Y, write $L_Y = RB_2 + \rho E_2$. Its bounded even E_2 vanishes exactly at contact, and both B_2, E_2 have identically zero leading part. Monic division gives factors $x, \rho u, \rho W$, or the same zero-leading-function quotient. Thus the entire normal restoration is bounded and has zero residue.

The soft residue has the form

$$q(k_0)R^2 + \kappa_0^2 c(k_0)S^2 + A_1' RU + B_1' ST + E_1' U^2.$$

It satisfies

$$E_1' > \frac{(A_1')^2}{4q(k_0)} + \frac{(B_1')^2}{4\kappa_0^2 c(k_0)}.$$

To prove this strict inequality, the leading scalar at $u = \rho s, W = \rho w$ is $\Phi = qs^2w^2+bs^2+cw^2-dsw+e$. Complete its w square:

$$\Phi = (qs^2 + c)\left(w - \frac{ds}{2(qs^2 + c)}\right)^2 + \frac{\mathcal{D}(s^2, v^2)}{qs^2 + c}.$$

The global Laurent calculation gives a positive definite Hessian of the second term in (s, v) at $(1, k_0)$; both coordinate derivatives are nonzero there. The first square supplies the missing w direction. Hence the full three-variable Hessian is positive definite. On first variations $T = U$; the displayed inequality is exactly its Schur complement after the invertible (R, S) change. Choose a real δ strictly between the second summand and $E_1' - (A_1')^2/(4q(k_0))$. Add δ times the exact zero

$$T^2 - U^2 - ST(v - k) - \kappa^{-1}RU(v - k) = 0.$$

The two blocks (R, U) and (S, T) are now strictly positive, and so is the full six-root residue Gram.

If the ratio W/Z is not infinitesimal, instead normalize $W = 1$; then Z is bounded. The earlier scalar is aP^2 plus a term of positive residue $c(k_0)$; its normal lifts have zero residue on $(P, 1)$. This is strict. The case $Z = 0$ is a literal sum of the event squares and the three positive core squares. The simultaneous sign change $(u, v, X, Y) \mapsto (-u, -v, -X, -Y)$ covers $v_0 = -k_0$.

The double-zero corner when $n_0 = 0$. Use instead the pure coefficient quantities

$$H_c = 4n + 4m + 2, \quad \xi_c = 4m/H_c, \quad \zeta = (4m + 2)/H_c, \quad \eta_c = C/(C + A),$$

and graph $X_g = \xi_c vW - \zeta uZ$, $Y_g = \eta_c uW$. Put $\epsilon = 4\alpha + 4n(4m+2)/H_c > 0$. Then ϵ is infinitesimal, while n/ϵ and x/ϵ are bounded. Use the hard roots as before and scalar roots

$$P = uW/\rho, \quad V = vW, \quad E = \rho Z, \quad B_\epsilon = \sqrt{\epsilon}\, uZ, \quad T = uvZ.$$

Direct packet expansion gives exactly

$$f_{zz} = 4\alpha + \epsilon u^2 + 4n\zeta^2u^4 + 4Dv^2 + u^2v^2B_b,$$

where B_b is bounded with residue 2. Its first four terms give coefficients $4\alpha/x$, $1 + (4n\zeta^2/\epsilon)u^2$, and $(4D/x)v^2$ on E^2, B_ϵ^2, E^2 respectively. The W^2 contribution has bounded coefficients on P^2, V^2 with positive residues Q and $4m_0/(2m_0 + 1)$. The WZ contribution is $uvWZ$ times a bounded function; use $uvWZ = vPE$ to give it zero cross residue. Thus the five scalar roots have strict diagonal residue $(Q, 4m_0/(2m_0 + 1), 4E_*, 1, 2)$.

For completeness, L_X has a zero-residue bounded coefficient on V. At $v = 0$ its Z coefficient is exactly $-8n\zeta u^3$; on B_ϵ this is $-8n\zeta u^2/\sqrt{\epsilon}$, which is infinitesimal because n/ϵ is bounded. The remaining Z coefficient has a factor uv^2, giving a zero-residue multiple of T. For L_Y, the large-packet coefficient is now exactly $B_2 = 8D_v(xA)\eta_c u^2$; the remaining terms lift as

$$L_Y = P(B_2 + xe_w) + T(\rho u e_z) + E\, 8v(D - \alpha),$$

with bounded e_w, e_z. All coefficients have zero residue. The hard residue is $\mathrm{diag}(4m_0+2, 4(C_* + A_*))$. This includes literal $n = 0$ without deleting a coordinate.

The remaining commuting charts. The following exact core identities retain all unused event packets. Every retained coefficient is nonnegative; in particular the n packet can be retained even when $n = 0$. They prove strict bounded Grams without dividing by a testing infinitesimal.

If u is a finite unit and $v \to 0$, take

$$R = (uW - Y)/\rho, \quad L = Y/\rho, \quad B = X + uZ, \quad V = vZ.$$

Then $M = -B + (\rho v/u)(R + D_u L)$, and exactly

$$H/2 = (1 + D_u v^2)B^2 + u^2 D_u V^2 - 2uvD_u BV$$

$$+ 2\rho R(uvB - D_u V) + xR^2 - 2D_u \rho LV + D_u x v^2 L^2.$$

Retain n, D, α. The residue on (B, V, R, L) is diagonal with entries $(4m_0 + 2, 2u_0^2 D_{u,0}, 4C_*, 4A_* D_{u,0})$.

If $u \to \infty$ and v is finite, put $p = 1/u$ and use $p^2 F$. Take $R = (W - pY)/\rho$, $L = Y/\rho$, $B = Z + pX$, $V = vX$. The homogenized m amplitude is $-B + \rho pvR + \rho v D_p L$, and the exact core is

$$\begin{aligned}\widehat{H}/2 = B^2 + D_p V^2 + 2\rho D_p RV - 2\rho pvRB + xR^2 \\ + 2D_p \rho pLV - 2D_p \rho vLB + D_p x v^2 L^2.\end{aligned}$$

Retain n, D, α. The residue on (B, V, R, L) is diagonal with entries $(4m_0 + 2, 2, 4C_* D_{v,0}, 4A_* D_{v,0})$.

If $v \to \infty$ and u is bounded, put $q = 1/v$ and use $q^2 F$. Take $R = (uW - Y)/\rho$, $L = Y/\rho$, $T = uqZ$, $N_1 = \rho Z$. Exactly,

$$\begin{aligned}\widehat{H}/2 &= (D_u + q^2)X^2 + D_u x L^2 + T^2 + 2qXT + q^2 xR^2 + 2uq\rho XR - 2qRN_1 - 2D_u qLN_1, \\ 4D\widehat{J}^2 &= 4(D/x)(-N_1 + \rho uX + xqR)^2.\end{aligned}$$

Retain n, m, α. The residue on (X, R, L, T, N_1) is diagonal with entries

$$(2D_{u,0}, 4C_*, 4A_* D_{u,0}, 2, 4D_*).$$

Finally if both angles are infinite, use $p = 1/u, q = 1/v$ and $p^2 q^2 F$. Set $R = (W - pY)/\rho$, $L = Y/\rho$, $T = qZ$. Then

$$\widehat{H}/2 = (1 + p^2 D_q)X^2 + xq^2 R^2 + xD_p L^2 + T^2 + 2\rho qXR + 2pqXT - 2\rho pRT - 2\rho D_p LT.$$

Retain n, m, D, α. The residue on (X, R, L, T) is diagonal with entries $(2, 4C_*, 4A_*, 2)$.

Exhaustion and independence. These charts cover every commuting angle residue: the whole finite $u = 0$ axis, the finite-unit $v = 0$ axis, either infinite axis, and their intersections. When both angle residues are noncommuting, apply the inherited physical noncommuting-residue theorem at arbitrary coefficient scales. Complex states require no additional chart: if $Z \neq 0$, multiply by a phase and normalize $Z = 1$; the real parent is the sum of its real-state value and its imaginary-state value, the latter having $Z = 0$ and hence being a literal square sum. If $Z = 0$ this is already true of both parts. Homogeneity covers all projective state charts. Roots may be unbounded throughout. All inverses or radicals used in coefficients are pure coefficient or pure testing operations on units, except $\sqrt{\epsilon}$ in roots and its controlled coefficient ratio $n/\sqrt{\epsilon}$. No mixed inverse and no comparison of independent infinitesimals occurs. Applying the strict bounded-tensor Gram lemma completes the proof.

Chapter 106

Ordinary-angle C1 with arbitrary positive shape openings

Component p64b054e815d0
Source: R13/baseline/baseline/baseline/baseline/baseline/baseline/baseline/advances/c1/arbitrary_shape_openings.tex
SHA256: 64b054e815d0c2e05a4f9af55664795c46fdfd4b528544b74a0add8fb3953946
Inventory use: AO1–AO12 | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

106.1 Ordinary-angle C1 with arbitrary positive shape openings

The unit assumptions on a, t, M, J in Theorem 108.1 can all be removed. The large positive packets must be retained before taking a bounded Gram residue.

Theorem 106.1 (Ordinary angles and arbitrary positive openings)**.** *Use the unchanged native C1 source (OA2), with $c = 2$ and $\lambda = n = 0$. Assume only that $k, s = g/f$ are positive bounded units, $\eta = f^2 > 0$ is infinitesimal, and*

$$\begin{gathered}\zeta = g^2 = s^2\eta, \quad K = 1 + k^2, \quad R = K - ak^2\eta, \quad T = K - k^2s^2\eta t, \\ a, t > 0, \quad M = R - t > 0, \quad J = T - a > 0.\end{gathered} \tag{AO1}$$

Then the full original parent is tensor SOS at every independent projective test and every real or complex state. No lower bound on a, t, M, J, and no boundedness assumption on the original event coefficients, is required.

Since $a < T < K$ and $t < R < K$, both shapes are bounded, and R, T have residue K. In particular

$$p = t/R \in (0, 1), \qquad h = a/T \in (0, 1) \tag{AO2}$$

are bounded at every opening rate. The seven native coquotients have the same residues as in (OA3). All original events are positive: A/a and α/a are η^2 times positive units, while d/a is η times a positive unit; positivity of the remaining events also follows from the explicit bounds below.

Retain the four large packets exactly. Put $\Delta = s/k$, $\Lambda = RTc_3/(2a)$, and

$$H_2 = 4Tc_2, \quad H_b = 4Tc_b, \quad H_3 = 4Rc_3, \quad H_5 = 4Rc_5.$$

All four have residue $8K^2$. For a bounded finite testing angle define

$$D_u = 1 + u^2, \quad D_v = 1 + v^2, \quad x = X + uZ, \quad y = uW - Y, \quad q_A = v(W + uY), \quad q_B = v(Z - uX),$$

$$N = x - pq_A, \qquad L = y - hq_B.$$

The four nonnegative packets are

$$\Pi_1 = H_2\frac{t}{M}N^2, \quad \Pi_2 = H_b\frac{M}{t}D_vN^2, \quad \Pi_3 = \Delta^2H_3\frac{J}{a}D_vL^2, \quad \Pi_4 = \Delta^2H_5\frac{a}{J}L^2. \tag{AO3}$$

Their possibly unbounded pure source coefficients stay in their separate square roots. They never enter the bounded remainder Gram. The exact whole-parent identity is

$$\begin{aligned} G := \Delta^2\Lambda F - \sum_{i=1}^{4}\Pi_i &= H_2\{N^2 - 2q_AN + (1-p)q_A^2\} + H_bD_v\{N^2 + 2q_AN + pq_A^2\} \\ &\quad + \Delta^2H_3D_v\{L^2 + 2q_BL + hq_B^2\} + \Delta^2H_5\{L^2 - 2q_BL + (1-h)q_B^2\} + \mathcal{W}, \\ \mathcal{W} &= 4RT\Delta^2\eta^2c_4D_uD_vY^2 + 4RT\Delta^4k^6\eta^2c_\alpha D_u(vY-Z)^2 \\ &\quad + 2RT\Delta^2\eta c_dD_u\{(vY-Z)^2 - D_vZ^2\}. \end{aligned} \tag{AO4}$$

This is the original four-packet identity (SD3)–(SD4), valid without unit shapes. For example, the first subtraction uses only $R = M + t$ and $x = N + (t/R)q_A$; its result is $H_2\{N^2 - 2q_AN + (M/R)q_A^2\}$. The other three follow in the same way from $R = M + t$ and $T = J + a$.

A bounded coordinate change with no mixed inverse. Let $w = (W + uY)/D_u$, which lies entirely in the testing field. Then

$$\begin{gathered} q_A = D_uvw, \quad X = N + pD_uvw - uZ, \quad q_B = v(D_uZ - uN - puD_uvw), \\ W = w + \frac{u(L + hq_B)}{D_u}, \qquad Y = uw - \frac{L + hq_B}{D_u}. \end{gathered} \tag{AO5}$$

These are exact inverse formulas. Their only testing denominator is the positive pure testing unit D_u; their source coefficients use only R^{-1}, T^{-1}. In particular no determinant involving p, h, u, v is inverted. Set $\widehat{G} = D_uG$. It is a polynomial in N, L, u, w, v, Z with bounded coefficients, since this multiplication clears the denominators in (AO5).

Suppose v is infinitesimal and u is a finite positive unit. The exact polynomial $\widehat{G}$ has a bounded Gram on

$$N,\ L,\ vw,\ vZ,\ \eta w,\ \eta Z. \tag{AO6}$$

After expanding only the squares in (AO4), every term contains a product of two listed roots. The coefficient errors carry v, η, or a native coquotient difference of zero residue. To display the Gram residue, use the purely formal substitutions $N = \eta\xi$, $L = \eta\chi$, $v = \eta\nu$ and read the coefficient of η^2. It is

$$\mathcal{G} = 16K^2D_u\xi^2 + 16\Delta^2K^2D_u\chi^2 + 8\Delta^2K^3D_u^2S, \tag{AO7}$$

where

$$S = u^2w^2 + s^2k^4Z^2 + \frac{D_u}{K}\nu^2\left(\frac{k^2}{s^2}w^2 + Z^2\right) - 2Ku\nu wZ. \tag{AO8}$$

There is no shape parameter in this residue other than the fixed units k, s. In particular no assertion that v/η is bounded has been made. Formula (AO7) follows directly from $H_2, H_b, H_3, H_5 \sim 8K^2$, $c_4, c_\alpha \sim 2K$, $c_d \sim 4K^2$: the p and h terms in the two scalar pairs sum to one, while the two linear normal coefficients have zero residue.

Finite unit Alice angles: the same scalar, fewer hard roots. The capacity identity (OA10), with W replaced by w, proves a strict six-root Gram unless $\bar{u} = \bar{k}$. At the matched angle,

$$S = (\nu Z - kw)^2 + \left(\frac{k}{s}\nu w - sk^2Z\right)^2.$$

Define

$$r_1 = vZ - \eta kw, \qquad r_2 = (k/s)vw - \eta sk^2Z.$$

The exact relation

$$wr_1 - (s/k)Zr_2 = \eta k(s^2Z^2 - w^2) \tag{AO9}$$

closes every state with $\overline{s^2Z^2 - w^2} \neq 0$ by adding its identically zero squared relation to the Gram on (N, L, r_1, r_2, η). The multiplier in this relation is never inverted.

Normalize the testing state. If N or L has nonzero residue, $\widehat{G}$ has positive residue and is already certified. Otherwise the inverse formulas force w or Z to be a unit. When (AO9) has zero multiplier residue, Z is a unit; normalize $Z = 1$. The negative-core branch has $w > 0$ by the pure testing core Schur calculation printed in the ordinary-angle proof. Thus only $\bar{w} = \bar{s}$ remains. The nonnegative-core branch and literal axes are direct original-event SOS exits.

The exact contact with arbitrary shape rates. The native contact (OA4), under (AO5), is

$$(u_c, w_c, v_c, N_c, L_c) = \left(k, \frac{sR}{K}, ks\eta, 0, 0\right). \tag{AO10}$$

All four packets (AO3) and all their first derivatives vanish there. The original native contact identities consequently give exact value and first-derivative zero for $\widehat{G}$ as well. This remains true when a packet coefficient is unbounded: each coefficient is an element of the source field multiplying a square that vanishes exactly.

Use the five roots

$$N,\ L,\ v - v_c,\ \eta(w - w_c),\ \eta(u - k). \tag{AO11}$$

The complete polynomial $\widehat{G}$ belongs to $(N, L, v)^2 + \eta(N, L, v) + (\eta^2)$, by (AO4)–(AO5). Translation by $v_c = \eta ks$ preserves this inclusion. Its constant normal coefficient is η^2 times a bounded polynomial in the two soft deviations; its linear normal coefficients are η times such polynomials. Exact contact and stationarity make their constant and the required first coefficients vanish. Finite monic Taylor division therefore gives an exact bounded Gram on (AO11), just as in (OA14), with no division by any testing/source comparison. Higher Taylor terms have zero residue after this allocation.

The residue Gram is now particularly simple. Its first two blocks are $16K^3$ and $16\Delta^2K^3$. Its remaining block is $8\Delta^2K^5$ times the half-Hessian of S at $(\nu, w, u) = (ks, s, k)$ and $Z = 1$. The latter is exactly (OA15), with positive leading principal minors

$$K, \qquad 4k^4, \qquad 12s^2k^6/K.$$

Thus the entire five-root Gram is strict independently of all four shape-opening rates.

The outer charts without bounded event weights. These charts admit a simpler uniform retained-packet comparison. The exact source formulas give

$$\begin{aligned} m/a &\geq \frac{2k^2c_2}{s^2Rc_3}, & c/a &\geq \frac{2}{T}, \\ D/a &\geq \frac{2c_5}{Tc_3} - \frac{\eta c_d}{c_3}, & \beta/a &\geq \frac{2k^2c_b}{s^2Rc_3} - \frac{\eta c_d}{c_3}. \end{aligned} \tag{AO12}$$

All four right sides are positive bounded units. Choose a positive real μ smaller than their residues. In F/a, retain the four nonnegative event packets with weights $m/a - \mu, c/a - \mu, D/a - \mu, \beta/a - \mu$. The remaining complete parent has the four corresponding weights all equal to μ, and the unchanged weights

$$d/a = \eta\, d_1, \quad A/a = \eta^2 A_2, \quad \alpha/a = \eta^2\alpha_2,$$

where d_1, A_2, α_2 are positive units. This is an exact decomposition at the present test, not a claim that the capped source is globally physical.

If the homogeneous Bob sine has nonzero residue, the four event amplitudes have trivial common state kernel, as proved in the ordinary-angle theorem. Their bounded positive Gram absorbs the infinitesimal signed core. If Bob is small and u is infinitesimal, use the six roots $(x, y, vW, vZ, \eta uW, \eta Z)$

of the same theorem. The four hard weights μ give a strict four-root block and the two weak diagonal coefficients are positive units. The signed soft cross term has an additional factor u, so its Gram coefficient has zero residue. At the reciprocal Alice endpoint put $q = 1/u$ and use

$$x' = qX + Z,\ y' = W - qY,\ vY,\ vX, \eta Y,\ \eta qX.$$

The hard residue is $4\mu\{(vY - x')^2 + x'^2 + y'^2 + (y' + vX)^2\}$; the weak diagonals have positive unit coefficients. Assign the signed soft cross term as a multiple of $q(vX)(\eta Y)$, which has zero residue. These are the exact finite and reciprocal monomial allocations already proved in the ordinary-angle theorem; they use no shape inverse. Literal infinity follows in the homogeneous chart.

These outer charts and the finite-unit Alice analysis exhaust all independent tests. Restore the retained packets, then the positive pure factors D_u, $\Delta^2\Lambda$, or a used in the respective charts. Every radical and inverse is confined to one field. Complex states split into their real and imaginary parts.

Remaining scope. This closes every positive opening a, t, M, J on the ordinary-unit-k, ordinary-unit-$s = g/f$, infinitesimal-η row. It does not address nonunit k or s, non-small η, or literal zero openings that are not defined by the native formulas. Actual-source entry retains exactly the inherited contact-sign and nonzero-minor domain of (CE1).

Chapter 107

C1 native formulas and exact stationarity

Component `pfecc204a8025`
Source: `CORE/c1/native_formulas.tex`
SHA256: `fecc204a8025e222eae91494308071a0997c95be28e9f123893b632a9f807c0f`
Inventory use: All positive native parameters: exact seven-coquotient identities and six contact jets

107.1 C1 native formulas and exact stationarity

These are polynomial identities for arbitrary positive k, f, g; the small-k assumptions of the earlier conductor application are not needed for these identities. For the ordinary-angle theorem set $f^2 = \eta$, $g^2 = \zeta = s^2\eta$, and $\Delta = s/k$. Write $K = 1 + k^2$, $R = K - ak^2\eta$, $T = K - k^2\zeta t$, $M = R - t$, and $J = T - a$.

$$\kappa = k^2, \quad \eta = f^2, \quad \zeta = g^2, \quad V_c = \kappa\eta\zeta, \quad L_c = 1 - a\eta, \quad H_c = 1 + (1-a)\eta, \quad Q_c = 1 - \kappa\zeta + aV_c.$$

Define

$$\begin{aligned}
P_4 &= 1 + \kappa + \eta\kappa(a - \kappa - 1) - \zeta\kappa t + V_c[-\zeta\kappa t^2 + 2\kappa t + a(1-t) + t],\\
P_\alpha &= 1 + \kappa - a\eta\kappa + \zeta[(1+\kappa) - (2+\kappa)t] + V_c[-\zeta t^2 + a(t-1) + t],\\
P_d &= (1+\kappa)^2 - a\eta\kappa^2 + \zeta\kappa[(\kappa+1) - (\kappa+2)t]\\
&\quad + V_c[-\zeta\kappa t^2 + \kappa a(t-1) + \kappa t + \kappa + 1].
\end{aligned}$$

The complete seven coquotients are

$$\begin{aligned}
c_2 &= 2(1+\kappa)(1-t\zeta)L_c(1+V_c)H_cQ_c,\\
c_b &= 2(1+\kappa)L_c[1 + (1-t)\zeta]H_cQ_c,\\
c_3 &= 2(1+\kappa)L_cH_cQ_c(1 - \kappa\eta + tV_c),\\
c_5 &= 2(1+\kappa)L_cH_cQ_c(1+V_c),\\
c_4 &= 2Q_cP_4, \qquad c_\alpha = 2L_cH_cP_\alpha, \qquad c_d = 4L_cQ_cP_d.
\end{aligned}$$

Use the source coefficients (OA2), and let $\Lambda = RTc_3/(2a)$, $\Phi = \Delta^2\Lambda F$. For the following verification only, write $r = uW - Y$, $B = X + uZ$, $r_2 = vW - X + uvY - uZ$, $r_5 = uW + uvX - Y - vZ$, and $D = 1 + u^2$. The symbol r in this paragraph is not the C2 source gap.

Short native proof of the exact contact identities. Write $V = 1 + V_c$ and $K = 1 + \kappa$ in this paragraph. The factored coquotients above give immediately

$$c_b - c_2 = \zeta c_3, \qquad c_3 - c_5 = -\kappa\eta c_b, \qquad c_2 = (1 - t\zeta)c_5. \tag{SD9}$$

Hence $Vc_b - c_2 = \zeta c_5$ and $Vc_3 - c_5 = -\kappa\eta c_2$. The remaining three balances are the following short polynomial identities; substituting the displayed P_4, P_α, P_d proves them by multiplication and collecting terms:

$$\begin{aligned}
\mathfrak{Y} &:= 2\kappa c_2 + 2VL_c c_4 - 2\kappa^2\zeta Q_c c_\alpha - Q_c c_d = 0,\\
\mathfrak{S} &:= \kappa L_c c_2 + c_5 + VL_c^2 c_4 + \kappa Q_c^2 c_\alpha + \tfrac{1}{2}(\kappa\zeta L_c^2 - 2L_c - \eta)c_d = 0,\\
\mathfrak{V} &:= \kappa R c_2 + t\kappa V_c c_b + aV_c c_3 + Tc_5 + KV_c L_c^2 c_4 - \kappa^2\zeta K Q_c L_c c_\alpha - \tfrac{K}{2}(Q_c L_c + \eta)c_d = 0. \qquad \text{(SD10)}
\end{aligned}$$

For example, after the common factor $4L_cQ_c$ is removed, the first balance is just

$$\kappa K(1 - t\zeta)VH_c + VP_4 - \kappa^2\zeta H_c P_\alpha - Q_c P_d = 0.$$

The two other balances are equally direct identities in the same three short polynomials; no expanded scalar or ideal calculation enters.

The exact contact, in the original state coordinates, is

$$u = k, \quad v = \Delta k^2\eta, \qquad (W, X, Y, Z) = (\Delta k, -k(1 - t\zeta), \Delta k^2 L_c, 1). \qquad \text{(SD11)}$$

At this point

$$r_2 = Mk\zeta, \quad B = tk\zeta, \quad r = av, \quad r_5 = -Jv, \quad vY - 1 = -Q_c, \quad D = K, \quad D_v = V.$$

Set $\Psi = \Phi/(4RT)$ and let $E_s = \frac{1}{2}\partial_s\Psi$, with all other original testing coordinates held fixed. Direct differentiation of the seven terms of the full parent (OA5), followed by (SD9), gives at (SD11)

$$\begin{aligned}
\Psi &= K\zeta^2\mathfrak{S},\\
E_W &= kv\{\zeta c_2 + \Delta^2(Vc_3 - c_5)\} = 0,\\
E_X &= k\zeta\{Vc_b - c_2 - \zeta c_5\} = 0,\\
E_Y &= \tfrac{1}{2}K\Delta^3\kappa\eta^2\mathfrak{Y},\\
E_u &= k\zeta^2\{\mathfrak{S} - c_2 + (1 - t\zeta)c_5\},\\
E_v &= v\Delta^2\mathfrak{V}.
\end{aligned}$$

They all vanish by (SD9)–(SD10). Homogeneity in (W, X, Y, Z) also gives $E_Z = 0$. This proves contact and full first-order stationarity directly in the native source.

Chapter 108

An ordinary-angle C1 family: all independent tests

Component p18a57c9c271d
Source: CORE/c1/ordinary_angle.tex
SHA256: 18a57c9c271dac00301b509f8f383ecde364a8f9a07a259928042e36892bbc22
Inventory use: OA2–OA16 and outer-angle proof; no need to import its stricter shape hypotheses into AO | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

108.1 An ordinary-angle C1 family: all independent tests

This section closes an actual source family outside the small-k uniform theorem. All inverses below are either real constants or bounded units in one of the two independent real closed fields. In particular no quotient of a testing infinitesimal by a source infinitesimal is used.

Theorem 108.1 (Regular ordinary-angle C1 source)**.** *Use the seven native C1 coquotients and original coefficients of the source-definition appendix, with $c = 2$ and $\lambda = n = 0$. Suppose*

$$k, s, a, t, M, J \text{ are positive bounded units}, \qquad \eta > 0 \text{ is infinitesimal}, \qquad f^2 = \eta, \quad g^2 = s^2\eta,$$

$$K = 1 + k^2, \quad R = K - ak^2\eta, \quad T = K - k^2s^2\eta t, \quad M = R - t > 0, \quad J = T - a > 0. \tag{OA1}$$

Then the full original parent is tensor SOS at every independent projective test, for every real or complex state.

Here $s = g/f$ is the actual contact state ratio. Its being a unit is part of the hypothesis. The source normalization and all six original retained event coefficients are preserved. In particular this theorem includes $k = s = 1$, $a = t = 1/2$, and arbitrary positive infinitesimal η.

Exact source and its leading scales. Put $e = k/s$ and $\Delta = s/k$ in the native formulas. To fix completely which source is meant, they are

$$\begin{gathered} d = \frac{2a\eta c_d}{c_3}, \quad m = \frac{2ak^2c_2}{s^2Mc_3}, \quad A = \frac{2a\eta^2c_4}{c_3}, \quad \alpha = \frac{2as^2k^4\eta^2c_\alpha}{c_3}, \\ D + \frac{d}{2} = \frac{2ac_5}{Jc_3}, \qquad \beta + \frac{d}{2} = \frac{2ak^2c_b}{s^2tc_3}, \qquad c = 2. \end{gathered} \tag{OA2}$$

The letter D here denotes the fifth event, whereas $D_u = 1 + u^2$. Setting $\eta = \zeta = 0$ in the short native polynomials gives $c_2 = c_b = c_3 = c_5 = c_4 = c_\alpha = 2K$ and $c_d = 4K^2$. Consequently the following positive bounded source expressions have the same residues as the indicated original coefficients:

$$\begin{gathered} m_0 = \frac{2ak^2}{s^2(K-t)}, \quad c_0 = 2, \quad D_0 = \frac{2a}{K-a}, \quad \beta_0 = \frac{2ak^2}{s^2t}, \\ d/\eta \sim 4aK, \qquad A/\eta^2 \sim 2a, \qquad \alpha/\eta^2 \sim 2as^2k^4. \end{gathered} \tag{OA3}$$

The symbol $\sim$ here means equality of residues, not equality of the source. Each displayed inverse is a pure source unit inverse. All original events are positive under (OA1), as follows from their positive leading coefficients.

The exact contact is

$$u_c = k, \quad v_c = ks\eta, \quad (W_c, X_c, Y_c, Z_c) = (s, -k(1 - ts^2\eta), ks(1 - a\eta), 1). \tag{OA4}$$

Its value and all first derivatives vanish. This assertion uses the native identities (SD9)–(SD11), which are polynomial identities valid for arbitrary k, not an asymptotic extension of a small-k assertion. The accompanying small checker repeats all six native balances. For actual canonical C1 contacts, the source transport and nonzero selected contact minor in (CE1)–(CE2) identify (OA2) with the original source; no new source-entry assumption is implicit here.

A four-square reduction. Write

$$x = X + uZ, \quad y = uW - Y, \quad D_u = 1 + u^2, \quad D_v = 1 + v^2.$$

The exact complete parent, including every signed term, is

$$\begin{aligned} F = {} & 4m\{v(D_uW - uy) - x\}^2 + 4cD_vy^2 + 4AD_uD_v(uW - y)^2 \\ & + (4D + 2d)\{y + v(ux - D_uZ)\}^2 + 4\alpha D_u\{v(uW - y) - Z\}^2 \\ & + (4\beta + 2d)D_vx^2 + 2dD_u(\{v(uW - y) - Z\}^2 - D_vZ^2). \end{aligned} \tag{OA5}$$

The last line includes the essential $2dD_u(vY - Z)^2$ term. No event is capped or discarded.

For a moment use formal variables ξ, χ, ν and substitute $x = \eta\xi$, $y = \eta\chi$, $v = \eta\nu$ into (OA5). The coefficient of η^2 is

$$\begin{aligned} \mathcal{F} = {} & 4m_0(\nu D_uW - \xi)^2 + 4\beta_0\xi^2 + 4c_0\chi^2 + 4D_0(\chi - \nu D_uZ)^2 \\ & + 8aD_uu^2W^2 + 8as^2k^4D_uZ^2 - 16aKD_uu\nu WZ. \end{aligned} \tag{OA6}$$

This is only a polynomial way to read a bounded Gram residue; it does not substitute or compare any actual testing and coefficient rates. Completing the first four squares gives

$$\mathcal{F} = 4(m_0 + \beta_0)\left(\xi - \frac{t}{K}\nu D_uW\right)^2 + 4(c_0 + D_0)\left(\chi - \frac{a}{K}\nu D_uZ\right)^2 + 8aD_uS, \tag{OA7}$$

where

$$S = u^2W^2 + s^2k^4Z^2 + \frac{D_u}{K}\nu^2\left(\frac{k^2}{s^2}W^2 + Z^2\right) - 2Ku\nu WZ. \tag{OA8}$$

For bounded u, v with v infinitesimal, (OA5) is an exact bounded quadratic Gram on

$$x,\ y,\ vW,\ vZ,\ \eta W,\ \eta Z. \tag{OA9}$$

To see the assertion directly, expand only the displayed squares in (OA5). Each monomial contains a product of two listed roots; all remaining factors are bounded. Terms omitted in (OA6), after this allocation, carry v, η, or a source coefficient difference of zero residue. Hence its Gram residue is exactly (OA6)–(OA8), with $\eta\nu$ read as v. This observation also applies to the modified endpoint root lists below.

Angles away from the actual contact. Sign gauges allow $u, v \geq 0$. Suppose first that u has positive finite residue. In the four roots $\nu W, \nu Z, W, Z$, split the cross term in (OA8) between the pairs $(\nu W, Z)$ and $(\nu Z, W)$, using their identical products. The two positive cross-term capacities are

$$\sqrt{D_u/K}\,k^3, \qquad \sqrt{D_u/K}\,u.$$

Their sum is strictly larger than Ku unless $\bar{u} = \bar{k}$, since the exact factorization is

$$(1+u^2)(u+k^3)^2 - K^3u^2 = (u-k)^2(k^4 + 2k^3u + 2ku + u^2). \tag{OA10}$$

Thus, when $\bar{u} \neq \bar{k}$, choose a real splitting constant strictly within the two residue capacities. The resulting two two-by-two blocks, together with the two completed hard squares, give a strict residue Gram on (OA9).

If u is infinitesimal, replace ηW in (OA9) by $\eta u W$. Assign the entire cross term to $(vW, \eta Z)$; its coefficient has zero residue. The four soft diagonal coefficients are positive units. The same monomial allocation in (OA5) proves an exact strict Gram.

The reciprocal Alice endpoint requires no source reflection. Put $h = 1/u$, multiply the homogeneous parent by the pure testing square h^2, and use $x' = hX + Z$, $y' = W - hY$. Its six roots are

$$x',\ y',\ vY,\ vX,\ \eta Y,\ \eta hX. \tag{OA11}$$

When h is infinitesimal, the four strong residue terms are $4m_0(vY - x')^2 + 4\beta_0 x'^2 + 4c_0 y'^2 + 4D_0(y' + vX)^2$. The two weak diagonal reserves are $8a(\eta Y)^2$ and $8as^2k^4(\eta hX)^2$. The signed cross term is assigned as a multiple of $h(vX)(\eta Y)$, so has zero residue; all other corrections also have zero residue. This is a strict six-root Gram, including the literal endpoint $h = 0$ by the homogeneous formula.

Matched angle but unmatched state. It remains to consider $\bar{u} = \bar{k}$. At $u = k$, (OA8) factors as

$$S = (\nu Z - kW)^2 + \left(\frac{k}{s}\nu W - sk^2 Z\right)^2. \tag{OA12}$$

Use the exact roots

$$r_1 = vZ - \eta kW, \qquad r_2 = \frac{k}{s}vW - \eta sk^2 Z,$$

together with the two completed hard roots $h_1 = x - (t/K)D_u vW$, $h_2 = y - (a/K)D_u vZ$, and η. All six roots in (OA9) are bounded linear combinations of these five. The exact relation

$$Wr_1 - \frac{s}{k}Zr_2 = \eta k(s^2Z^2 - W^2) = p\eta \tag{OA13}$$

shows that, if $\bar{p} \neq 0$, adding the identically zero square $(p\eta - Wr_1 + (s/k)Zr_2)^2$ makes the five-root Gram strict. This operation never inverts p.

After pure testing state normalization, any state with positive parent residue already has a one-root certificate. Otherwise the four strong events force $\bar{x} = \bar{y} = 0$, so W or Z is a unit. If $\bar{p} = 0$, then Z is a unit and we may put $Z = 1$. The remaining choices are $\bar{W} = \pm\bar{s}$. The negative choice is a direct nonnegative-core exit. Indeed set $w = (W + uY)/D_u = W - uy/D_u$. The pure testing core Schur calculation has hard block

$$\begin{pmatrix} 1 + v^2 + u^2v^2 & uv \\ uv & (D_u + v^2)/D_u \end{pmatrix} \succ 0$$

and positive Schur denominator $\mathcal{D} = u^2v^4 + u^2v^2 + u^2 + v^4 + 2v^2 + 1$. Its remaining scalar has positive factor $uvD_uD_v/\mathcal{D}$ and is proportional to

$$D_u uvw^2 - 2D_uD_v w + uv.$$

For $u, v > 0$ it is nonnegative if $w \leq 0$. Hence a negative-core test must have $w > 0$; with $\bar{y} = 0$ it cannot have $\bar{W} = -\bar{s}$. A nonnegative core is itself a pure testing square after evaluation, and each original event is already a square. Literal $u = 0$ or $v = 0$ is included in this direct exit.

The contact: five roots and a finite Taylor argument. We have reduced to $Z = 1$ and

$$u - k,\ W - s,\ v,\ x,\ y \text{ infinitesimal.}$$

Set

$$x_c = \eta k t s^2, \quad y_c = \eta k s a, \quad v_c = \eta k s, \quad U = u - k, \quad B = W - s,$$
$$N_1 = x - x_c, \quad N_2 = y - y_c, \quad N_3 = v - v_c.$$

The exact roots for this chart are

$$N_1,\ N_2,\ N_3,\ \eta U,\ \eta B. \tag{OA14}$$

Here is the full allocation argument, including the normal terms. As a polynomial in x, y, v, (OA5) belongs to

$$(x, y, v)^2 + \eta(x, y, v) + (\eta^2)$$

over the bounded polynomial coefficient ring in u, W and the source unit inverses already listed. Since x_c, y_c, v_c are η times bounded source elements, the shifted polynomial belongs to $(N_1, N_2, N_3)^2 + \eta(N_1, N_2, N_3) + (\eta^2)$. Its constant normal coefficient is η^2 times a polynomial in U, B whose constant and linear terms vanish by exact contact and angle stationarity. Its linear normal coefficients are η times polynomials in U, B whose constant terms vanish by exact state and Bob stationarity. These conclusions can be read in the coefficient field, where $\eta \neq 0$; the resulting quotients are bounded by the displayed factorization, not by cancellation of a mixed element. Monic Taylor division by U, B therefore allocates these terms to products of the roots (OA14). Terms of normal degree at least two are allocated to their first two normal factors. This is an exact finite Gram construction with bounded entries.

All terms beyond weighted degree two carry one of the infinitesimal deviations N_i, U, B, or a coefficient difference of zero residue. The Gram residue is consequently the weighted Hessian of (OA6) at

$$(u, W, \nu, \xi, \chi) = (k, s, ks, kts^2, ksa), \qquad Z = 1.$$

Its ξ, χ block is positive definite by (OA7). The Schur complement is $8aK$ times the half-Hessian of S in (ν, W, u) at this point. That half-Hessian is

$$\begin{pmatrix} K & k(k^2-1) & s(k^2-1) \\ k(k^2-1) & k^2 K & sk(k^4+1)/K \\ s(k^2-1) & sk(k^4+1)/K & s^2 K \end{pmatrix}, \tag{OA15}$$

with leading principal minors

$$K, \qquad 4k^4, \qquad 12 s^2 k^6 / K. \tag{OA16}$$

These are positive units. The five-root Gram is strict, which proves the contact chart without assuming that v/η, x/η, or y/η is bounded.

Exhaustion and restoration. For completeness, if the homogeneous Bob sine v_1 has nonzero residue, the four strong packet amplitudes have trivial common state kernel at every projective Alice angle. Indeed $x = y = 0$ and the other two amplitudes then give $v_1(u_0 W + u_1 Y) = v_1(u_1 X - u_0 Z) = 0$; since $u_0^2 + u_1^2$ is a testing unit, all state coordinates vanish. Their four-root Gram absorbs the infinitesimal signed coefficient; all other event squares may be retained. Thus only the Bob-small chart used above remains. The finite and reciprocal Alice charts, the two state branches, and the exact contact chart exhaust it. Every strict bounded residue Gram is tensor SOS by the established strict-Gram lemma. All projective multipliers are pure testing squares and are restored by separate field square roots. Complex states split into their real and imaginary parts.

Exact advance and remaining scope. The theorem closes the regular ordinary-k, comparable-f, g source slice (OA1), including the previous explicit $k = 1$ entry obstruction. It supplies a general mechanism: two completed strong squares, a scalar capacity factorization locating the unique angular degeneracy, and finite weighted contact division with a strict five-root Hessian. It makes no claim for a nonunit s, a nonunit shape or gap among a, t, M, J, non-small η or ζ, or the still-unclosed small-k unit/unbounded-η source rows.

Chapter 109

Two weak complementary events force the C2 radial certificate

Component p534f57e1da6f
Source: CORE/contact/chart_free_c2.tex
SHA256: 534f57e1da6f10b1f7e4c37aff830d3ad6bea003ec6350fabd1bdf5cedd3e7b7
Inventory use: Only exact physical witnesses, contact amplitude estimate, relative coalescence, scaled stationary signs and CT8 radial source entry; its all-tests calls are resolved by the actual radial/ordinary bodies listed here

109.1 Two weak complementary events force the C2 radial certificate

Here weak means infinitesimal after division by the original positive signed coefficient d. All source calculations take place in one real closed coefficient field. Testing fields remain independent.

Theorem 109.1 (C2 entry without a contact-chart hypothesis)**.** *Let the physical canonical parent have $\lambda = \beta = 0$, $d > 0$, and nonnegative original event coefficients. Suppose it has an actual finite noncommuting stationary zero contact, and*

$$C/d,\ \alpha/d \quad \textit{are infinitesimal.}$$

Then its full parent is tensor SOS at every independent real or complex projective test. No hypothesis on the other four event weights, on the contact-angle ratio, or on the signs of the source gaps is imposed.

More precisely, normalize $d = 1$. After local real sign gauges and at most the exact C2 reflection, the actual contact enters the positive rational C2 source with

$$c \textit{ bounded}, \qquad a, b, t \longrightarrow 1, \qquad r, h, g, j > 0 \textit{ infinitesimal.}$$

In particular all four matching weights are unbounded and both complementary weights are strictly positive on this finite-contact domain.

The proof has three ingredients: a finite physical witness makes the matching weights large; the quadratic CHSH excess controls the contact amplitudes at the scale of the actual angles; stationarity fixes their signs. This last step establishes the source chart rather than assuming it.

109.1.1 The physical bound at the correct angular scale

For $u, v > 0$, put $D_u = 1 + u^2, D_v = 1 + v^2$, and let $M(u, v)$ be the real symmetric matrix of the signed core

$$H = 2\{R_D^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_u D_v Z^2\},$$

where

$$R_m = vW - X + uvY - uZ, \qquad R_D = uW + uvX - Y - vZ.$$

Write $L = D_u D_v$. Direct matrix multiplication gives

$$(M - LI)^2 = L^2 I + 4uvLJ, \qquad J = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \qquad J^2 = I. \tag{109.1}$$

For a unit real state ψ, Cauchy–Schwarz therefore gives

$$\psi^T M \psi \geq L - \sqrt{L^2 + 4uvL} \geq -2uv.$$

The last inequality follows by squaring the positive quantities $L + 2uv$ and $\sqrt{L^2 + 4uvL}$; their squared difference is $4u^2v^2$. Homogeneity proves

$$H \geq -2uv \, \|\psi\|^2. \tag{109.2}$$

This is an exact ordered-field inequality, not a truncated expansion. It is the usual quadratic CHSH excess in the present normalization.

At a normalized zero contact, every event term is nonnegative. Define the four matching amplitudes, in the fixed order (n, A, m, D), by

$$\mathcal{L} = (\sqrt{D_u D_v}X, \sqrt{D_u D_v}Y, R_m, R_D).$$

Equation (109.2) gives, for each matching event,

$$e_i \mathcal{L}_i^2 \leq uv/2. \tag{109.3}$$

All square roots and inequalities in this paragraph are coefficient-side.

109.1.2 Matching weights and the compact contact geometry

First every matching weight is unbounded. To see the mechanism for D, use the coefficient-field experiment

$$u = 1, \quad v = \sqrt{1 + \delta}, \quad (W, X, Y, Z) = (1, 0, 0, \sqrt{1 + \delta}), \qquad \delta > 0.$$

The n, A, m packets vanish, and the full eight-event parent is exactly

$$4D\delta^2 - 2\delta + 8\lambda + 4(2 + \delta)C + 8(1 + \delta)\alpha + 4(2 + \delta)(1 + \delta)\beta.$$

If $D \leq B$ for a positive real B, choose the positive real $\delta = 1/[4(B + 1)]$. The first two terms have a strictly negative real upper bound, whereas the complementary terms are infinitesimal. This contradicts physicality. The exact even rotations of the event cycle give the same argument for n, A, m, preserving the sum of the four complementary weights. The accompanying checker also supplies the four witnesses directly. Hence $1/n, 1/A, 1/m, 1/D$ are infinitesimal.

Normalize the actual state to norm one and use positive actual half-angles, obtainable by local real basis gauges. The weaker bound $H/(D_u D_v) \geq -2$ shows that all normalized matching probabilities vanish in residue. In particular $X, Y \to 0$. Put

$$c_u = (1 + u^2)^{-1/2}, \; s_u = uc_u, \qquad c_v = (1 + v^2)^{-1/2}, \; s_v = vc_v.$$

If $\sigma = \mathrm{res}(c_u s_v)$ and $\tau = \mathrm{res}(s_u c_v)$, the two remaining matching equations read

$$\sigma\overline{W} - \tau\overline{Z} = 0, \qquad \tau\overline{W} - \sigma\overline{Z} = 0.$$

At least one of $\overline{W}, \overline{Z}$ is nonzero. Thus $\sigma = \tau \geq 0$. If positive, the two actual half-angles are bounded units with the same residue. If zero, both are infinitesimal or both unbounded, by the two unit-circle identities. The exact C2 reflection sends (u, v) to $(1/v, 1/u)$ and exchanges (C, α) and (m, D), preserving the hypotheses. We may therefore assume either ordinary bounded-unit angles or two infinitesimal angles.

109.1.3 Relative coalescence when both angles vanish

Suppose u, v are infinitesimal. From (109.3), all four $\mathcal{L}_i/\sqrt{uv}$ are infinitesimal. In particular $X/\sqrt{uv}, Y/\sqrt{uv} \to 0$. Set $a = v/u > 0$. Dividing the two other matching amplitudes by $\sqrt{uv}$ gives

$$\sqrt{a}\,W - \frac{Z}{\sqrt{a}} \longrightarrow 0, \qquad \frac{W}{\sqrt{a}} - \sqrt{a}\,Z \longrightarrow 0.$$

The discarded terms are $X/\sqrt{uv}, Y/\sqrt{uv}$ and $\sqrt{uv}X, \sqrt{uv}Y$, all infinitesimal. If a were unbounded, these equations would force both W, Z infinitesimal; the same follows if a is infinitesimal. Both contradict normalization. Thus a is a bounded unit. Taking residues now yields $a \to 1$ and $W/Z \to 1$, with both W, Z units. Consequently

$$X/u,\ Y/u,\ R_m/u,\ R_D/u \longrightarrow 0.$$

This is relative coalescence, which the compact-angle argument alone does not establish.

Normalize $W = 1$ by a pure coefficient factor. Write

$$x = X/u, \quad y = Y/u, \quad q = v/u, \qquad (u, x, y, Z, q) \longrightarrow (0, 0, 0, 1, 1).$$

The core divided by u^2 is the polynomial

$$\begin{aligned}\mathcal{H} := H/u^2 = 2\{&(1 - y - qZ + u^2qx)^2 + (1 + u^2q^2)(x + Z)^2 \\ &+ (1 + u^2)(u^2q^2y^2 - 2qyZ - q^2Z^2)\}.\end{aligned} \tag{109.4}$$

It has zero residue at this point. The nonnegative alpha packet and the zero-contact equation give

$$0 \le 4\frac{\alpha}{u^2}(1 + u^2)(u^2qy - Z)^2 \le -\mathcal{H}.$$

Its square factor is a positive unit. Hence

$$\alpha/u^2 \longrightarrow 0. \tag{109.5}$$

This estimate is derived before the scaled stationary equations are used.

109.1.4 Stationarity fixes the source signs

In the small-angle case introduce the four exact local coordinates

$$\ell = \left(\sqrt{D_uD_v}\,x, \sqrt{D_uD_v}\,y, q - x + u^2qy - Z,\ 1 + u^2qx - y - qZ\right).$$

Their Jacobian with respect to (x, y, Z, q) has residue

$$J_0 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & -1 & 1 \\ 0 & -1 & -1 & -1 \end{pmatrix}, \qquad \det J_0 = 2.$$

Thus its inverse and the transformed derivatives are bounded at the actual contact. Only this pointwise coefficient-field inverse is used. Direct differentiation of (109.4) gives

$$\operatorname{res}(\nabla_\ell \mathcal{H}) = 2(1, -1, -1, 1).$$

The full parent divided by u^2 is exactly

$$4\sum_i e_i\ell_i^2 + 4C(1 + u^2q^2)(1 - y)^2 + 4(\alpha/u^2)(1 + u^2)(u^2qy - Z)^2 + \mathcal{H}.$$

The two complementary-gradient terms have zero residue, by weak C and (109.5). The original stationary equations imply stationarity in ℓ, with u fixed. These equations first show that every $e_i\ell_i$ is bounded, so its residue is defined. Consequently

$$\operatorname{res}(n\ell_n, A\ell_A, m\ell_m, D\ell_D) = \tfrac{1}{4}(-1,1,1,-1). \tag{109.6}$$

Every matching weight is positive, so $X<0, Y>0, R_m>0, R_D<0$ as exact source inequalities.

For ordinary angles the argument is shorter. Normalize $W=1$; the actual contact has residue $(X,Y,Z,v)=(0,0,1,u)$, where u is a bounded unit. Use the unscaled $\mathcal{L}$ coordinates. Their Jacobian determinant at that residue is $2u(1+u^2)^2$, and

$$\nabla_{\mathcal{L}} H = 2u(1,-1,-1,1)$$

there. All complementary gradients are bounded times C,α, hence infinitesimal. Stationarity gives $\operatorname{res}(e_i\mathcal{L}_i) = \overline{u}\,(-1,1,1,-1)/4$, with the same exact signs.

109.1.5 The original source and all independent tests

In either case $W/Z\to 1$, $v/u\to 1$, and $X/u, Y/u\to 0$. Set, using the original actual contact,

$$c=u,\quad a=v/u,\quad b=aW/Z,\quad r=-X/(uZ),\quad t=(uW-Y)/(vZ).$$

Then $a,b,t\to 1$ and

$$h=b-a^2t=\frac{aY}{uZ}>0,\qquad g=\frac{R_m}{uZ}>0,\qquad j=-\frac{R_D}{vZ}>0.$$

All four r,h,g,j are infinitesimal, so $0<r<1$. The remaining chart boundaries are positive units:

$$1-c^2h=\frac{Z-vY}{Z}>0,\qquad b-a^2c^2r=\frac{a(W+vX)}{Z}>0.$$

These definitions give exactly

$$h=b-a^2t,\qquad g=(1+c^2)b-(1-r+a^2c^2t),\qquad j=1+c^2r-t.$$

There is no remaining sign-entry assumption.

Finally the stationary coefficient matrix has the nonzero determinant

$$4^6[-64ac^6rgt\,hj(1-c^2h)(1+c^2)^2(1+a^2c^2)^2\Delta].$$

The inherited short three-by-three reduction proves this formula. For $z=c^2$, its positive expansion is

$$\Delta = 1+(1+z)b+z(1+z)b^2+2z(1+z)br+z^2r^2+2zr+a^2z\{(1+z)b+1+zr(2-r)\}\geq 1.$$

The original coefficients therefore equal the unique positive rational C2 source formulas. In particular their positive numerator identity forces $C,\alpha>0$. If c is infinitesimal, the complete small-angle radial theorem, including its outer-test corollary, applies. If c is a unit, the inherited ordinary-angle radial corollary applies. The precise source files are identified in the reproduction paragraph below. These certificates concern the full parent and arbitrary independent testing fields. Reflect back when required and restore d by a pure coefficient square root. This proves Theorem 109.1.

The conclusion closes the full two-weak-complementary-event *finite-contact* C2 branch, not just a preselected rational chart. It does not assert that every weak-C source has weak α. The explicit one-weak escape family in the following supplement shows why that remaining implication fails.

Chapter 110

A short determinant check for exact source entry

Component `p61209c7390b0`
Source: `CORE/contact/source_determinant.tex`
SHA256: `61209c7390b099153809a244a5ea3a0f1060bd13f975a467b66ca9511b218635`
Inventory use: Actual C2 stationary-source uniqueness determinant | Complete algebraic stationary determinant reduction and uniqueness; Actual C2 stationary-source uniqueness determinant | Complete algebraic stationary determinant reduction and uniqueness

110.0.1 A short determinant check for exact source entry

Here is a reproducible uniqueness check for the six source coefficients. Let M have rows indexed by (W, X, Y, Z, u, v) and columns equal to the derivatives of

$$D_u D_v X^2, \quad R_2^2, \quad D_v R_3^2, \quad D_u D_v Y^2, \quad R_5^2, \quad D_u R_\alpha^2,$$

evaluated at $(u, v; W, X, Y, Z) = (c, ac; b/a, -cr, c(b/a - at), 1)$. Write $z = c^2$, $D = 1 + z$, $E = 1 + a^2 z$, $h = b - a^2 t$, $g = Db - (1 - r + a^2 zt)$, and $j = 1 + zr - t$. Factoring the six columns by

$$2cr, \quad 2cg, \quad 2act, \quad -2ch/a, \quad 2acj, \quad 2(1 - zh)$$

and the first row by c leaves the matrix

$$B = \begin{pmatrix} 0 & a & E & 0 & -1 & 0 \\ -DE & -1 & 0 & 0 & -az & 0 \\ 0 & az & -E & -DE & 1 & -acD \\ 0 & -c & 0 & 0 & ac & D \\ zrE & zh - 1 & (b/a)E & -zhE/a & azr - b/a & c(1 - zh) \\ azrD & b/a + zh/a & a^2 zt & -zDh & 1 + zr & -cDh/a \end{pmatrix}.$$

Use the pivots in row 2, column 1; row 1, column 3; and row 4, column 6. Eliminate those three columns, multiply the remaining second and third rows by D and E, and factor $-DE$ from their middle column. The determinant reduces to $\det B = D^2 E^2 \det T$, where

$$T = \begin{pmatrix} a & 1 & a^2 z \\ -1 - b - z(r + a^2 t) & zh/a & az(r - 1 + zh) \\ b/a + az(h - r) & zh & E(1 + zh) + z(r + a^2 t) \end{pmatrix}.$$

Subtract (zh/a) times the first row from the second and zh times the first row from the third. Expanding along the middle column gives

$$\det T = [1 + b + z(b + r)][1 + z(a^2 + b + r)] - z(1 - r)(b - a^2 zr) = \Delta.$$

The last equality follows by expanding the two products and is exactly the source definition of Δ. Restoring the factors proves

$$\det M = -64ac^6rgthj(1-zh)D^2E^2\Delta. \tag{110.1}$$

Each source coefficient occurs with a factor 4 in the parent, so the stationary linear system in $(e_1, e_2, e_3, e_4, e_5, \alpha)$ has determinant $4^6 \det M$. On the positive source domain this is nonzero. Since the six source formulas above satisfy all six stationary equations, they are their unique solution. This checks exact coefficient entry, independently of the subsequent asymptotic Gram argument.

Chapter 111

Exact C2 source and its uniqueness

Component `p6e5d41ed80b3`
Source: `CORE/contact/source_formulas.tex`
SHA256: `6e5d41ed80b37604ed7b057c73fddb378b4da3c993233566bf3c50b41176d477`
Inventory use: The six actual native C2 source formulas used in RC3-RC4; formulas only | Actual C2 source rational coefficients; The six actual native C2 source formulas used in RC3-RC4; formulas only | Actual C2 source rational coefficients

111.1 Exact C2 source and its uniqueness

In this appendix, c denotes the actual half-angle and C the event weight previously denoted c. Set

$$z = c^2, \quad A_c = 1 + zr, \quad h = b - a^2 t, \quad g = (1+z)b - (1 - r + a^2 zt), \quad j = 1 + zr - t,$$

$$\begin{aligned}\Delta &= (1+z)zb^2 + 2z(1+z)br + (1+z)(1+a^2z)b \\ &\quad + z^2(1-a^2)r^2 + 2z(1+a^2z)r + 1 + a^2z, \\ N &= (1+z)(b+r)(b+a^2(1-t)) - a^2(1+a^2z)(1-r)t.\end{aligned}$$

The unchanged $d = 1$, $\lambda = \beta = 0$ source is

$$n = \frac{(b+r)A_c(A_c + a^2zt) - r\Delta}{2r\Delta}, \qquad m = \frac{(1+a^2z)(1-r)A_c(A_c+a^2zt)}{2g\Delta},$$

$$C = \frac{A_c N}{2a^2 t\Delta}, \qquad A = \frac{a^2 A_c(b+r)(A_c + a^2zt)}{2h\Delta},$$

$$D = \frac{(1+z)(b+r)(b-a^2zr)(A_c+a^2zt)}{2a^2 j\Delta}, \qquad \alpha = \frac{z^2(b+r)N}{2(1-zh)\Delta}.$$

Its exact contact is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(c, ac; \frac{b}{a}, -cr, c\left(\frac{b}{a} - at\right), 1\right).$$

The positive chart requires $a, c, b, t > 0$, $0 < r < 1$, $h, g, j > 0$, $1 - zh > 0$, $b - a^2zr > 0$, and nonnegative original event weights. The source identities and the full independent-test radial certificates are included in the accompanying inherited modules; the source-entry proof above imposes no hidden shape-unit hypothesis.

The numerator has the useful positive identity

$$N = (1+c^2)(b+r)h + a^2\{g + j + rt(1 + a^2c^2)\} > 0.$$

It follows by substituting the three source-gap definitions; every term is positive in the domain used by the contact-entry theorem. For the determinant calculation use the aliases

$$R_2 = R_m, \quad R_3 = uW - Y, \quad R_5 = R_D, \quad R_\alpha = vY - Z, \qquad (e_1, e_2, e_3, e_4, e_5) = (n, m, C, A, D).$$

Source inclusion. The original input is reproduced once as Component p61209c7390b0 (Chapter 110).

Chapter 112

A natural weak-event entry into the complete uniform C1 theorem

Component `pa133fae35e52`
Source: `CORE/prior/c1/entry_and_remainder.tex`
SHA256: `a133fae35e5258c0730ee18f58ffefe20e959ddba3c6fef3905847d291e63593`
Inventory use: CE1–CE7 only | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

112.1 A natural weak-event entry into the complete uniform C1 theorem

Throughout, d is the original positive signed coefficient. The event coefficient at the fourth cycle position is denoted A_{ev}, and $D_{\mathrm{ev}} = e_5$. These must not be confused with actual state coordinates. The source has the canonical omissions $\lambda = n = 0$ and is normalized by $c = e_3 = 2$. We use the canonical actual-contact sign entry, including its admitted single-zero D_{ev} or β boundaries. Explicitly, the retained $m, c, A_{\mathrm{ev}}, \alpha$ are positive, and D_{ev}, β are positive or exactly one of them is zero. The noncoercive further-zero faces are already separate exits. The inherited sign and nonsingularity input is `C1_GLOBAL_SIGN_ENTRY.md`, Sections 6–7, equations (9)–(10), on every finite noncommuting physical contact in precisely this coefficient domain; that input is not reproved here.

Exact transport from the actual contact. Write its positive-angle contact as

$$u_c = k, \quad v_c = \ell, \qquad \psi_c = (S, k(\theta - 1), kSr, 1), \quad S > 0, \quad 0 < r, \theta < 1.$$

The exact C1 reflection lets us orient $0 < k \le 1$. Define source-only parameters

$$\eta = \frac{\ell}{kS}, \quad \zeta = \frac{S\ell}{k}, \quad f = \sqrt{\eta}, \quad g = \sqrt{\zeta}, \quad e = \frac{k}{S} = \frac{kf}{g}, \quad a = \frac{1 - r}{\eta}, \quad t = \frac{\theta}{\zeta}. \tag{CE1}$$

Thus the classifier $\mathfrak{T} = k/(S\ell)$ is exactly ζ^{-1}, and its old state-ratio variable B is exactly $\eta^{-1/2}$. Put $\kappa = k^2$, $K = 1 + \kappa$. The original amplitude signs imply

$$R = 1 + \kappa r, \quad T = 1 + \kappa(1 - \theta), \quad M = R - t > 0, \quad J = T - a > 0,$$

$$E = 1 - \kappa\eta(1 - \theta) > 0, \qquad Q_c = 1 - \kappa\zeta r > 0. \tag{CE2}$$

Indeed the positive second amplitude is $k\zeta M$, the negative fifth amplitude is $-\ell J$, the state gap is SE, and the remaining state gap is Q_c. In particular $0 < a, t < 2$, without any shape-unit hypothesis.

The seven native coquotients used by the uniform theorem are valid under (CE1), not just asymptotically. Their constructed contact is

$$(W, X, Y, Z) = (\Delta k, -k(1 - t\zeta), \Delta k^2(1 - a\eta), 1), \qquad \Delta = e^{-1} = S/k,$$

which is exactly the original contact. The native contact identities prove all six kernel and angular equations. To make the nonsingularity premise explicit, put $Y_0 = Sr$, $h_0 = S + k\ell(\theta - 1)$, $R_2 = \ell(S + k^2 Y_0) - k\theta$, and $R_5 = k(S - Y_0) + k^2\ell(\theta - 1) - \ell$. The inherited selected six-column contact system, on $(d, m, A_{\mathrm{ev}}, D_{\mathrm{ev}}, \alpha, \beta)$ with $c = 2$, has determinant

$$-8192\theta k^2 \ell (1 + k^2)^2 (1 + \ell^2)^2 \mathcal{D},$$

$$\mathcal{D} = Y_0 (1 + k^2)(kY_0 + \ell)(1 - k\ell Y_0) h_0 R_2 (-R_5) > 0.$$

Each factor is strictly positive by the stated source sign theorem. Thus the normalized coefficients equal the unique original source coefficients in that inherited system. This establishes full original-coefficient transport, including the exact zero openings; it does not use an archived completion label.

A bounded polynomial consequence of the original fifth event. Assume now

$$k \text{ infinitesimal}, \qquad \zeta = \mathfrak{T}^{-1} \text{ infinitesimal}. \tag{CE3}$$

Set $q = \kappa\eta$. Since $t < 2$ and $E = 1 - q(1 - t\zeta) > 0$, q is bounded and $\bar{q} \le 1$. The original fifth-event formula is

$$D_{\mathrm{ev}} = \frac{a}{Jc_3}(2c_5 - J\eta c_d).$$

Using $L_c = r$, $H_c = r + \eta$ and $V_c = q\zeta$, its sign is precisely

$$\mathcal{B}_5 := K(1 + q\zeta)(r + \eta) - J\eta P_d \ge 0. \tag{CE4}$$

Although η need not be bounded, this whole expression has the following bounded polynomial form. Define

$$\begin{aligned} H &= 2 + \kappa r + \zeta[K - (\kappa + 2)t] + q\zeta(-\zeta t^2 + t + 1) + \kappa\zeta(1 - r)(t - 1), \\ P_d &= 1 + q\zeta + \kappa H. \end{aligned}$$

Then exact cancellation using $a\eta = 1 - r$ and $q = \kappa\eta$ gives

$$\mathcal{B}_5 = Kr(1 + q\zeta) + (1 - r)P_d + q\{\zeta t(1 + q\zeta) - (K - \kappa\zeta t)H\}. \tag{CE5}$$

Every variable on the right is bounded. Hence (CE4) gives

$$1 - 2\bar{q} \ge 0, \qquad 0 \le \bar{q} \le \tfrac{1}{2}. \tag{CE6}$$

This is obtained before dividing by $1 - q$ or by a coefficient opening. In particular E is a positive unit with residue at least $1/2$.

The fourth coquotient polynomial likewise has the bounded expression

$$\begin{aligned} P_4 = {} & K(1 - q) + \kappa(1 - r) - \kappa\zeta t \\ & + q\zeta(-\kappa\zeta t^2 + 2\kappa t + t) + \kappa\zeta(1 - r)(1 - t). \end{aligned} \tag{CE7}$$

Thus $\bar{P}_d = 1$ and $\bar{P}_4 = 1 - \bar{q} \ge 1/2$. These exact low-degree identities are the physicality input needed for the entry, rather than an estimate on a truncated coefficient.

Theorem 112.1 (Weak fourth event gives the uniform entry)**.** *For an actual canonical C1 source satisfying (CE3), the following conditions are equivalent:*

$$A_{\mathrm{ev}}/d \text{ is infinitesimal}, \qquad \eta = f^2 \text{ is infinitesimal}, \qquad B \text{ is unbounded}. \tag{CE8}$$

Under any of these conditions the full original parent is tensor SOS at every independent projective test and every real or complex state, with no restriction on e.

Proof. The exact original source gives

$$\frac{A_{\mathrm{ev}}}{d} = \eta\frac{c_4}{c_d} = \frac{\eta P_4}{2rP_d}. \tag{CE9}$$

The ratio P_4/P_d is a positive bounded unit by (CE6)–(CE7). If (CE9) is infinitesimal then η/r is infinitesimal; since $r < 1$, η is infinitesimal. Conversely, if η is infinitesimal then $r = 1 - a\eta$ has residue one because $a < 2$, and (CE9) is infinitesimal. The final equivalence follows from $B = \eta^{-1/2}$. Now (CE1)–(CE3) give k, f, g positive infinitesimals and the unchanged positive bounded shapes and gaps of the uniform C1 source. The complete uniform theorem, including its newly proved unbounded-e branch, applies for every positive $e = kf/g$. All quantities used for entry lie in the coefficient field; no source/test rate comparison occurs. □

What the other weak templates mean on this same source row. The same calculation gives useful exact exclusions, not further SOS claims. In (CE3),

$$\frac{d}{c} = \frac{2(1-r)P_d}{K(r+\eta)E}, \qquad \frac{D_{\mathrm{ev}}}{d} = \frac{\mathcal{B}_5}{2J\eta P_d}, \qquad \frac{\alpha}{d} = \frac{\kappa^2\zeta(r+\eta)P_\alpha}{2Q_cP_d}. \tag{CE10}$$

Since $1 = r + a\eta \le 2(r+\eta)$, the first ratio is bounded: c/d is never weak in this orientation. Also P_α, Q_c are positive units, and $\kappa^2\zeta(r+\eta) = \kappa^2\zeta r + \kappa\zeta q$ is infinitesimal; hence α/d is always weak. If η is bounded, q is infinitesimal and $\mathcal{B}_5$ is a unit, so D_{ev}/d cannot be infinitesimal. If η is unbounded, $a = (1-r)/\eta$ is infinitesimal, J is a unit, and (CE10) shows D_{ev}/d infinitesimal. Thus

$$D_{\mathrm{ev}}/d \text{ weak} \quad\Longleftrightarrow\quad \eta \text{ unbounded}. \tag{CE11}$$

The complete remaining endpoint ratio is

$$\frac{\beta}{d} = \frac{K(\kappa r+q)[1+(1-t)\zeta] - t\zeta P_d}{2t\zeta P_d}. \tag{CE12}$$

On the unbounded-η branch, if β/d is weak too, this forces $q/(t\zeta)$ to have residue one. Equivalently e^2/t has residue one; in particular q is infinitesimal. This is an exact necessary critical balance, not an all-tests certificate for that branch. For completeness,

$$\frac{m}{d} = \frac{K(\kappa r+q)(1-t\zeta)(1+q\zeta)}{2M\zeta P_d}. \tag{CE13}$$

It records all weak-m possibilities without comparing source scales.

Precise complement and a genuine entry obstruction. Within (CE3), the new natural criterion closes exactly the infinitesimal η slice. The bounded-unit and unbounded η slices remain outside the uniform theorem. The C1 stabilizer exchanges A_{ev}, α, c, m, and D_{ev}, β, but sends k to $1/k$. Thus reflecting one of these small-k complements does not give a uniform small-k entry. In particular the fact that α/d is always weak does not close the row by a second use of (CE8).

Beyond (CE3), neither an ordinary actual Alice angle nor bounded $\mathfrak{T}$ has been covered by this entry theorem. This distinction cannot be removed by the weak-event hypothesis alone. Let $\varepsilon > 0$ be infinitesimal and take the exact native source

$$k = 1, \qquad f^2 = g^2 = \varepsilon, \qquad a = t = \tfrac{1}{2}. \tag{CE14}$$

Every original retained event is positive: their leading values are

$$c = 2, \quad m \sim \tfrac{2}{3}, \quad D_{\mathrm{ev}} \sim \tfrac{2}{3}, \quad \beta \sim 2, \qquad d \sim 4\varepsilon, \quad A_{\mathrm{ev}} \sim \varepsilon^2, \quad \alpha \sim \varepsilon^2.$$

The displayed native contact has kernel and angle stationarity by the six short native identities. On $Z = 0$ the signed core is a sum of squares, and its positive d, c, A_{ev} terms make the state restriction

positive definite, so the contact matrix has three positive complementary eigenvalues. The inherited Laurent determinant has nonnegative outer coefficients and two-dimensional support; its logarithmic convexity makes this stationary zero a global minimum. The full-parent inertia criterion, including the projective limits, therefore makes this an actual physical source. These are exactly the hypotheses of the previously proved stationary-contact sufficiency argument, not a numerical positivity test. Its normalized signed coefficient is infinitesimal, and $A_{\mathrm{ev}}/d, \alpha/d \sim \varepsilon/4$ are weak, so it belongs to the weak-A template. But $u_c = 1$, fixed by the C1 stabilizer, and $v_c = \varepsilon$. Neither orientation has the uniform requirement u_c infinitesimal. This is an obstruction to an exhaustive uniform entry claim, not a counterexample to tensor SOS.

The resulting source frontier is therefore explicit: the weak-A template is newly certified on the entire small-u_c/unbounded- $\mathfrak{T}$ row, with arbitrary e; its other actual-angle rows, and the unit/unbounded-η portions of that row after other weak orientations, still require additional analytical certificates or exits. No archived all-tests label has been promoted to a proof here.

Chapter 113

Global sign entry for the adjacent contact family

Component `pe31b9ef5ec43`
Source: `CORE/prior/inherited/C1_GLOBAL_SIGN_ENTRY.md`
SHA256: `e31b9ef5ec430cd15faef01eb6b7a33a604ee73f342def5fc245a7aa4a020ccc`
Inventory use: Retained exact cofactor, selected minor, Cramer identity, seed and C1 sign proof; analytical foundational calls replaced by SE1-SE10

113.1 Global sign entry for the adjacent contact family

113.1.1 Statement and exact scope

Use the canonical full parent in coefficient order $c = (d, \lambda, e_1, e_2, e_3, e_4, e_5, \alpha, \beta)$. On the adjacent face $\lambda = e_1 = 0$, assume d>0 and initially that the other six event coefficients are strictly positive. Normalize $e_3 = 2$. At a physical noncommuting zero take the angle gauges $u, v > 0$ and normalize the last kernel coordinate to one:

$$\psi = (A, u(\eta - 1), uY, 1)^T.$$

Then

$$\boxed{A > 0, \qquad 0 < Y < A, \qquad 0 < \eta < 1.} \tag{1}$$

These are actual ordered-field inequalities, with no residue or rate assumptions. The assertion holds over every real closed extension of the reals. It applies to every such boundary contact, without an extremality assumption. Section 6 extends the result to the exact three-consecutive-omission boundary. This is a sign theorem, not an SOS certificate for the full family.

The proof uses the established full-parent determinant, strict curvature, and contact sufficiency theorems. Its only new continuity argument is spelled out below; connectedness is used on a specified positive parameter orthant, not assumed for an unspecified contact locus.

113.1.2 1. Every positive event tuple has one analytic maximal-signed contact

Put $w = (e_2, e_4, e_5, \alpha, \beta) \in \mathbb{R}^5_{>0}$, retaining $e_3 = 2$, and define

$$d_*(w) = \max\{d \geq 0 : (d, 0, 0, e_2, 2, e_4, e_5, \alpha, \beta) \in C\}. \tag{2}$$

This maximum exists, is strictly positive, and is finite. Closedness and convexity give a closed interval of admissible nonnegative d. At d=0 all outer determinant coefficients are nonnegative, and the central coefficient is strictly positive: its nonnegative expansion contains $16e_2e_3e_4e_5$. For sufficiently small d>0 the central coefficient remains positive and all outer coefficients remain nonnegative, proving physicality by full-parent inertia. Testing a fixed strictly CHSH-violating realization bounds d above.

For these six positive events the determinant support is a fixed two-dimensional support with zero in its interior. The logarithmic Laurent determinant therefore is strictly convex and coercive. At d=d_* its minimum must be zero: a strictly positive minimum is stable under small increases of d, by the uniform supported-perturbation estimate in the routing proof. Thus the minimum supplies unique squared contact angles $U = u^2, V = v^2 > 0$. The parent has three positive eigenvalues, so its kernel is simple. The established packet argument shows that its last coordinate is nonzero. Consequently the chosen positive angles and normalized kernel are unique.

At this contact the core expectation is strictly negative. Indeed,

$$0 = d_* \, \psi^T H \psi + \sum_i w_i \, \psi^T E_i \psi.$$

If all retained expectations vanished, all retained event matrices would annihilate the state and the kernel equation would give H psi=0. The core is invertible at a finite noncommuting pair, a contradiction. At least one expectation is positive, and d_*>0, giving $\psi^T H \psi < 0$.

Let z consist of the three affine state coordinates and two positive angles, and let fd, w, z be the full polynomial parent slack. Its complete five-variable contact Hessian is positive definite. The equations

$$\nabla_z f = 0, \qquad f = 0$$

have Jacobian with respect to z, d

$$\begin{pmatrix} \nabla_z^2 f & \partial_d \nabla_z f \\ 0 & \partial_d f \end{pmatrix},$$

which is invertible because the upper block is positive definite and $\partial_d f = \psi^T H \psi < 0$. The Nash implicit function theorem gives a local analytic branch of d and z. Stationary-contact sufficiency makes that branch physical; its negative core expectation prevents increasing d at the fixed test. Thus it equals d_*w, with the unique chosen contact. These branches agree on overlaps, giving analytic functions on the entire connected orthant $\mathbb{R}^5_{>0}$.

113.1.3 2. A>0 from the negative CHSH cone

Write an arbitrary real state as A, X, Y_s, Z. Set $h = 1 + v^2 + u^2 v^2$. The X,Y_s block of the raw core H is

$$2 \begin{pmatrix} h & -uv \\ -uv & h \end{pmatrix} \succ 0.$$

Its Schur complement on A,Z is

$$\frac{2(1+u^2)(1+v^2)}{h^2 - u^2 v^2} \begin{pmatrix} u^2 v^2 & -uvh \\ -uvh & u^2 v^2 \end{pmatrix}. \tag{3}$$

For AZ<=0 its quadratic form is nonnegative. Completing the X,Y_s square then gives psi^T H psi>=0. Every contact above has negative core expectation, so AZ>0. With Z=1, this proves A>0.

Every physical noncommuting contact with this fixed event tuple is on the maximal graph (2): its negative core expectation means that increasing d at the fixed test immediately produces a negative slack.

113.1.4 3. eta>0 from a positive cofactor identity

Let B=adj$P$. At contact B=gamma psi psi^T for gamma>0, and hence

$$u B_{44} + B_{24} = \gamma u \eta.$$

Here indices are one-based. Direct full-parent expansion gives

$$u B_{44} + B_{24} = 16 u v^2 (1+u^2)^2 (1+v^2) \, F_\eta, \tag{4}$$

where, writing Dv=1+vˆ2,

$$\begin{aligned} F_\eta = (d+2e_5)\big[(d+2\alpha)(e_2Dv+e_3u^2v^2)+2e_4Dv(e_2+e_3u^2)\big] \\ +2e_2e_3\big[d(u^2v^2+Dv)+2\alpha+2e_5v^2(1+u^2)\big]. \end{aligned} \tag{5}$$

Every term is nonnegative, and d,e2,u,v>0 give strict positivity. Thus eta>0 throughout the contact graph.

113.1.5 4. The three remaining sign walls cannot be crossed

At any physical zero, a retained event with zero amplitude can be removed while preserving the kernel and angular stationarity equations: the event value and every first derivative of its squared amplitude vanish there. The stationary-contact theorem makes the resulting nonnegative coefficient vector physical. A physical d>0 vector with noninterior Newton support, however, cannot have a noncommuting zero, by the routing theorem.

If Y=0, the retained e4 event vanishes. Removing it gives omissions lambda,e1,e4, including the distant pair lambda,e4. If A-Y=0, the retained e3 event vanishes, and its removal gives the distant omitted pair e1,e3. Both contradict the preceding paragraph. Therefore Y and A-Y never vanish.

For eta=1, use the exact cross-multiplied contact identity

$$e_2\,[A+uv(\eta-1)]\,[v(A+u^2Y)-u\eta] = 2u^3(A-Y)(1-\eta)(1+v^2). \tag{6}$$

To justify (6) even when a selected Cramer minor vanishes, first derive it on the nonzero-minor locus from the verified rational contact solution. The selected minor is an analytic function on the connected positive event orthant. It is not identically zero, as the exact seed in Section 5 shows. A nonzero analytic function on a connected open set cannot vanish on an open subset; hence the nonzero-minor locus is dense. Both sides of (6) are continuous on the entire analytic contact graph, so (6) extends everywhere. This explicitly handles exceptional Cramer points.

At eta=1, A>0 and e2>0 make (6) force $v(A+u^2Y)-u=0$, which is precisely the retained e2 amplitude. Removing e2 gives the distant omitted pair lambda,e2, again impossible. Hence eta-1 never vanishes.

The continuous nonzero functions Y, A-Y, and eta-1 have fixed signs on the connected positive event orthant. The exact seed gives their signs.

113.1.6 5. Exact positive seed and transfer

Take

$$\begin{gathered} u=v=A=1, \quad Y=\tfrac{1}{4}, \quad \eta=\tfrac{1}{2}, \\ c=\left(\tfrac{27}{2},0,0,4,2,9,\tfrac{5}{4},\tfrac{13}{4},\tfrac{9}{4}\right). \end{gathered} \tag{7}$$

The complete parent kernel and both angle stationarity equations hold exactly; the selected Cramer minor is nonzero. All six event weights are positive. Contact sufficiency therefore places it on the graph (2). It has Y>0, A-Y>0, and eta-1<0. The previous section proves (1) over the reals.

The final implication is a finite first-order assertion in the coefficient, state and angle variables: physicality is expressed by quantified polynomial PSD inequalities; the kernel and stationarity equations are polynomial; positivity of u,v,d and the six retained events and the desired signs are ordered-field formulas. Transfer proves (1) over every real closed extension. No logarithm or analytic continuation is used within a non-Archimedean field.

113.1.7 6. Exact three-consecutive omissions

After choosing the adjacent omitted pair lambda=e1=0, a three-consecutive omission has either beta=0 or e5=0. The other five events are positive. At any real finite noncommuting contact with d>0, the full Hessian remains positive definite and the core expectation remains strictly negative by

the same established theorems. Apply the implicit function argument of Section 1 at this boundary tuple and open its third omitted coefficient to epsilon>0, keeping the other event weights fixed. For sufficiently small epsilon the signed coefficient stays positive and the branch has nonnegative coefficients. Stationary-contact sufficiency makes it physical. Thus it belongs to the six-positive-event graph already treated.

Taking the analytic limit gives Y>=0, A-Y>=0, and eta<=1. Strict A>0 and eta>0 follow directly from Sections 2 and 3; the cofactor formula stays strict when beta=0 or e5=0. The still-positive e4 and e3 coefficients make the same distant-omission removal arguments exclude Y=0 and A-Y=0. Equation (6) passes to the limit along the analytic branch; e2 remains positive, so eta=1 again forces the forbidden e2 removal. This proves strict inequalities (1) at the three-omission boundary as well.

This limit is taken at a fixed finite real noncommuting contact, where the implicit-function hypotheses are verified. It does not assume convergence of an optimizer escaping to infinity. The resulting first-order implication then transfers to every real closed extension.

113.1.8 7. All amplitude signs and a single rational chart

There is a physical Bell reflection preserving the adjacent omitted pair. It exchanges e2 with e3, e4 with alpha, and e5 with beta (and lambda with e1). After real local basis gauges its angle action is u'=1/u, v'=v and its state action, up to a positive normalization, is

$$\psi' = (vA - u(\eta - 1),\ -A - uv(\eta - 1),\ 1 - uvY,\ uY + v)^T. \tag{8}$$

The matrix implementing this state action is

$$G = \begin{pmatrix} v & -1 & 0 & 0 \\ -1 & -v & 0 & 0 \\ 0 & 0 & -v & 1 \\ 0 & 0 & 1 & v \end{pmatrix}, \qquad G^T G = (1 + v^2) I.$$

If c' denotes the displayed coefficient permutation, direct full-parent congruence gives $u^2(1 + v^2)P_{c'}(1/u, v) = GP_c(u, v)G^T$. The transformed event weights satisfy the same positivity or single-zero conditions. Apply (1) to the transformed contact and divide its state by its positive last coordinate uY+v. The inequalities Y'>0, eta'<1, eta'>0, and Y'<A' give, respectively,

$$1 - uvY > 0, \qquad h_0 := A + uv(\eta - 1) > 0,$$

$$R_5 := u(A - Y) + u^2 v(\eta - 1) - v < 0, \qquad R_2 := v(A + u^2 Y) - u\eta > 0. \tag{9}$$

Thus every one of the eight event amplitudes has a fixed, strictly nonzero sign in this canonical state gauge. In particular, the denominator product in the established selected-minor identity is strictly positive:

$$\mathcal{D} = Y(1 + u^2)(uY + v)(1 - uvY)h_0 R_2(-R_5) > 0,$$

$$\det J_{(d,e_2,e_4,e_5,\alpha,\beta)} = -8192\eta u^2 v(1 + u^2)^2(1 + v^2)^2 \mathcal{D} < 0. \tag{10}$$

Consequently the same e3=2 rational Cramer formula is valid on both the exact two-omission family and its routed three-consecutive-omission boundaries. It is legitimate here to include a zero coefficient's column in the selected square matrix: (10), rather than an active-column deletion argument, proves its invertibility. This replaces the earlier need for six separate five-row templates on the adjacent three-omission boundary.

For six positive events, (10) also proves that every maximal-signed boundary point above is an extreme ray of the full canonical cone. The seven active columns have rank six and their kernel is one-dimensional. For a single additional event omission, the six active columns have rank five: the unique kernel of all seven columns has zero in that omitted coordinate and is strictly positive on the remaining active coordinates. The exact contact-face description then proves extremality as well.

113.1.9 8. What this closes

The adjacent exact-two-omission contact formula now enters its positive shape domain globally: A>0, t=u/A>0, 0<r=Y/A<1, and 0<eta<1. The same conclusion holds for the routed three-consecutive-omission contact families, with the same rational formula and a strictly signed denominator. Those quantities need not have positive real residues or be bounded. Consequently this theorem closes sign entry, but not the remaining joint rate and optimizing-angle boundary certificates. The separated omission face remains outside this theorem.

113.1.10 Verification and review

The checker verify_sign_entry.py reconstructs the full parent and verifies the negative-core Schur complement, the positive cofactor identity, the cross-multiplied contact identity on the Cramer locus, the exact seed kernel, both angular derivatives, its selected minor -8640, and the positive d=0 central coefficient. The analytic graph, dense extension, no-wall arguments, and real-closed-field transfer are proved in the text. Two same-system agents reviewed the argument and its three-omission extension; this is not external or proof-assistant verification.

Chapter 114

The unbounded-e C1 source away from the zero-W state

Component `p7ef919c24082`
Source: `CORE/prior/inherited/c1_unbounded_e_w_unit.tex`
SHA256: `7ef919c2408221436586e92643ba0659249ec05a53524eabfb2eb4f6c0ab39d1`
Inventory use: UW1–UW3 only; old source and state restrictions are not silently imported | C1_BARE_ OTHER_TERMINALS; precise subordinate uses retained below

114.1 The unbounded-e C1 source away from the zero-W state

The complementary state chart has a short retained-current proof. It does not require a contact division or a comparison of coefficient and testing rates.

Theorem 114.1 (Unbounded-e, W a unit)**.** *Use the unchanged canonical uniform C1 source of Section 123.1, with $k, f, g > 0$ infinitesimal, $e = kf/g$ unbounded, and the original positive bounded shapes and gaps $a, t, M = R - t, J = T - a$. Retain all original nonnegative event coefficients. Then the entire parent is tensor-SOS for every independent test with u, v, X, Y infinitesimal, Z bounded, and W a unit.*

Proof. Normalize $W = 1$ using the testing field and put $\Delta = 1/e$. Evaluate the seven short source polynomials with their native $K = k^2$, $\eta = f^2$, $\zeta = g^2$ and write $c_b = c_\beta$. Thus

$$R = 1 + k^2(1 - af^2), \qquad T = 1 + k^2 - k^2 g^2 t,$$

and their residues are $2, 2, 2, 2, 2, 2, 4$ in the original order, irrespective of the size of e. These are the same polynomial source identities as before, now used on the unbounded-e domain. Write $D_u = 1 + u^2$, $D_v = 1 + v^2$ and

$$Q = W + uY, \quad B = X + uZ, \quad r = uW - Y, \quad r_2 = vQ - B, \quad r_5 = r - v(Z - uX).$$

Thus Q is a pure-testing unit. All appearances of Q^{-1} below are inverses in the testing field, with bounded residue one. Let $H_2 = 4Tc_2$, $H_b = 4Tc_b$, $H_3 = 4Rc_3$, $H_5 = 4Rc_5$. The exact original source, with $g = \Delta kf$, is

$$\begin{gathered} c = 2, \quad d = \frac{2af^2 c_d}{c_3}, \quad m = \frac{2ae^2 c_2}{Mc_3}, \quad A = \frac{2af^4 c_4}{c_3}, \quad \alpha = \frac{2a\Delta^2 k^6 f^4 c_\alpha}{c_3}, \\ D + \frac{d}{2} = \frac{2ac_5}{Jc_3}, \qquad \beta + \frac{d}{2} = \frac{2ae^2 c_b}{tc_3}, \qquad \Lambda = \frac{RTc_3}{2a}. \end{gathered} \tag{UW1}$$

The letter D in (UW1) is the original coefficient e_5; it is distinct from the displayed testing factors D_u, D_v. Directly combining the three positive signed-core squares with their matching events gives

the full-parent identity

$$\begin{aligned}\Delta^2\Lambda F = {} & H_2\frac{R}{M}r_2^2 + H_b\frac{R}{t}D_vB^2 + \Delta^2H_3\frac{T}{a}D_vr^2 + \Delta^2H_5\frac{T}{J}r_5^2\\ & + 4RT\Delta^2f^4c_4D_uD_vY^2 + 4RT\Delta^4k^6f^4c_\alpha D_u(vY-Z)^2\\ & + 2RT\Delta^2f^2c_dD_u(v^2Y^2 - 2vYZ - v^2Z^2).\end{aligned} \tag{UW2}$$

Use $R/M = 1 + t/M$, $R/t = 1 + M/t$, $T/a = 1 + J/a$ and $T/J = 1 + a/J$. Retain the four positive excess packets, the complete alpha packet, and the positive v^2Y^2 part of the last line. The remaining expression has the five-root frame

$$z = (r_2, B, \Delta r, \Delta r_5, \Delta f^2 Y).$$

Its diagonal part is

$$H_2z_1^2 + H_bD_vz_2^2 + H_3D_vz_3^2 + H_5z_4^2 + 4RTc_4D_uD_vz_5^2.$$

The two complete negative terms are, exactly,

$$-4RTc_dD_u\frac{\Delta Z}{Q}\,z_5(z_1+z_2) - 2RTc_dD_u\frac{\Delta^2f^2Z^2}{Q^2}(z_1+z_2)^2, \tag{UW3}$$

because $vQ = z_1 + z_2$. Every displayed Gram coefficient is bounded. The correction coefficients in (UW3) have zero residue, while the diagonal residue is $8I_5$. The bounded strict-residue Gram lemma therefore applies. Restoring the six retained nonnegative packets and undoing the positive pure-coefficient factor $\Delta^2\Lambda$ proves the assertion. No shape or gap is inverted in the bounded Gram; the possibly unbounded gap ratios occur only in retained squares. □

Corollary 114.2 (Precise remaining unbounded-e test domain)**.** *For this source, the only independent testing domain not covered by the preceding theorem and the existing uniform outer-angle identities is*

$$Z = 1, \qquad u, v, W, X, Y \text{ infinitesimal.}$$

Proof. The two strength ratios $d/c, d/D$ and the original harmonic mass formula (UA1) depend only on k, f, g being infinitesimal. In particular the mass has residue $1/4$ for unbounded e as well. Hence the two outer-angle lemmas apply without change, including homogeneous infinite-angle endpoints. The literal axes give a nonnegative pure-testing core. Normalize a nonzero state by a coordinate of maximal absolute value. If X or Y is a unit, that core has positive residue $2(X^2+Y^2)$. Otherwise X, Y are infinitesimal; the theorem covers W a unit, and its complement has Z a unit and W infinitesimal. Pure-testing Z normalization gives the stated domain. □

Chapter 115

All tests for the uniform infinitesimal C1 source

Component `p6cad7544d1b8`
Source: `CORE/prior/inherited/c1_uniform_all_tests.tex`
SHA256: `6cad7544d1b8537e36bef0527c8f2a6dd53113d9a6ba10856bea98dc9142a113`
Inventory use: Two strong currents and Critical current mass lemmas, before the uniform theorem | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

115.1 All tests for the uniform infinitesimal C1 source

The two state charts can be joined without importing an all-tests atlas label. We first record two short outer-angle identities.

Lemma 115.1 (Two strong currents)**.** *Suppose the original event coefficients are nonnegative, $d > 0$, and $d/c, d/D$ are infinitesimal, where $c = e_3$, $D = e_5$. Then every testing chart with $1/v$ bounded is tensor-SOS, for every other angle and every state.*

Proof. Put $g = cD/(c+D)$, $h = D/(c+D)$, $R = uW - Y$, $C = v(uX - Z)$, $B = X + uZ$, and $N = R + hC$. Let $D_u = 1 + u^2$, $D_v = 1 + v^2$ and set

$$K_0 = 2\sqrt{g}\,C, \qquad B_0 = \sqrt{2d}\sqrt{D_v/D_u}\,B, \quad c_u = D_u^{-1/2}, \quad t_u = uD_u^{-1/2}, \quad \eta_v = \sqrt{D_v}/v.$$

The complete selected parent is exactly

$$\begin{aligned} 4cD_vR^2 + 4D(R+C)^2 + d\mathcal{H} = {} & 4(c+D)N^2 + 4cv^2R^2 + 2d(R+C)^2 \\ & + 2dD_u(vY - Z)^2 + (K_0, B_0)Q(K_0, B_0)^T, \end{aligned}$$

where

$$Q = \begin{pmatrix} 1 - \dfrac{d}{2g}c_u^2\eta_v^2 & \sqrt{\dfrac{d}{2g}}t_u\eta_v \\ \sqrt{\dfrac{d}{2g}}t_u\eta_v & 1 \end{pmatrix}.$$

To check the identity, use $Z = (uB - C/v)/D_u$ and complete the two current squares. Since $d/g = d/c + d/D$ is infinitesimal and c_u, t_u, η_v are bounded, $\bar{Q} = I_2$. Restore every unselected positive event. All roots separate between fields. Clearing the original pure-testing homogeneous angle factors gives the same identity at literal infinity: $\eta_v^2 = 1 + (1/v)^2$, and c_u, t_u are the normalized homogeneous Alice coordinates. Thus no infinite-angle endpoint is excluded. □

Lemma 115.2 (Critical current mass outside the small Alice chart)**.** *Under the preceding strength assumptions, suppose*

$$\gamma = \frac{Ag}{d^2} \quad \textit{is bounded}, \qquad \bar{\gamma} \geq \tfrac{1}{4},$$

where $A = e_4$ is the original event coefficient. Then every independent test with v infinitesimal and $1/u$ bounded is tensor-SOS, for every state, including the homogeneous Alice-infinite endpoint.

Proof. Keep $R, C, B, N, g, h, c_u, t_u$ from the preceding proof, and put $\epsilon = d/g$, $q = \sqrt{\epsilon}$ and

$$K_1 = 2\sqrt{g}\,C + \sqrt{d\epsilon}\,Y, \quad B_1 = \sqrt{d}(B - uvY), \quad V_0 = \sqrt{d}\,vY, \quad Y_0 = \sqrt{d\epsilon}\sqrt{D_u}\,Y.$$

Define

$$L_0 = \frac{qc_u}{2}(K_1 - c_u Y_0) - t_u v B_1 + c_u v V_0.$$

Direct multiplication gives $L_0 = \sqrt{dD_u}\,v(vY - Z)$ and the exact full selected-parent identity

$$\begin{aligned} &4cD_v R^2 + 4D(R + C)^2 + 4AD_u D_v Y^2 + d\mathcal{H} \\ &\qquad = 4(c + D)N^2 + 4cv^2R^2 + 2d(R + C)^2 + \mathcal{G}, \\ \mathcal{G} &= K_1^2 + 2D_v(B_1^2 + V_0^2) + (4\gamma D_v - c_u^2)Y_0^2 \\ &\quad + 2qv(K_1 - c_u Y_0)V_0 - 2L_0^2. \end{aligned}$$

The four-root Gram of $\mathcal{G}$ has residue $\operatorname{diag}(1, 2, 2, 4\bar{\gamma} - \bar{c}_u^2)$. Its last entry is at least $1 - \bar{c}_u^2 = \bar{t}_u^2 > 0$ because $1/u$ is bounded. Every correction is polynomial in bounded q, v, c_u, t_u, γ and has zero residue. The strict-residue lemma and restoration of the other events prove the claim. The displayed bounded homogeneous factors also cover literal Alice infinity. □

Theorem 115.3 (Whole uniform infinitesimal C1 domain)**.** *Use the unchanged canonical uniform C1 source, including all exact coquotients and original coefficient zeros. Suppose*

$$k, f, g, e > 0 \text{ are infinitesimal}, \qquad e = kf/g,$$

and the positive bounded shapes satisfy

$$R = 1 + k^2(1 - af^2), \quad T = 1 + k^2 - k^2 g^2 t, \qquad M = R - t > 0, \quad J = T - a > 0.$$

Assume every original event coefficient is nonnegative. Then the entire parent is tensor-SOS at every independent projective measurement pair and every real or complex testing state. There is no restriction on $b = g/k$ or on $A_c = g/f$.

Proof. First derive the two outer-angle premises from the original coefficients. Write $B_c = 2c_5 - Jf^2 c_d$. The exact source gives

$$D = \frac{aB_c}{Jc_3}, \qquad \frac{d}{D} = \frac{2Jf^2 c_d}{B_c}, \qquad \frac{d}{c} = \frac{af^2 c_d}{c_3}.$$

Since $\bar{B}_c = 4$, both ratios are infinitesimal, independently of any loss of residue of a or J. Also, for the original harmonic mass $g_B = cD/(c + D)$,

$$\frac{Ag_B}{d^2} = \frac{c_4 c_3 B_c}{c_d^2(2Jc_3 + aB_c)}. \tag{UA1}$$

The denominator is a unit: its residue is $16 \cdot 4(\bar{J} + \bar{a}) = 64$, since $J + a = T$ has residue one. The numerator has residue 16. Thus the original mass in (UA1) has residue exactly $1/4$. This argument uses neither a strong first pair nor beta positivity beyond the original source hypotheses.

The two preceding lemmas handle every test except u, v infinitesimal. Literal $u = 0$ or $v = 0$ also gives a nonnegative pure-testing core immediately. Normalize a nonzero state by a coordinate of largest absolute value; all coordinates are then bounded and at least one is a unit. If X or Y is a unit, the pure-testing core has positive residue $2(X^2 + Y^2)$, so its product with the positive coefficient d is itself a square and all other events are positive squares. It remains that X, Y are infinitesimal.

The following coefficient-only split is exhaustive. If $b = g/k$ is bounded, then $f = eb$ and $g = kb$, so the bounded-b Z theorem treats Z a unit and arbitrary bounded W. If Z is infinitesimal, W is a unit, and the bounded-b all-rho W endpoint applies after $W = 1$ normalization.

If b is unbounded, put $A = f$, $H = g$, $K = k/g = 1/b$. All three are infinitesimal. Original beta nonnegativity gives exactly

$$\beta + \frac{d}{2} = \frac{2ae^2c_b}{tc_3}, \qquad \frac{d}{2} = \frac{af^2c_d}{c_3} \quad \Longrightarrow \quad E := tb^2 \le \frac{2c_b}{c_d}.$$

Hence $E > 0$ is bounded with $\bar{E} \le 1$ and the exact source map is

$$(f, g, k, e, a, t) = (A, H, HK, AK, a, K^2E).$$

Both original gaps, every coefficient and every literal coefficient zero are preserved. The new inverse-b Z theorem handles Z a unit; the finite ten-root W theorem handles the complementary W-unit, Z-infinitesimal chart. Thus all state cases are exhausted.

The zero state is trivial. For a complex state the real parent matrix splits its evaluation into the sum of its real and imaginary quadratic evaluations. All state normalizations and homogeneous projective factors are pure testing elements. No comparison of an independent testing quantity with a coefficient scale was used. □

Chapter 116

Matching both weak C2 packets to the four strong packets

Component `p8b53b68d8af6`
Source: `CORE/prior/inherited/c2_matching_strength.tex`
SHA256: `8b53b68d8af6a308d041e96a3bf10b86a1c991010146e95da907ad2d40e15ec0`
Inventory use: Exact complete-parent reflection, both product-bound corner joins, and full ordinary-angle radial source proof

116.1 Matching both weak C2 packets to the four strong packets

A coefficient condition symmetric in the two weak packets closes both commuting angle corners at once. It also gives a second exact radial source family whose contact angles have ordinary positive residue.

Theorem 116.1 (Two matched weak packets)**.** *Let the original physical C2 parent have $d = 1$, $\lambda = \beta = 0$, and event weights (n, m, C, A, D, α). Suppose $C, \alpha > 0$ are infinitesimal, all four other weights are positive, and each of*

$$Cn,\ Cm,\ CA,\ CD, \qquad \alpha n,\ \alpha m,\ \alpha A,\ \alpha D$$

has bounded reciprocal in the coefficient field. Then the full parent is tensor SOS at every independent projective test and every real or complex state.

Proof. Lemma 117.1 applies directly to every commuting residue test except the finite zero-zero corner. For that corner use the exact reflection

$$(u, v; W, X, Y, Z; n, m, C, A, D, \alpha) \longmapsto (1/v, 1/u; Z, X, Y, W; n, D, \alpha, A, m, C). \tag{116.1}$$

More explicitly, if F denotes the full parent with its indicated arguments, then

$$F(u, v; W, X, Y, Z; n, m, C, A, D, \alpha) = u^2 v^2 F(1/v, 1/u; Z, X, Y, W; n, D, \alpha, A, m, C). \tag{116.2}$$

This is a homogeneous polynomial identity, so its interpretation in reciprocal charts includes literal zero and infinity endpoints.

For a short verification, the reflected R_2, R_5 amplitudes, after multiplication by uv, are $-R_5, -R_2$. The reflected C and α packets exchange, while the n, A packets are fixed. If $\mathcal{H}$ is the signed core, its common expanded form is

$$\begin{aligned}\mathcal{H}/2 = {} & u^2(W^2 + Z^2) + (1 + v^2 + u^2 v^2)(X^2 + Y^2) \\ & + 2u^2 v(WX - YZ) + 2u(XZ - WY) - 2uv(WZ + XY).\end{aligned}$$

Multiplying its reflected expression by u^2v^2 fixes the first, second and last terms, and exchanges the middle two. This proves the whole-parent identity, including its signed part.

The reflected weak weight is α, its four strong weights are n, D, A, m, and its other nonnegative weak packet is C. Thus all hypotheses of Lemma 117.1 hold for the reflected parent. Its infinity-infinity chart is precisely the original zero-zero chart. Restore the homogeneous pure-testing factors to obtain the certificate there as well.

Every noncommuting residue test is covered by the already-established physical-source theorem, after positive pure-coefficient normalization if a coefficient is unbounded. Undoing that normalization preserves SOS. These alternatives exhaust the tests, and real/imaginary splitting handles complex states. No coefficient scale was compared with an independent testing scale. □

Corollary 116.2 (Radial C2 source with ordinary contact angles)**.** *Use the same rational source* (119.1)*, now with source half-angle c a positive bounded unit. Suppose $h = b - a^2t, r, g, j$ are positive infinitesimals and $a, 1 - r, t$ are positive bounded units. Then the full parent is tensor SOS at every independent projective test and every real or complex state.*

Proof. Put $z = c^2$ and continue to write $C = e_3$ for the weak event weight. The exact source identities

$$b - 1 = -r + g - zh, \qquad t - 1 = zr - j, \qquad a - 1 = \frac{g + j - (1 + z)(r + h)}{t(a + 1)}$$

remain valid for this ordinary source angle. Thus a, b, t have residue one. The positive source identity

$$N = (1 + z)(b + r)h + a^2\{g + j + rt(1 + a^2z)\}$$

shows that $N > 0$ is infinitesimal and $h/N, r/N, g/N, j/N$ are bounded: each of the four positive coefficients in this sum is a bounded unit.

Here is the entire normalized coefficient calculation. Write $z_0 = \overline{z} > 0$. Substitution of $a = b = t = 1, r = 0$ in the bounded source numerators and unit denominators gives

$$\overline{\Delta} = 2(1 + z_0)^2, \qquad \overline{C/N} = \frac{1}{4(1 + z_0)^2}, \qquad \overline{\alpha/C} = z_0^2,$$
$$\overline{rn} = \overline{hA} = \frac{1}{4(1 + z_0)}, \qquad \overline{gm} = \overline{jD} = \frac{1}{4}. \tag{116.3}$$

For example, the exact quotients used in the first line are

$$\frac{C}{N} = \frac{1 + zr}{2a^2t\Delta}, \qquad \frac{\alpha}{C} = \frac{z^2(b + r)a^2t}{(1 - zh)(1 + zr)}.$$

The four strong normalized products use the numerators of (119.1) after multiplication by r, h, g, j, respectively; no vanishing source gap is inverted in this calculation. All six weights are therefore positive, C, α are infinitesimal, and their normalized quotients displayed above are bounded units. For instance

$$\frac{1}{Cn} = \frac{r/N}{(C/N)(rn)}, \qquad \frac{1}{\alpha n} = \frac{C/\alpha}{Cn};$$

the other six product bounds follow in the same way from $g/N, h/N, j/N$. These are exactly the hypotheses of Theorem 116.1.

For completeness, the source is physical by the unchanged exact contact argument. The rational stationary identities (119.5) are valid at this value of c as well. At that noncommuting contact the state matrix has a nonzero kernel vector. Its restriction to $Z = 0$ is positive definite: the signed core is nonnegative there and the positive event packets have zero common kernel. Hence the other three eigenvalues are positive. The exact contact and angular stationarity give the stationary zero of the Laurent determinant; its constant-plus-positive-exponentials form makes this its global minimum. The determinant criterion and homogeneous continuity prove physicality, exactly as for the small-angle source. Theorem 116.1 now applies.

This corollary concerns the stated radial subfamily of the ordinary source-angle cell. It does not assert that every ordinary-angle C2 source has the required matching strengths or enters this subfamily. □

Chapter 117

All outer tests for the all-small C2 radial family

Component `pc7dacaa3258e`
Source: `CORE/prior/inherited/c2_radial_all_tests.tex`
SHA256: `c7dacaa3258ed63c64024d672c622717ee267fefaebc871a4becf6f9f789cce5`
Inventory use: Complete four-strong/weak-pivot outer theorem and radial-source corollary; the broader unit-N/bounded-shape final corollary is not imported

117.1 All outer tests for the all-small C2 radial family

Scope of retained extract. This component retains the complete four-strong outer theorem and the radial-source corollary with both proofs. The final broader bounded-shape corollary, whose unit-N branch is not used in the current source routing, is outside the mathematical scope used in this rederivation and is omitted here. The entire original file and the exact omitted span remain in the source archive and presentation manifest.

The radial corner has four strong event packets and one weak packet. The weak packet must be retained at its exact scale at the opposite commuting corner. An elementary conductor identity makes that scale sufficient; it does not compare coefficient and testing infinitesimals. Throughout this section C denotes the event weight e_3, whereas c continues to denote the source half-angle in (119.1).

Lemma 117.1 (Four strong packets and a weak opposite-corner pivot)**.** *Consider the full parent* (119.4)*, with*

$$(e_1, e_2, e_3, e_4, e_5) = (n, m, C, A, D), \qquad \alpha \geq 0.$$

Suppose $C > 0$ is infinitesimal and the four positive pure coefficient quantities Cn, Cm, CA, CD have bounded reciprocals. Then the full parent is tensor SOS at every commuting residue test outside the finite zero-zero corner, for every real or complex projective state. The four strong coefficients and the roots in the proof may be unbounded.

Proof. The hypotheses imply that $1/n, 1/m, 1/A, 1/D$ are infinitesimal. First consider the outer axes other than the infinity-infinity corner. Choose $\mu = \min(1, n, m, A, D)$, a positive bounded unit. Retain the nonnegative excess of each strong packet, together with the whole C and α packets. The remaining parent consists of the signed core and the four packets with common weight μ.

Here are the four kernel calculations, with positive factors $1 + u^2$ or $1 + v^2$ omitted only from the packet amplitudes. At $u = 0$, finite nonzero v, the four amplitudes are $(X, vW - X, Y, -Y - vZ)$; at $v = 0$, finite nonzero u, they are $(X, -X - uZ, Y, uW - Y)$. Their common kernels are zero, and half the signed core is, respectively,

$$D_v(X^2 + Y^2), \qquad (uW - Y)^2 + (X + uZ)^2.$$

At $u=\infty$, finite v, the four homogenized amplitudes are $(X, vY-Z, Y, W+vX)$; at $v=\infty$, finite u, they are $(X, W+uY, Y, uX-Z)$. These common kernels are zero even when the finite angle is zero. Half the homogenized signed core is, respectively,

$$(W+vX)^2+(vY-Z)^2, \qquad D_u(X^2+Y^2).$$

Thus the selected state Gram has strict positive residue. Its coefficients are bounded in each indicated local testing chart, so the strict-residue Gram lemma proves the whole chart. This includes the mixed zero-infinity endpoints and either sign of every finite angle.

It remains to treat $u=v=\infty$ at residue. Put $p=1/u$, $q=1/v$, using the homogeneous reciprocal coordinates at exact infinity, and write $D_p=1+p^2$, $D_q=1+q^2$. Thus p,q are infinitesimal or zero. The homogeneous full parent in this chart is exactly

$$\begin{aligned}
\widehat{F} &= 4nD_pD_qX^2+4mM^2+4CD_q(W-pY)^2+4AD_pD_qY^2+4DJ^2+4\alpha D_p(Y-qZ)^2+H,\\
M &= pW-pqX+Y-qZ, \qquad J=qW+X-pqY-pZ,\\
H &= 2\{J^2+D_p(Y-qZ)^2+D_q(pX+Z)^2-D_pD_qZ^2\}. \qquad (117.1)
\end{aligned}$$

For finite nonzero u,v this is p^2q^2F; homogeneous interpretation also gives the exact endpoint polynomial.

Set $L=1+p^2+p^2q^2$. The needed conductor identity is

$$\frac{H}{2} = qWJ-qZM+L(X^2+Y^2)-2pqXY+q(W+pqZ)X-q(pqW+Z)Y. \qquad (117.2)$$

For a direct check, put $M_0=pW-qZ$, $J_0=qW-pZ$. The scalar part of $H/2$ is

$$J_0^2+(q^2-p^2)Z^2 = qWJ_0-qZM_0.$$

Its linear X,Y part is $2q(W+pqZ)X-2q(pqW+Z)Y$, and its quadratic part is $L(X^2+Y^2)-2pqXY$. Substitution of $M=M_0-pqX+Y$, $J=J_0+X-pqY$ gives (117.2).

First normalize $W=1$ and assume Z bounded. No bound is required on X or Y. Retain the α packet and use the five roots

$$K_1=\sqrt{n}X, \quad K_2=\sqrt{m}M, \quad K_4=\sqrt{A}Y, \quad K_5=\sqrt{D}J, \quad T_C=\sqrt{C}\,W.$$

Here $T_C=\sqrt{C}$ in this state chart. Equations (117.1)–(117.2) give the following *exact* Gram expression for the remainder:

$$\begin{aligned}
\widehat{F}-4\alpha D_p(Y-qZ)^2 &= 4D_pD_q(K_1^2+K_4^2)+4K_2^2+4K_5^2+4D_q\left(T_C-p\sqrt{C/A}\,K_4\right)^2\\
&\quad +2L\left(\frac{K_1^2}{n}+\frac{K_4^2}{A}\right)-\frac{4pq}{\sqrt{nA}}K_1K_4\\
&\quad +2qT_C\left\{\frac{K_5}{\sqrt{CD}}-\frac{ZK_2}{\sqrt{Cm}}+\frac{(1+pqZ)K_1}{\sqrt{Cn}}-\frac{(pq+Z)K_4}{\sqrt{CA}}\right\}. \qquad (117.3)
\end{aligned}$$

Every denominator is pure coefficient. The four inverse square roots $1/\sqrt{Ce_i}$ are bounded by hypothesis, and $\sqrt{C/A}=C/\sqrt{CA}$ is infinitesimal. All coefficients of this Gram are therefore in the bounded tensor ring. Every displayed correction has zero residue, whereas its five positive packet pivots have residue 4. Thus its residue is exactly

$$4(K_1^2+K_2^2+K_4^2+K_5^2+T_C^2)\succ 0.$$

The strict-residue Gram lemma applies without any bound on the roots. In particular it makes no comparison between C and p or q.

The other state chart is $Z = 1$, W infinitesimal or zero. Here the exact pure-testing conductor is a unit:

$$q = \Gamma := \frac{M + WJ + (pq - W)X + (Wpq - 1)Y}{W^2 - 1}. \tag{117.4}$$

Indeed $M + WJ = q(W^2 - 1) + (W - pq)X + (1 - Wpq)Y$. Substitution in (117.2) gives

$$\frac{H}{2} = L(X^2 + Y^2) - 2pqXY + \Gamma\{WJ - M + (W + pq)X - (pqW + 1)Y\}. \tag{117.5}$$

This is an exact quadratic form on X, M, Y, J, with bounded pure-testing coefficients. Retain the whole C and α packets separately. Scale the four remaining roots as before. Every coefficient of the scaled core has zero residue, because each contains two inverse strong packet square roots. Adding the four strong packets consequently produces the strict residue $4(K_1^2 + K_2^2 + K_4^2 + K_5^2)$. Again only the Gram coefficients, not X, Y or the packet roots, needed boundedness.

These two state charts are exhaustive when $(W, Z) \neq (0, 0)$: if $W \neq 0$ and Z/W is bounded use the first; otherwise $Z \neq 0$ and W/Z is infinitesimal, giving the second. Normalization and restoration use pure-testing factors. When $W = Z = 0$, (117.2) is just the bounded quadratic form $H/2 = L(X^2 + Y^2) - 2pqXY$; after the same four-root scaling its coefficients have zero residue, so the four retained strong packets again give a strict Gram. This includes the zero state. Splitting a complex state into real and imaginary parts completes the proof. □

Corollary 117.2 (The radial source has no missing outer commuting chart)**.** *Use the unchanged exact source* (119.1) *under the hypotheses of Theorem 118.1: h, r, g, j are positive infinitesimals and $a, 1 - r, t$ are positive bounded units. Then the full parent is tensor SOS at every commuting residue test and every real or complex state. Together with the physical-source noncommuting-residue theorem, this proves all tests for this exact radial source family.*

Proof. The source identities and positive decomposition of N proved in Section 118.1 give

$$N \asymp h + r + g + j, \qquad U_1 = re_1,\ U_2 = ge_2,\ U_4 = he_4,\ U_5 = je_5 > 0 \quad \text{bounded units},$$

where $N/(h + r + g + j)$ has residue 1. Moreover, directly from (119.1),

$$\frac{C}{N} = \frac{A_c}{2a^2 t\Delta} > 0 \quad \text{is a bounded unit.}$$

For example,

$$\frac{1}{Ce_1} = \frac{r/N}{(C/N)U_1},$$

and the three analogous formulas use $g/N, h/N, j/N$. Thus all four product reciprocals in Lemma 117.1 are bounded, while C is positive infinitesimal. That lemma covers every outer commuting chart, and Theorem 118.1 covers the zero-zero chart.

The final noncommuting join uses only the already-established physicality of the exact source and the physical-source noncommuting-residue theorem. In particular the exact contact and Laurent-determinant physicality argument used for the unit-N source requires $N > 0$, not that N have positive residue. All five event weights are positive. On the three-dimensional state subspace $Z = 0$ the signed core is nonnegative, and the positive event packets have zero common kernel at the noncommuting contact; hence the state restriction is positive definite. The exact stationary state is a nonzero kernel vector, so the other three state eigenvalues are positive. Determinant stationarity remains exact, and the same Laurent-determinant argument proves physicality. No coefficient-entry claim is made here for C2 sources outside the stated radial family. □

Chapter 118

The last bounded-shape C2 corner: a weighted radial root

Component `p163e0ebf59f6`
Source: `CORE/prior/inherited/c2_radial_corner.tex`
SHA256: `163e0ebf59f6150a66255b45f6633a7b5acb96cefb52013d74b7bac54eecf672`
Inventory use: Complete all-small source-ideal and weighted-radial correction theorem; complete normal/cross/radial coefficients and both projective states.

118.1 The last bounded-shape C2 corner: a weighted radial root

It remains to treat the corner in which h, r, g, j are all positive infinitesimals. The radial coefficient then tends to zero, but the positive source identity gives exactly the weight needed to normalize it.

Theorem 118.1 (The all-small radial corner)**.** *Use the unchanged source* (119.1)*. Suppose that* $c > 0$ *and* $h = b - a^2 t, r, g, j > 0$ *are infinitesimal, while* $a, 1 - r, t$ *are positive bounded units. For every independent infinitesimal or zero testing pair* u, v *and every real or complex projective state, the full parent is a finite sum of squares in* $K_c \otimes_{\mathbb{R}} K_t$*.*

118.1.1 The small source ideal controls all coefficients

Write $z = c^2$ and $D_c = 1 + c^2$. The source identities give

$$b - 1 = -r + g - zh, \qquad t - 1 = zr - j, \qquad a - 1 = \frac{g + j - D_c(r + h)}{t(a + 1)}. \tag{118.1}$$

Thus a, b, t all have residue 1. Let $I = (h, r, g, j)$ in the ring of bounded source coefficients used below. By (120.2), $N > 0$ and $N/(h + r + g + j)$ has residue 1. In particular

$$h/N, \quad r/N, \quad g/N, \quad j/N \text{ are bounded}, \qquad 4\alpha/(c^4 N) \text{ has residue 1.} \tag{118.2}$$

All rational coefficient functions in the retained remainder have denominators with nonzero residue: the possibly small denominators r, g, j, h have been absorbed into U_1, U_2, U_5, U_4. Specialize a bounded rational coefficient by putting $h = r = g = j = 0$ and $a = b = t = 1$, leaving c untouched. Its difference from this specialization lies in I, with bounded coquotients. Indeed finite polynomial subtraction factors the differences by $r, g, j, a - 1$, and (118.1) replaces $a - 1$ by bounded multiples of the four generators. Subtraction of fractions only uses unit denominators. The monic polynomial divisions in the preceding proofs preserve this property. Pure-testing denominators D_u, D_v do so as well. Consequently every coefficient in I, divided by the pure coefficient N, remains bounded. This observation involves no comparison with the independent testing infinitesimals.

118.1.2 An explicit additional zero correction

Use the four retained packets and six-root Gram from Section 120.1, including its zero correction (120.9). That construction required only that e_3 be bounded; it still gives an exact bounded Gram here. Two of its current coefficients simplify exactly to

$$Q(u^2) = h - a^2 r u^2, \qquad Q(u^2) + a^2 D_u r = h + a^2 r. \tag{118.3}$$

Both belong to I. The coefficient of T in the exact relation ζ is therefore $u(h + a^2 r)$.

We record the one base calculation needed for the new correction. On the scalar normal graph, after specialization modulo I, put

$$Y_* = uW - v, \qquad G_* = D_u v W - D_v u.$$

The entire bounded scalar remainder becomes

$$f_* = 2uG_* + \frac{2}{D_c} D_u D_v u Y_* + 2\{D_u(vY_* - 1)^2 + D_v u^2 - D_u D_v\}. \tag{118.4}$$

Here $U_2 = 1/4$, $U_4 = 1/(4D_c)$; all terms containing a generator of I, e_3, or α vanish. Direct differentiation gives

$$\begin{gathered} f_*(u,u,1) = 0, \\ (f_*)_W(u,u,1) = \frac{2u^2(u^2-c^2)D_u}{D_c}, \qquad (f_*)_v(u,u,1) = -\frac{2u(u^2-c^2)D_u}{D_c}. \end{gathered} \tag{118.5}$$

Use the division order of the preceding proof: first restrict to $v = au$, then expand at W_c, and retain the finite second-order remainder in $v - au$ as an R^2 coefficient. The first two derivatives in (118.5) imply that the ST coefficient is $2uD_u/D_c$ and the direct RT coefficient is $2uD_u/D_c$. Modulo I, elimination of S reads

$$S = \frac{H_2 + (1 - u^2 v^2)H_1 + uvH_5 + (D_u W - u(v+u))R}{D_u}.$$

Hence the full scalar RT coefficient after elimination is exactly

$$\frac{2u}{D_c}(D_u W + 1 - uv). \tag{118.6}$$

The earlier zero correction adds no RT term at this specialization, because (118.3) vanishes there. Meanwhile the coefficient of R in ζ specializes to $-(D_u W + 1 - uv)$. Add one more exact zero:

$$\frac{2u}{D_c}\,\zeta T = 0. \tag{118.7}$$

By (118.6), the corrected RT coefficient now lies in I. All radial cross coefficients still contain an explicit factor u.

118.1.3 Weighted boundedness and the strict residue

Replace the radial root by the pure-coefficient rescaling

$$T_N = \sqrt{N}\,(u^2 - c^2).$$

We check every type of coefficient of the resulting Gram. The hard block and the hard–R block were already bounded. A hard–T coefficient contains $u/\sqrt{e_i}$, times bounded factors, for $i = 1, 2, 4, 5$: this follows from the odd-parity normal division, the pure current elimination, and the two displayed zero corrections. Dividing it by $\sqrt{N}$ remains bounded, with zero residue, because

$$\frac{1}{\sqrt{Ne_1}} = \frac{\sqrt{r/N}}{\sqrt{U_1}}, \quad \frac{1}{\sqrt{Ne_2}} = \frac{\sqrt{g/N}}{\sqrt{U_2}}, \quad \frac{1}{\sqrt{Ne_4}} = \frac{\sqrt{h/N}}{\sqrt{U_4}}, \quad \frac{1}{\sqrt{Ne_5}} = \frac{\sqrt{j/N}}{\sqrt{U_5}}.$$

The corrected RT coefficient lies in I and contains u, so its quotient by $\sqrt{N}$ is bounded and has zero residue.

For the T^2 coefficient, the graph identity $f_*(u,u,1)=0$ says that its monic scalar quotient $C_2(u^2)$ vanishes modulo I. Every other T^2 contribution contains Q or $h+a^2r$, hence belongs to I too. Its quotient by N is therefore bounded. All such extra terms have an explicit factor u^2. At $u=0$ the scalar quotient is exactly $4\alpha/c^4$, so (118.2) makes the weighted radial pivot equal to 1 at residue. In particular the new correction adds $2u^2(h+a^2r)/(D_cN)$ to that coefficient, which is bounded and has zero residue.

The residue of the final Gram, in the roots (K_1,K_2,K_4,K_5,R,T_N), is therefore

$$4K_1^2+4K_2^2+4K_4^2+4K_5^2+(1+\overline{W})^2R^2+T_N^2\succ 0. \tag{118.8}$$

Here the relation ζ has residue $-(1+\overline{W})R$, since all four inverse packet scales are infinitesimal. The strict-residue Gram lemma proves the full bounded $Z=1, W\geq 0$ chart. The $W=1, Z$ infinitesimal proof of the preceding section also applies unchanged: it uses boundedness, not a positive residue, of e_3. Its four packet roots have strict residue. Sign dispatch and homogeneous restoration therefore prove the theorem for all projective states.

Corollary 118.2 (Complete bounded-shape C2 source family)**.** *In the unchanged C2 source let $c>0$ be infinitesimal, $r,g,j>0$ bounded, and $a,1-r,t$ positive bounded units. Assume only $h=b-a^2t>0$. Then every independent zero-zero test and every projective state have tensor-SOS parent.*

Proof. The gap h is bounded. If it is a unit, (120.2) makes N a unit, so Theorem 119.1 applies. If h is infinitesimal and one of r,g,j is a unit, use Theorem 120.1. If all four are infinitesimal, use Theorem 118.1. These alternatives are exhaustive. □

Chapter 119

The C2 boundary: one uniform five-root proof

Component `p9f1235511111`
Source: `CORE/prior/inherited/c2_unified.tex`
SHA256: `9f1235511111e57d1ccceec8476e65f1973266a17f78189ee10c9e3bf015e2f8`
Inventory use: Only exact source/complete parent, centered packet differences, whole scalar/both normal parity and monic division, current elimination, reference Gram computation and complete state/sign join; its unit-N theorem is not imported on the radial domain.

119.1 The C2 boundary: one uniform five-root proof

The argument below proves the entire original parent for one explicit C2 source domain and every independent zero-zero test. Three source parameters may approach their boundary at arbitrary relative rates. Exact contact, parity, and a retained current replace polynomial tables.

119.1.1 The unchanged source and its domain

Let K_c, K_t be independent real closed extensions of $\mathbb{R}$. In the coefficient field take $c > 0$ infinitesimal and $a, r, g, j > 0$ bounded. Define

$$z = c^2, \quad k = 1 + z, \quad t = 1 + zr - j, \quad B = 1 - r + g + a^2 zt, \quad b = B/k, \quad A_c = 1 + zr,$$

and define

$$\begin{aligned}\Delta &= (1+z)zb^2 + 2z(1+z)br + (1+z)(1+a^2z)b \\ &\quad + z^2(1-a^2)r^2 + 2z(1+a^2z)r + 1 + a^2z, \\ N &= (1+z)(b+r)(b+a^2(1-t)) - a^2(1+a^2z)(1-r)t.\end{aligned}$$

Take $d = 1$, $\lambda = \beta = 0$, and

$$\begin{aligned} e_1 &= \frac{(b+r)A_c(A_c+a^2zt) - r\Delta}{2r\Delta}, & e_2 &= \frac{(1+a^2z)(1-r)A_c(A_c+a^2zt)}{2g\Delta}, \\ e_3 &= \frac{A_cN}{2a^2t\Delta}, & e_4 &= \frac{a^2A_c(b+r)(A_c+a^2zt)}{2(b-a^2t)\Delta}, \\ e_5 &= \frac{(1+z)(b+r)(b-a^2zr)(A_c+a^2zt)}{2a^2j\Delta}, & \alpha &= \frac{z^2(b+r)N}{2(1-z(b-a^2t))\Delta}. \end{aligned} \tag{119.1}$$

Require the following unit conditions:

$$\begin{gathered} c > 0 \text{ is infinitesimal,} \\ r, g, j > 0 \text{ are bounded,} \\ a, \quad 1-r, \quad t, \quad b-a^2t, \quad N \text{ are positive bounded units.} \end{gathered} \tag{119.2}$$

Any of r, g, j may have zero or positive residue; no relative-rate comparison is assumed.

The quantities b, Δ are positive bounded units. The source formulas show that

$$U_1 = re_1, \qquad U_2 = ge_2, \qquad U_5 = je_5, \qquad e_3, e_4, \qquad \alpha/c^4 \tag{119.3}$$

are positive bounded units. For example, after setting $c = 0$ their first three values are

$$\frac{b_0(1-r)}{2(b_0+1)}, \qquad \frac{1-r}{2(b_0+1)}, \qquad \frac{b_0(b_0+r)}{2a^2(b_0+1)}, \qquad b_0 = 1-r+g.$$

Consequently e_1, e_2, e_5 have positive bounded reciprocals, whether or not the coefficients themselves are bounded. All six event coefficients are positive.

Theorem 119.1 (Uniform C2 unit-gap zero-zero theorem)**.** *Assume* (119.2)*. For every independent testing pair $u, v \in K_t$ that is infinitesimal or zero, and every independent real or complex projective state, the entire parent* (119.4) *is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$.*

119.1.2 Full parent, contact, and retained squares

Write $D_u = 1 + u^2, D_v = 1 + v^2$, and

$$R_2 = vW - X + uvY - uZ, \qquad R_3 = uW - Y,$$
$$R_5 = uW + uvX - Y - vZ, \qquad R_\alpha = vY - Z, \qquad R_\beta = X + uZ.$$

The full homogeneous parent is

$$\begin{aligned} F = {} & 4e_1 D_u D_v X^2 + 4e_2 R_2^2 + 4e_3 D_v R_3^2 + 4e_4 D_u D_v Y^2 + 4e_5 R_5^2 + 4\alpha D_u R_\alpha^2 \\ & + 2(R_5^2 + D_u R_\alpha^2 + D_v R_\beta^2 - D_u D_v Z^2). \end{aligned} \tag{119.4}$$

Its exact coefficient-field stationary zero is

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(c, ac; \frac{b}{a}, -cr, c\left(\frac{b}{a} - at\right), 1\right). \tag{119.5}$$

Substitution of (119.1) in the four state derivatives and two angle derivatives of (119.4) gives zero. These are six rational identities: one can check them before substituting for b by using $g = (1+c^2)b - (1 - r + a^2c^2t)$ and $j = 1 + c^2r - t$. The four state identities imply $F = 0$ by homogeneity. Thus this contact statement is a finite direct calculation, without an asymptotic premise.

First work in the chart $Z = 1$, $W \geq 0$ bounded. Center three event amplitudes:

$$H_1 = X + ru, \qquad H_2 = R_2 - gu, \qquad H_5 = R_5 + jv.$$

Put $n_x = H_1, n_y = H_5$. Then, exactly,

$$X = -ru + n_x, \qquad Y = uW - (1 - j + ru^2)v + uvn_x - n_y, \tag{119.6}$$
$$H_2 = G - (1 - u^2v^2)n_x - uvn_y, \qquad G = D_u vW - [1 - r + g + (1 - j + ru^2)v^2]u. \tag{119.7}$$

Define the *entire* remainder

$$\mathcal{R} = F - 4e_1 D_u D_v H_1^2 - 4e_2 H_2^2 - 4e_5 H_5^2. \tag{119.8}$$

It has bounded coefficient-field coefficients. Indeed the three packet differences are

$$-8re_1 D_u D_v u H_1 + 4r^2 e_1 D_u D_v u^2,$$
$$8ge_2 u H_2 + 4g^2 e_2 u^2, \qquad -8je_5 v H_5 + 4j^2 e_5 v^2,$$

and (119.3) bounds their coefficients. All remaining packets already have bounded coefficients. Quadratic state degree gives the exhaustive expansion

$$\mathcal{R} = f + L_x n_x + L_y n_y + Q_{xx} n_x^2 + Q_{xy} n_x n_y + Q_{yy} n_y^2, \tag{119.9}$$

where all six coefficients are polynomials in u, v, W with bounded coefficient-field coefficients. There is no higher-normal remainder.

119.1.3 Exact parity and Taylor division

The three soft roots will be

$$R = au - v, \qquad S = u(W - W_c), \qquad T = u^2 - c^2. \tag{119.10}$$

We prove, with bounded polynomial coefficients,

$$f \in (R, S, T)^2, \qquad L_x, L_y \in (R, S, T). \tag{119.11}$$

The parent and its centered substitutions are invariant under $(u, v, n_x, n_y) \mapsto (-u, -v, -n_x, -n_y)$ with W fixed. Thus $f(u, au, W) = f_0(u^2, W)$ is even in u, whereas $f_v(u, au, W)$ and $L_i(u, au, W)$ are odd. The polynomial $f_0(\xi, W)$ has degree at most two in W. At $u = v = n_x = n_y = 0$, the parent is 4α, independent of W. Hence the nonconstant W coefficients of f_0 are divisible by ξ.

At (119.5) every H_i vanishes. Consequently f and all its first derivatives vanish there, as do both L_i. Write

$$f_0(\xi, W) = A(\xi)(W - W_c)^2 + B_1(\xi)(W - W_c) + C_1(\xi).$$

Then $\xi \mid A, B_1$, $B_1(c^2) = 0$, and $C_1(c^2) = C_1'(c^2) = 0$. Since $c \neq 0$, there are polynomials with bounded coefficients such that

$$A = \xi A_1, \qquad B_1 = \xi(\xi - c^2)B_2, \qquad C_1 = (\xi - c^2)^2 C_2.$$

All divisions here are by a monomial or a monic polynomial. Synthetic division uses only powers of the bounded coefficient c^2; in particular no quotient contains $1/c$. It follows that

$$f(u, au, W) = A_1 S^2 + uB_2 ST + C_2 T^2.$$

Next write $f_v(u, au, W) = u\ell(u^2, W)$. Contact stationarity gives $\ell(c^2, W_c) = 0$. Subtracting the value at W_c and dividing that value by $\xi - c^2$ gives

$$f_v(u, au, W) = S\ell_1 + uT\ell_2$$

with bounded coefficients. Exact finite Taylor expansion in $v = au - R$ proves the scalar assertion of (119.11). The same odd-parity argument applies to each $L_i(u, au, W)$, and finite division by $v - au$ proves the normal assertions. This accounts for all scalar, first-normal, and quadratic-normal terms in the original parent.

119.1.4 Eliminate one soft root by a pure tensor unit

Put $K = 1 - j + ru^2$ and

$$Q(\xi) = aW_c - a^2(1 - j) - a^2 r(\xi + c^2).$$

The source definition of b gives the exact identity

$$G = -[D_u W - Ku(v + au)]R + aD_u S + uTQ(u^2). \tag{119.12}$$

Indeed $(\xi - c^2)Q(\xi)$ equals $a(1 + \xi)W_c - (1 - r + g) - a^2(1 - j + r\xi)\xi$. Together with (119.7), this yields

$$S = \frac{H_2 + (1 - u^2 v^2)H_1 + uvH_5 + [D_u W - Ku(v + au)]R - uTQ(u^2)}{aD_u}. \tag{119.13}$$

The denominator is a product of a pure coefficient unit and a pure testing unit. Thus this is an allowed exact substitution in the tensor ring. Take the five roots

$$K_1 = \sqrt{e_1}H_1, \quad K_2 = \sqrt{e_2}H_2, \quad K_5 = \sqrt{e_5}H_5, \quad R, \quad T. \tag{119.14}$$

All inverse square roots used to substitute for the H_i are bounded pure coefficient quantities. Adding the three retained packets back therefore gives a bounded exact Gram matrix for the entire parent on (119.14).

119.1.5 A uniform positive residue calculation

Here is a calculation using only small quadratic forms. It proves strictness even when any of the first three source coefficients diverges. Set $c = 0$ formally in the bounded source formulas, and put

$$b_0 = 1 - r + g, \qquad t_0 = 1 - j, \qquad w_0 = b_0/a,$$
$$N_0 = (b_0 + r)(b_0 + a^2(1 - t_0)) - a^2(1 - r)t_0.$$

Define pure coefficient quantities

$$n = \frac{b_0(1-r)}{2r(b_0+1)}, \qquad m = \frac{1-r}{2g(b_0+1)}, \qquad C = \frac{N_0}{2a^2t_0(b_0+1)},$$
$$A = \frac{a^2(b_0+r)}{2(b_0-a^2t_0)(b_0+1)}, \qquad D = \frac{b_0(b_0+r)}{2a^2j(b_0+1)}.$$

The letters n, m, C, A, D in this calculation denote these reference weights, not additional assumptions or altered source coefficients. The quantities C, A, w_0 are positive bounded units; n, m, D and their reciprocals are positive, with the reciprocals bounded.

The quadratic angular and hard-state part of the parent at $c = 0$ is

$$F_2 = 4nX^2 + 4m(vW - X - u)^2 + 4C(uW - Y)^2 + 4AY^2 + 4D(uW - Y - v)^2 + 2\big((uW - Y)^2 - 2uvW + X^2 + 2uX + u^2\big). \tag{119.15}$$

Write

$$h_x = 4n + 4m + 2, \qquad h_y = 4C + 4A + 4D + 2,$$
$$p = \frac{(4C + 4D + 2)4A}{h_y}, \quad q = \frac{4m(4n+2)}{h_x}.$$

Completing the two hard squares and using the source identities gives

$$F_2 = h_x\left[n_x + \frac{4m}{h_x}(WR - aS)\right]^2 + h_y\left[-n_y + \frac{4A}{h_y}(S + w_0R/a)\right]^2 + p(S + w_0R/a)^2 + q(WR - aS)^2. \tag{119.16}$$

Here the linearized centered substitutions are $X = -ru + n_x$, $Y = uW - t_0v - n_y$, $R = au - v$, and $S = u(W - w_0)$. This identity can be checked by two square completions and the following five elementary rational identities:

$$\frac{(4m+2)4n}{h_x} = qa^2w_0^2, \qquad \frac{4D(4C+4A+2)}{h_y} = pw_0^2/a^2,$$
$$4m + 4D + 2 - \frac{(4m+2)4m}{h_x} - \frac{(4C+4D+2)4D}{h_y} = w_0(p/a + qa),$$
$$(4m+2) - 4maw_0 = rh_x, \qquad \frac{4A}{h_y}w_0 + \left(\frac{4D}{h_y} - t_0\right)a = 0.$$

Their denominators are explicit pure coefficient factors, and direct substitution of the five displayed weights proves each identity.

At residue level the exact current relation says $H_2 = -WR + aS - n_x$. Substituting $S = (H_2 + n_x + WR)/a$ in (119.16) cancels the hard-completion denominators and gives the particularly simple form

$$(4n+2)n_x^2 + 4mH_2^2 + (4C + 4D + 2)n_y^2 + 4A\left(\frac{(W + w_0)R + n_x + H_2}{a} - n_y\right)^2. \tag{119.17}$$

Scale the reference roots by $\sqrt{n}, \sqrt{m}, \sqrt{D}$ and denote the resulting abstract Gram coordinates by K_1, K_2, K_5. The reference scaled Gram is

$$\mathcal{E} = (4 + 2/n)K_1^2 + 4K_2^2 + \big(4 + (4C+2)/D\big)K_5^2$$

$$+\frac{4A}{a^2}\left[(W+w_0)R+\frac{K_1}{\sqrt{n}}+\frac{K_2}{\sqrt{m}}-\frac{aK_5}{\sqrt{D}}\right]^2. \tag{119.18}$$

Every coefficient of this Gram matrix is bounded, including when r, g, j have zero residue. Its residue is strictly positive definite: the first three square coefficients are at least 4, and after those roots vanish the last square has strictly positive coefficient on R because A, a, w_0 are positive units and $W \geq 0$.

We justify carefully that this is the residue of the exact Gram just constructed. First $re_1 - rn$, $ge_2 - gm$, and $je_5 - jD$ are infinitesimal, as are $e_3 - C$, $e_4 - A$, and $W_c - w_0$. Moreover $e_1/n = U_1/(rn)$, $e_2/m = U_2/(gm)$, and $e_5/D = U_5/(jD)$ are positive bounded units with residue 1. Thus the reference root scaling and the actual scaling in (119.14) induce the same residue coordinates. After retention and scaling, every coefficient involving the possibly unbounded packets is a polynomial combination of bounded quantities such as

$$U_1, \quad rU_1, \quad \frac{1}{\sqrt{e_1}} = \frac{\sqrt{r}}{\sqrt{U_1}}, \quad \frac{U_1}{\sqrt{e_1}} = \sqrt{r}\sqrt{U_1},$$

and the corresponding quantities for g, j. All square-root denominators here have positive residue. Their residues thus agree with those obtained using rn, gm, jD. Monic division, followed by (119.13), also commutes with residue, since its only further denominator is aD_u. Hence the four-root block is exactly the residue of (119.18).

The radial pivot has an equally short exact computation. On the scalar graph $v = au, W = W_c$ the remainder is $(u^2 - c^2)^2 C_2(u^2)$. At $u = 0$ it equals 4α; hence $C_2(0) = 4\alpha/c^4$. The latter is a positive bounded unit by (119.3). All radial cross coefficients contain an explicit factor u, by the parity division above; the elimination (119.13) adds only further such factors. Thus the entire five-root residue is

$$\overline{\mathcal{E}} + 4\overline{\alpha/c^4}\, T^2 \succ 0. \tag{119.19}$$

The strict-residue Gram lemma proves the full bounded $Z = 1, W \geq 0$ chart, without any restriction on the size of X, Y or of the roots.

119.1.6 The other state chart and endpoint dispatch

For $W = 1$, Z infinitesimal, retain the α packet and the three original squares in X, R_2, R_5. The exact identities

$$\begin{aligned} Y &= uvX - R_5 + u - vZ, \qquad R_3 = -uvX + R_5 + vZ, \\ v &= \frac{R_2 + (1 - u^2v^2)X + uD_vZ + uvR_5}{D_u} \end{aligned} \tag{119.20}$$

express the entire bounded remainder as a bounded quadratic form on X, R_2, R_5, u. For clarity, this does not require a division by any root: D_u is pure testing. The exact signed-core completion

$$\begin{aligned} H &= 2\left[C_*^2 + D_vQ_*^2 + D_uv^2Y^2 + \frac{D_uu^2v^2(1-u^2v^2)}{D_v}Z^2\right] - 4D_uuvZ, \\ C_* &= u + uvR_\beta - Y, \qquad Q_* = R_\beta - \frac{D_uuv^2}{D_v}Z \end{aligned}$$

shows quadraticity directly after (119.20): every displayed amplitude is linear in the four roots, and the last indefinite term is u times the same linear expression for v. All coefficients are bounded. Adding the three retained packets and using roots $K_1 = \sqrt{e_1}X$, $K_2 = \sqrt{e_2}R_2$, $K_5 = \sqrt{e_5}R_5$, and u, the residue is the residue of

$$(4 + 2/e_1)K_1^2 + 4K_2^2 + (4 + (4e_3 + 2)/e_5)K_5^2 + 4e_4\left(u - \frac{K_5}{\sqrt{e_5}}\right)^2.$$

It is strictly positive because e_4 is a positive bounded unit. This proves the whole chart, including arbitrary X, Y: only the Gram coefficients, not the roots, needed boundedness.

Finally the testing-only involutions

$$(u, W, X, Y, Z) \mapsto (-u, -W, -X, Y, Z), \qquad (v, W, X, Y, Z) \mapsto (-v, -W, X, -Y, Z)$$

preserve the full parent and allow $u, v \geq 0$. Put $L = 1 + v^2 + u^2 v^2$. The X, Y block of half the signed core has eigenvalues $L \pm uv > 0$, and its exact Schur complement is

$$\frac{uvD_uD_v[uv(W^2 + Z^2) - 2LWZ]}{L^2 - u^2v^2}. \tag{119.21}$$

If $WZ \leq 0$, the pure-testing core is nonnegative and is a square in K_t; every event is already a tensor square. This includes exact zero state or angle endpoints. If $WZ > 0$, either W/Z is bounded, giving the first chart, or Z/W is infinitesimal, giving the second. Restoring the state scale is pure testing multiplication. Splitting a complex state into real and imaginary parts completes the proof of Theorem 119.1.

Scope. This theorem includes all eight ordinary or infinitesimal combinations of r, g, j under (119.2); S38 is one special case. A vanishing residue of a, $1-r$, t, $b-a^2t$, or N requires an additional argument. Complementary testing-angle corners and entry of every C2 source into this domain are separate obligations. The proof uses no archived scalar or normal coefficient arrays.

Chapter 120

The next C2 boundary: retain the fourth event current

Component p53752031d16c
Source: CORE/prior/inherited/c2_y_current.tex
SHA256: 53752031d16c6691f061be1e7f5980fbd300cd3e60d5c03e599ed99f4179408a
Inventory use: Fourth centered current and bounded retained remainder, exact six-root zero relation/correction, full W-unit state identity. Its N-unit theorem is not imported on the radial domain.

120.1 The next C2 boundary: retain the fourth event current

The gap $h = b - a^2 t$ need not stay a unit. When it becomes infinitesimal, the fourth event is strong. Its centered current supplies an exact zero relation which repairs the preceding Gram matrix.

120.1.1 Source positivity and the precise remaining corner

Keep the exact source (119.1), with $c > 0$ infinitesimal, $r, g, j > 0$ bounded, and $a, 1 - r, t$ positive bounded units. This section assumes

$$h = b - a^2 t > 0 \text{ infinitesimal}, \qquad N \text{ a positive bounded unit.} \tag{120.1}$$

There is a useful exact positive identity:

$$N = (1 + c^2)(b + r)h + a^2\big[g + j + rt(1 + a^2 c^2)\big]. \tag{120.2}$$

It follows by writing $b + a^2(1 - t) = h + a^2$ in the definition of N and using the definitions of b, t. Thus all source coefficients are positive. Under the other unit hypotheses, N is a unit exactly when at least one of h, r, g, j has positive residue. In (120.1), therefore, at least one of r, g, j is ordinary. As before, re_1, ge_2, je_5, e_3 are positive bounded units, and now

$$U_4 = he_4 = \frac{a^2 A_c (b + r)(A_c + a^2 c^2 t)}{2\Delta}, \qquad \frac{\alpha}{c^4} = \frac{(b + r)N}{2(1 - c^2 h)\Delta} \tag{120.3}$$

are positive bounded units. Hence e_4 is strong and $1/\sqrt{e_4}$ is infinitesimal.

Theorem 120.1 (Full parent at the fourth-event boundary)**.** *For the exact source under* (120.1)*, every independent testing pair u, v that is infinitesimal or zero, and every real or complex projective state, the entire parent* (119.4) *is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$.*

120.1.2 The exact zero relation

First take $Z = 1$ and bounded $W \geq 0$. In addition to the three currents already defined, put

$$H_4 = Y - \frac{h}{a}u.$$

This current vanishes at the exact stationary contact, since $W_c - at = h/a$. Retain the extra packet $4e_4 D_u D_v H_4^2$. Its difference from the original fourth event is

$$4e_4 D_u D_v (Y^2 - H_4^2) = \frac{8U_4}{a} D_u D_v u H_4 + \frac{4hU_4}{a^2} D_u D_v u^2.$$

Thus the entire remainder after retaining four packets has bounded coefficients. Its exact contact, parity, quadratic state degree, and value 4α at $u = v = H_1 = H_5 = 0$ are unchanged. The complete monic-division argument in Section 119.1 therefore gives a bounded quadratic form on (H_1, H_5, R, S, T). Eliminate S by (119.13) and restore the four retained packets. This gives a bounded exact Gram on the six roots

$$K_1 = \sqrt{e_1} H_1, \quad K_2 = \sqrt{e_2} H_2, \quad K_4 = \sqrt{e_4} H_4, \quad K_5 = \sqrt{e_5} H_5, \quad R, \quad T. \tag{120.4}$$

There is one exact relation among these roots. With $K = 1 - j + ru^2$ and Q as in Section 119.1, direct substitution gives

$$H_4 = S + KR - aruT + uvH_1 - H_5.$$

Combining this with the current formula for S yields

$$\begin{aligned} \zeta = {} & aD_u H_4 - H_2 - (1 - u^2 v^2 + aD_u uv) H_1 + (aD_u - uv) H_5 \\ & - [D_u W + (a - uv)K] R + u[Q(u^2) + a^2 D_u r] T = 0. \end{aligned} \tag{120.5}$$

In the scaled roots every coefficient here is bounded. Its residue is

$$\overline{\zeta} = -\frac{K_2}{\sqrt{m}} - \frac{K_1}{\sqrt{n}} + \frac{aK_5}{\sqrt{D}} - (W + at_0) R, \tag{120.6}$$

where, here and in residue calculations below, coefficient expressions mean their residues; n, m, C, A, D, t_0, w_0 are the reference quantities defined in the preceding section. The K_4 and T coefficients vanish at residue. The coefficient of R is a nonzero real number, since $W \geq 0$ and a, t_0 are positive units.

120.1.3 The corrected residue is visibly strict

Let $h_0 = b_0 - a^2 t_0$. The definitions give

$$h_0 = (1 + c^2) h + a^2 c^2 r > 0, \tag{120.7}$$

so h_0 is infinitesimal too. The reference fourth weight satisfies $h_0 A > 0$ with unit residue, and $U_4 - h_0 A$ is infinitesimal. The ratio e_4/A need not have residue one. It is unnecessary: the bounded retained remainder depends on U_4, h, while the fourth scaled packet has coefficient $4D_u D_v$ and $1/\sqrt{e_4} = \sqrt{h}/\sqrt{U_4}$ has zero residue. These facts justify the reference comparison even when $h \ll c^2 r$.

For a transparent calculation, start from (119.17). The linearized fourth current is

$$H_{4,0} = S + t_0 R - H_5 = \frac{(W + at_0) R + H_1 + H_2 - aH_5}{a}.$$

The last term of that reference quadratic form is exactly

$$4A \left(H_{4,0} + \frac{h_0}{a^2} R \right)^2.$$

Subtract the fourth retained square $4AH_{4,0}^2$ and restore $4K_4^2$. The resulting four-hard-root part is

$$(4+2/n)K_1^2+4K_2^2+4K_4^2+(4+(4C+2)/D)K_5^2 \\ +\frac{8h_0A}{a^2}H_{4,0}R+\frac{4h_0^2A}{a^4}R^2.$$

At residue, the last square coefficient vanishes, $H_{4,0}=-\overline{\zeta}/a$, and h_0A has the same residue as U_4. The radial calculation is unchanged: its pivot is $4\overline{\alpha/c^4}>0$ and all radial cross coefficients vanish. Thus the entire Gram residue is the residue of

$$\mathcal{D}+4(\alpha/c^4)T^2-\frac{8U_4}{a^3}\zeta R,\quad \mathcal{D}=(4+2/n)K_1^2+4K_2^2+4K_4^2+\big(4+(4C+2)/D\big)K_5^2. \tag{120.8}$$

All these residue comparisons use bounded normalized coefficients and monic division, exactly as in the preceding proof.

Now add the following identically zero polynomial to the exact Gram:

$$\frac{8U_4}{a^3}\zeta R+\zeta^2=0. \tag{120.9}$$

Its coefficients are bounded, and the equality follows from the exact relation (120.5). The corrected residue is

$$\overline{\mathcal{D}}+4\overline{\alpha/c^4}\,T^2+\overline{\zeta}^{\,2}\succ 0.$$

Indeed the first four diagonal coefficients are at least 4, the radial pivot is positive, and after those five roots vanish $\overline{\zeta}^{\,2}$ has positive coefficient on R by (120.6). The strict-residue Gram lemma proves the full bounded $Z=1, W\geq 0$ chart.

120.1.4 The other projective chart

For $W=1$, Z infinitesimal, retain the original packets in X, R_2, Y, R_5 and the α packet. The exact pure-testing identity

$$u=\frac{D_uY+[Z(1-u^2v^2)-D_uuv]X+(D_u+uvZ)R_5+ZR_2}{D_u-D_vZ^2} \tag{120.10}$$

follows from the two amplitude definitions. Its denominator is a positive pure-testing unit. Together with (119.20), it expresses u, v linearly in X, R_2, Y, R_5, with bounded testing coefficients. The signed-core completion in the preceding section then makes the whole remaining part a bounded quadratic form on the four scaled packet roots. Its residue is

$$(4+2/e_1)K_1^2+4K_2^2+4K_4^2+(4+(4e_3+2)/e_5)K_5^2\succ 0.$$

This proves the chart for arbitrary X, Y. The same testing-only sign gauges, signed-core Schur complement, homogeneous restoration, and real/imaginary splitting complete all projective states.

The only remaining corner of this particular bounded-shape source family has h, r, g, j all infinitesimal. Then a^2 has residue 1, N is infinitesimal, and the radial pivot used here vanishes. Neither this theorem nor the preceding unit-gap theorem asserts a certificate for that further degeneration.

Chapter 121

Endpoint hermite

Component `p935c9d6318ea`
Source: `CORE/prior/inherited/endpoint_hermite.tex`
SHA256: `935c9d6318ea9a43321c6a0f3ca864560d6430b9209388fe4362319f16b5ddee`
Inventory use: actual bounded-b graph and complete normal monic divisions

Deriving the scalar and first-normal inputs by monic division. Here is a finite construction of every coefficient in the preceding table. It uses the seven factored packets, differentiation, and division by monic polynomials. Thus neither a stored expansion nor a rational-function ideal calculation is an input.

Let $\mathcal{R} = \mathbb{Q}[a, t, b, e, k]$ and use an indeterminate U for u^2 in this paragraph only. Put $K = k^2$. Substitute $(X, Y, v) = (X_g, Y_g, v_g)$ in the peeled polynomial E. Directly from the seven packets, the resulting expression has the form

$$\frac{E_g}{e^2 b^4} = e^2 \mathcal{A}(U) + ekZ\mathcal{B}(U) + k^2 Z^2 \mathcal{C}(U), \qquad \mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathcal{R}[U].$$

The displayed division means coefficientwise polynomial extraction: the constant, linear and quadratic coefficients in Z have the factors e^2, ek and k^2, respectively. No evaluated tensor is divided. Substituting the graph into the factored packets gives the five short contact and axis identities

$$(\mathcal{A} + \mathcal{B} + \mathcal{C})(K) = 0, \qquad (\mathcal{B} + 2\mathcal{C})(K) = 0, \qquad (\mathcal{A} + \mathcal{B} + \mathcal{C})'(K) = 0,$$
$$\mathcal{A}(0) = \mathcal{B}(0) = 0.$$

Consequently the following monic divisions have zero remainder:

$$\mathcal{N} = \frac{\mathcal{A} + \mathcal{B} + \mathcal{C}}{(U - K)^2}, \qquad \mathcal{D} = \frac{\mathcal{B} + 2\mathcal{C}}{U - K},$$
$$\mathcal{R}_0 = \frac{\mathcal{C} + K\mathcal{D} + K^2\mathcal{N}}{U}, \qquad \mathcal{J} = \frac{-\mathcal{D} - 2K\mathcal{N}}{U}.$$

For the last two divisions, evaluate the numerators at $U = 0$ and use the first two division identities together with $\mathcal{A}(0) = \mathcal{B}(0) = 0$. This is a polynomial-ring argument; the symbols K and U are not inverted after evaluation. Set

$$A_{00} = \mathcal{R}_0(u^2), \qquad A_{01} = u\mathcal{J}(u^2), \qquad A_{11} = \mathcal{N}(u^2).$$

Comparing the three coefficients of $1, Z, Z^2$ proves exactly

$$\frac{E_g}{e^2 b^4} = A_{00}\,[u(e - kZ)]^2 + A_{01}\,u(e - kZ)(eu^2 - k^3 Z) + A_{11}(eu^2 - k^3 Z)^2.$$

Multiplying by $e^2 b^4$ gives the scalar identity used above.

For each $\xi \in \{X, Y, v\}$, let R_ξ be the graph restriction of $\partial_\xi E$, divided coefficientwise by e^2b^2, eb^2, e^2b^2, respectively. The first two are linear in Z. The third has one quadratic coefficient, supplied solely by the signed packet:

$$[Z^2]R_v = -4R_cT_0c_d\, ekb^2u(1+u^2).$$

Define

$$C_X = C_Y = 0, \qquad C_v = 4R_cT_0c_d\, eb^2(1+u^2).$$

Then $\widetilde{R}_\xi = R_\xi - C_\xi Z\, u(e-kZ)$ is linear in Z and has the polynomial parity form

$$\widetilde{R}_\xi = eu\, L_{\xi,0}(U) + kuZ\, L_{\xi,1}(U), \qquad L_{\xi,i} \in \mathcal{R}[U].$$

In particular, subtracting the quadratic term also leaves its corrected linear coefficient divisible by ku; this follows upon collecting the signed packet with the two current packets. The contact identity is $(L_{\xi,0} + L_{\xi,1})(K) = 0$. Form the monic quotient

$$J_\xi = \frac{L_{\xi,0} + L_{\xi,1}}{U - K}.$$

The promised full first-normal coefficients are now defined by the short formulas

$$P_\xi = (L_{\xi,0} - UJ_\xi)(u^2) + C_\xi Z, \qquad Q_\xi = uJ_\xi(u^2).$$

Indeed direct multiplication gives $R_\xi = P_\xi u(e-kZ) + Q_\xi(eu^2 - k^3Z)$. This proves the full first-normal identities, including the quadratic-in-Z term; no first-normal term has been omitted.

For completeness, the weighted splits have an equally explicit finite rule. In a polynomial expanded in (e, k, b, u), place a monomial with k-exponent at least r in the first bucket and divide it by k^r; place every remaining monomial in the second bucket and divide it by e^sb^s. Use $(r, s) = (4, 2)$ for A_{00} and $(r, s) = (2, 1)$ for

$$A_{01}, \quad P_X, \quad P_Y, \quad P_v - 16kP_0.$$

Substitution of the seven short packets into the monic formulas shows that every monomial left for the second bucket has the required e^sb^s factor. Equivalently, the only polynomial remainders to check are

$$A_{00} \equiv 0 \pmod{(k^4, e^2b^2)}, \qquad A_{01}, P_X, P_Y, P_v - 16kP_0 \equiv 0 \pmod{(k^2, eb)}.$$

These are ordinary finite polynomial truncations: discard the monomials divisible by the displayed generators. The two bucket quotients have respectively residues $(-8, -8)$ for A_{00} and $(0, 0)$ for each of the other four entries; A_{11} has residue 8, and all Q_ξ have residue zero. To obtain these numbers, after the divisions set $e = k = u = Z = 0$, keeping a, t, b free. In particular the new term C_vZ belongs to $eb\,\mathcal{R}[u, Z]$ and its second-bucket quotient contains the infinitesimal testing factor Z. No assumption $\overline{b} = 0$ is used.

This is a reproducible coefficient construction: the source consists of seven short packets, the scalar construction of four monic quotients, and each first-normal construction of one monic quotient. The finite checks accompany these formulas as a convenience; the definitions and the multiplication identities, rather than expanded coefficient arrays, are the proof objects.

Chapter 122

Endpoint source

Component `p7d4e5164e84a`
Source: `CORE/prior/inherited/endpoint_source.tex`
SHA256: `7d4e5164e84a9b54940b4eea47845ca6533efe48ff8971e7e374b1dd4910b5b8`
Inventory use: actual seven source coquotients

Seven short source polynomials. The following definitions make the endpoint input reproducible without a stored expansion. Put

$$K = k^2, \quad \eta = e^2 b^2, \quad \zeta = k^2 b^2, \quad V_c = K\eta\zeta.$$

$$L_c = 1 - a\eta, \qquad H_c = 1 + (1-a)\eta, \qquad Q_c = 1 - K\zeta + aV_c.$$

Define

$$\begin{aligned}
P_4 &= 1 + K + \eta K(a - K - 1) - \zeta K t + V_c[-\zeta K t^2 + 2Kt + a(1-t) + t],\\
P_\alpha &= 1 + K - a\eta K + \zeta[(1+K) - (2+K)t] + V_c[-\zeta t^2 + a(t-1) + t],\\
P_d &= (1+K)^2 - a\eta K^2 + \zeta K[(K+1) - (K+2)t]\\
&\quad + V_c[-\zeta K t^2 + Ka(t-1) + Kt + K + 1].
\end{aligned}$$

The seven coquotients are

$$\begin{aligned}
c_2 &= 2(1+K)(1-t\zeta)L_c(1+V_c)H_cQ_c,\\
c_\beta &= 2(1+K)L_c[1 + (1-t)\zeta]H_cQ_c,\\
c_3 &= 2(1+K)L_cH_cQ_c(1 - K\eta + tV_c),\\
c_5 &= 2(1+K)L_cH_cQ_c(1+V_c),\\
c_4 &= 2Q_cP_4, \qquad c_\alpha = 2L_cH_cP_\alpha, \qquad c_d = 4L_cQ_cP_d.
\end{aligned}$$

Their residues are respectively $2, 2, 2, 2, 2, 2, 4$, so all are positive coefficient units on this domain. With $M = R_c - t$ and $J = T_0 - a$, the unchanged source is

$$\lambda = n = 0, \quad c = 2, \qquad d = \frac{2a\eta c_d}{c_3}, \quad m = \frac{2ae^2c_2}{Mc_3}, \quad A = \frac{2a\eta^2 c_4}{c_3}, \quad \alpha = \frac{2aK^2\eta\zeta c_\alpha}{c_3},$$

$$D + \frac{d}{2} = \frac{2ac_5}{Jc_3}, \qquad \beta + \frac{d}{2} = \frac{2ae^2c_\beta}{tc_3}, \qquad \Lambda = \frac{R_cT_0c_3}{2a} > 0.$$

The theorem retains the hypothesis $D, \beta \geq 0$. Write

$$r_2 = v - X + uvY - uZ, \quad r_\beta = X + uZ, \quad r_3 = u - Y, \quad r_5 = u + uvX - Y - vZ, \quad r_\alpha = vY - Z,$$

and put $q_0 = v_g$, $z_0 = \eta u$. Then G is exactly the sum of

$$G_2 = 4T_0e^2c_2[(r_2 + tq_0)^2 - tR_cq_0^2],$$

$$
\begin{aligned}
G_\beta &= 4T_0 e^2 c_\beta (1+v^2)[(r_\beta + Mq_0)^2 - R_c M q_0^2],\\
G_3 &= 4R_c c_3 (1+v^2)[(r_3 + Jz_0)^2 - T_0 J z_0^2],\\
G_5 &= 4R_c c_5 [(r_5 - az_0)^2 - aT_0 z_0^2],\\
G_4 &= 4R_c T_0 \eta^2 c_4 (1+u^2)(1+v^2) Y^2,\\
G_\alpha &= 4R_c T_0 K^2 \eta\zeta c_\alpha (1+u^2) r_\alpha^2,\\
G_d &= 2R_c T_0 \eta c_d (1+u^2)[r_\alpha^2 - (1+v^2) Z^2].
\end{aligned}
$$

These seven factored expressions agree separately with the defining source. Their proof requires multiplication of short polynomials only.

Chapter 123

A product identity repairs the endpoint at every coefficient rate

Component p05180564ed05
Source: CORE/prior/inherited/product_repair.tex
SHA256: 05180564ed0508f8b2ad17b0027feb9f0cbd47ec865e82f3ee046c30c5b545cf
Inventory use: actual bounded-b W-state nine-root proof

123.1 A product identity repairs the endpoint at every coefficient rate

The new boundary mechanism is algebraic. A troublesome root is retained, and a second relation controls its product with an additional root. This supplies two positive pivots without comparing an infinitesimal in one field with an infinitesimal in the other.

Lemma 123.1 (Product-relation completion)**.** *Let z be a root vector over T, and $h, r \in T$. Suppose*

$$F = (z,h,r)^T Q(z,h,r), \qquad Q \in \mathrm{Sym}(B), \qquad \bar{Q} = \mathrm{diag}(Q_0, -a, 0),$$

where Q_0 is a real positive-definite matrix and $a \in \mathbb{R}_{\geq 0}$. Suppose $p \in B$ has nonzero residue, A is a row with entries in $\mathfrak{n}$, and

$$ph = Az + r, \qquad phr = (z,h,r)^T E(z,h,r), \quad E \in \mathrm{Sym}(\mathfrak{n}).$$

Then $F \in \Sigma T^2$.

Proof. For a sufficiently large positive real M, add the exact zero

$$M\left[p^2h^2 + r^2 - (Az)^2 - 2phr\right] = 0.$$

The residue Gram becomes $\mathrm{diag}(Q_0, M\bar{p}^2 - a, M) \succ 0$. The subtracted terms have zero residue by hypothesis. Lemma 1.2 applies. In particular p is never inverted. □

123.1.1 Exact contact scope and full-polynomial input

This application uses the canonical C1 contact model, with $\lambda = n = 0$. Its source parameters are positive e, k, b, a, t satisfying

$$e, k \in \mathfrak{m}_c, \quad b \in B_c, \qquad R_c = 1 + k^2(1 - ae^2b^2), \quad T_0 = 1 + k^2 - tk^4b^2,$$

$$0 < a < T_0, \qquad 0 < t < R_c.$$

The signed weight satisfies $d > 0$, and all eight event coefficients are nonnegative original physical C1 contact coefficients. This is a specified contact chart, not every coefficient vector with the same

omissions. The independent testing chart is $W = 1$ with u, v, X, Y, Z infinitesimal. In this section there is no restriction on k^3/e or k/e.

The complete seven-packet contact polynomial is denoted G. Its short factored definition and unchanged original coefficient vector are printed below. The accompanying checkers compare the entire polynomials, including the scalar and normal terms; no global coverage claim is inferred from those checks.

Define

$$\begin{gathered} v_g = ekb^2u, \quad X_g = -uZ + tv_g, \quad Y_g = (1 - ae^2b^2)u, \\ N_1 = X - X_g, \quad N_2 = Y_g - Y, \quad \delta v = v - v_g, \quad D_t = Z - uX, \\ U_0 = 1 + u^2, \quad c_0 = 1 - tk^2b^2, \quad P_0 = T_0 + u^2 = U_0 + k^2c_0, \\ S = u + uvX - Y - vZ + (T_0 - a)e^2b^2u, \\ s_0 = eb^2u(e - kZ), \qquad q = eb^2(eu^2 - k^3Z). \end{gathered}$$

Source inclusion. The original input is reproduced once as Component p7d4e5164e84a (Chapter 122).

Direct substitution gives the exact nonlinear relation

$$L = P_0s_0 - S + N_2 - D_t\delta v + uv_gN_1 - uc_0q = 0. \tag{E1}$$

Peel the two complete current squares:

$$E = G - H_3(1 + v^2)N_2^2 - H_5S^2, \qquad H_3 = 4R_cc_3, \quad H_5 = 4R_cc_5, \quad \bar{H}_3 = \bar{H}_5 = 8.$$

Each peeled packet divided by e^2 is polynomial in bounded variables. The full graph scalar has the identity

$$E_g = A_{00}s_0^2 + A_{01}s_0q + A_{11}q^2,$$

and the entire first normals, with $g_0 = s_0/(eb^2)$ and $g_2 = q/(eb^2)$ denoting their displayed polynomial expressions, satisfy

$$E_X = e^2b^2(P_Xg_0 + Q_Xg_2), \quad E_Y = eb^2(P_Yg_0 + Q_Yg_2), \quad E_v = e^2b^2(P_vg_0 + Q_vg_2).$$

The exact polynomial splits and residues needed are

Quantity	Polynomial split	Quotient residues
A_{00}	$k^4a_0 + e^2b^2a_1$	$-8, -8$
A_{01}	$k^2c_1 + ebc_2$	$0, 0$
P_X	$k^2p_X + ebq_X$	$0, 0$
P_Y	$k^2p_Y + ebq_Y$	$0, 0$
$P_v - 16kP_0$	$k^2p_v + ebq_v$	$0, 0$

Also $\bar{A}_{11} = 8$ and $\bar{Q}_X = \bar{Q}_Y = \bar{Q}_v = 0$. These equalities remain valid with b symbolic; boundedness, rather than infinitesimality, suffices for b.

Source inclusion. The original input is reproduced once as Component p935c9d6318ea (Chapter 121).

Exact finite Taylor expansion in $(N_1, -N_2, \delta v)$ accounts for every higher-normal term. Because E/e^2 is polynomial, every term of normal degree at least two allocates to $x = eN_1$, $y = e\delta v$, N_2 with bounded coefficients. Terms of degree at least three have an extra infinitesimal normal. Terms involving N_2 retain an e factor. The only additional nonzero residue is

$$B_0 = \begin{pmatrix} 16 & -8 \\ -8 & 8 \end{pmatrix} \quad \text{on } (x, y).$$

It is visible without expansion of the full Hessian: at $u = Y = Z = 0$, the quadratic residue of E/e^2 is $8(X - v)^2 + 8X^2$, with b still free.

123.1.2 The new identity and its nine-root proof

If the signed testing core is nonnegative, its pure testing square root and the original event packets already give SOS. Otherwise the sign gauges $u, v \geq 0$ preserve every packet, and the core Schur complement implies $u, v > 0$ and $D_t > 0$. To check the latter, after $Z = D_t + uX$ the Schur complement on (W, D_t) is

$$\frac{2(1+v^2)}{N_0} \begin{pmatrix} u^2v^2(1+u^2) & -uv(1+u^2)(1+v^2) \\ -uv(1+u^2)(1+v^2) & u^2v^2 \end{pmatrix},$$

where $N_0 = u^2v^4 + u^2v^2 + u^2 + v^4 + 2v^2 + 1 > 0$. The eliminated block is positive definite. For $W = 1, D_t \leq 0$, the remaining form is nonnegative, proving the claim.

Consequently $w = \sqrt{ek}\sqrt{D_t}\,\delta v$ is a legal separated root. Set

$$h = k^2 s_0 - uq = e^2b^2u(k^2 - u^2), \qquad z = ebs_0, \qquad r = k^2 D_t \delta v,$$
$$A_0 = k^2(S - N_2) - k^3b^2u^2x - uU_0q.$$

Multiplication of (E1) by k^2 gives $P_0h = A_0 + r$. The additional exact identity is

$$\begin{aligned} P_0hr = {} & ku^2(P_0 - k^2)y(S - N_2) - ku(P_0 + c_0k^2u^2)yq \\ & + u^2(P_0 - k^2)w^2 - k^4b^2u^2T_0xy. \end{aligned} \tag{E2}$$

Every displayed coefficient has zero residue and is a polynomial in separately bounded factors. To derive (E2), substitute the two soft roots to obtain

$$u(P_0 - k^2)s_0 - (1 + c_0u^2)q = ekb^2(k^2 - u^2)(D_t + uN_1).$$

Then use $D_t = U_0Z - uN_1 - tuv_g$, $T_0 = 1 + k^2c_0$, and (E1) to replace $S - N_2$ in (E2). The terms in $D_t(\delta v)^2$ cancel; the remaining terms give P_0hr. This is a whole-polynomial calculation.

Let J_0 denote the right-hand side of (E2). Thus

$$Z_0 = P_0^2h^2 + r^2 - A_0^2 - 2J_0 = 0.$$

Add to the entire G the exact zero expression

$$-16kyL + 16(ebL)^2 + 16Z_0.$$

The first term cancels the exceptional $16kP_0$ first normal and supplies $16w^2$. Use $k^2s_0 = h + uq$ in every other scalar and first-normal term. The scalar becomes

$$a_0(h + uq)^2 + a_1z^2 + [c_1(h + uq) + c_2z]q + A_{11}q^2.$$

Its only negative residue diagonals are -8 on h, z. The safe relation

$$ebL = P_0z - ebS + ebN_2 - bD_ty + ekb^3u^2x - ebuc_0q$$

has only infinitesimal coefficients apart from P_0; hence $16(ebL)^2$ changes the z diagonal to 8. Every coefficient of A_0 is infinitesimal, as is every coefficient in (E2). Thus $16Z_0$ changes the h diagonal to 8 and supplies a new r diagonal 16, with no other residue change.

The full Gram on

$$\boxed{(x, y, N_2, S, w, q, h, z, r)}$$

therefore has residue

$$\boxed{B_0 \oplus \operatorname{diag}(8, 8, 16, 8, 8, 8, 16) \succ 0.}$$

Its first two LDL pivots are $16, 4$; the other seven pivots are printed above. The strict-residue lemma proves SOS of the complete seven-packet polynomial for every coefficient ratio k^3/e.

123.1.3 Restoration and coverage

The complete original-parent restoration is

$$\Lambda F = G + 4T_0e^2c_2\frac{t}{M}(r_2 - Mq_0)^2 + 4T_0e^2c_\beta\frac{M}{t}(1+v^2)(r_\beta - tq_0)^2$$
$$+ 4R_cc_3\frac{J}{a}(1+v^2)(r_3 - az_0)^2 + 4R_cc_5\frac{a}{J}(r_5 + Jz_0)^2.$$

Every additional packet is a nonnegative coefficient factor times a pure-testing factor times a square. This identity is four instances of

$$S^2 + 2(A+B)qS + (A+B)Aq^2 + \frac{B}{A}S^2 = \frac{A+B}{A}(S+Aq)^2.$$

Use respectively $(A,B,q) = (M,t,q_0), (t,M,q_0), (a,J,z_0), (J,a,-z_0)$ and $S = r_2 - Mq_0, r_\beta - tq_0, r_3 - az_0, r_5 + Jz_0$. Comparing the seven resulting coefficients with the displayed source proves the identity. Since Λ is a positive pure coefficient factor, the pure-factor restoration in RFA13 (p. 7) recovers the original parent. Normalizing the original state to $W = 1$ and undoing that normalization use only testing factors.

This removes the former k^3/e restriction and includes positive bounded b. It does not prove entry of every physical C1 or C2 source into this domain. For unbounded b, the recovered inverse-b theorem remains a different input with a ten-root, large polynomial certificate. The new short identity should not be presented as a replacement for that certificate or for the full source cover.

Chapter 124

All matching-strong C2 sources in the positive contact chart

Component `pf8e9d3c47c72`
Source: `CORE/prior/matching/matching_source_entry.tex`
SHA256: `f8e9d3c47c72299b408f54ca1cf43acdeae95bba13578f285cb562c8cf1fee35`
Inventory use: Only exact rational source reflection and all preserved strict source gaps

124.1 All matching-strong C2 sources in the positive contact chart

We now remove the bounded-angle and bounded-ratio assumptions from the weak-event entry lemma, provided the four matching packets are strong. All source hypotheses refer to the same exact positive rational C2 chart; this is not an assertion that every canonical C2 source has that chart. Write C for its event weight and c for its contact half-angle.

Theorem 124.1 (A complete matching-strong C2 source domain)**.** *Use the physical C2 source with $d = 1$, $\lambda = \beta = 0$,*

$$c, a, b, t > 0, \qquad 0 < r < 1, \qquad h = b - a^2 t > 0,$$
$$g = (1 + c^2)b - (1 - r + a^2 c^2 t) > 0, \qquad j = 1 + c^2 r - t > 0,$$

and the original positivity conditions $1 - c^2 h > 0$, $b - a^2 c^2 r > 0$, with $n, m, A, D > 0$. Suppose C, α are infinitesimal and

$$1/n, \quad 1/m, \quad 1/A, \quad 1/D \quad \textit{are bounded.}$$

Then every independent projective test and real or complex state has tensor-SOS full parent. No bound is assumed on a, c, b, g and no comparability is assumed between C and α. After at most the exact C2 reflection, its actual source has

$$c \textit{ bounded}, \qquad a, b, t \longrightarrow 1, \qquad r, h, g, j \longrightarrow 0.$$

The arrows denote coefficient residues.

Proof. Let $\mu = \min(n, m, A, D)$. The previously proved matching budget, applied with $T = C + \alpha$, makes $1/[\mu(C+\alpha)]$ bounded. Since $C+\alpha$ is infinitesimal, $1/\mu$ is infinitesimal. Thus all four matching weights are unbounded.

Normalize the actual contact state to have Euclidean norm one. At the exact contact,

$$0 = 4\sum_i e_i p_i + \frac{H}{D_u D_v}, \qquad \frac{H}{D_u D_v} \geq -2Z^2 \geq -2,$$

where $p_n = X^2$, $p_A = Y^2$ and the other p_i are the normalized squared event amplitudes. Thus $0 \le p_i \le 1/(2e_i)$ for each matching event. Every matching probability has zero residue; in particular X, Y have zero residue. Put

$$c_u = (1+u_c^2)^{-1/2}, \quad s_u = u_c c_u, \qquad c_v = (1+v_c^2)^{-1/2}, \quad s_v = v_c c_v.$$

These are bounded positive coefficient quantities with $c_u^2 + s_u^2 = c_v^2 + s_v^2 = 1$. Write $\sigma = \overline{c_u s_v}$ and $\tau = \overline{s_u c_v}$. The two other vanishing matching probabilities give

$$\sigma\overline{W} - \tau\overline{Z} = 0, \qquad \tau\overline{W} - \sigma\overline{Z} = 0.$$

At least one of $\overline{W}, \overline{Z}$ is nonzero, so $\sigma^2 = \tau^2$. Both are nonnegative, hence $\sigma = \tau$. If they are positive, both contact half-angles are bounded units with the same residue. If they vanish, either both contact half-angles are infinitesimal or both are unbounded. Indeed $c_u s_v = s_u c_v = 0$ in residue, together with the two unit-circle identities, permits precisely these two endpoint alternatives.

The exact reflection sends (u_c, v_c) to $(1/v_c, 1/u_c)$. It therefore reduces the two endpoint cases to the infinitesimal one. It preserves the positive source domain: for $z = c^2$ its parameters are

$$c' = \frac{1}{ac}, \quad a' = a, \quad b' = \frac{a^2}{b}, \quad r' = \frac{a^2 z r}{b}, \quad t' = \frac{1 - zh}{b},$$

$$h' = \frac{a^2 z h}{b}, \quad g' = \frac{a^2 j}{b}, \quad j' = \frac{g}{b}.$$

In particular

$$1 - r' = \frac{b - a^2 z r}{b} > 0, \qquad 1 - z'h' = \frac{a^2 t}{b} > 0, \qquad b' - a^2 z' r' = \frac{a^2(1-r)}{b} > 0.$$

Direct substitution in the rational source gives

$$\Delta' = \frac{\Delta}{b^2 z^2}, \quad N' = \frac{a^2 N}{b^2}, \qquad (n', m', C', A', D', \alpha') = (n, D, \alpha, A, m, C).$$

Consequently every hypothesis is retained. These are pure-coefficient operations. In testing coordinates the reflection is the full homogeneous parent identity with factor $u^2 v^2$, interpreted in projective charts.

It remains to bound a after this orientation. At an ordinary contact residue $a = v_c/u_c$ is a bounded unit. In the other case both c and ac are infinitesimal. Suppose for contradiction that $q = a^2$ is unbounded. Then z and qz are infinitesimal; also $0 < t < 1 + zr$ is bounded. The numerator in

$$m = \frac{(1+qz)(1-r)(1+zr)(1+zr+qzt)}{2g\Delta}$$

is bounded. Since $1/m$ is bounded, $g\Delta$ is bounded. The positive expansion of Δ in the entry lemma gives $\Delta \ge 1$. Thus g is bounded, and

$$b = \frac{1 - r + g + qzt}{1 + z}$$

is bounded. Positivity of $h = b - qt$ gives $0 < t < b/q$, so t is infinitesimal and $j = 1 + zr - t$ has residue one. But then

$$D = \frac{(1+z)(b+r)(b-qzr)(1+zr+qzt)}{2qj\Delta}$$

is infinitesimal: the numerator is bounded, since $0 < b - qzr < b$, whereas $1/q$ is infinitesimal, j is a unit, and $\Delta \ge 1$. This contradicts boundedness of $1/D$.

We have proved that a, c are bounded in the oriented source. Its weak event C is infinitesimal, so the weak-event entry lemma applies: a, b, t have residue one and r, h, g, j are infinitesimal. If c is infinitesimal, the completed small-angle radial theorem gives all independent tests. If c is a bounded unit, the completed ordinary-angle radial theorem gives all tests. Reflect back when necessary. Restore the positive pure coefficient normalization if the original d was not one. □

The conclusion strengthens the earlier comparable-weak-packet corollary on this exact positive contact chart. It covers unbounded ratios α/C and C/α, while preserving independence of actual and testing parameters. Within the branch where C, α are both infinitesimal, the remaining C2 sources have at least one weak matching event or have not yet been entered into this positive contact chart. The theorem does not remove that source-chart obligation.

Chapter 125

A short contact–conductor proof of the C1 memberships

Component `pfabc29e28ab3`
Source: `CORE/prior/prior_core/c1/contact_conductor_proof.tex`
SHA256: `fabc29e28ab3d39610622d197401f8d3487d093c77d8298f647894903029ed03`
Inventory use: CC1–CC8; full factored scalar and both first normals | CC1-CC6; bounded/weighted application discussion | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

125.1 A short contact–conductor proof of the C1 memberships

Write $d = \delta$ in this section; this local parameter is distinct from the global Bell coefficient d_{B}. Put

$$R_0 = \mathbb{Q}[d, k, b, a, t], \qquad R = R_0[u, w, v], \qquad J = (u - k, w - dk, v - dk^2b^2).$$

For $F \in R$, write $F_c = F(k, dk, dk^2b^2)$. Define

$$A = wv, \quad B = dv, \quad C = db^2uw, \quad D_0 = d^2k^3b^2, \qquad M = (A, B, C),$$

$$x = B - C, \qquad y = A - kC, \qquad z = uB - kC, \qquad h = kC - D_0.$$

These are the original roots $g_0, h_{13}, h_{23}, h_{34}$; in particular, $D_0 = a_4^2$, whereas the source monomial in g_4 is kD_0. Adjoin the separated radicals used in the original four-monomial circuit,

$$a_1 = \sqrt{wv}, \quad a_2 = \sqrt{duv}, \quad a_3 = b\sqrt{dkuw}, \quad a_4 = dk^{3/2}b, \qquad q = a_1a_2 - a_3a_4.$$

Only $q^2 \in R$ is needed in the scalar lift. The following three identities are the entire conductor calculation:

$$C(u - k) = z - ux, \tag{125.1}$$

$$C(w - dk) = dy - wx, \tag{125.2}$$

$$C(v - dk^2b^2) = b^2(wz + dky - kwx). \tag{125.3}$$

Thus $CJ \subset (x, y, z)$ without inverting any parameter. For

$$r_* = y + z - h - 2kx$$

we also have the two exact identities

$$r_* = (v - dk^2b^2)((w - dk) + d(u - k)) - dkb^2(u - k)(w - dk), \tag{125.4}$$

$$kCr_* = q^2 - yz. \tag{125.5}$$

The first says $r_* \in J^2$. The second supplies the otherwise missing conductor factor through one of the original circuit squares.

Explicit monic division. Every $S \in R$ has a polynomial decomposition

$$S = S_c + (u-k)T_u + (w-dk)T_w + (v-dk^2b^2)T_v. \tag{CC1}$$

One obtains it by substituting $u = k$, then $w = dk$, then $v = dk^2b^2$, and using $X^n - Y^n = (X-Y)\sum_{j=0}^{n-1} X^{n-1-j}Y^j$ at each step. There are no parameter denominators. If $S_c = 0$, equations (125.1)–(125.3) give

$$CS = x(-uT_u - wT_w - b^2kwT_v) + y(dT_w + dkb^2T_v) + z(T_u + b^2wT_v). \tag{CC2}$$

Lemma 125.1 (First-order and restricted second-order contact lifts)**.** *In the above ring, the following assertions hold, with polynomial coefficients in R:*

$$M \cap J \subset (x,y,z), \qquad \big(M^2 + (D_0^2)\big) \cap J^2 \subset (x,y,z,h,q)^2. \tag{CC3}$$

Here (D_0^2) is the principal ideal generated by D_0^2; the second assertion does not replace it by all mixed products involving D_0.

Proof. For the first assertion, write $F = R_AA + R_BB + R_CC$. Since $A = y + kC$ and $B = x + C$,

$$F = R_Ay + R_Bx + CS, \qquad S = kR_A + R_B + R_C.$$

The contact value is $F_c = C_cS_c$, where $C_c = d^2k^2b^2$ is a nonzero polynomial. Since R_0 is a domain, $F_c = 0$ implies $S_c = 0$. Apply (CC2). This cancellation proves a zero identity; it introduces no coefficient $1/C_c$.

For the second assertion choose any displayed representation $F = \sum_{i\le j} f_{ij}m_im_j + \rho D_0^2$, with $(m_1,m_2,m_3) = (A,B,C)$. Substitute

$$A = y + kC, \qquad B = x + C, \qquad D_0 = kC - h.$$

This gives

$$F = Q_2 + C(xH_x + yH_y + zH_z + hH_h) + C^2S, \qquad H_h = -2k\rho, \tag{CC4}$$

where $Q_2 \in (x,y,h)^2$ and initially $H_z = 0$. The contact value again implies $S_c = 0$. Use (CC2) on CS and absorb C^2S into the four coefficients in (CC4). This leaves $H_h = -2k\rho$ unchanged and gives the same formula without C^2S.

The first derivatives of $F \in J^2$ vanish at contact. The first three root gradients have determinant

$$\det\left(\frac{\partial(x,y,z)}{\partial(u,w,v)}\right)_c = -d^4k^4b^4 \neq 0. \tag{CC5}$$

Equation (125.4) gives a gradient relation with coefficient vector $(-2k,1,1,-1)$. Its coefficient at h is -1. Set $\lambda = 2k\rho_c$. Subtract $\lambda(-2k,1,1,-1)$ from (H_x,H_y,H_z,H_h). The last resulting contact value is zero, and the first three satisfy the homogeneous linear system with determinant (CC5). Multiplication by its adjugate and cancellation in the domain R_0 show that those three contact values are zero too. No inverse of the determinant is used in a coefficient: each resulting polynomial J_i lies in J and is divided only by the monic procedure (CC1). Consequently

$$F = Q_2 + 2k\rho_cCr_* + C\sum_{i=1}^{4} r_iJ_i = Q_2 + 2\rho_c(q^2 - yz) + \sum_{i=1}^{4} r_i(CJ_i), \tag{CC6}$$

where $(r_1,r_2,r_3,r_4) = (x,y,z,h)$. Each CJ_i is explicitly a combination of x,y,z by (CC2), which proves the assertion in the unlocalized polynomial ring. □

Why the restriction matters. The unrestricted claim obtained by replacing $M^2 + (D_0^2)$ with $(A,B,C,D_0)^2$ is false. Indeed $Cr_* \in (A,B,C,D_0)^2 \cap J^2$. Under the positive weights below it has weight 42. Every weight-42 member of the square of the original soft-root ideal must be divisible by x, since x has weight 20 and every other original soft root has weight at least 22. But Cr_* is not divisible by x: setting $d = k = b = w = 1$, $u = 2$, $v = 2$ gives $x = 0$ and $Cr_* = 2$. The factor k supplied by D_0^2 in (CC4) is therefore substantive.

Application to the complete source scalar and both normals. Use precisely the seven source coquotients and the definitions in Appendix 126.1, renaming their clearing multiplier $\mathcal{C} = K^6Q^4T_n$ to distinguish it from the present conductor monomial C. In particular

$$K = 1 + k^2, \quad Q = K + ak^2b^2, \quad T_n = K^2 - td^2k^4Qb^2, \quad D = 1 + u^2, \quad D_v = 1 + v^2,$$

$$V_n = K^2 - tQuvw, \quad Y_n = KT_nuw - avV_n, \quad B_0 = vY_n - KT_n, \quad A_c = C_5T_n + aK(C_3D_v - C_5).$$

The exact scalar F_0 has the following five-term form:

$$\begin{aligned} F_0 = {} & 4KQT_n^2[C_2(K^2 - tQ) + C_bD_vtQ]D^2A^2 \\ & + 4Q^2A_cD^2V_n^2B^2 \\ & + 4K^3C_4DD_v(KT_nC - ab^2BV_n)^2 \\ & + 4KQ^2C_aDB_0^2D_0^2 \\ & + 2K^2QC_dD\{b^2B^2(Y_n^2 - K^2T_n^2 + 2aKT_nV_n) - 2K^2T_n^2BC\}. \end{aligned}$$

Also the exact leading polynomial is

$$P = D(A^2 + B^2) + K(C^2 + D_0^2) - 2K^2BC.$$

Thus $E_0 = F_0 - 8K^2D(K^6Q^4T_n)P$ belongs to $M^2 + (D_0^2)$ directly, without ideal division.

For the first normals put

$$\begin{aligned} S_B = {} & -8aK^2b^4C_4D_vV_n + 8d^2Q^2k^6b^4C_aB_0 + 4KQb^2C_dB_0, \\ S_C = {} & 8K^3T_nb^2C_4D_v. \end{aligned}$$

The normal numerator satisfies $S_n/d = BS_B + CS_C$. Hence

$$\begin{aligned} L_1 = {} & A\,[8K^3QT_n^2(C_bD_v - C_2)D] \\ & + B\,[-8dK^2Q^2uDvV_nA_c + dK^3auvS_B] + C\,[dK^3auvS_C], \\ L_2 = {} & B\,[8K^2Q^2T_nDV_n(C_3D_v - C_5) - K^2T_nS_B] - C\,K^2T_nS_C. \end{aligned}$$

These formulas prove $L_1 \in M$ and $L_2 \in (B, C)$. The native contact and derivative identities (SD9)–(SD11) in Appendix 126.1 prove

$$(E_0)_c = (\partial_uE_0)_c = (\partial_wE_0)_c = (\partial_vE_0)_c = 0, \qquad (L_1)_c = (L_2)_c = 0. \tag{CC7}$$

These are source polynomial identities, with no positivity hypothesis or parameter inversion. Their proof uses only six short native balances and the chain rule. The supplied small checker confirms the identities and multiplies back the complete lifts from (CC2) and (CC6). By (CC3),

$$E_0 \in (x, y, z, h, q)^2, \qquad L_1, L_2 \in (x, y, z). \tag{CC8}$$

Thus all three memberships previously expressed by the finite Gröbner-basis witness have a single contact–conductor mechanism.

Bounded coefficients and the original Gram argument. Give (d, k, b, a, t, u, w, v) weights $(2, 2, 6, 0, 0, 2, 4, 18)$. Then A, y, z, h, q have weight 22, while B, C, x have weight 20. The three substitutions defining contact preserve weight, as do the monic geometric-sum divisions after accounting for the weight of their linear factor. The factored weight argument in Appendix 126.1 gives, without a large expansion,

$$\mathrm{wt}(E_0) \geq 52, \quad \mathrm{wt}(L_1) \geq 42, \quad \mathrm{wt}(L_2) \geq 32.$$

Consequently the construction gives lower bounds $8, 20, 10$ for the three families of correction coefficients, respectively. The optional exact checker obtains the stronger scalar minimum 56 and

coefficient bounds $12, 20, 12$ for its particular monomial assignment. All are positive. Every coefficient has nonnegative exponents, including in the bounded shapes a, t; every correction therefore has zero residue on the stated C1 branch. The full hard-normal block and its positive residue are printed in Appendix 126.1. Restoration is exactly that of the source theorem: restore $\mathcal{C} = K^6Q^4T_n$ times the four positive packets subtracted there, then undo the positive multiplier $\mathcal{C}\Delta^2\Lambda$. The present argument proves the complete memberships on that branch and does not assert a global source cover.

Chapter 126

Self-contained C1 source definitions and contact verification

Component `p0acdb66bba2f`
Source: `CORE/prior/prior_core/c1/source_definitions.tex`
SHA256: `0acdb66bba2fea9b217d8089bcf1df5a1fb50380b01a1a693bbb1b9b1295f241`
Inventory use: SD1–SD11; full quadratic normal block | SD1-SD11; especially SD2-SD5 | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

126.1 Self-contained C1 source definitions and contact verification

The letter d in the contact–conductor proof denotes the new parameter δ, not the Bell coefficient. Here the latter is denoted d_{B}, and the original event coefficient e_5 is denoted D_{e}. The testing factor is always $D = 1 + u^2$. The conductor monomial $C = \delta b^2 uw$ is distinct from the clearing multiplier $\mathcal{C} = K^6 Q^4 T_n$.

Domain and seven native coquotients. The coefficient field and testing field are independent real closed fields. On this C1 source branch, $k, f, g > 0$ are coefficient infinitesimals, $e = kf/g$ is positive unbounded, and $\Delta = e^{-1}$. The shapes a, t are positive bounded elements;

$$R = 1 + k^2(1 - af^2), \qquad T = 1 + k^2 - k^2 g^2 t, \qquad M = R - t > 0, \quad J = T - a > 0.$$

All original event coefficients are assumed nonnegative. In particular, the formulas below retain the requirements $D_{\mathrm{e}}, \beta \geq 0$. No relation compares a testing rate with a coefficient rate. Put

$$\kappa = k^2, \quad \eta = f^2, \quad \zeta = g^2, \quad V_c = \kappa\eta\zeta, \quad L_c = 1 - a\eta, \quad H_c = 1 + (1 - a)\eta, \quad Q_c = 1 - \kappa\zeta + aV_c.$$

Define

$$\begin{aligned}
P_4 &= 1 + \kappa + \eta\kappa(a - \kappa - 1) - \zeta\kappa t + V_c[-\zeta\kappa t^2 + 2\kappa t + a(1 - t) + t],\\
P_\alpha &= 1 + \kappa - a\eta\kappa + \zeta[(1 + \kappa) - (2 + \kappa)t] + V_c[-\zeta t^2 + a(t - 1) + t],\\
P_d &= (1 + \kappa)^2 - a\eta\kappa^2 + \zeta\kappa[(\kappa + 1) - (\kappa + 2)t]\\
&\quad + V_c[-\zeta\kappa t^2 + \kappa a(t - 1) + \kappa t + \kappa + 1].
\end{aligned}$$

The complete seven coquotients are

$$\begin{aligned}
c_2 &= 2(1 + \kappa)(1 - t\zeta)L_c(1 + V_c)H_c Q_c,\\
c_b &= 2(1 + \kappa)L_c[1 + (1 - t)\zeta]H_c Q_c,\\
c_3 &= 2(1 + \kappa)L_c H_c Q_c(1 - \kappa\eta + tV_c),\\
c_5 &= 2(1 + \kappa)L_c H_c Q_c(1 + V_c),
\end{aligned}$$

$$c_4 = 2Q_cP_4, \qquad c_\alpha = 2L_cH_cP_\alpha, \qquad c_d = 4L_cQ_cP_d.$$

Their residues, in this order, are $2,2,2,2,2,2,4$. Here c_b is the original c_β, and the subscript on c_d is a name rather than the conductor parameter.

Original source and full parent. The original event coefficients and multiplier are

$$\begin{aligned} &\lambda = n = 0, \qquad c = 2, \qquad d_{\mathrm{B}} = \frac{2a\eta c_d}{c_3}, \qquad m = \frac{2ae^2c_2}{Mc_3}, \qquad A = \frac{2a\eta^2c_4}{c_3}, \\ &\alpha = \frac{2a\Delta^2k^6\eta^2c_\alpha}{c_3}, \qquad D_{\mathrm{e}} + \frac{d_{\mathrm{B}}}{2} = \frac{2ac_5}{Jc_3}, \qquad \beta + \frac{d_{\mathrm{B}}}{2} = \frac{2ae^2c_b}{tc_3}, \qquad \Lambda = \frac{RTc_3}{2a}. \end{aligned} \tag{SD1}$$

Let F denote the original complete parent, including its signed core. For a testing state (W,X,Y,Z) define

$$D = 1+u^2, \quad D_v = 1+v^2, \quad B = X+uZ, \quad Q_A = W+uY,$$
$$r = uW - Y, \quad r_2 = vQ_A - B, \quad r_5 = r - v(Z-uX).$$

With $H_2 = 4Tc_2$, $H_b = 4Tc_b$, $H_3 = 4Rc_3$, $H_5 = 4Rc_5$, the exact entire parent is

$$\begin{aligned} \Phi := \Delta^2\Lambda F = {} & H_2\frac{R}{M}r_2^2 + H_b\frac{R}{t}D_vB^2 + \Delta^2H_3\frac{T}{a}D_vr^2 + \Delta^2H_5\frac{T}{J}r_5^2 \\ & + 4RT\Delta^2\eta^2c_4DD_vY^2 + 4RT\Delta^4k^6\eta^2c_\alpha D(vY-Z)^2 \\ & + 2RT\Delta^2\eta c_dD\{(vY-Z)^2 - D_vZ^2\}. \end{aligned} \tag{SD2}$$

Thus every original source event is accounted for.

Retained packets and state coordinates. In the remaining chart $Z=1$ with u,v,W,X,Y infinitesimal, put

$$p = t/R, \quad h = a/T, \quad q_A = v(W+uY), \quad q_B = v(1-uX), \quad N = B - pq_A, \quad M_B = r - hq_B.$$

Retain the four nonnegative packets

$$\Pi_1 = H_2\frac{t}{M}N^2, \quad \Pi_2 = H_b\frac{M}{t}D_vN^2, \quad \Pi_3 = \Delta^2H_3\frac{J}{a}D_vM_B^2, \quad \Pi_4 = \Delta^2H_5\frac{a}{J}M_B^2. \tag{SD3}$$

Their subtraction is the exact identity

$$\begin{aligned} G = \Phi - \sum_j \Pi_j = {} & H_2\{N^2 - 2q_AN + (1-p)q_A^2\} + H_bD_v\{N^2 + 2q_AN + pq_A^2\} \\ & + \Delta^2H_3D_v\{M_B^2 + 2q_BM_B + hq_B^2\} + \Delta^2H_5\{M_B^2 - 2q_BM_B + (1-h)q_B^2\} + \mathcal{W}, \\ \mathcal{W} = {} & 4RT\Delta^2\eta^2c_4DD_vY^2 + 4RT\Delta^4k^6\eta^2c_\alpha D(vY-1)^2 \\ & + 2RT\Delta^2\eta c_dD\{(vY-1)^2 - D_v\}. \end{aligned} \tag{SD4}$$

The testing coordinate $w = (W+uY)/D$ has the full inverse

$$\begin{aligned} &X = N + pDvw - u, \qquad q_B = v(D - uN - puDvw), \\ &W = w + \frac{u(M_B+hq_B)}{D}, \qquad Y = uw - \frac{M_B+hq_B}{D}. \end{aligned} \tag{SD5}$$

Its only testing denominator is the positive unit D. On the negative-core branch, sign gauges give $u,v>0$, and $w>0$: after eliminating the two positive core squares the remaining scalar is

$$\frac{uvDD_v}{\mathcal{D}}\{Duvw^2 - 2DD_vw + uv\}, \qquad \mathcal{D} = u^2v^4 + u^2v^2 + u^2 + v^4 + 2v^2 + 1 > 0,$$

which is nonnegative if $w \le 0$. The eliminated matrix is $\left(\begin{smallmatrix} 1+v^2+u^2v^2 & uv \\ uv & (D+v^2)/D \end{smallmatrix}\right) \succ 0$. The nonnegative-core branch and literal axes already give direct SOS. Thus the radicals in the conductor proof separate between the two fields.

Coefficient substitution and complete first normals. Put

$$K = 1 + k^2, \quad \delta = \Delta R/K, \quad b^2 = \eta K/R, \quad Q = K + ak^2b^2, \quad T_n = K^2 - t\delta^2 k^4 Q b^2.$$

Then δ, k, b are positive infinitesimals, and

$$\eta = Kb^2/Q, \quad \Delta = \delta Q/K, \quad \zeta = \delta^2 k^2 Q b^2/K, \quad R = K^2/Q, \quad T = T_n/K. \tag{SD6}$$

All displayed denominators are positive pure-coefficient units. Set $C_i = K^2Q^3c_i$ after this substitution. Explicitly, a native monomial $c\,k^j f^{2r} g^{2s} a^\alpha t^\tau$ becomes

$$c\,\delta^{2s} k^{j+2s} b^{2(r+s)} a^\alpha t^\tau K^{2+r-s} Q^{3-r+s}. \tag{SD7}$$

The native coquotients have $-2 \le r - s \le 3$, so these are polynomials with no negative exponents or shape inverses. Define V_n, Y_n, B_0, A_c as in the conductor proof and $\mathcal{C} = K^6Q^4T_n$. The factored scalar printed there is exactly $F_0 = (\mathcal{C}G)|_{N=M_B=0}$. For completeness put

$$S_n = 8\delta^2 K^2 b^4 C_4 D_v Y_n + 8\delta^4 Q^2 k^6 b^4 C_\alpha v B_0 + 4\delta^2 KQ b^2 C_d v B_0.$$

The complete coefficients on the hard normals $N, \delta M_B$ are

$$\begin{aligned} L_1 &= 8K^3 Q T_n^2 (C_b D_v - C_2) Dvw - 8\delta^2 K^2 Q^2 u Dv^2 V_n A_c + K^3 auv S_n, \\ L_2 &= 8\delta K^2 Q^2 T_n Dv V_n (C_3 D_v - C_5) - K^2 T_n S_n/\delta. \end{aligned} \tag{SD8}$$

The quotient in the last line is polynomial because S_n contains δ^2. These formulas follow by substituting (SD5) into (SD4), channel by channel; they include all normal terms of degree one.

Short native proof of the exact contact identities. Write $V = 1 + V_c$ and $K = 1 + \kappa$ in this paragraph. The factored coquotients above give immediately

$$c_b - c_2 = \zeta c_3, \qquad c_3 - c_5 = -\kappa\eta c_b, \qquad c_2 = (1 - t\zeta)c_5. \tag{SD9}$$

Hence $Vc_b - c_2 = \zeta c_5$ and $Vc_3 - c_5 = -\kappa\eta c_2$. The remaining three balances are the following short polynomial identities; substituting the displayed P_4, P_α, P_d proves them by multiplication and collecting terms:

$$\begin{aligned} \mathfrak{Y} &:= 2\kappa c_2 + 2VL_c c_4 - 2\kappa^2 \zeta Q_c c_\alpha - Q_c c_d = 0, \\ \mathfrak{S} &:= \kappa L_c c_2 + c_5 + VL_c^2 c_4 + \kappa Q_c^2 c_\alpha + \tfrac{1}{2}(\kappa\zeta L_c^2 - 2L_c - \eta)c_d = 0, \\ \mathfrak{V} &:= \kappa R c_2 + t\kappa V_c c_b + aV_c c_3 + Tc_5 + KV_c L_c^2 c_4 - \kappa^2 \zeta K Q_c L_c c_\alpha - \tfrac{K}{2}(Q_c L_c + \eta)c_d = 0. \end{aligned} \tag{SD10}$$

For example, after the common factor $4L_cQ_c$ is removed, the first balance is just

$$\kappa K(1 - t\zeta)VH_c + VP_4 - \kappa^2 \zeta H_c P_\alpha - Q_c P_d = 0.$$

The two other balances are equally direct identities in the same three short polynomials; no expanded scalar or ideal calculation enters.

The exact contact, in the original coordinates of (SD2), is

$$u = k, \quad v = \Delta k^2 \eta, \qquad (W, X, Y, Z) = (\Delta k, -k(1 - t\zeta), \Delta k^2 L_c, 1). \tag{SD11}$$

At this point

$$r_2 = Mk\zeta, \quad B = tk\zeta, \quad r = av, \quad r_5 = -Jv, \quad vY - 1 = -Q_c, \quad D = K, \quad D_v = V.$$

Set $\Psi = \Phi/(4RT)$ and let $E_s = \frac{1}{2}\partial_s \Psi$, with all other original testing coordinates held fixed. Direct differentiation of the seven terms in (SD2), followed by (SD9), gives at (SD11)

$$\Psi = K\zeta^2 \mathfrak{S},$$

$$
\begin{aligned}
E_W &= kv\{\zeta c_2 + \Delta^2(Vc_3 - c_5)\} = 0,\\
E_X &= k\zeta\{Vc_b - c_2 - \zeta c_5\} = 0,\\
E_Y &= \tfrac{1}{2}K\Delta^3\kappa\eta^2\mathfrak{Y},\\
E_u &= k\zeta^2\{\mathfrak{S} - c_2 + (1 - t\zeta)c_5\},\\
E_v &= v\Delta^2\mathfrak{V}.
\end{aligned}
$$

They all vanish by (SD9)–(SD10). Homogeneity in (W, X, Y, Z) also gives $E_Z = 0$. This proves contact and full first-order stationarity directly in the native source.

In (SD5)–(SD6), this point is exactly

$$(u, w, v, N, M_B) = (k, \delta k, \delta k^2 b^2, 0, 0).$$

All four packets (SD3) and their first derivatives vanish there. Changing coordinates and multiplying by $\mathcal{C}$ therefore gives $(F_0)_c = (\partial_u F_0)_c = (\partial_w F_0)_c = (\partial_v F_0)_c = (L_1)_c = (L_2)_c = 0$. The leading polynomial P is a sum of circuit squares vanishing at this contact, so $P \in J^2$. This proves every identity (CC7) without using the expanded-source checker as a premise.

Positive correction weight without a large expansion. Use the weights $(2, 2, 6, 0, 0, 2, 4, 18)$ on $(\delta, k, b, a, t, u, w, v)$. Write $O_{\mathrm{wt}}(s)$ for a polynomial all of whose monomials have weight at least s. Native η and ζ have weights at least 12 and 20. Setting them to zero in the seven short coquotients, and then applying (SD7), gives

$$C_i = 2K^6 + O_{\mathrm{wt}}(12) \quad (i = 2, b, 3, 5, 4, \alpha), \qquad C_d = 4K^7 + O_{\mathrm{wt}}(12).$$

Also

$$Q = K + O_{\mathrm{wt}}(16), \quad T_n = K^2 + O_{\mathrm{wt}}(24), \quad V_n = K^2 + O_{\mathrm{wt}}(24), \quad D_v = 1 + O_{\mathrm{wt}}(36),$$

$$Y_n = O_{\mathrm{wt}}(6), \quad B_0 = -K^3 + O_{\mathrm{wt}}(24), \quad A_c = 2K^8 + O_{\mathrm{wt}}(12), \quad \mathcal{C} = K^{12} + O_{\mathrm{wt}}(16).$$

In the five factored scalar terms, discard only coefficients carrying weight at least 12. The surviving expression is exactly

$$8K^{14}D\{D(A^2 + B^2) + K(C^2 + D_0^2) - 2K^2BC\} = 8K^{14}DP.$$

Here A^2, D_0^2 have weight 44, while B^2, C^2, BC have weight 40. Consequently $E_0 = O_{\mathrm{wt}}(52)$, sufficient for every scalar lift coefficient to have weight at least 8. For the normals use (SD9) in its polynomial forms

$$K(C_b - C_2) = \delta^2 k^2 Q b^2 C_3, \qquad Q(C_3 - C_5) = -k^2 K b^2 C_b.$$

Since K, Q have constant term one, these show weights at least $20, 16$ for the two differences, respectively, without putting their inverses in any lifting coefficient. The displayed S_B, S_C in the conductor proof have weight at least 12. Its factored normal formulas now give $L_1 = O_{\mathrm{wt}}(42)$ and $L_2 = O_{\mathrm{wt}}(32)$. The monic lift therefore has positive coefficient weights at least $8, 20, 10$. The stronger exact checker bounds $12, 20, 12$ are optional.

Full hard block and restoration. There are no state-normal terms above degree two. Put

$$C_B = H_3 D_v h + H_5(1 - h), \qquad B_Y = 4RT\Delta^2\eta^2 c_4 D D_v + 4RT\Delta^4 k^6 \eta^2 c_\alpha D v^2 + 2RT\Delta^2 \eta c_d D v^2.$$

Before multiplication by $\mathcal{C}$, the exact quadratic block on $(N, \delta M_B)$ has entries

$$
\begin{aligned}
A_{11} &= H_2 + H_b D_v + \Delta^2 C_B u^2 v^2 + B_Y h^2 u^2 v^2 / D^2,\\
A_{12} &= -\frac{\Delta^2}{\delta}(H_3 D_v - H_5)uv - \frac{B_Y}{\delta D^2} huv,
\end{aligned}
$$

$$A_{22} = \frac{\Delta^2}{\delta^2}(H_3 D_v + H_5) + \frac{B_Y}{\delta^2 D^2}.$$

The displayed ratios cancel coefficient monomials exactly: $\Delta/\delta = Q/K$ is a positive unit and B_Y/δ^2 is bounded infinitesimal. Thus this full block has residue $16I_2$, also after multiplication by $\mathcal{C}$. The conductor proof and the positive circuit frame give the bounded strict-residue Gram for $\mathcal{C}G$. Restore the four full positive packets $\mathcal{C}\Pi_j$ from (SD3), then undo $\mathcal{C}\Delta^2\Lambda$. This recovers the entire original parent; retained gap ratios need not be bounded and are never placed in that bounded Gram. The complementary state and outer-angle charts require the already proved C1 chart arguments. This appendix makes no claim of an exhaustive global source cover.

Chapter 127

The C1 circuit needs six soft directions

Component `p58894e12bcef`
Source: `CORE/prior/prior_core/circuit/circuit_frame.tex`
SHA256: `58894e12bcef3137ea80fb46ac9c63596adbf4d664874009ded5c97ff818dc77`
Inventory use: compact-circuit identity and its 3x3 contrast Gram; g4 identity only with importing-row coefficient bounds | C1_BARE_OTHER_TERMINALS; precise subordinate uses retained below

127.1 The C1 circuit needs six soft directions

Let

$$a_1 = \sqrt{wv}, \quad a_2 = \sqrt{\delta uv}, \quad a_3 = b\sqrt{\delta kuw}, \quad a_4 = \delta k^{3/2} b.$$

The radicals are separated coefficient/testing radicals. Put

$$h_{13} = a_1^2 - a_3^2, \quad h_{23} = a_2^2 - a_3^2, \quad h_{34} = a_3^2 - a_4^2, \qquad q_1 = a_1a_2 - a_3a_4, \quad q_2 = a_1a_3 - a_2a_4.$$

Averaging the three-square arithmetic–geometric-mean identities for pairings 12|34 and 13|24 gives exactly

$$\sum_{i=1}^{4} a_i^4 - 4a_1a_2a_3a_4 = h_{13}^2 + h_{23}^2 + h_{34}^2 - h_{13}h_{23} + h_{23}h_{34} + q_1^2 + q_2^2. \tag{127.1}$$

The contrast block is

$$G = \begin{pmatrix} 1 & -1/2 & 0 \\ -1/2 & 1 & 1/2 \\ 0 & 1/2 & 1 \end{pmatrix} \succ 0; \qquad \det G_{[1]} = 1, \quad \det G_{[2]} = 3/4, \quad \det G = 1/2.$$

Unlike a single pairing, the two pairings control all three independent contrasts of the four squared monomials.

Write $K = 1 + k^2$, $D = 1 + u^2$ and

$$g_0 = \delta(v - b^2uw), \qquad g_4 = uwv - \delta^2k^4b^2.$$

The leading scalar used in the complete source is

$$P = D(w^2 + \delta^2)v^2 + \delta^2Kb^4u^2w^2 + \delta^4Kk^6b^4 - 2\delta^2K^2b^2uwv = g_0^2 + g_4^2 + \sum_i a_i^4 - 4\prod_i a_i.$$

There is a second exact simplification:

$$g_4 = uh_{13} + kh_{23} - kug_0 + kh_{34}. \tag{127.2}$$

All four displayed coefficients are infinitesimal in the allowed chart. Thus g_4^2 can be retained as a positive square or absorbed as a zero-residue Gram correction. Equations (127.1) and (127.2) give a strict residue Gram on just

$$\boxed{(g_0, h_{13}, h_{23}, h_{34}, q_1, q_2).}$$

The first-order and restricted second-order conductor theorem places every source correction in the first five of these roots. The two hard state normals therefore give an *eight-root* bounded strict-residue Gram for the normalized remainder after retaining the four positive packets, replacing its previous thirteen-root frame. Restoring those packets gives the entire parent. Their possibly unbounded gap ratios are kept outside this bounded Gram. This is a local Gram-frame dimension, not a global SDP block-size bound or a claimed optimal number of tensor squares.

One pairing alone does not suffice. At the formal complex point $(a_1, a_2, a_3, a_4) = (1, 1, i, -i)$, its roots $a_1^2 - a_2^2$, $a_3^2 - a_4^2$, $a_1 a_2 - a_3 a_4$ all vanish, whereas $h_{13} = 2$. Hence their ideal does not contain h_{13}, and their squared ideal cannot absorb every infinitesimal multiple of h_{13}^2. This is an algebraic obstruction to that proposed reduction, not a physical complex-state counterexample. The second pairing supplies exactly the missing strict contrast direction.

Chapter 128

An exhaustive positive-d source partition

Component `pfb270019c56f`
Source: `CORE/prior/prior_core/inherited/positive_d_source_partition.tex`
SHA256: `fb270019c56f27e2c8a4ea33f4d82e8514970cd4c4e2a28aca98ffb85a4d502b`
Inventory use: Only displayed SP1-SP3 coefficient factorization and angular-balance proofs; missing terminal references are replaced by AU15-AU18 | Only actual SP2-SP3 equations and their bounded-source proof; no missing terminal reference is imported

128.1 An exhaustive positive-d source partition

Presentation scope note. This manuscript imports this retained component only for the source partition (SP1)–(SP3) and its derivation. Historical all-tests terminal calls below are supplied by the complete replacement arguments named at their points of use; they are not additional imported premises. The original source file is unchanged.

Throughout this section the coercive canonical source is bounded and physical, d is a positive unit, and an actual finite noncommuting stationary zero contact exists. Write $U = u_c^2, V = v_c^2$. For nonnegative pure source quantities, $a \asymp b$ means that a/b and b/a are bounded positive units. This notation is used only inside K_c.

Lemma 128.1 (Two forbidden weak faces)**.** *Every positive-d C2 source has A a positive unit. In C1, at least one of m, c is a positive unit.*

Proof. The following identities for the central determinant coefficient follow by substituting into the displayed four-by-four parent:

$$h\big|_{\lambda=\beta=A=0} = -d(d+2n)(2D+2c+d)(2\alpha+d+2m),$$
$$h\big|_{\lambda=n=m=c=0} = -d(2D+d)(2\beta+d)(2A+2\alpha+d).$$

Both are strictly negative when $d > 0$. If $\bar{A} = 0$ in C2, the residue Laurent polynomial has only the terms $h + q_{01}/U + q_{02}V/U + q_{12}V$. Substituting $U = t^2, V = 1/t$ and sending the real number t to infinity gives the negative limit h, contradicting physicality. If $\bar{m} = \bar{c} = 0$ in C1, its residue Laurent polynomial is $h + q_{10}/V$; letting V tend to infinity gives the same contradiction. These are arguments over the real residue source and involve no tensor division. □

Proposition 128.2 (The positive-d C2 source partition)**.** *The C2 source is covered by the existing small-α all-tests theorem or its exact stabilizer image unless α, c, A are units. In the latter case the actual contact satisfies*

$$U \text{ bounded}, \qquad V^{-1} \text{ bounded}, \qquad U^2 \asymp m, \qquad DV^2 \asymp 1. \tag{SP1}$$

Consequently its four exhaustive source types are:

1. *m, D units: u_c, v_c bounded nonzero in residue;*

2. *m weak, D a unit: u_c infinitesimal and v_c a bounded unit;*

3. *m a unit, D weak: u_c a bounded unit and v_c unbounded;*

4. *m, D weak: u_c infinitesimal and v_c unbounded.*

The first three types have all-tests certificates from the preceding modules. The fourth, with n arbitrary, is the common reciprocal type completed by RC1–RC25 (p. 739), fitted to this source in AU16 and AU19 (p. 695).

Proof. The stabilizer $(\lambda\ \beta)(m\ D)(c\ \alpha)$ exchanges c, α and acts on the absolute half-angles by $(|u|, |v|) \mapsto (1/|v|, 1/|u|)$. At measurement level it exchanges the parties and sends $(A_0, A_1; B_0, B_1)$ to $(-B_0, B_1; -A_0, A_1)$, followed by local basis changes. Thus either weak residue enters the exhaustive CU/WA/WN union in AU15 (p. 695). Lemma 128.1 gives the unit A. In the remaining branch write

$$q_{01} = ma_1, \quad q_{02} = mDa_2, \quad q_{10} = a_3, \quad q_{12} = Da_4, \quad q_{20} = na_5, \quad q_{21} = a_6,$$

where all a_i are bounded positive units. At the actual contact, $L(U, V) = 0$ and every noncentral term is nonnegative. Since h is bounded, $U, V^{-1}, m/U, DV$ are bounded. Angular stationarity is

$$(q_{02}/U + q_{12})V^2 = q_{10} + q_{20}U, \qquad U^2(q_{20} + q_{21}V) = mV(a_1 + a_2DV).$$

In the first equation the right side is a positive bounded unit, and the left side is $DV^2(a_4 + a_2m/U)$ with a positive-unit parenthesis. Thus $DV^2 \asymp 1$. Dividing the second equation by V gives $U^2(a_6 + na_5/V) = m(a_1 + a_2DV)$; both parentheses are positive bounded units, so $U^2 \asymp m$. This proves (SP1) and the exhaustive partition. Coercivity ensures $m, D > 0$ as actual source quantities; the same fact also follows from the two equations.

For the first type, the direct all-tests argument in Lemma 128.4 below applies. The second has the natural pitchfork entry of S (p. 584) under the source fit and all-tests argument in AU7–AU9 (p. 695); Lemma 128.4 promotes that terminal to all tests. The stabilizer exchanges m, D, so the third type is the exact image of the second. No restriction on n has been used. □

Proposition 128.3 (The positive-d C1 source partition)**.** *For C1, weak α or weak A is already covered by an all-tests theorem and its stabilizer image. Otherwise α, A are units. After the C1 stabilizer, c may be taken to be a unit. The remaining exhaustive source alternatives are:*

1. *m is a unit and D or β is a unit: the source has an all-tests certificate;*

2. *m is weak and D is a unit: the source has an all-tests pitchfork certificate;*

3. *m is a unit and both D, β are weak: the actual contact has u_c a bounded unit and v_c unbounded;*

4. *m, D are weak: the actual contact has u_c infinitesimal and v_c unbounded, with β arbitrary.*

The third type is now covered at every test by PP1–PP19 (p. 725), fitted to this source in AU18–AU19 (p. 695). The fourth is covered at every test when β is a unit by Corollary 128.5 below. Thus only its weak-β subcase remains after these reductions; it is completed by CR1–CR39 (p. 674), fitted to this source in AU18–AU19 (p. 695).

Proof. The C1 stabilizer $(\lambda\ n)(m\ c)(A\ \alpha)(D\ \beta)$ exchanges A, α. Hence WN1–WN44 (p. 718), together with the general entry in WA1–WA10 (p. 713) and AU18 (p. 695), cover either weak case. The forbidden-face lemma permits an orientation with c a unit in the remaining branch. Now

$$q_{01} = ma_1, \quad q_{02} = mDa_2, \quad q_{10} = a_3, \quad q_{12} = \beta ma_4 + Da_5, \quad q_{21} = a_6, \quad q_{22} = \beta a_7,$$

where all a_i are positive bounded units. The contact equality bounds each positive Laurent term, so $U, V^{-1}, m/U, DV$ are bounded. The U stationarity equation reads

$$U^2(a_6 + \beta a_7 V) = m(a_1 + a_2DV). \tag{SP2}$$

Its right parenthesis is a positive bounded unit, and its left parenthesis is bounded below by a positive unit. Therefore weak m forces U infinitesimal. If m is a unit, boundedness of m/U forces U to be a unit. The other stationarity equation is

$$a_3 = V^2\{mDa_2/U + \beta m a_4 + Da_5 + \beta a_7 U\}. \tag{SP3}$$

If m is a unit, so is U, and the braces are comparable with $D + \beta$. Thus V is a unit exactly when D or β is a unit, and is unbounded exactly when both are weak. If m is weak, U is infinitesimal. If V is bounded, (SP2) gives m/U weak, and (SP3) forces D to be a unit. Conversely a unit D makes V bounded in (SP3), so the remaining branch has V unbounded. These alternatives exhaust all cases.

In the first type apply the stabilizer if necessary to arrange D a unit; then d, c, A, D, α, m are units and Lemma 128.4 applies. In the second, AU17–AU18 (p. 695) supplies the exact finite pitchfork entry for weak m and unit D; AU7–AU9 (p. 695) with the complete terminal S (p. 584) supplies the all-tests conclusion. □

Lemma 128.4 (All-tests promotion for strong and pitchfork sources)**.** *Let C1 or C2 have d, c, A, D, α units. If m is a unit, it has a tensor certificate at every independent projective test. If m is weak and its actual contact has u_c infinitesimal and v_c a bounded unit, the same conclusion follows from its finite-angle pitchfork theorem.*

Proof. All noncommuting residue tests are covered by Theorem 11.2. At the zero-zero angle corner, a zero state is the W ray because α is a unit. Normalize $W = 1$ and retain the entire nonnegative m packet. Every remaining monomial is in $(u, X, Y, Z)^2$. Its quadratic residue is

$$2(d + 2\beta + 2n)X^2 + (4c + 4D + 2d)(u - Y)^2 + 4AY^2 + 4\alpha Z^2,$$

which is positive definite in (u, X, Y, Z). Finite polynomial allocation therefore gives an exact strict-residue Gram; the retained packet is restored. This works whether m is weak or a unit and includes every bounded projective state chart near the unique zero ray.

At $v = 0, u \neq 0$, the zero-state classification excludes zeros because A, α are units. At $u = 0, v \neq 0$, it excludes zeros if m is a unit. If m is weak, the sole zero state is the W ray. The appropriate pitchfork theorem covers the matched remote residues $v = \pm\bar{v}_c$; its printed positive quartic boundary factor covers every unmatched finite remote residue by the strict four-root Gram. Its projective remote endpoint is also strict.

All other projective angles use the exact homogeneous chart identities in AU9 (p. 695) and the complete single-infinity identities immediately following it. Their hypotheses here are precisely the units d, c, A, D: at $(\infty, 0)$ use $(pX + Z, Y, W, vX)$, at $(0, \infty)$ use (X, Y, uW, Z), and at (∞, ∞) use the five-root quadratic form on (W, X, Y, p, q) after retaining the β, α, n packets. Those displayed residue forms are positive definite independently of the residues of m, n, α, β. One infinite angle with the other finite nonzero has no kernel by the same c, A, D square identities. This exhausts both projective angle lines and all state charts; positive pure testing factors restore the original slack. □

Corollary 128.5 (All tests for the strong-β reciprocal source)**.** *Under the unit-β subcase of CR1–CR39 (p. 674) for weak m, D, fitted to the source in AU18–AU19 (p. 695), the C1 source has a tensor certificate at every independent projective test, not only at its reciprocal testing corner.*

Proof. That source fit gives d, c, A, α, β units, m, D weak, $\bar{h} = 0$, and every state at the corner $(u, v) = (0, \infty)$. Theorem 11.2 handles every noncommuting residue test. At $(0, 0)$ use the strict four-root Gram in Lemma 128.4; its displayed residue remains positive when D is weak, since c, A, α, d are units. At $v = 0, u \neq 0$ the units A, α exclude a zero state.

At $u = 0$ and finite nonzero v, the only zero state is the W ray. Retain the whole m packet and use (u, X, Y, Z) as roots. The real hard block is the positive X mass together with

$$\{4(c + A) + 2d\}(1 + v^2)Y^2 + 4\alpha(vY - Z)^2,$$

which is positive definite. The scalar angular Schur coefficient is the pitchfork boundary expression

$$f(v)=\frac{8(1+v^2)}{t_7\det B(v)}(q_{10}+\bar{h}v^2+\bar{q}_{12}v^4)=\frac{8(1+v^2)q_{10}}{t_7\det B(v)}>0.$$

Thus the exact four-root Gram has positive definite residue. All divisions here describe a real residue calculation, not an operation in the tensor ring.

At $(\infty,0)$ the existing four-root identity has residue

$$2dR^2+2d(W+V)^2+4cW^2+4AY^2\succ 0,$$

so it does not need a unit D. At (∞,∞) the residue state matrix is positive definite since c,β,d are units. For $u=\infty$ and finite nonzero v, the signed core together with the c,A,β packets contains

$$2d\{(W+vX)^2+(vY-Z)^2\}+4c(1+v^2)W^2+4A(1+v^2)Y^2+4\beta(1+v^2)Z^2\succ 0.$$

For $v=\infty$ and finite nonzero u, use instead

$$2d(1+u^2)(X^2+Y^2)+4c(uW-Y)^2+4\beta(X+uZ)^2\succ 0.$$

These checks exhaust the projective angle charts. The stated reciprocal theorem supplies the sole remaining double corner. □

Chapter 129

The joint small-a, small-e corner: a bounded-ratio closure

Component `p26da50239f47`
Source: `R16/inherited_r14/proofs/weak_local/bounded_ratio_closure.tex`
SHA256: `26da50239f4701b3f3d5afaaa21060d5a09e8919997fdbcd541f2d4bb20b0f79`
Inventory use: R9B2 and its joint monic source-regularity proof

129.1 The joint small-a, small-e corner: a bounded-ratio closure

Theorem 129.1. *Retain the full physical and stationary source domain R9 of (WF3), including all inherited exclusions, $n \geq 0$, and independent source and testing fields. For every local testing pair $u, v \neq 0$ for which $w = v/u$ is bounded in the testing field, the entire original C2 parent is finite tensor SOS, with every state component retained. Literal axes are also covered. The theorem does not cover the remaining local chart v/u unbounded, and therefore is not an all-tests admission of R9.*

The proof uses the exact identities (WC3)–(WC17a), not the unit-a contact-lifting argument. Take $u, v > 0$ using the inherited exact sign gauges. The source quantities B, a, h, e are positive infinitesimals, t is a unit with $t_0 > 1$, and $r = 1 - e$. Put

$$c = \sqrt{a/t}, \quad k = \sqrt{aB}, \quad A_1 = uv, \quad A_2 = ck\frac{v^2}{u}, \quad A_3 = \frac{k^2}{c}, \quad G = kv, \quad F_v = v^2 - tB.$$

All divisions and radicals are pure-field operations. In particular $w = v/u$ is bounded and c is infinitesimal. The crucial identity is

$$\boxed{G^2 = (cw)A_1A_3.} \tag{R9B1}$$

Thus the negative circuit term is an infinitesimal off-diagonal coefficient in the (A_1, A_3) frame. No circuit conductor and no comparison of source and testing rates is needed.

Joint source regularity at $a = 0$. Use the unchanged definitions $q_s, \widetilde{m}, c_F, f_A, f_B, f_C, f_G$ from (WC1), (WC6), and (WC8), and put $f_{C0} = f_C|_{U=0}$. Unlike $\partial_U f_C$, all coefficients needed below are jointly regular at $B = a = h = 0, r = 1$. More explicitly,

expression	joint residue
c_F	$1/(2t_0)$
$f_A\|_{U=V=0},\ f_B,\ f_{C0}$	$1/2$
f_G	$3/2$
$\widetilde{m}/B$	0
$K_X/B,\ K_E/B,\ K_Y/B$	$t_0/2,\ -1/2,\ 0$
$d_4,\ b_4$	$t_0/[2(t_0-1)],\ 0$
α/B^2	$t_0/2$

(R9B2)

These are finite rational identities. To verify them, use the unit common denominator

$$\mathcal{D} = 2rg(1-h)(1-hB)(t-1+h+aBtr)\Delta$$

from (WC3), insert the exact source numerators, and divide numerator and denominator by each explicit monic factor B and a. The required numerator remainders vanish identically, not just at the corner. Each remaining denominator has a nonzero corner value. The accompanying focused checker performs precisely these divisions and prints the denominator and residue for every row. There is no e/B, h/a, or mixed-rate regularity assumption. All remaining source coefficients of (WC17a) are rational with the same unit denominators and are therefore bounded on this chart.

A strict scalar Gram without a contact lift. Define the legal separated roots

$$T_q = \sqrt{q_s}\, u, \qquad T_n = \sqrt{n}\, ru^2, \qquad T_\alpha = \sqrt{\alpha}\, u, \qquad d = 1 - \frac{a}{t}w^2.$$

Here $q_s > 0$ and $n \geq 0$; literal $n = 0$ gives the zero root T_n and requires no division. Also d has residue one. The exact scalar identity (WC9) becomes

$$\mathcal{S} = d^2(T_q^2 + T_n^2) + c_F F_v^2 + f_A A_1^2 + f_B A_2^2 + f_{C0} A_3^2 - f_G(cw)A_1A_3 + T_\alpha^2. \tag{R9B3}$$

Indeed $H_u = -du$ and the formerly problematic term is exactly

$$(f_C - f_{C0})A_3^2 = \alpha u^2.$$

It is retained as a square, not treated as a bounded perturbation. Since $f_A = f_A|_{U=V=0} + d_4U + b_4V/t^2$, (R9B2) proves that (R9B3) has a bounded Gram with strictly positive diagonal real residue on

$$T_q, T_n, F_v, A_1, A_2, A_3, T_\alpha.$$

Its only displayed off-diagonal entry is infinitesimal by (R9B1).

Every original state normal. Use exactly the full-state frame (WC4),

$$E = tuW - v, \qquad N_y = (Y - huW)/\sqrt{h}, \qquad \xi = X + ru,$$

initially at $Z = 1$. The original hard block has the unchanged strict residue (WC5), because its rational coefficients have no negative power of a and (WF6)–(WF7) supply its source limits. The extracted angular contributions in (WC15) are infinitesimal linear combinations of T_q, T_n: simply replace $R_q = -dT_q$ and $R_n = -dT_n$. Their coefficients remain bounded infinitesimals even when $n = 0$.

For the complete residual normals, use their exact polynomial arrays (WC17a). Every term is either Buw^j with $0 \leq j \leq 3$ or $u^i w^j$ with $i \geq 3, j \geq 1$, with bounded pure-source coefficients by (R9B2). The following exact allocations have only infinitesimal coefficients:

$$\boxed{Buw^j = \frac{uw^{j+1}}{t}A_1 - \frac{uw^j}{t}F_v, \qquad u^i w^j = u^{i-2}w^{j-1}A_1.} \tag{R9B4}$$

The first identity follows from $tB = v^2 - F_v$ and $A_1 = uv$; the second is immediate. These identities include the formerly problematic Bu term, and never divide by a, k, or a mixed element. Hence all three full first normals are infinitesimal linear combinations of roots already carrying strict scalar pivots. The $\sqrt{h}$ multiplying the N_y normal is bounded and causes no difficulty. Joining them with the hard block gives an exact finite bounded Gram with strictly positive real residue. The inherited bounded-Gram lemma now proves tensor SOS for the complete parent.

Homogenize by Z^2 exactly as after (WC28), so all roots become homogeneous state-linear expressions and no Z inverse remains. The resulting identity includes $Z = 0$. The original sign gauges handle all signs, the two exact positive-axis identities (WC28) handle literal axes, and real/imaginary splitting handles complex states. No norm, original packet, or hard normal was discarded.

Exact remaining scope. Together with the R13 remote reduction, this leaves on R9 only $u, v \neq 0$ both infinitesimal with $\omega = u/v$ infinitesimal. The new identity (R9B1) does not claim cw bounded when w is unbounded. That reciprocal local chart remains an explicit proof obligation. The accompanying 23 exact checks (including twelve joint source regularity checks) verify the new finite identities; they are not a replay of an older suite.

Chapter 130

Two exact conductors for the remaining small-η chart

Component `peea9e9c546c5`
Source: `R16/inherited_r15/proofs/c1_contact/small_eta_conductors.tex`
SHA256: `eea9e9c546c59dd6fd32b36f9b1fa097fdef72e19bf518cd5797e3a2f1ba7bbe`
Inventory use: SC1–SC5 and Exact full angular cancellation paragraph; stop before The complete radial group | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

130.1 Two exact conductors for the remaining small-η chart

These identities resolve specified higher-angle terms; they do not constitute an admission of C14.3b. Keep

$$c^2 = 1/(\eta\zeta), \quad k^2 = x, \quad y = x/c^2, \quad A = uv, \quad B = ckv^2/u, \quad C = k^2/c, \quad G = kv,$$

$$d = B - G, \quad J = A - C, \quad F = V - y, \quad p = ckv/u, \quad R_c = pE + d.$$

The elementary conductor is

$$ud = k(cF - J). \tag{SC1}$$

Consequently, for the pure-source coefficient $e_4 = a_{UVV}$,

$$e_4 xvEF = e_4(k/c)uR_cF - e_4xF^2 + e_4(x/c)JF. \tag{SC2}$$

This is useful after the full angular completion. The exact scalar remainder obeys

$$(c_F + e_4x)F^2 + e_4vR_DF = (c_F + e_4(U + x))F^2 - e_4yuH_uF.$$

Substituting $H_u = Q/\sqrt{q} - a(v/u)E/\eta + a\xi$ and using SC2 gives

$$\begin{aligned}(c_F + e_4U)F^2 - e_4yuQF/\sqrt{q} - e_4yau\xi F \\ + e_4(k/c)uR_cF + e_4(x/c)JF.\end{aligned} \tag{SC3}$$

Here $a = 1/\zeta$. On the source chart with η, x infinitesimal and ζ a positive unit,

$$\eta\lambda \sim \tfrac{1}{2}, \quad \eta^2 c_F \sim \tfrac{1}{2}, \quad \eta^2 e_4 \sim \tfrac{1}{2}, \quad q/x^2 \text{ is a positive bounded unit.}$$

Thus the normalized coefficients of the last two terms on $(\sqrt{\lambda}R_c, \sqrt{\lambda}J, \sqrt{c_F}F)$ are $O(\sqrt{x}\,u)$ and $O(x)$. The Q, F coefficient is $O(u)$; the ξ, F coefficient is $O(xu)$, including normalization by $\sqrt{B_x}$ since $B_x \geq 1/2$. Retain e_4UF^2 as an exact square. Finally $\sqrt{\lambda}R_c = R + \sqrt{\lambda}kE$ is bounded on the coupled root R and the weighted hard direction $E/\sqrt{\eta}$. No bound on the bare d is used.

For the complete divergent D event, write the source block in terms of $P = (1 - r)E/\eta - \nu$, $Q_t = rE/\eta + \nu$, and $b = Ar/\eta - (m\theta + \alpha + 1/2) = O(x)$. Put $T_D = R_D - uv\xi/h$. Then the

complete original $D_p = D + 1/2$ event is $D_p\{P - hT_D - (j/\eta)vZ\}^2$. Its centered quadratic packet is $D_p(P - hT_D)^2$; the contact-linear and scalar terms involving $(j/\eta)vZ$ must remain in the exact contact balances. Replacing R_D by T_D in the centered cross term $2bQ_tR_D$ leaves $2(b/h)uvQ_t\xi$. The exact allocation is

$$uvQ_t = \frac{rU}{\eta ck}R_c - \frac{ru}{\eta}F + \frac{ru}{\eta c}J + uv\nu. \tag{SC4}$$

After multiplication by b, the coefficients on $\sqrt{\lambda}R_c, \sqrt{c_F}F, \sqrt{\lambda}J, \nu/\sqrt{\eta}$ are respectively

$$O(\sqrt{x}\,U), \quad O(xu), \quad O(xu), \quad O(xuv\sqrt{\eta}).$$

Thus this particular full-D cross term is harmless without comparing uv to $\sqrt{\eta}$.

Exact full angular cancellation. Put $D_0 = u - cv$ and write the full angular root as $\mathcal{Q}$. There are exact identities

$$R_c = kw(cE - D_0), \qquad \mathcal{Q}/\sqrt{q} = -D_0 + (c/k)R_c - a\xi.$$

Thus the complete principal ξ row cancels to

$$-2\eta\lambda ckR_c\xi + 2\eta\lambda c(R_D - uJ)\xi = -2\eta\lambda x\mathcal{Q}\xi/\sqrt{q} - 2\eta\lambda xa\xi^2. \tag{SC5}$$

Together with the singular rank's $+\mu\xi^2$, the last term leaves $-\mu\xi^2$, since $\mu = \eta\lambda ax$; this is infinitesimal relative to $B_x\xi^2$. The $\mathcal{Q}, \sqrt{B_x}\xi$ block has strictly positive residue. Indeed, the exact source limits give

$$B_xq/x^2 \sim \frac{B_x\zeta(1+\zeta)}{2} + \frac{(1+\zeta)^2}{4} > \frac{1}{4} \sim (\eta\lambda)^2,$$

where for unbounded B_x the normalized cross tends to zero. For bounded $B_x \geq 1/2$, the difference has the explicit positive lower bound $\zeta(3+2\zeta)/4$. No inequality $B_x > 1/(2\zeta)$ is needed.

The complete radial group, and the remaining dependency. Define $\overline{F} = F + vE$ and $H_{\text{full}} = \mathcal{Q}/\sqrt{q} + a\xi = H_u + c^2wE$. Then exactly

$$uR_c = k(c\overline{F} - J), \qquad U - x = c^2\overline{F} - uH_{\text{full}}. \tag{SC6}$$

The entire c_F group is $c_F\overline{F}^2$. The entire e_4 group, including both hard and first-normal terms, has the following exact simplification:

$$\begin{aligned} &e_4\{U\overline{F}^2 + y(U-x)(\overline{F} + vE)\} \\ &= e_4(U+x)\overline{F}^2 - e_4yu(\mathcal{Q}/\sqrt{q} + a\xi)\overline{F} + e_4yER_D. \end{aligned} \tag{SC7}$$

The normalized $\mathcal{Q}, \overline{F}$ and $\xi, \overline{F}$ coefficients are $O(u)$ and $O(xu)$. The last term is an $O(x)$ coefficient perturbation of the weighted source block because $e_4y\eta = O(x)$. It should be incorporated before applying the full D_p state shift.

There remains a specific root dependency: SC4 allocates the shifted $uvQ_t\xi$ term on the *bare* F root. Completing the full radial square retains $\overline{F}$, not an independent strict F. Replacing $F = \overline{F} - vE$ in SC4 merely yields a tautology by SC6. Therefore SC4 and SC7 cannot simply be joined while counting both radial pivots. A possible completion must center the m event jointly with the other packets. The whole original mR_m^2 cannot be extracted as a standalone positive square: at actual source contact $R_m = \zeta MuZ \neq 0$, although the parent vanishes. The relevant exact centered identity is

$$R_m - \zeta MuZ = \frac{\mathcal{Q}}{a\sqrt{q}} + \frac{r}{\eta}u\widetilde{F}, \qquad \widetilde{F} = \frac{\eta}{r}vY - yZ, \quad M = 1 + xr - a\theta. \tag{SC8}$$

It preserves contact and indicates where the uvQ_t control could enter a simultaneous completion; the favorable source quotient $b/\sqrt{m} = O(\sqrt{x})$ alone does not supply that completion. The contributions $2mr/\eta^2$ and mr^2/η^2 already counted in the radial groups must also be reconciled exactly. Such a contact-centered joint reassembly has not been proved here. The displayed identities resolve the principal angular and full radial obstructions but do not yet admit C14.3b.

Chapter 131

The weak-C2 R9 contact corner: a full-parent coupled-circuit certificate

Component `pb18dc55f4f2f`
Source: `R16/inherited_r15/proofs/r9_contact/reciprocal_contact_closure.tex`
SHA256: `b18dc55f4f2fb27fe0c98a7478243f59e43e1b958a2b565dd8e423ff64bf9dc0`
Inventory use: RC1-RC13

131.1 The weak-C2 R9 contact corner: a full-parent coupled-circuit certificate

Scope. Retain the original R9 physical stationary source and all inherited intersections. Thus B, a, h, e are positive infinitesimals, $r = 1 - e$, t and j are units with $t_0 > 1$, and $n \geq 0$. Consider arbitrary independent $u, v > 0$ infinitesimal, with $\omega = u/v > 0$ unrestricted. The following is a certificate for the entire original parent, not only its contact graph. Its bounded Gram coefficients contain no ω or $1/\omega$, so it covers every positive testing ratio at once. The original axes/sign identities and R13 remote theorem then close all tests on R9.

All formulas below refer to $F/4$. Use the unchanged WC3, WC6, WC13–WC17a coefficients, and put

$$x = aB = k^2, \quad y = tB = l^2, \quad c = \sqrt{a/t} = k/l, \quad U = u^2,\ V = v^2, \quad q = q_s + \alpha,$$

$$\mu = m - a^2 q, \quad \lambda = \frac{\mu}{ta^2 B}, \quad K'_X = K_X - a\alpha, \quad K'_E = K_E + a\alpha/t.$$

Indeed $\lambda = f_B - \alpha/y$, where the R14 exact joint-source divisions give $f_B \sim 1/2$ and $\alpha/B^2 \sim t_0/2$. Since $\alpha/y = (B/t)(\alpha/B^2)$ is infinitesimal, $\lambda \sim 1/2 > 0$. The same exact reductions give $c_F \sim 1/(2t_0)$, $d_4 \sim t_0/[2(t_0 - 1)]$ and b_4 bounded. These reductions hold on the literal $n = 0$ face, and no division by n is used. In particular all source denominators used below are positive; no comparison of source and testing rates occurs.

An exact scalar completion. At $Z = 1$ use

$$A = uv, \quad B_c = ck\, v^2/u, \quad C_c = k^2/c, \quad G = kv, \quad d = B_c - G, \quad J = A - C_c,$$

$$F_v = V - y, \qquad R_D = v(U - x), \qquad H_u = (a/t)V/u - u.$$

Thus $AB_cC_c = G^3$. Completing the entire αU term into qH_u^2 changes f_B to $\lambda = f_B - \alpha/y$ and f_G to $f'_G = f_G - 2\alpha/y$. The exact source contact balances imply

$$f_{A0} = \lambda - 2d_4 x - b_4 y/t^2, \qquad f_{C0} = \lambda + b_4 y/t^2, \qquad f'_G = 3\lambda - d_4 x + b_4 y/t^2. \tag{RC1}$$

These identities follow from the value and both first derivatives of the original scalar at $(u,v)=(k,l)$, since the extracted H_u, F_v squares vanish there; they can also be checked by the finite source formulas. Using $A^2=UV$, $G^2=xV$ and $C_c^2=xy$ now gives the exact scalar identity

$$\mathcal{S}=qH_u^2+nr^2u^2H_u^2+\left(c_F+\frac{b_4x}{t^2}\right)F_v^2+\lambda L_3+d_4R_D^2+\frac{b_4}{t^2}vR_DF_v, \tag{RC2}$$

where $L_3=A^2+B_c^2+C_c^2-3G^2$. For example the two nontrivial remainders are precisely

$$(U-2x)A^2+xG^2=V(U-x)^2,$$

$$(V-y)A^2+y(C_c^2-G^2)=V(U-x)(V-y)+x(V-y)^2.$$

This uses no limiting expansion or approximate scalar positivity.

The complete original hard block. Keep the full WC4 state frame

$$E=tuW-vZ,\qquad N_y=(Y-huW)/\sqrt{h},\qquad \xi=X+ruZ.$$

Work initially at $Z=1$. Complete the two angular squares as

$$Q=\sqrt{q}\left(H_u+\frac{a}{t\omega}E-a\xi\right)=\sqrt{q}\,(avW-uZ-a\xi),$$

$$N=\sqrt{n}\{ruH_u+u\xi\}. \tag{RC3}$$

The second formula is homogenized by replacing uH_u by $[(a/t)V-U]Z$. It includes $n=0$ without division. Subtracting Q^2+N^2 from the entire parent leaves the singular hard quadratic exactly as the single positive rank-one packet

$$\mu\left(\frac{E}{t\omega}-\xi\right)^2. \tag{RC4}$$

All other hard coefficients form a bounded quadratic $\mathcal{H}_0(E,N_y,\xi)$ with residue

$$\operatorname{diag}\left(\frac{1}{2(t_0-1)},\frac{t_0}{2},\frac{1}{2}\right). \tag{RC5}$$

This assertion concerns the full parent. To check that no hidden $1/\omega$ remains, its only such term before subtraction is the m-event coefficient $(1+hU)/(t\omega)$. After removing its leading $m/(t\omega)$ part, the E^2 error is $m(2hV+h^2UV)/t^2$, the $E\xi$ error is $-2mhuv/t$, and the EN_y coefficient is bounded by a bounded source factor times $m\sqrt{h}V$. All remaining event coefficients have no negative power of ω: $uW=(v+E)/t$ and $Y=h(v+E)/t+\sqrt{h}N_y$ make this explicit. The A event uses Ah, so no inverse power of h survives. Subtracting Q^2,N^2 changes its bounded entries only by infinitesimals after RC4 is removed. This proves RC5 directly from WC3, without a mixed inverse.

Exact centering of every first normal. Set

$$a_E=2Ah\,h/t^2-2e_1/t,\quad b_E=2b_2/t^2,\quad c_E=2b_4/t^2.$$

Let a_Y,b_Y,c_Y be the three coefficients of u^3w,u^3w^3,u^5w^3 in $\mathcal{R}_Y$ in WC17a. All are bounded. Original contact of the complete normals gives

$$2K_E'+a_Ex+b_Ey+c_Exy+2\lambda x=0,$$

$$2K_X'+x_3y+x_5xy-2t\lambda x=0,\qquad 2K_Y+a_Yx+b_Yy+c_Yxy=0. \tag{RC6}$$

These are source identities. With $p=ck/\omega$ (possibly unbounded), one has $pG=kB_c$ and hence

$$pB_c-kG=(p+k)d.$$

The entire residual normal coefficients after RC3 are exactly

$$\mathcal{R}'_E = 2\lambda(p+k)d + (a_E + c_E V)R_D + (b_E + c_E x)vF_v, \tag{RC7}$$

$$\mathcal{R}'_X = -2t\lambda ck\, d + (2t\lambda c + x_5 uv)R_D - 2t\lambda cuJ + u(x_3 + x_5 x)F_v, \tag{RC8}$$

$$\mathcal{R}_Y = (a_Y + c_Y V)R_D + (b_Y + c_Y x)vF_v. \tag{RC9}$$

The actual N_y coefficient is $\sqrt{h}\mathcal{R}_Y$. The identity underlying RC8 is particularly useful:

$$xu - ckG = cR_D - cuJ.$$

The identities $UV - xy = V(U-x) + x(V-y)$ and RC6 prove all three formulas. Thus no normal is inferred merely from its vanishing.

Absorb the unbounded ratio without comparing it. The elementary circuit identity and its exact coupled completion are

$$\begin{aligned} L_3 &= J^2 + (1 + 2k/p)d^2, \\ \lambda L_3 + \lambda p^2 E^2 + 2\lambda(p+k)Ed &= \lambda J^2 + \frac{2\lambda k}{p}(pE+d)^2 \\ &\quad + \lambda\{(p-k)E + d\}^2 - \lambda k^2 E^2. \end{aligned} \tag{RC10}$$

All radicals in its positive squares are separated: $k/p = \omega/c$. No boundedness assumption on p has been used. Put $R = \sqrt{\lambda}\{(p-k)E + d\}$. The singular cross term in RC4 must be kept with RC8. Together they are

$$-2t\lambda ck\,\xi(pE+d) = -2t\sqrt{\lambda}ck\,\xi R - 2t\lambda ck^2 E\xi. \tag{RC11}$$

Both displayed coefficients are infinitesimal. The remaining $\mu\xi^2$ term is bounded and infinitesimal. This exact cancellation is the step that permits arbitrary independent rates of c and ω.

The two coupled roots simplify after homogenization to

$$\begin{aligned} pE + dZ = kv(tcW - Z), \qquad R = \sqrt{\lambda}\, tk(cv - u)W, \\ \sqrt{2\lambda k/p}\,(pE + dZ) = \sqrt{2\lambda/c}\, k\sqrt{uv}\,(tcW - Z). \end{aligned} \tag{RC13}$$

In particular their final forms contain no testing-ratio inverse. Retain $Q^2, N^2, (2\lambda k/p)(pE+d)^2$ as exact positive squares. Every remaining term is a bounded Gram on

$$(E, R_D, N_y, \xi, F_v, J, R).$$

Its only nonzero off-diagonal residue is between E and R_D: $a_E \to -1/(t_0 - 1)$. Their full residue block is

$$\frac{1}{2(t_0-1)}\begin{pmatrix} 1 & -1 \\ -1 & t_0 \end{pmatrix} \succ 0, \qquad \det = \frac{1}{4(t_0-1)}. \tag{RC12}$$

The other diagonal residues are $t_0/2, 1/2, 1/(2t_0), 1/2, 1$. All other entries are infinitesimal: this is explicit in RC7–RC11, using $\sqrt{h}, c, u, v, k$ infinitesimal and all printed source arrays bounded. The subtraction $-\lambda k^2 E^2$ in RC10 does not change the positive residue. The bounded-Gram lemma therefore supplies finite tensor SOS for the entire parent.

Restoration and all-tests join. Homogenize all scalar roots by Z. The expressions RC3 are already state-linear, and every other root becomes a state-linear expression; no Z inverse remains. Hence $Z = 0$ is included. All divisions are pure source or pure testing. Although p and k/p may be unbounded, they occur only inside the printed exact positive square or the root R, never as an entry of the bounded Gram. No ratio hypothesis on u/v has been used. The exact original gauges restore both angle signs; real and imaginary parts give complex states. The final roots in RC3 and RC13 and all other Gram roots have no testing inverse, so substitution extends the identity to either literal axis directly. The inherited R13 R9 remote theorem covers the complement of both angles infinitesimal. These are exhaustive pure-testing cases; no testing-ratio split remains. Thus this coupled certificate closes the whole R9 source shape, with every original source gap and literal $n = 0$ retained. No other weak-C2 source shape is claimed here.

Chapter 132

All-tests closure of the weak-C2 source shape R10

Component `p526dc49d5fcb`
Source: `R16/proofs/r10/r10_all_tests_join.tex`
SHA256: `526dc49d5fcb8320e449c7a25479dd9b0e328d90fc6d993df59cc9c944e66488`
Inventory use: entire theorem

132.1 All-tests closure of the weak-C2 source shape R10

Theorem 132.1. *Retain the unchanged original physical weak-C2 source, $d = 1$, $\Lambda = \beta = 0$, $n \geq 0$, every strict source gap, the total-scale intersection and all inherited exclusions defining R10:*

$$t, g, j \text{ are positive bounded units}, \qquad B, h, e, x = aB \in \mathrm{Inf}, \qquad a \text{ unbounded}.$$

For every independent projective testing angle pair and every real or complex testing state, the complete original parent is finite tensor SOS. Thus the whole R10 source shape is admitted. No comparison of a source rate with a testing rate, and no mixed inverse, is used.

Exact source entry and remote tests. Use `r10_source_and_weighted_closure.tex`. Original physicality proves, with $s = 1/a$, $w = ae$ and $b = a^2 B$,

$$b/w \text{ bounded}, \qquad n = O(s^2 w), \qquad m/w \sim 1/[2(1-w)],$$

and all its opening-uniform source bounds. Its full-parent remote argument covers every testing pair outside both angles infinitesimal, including projective endpoints. It permits bounded m and retains the entire positive m row where needed.

The three exhaustive local source cases. If n/e^2 is noninfinitesimal, the strict weighted angular pivot R10.11–R10.12 and the complete identity R10.9 give the local certificate for every testing state. This is the first component's weighted local admission.

If $n = 0$, the exact physical OP4 identification has native C1 $\eta_1 = t$ a unit, $\zeta_1 = s$, $r_1 = h$ infinitesimal, and $e^2_{1,\mathrm{scale}} = tb$ bounded. Its infinitesimal and positive-unit types are covered respectively by the inherited small-e and centered-A theorems, retaining literal $\beta_1 = 0$. This is the stated C1.2/C1.3 routing, not a continuation through negative formal beta.

It remains that $0 < n/e^2 \in \mathrm{Inf}$. On both-small-angle tests, `../root/r10_unit_W_join.tex` applies directly to the actual C2 coefficients. The exact pure-source critical correction obeys $(C_* - C)/m = sR$ with R bounded; SEO11–SEO16 consequently prove every normalized unit-W state, including $Z = 0$. On a negative pure-testing-core branch, normalized X, Y are infinitesimal.

Any state not already covered therefore has Z a unit and W infinitesimal. Normalize $Z = 1$. The inherited testing Schur identity and sign gauges then give exactly

$$u, v, \mathfrak{w} = (W + uY)/(1 + u^2) > 0 \quad \text{all infinitesimal.}$$

This is the contact chart proved in `r10_thin_layer_contact.tex`. Its comparison with formal C1 is implemented by exact full-parent coefficient identities, a new positive contact-zero n row, and the restricted CA/SE finite contact conductors. It does not invoke a physical C1 theorem at negative beta. The strict eight-root base plus its infinitesimal bounded stress proves the actual C2 parent on this remaining chart.

Restoration and scope. The nonnegative-core branch and literal testing axes retain their original event-square exits. The projective remote identities and unit-W join cover the required endpoints; the local conductor uses only pure-testing and pure-source inverses. Real and imaginary state evaluations give the complex assertion. These cases exhaust all original R10 sources and independent tests, so no local R10 obligation remains.

The focused independent audits are `../joint_frame/r10_unit_W_audit.tex` and `../joint_frame/r10_thin_layer_audit.tex`. The three components establish R10 only. They do not admit another weak-C2 shape, a remaining C1 branch, or the global theorem.

Chapter 133

R10: forced source scale, every remote test, and a weighted local admission

Component `p9ee4ba7326a7`
Source: `R16/proofs/r10/r10_source_and_weighted_closure.tex`
SHA256: `9ee4ba7326a7a5d8b45598ae06d8c427604c26be4ba94836adccf0fa638d82cf`
Inventory use: R10.1-R10.12 and complete remote join

133.1 R10: forced source scale, every remote test, and a weighted local admission

Scope and result. Retain the original weak-C2 R10 physical stationary source, every strict source gap, $n \geq 0$, the positive-infinitesimal total-scale intersection, and every inherited exclusion. Thus t, g, j are positive bounded units, B, h, e are infinitesimal, a is unbounded, and $x = aB$ is infinitesimal. The new all-tests admission proved here is the source subdomain

$$n/e^2 \notin \mathrm{Inf}\,. \tag{R10.1}$$

Every nonlocal independent testing chart is also proved on all of R10. The positive-n, infinitesimal-n/e^2 local complement is handled by the following thin-layer and state-entry sections; it is not covered by this weighted pivot. The literal $n = 0$ boundary is already an inherited C1 admission, by the exact map specified below. These statements retain all independent rates.

The decisive exact source scale. Use the pure-source variables

$$s = 1/a, \qquad w = ae, \qquad b = a^2B = x/s, \quad B = s^2b, \quad x = sb, \quad e = sw, \quad r = 1 - sw.$$

Initially b is not assumed bounded. The gap $g = 1 - w + hx$ makes w bounded and $1 - w$ a positive unit in residue. In the exact positive polynomials of BS2 put

$$Q_* = 1 + B(1 + t + ah + 2ar + at) + B^2\{a(h + t) + a^2[h(r + 1) + t(1 + r - r^2)]\},$$

$$K_* = (1 + s)(h + t)(1 + x), \qquad \Delta_* = s\Delta.$$

Before imposing a bound on b, $Q_* \sim 1$ and $\Delta_* \sim 1$: write them using bounded w, infinitesimal s, x, h; their only nonconstant monomials contain s or x. BS1 gives exactly

$$\boxed{2r\Delta_*\, n/s^2 = wQ_* - bK_*.} \tag{R10.2}$$

Consequently

$$0 < b/w \leq Q_*/K_*, \qquad 0 \leq n/s^2 \leq wQ_*/(2r\Delta_*). \tag{R10.3}$$

In particular $b = O(w)$ is bounded, $n = O(s^2 w)$, an is infinitesimal, and $ck = \sqrt{b/t}$ is bounded for $c = \sqrt{a/t}$, $k = \sqrt{x}$. This is an original-source bound, including equality $n = 0$, not a comparison with either testing angle.

After this bound every displayed source denominator is a unit in (s, b, w, h, t) except the explicit positive monomials already canceled. Direct finite rational cancellation in the source formulas gives

$$\begin{aligned} m/w &\sim \frac{1}{2(1-w)}, \quad C \sim Ah \sim t/2, \quad D \sim \frac{1}{2(t-1)}, \quad \alpha/(s^3 b^2) \sim t/2, \\ \lambda &= \frac{m - a^2 q}{t a^2 B} \sim \frac{1}{2}, \qquad c_F \sim \frac{t - w + tb}{2t^2} > 0, \qquad q = nr^2 + (m + \tfrac{1}{2})e^2 + \alpha. \end{aligned} \tag{R10.4}$$

The last strictness follows from $t_0 > 1$ and $w_0 < 1$. These are joint residue identities in bounded letters; they do not set w or b to a positive unit. In particular they hold when either is infinitesimal. The cancellation defining λ leaves no b or s denominator; the new checker verifies this explicitly. We also have m bounded, $n, \alpha, A^{-1/2}$ infinitesimal, and

$$d_4 \sim \frac{t}{2(t-1)}, \quad b_4 \sim 0, \quad a_E \sim -\frac{1}{t-1}, \quad x_3 \sim -1. \tag{R10.5}$$

All first-normal arrays of the R15 source appendix are bounded: the only newly unbounded letter a occurs there through an or aq, both infinitesimal. The factors $a^2 n$ in c_F are bounded.

All remote tests on all of R10. The exact full-parent identities GR2–GR7 and WE8/WE10 apply. Their R9 source use required m infinitesimal, but their algebra does not. Under R10.4 the entire nonnegative m packet is retained. For GR2 it adds a positive rank-one form to the same strict six-root residue GR3; all its coefficients are bounded. For the chart u infinitesimal, v a bounded unit, keep $S = \sqrt{m}W$. The complete (S, X) residue is now

$$(v_0 S - \sqrt{m_0} X)^2 + \frac{1 + v_0^2}{2} X^2, \qquad \det = \frac{v_0^2(1 + v_0^2)}{2} > 0. \tag{R10.6}$$

The (uW, Z) block remains GR5 and the normalized Y pivot remains strict; the extra uvY, uZ terms in the m packet are infinitesimal. For unbounded v its reciprocal identity has the same diagonal residue GR6 because the potential $\sqrt{m}X$ term carries $1/v$. On the remaining remote charts retain mR_m^2 as a full positive square and apply the unchanged strict residue to the remainder. The original homogeneous identities include either literal infinity, and the GR2/WE8 axes retain $n = 0$ without a division. The original sign gauges and real/imaginary split are unchanged. Hence every test outside both angles infinitesimal is covered for the entire R10 source.

The exact local full-parent identity. Use the R15 generic contact balances RC1, RC6 and exact normal identities RC7–RC9. These are rational identities of the original source and never require bounded a. Set $U = u^2, V = v^2$, $y = tB$,

$$\begin{gathered} E = tuW - vZ, \quad N_y = (Y - huW)/\sqrt{h}, \quad \xi = X + ruZ, \\ F_v = (V - y)Z, \quad R_D = v(U - x)Z, \quad J = (uv - x/c)Z, \\ Q = \sqrt{q}(avW - uZ - a\xi), \qquad N = \sqrt{n}\{r[(a/t)V - U]Z + u\xi\}, \\ R = \sqrt{\lambda}\, tk(cv - u)W, \qquad T = \sqrt{2\lambda/c}\, k\sqrt{uv}(tcW - Z), \quad \mu = m - a^2 q = ta\lambda x. \end{gathered}$$

At positive nonzero angles these are exactly the R15 coupled roots. They have no testing-ratio inverse. Let $\mathcal{H}_0$ denote the hard quadratic after subtracting $Q^2 + N^2$ and the exact rank packet $\mu\{vE/(tu) - \xi\}^2$ from the original parent. Independent full-parent differentiation in the new checker verifies that every entry of $\mathcal{H}_0$ has no a dependence and no testing inverse. Its R10 local residue is

$$\operatorname{diag}\left(\frac{1}{2(t_0 - 1)}, \frac{t_0}{2}, \frac{1}{2}\right) \quad \text{on } (E, N_y, \xi). \tag{R10.7}$$

This permits bounded m; every occurrence of m in the remaining hard entries carries a positive testing degree or h.

The principal angular–ξ coupling is resolved as a whole. Writing $R_c = kv(tcW - Z)$, direct multiplication gives

$$-2t\lambda ckR_c\xi + 2t\lambda c(R_D - uJ)\xi = -2t\lambda xQ\xi/\sqrt{q} - 2\mu\xi^2. \tag{R10.8}$$

Thus neither cu nor cR_D is left as a Gram coefficient. Combining RC2, RC7–RC10, the full rank packet and R10.8 gives the entire original $F/4$ exactly as

$$\begin{aligned} &Q^2 + N^2 + T^2 + R^2 + \lambda J^2 + d_4R_D^2 + (c_F + b_4x/t^2)F_v^2 + (b_4/t^2)vR_DF_v \\ &\quad + \mathcal{H}_0 - \lambda xE^2 - \mu\xi^2 - 2(t\lambda x/\sqrt{q})Q\xi \\ &\quad + E\{(a_E + c_EV)R_D + (b_E + c_Ex)vF_v\} \\ &\quad + \sqrt{h}N_y\{(a_Y + c_YV)R_D + (b_Y + c_Yx)vF_v\} \\ &\quad + \xi\{x_5uvR_D + u(x_3 + x_5x)F_v\}. \end{aligned} \tag{R10.9}$$

All coefficients refer to the exact source arrays in the R15 appendix. This identity retains every scalar, first-normal and hard-normal term.

A strict weighted pivot when n/e^2 is noninfinitesimal. Put $\rho = n/e^2$, $H = m + 1/2$, $\chi = m/w$, and $\overline{q} = a^2q$. Then

$$\overline{q} = w^2(\rho r^2 + H) + a^2\alpha, \qquad \chi - H \in \mathrm{Inf}\,. \tag{R10.10}$$

The Q, ξ source block in R10.9 has Schur pivot

$$S = \frac{1}{2} - \mu - \frac{(t\lambda x)^2}{q} = H - \frac{m^2}{\overline{q}}. \tag{R10.11}$$

The equality uses $a^2q + \mu = m$ exactly. Since $\alpha \geq 0$,

$$S \geq H - \frac{\chi^2}{\rho r^2 + H} = \frac{H\rho r^2}{\rho r^2 + H} + \frac{H^2 - \chi^2}{\rho r^2 + H}. \tag{R10.12}$$

Here H is a positive bounded unit. Under R10.1 the first summand has positive residue and the second is infinitesimal, also when ρ is unbounded. Hence S is a positive bounded unit. Moreover b/w is bounded and $\overline{q} \geq Hw^2$, so $x/\sqrt{q} = b/\sqrt{\overline{q}}$ is bounded. All entries of this strict source block are therefore bounded.

In the rest of R10.9 the only nonzero off-diagonal residue is a_E between E and R_D, giving

$$\frac{1}{2(t_0 - 1)}\begin{pmatrix} 1 & -1 \\ -1 & t_0 \end{pmatrix} \succ 0.$$

The N_y, F_v, J, R pivots have positive residues $t_0/2, (t_0 - w_0 + t_0b_0)/(2t_0^2), 1/2, 1$. Every other entry is infinitesimal by R10.4–R10.5 and the explicit positive testing degrees in R10.9. Retain N^2, T^2 as exact positive squares. The bounded-Gram lemma supplies finite tensor SOS for the entire parent on R10.1. No source/testing rate comparison occurs. All roots homogenize without Z or testing-ratio inverses, so literal axes and $Z = 0$ are included directly. The original signs, complex-state split and the remote theorem complete the all-tests admission.

Literal $n = 0$ and the exact remaining complement. At $n = 0$, OP4 is a physical C1 identification with

$$x_1 = x, \quad \eta_1 = t, \quad \zeta_1 = s, \quad r_1 = h, \quad \theta_1 = e, \quad e_{1,\mathrm{scale}} = \sqrt{tb}.$$

Thus k_1, ζ_1, r_1 are infinitesimal, η_1 a unit, and $e_{1,\mathrm{scale}}$ bounded. Its infinitesimal type is covered by `baseline/baseline/baseline/advances/c1_scales/small_e_all_tests.tex`; its positive-unit type is covered by the neighboring `centered_a_bounded_delta.tex`. Both retain $\beta = 0$. This is C1.2/C1.3 routing, not the unbounded-η C1.1 theorem. For $n > 0$ OP5 gives formal $\beta_1 < 0$, so this physical C1 theorem cannot be extended by continuity or imported into the neighborhood. The complement left for the companion thin-layer proof is exactly

$$0 < n/e^2 \in \mathrm{Inf}, \qquad u, v \in \mathrm{Inf},$$

with every original physical and total-scale intersection retained. In this complement R10.11 can have zero residue: at a unit w and $b_0 = w_0/t_0$, its residue is zero. Thus R10.9 is a useful full-parent identity there, but its stated strict-Gram argument does not prove that complement. The companion thin-layer proof and direct state join complete this complement; the final join states the whole-R10 theorem.

Chapter 134

The positive-n thin R10 layer: an exact extra row and a bounded conductor perturbation

Component `pef37475d831e`
Source: `R16/proofs/r10/r10_thin_layer_contact.tex`
SHA256: `ef37475d831e144531c59d8d80c245a3f315a6c02fdf0db156d80f66e82e481c`
Inventory use: TL1-TL10

134.1 The positive-n thin R10 layer: an exact extra row and a bounded conductor perturbation

Domain. All polynomial identities below use the convention $F/4$, as in R10.9; in particular the original n packet is $nD_uD_vX^2$. All parent and packet identities in this section use the common normalization $F/4$. Keep every original R10 condition and assume $\rho = n/e^2 > 0$ infinitesimal. Use s, w, b, x, B from R10.2. Dividing that exact identity by w gives

$$Q_* - (b/w)K_* = 2r\Delta_*\rho w, \qquad b/w \sim 1/t. \tag{TL1}$$

Thus $\psi = w/b$ is a positive bounded unit. No testing rate occurs. The formal native C1 map OP4 is

$$x_1 = x, \quad \eta_1 = t, \quad \zeta_1 = s, \quad r_1 = h, \quad \theta_1 = e, \quad t_1 = \theta_1/\zeta_1 = w.$$

Write $P_d = P_n/B$. OP3–OP5 give the exact slack

$$s_n = 2r\Delta_*\rho w/P_d, \quad \beta_1 = -s_n/2, \quad \pi = 1 - s_n = 1 + 2\beta_1 > 0, \quad \kappa = (1+2n)/\pi. \tag{TL2}$$

Hence $s_n/w = O(\rho)$, β_1 is infinitesimal, and $\kappa \sim 1$. The formal C1 parent is used through its explicit identities, not imported as a physical C1 source theorem.

Exact coefficient comparison. Let F_1 be the native C1 parent at this same contact, normalized by $d_1 = 1$, and F_2 the actual C2 parent, with $d_2 = 1$. For a coefficient L put $\delta L = L_2 - \kappa L_1$. Set $K = 1 + x$, $c_a = 1 + xr$, $S_c = 1 + B(1-h) + xr$, $Q_c = 1 - hB$, and $\Theta = n/B = \rho w\psi$. The following short polynomials are bounded with the displayed unit denominators:

$$P_\kappa = c_a^2 + sx(1+t) + sx^2(r + he + t),$$

$$P_D = tx^2r^2 + x\{t - 1 + e(1 - h - 2t)\} - 1, \quad P_\alpha = 1 + s - 2e + x\{r^2 + s(r + he)\}.$$

Exact cancellation of the original C2 and native C1 formulas gives

$$\begin{array}{c|l} \delta m & 0 \\ \delta C & -\Theta c_a/(1-h) \\ \delta(Ah) & -\Theta t c_a S_c/[K(h+t)] \\ \delta D & \Theta S_c P_D/[K(h+t)tj] \\ \delta\alpha & -nxP_\alpha/(KQ_c) \\ \delta d = 1-\kappa & -2\Theta P_\kappa/[K(h+t)] \\ \delta\beta + \delta d/2 & -n. \end{array} \tag{TL3}$$

The new checker verifies all seven identities and TL2 exactly. In particular $\delta C, \delta D, \delta d, \delta(Ah) = O(\rho w)$, while $\delta\alpha = O(nx)$. There is no surviving $1/h$ after the large A event is centered.

The formal C1 local identities still supply a strict base. The native η_1 is a unit, k_1, ζ_1, r_1 are infinitesimal, and $e^2_{1,\mathrm{scale}} = tb$ is bounded. Separate its positive-unit and infinitesimal types. Use respectively the exact centered-A CA5/CA8 construction (or its CS1 scaling) and the SE4/SE9–SE11 construction. This use concerns the printed formulas and conductor lemmas; the physical-source theorem is not being invoked at $\beta_1 < 0$. Here are all changes to their source premises:

The formulas and native stationary identities are rational identities, valid at the formal OP4 contact. All native gaps remain positive; $M = g$ and $J = j$ are units. The other native event coefficients are positive, with $C_1, D_1, A_1 h$ positive units and A_1 unbounded. The opening ratio $t_1/e^2_{1,\mathrm{scale}} = w/(tb)$ is a positive bounded unit by TL1, which supplies the only scale bound previously obtained from $\beta_1 \geq 0$. All four retained packets SD3 involve the positive paired coefficient $\beta_1 + 1/2 = \pi/2$, never a standalone β_1 square. In SE4, the coefficient of the reserved normal packet is $(2+4\beta_1)\nu$, so its subtraction of ν is still positive, since β_1 is infinitesimal. The remaining source bounds, finite coquotients and strict residue matrices are unchanged. A positive-unit normalization to $C_1 = 2$, when used by those formulas, changes none of these bounds. Consequently their exact local construction yields the same strict bounded eight-root Gram and positive retained packets for κF_1 on its contact chart.

A new positive contact-zero n row. Use the inherited pure-testing coordinate and normals

$$\mathfrak{w} = (W + uY)/D_u, \quad A_t = v\mathfrak{w}, \quad X = N + pD_u A_t - uZ, \quad p = t_1/R_1,$$

with $D_u = 1 + u^2, D_v = 1 + v^2$. In the exact CA chart the contact is

$$(u, \mathfrak{w}, v) = (k, \delta k, \delta k^2 E), \quad K_c = \delta^2 E k^2 = \zeta_1 K/Q.$$

Here Q, E denote the inherited CA source letters, not the angular weight of R10. The SE chart has $k = \varepsilon j$, $\delta = 1/\varepsilon$, so $K_c = j^2 E$. Put $\gamma = r/K_c$ and retain the entire state-linear square

$$\boxed{\mathcal{N}^2 = nD_u D_v (X + \gamma A_t)^2.} \tag{TL4}$$

At the actual contact $A_t = K_c k$ and $X = -rk$, so its value and all first derivatives vanish exactly. The square root of $D_u D_v$ is pure testing; every other radical or inverse is pure source or pure testing. The following entire-parent identity includes its full normal:

$$\begin{aligned} nD_u D_v X^2 - \mathcal{N}^2 = {} & 2n\gamma D_u D_v u A_t Z - (2n\gamma p D_u + n\gamma^2) D_u D_v A_t^2 \\ & - 2n\gamma D_u D_v A_t N. \end{aligned} \tag{TL5}$$

There is no remaining hard quadratic from this subtraction. This row is necessary: without it the scalar nu^2 lies outside the old raw square ideal, even if n/k^4 is infinitesimal.

Exact bounded stress after the positive packets. Let $A_{*,1}$ be the inherited bounded centered-A cap. In the actual parent retain $(A_2 - \kappa A_{*,1}) D_u D_v (Y - huW)^2$, which is positive because A_2 is unbounded and $\kappa A_{*,1}$ bounded. Comparing with the corresponding retained packet of κF_1 replaces the whole A difference by

$$\delta(Ah) D_u D_v (2uWY - hu^2 W^2).$$

After retaining TL4, the full difference of the bounded remainders is exactly

$$\begin{aligned} \mathscr{E} = {} & [nD_u D_v X^2 - \mathcal{N}^2] - nD_v (X + uZ)^2 + \delta C D_v (uW - Y)^2 \\ & + (\delta D + \delta d/2) R_{D,\mathrm{event}}^2 + \delta\alpha D_u (vY - Z)^2 \\ & + (\delta d/2) D_u \{(vY - Z)^2 - D_v Z^2\} \\ & + \delta(Ah) D_u D_v (2uWY - hu^2 W^2), \end{aligned} \tag{TL6}$$

where $R_{D,\mathrm{event}} = uW + uvX - Y - vZ$ is the original event row. This follows from TL3; in particular the paired beta difference is exactly $-n$. Every scalar, first-normal and hard-normal term is present. The full expression and its first derivatives vanish at the unchanged source contact, because both full parents and every retained square do.

Finite support and bounds for the unchanged conductor. Work on the local chart $Z = 1$, with $u, v, \mathfrak{w}$ positive infinitesimal. At zero normals set

$$V_0 = 1 - puv\mathfrak{w}, \quad q_B = D_u v V_0, \quad uW = u\mathfrak{w} + u^2 h_1 v V_0, \quad Y = u\mathfrak{w} - h_1 v V_0, \quad h_1 = a_1/T_1. \tag{TL7}$$

These are exact inherited state inverses. The rows $uW - Y$ and $R_{D,\mathrm{event}}$ are in (v) at zero normals; $X + u = pD_u A_t$. The centered A expression in TL6 is a quadratic in $(v, u\mathfrak{w})$ with bounded coefficients. The signed-core line has no constant: $(vY - 1)^2 - D_v$ is explicitly $(vY)^2 - 2vY - v^2$. The sole state-constant term is $\delta\alpha D_u$. Finally TL5 is a combination of $uA_t = (v)(u\mathfrak{w})$ and A_t^2, with first normal in (A_t). Thus this is an explicit raw allocation, not an ideal-membership guess.

For the unit-$e_{1,\mathrm{scale}}$ chart use

$$(A, B_c, C_c, D_0) = (\mathfrak{w} v, \delta v, \delta E u \mathfrak{w}, \delta^2 k^3 E).$$

Every scalar coefficient of $\mathscr{E}/\rho$ in $(A, B_c, C_c)^2 + (D_0^2)$ is bounded with unit denominators. Both complete first normals belong to (A, B_c, C_c) with bounded coefficients, and the hard-normal entries are bounded. The decisive source coquotients are

$$\Theta/\rho = \delta^2 Q E w^2/K, \quad n\gamma^2/\rho = w^2 r^2 Q^2/K^2, \quad nx/(\rho D_0^2) = Q^2 w^2/K^2. \tag{TL8}$$

The remaining $n\gamma$ coquotient has an additional $\delta^2 k^2 E$. All apparent negative k powers have therefore canceled before the conductor is used.

For the infinitesimal-$e_{1,\mathrm{scale}}$ chart use the exact SE parameters $w = t_1 = \varepsilon^2\tau$, $\zeta_1 = j^2 QE/K$, $k = \varepsilon j$, and

$$(A, B_c, C_c, D_0) = (\varepsilon \mathfrak{w} v, v, E u \mathfrak{w}, \varepsilon^2 j^3 E).$$

The full raw scalar allocation of $\mathscr{E}/\rho$ is

$$f_A A^2 + f_{BB} B_c^2 + f_{BC} B_c C_c + f_{CC} C_c^2 + f_0 D_0^2. \tag{TL9}$$

All five coefficients are bounded; after division by ρ each has a factor ε^2. The A^2 coefficient has this factor by TL10; all its other terms carry $p = \varepsilon^2\tau/R_1$, because $\delta m = 0$. The coefficients of $B_c^2, B_c C_c, C_c^2$ have it from Θ/ρ, and the constant coefficient from the final identity in TL10. In particular

$$\begin{gathered} \Theta/\rho = \varepsilon^2 \tau^2 QE/K, \\ \frac{n\gamma^2}{\rho\varepsilon^2} = \frac{\varepsilon^2\tau^2 r^2 Q^2}{K^2}, \quad \frac{n\gamma}{\rho\varepsilon} = \frac{\varepsilon^3 j^2 E \tau^2 r Q^2}{K^2}, \quad \frac{nx}{\rho D_0^2} = \frac{\varepsilon^2\tau^2 Q^2}{K^2}. \end{gathered} \tag{TL10}$$

The first-normal coefficients are in (A, B_c, C_c) with bounded arrays, also divisible by ε^2 after division by ρ; the new A coefficient has the stronger $\varepsilon^3 j^2$ factor printed in TL10. The extra term $-nD_v(X+u)^2$ has scalar coefficient $-np^2D_u^2D_v$ on A_t^2, first normal $-2npD_uD_vA_tN$, and hard term $-nD_vN^2$; TL10 and $p = \varepsilon^2\tau/R_1$ bound each normalized coefficient.

Exactly as in SE8, all dependence of the raw coefficients on $\mathfrak{w}$ comes through V_0 and has a factor $p = O(\varepsilon^2)$. The additional TL5 coefficients are independent of $\mathfrak{w}$. Modulo ε, scalar raw coefficients are even in u, v; the normal raw coefficients are independent of $\mathfrak{w}$. Clearing powers of D_u and the source unit denominators preserves these properties. These are precisely the finite divisibility premises of SE7: the first scalar conductor has $T_{\mathfrak{w}} \in (\varepsilon)$, the second-stage remainders have $T_{\mathfrak{w}} \in (\varepsilon, v)$, and each full normal has $T_{\mathfrak{w}} \in (\varepsilon)$.

For either chart, TL6 has exact contact value and first-jet zero. Apply the inherited monic contact lift to $\mathscr{E}/\rho$: CC3–CC6 in the first chart and the division-free SE7 lift in the second. The raw support restriction retains only D_0^2, not arbitrary mixed D_0 products. The proofs use only finite monic division and bounded unit denominators; extending their coefficient ring by the bounded source letters in TL8–TL10 changes nothing. They give a bounded Gram matrix on the same eight inherited roots for $\mathscr{E}/\rho$. Hence the exact stress $\mathscr{E}$ has an infinitesimal Gram matrix on that list. No inverse of ρ occurs in its final coefficients: division by ρ here records the common positive source factor before a linear finite lift.

Adding this infinitesimal matrix to the strict formal-C1 base Gram preserves its positive residue. Restore all its retained packets, the modified positive centered-A packet, and TL4. This proves the full actual C2 parent on the thin-layer contact chart, with both complete normals, all hard couplings and every independent rate. The direct actual-C2 outer/state argument and the all-R10 remote argument are required for the all-tests join; no negative formal-beta packet is used in either join.

New exact receipts. `check_thin_stress.py` verifies fourteen exact source, row and scaling identities. `check_thin_full_support.py` verifies five further statements: reconstruction of the entire original-parent stress, complete scalar restricted raw support, each full first-normal support, and absence of higher normal degree. These are new affected checks, not reruns of the inherited C1 suites. The finite conductor application is the analytic argument above.

Chapter 135

R10: a direct critical-wall certificate for every unit-W state

Component `pf1aac1dff7c7`
Source: `R16/proofs/root/r10_unit_W_join.tex`
SHA256: `f1aac1dff7c7e7d6e81668ec7d431739b00dd7021043fb193fc39fc82a794233`
Inventory use: entire exact source coquotient and whole-state argument

135.1 R10: a direct critical-wall certificate for every unit-W state

Statement. On the entire original R10 source, every independent test with u, v infinitesimal and normalized W a testing unit is tensor SOS. This includes bounded Z of every sign, infinitesimal Z, and $Z = 0$. The theorem uses the actual physical C2 coefficients. No continuation of a negative-beta C1 source is involved.

Use the R10 source scale $s = 1/a$, $w = ae$, $b = a^2B$, and put $z = b/w$. The source constraint proves z bounded. Write $A_h = Ah$. The source estimates give A_h, C, D positive units, h infinitesimal, m/w a positive unit, and $t_0 > 1$, $w_0 < 1$. Define the positive pure-source critical coefficient

$$C_* = \frac{(h + 2A_h)(1 + 2D)}{2(4A_hD - h)} = \frac{(1 + 2A)(1 + 2D)}{2(4AD - 1)}. \tag{UW1}$$

Its denominator after the first normalization is a positive unit. We prove the opening-uniform estimate

$$\frac{C_* - C}{m} = s\,R, \qquad R \text{ bounded.} \tag{UW2}$$

A short polynomial proof of the source estimate. All the following letters are polynomials in bounded source variables (s, w, z, h, t) after $b = wz$, with $x = sb$, $B = s^2b$, $r = 1 - sw$:

$$c_a = 1 + xr, \quad S_c = 1 + B(1 - h) + sbr, \quad g = 1 - w + hx, \quad J = tc_a - (1 - h), \quad \Delta_* = s\Delta,$$

$$M_* = (1 + x)(s + r)(h + t) - sw(1 + tB)(1 - h).$$

Use the exact source numerators and denominators

$$\begin{aligned} N_C &= c_aM_*, & D_C &= 2(1 - h)\Delta_*, \\ N_A &= tc_a(s + r)S_c, & D_A &= 2\Delta_*, \\ N_D &= (1 + x)(s + r)(1 - txr)S_c, & D_D &= 2J\Delta_*, \\ N_m &= w(1 + tB)c_aS_c, & D_m &= 2g\Delta_*. \end{aligned}$$

Thus $C = N_C/D_C$, $A_h = N_A/D_A$, $D = N_D/D_D$, and $m = N_m/D_m$. Clear the denominator in UW1:

$$\begin{aligned} P &= (hD_A + 2N_A)(D_D + 2N_D)D_C \\ &\quad - 2N_C(4N_AN_D - hD_AD_D), \\ Q &= 2(4N_AN_D - hD_AD_D)D_C. \end{aligned} \tag{UW3}$$

Then $C_* - C = P/Q$. At $s = 0$ one has

$$C = \frac{t+h}{2(1-h)}, \quad A_h = t/2, \quad D = \frac{1}{2(t-1+h)},$$

and substitution gives $C_* = C$ identically. At $w = 0$ (hence $b = 0$), the same three formulas hold for every s, because all factors $1 + s$ cancel. Therefore the polynomial P is divisible by both s and w, hence $P = swP_1$. Since $N_m = wN_{m,1}$,

$$\frac{C_* - C}{m} = s\,\frac{P_1 D_m}{QN_{m,1}}.$$

The denominator $QN_{m,1}$ at $s = h = 0$ is $16t > 0$. This proves UW2 by finite polynomial cancellation, without comparing w to s or h. The new checker verifies exactly this divisibility and denominator. Its displayed residue also gives $R \sim t(1-w)(tz+2)$, though the proof only needs boundedness.

Complete parent and state allocation. Set $\epsilon = (C_* - C)_+$. By UW2, ϵ/m is infinitesimal. If $C > C_*$, retain the nonnegative excess $C - C_*$. Retain the entire original n and α packets; beta is zero. The remaining signed core and C_*, A, D packets are exactly the critical parent SEO11, with $c = C_*$ and $d = 1$, because

$$2C_*(4AD - 1) = (1 + 2A)(1 + 2D).$$

The exact critical-wall identities SEO12–SEO15 give a bounded strict Gram on

$$B_t = X + uZ, \quad H_A = \sqrt{A}(Y - yV), \quad H_D = \sqrt{4D+2}\{uW - Y + uvB_t - (b_* + u^2)V\}, \quad uV,\ vV,$$

where $V = vZ$, $y = 1/(2A)$ and $b_* = (4AD-1)/(A(4D+2))$. Here y, b_* are bounded; C_*, D are units and A may be unbounded. The hard residue is strict and the two radial masses have positive residues 2 and $2/t_0$. This is an application of those full polynomial identities, not of their original C1 source theorem.

Normalize $W = 1$ by a pure-testing factor. The entire m event obeys

$$R_m = v(1 + uY) - B_t, \qquad V = \frac{Z}{1+uY}(R_m + B_t). \tag{UW4}$$

The denominator is a positive testing unit. From the displayed H_D root,

$$uW - Y = (b_* + u^2)V - uvB_t + H_D/\sqrt{4D+2}.$$

Thus $\sqrt{\epsilon}(uW - Y)$ is a linear combination of the strict critical roots and $\sqrt{m}R_m$ with infinitesimal coefficients. The only potentially small denominator is accounted for by $\sqrt{\epsilon/m}$, infinitesimal by UW2. Consequently the complete negative correction $-4\epsilon(1+v^2)(uW-Y)^2$ has an infinitesimal bounded Gram. The m packet supplies a strict additional pivot. Restore all retained original packets to obtain the entire C2 parent, including every hard state direction.

State entry for the remaining local work. Normalize a real testing state by a maximal coordinate. On the negative pure-testing-core branch at u, v infinitesimal, X, Y are infinitesimal. If W is a unit the theorem applies. Otherwise Z is a unit; set $Z = 1$, and then $w_{\text{test}} = (W + uY)/(1+u^2)$ is infinitesimal. The inherited exact Schur sign identity gives the positive contact chart. Literal axes have the original event-square exits; complex states split into real and imaginary parts. Together with the R10 remote theorem, the only remaining task is the contact chart $u, v, w_{\text{test}} > 0$ infinitesimal, $Z = 1$, on any source not already admitted locally.

Chapter 136

C14.3b: the complete gamma–circuit join and its negative soft source pivot

Component `pd6aedcfef8b0`
Source: `R17/proofs/c1_small_eta/coupled_gamma_soft_obstruction.tex`
SHA256: `d6aedcfef8b038e63739f4fe9c07831f1f1b8e419259cf37a2ab4410611f73cc`
Inventory use: GS1 exact polarization only | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

136.1 C14.3b: the complete gamma–circuit join and its negative soft source pivot

Status and unchanged domain. This is a diagnostic of the fully conducted frame. The damped DG/DS certificates elsewhere in this revision bypass it on their admitted domains; it creates no additional obligation beyond the current frontier. This note does not admit C14.3b. It gives an exact simultaneous join of its remaining full normal with both coupled-circuit rows, and identifies an additional obstruction after the former order-one angular obstruction has been removed. Retain $x, \eta > 0$ infinitesimal, $a = 1/\zeta$ a positive bounded unit, both original gap openings, $\beta \geq 0$, independent infinitesimal testing angles u, v, and the normalized W-unit state chart. No source/testing rate comparison or mixed inverse is used. The completed R16 bounded-$W/Z, X/Z$ chart and literal $Z = 0$ endpoint are unchanged.

Use the homogeneous JF1 notation, and abbreviate $Q = Q_t$, $T = T_D$, $G = \gamma h$, $x = k^2$, $a = \eta c^2$. Put

$$H = avW - uZ, \quad \tau = \eta cW - Z, \quad D_0 = u - cv, \quad J = (uv - x/c)Z,$$
$$E = \eta uW - vZ = \eta(P + Q), \quad R_c = kv\tau, \quad R = -\sqrt{\lambda}\,\eta k D_0 W, \quad \mathcal{T}^2 = (2\lambda x/c)uv\tau^2.$$

The inherited gauges allow $u, v \geq 0$; the identities below themselves are polynomial and do not require a testing ratio inverse. In particular no inverse of Z is used.

Join the entire remaining normal before extracting the circuit. The exact conductor and polarization are

$$uv\xi = h\{uJ + kR_c - (x/c)H - T\}, \qquad EH = uv\tau^2 - \eta D_0^2 WZ. \tag{GS1}$$

Define the source-only shift

$$d = \frac{G}{2\lambda}, \quad \widehat{R} = R + \frac{Gk}{2\sqrt{\lambda}}Q, \quad b_\gamma = b' - G/2,$$

and the complete modified source packet

$$\begin{aligned} S_\gamma = {} & CP^2 + AQ^2 + D_p(P - hT)^2 + 2b_\gamma QT \\ & - \lambda x(E - dQ)^2. \end{aligned} \tag{GS2}$$

Then the entire affected block has the exact identity

$$\begin{aligned} &CP^2 + AQ^2 + D_p(P - hT)^2 + 2b'QT + R^2 + \mathcal{T}^2 - \lambda x E^2 + \gamma uvQ\xi \\ &= S_\gamma + \widehat{R}^2 + GuQJ + (Gx/c)PH \\ &\quad + \frac{x}{c}(2\lambda - G/\eta)uv\tau^2 + (Gx/c)D_0^2 WZ. \end{aligned} \tag{GS3}$$

This joins the full R_c coupling, its R companion, the negative E correction and the extra circuit row at once. The completion

$$R^2 + (Gk/\sqrt{\lambda})QR + GxQE - \lambda x E^2 = \widehat{R}^2 - \lambda x(E - dQ)^2$$

is exact. The whole angular EH term has canceled in GS3. Thus the order-one negative angular residue JR7 is not being reused as the obstruction in this note. All the remaining JF1 terms are left unchanged, including λJ^2, the complete radial rows, the angular block, the full D_p shift and every higher-angle square.

The newly exposed source block is strictly indefinite. First omit just the negative last square in GS2, and call the result S_0. Its pure-source P completion is

$$S_0 = (C + D_p)\left(P - \frac{hD_p}{C + D_p}T\right)^2 + AQ^2 + h^2KT^2 + 2b_\gamma QT, \qquad K = \frac{CD_p}{C + D_p}. \tag{GS4}$$

On $(\sqrt{\eta}Q, T/\sqrt{\eta})$ the remaining determinant is

$$\Delta = h^2AK - b_\gamma^2. \tag{GS5}$$

There is a particularly simple exact native-source identity:

$$b_\gamma = -\frac{hC_n}{r}, \qquad C_n = (1 - r)C = \frac{(1 + x)(r + \eta)E_s}{2P_d}, \quad E_s = 1 - x\eta h. \tag{GS6}$$

Indeed $b_\gamma = (h/r)(\eta B_x(1 - h) - D_p j)$, and the original formulas give $\eta B_x(1 - h) - D_p j = -C_n$. This identity holds at literal $\beta = 0$ as well.

For a gap-free expression put

$$T_s = 1 + xh, \quad j = \eta T_s + r - 1, \quad L = T_s - xhj + x(1 - r)/a, \quad P_4 = T_s E_s + x(1 - r)(h + 1/a).$$

The unchanged source has $A = \eta P_4/(2rP_d)$ and $K = C_n(1 + y)/(\eta L)$, hence

$$\boxed{\Delta = \frac{h^2 C_n}{2r^2 P_d L}\{r(1 + y)P_4 - (1 + x)(r + \eta)E_s L\}.} \tag{GS7}$$

Here r, h, L, P_d are positive source units. Neither original gap appears in a denominator.

Write $h = 1 - x\pi$, $r = 1 - \eta\sigma$ with the inherited bounded nonnegative π, σ. Exact finite source division, not a comparison of x and η, gives

$$\boxed{\Delta = -xd_x - \eta d_\eta, \qquad d_x \sim \tfrac{1}{4}, \quad d_\eta = \frac{r + \eta}{4r^2} \sim \tfrac{1}{4}.} \tag{GS8}$$

For an explicit definition, set $d_x = -\{\Delta - \Delta|_{x=0}\}/x$; the numerator vanishes identically at $x = 0$. Formula GS7 has a unit denominator there, so this is a bounded source rational function. Also $\Delta|_{x=0} = -\eta(r + \eta)/(4r^2)$ and $\partial_x \Delta|_{x=\eta=0} = -1/4$, proving all of GS8 uniformly through the gap openings. In particular $\Delta < 0$ on the entire unchanged source chart, regardless of x/η.

This is an actual independent-root negative direction. Choose $Q = 1$, $T = -b_\gamma/(h^2K)$ and $P = hD_pT/(C + D_p)$. Then

$$S_0 = \Delta/(h^2K) < 0.$$

Subtracting the last nonnegative square in GS2 cannot repair it; $S_\gamma < 0$ at this same independent-root direction. Therefore one cannot extract the two new circuit terms in GS3, complete $\widehat{R}$ independently, and claim that the remaining source block has a nonnegative, let alone strict, bounded Gram. This obstruction survives the simultaneous R completion, which was not performed in the earlier failed angular-residue calculation.

The signed circuit remainder also persists. The other coefficient in GS3 has the exact finite source division

$$2\eta\lambda - G = -xd_c - \eta d_{\eta c}, \qquad d_c \sim \pi + 1 + \frac{1}{2a} > 0, \quad d_{\eta c} = \sigma(r+\eta)/r \geq 0. \tag{GS9}$$

The quotient has only source-unit denominators. The numerator used for the x division vanishes at $x = 0$ after subtracting the exact value $-\eta\sigma(r+\eta)/r$. Consequently the last line of GS3 is

$$-\frac{x(xd_c + \eta d_{\eta c})}{\eta c} uv\tau^2 + \frac{Gx}{c} D_0^2 WZ. \tag{GS10}$$

On the pure-testing sign chart $WZ \geq 0$ the second term is a literal positive square, but the first is negative. On $WZ < 0$ that second term changes sign too. Thus no sign of the state or source is being discarded in the proposed simultaneous join.

Two exact unit-W dependencies that a further join must preserve. Write $f_b = F_b/\eta = uvW - (x/a)Z$. Then

$$Zf_b - WJ = \frac{x}{a} Z\tau, \qquad \sqrt{x}\frac{E}{\sqrt{\eta}} = -\frac{R}{\sqrt{\lambda\eta}} + \sqrt{\frac{x}{\eta}} v\tau. \tag{GS11}$$

The first J in GS11 is the homogeneous root $(uv - x/c)Z$. These identities use W, Z multiplicatively and remain valid at $Z = 0$. They are relevant to the soft scale exposed in GS8, but do not by themselves provide a strict independent $\sqrt{x\eta}Q$ pivot or authorize another bare radial square. In particular the positive roots and state coordinates are dependent; the negative independent-root direction above is not a counterexample to the original parent.

At the original contact all roots in GS1–GS3 vanish, including P, Q, T, J, R, τ; equality and every previously retained normal are preserved. The exact full-parent identity is obtained by adding unchanged JF1 terms to GS3, so there is no dropped normal or packet. The remaining obligation is now concrete: compensate both the negative source scale $x + \eta$ in GS8 and the signed two-row remainder GS10 using the actual circuit/radial/state dependencies. No such compensation, and no all-tests admission, is asserted here.

The focused checker `check_coupled_gamma_soft.py` verifies GS1–GS3, the source completion, the gap-free native-source reduction, the two finite divisions and residues, and GS11. It does not rerun inherited suites or compare source/testing rates numerically.

Chapter 137

C14.3b: a complete source subchart from a damped two-circuit completion

Component p024c821bfb59
Source: R17/proofs/c1_small_eta/damped_gamma_source_subchart.tex
SHA256: 024c821bfb59309356b1562279a3ead111f5821540325d92c293cf2376932597
Inventory use: DG2–DG9 and all allocations except DG10 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

137.1 C14.3b: a complete source subchart from a damped two-circuit completion

Theorem 137.1. *Normalize $d = 1$ and retain the unchanged physical C14.3b source, with $x, \eta > 0$ infinitesimal, ζ a positive bounded unit, every original j and $1 - r$ opening, $\beta \geq 0$ including literal $\beta = 0$, and all inherited intersections. On the additional* pure-source *subchart*

$$x/\eta \text{ bounded}, \tag{DG1}$$

the entire original parent is finite tensor SOS for every independent projective angle and state test. No relation between a source rate and a testing rate is assumed. The proof below treats the whole local corner; the inherited remote theorem gives its complement. The source subchart x/η unbounded is not admitted by this theorem.

The sign reduction and the complete frame. Use the sign gauges to take $u, v \geq 0$. For real states with $WZ \leq 0$ the independent half-core sign lemma proved in this revision gives a direct tensor-SOS decomposition of the original signed half core; all remaining original event packets are positive source multiples of squares. Thus only $WZ > 0$ needs the following construction. No normalization or inverse of a state coordinate is used here.

Use all homogeneous JF1 definitions, and abbreviate $Q = Q_t$, $T = T_D$, $B = B_x$, $a = 1/\zeta$, $x = k^2$, $a = \eta c^2$. In particular

$$E = \eta uW - vZ = \eta(P + Q), \quad H = avW - uZ, \quad \tau = \eta cW - Z, \quad D_0 = u - cv,$$
$$J = (uv - x/c)Z, \quad R = -\sqrt{\lambda}\,\eta k D_0 W, \quad f_t = F_t/\eta, \quad \mathcal{Q} = \sqrt{q}(H - a\xi), \quad N_x = \sqrt{B}\xi.$$

The full radial packet $\mathscr{R}$ is JF2, with the strict two-root completion JF3. Its roots will be denoted R_b, f_t; both normalized pivots have positive unit residue. The angular packet is exactly

$$\mathscr{A} = \mathcal{Q}^2 - \frac{2\mu}{a\sqrt{qB}}\mathcal{Q}N_x + (1 - \mu/B)N_x^2. \tag{DG2}$$

Its normalized Schur pivot is strict by the unchanged R15/R16 source margin, including $\beta = 0$ and unbounded B. Those previously proved angular and radial blocks are used once, as their complete packets.

Damp the remaining normal and complete both circuit companions. Set the positive source unit

$$G_0 = 2\eta\lambda - 2\eta \sim 1, \qquad \epsilon = \gamma - G_0/h,$$

$$\widehat{R} = R + \frac{G_0 k}{2\sqrt{\lambda}}Q, \qquad \widehat{J} = \sqrt{\lambda}J + \frac{G_0 u}{2\sqrt{\lambda}}Q, \qquad c_U = A - \frac{G_0^2}{4\lambda}.$$

Use the exact full-D conductor and polarization GS1 to conduct only $(G_0/h)uvQ\xi$, leaving $\epsilon uvQ\xi$ unaltered. Complete *both* R and J, incorporating the entire native AUQ^2 row into the latter completion. Since $G_0/(2\lambda) = \eta - \eta/\lambda$, the complete original parent has the following exact identity:

$$\begin{aligned} F/4 = {} & \mathscr{A} + \mathscr{R} + S_0 - \lambda x(\eta P + \eta Q/\lambda)^2 + \widehat{R}^2 + \widehat{J}^2 \\ & + c_U UQ^2 + e_4 UF_t^2 + BV\xi^2 + \frac{2x}{c}uv\tau^2 + \frac{G_0 x}{c}D_0^2 WZ \\ & - e_4 yuHF_t + \kappa uF_t\xi + \frac{G_0 x}{c}PH + \epsilon uvQ\xi, \end{aligned} \tag{DG3}$$

where

$$S_0 = CP^2 + AQ^2 + D_p(P - hT)^2 + 2b_0 QT, \qquad b_0 = b' - G_0/2. \tag{DG4}$$

This is a simultaneous completion of the entire remaining normal and the coupled circuit. It retains every full-D, full-Y and higher normal of JF1. There is no second bare radial pivot and no extracted uncentered m event. The coefficient of the extra $uv\tau^2$ row is exactly $2x/c > 0$, because $2\lambda - G_0/\eta = 2$.

Two strictly positive source deficits replace the failed soft pivot. Put $K = CD_p/(C + D_p)$, $s = x + \eta$, and

$$\Delta_0 = h^2 AK - b_0^2.$$

The original source formulas imply exact finite divisions

$$\Delta_0 = xd_x + \eta d_\eta, \qquad \frac{c_U}{\eta} = xe_x + \eta e_\eta, \tag{DG5}$$

with bounded source coefficients having the following residues:

$$\begin{array}{c|c} d_x & \pi/2 + 1/4 + 1/(4a) \\ d_\eta & \sigma/2 + 3/4 \\ e_x & \pi + 1 + 1/(2a) \\ e_\eta & \sigma + 3/2 \end{array} \qquad h = 1 - x\pi, \quad r = 1 - \eta\sigma. \tag{DG6}$$

All four residues are strictly positive. Here $\pi, \sigma \geq 0$ are bounded by the unchanged physical source estimates. Thus

$$\Delta_0/s \text{ is a positive bounded unit}, \qquad c_U/(\eta s) \text{ is a positive bounded unit}. \tag{DG7}$$

These assertions compare source quantities only and do not use DG1.

For clarity, the divisions in DG5 are opening-safe. Write $C_n = (1 - r)C$, $D_j = jD_p$ and $\Theta_B = B(1 - h)$, all given by their unchanged unit-denominator source formulas. The identity $\eta\Theta_B - D_j = -C_n$ gives $b' - \gamma h/2 = -hC_n/r$. The gap-cleared harmonic coefficient is

$$\eta K = \frac{C_n D_j}{C_n(1 + xh - \sigma) + D_j\sigma}.$$

Its denominator is a positive unit, even at either opening. The quantities $\eta\lambda$, b', A/η and γh also have unit-denominator expressions after their displayed monomial factors are canceled. Both functions on the left of DG5 vanish at $x = \eta = 0$. Define their η coefficients by their exact $x = 0$ values divided by η, and then divide the remainder by x. This gives DG5 with only source-unit denominators; differentiation at the origin gives DG6. No assumption on x/η enters these exact finite divisions.

The source roots control the complete soft direction. There is the exact pure-source completion

$$S_0 = N_s^2 + N_T^2 + N_Q^2,$$

$$N_s = \sqrt{C + D_p}\left(P - \frac{hD_p}{C + D_p}T\right), \quad N_T = h\sqrt{K}\left(T + \frac{b_0}{h^2K}Q\right), \quad N_Q = \sqrt{\frac{\Delta_0}{h^2K}}Q. \tag{DG8}$$

As $\eta K \sim 1/2$, DG7 gives $N_Q/\{\sqrt{\eta s}\,Q\}$ a positive source unit. The inverse identities show

$$T = \frac{N_T}{h\sqrt{K}} - \frac{b_0}{h^2K}Q, \qquad P = \frac{N_s}{\sqrt{C + D_p}} + \frac{hD_p}{C + D_p}T. \tag{DG9}$$

Consequently P has bounded coefficients on (N_s, N_T, N_Q): the first two coefficients are $O(\sqrt{\eta})$ and the last is $O(\sqrt{\eta/s})$, which is bounded because $\eta \le s$. This is the complete P control after the full D shift; it is not an assumption that the old $E/\sqrt{\eta}$ pivot remains strict.

The negative square in DG3 is an infinitesimal Gram on these three roots. Its root is

$$\sqrt{\lambda x}(\eta P + \eta Q/\lambda) = \eta\sqrt{\lambda x}\,P + \eta\sqrt{x/\lambda}\,Q.$$

Since $\eta\lambda \sim 1/2$, the first coefficient is $O(\sqrt{\eta x})$ on the bounded allocation of P; the second, after normalizing Q by N_Q, is $O(\eta\sqrt{x/s})$. Both are infinitesimal, without DG1.

Every remaining complete normal is small on this frame. The angular-radial term in DG3 is

$$-e_4 y\eta u\left(\mathcal{Q}/\sqrt{q} + aN_x/\sqrt{B}\right) f_t.$$

Its coefficients are $O(u)$ and $O(xu)$, using $e_4\eta^2 \sim 1/2$ and q/x^2 a positive bounded unit. The term $\kappa u F_t \xi$ has coefficient $(\eta\kappa)u/\sqrt{B}$, again infinitesimal. The new PH term has coefficients

$$\frac{G_0 x}{c\sqrt{q}} \quad \text{and} \quad \frac{G_0 a x}{c\sqrt{B}}$$

on $P\mathcal{Q}$ and PN_x; these are respectively $O(\sqrt{\eta})$ and $O(x\sqrt{\eta})$, so DG9 allocates the whole term infinitesimally.

Finally $\epsilon = O(s)$ by the same gap-free source expressions. The complete, still unconducted normal has coefficient

$$\frac{\epsilon u v}{\sqrt{\eta s}\sqrt{B}} = O\left(\sqrt{s/\eta}\,uv\right) \tag{DG10}$$

on $N_Q N_x$. DG1 is used *exactly here*: it makes s/η bounded, and hence the coefficient infinitesimal for independent infinitesimal u, v. No testing comparison is imposed.

Thus the remainder on

$$(N_s, N_T, N_Q, R_b, f_t, \mathcal{Q}, N_x)$$

has a bounded Gram with strictly positive residue. The bounded-Gram lemma makes it finite tensor SOS. Add the whole positive rows $\widehat{R}^2$, $\widehat{J}^2$, $c_U U Q^2$, $e_4 U F_t^2$, $BV\xi^2$, $(2x/c)uv\tau^2$ and $(G_0 x/c)D_0^2 WZ$. The last is a literal tensor square on the current pure-testing sign chart; all radicals are source-only or testing-only. This proves the complete local parent, not merely its scalar restriction.

Equality, endpoints and the global join. At original contact $P, Q, T, J, R, \tau, F_t, \xi$ vanish, so all newly completed rows vanish too. DG3 preserves all contact normals exactly. The row formulas have no angle ratio or state-coordinate inverse; $\sqrt{uv}$ is defined as zero on a literal axis. Thus they include both literal local axes. The inherited gauges cover every angle sign. For a complex state, split the real-coefficient Hermitian parent into its real and imaginary evaluations and apply either the $WZ \le 0$ core lemma or this $WZ > 0$ proof separately to each real state. There is no assumption that both real and imaginary state components have a unit Z or W. Combine the local proof with the unchanged remote all-state theorem to obtain the all-tests assertion on DG1.

The focused checker `check_damped_gamma_closure.py` verifies the new two-circuit completion, exact source square roots, and the four source coefficients in DG6. The companion GS checker certifies the full affected-block identity. This is a genuine source subchart admission; it leaves the source chart x/η unbounded open and makes no inference there from DG10.

Chapter 138

The damped frame also enlarges the pure-testing state chart

Component p8cfada778490
Source: R17/proofs/c1_small_eta/damped_state_extension.tex
SHA256: 8cfada778490ddc7e1f29ebd9c7ef1007bb5c294c670c2f2a38d491b1309e5dd
Inventory use: DS2–DS3 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

138.1 The damped frame also enlarges the pure-testing state chart

Proposition 138.1. *On the entire unchanged C14.3b source, with no condition on x/η, the complete local parent is tensor SOS on*

$$Z \neq 0, \qquad X/Z \text{ bounded}, \qquad u^2 W/Z \text{ bounded}. \tag{DS1}$$

No bound on Y/Z is imposed. The literal $Z = 0$ endpoint is already covered by the direct sign-core theorem. This extends R16's local bounded-$W/Z, X/Z$ state chart, since u is infinitesimal.

Use the exact damped identity DG3 with $s = x + \eta$. Its strictly positive source frame and all residual allocations except DG10 are valid without the source restriction x/η bounded. Thus only $\epsilon uvQ\xi$ requires a new allocation. Put $\phi = \xi/Z = X/Z + hu$, which is bounded on DS1. There is the exact identity

$$\boxed{\epsilon uvQ\xi = \epsilon U(W/Z)E\xi - \epsilon u\phi E^2/\eta - \epsilon uvP\xi.} \tag{DS2}$$

It follows from $Q = E/\eta - P$ and $vZ = \eta uW - E$; its only testing inverse is Z^{-1}.

On the DG8 source frame, P has a bounded allocation and $\sqrt{\eta s}\,Q$ has a bounded allocation with unit coefficient. Use $E = \eta P + \eta Q$ and $\epsilon = O(s)$. The first term of DS2 has coefficients $O(\eta s)U(W/Z)$ and $O(\sqrt{\eta s})U(W/Z)$ on PN_x and $N_Q N_x$, respectively; these are infinitesimal under DS1. In the second term, the three coefficients after expanding E^2/η are

$$O(\eta s)u\phi, \qquad O(\sqrt{\eta s})u\phi, O(1)u\phi$$

on P^2, PN_Q, and N_Q^2. They are all infinitesimal, since u is infinitesimal and ϕ bounded. The last term of DS2 has coefficient $O(suv)$ on PN_x, also infinitesimal. This allocates the complete missing normal on the existing strict frame; no old $E/\sqrt{\eta}$ pivot is reintroduced. The same retained positive rows and sign-core join used in DG3 prove the assertion.

A direct state exit and the resulting frontier. For $Z \neq 0$, if X/Z is unbounded, then $(X + uZ)/Z$ is unbounded. The following portion of the original signed half core is strictly positive in the pure testing field:

$$\frac{1+v^2}{2}\{(X+uZ)^2 - (1+u^2)Z^2\} > 0. \tag{DS3}$$

Every remaining original packet has a nonnegative pure-source coefficient, including the other two half-core squares. A real closed testing field therefore gives a direct tensor-SOS certificate. This exit uses no contact or source estimate. Consequently the remaining nontrivial real state chart, after the sign gauges, can be restricted to

$$WZ > 0, \quad Z \neq 0, \quad X/Z \text{ bounded}, \quad u^2W/Z \text{ unbounded}. \tag{DS4}$$

After normalizing a W-unit state by $W = 1$, DS4 says Z/u^2 is positive infinitesimal. In addition, the separate DG source admission removes every source with x/η bounded. These are pure-source and pure-testing reductions, not a comparison of their rates.

For a complex state satisfying the modulus version of DS1, use a common testing phase to make nonzero Z real. Its real part satisfies DS1 and its imaginary part has $Z = 0$; each has the indicated complete certificate. All displayed formulas include the local axes by their literal values. The focused checker verifies DS2 and the exact original-parent packet accounting used in DS3. No all-tests claim is made for the remaining source chart x/η unbounded.

Precise remaining normalized corner and residual. Combine DG1, DS3–DS4, the inherited reciprocal-angle local theorem, and the negative-core consequence uvW/Z bounded from this revision's sign-core Schur formula. On a normalized W-unit state, the only remaining C14.3b corner has

$$\begin{gathered} x/\eta \text{ unbounded}, \qquad u, v > 0 \text{ infinitesimal}, \quad v/u \text{ infinitesimal}, \\ W = 1, \quad Z > 0, \quad Z/u^2 \text{ infinitesimal}, \quad X/Z \text{ bounded}, \quad uv/Z \text{ bounded}. \end{gathered} \tag{DS5}$$

All original source openings and $\beta \geq 0$ remain. The two last terms of DS2 are already infinitesimal Grams on the strict damped frame throughout DS5. The sole unallocated expression is exactly

$$\epsilon U(W/Z)E\xi. \tag{DS6}$$

As $s \asymp x$ on this pure-source branch, its coefficients after $E = \eta P + \eta Q$ have orders $\eta x\, UW/Z$ and $\sqrt{\eta x}\, UW/Z$ on PN_x and $N_Q N_x$. The negative-core bound on uvW/Z only rewrites the second as $O(\sqrt{\eta x}\, u/v)$; it does not bound it in independent source/testing fields. Thus DS5 is a genuine remaining chart and DS6 a precise remaining complete normal. No source/testing ordering or additional strict pivot is claimed to remove it.

Chapter 139

The entire unit-η large-ζ source, through the actual-j opening

Component p60eca5442075
Source: R17/proofs/c1_source_ends/large_zeta_unit_eta_all_tests.tex
SHA256: 60eca54420756dab5e626cf4c689eb44616412a4845bd4d0b631b4c046a881c4
Inventory use: EU1–EU12 and projective join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

139.1 The entire unit-η large-ζ source, through the actual-j opening

Theorem 139.1. *Retain the original physical C1 source and every original contact, strict-gap, nonnegative-weight, total-scale and earlier-exclusion intersection. Suppose $x = k^2, L = x\zeta$ are positive infinitesimals, ζ is unbounded and η is a positive bounded unit. Then the complete original parent is finite tensor SOS for every independent projective test and every real or complex state. No positivity or noninfinitesimal lower bound is imposed on $j_0 = \eta + r - 1$: only the actual original gap $j = \eta(1 + xh) - 1 + r > 0$ is used. Both r openings, simultaneous r, j openings and literal $\beta = 0$ are included.*

This extends R15.LZA to the whole η-unit source portion of C14.3a. It does not admit the infinitesimal-η remainder. Every identity below uses the normalized quarter-parent $\mathcal{P} = F/4$. The complete parent identity JF1–JF3 from R16 is algebraic in the original stationary source; none of its identities assumes unit ζ. We retain that identity and change its source normalization.

Opening-cleared source bounds. Put

$$a = 1/\zeta, \quad x = aL, \quad y = \eta L, \quad \theta = 1 - h = L\pi, \quad B = \beta + \tfrac{1}{2}, \quad D_p = D + \tfrac{1}{2}, \quad \rho_0 = r + \eta.$$

The original beta numerator gives bounded $\pi > 0$; $h, M, E, Q, P_d, P_4 \sim 1$ as in SB1–SB3. In particular ρ_0 is a positive bounded unit. All formulas here preserve actual $j > 0$, rather than replacing it by j_0. Define, exactly as in JF1,

$$\begin{gathered} q = \alpha + (m + B)\theta^2, \quad \mu = m - a^2 q, \quad \lambda = \mu/(\eta a^2 L), \\ T_0 = A + \alpha + \tfrac{1}{2}, \quad T_m = T_0 + m, \quad e_4 = r^2 T_m/\eta^2, \\ b' = Ar/\eta - m\theta - \alpha - \tfrac{1}{2} + rT_m L/2, \\ \kappa = 2[B\theta - D_p j/\eta - mr/\eta], \qquad \gamma = 2b'/h + 2D_p j/r - 2\eta B\theta/r. \end{gathered} \tag{EU1}$$

Finite canceled source division gives

$$\begin{gathered} C(1 - r) \sim \rho_0/2, \quad Ar \sim \eta/2, \quad D_p j \sim \rho_0/2, \quad m/(a^2 L) \sim \rho_0/2, \\ \alpha/L^2 \sim \rho_0/2, \quad B\theta/L \sim \rho_0/2, \quad q/L^2 - \rho_0(1 + \pi)/2 \in \mathrm{Inf}, \\ \mu/(a^2 L) \sim \rho_0/2, \quad rT_m \sim \rho_0/2, \quad b'/L \sim -\rho_0/4, \quad \eta\kappa \sim -\rho_0. \end{gathered} \tag{EU2}$$

The canceled denominators at $a = L = 0$ are only $1, 2, 4$; there is no r, $1 - r$, j, θ or j_0 factor. Thus these are uniform through every opening in the theorem. In particular C, A, D_p are bounded below by positive real constants, although any of them may be unbounded. Also q/L^2 and λ are positive bounded units, $\mu > 0$, and $b' = O(L)$. The positive lower bound for D_p uses bounded positive actual j and the third estimate, not positivity of j_0.

Use the complete homogeneous parent. For nonnegative testing angles u, v put $U = u^2, V = v^2$ and define

$$\begin{gathered} E_t = \eta u W - v Z, \quad Q_t = Y - r v Z/\eta, \quad P = E_t/\eta - Q_t, \quad \xi = X + h u Z, \\ R_D = v(U - x)Z, \quad T_D = R_D - uv\xi/h, \quad H = avW - uZ - a\xi, \\ F_b = (V - y)Z + vE_t, \qquad F_t = (V - y)Z + (\eta/r)vQ_t. \end{gathered} \tag{EU3}$$

All inverses in these roots are pure source. The exact full parent is

$$\begin{aligned} \mathcal{P} = {} & qH^2 + \mathscr{C} + CP^2 + AQ_t^2 + D_p(P - hT_D)^2 + 2b'Q_tT_D + \mathscr{R} \\ & + e_4UF_t^2 - e_4yu(H + a\xi)F_t + AUQ_t^2 + B(1 + V)\xi^2 - 2(\mu/a)H\xi - \mu\xi^2 \\ & + \kappa u F_t \xi + \gamma uv Q_t \xi. \end{aligned} \tag{EU4}$$

This is the complete JF1 identity, including every hard row and every first normal. Set

$$\rho = m + rT_m x/2 = O(x), \qquad K_r = T_0 + 2\rho - \rho^2/C.$$

Then $K_r/T_m \sim 1$ and K_r is bounded below by a positive real constant. The complete radial row is exactly

$$\mathscr{R} = R_b^2 + R_t^2, \qquad R_b = \frac{\sqrt{C}}{\eta}[F_b - r(1 - \rho/C)F_t], \qquad R_t = \frac{r\sqrt{K_r}}{\eta}F_t. \tag{EU5}$$

It follows that

$$\frac{F_b}{\eta} = \frac{R_b}{\sqrt{C}} + \frac{1 - \rho/C}{\sqrt{K_r}}R_t \tag{EU6}$$

is a bounded pure-source allocation even when r is infinitesimal. No independent bare $(V - y)Z$ pivot has been inserted.

A strict source frame without either gap inverse bound. Use the three original weighted triangle roots

$$N_C = \sqrt{C}\,P, \quad N_A = \sqrt{A}\,Q_t, \quad N_D = \sqrt{D_p}(P - hT_D). \tag{EU7}$$

Their positive squares are exactly the three original source-triangle terms in EU4. They give the bounded pure-source allocations

$$P = \frac{N_C}{\sqrt{C}}, \quad Q_t = \frac{N_A}{\sqrt{A}}, \quad T_D = \frac{N_C/\sqrt{C} - N_D/\sqrt{D_p}}{h}, \quad E_t = \eta(N_C/\sqrt{C} + N_A/\sqrt{A}). \tag{EU8}$$

Consequently the entire $2b'Q_tT_D$ term is an infinitesimal Gram on this strict diagonal frame. This does not cap or remove a divergent original event and does not assert boundedness of its coefficient.

The complete two-normal cancellation. The source definitions in EU1 give the exact identity

$$\frac{\eta\kappa}{r} + \gamma = 2b'/h - 2m.$$

Since $F_t = F_b - vE_t + (\eta/r)vQ_t$, both complete normal rows in EU4 therefore satisfy

$$\boxed{\kappa u F_t \xi + \gamma uv Q_t \xi = \kappa u F_b \xi + uv\{(2b'/h - 2m)Q_t - \kappa E_t\}\xi.} \tag{EU9}$$

This is the crucial simultaneous allocation. It cancels the potentially unbounded $1/r$ coefficients before any estimate is made. It neither estimates the two original rows separately nor uses their contact vanishing as a substitute for the identity.

Set $N_x = \sqrt{B}\xi$ and $\mathcal{Q} = \sqrt{q}H$. By EU2, EU6 and EU8, every coefficient of EU9 on $(N_C, N_A, R_b, R_t, N_x)$ is infinitesimal: the first uses $\eta\kappa = O(1)$ and an extra u; the second uses $b', m \in \mathrm{Inf}$ and an extra uv; the third uses $\eta\kappa = O(1)$ and an extra uv. The factors $1/\sqrt{B}$, $1/\sqrt{A}$, $1/\sqrt{C}$ and $1/\sqrt{K_r}$ are bounded. No testing-state normalization occurs.

Every remaining scalar and hard term. The complete angular block on $(\mathcal{Q}, N_x)$ is

$$\mathcal{Q}^2 - \frac{2\mu}{a\sqrt{qB}}\mathcal{Q}N_x + (1-\mu/B)N_x^2.$$

Its off-diagonal coefficient is $O(a)$ and μ/B is infinitesimal:

$$\frac{\mu}{a\sqrt{qB}} = a\,\frac{\mu/(a^2L)}{\sqrt{q/L^2}\sqrt{B}}.$$

Thus its residue is the identity, including literal $\beta = 0$. The radial-angular cross has the exact normalized expression

$$-e_4yu(H+a\xi)F_t = -\frac{rT_mL}{\sqrt{K_r}}u\left(\frac{\mathcal{Q}}{\sqrt{q}} + \frac{aN_x}{\sqrt{B}}\right)R_t. \tag{EU10}$$

Both coefficients are infinitesimal because $rT_m = O(1)$, $L/\sqrt{q} = O(1)$ and u is infinitesimal.

The coupled circuit $\mathscr{C}$ is exactly the sum of the following three whole positive squares, followed by $-\lambda x E_t^2$:

$$\sqrt{\lambda}(uv - x/c)Z, \qquad \sqrt{\lambda}\,\eta k(cv-u)W, \qquad \sqrt{2\lambda/c}\,k\sqrt{uv}(\eta cW - Z), \qquad c^2 = a/\eta, \quad k^2 = x. \tag{EU11}$$

The last negative term is an infinitesimal Gram on EU8 since λ is bounded and x infinitesimal. All radicals in EU11 are separated pure-source and pure-testing radicals.

Retain EU11's three squares, $e_4UF_t^2$, AUQ_t^2 and $BV\xi^2$ as whole positive terms. Every other term of EU4 has now been assigned explicitly to the seven-root frame

$$(N_C, N_A, N_D, R_b, R_t, \mathcal{Q}, N_x). \tag{EU12}$$

The Gram on this frame has identity residue and bounded coefficients. The bounded-Gram lemma proves finite tensor SOS for the complete parent whenever both testing angles are infinitesimal. Every estimate is a product of a source bound and a testing bound; there is no source/testing order comparison and no inverse of a mixed tensor.

All tests, endpoints and states. The roots and the whole-parent identity are homogeneous in all four state coordinates. In particular no Z inverse or state restriction was used. The positive circuit rows EU11 are defined on literal $u = 0$ or $v = 0$ and their squared identity is polynomial there. The inherited exact sign gauges give all local sign charts. The unchanged small-L, bounded-η remote theorem covers the complement of both infinitesimal testing angles on this exact source domain, retaining all its projective endpoints and openings. Applying the real homogeneous identity to real and imaginary state parts proves the complex-state assertion. All of the source unit-η portion of C14.3a is therefore all-tests admitted.

Focused receipt and exact residual. The new checker `check_large_zeta_unit_eta_source.py` verifies 15 targeted assertions: the opening-cleared source residues and their unit denominators, the exact cancellation EU9 of both entire normal rows, and the radial-angular normalization EU10. The unchanged full parent identity is JF1, independently reconstructed in R16; no historical verification suite has been rerun. The remaining C14.3a sources have η infinitesimal (and hence $r \sim 1$), with all original $1-r, j$ openings retained. No complete C1 branch is claimed closed by this theorem alone.

Chapter 140

The positive-unit e subdomain of weak-C2 R4

Component `p78dfeedee062`
Source: `R17/proofs/c1_source_ends/r4_e_unit_all_tests.tex`
SHA256: `78dfeedee06220f42324bb95b6cd5822ad05cf45d1a52af808765ccc0cd70f14`
Inventory use: R4E1-R4E16

140.1 The positive-unit e subdomain of weak-C2 R4

Theorem 140.1. *Keep the original physical stationary C2 source, $d = 1$, literal $\lambda_{\text{event}} = \beta = 0$, $n \geq 0$, all strict source gaps, the positive-infinitesimal total-scale intersection, and every inherited prior-admission exclusion. Suppose the ordered source is in R4 and $e = 1 - r$ is a positive bounded unit. Then the entire original four-state parent is a finite tensor SOS for every independent projective testing pair and every real or complex state. This admits precisely the e-unit subdomain of R4; the e-infinitesimal part is not claimed.*

Source entry. Use the exact WE1 source parameters

$$a = z/B, \quad t = y/B, \quad h = H/B, \quad l = 1 - h, \quad g = 1 - ae + haB, \quad j = 1 + aBr - l/t.$$

On R4, B, z, y, j are positive infinitesimals and t, h, l positive bounded units. Since e is a unit, $g > 0$ forces a bounded. The exact j identity gives $t_0 = l_0$. In the following residue formulas the letters on the right denote their real residues, with $h + l = 1$, $e = 1 - r$, and $g = 1 - ae \geq 0$. Put $M_* = 1 + ar - ael = g + a(r + eh) > 0$. Joint regular reduction of WE1 gives

$$(nr)_0 = \frac{e}{2(1+a)}, \quad (mg)_0 = \frac{a^2 e}{2(1+a)}, \quad (Ah)_0 = \frac{l(1+ar)}{2(1+a)}, \quad (Dj)_0 = \frac{1+ar}{2l(1+a)}, \tag{R4E1}$$

$$C_0 = \frac{M_*}{2l(1+a)}, \qquad \alpha_{*,0} := (\alpha/B^2)_0 = \frac{(1+ar)M_*}{2(1+a)}. \tag{R4E2}$$

Thus A, C, α_*, Dj are positive bounded units, while D is unbounded. The coefficient n is either a positive unit or unbounded, according as r is a unit or infinitesimal. In particular literal $n = 0$ is impossible here by R4E1. No coefficient boundary was discarded. If g is a unit then $\mu = m/a^2$ is a positive bounded unit with

$$\mu_0 = \frac{e}{2g(1+a)}. \tag{R4E3}$$

If g is infinitesimal, a is a unit, m unbounded, and mg a positive unit. These are joint regular formulas: after clearing r, g, j by the displayed products, the remaining WE1 denominators are products of $t, l, 1 - hB, \Delta$, with $\Delta_0 = 1 + a > 0$. No relative gap rates are imposed. Write $N_* = n + 1/2$, so N_*^{-1} and $N_*^{-1/2}$ are bounded and nr is bounded in all source cases.

Exact removal of the divergent D channel. Use the complete WA3–WA8 identity in the scaled roots

$$\mathcal{R} = (\xi, \eta, R_1, R_2', T)^T, \quad \xi = X + ruZ, \quad \eta = Y - huW,$$
$$R_1 = uW - vZ/t, \quad R_2' = avW - uZ, \quad T = uvW - BZ, \qquad F/4 = \mathcal{R}^T G \mathcal{R}.$$

The exact full-matrix identity RD3–RD6, proved independently of any small-h hypothesis in the R17 small-r R8 lemma, gives

$$L_D = uv\xi - \eta + lR_1 + ruvR_2' - arvT = uW + uvX - Y - vZ + jvZ,$$
$$F/4 = DL_D^2 + \mathcal{R}^T(G|_{D=0} + DjH_D)\mathcal{R}. \tag{R4E4}$$

Here the only nonzero symmetric entries of H_D are $(13) = tuv$, $(15) = -tu$, $(34) = rtuv$, $(35) = -artv$. The underlying root syzygy is exact, and this identity contains the full hard block and both complete first normals. Formal $D = 0$ does not assert a physical modified source. Put $\widehat{D} = \sqrt{D}L_D$ and solve the source-unit coefficient l:

$$R_1 = \frac{1}{l}\left(D^{-1/2}\widehat{D} + \eta - uvN_*^{-1/2}\widehat{\xi} - ruvR_2' + arvT\right), \qquad \widehat{\xi} = \sqrt{N_*}\xi. \tag{R4E5}$$

Every coefficient is bounded and separated, with no mixed inverse.

All local tests when g is a unit. The D-free matrix after the $\widehat{\xi}$ normalization has bounded pure-source coefficients. Indeed all scalar occurrences of n are through nr^2, and all normal occurrences through nr; the hard coefficient n/N_* is bounded. A, C, m, α_*, μ are bounded, and t, l, g have unit inverse. The only possible a pole after using R_2' is

$$Q_* = \frac{b_2 - \alpha_*}{a}, \quad b_2 = 2a^2\mu h + Cl^2 + (Ah)h + (B^2\alpha_* + 1/2)h^2. \tag{R4E6}$$

The exact WE3 numerator vanishes at $a = 0$ for arbitrary B, h, r, t. After cancellation its denominator has only unit factors on this g-unit chart; no D or j occurs. All other scaled WA coefficients have the same regular rewrites as RD7. Thus R4E5 and bounded Dj give a bounded full five-root Gram on $(\widehat{\xi}, \eta, \widehat{D}, R_2', T)$.

At the local testing residue $u = v = 0$, $D^{-1/2}$ vanishes and $R_1 = \eta/l$. The complete residue has independent pivots A_0/l_0^2, 1, $\alpha_{*,0}$ on $\eta, \widehat{D}, T$. Its remaining $\widehat{\xi}, R_2'$ block is

$$\begin{pmatrix} 1 + a_0^2\mu_0\nu & -a_0\mu_0\sqrt{\nu} \\ -a_0\mu_0\sqrt{\nu} & \mu_0 \end{pmatrix}, \qquad \nu = (N_*^{-1})_0 \geq 0. \tag{R4E7}$$

The determinant is exactly $\mu_0 > 0$. This includes r infinitesimal ($\nu = 0$), a infinitesimal, and every independent relative rate. For reference, the η pivot is the exact cancellation

$$(A + C + 1/2) + \frac{2[-(C + 1/2)l + Ah]}{l} + \frac{(C + 1/2)l^2 + Ah^2}{l^2} = A/l^2.$$

No scalar or normal term is omitted in this computation.

All local tests when g is infinitesimal. Here a is a unit. Use the unscaled $R_2 = vW - uZ/a$, substituting $R_2' = aR_2$ in R4E4–R4E5, and the exact centered-m matrix identity R5.2–R5.4 before R4E4. Namely, retain

$$(m - K)R_{mc}^2, \qquad R_{mc} = -\xi + uv\eta + (1 - g)R_2 + huT,$$

and replace the remaining matrix by $G|_{m=K} + \tau H_m$, where $\tau = (m - K)g$ is bounded with positive residue and

$$(H_m)_{14} = -1, \quad (H_m)_{25} = v, \quad (H_m)_{44} = 2 - g, \quad (H_m)_{45} = hu.$$

Choose a sufficiently large positive real K; $m - K > 0$ for every such K since m is unbounded. This identity and R4E4 commute, as H_m has no D dependence. All remaining coefficients are bounded: a is a unit, m has been replaced by K, τ is bounded, and the only unbounded n, D are treated by R4E5. The same three residue pivots remain. The other block is now

$$\begin{pmatrix} 1+K\nu & -(K+\tau_0)\sqrt{\nu} \\ -(K+\tau_0)\sqrt{\nu} & K+2\tau_0 \end{pmatrix}, \qquad \det = K + 2\tau_0 - \tau_0^2\nu. \tag{R4E8}$$

Since $0 \le \nu \le 2$, any real $K > 2\tau_0^2 + 1$ makes it strict. No source opening is compared with a testing rate. Restore the exact retained square. Equations R4E7–R4E8 and WA10 therefore prove the entire local parent for all independent infinitesimal or zero u, v, with arbitrary states and either sign. The finite separated lift whitens the strict pure-source matrix and handles its finitely many infinitesimal testing monomials; it introduces no mixed radicals.

Finite remote tests. If both finite angles have nonzero residue, use the inherited all-noncommuting-residues theorem, at arbitrary coefficient scales (the precise entry is the inherited `penpaper_noncommuting.tex`). Its hypotheses apply to this unchanged physical parent. It remains to give the two one-unit-angle charts. In these remote charts, positive original event coefficients may be clipped at positive real lower bounds and their exact positive excess packets retained; no contact assertion is made for the clipped polynomial.

If u is a bounded unit and v infinitesimal or zero, clip n, A, C at positive real lower bounds n_*, A_*, C_*, and retain all their excesses, the entire m and α packets. Normalize the original D event by $\widehat{J} = \sqrt{D}(uW + uvX - Y - vZ)$ and solve

$$W = \frac{D^{-1/2}\widehat{J} + Y + vZ - uvX}{u}.$$

The whole remainder has bounded coefficients on $(\widehat{J}, X, Y, Z)$. Its residue is

$$\widehat{J}^2 + n_* D_u X^2 + \tfrac{1}{2}(X + uZ)^2 + A_* D_u Y^2 \succ 0, \qquad D_u = 1 + u_0^2. \tag{R4E9}$$

Here and in the remaining residue displays the tests denote their real residues. Positivity is strict because $u_0 \neq 0$. All source cases, including unbounded n, m, have been covered by exact excess packets.

If u is infinitesimal or zero, v a bounded unit, and a a unit, m has a positive real lower bound even when it is unbounded. Clip n, m, A, C at positive real bounds and retain their excesses and α. Solve instead $Z = (uW + uvX - Y - D^{-1/2}\widehat{J})/v$. The bounded remainder residue is

$$\widehat{J}^2 + (n_* + 1/2) D_v X^2 + m_*(vW - X)^2 + (A_* + C_* + 1/2) D_v Y^2 \succ 0. \tag{R4E10}$$

This is strict since $m_* > 0$ and $v_0 \neq 0$.

Finally let a be infinitesimal in this chart. Then $g_0 = 1$, $\mu = m/a^2$ is a unit, and

$$A_0 = \frac{l}{2h}, \qquad C_0 = \frac{1}{2l}. \tag{R4E11}$$

Keep $S = aW$ as an extra legal root and write $R = uW$ everywhere else. Clip only n at a positive real n_*, retaining its excess and the α packet. Retain the actual A, C, m, D and the whole signed core. Use

$$R = Y + vZ - uvX + D^{-1/2}\widehat{J}, \qquad mM^2 = \mu(vS - aX + auvY - auZ)^2. \tag{R4E12}$$

There is no inverse of a or u here. The complete remainder is a bounded matrix on $(S, X, Y, \widehat{J}, Z)$. Its separate residue pivots are $\mu_0 V$, $(n_* + 1/2) D_v$, and 1 on $S, X, \widehat{J}$. The remaining Y, Z matrix is

$$\begin{pmatrix} (l+V)/(2h) & -v/2 \\ -v/2 & V(h+V)/(2l) \end{pmatrix}, \qquad \det = \frac{V^2(1+V)}{4hl} > 0, \quad V = v_0^2 > 0. \tag{R4E13}$$

This exact computation retains the entire signed core; it is not a state-slice argument. The possibly dependent extra root S causes no difficulty for a positive Gram. WA10 at the real angle residues now proves each complete finite remote chart.

Projective endpoints and the join. The source bounds give positive real lower bounds for n, C, A, D. Clip them at $n_*, C_*, A_*, D_* > 0$, retain their whole positive excess packets, and retain the entire original m, α packets. All remaining source coefficients are real constants. The following three exact homogeneous charts give the complete endpoint join, without any further source hypothesis.

For u unbounded put $p = 1/u$ and use $p^2F/4$. If v is bounded, the selected residue at $p = 0$ is

$$n_*D_vX^2 + C_*D_vW^2 + A_*D_vY^2 + (D_* + 1/2)(W + vX)^2 + \tfrac{1}{2}(vY - Z)^2 \succ 0. \tag{R4E14}$$

It is strict on (X, W, Y, Z) for every real v, including zero. The full homogeneous selected matrix has bounded polynomial entries.

For v unbounded put $q = 1/v$ and use $q^2F/4$. If u is bounded, write $R = uW$; every remaining occurrence of W is through R. The selected residue at $q = 0$ is

$$(n_* + 1/2)D_uX^2 + C_*(R - Y)^2 + (A_* + 1/2)D_uY^2 + D_*(uX - Z)^2 \succ 0. \tag{R4E15}$$

It is strict on (X, R, Y, Z) for every real u, including zero, and the exact selected polynomial uses no inverse of u.

If both tests are unbounded use $p = 1/u, q = 1/v$ and $p^2q^2F/4$, with the legal roots $P = pZ, Q = qZ$. The exact half-core becomes

$$\tfrac{1}{2}\{(qW + X - pqY - P)^2 + (1 + p^2)(Y - Q)^2 + (1 + q^2)(p^2X^2 + 2XP - P^2)\}.$$

Together with the selected event packets its residue is

$$(n_* + 1/2)X^2 + C_*W^2 + A_*Y^2 + D_*(X - P)^2 + \tfrac{1}{2}(Y - Q)^2 \succ 0, \tag{R4E16}$$

strict on (X, W, Y, P, Q). Dependent roots do not affect positivity. All finite reciprocal deviations are pure testing infinitesimals, so the separated strict-Gram lift applies in all three charts. Restore the original positive excesses and m, α packets. Homogeneous restoration uses pure-testing squares off the endpoints; literal $p = 0$ or $q = 0$ uses the polynomial endpoint value itself.

The finite local, finite noncommuting, both finite one-unit-angle charts, and the homogeneous infinity charts exhaust all independent tests. All four original state coordinates and both normal directions were retained. Polynomial axis values, sign gauges, and application to real and imaginary state parts supply the literal axes, both signs, and complex states. The result is an all-tests admission of the positive-unit-e source subdomain of R4, with its original physicality, strict gaps, coefficient signs, and residual intersections unchanged.

Chapter 141

C14.1b: the entire parent on bounded testing state quotients

Component `pff0a29ebf284`
Source: `R17/proofs/c1_sphere/bounded_state_full_parent.tex`
SHA256: `ff0a29ebf284bdc413f495bd78ddcfaf19366a21b4e0130de5f183ca91bb4364`
Inventory use: Inherited C14.1b bounded-state and sign exits | BS complete bounded-state proof | C14_ ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

141.1 C14.1b: the entire parent on bounded testing state quotients

Theorem 141.1. *Retain every original C14.1b source hypothesis and intersection: $x, z, \sigma > 0$ are infinitesimal, $H = h/\sigma$ is unbounded, $0 < h < 1$, and all original stationary-contact, strict-gap, coefficient-sign and residual-complement conditions remain in force. Normalize $d = 1$ by its positive pure-source value, and retain the original positive-infinitesimal total-scale intersection. For every independently chosen bounded u, infinitesimal or zero w, and real testing state with $Z \neq 0$ and $f = W/Z$ bounded, the complete original reciprocal parent is a finite tensor SOS. Both testing signs and zero angles are included. The literal $Z = 0$ chart is covered by the original-core sign exit.*

This is a full-parent state-subchart admission, not a complete C14.1b source admission. After the sign exit, the remaining real states have $WZ > 0$ and unbounded W/Z. A complex state is covered here when each of its real and imaginary parts satisfies this state chart or the sign exit. No assertion for other complex states is implicit.

Exact parent entry. Use SP1–SP6 of the companion sphere-pole reduction, including all its generic JR/JB coefficients and retained centered excesses. In particular $a_* = \ell_* = 1/4$ are the fixed JR1 caps; their exact nonnegative centered excesses remain in the parent. Also $P_\beta = uZ - (x/\sigma)wW$, $X_e = X + (xh/\sigma)wW$, $A_\theta = q(u^2 - x)W$, $q^2 = z$, and $R = wZ/q - uqW$. Set

$$s = x/\sigma, \quad L_h = \lambda/H, \quad K_A = 4\kappa H - \lambda, \quad \rho_\lambda = \lambda x/\sigma.$$

The SP3 identity $\lambda = 2H + \lambda_0$, with λ_0 bounded, implies $\lambda > 0$ unbounded and $L_h \sim 2$. Also s is infinitesimal by SP11, and $\rho_\lambda = 2xh/\sigma^2 + \lambda_0 s$ is bounded and positive. The exact source cancellation SP10 makes K_A bounded. More precisely,

$$\operatorname{res} K_A = 2 - 4 \operatorname{res} h, \qquad \operatorname{res} \chi = 4(1 - \operatorname{res} h). \tag{BS1}$$

These follow from $\alpha_\kappa = 2H(2\kappa - 1)$, $K_A = \alpha_\kappa - \lambda_0$: $\operatorname{res} \alpha_\kappa = -2 \operatorname{res} h$ and $\operatorname{res} \lambda_0 = -2(1 - \operatorname{res} h)$. The polynomial source divisibility in SP10 proves these assertions uniformly in h, rather than multiplying an unspecified infinitesimal error by H.

Use SP7 with the *whole* coefficient λ. The last line of SP6 becomes exactly

$$K_A A_\theta^2 + \chi u A_\theta R + \underbrace{\lambda\sigma P_\beta^2 + \rho_\lambda (u^2 - x)(w^2 - \sigma z) W^2}_{\mathscr{P}}. \tag{BS2}$$

This step retains the signed contact product; it has not discarded one of its negative terms.

An exact bounded conductor for the contact product. All following state quotients belong solely to the testing field. Put

$$U_h = \sqrt{h}\, uZ, \qquad Q_\lambda = \sqrt{\rho_\lambda}\, uwW, \qquad M_\lambda = \sqrt{\lambda}\, xqW.$$

The roots are homogeneous in the testing state and explicitly separated. At $Z \neq 0$, the identity $W = fZ$ gives

$$\sqrt{h}\, P_\beta = (1 - szf^2) U_h - \sqrt{h}\, sqfR. \tag{BS3}$$

Expanding all four terms of the contact product gives the exact identity

$$\begin{aligned} \mathscr{P} = {} & L_h \{(1 - szf^2) U_h - \sqrt{h}\, sqfR\}^2 + Q_\lambda^2 + M_\lambda^2 \\ & - L_h szf^2 U_h^2 - L_h s^2 \{\sqrt{h}\, qfR + zf^2 U_h\}^2. \end{aligned} \tag{BS4}$$

Both negative lines are retained exactly. Their coefficients, expanded as a Gram on (U_h, R), are infinitesimal sums of separated tensors: s, z, q are source infinitesimals, $\sqrt{h}$ is bounded, and f is bounded pure testing. The first square has residue $2U_h^2$. No inverse of $1 - szf^2$, or of any other mixed quantity, is used. The two remaining displayed squares are retained with their exact positive source weights, through either sphere endpoint.

Every coupling to the collapsing Gamma direction. The only Γ cross terms in SP6 are $-4\nu N_x(u\Gamma)$ and $R_\gamma(N_y + bR)(qw\Gamma)$. They both have exact conductors. Define the bounded infinitesimal separated expression

$$B_f = (\sigma - xzf^2)/\sqrt{h}.$$

Indeed $\sigma/\sqrt{h} = \sqrt{\sigma}\sqrt{\sigma/h}$ is infinitesimal because H is unbounded, and $x/\sqrt{h} = (x/\sigma)(\sigma/\sqrt{h})$ is infinitesimal. Direct substitution of the original roots gives

$$u\Gamma = B_f U_h - uqfA_\theta - u^2 qfR, \tag{BS5}$$

$$qw\Gamma = qzfB_f U_h - uz^2 f^2 A_\theta + z(\sigma - ufw - u^2 zf^2) R. \tag{BS6}$$

Every coefficient on the right is infinitesimal and separated for bounded u and f, including arbitrary relative source and testing rates. Thus both full first-normal couplings, and the $R\Gamma$ coupling, become infinitesimal bounded Gram entries on $(N_x, N_y, U_h, A_\theta, R)$. No free positive pivot on Γ has been imposed.

The entire remaining diagonal in Γ is

$$\left(\frac{4D}{\sigma^2} + 4bpzu^2\right)\Gamma^2, \tag{BS7}$$

a sum of two exact squares with nonnegative pure-source weights. It is retained separately. The other already-positive extracted square $2(2D+1)(uX_e)^2$, both centered cap excesses, and $4\beta D_w N_x^2$ are also retained exactly.

All remaining full-parent entries. After BS2–BS7, the only other cross terms are exactly SP5 without its two Γ terms:

$$\begin{aligned} &-4uqw(2m+2D+1)N_xN_y + uqwR_xN_xR + (c_N+zR_y)N_yR \\ &\quad + qwB_xN_xA_\theta + uA_yN_yA_\theta. \end{aligned} \tag{BS8}$$

Their coefficients are infinitesimal. The opening-uniform source bounds are $p, l \sim 1/2$, $p-l \in z\mathcal{O}_c$, m, D, b_θ, C bounded, $B_x = 8\kappa e$ infinitesimal, and $A_y = O(z)$; u and f are bounded and w infinitesimal. The three corresponding hard/radial diagonals have residues

$$[2(1+w^2)+4mw^2]_0 = 2, \quad (H_y)_0 = 2(1+u_0^2), \quad (c_G+zK_r)_0 = 2(1+u_0^2). \tag{BS9}$$

Thus no unallocated full-parent entry is hidden in the scalar argument.

Two exhaustive pure-source cases. If h is infinitesimal, BS1 gives $(K_A)_0 = 2$ and $\chi_0 = 4$. Retain $Q_\lambda^2 + M_\lambda^2$ outside the remaining Gram. In the root order $(N_x, N_y, U_h, A_\theta, R)$ its exact bounded Gram has residue

$$\operatorname{diag}(2, 2(1+u_0^2), 2) \;\oplus\; \begin{pmatrix} 2 & 2u_0 \\ 2u_0 & 2(1+u_0^2) \end{pmatrix}. \tag{BS10}$$

The last determinant is exactly 4, so this residue is strictly positive for every bounded testing residue u_0, including zero.

If h is a positive unit, the potentially signed A_θ diagonal is instead removed using the exact root identity

$$A_\theta = \frac{qfu}{\sqrt{h}}\,U_h - \frac{1}{\sqrt{\lambda}}M_\lambda. \tag{BS11}$$

Both coefficients are infinitesimal. Keep M_λ^2 in the Gram and retain Q_λ^2 separately. Since K_A, χ, u are bounded, all the A_θ terms now contribute only infinitesimal entries. In the root order $(N_x, N_y, U_h, R, M_\lambda)$ the complete residue is

$$\operatorname{diag}(2, 2(1+u_0^2), 2, 2(1+u_0^2), 1) \succ 0. \tag{BS12}$$

This includes the $\theta = 1-h$ opening. The two cases require no comparison of a testing parameter with h.

The finite bounded positive-Gram criterion supplies legal tensor SOS in either case. Its strict real residue can depend on the real number u_0. Every error is a finite sum of separated bounded factors with an infinitesimal source or testing factor; the usual finite two-factor coefficient splitting applies. There is no mixed Cholesky operation.

Endpoints, signs, and scope. Every identity used above is exact at $u = 0$ or $w = 0$. The sources $D = 0$ and either sphere endpoint were never inverted. At the additional $D = 0$ null point $u = w = W = X = Y = 0$, all roots in the strict Gram vanish and the retained BS7 contribution is exactly $4DZ^2$. Thus the certificate respects that null point and does not restore the invalid old soft-coordinate cap.

The argument itself allows signed bounded f. For $Z = 0$ the original-core sign lemma applies since $uwWZ = 0$; all original event weights remain nonnegative pure source. That exit also covers $uwWZ \le 0$ regardless of the size of W/Z. After the exact testing sign gauges, the only real-state complement on the remaining tests is therefore $WZ > 0$ and unbounded W/Z. For completeness the reciprocal core at $Z = 0$ is the manifest sum $(uwW + uX - wY)^2 + D_uY^2 + D_wX^2$. At the literal reciprocal endpoint $w = 0$ it is exactly $K_w = D_u(X^2+Y^2)$. Thus this endpoint needs no finite-v Schur formula or continuity argument.

The focused checker verifies BS2–BS8 as an identity of the entire SP6 parent, both Gamma conductors, the contact-product expansion, BS11, both strict residue calculations, and the exact extra zero. It does not assert that the remaining unbounded-state quotient is covered, or count this result as a whole-source admission.

Chapter 142

A full-parent pole extraction and a larger comparison-model state chart

Component `pfeca652aef94`
Source: `R17/proofs/c1_sphere/sphere_pole_reduction.tex`
SHA256: `feca652aef94e1f77d3bd188266d5fbe2de9e7d0342306de03b1b1f463486610`
Inventory use: Source sphere and exact pole cancellation | SP exact identities and source cancellations | C14_ ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

142.1 A full-parent pole extraction and a larger comparison-model state chart

Status. The first result below is a new exact reduction of the complete generic JR5/JB1 parent. On C14.1b it removes every unbounded h/σ coefficient from that parent and leaves one explicitly stated bounded contact-product conductor problem. It is not a positivity proof of the resulting bounded form. The second result extends the R14 JU8 comparison-model certificate to a strictly larger pure-testing state chart and to $Z = 0$. Neither result admits a complete C1 source branch.

Notation and imported exact identity. Use the generic JR/JB coefficients, the exact retained centered excesses, and the source BE2–BE8 definitions. Put

$$t = u^2, \quad \xi = w^2, \quad \Delta = t - x, \quad \delta = \xi - \sigma z, \quad q^2 = z,$$
$$H = h/\sigma, \quad e = xh/\sigma, \quad b_\theta = \beta\theta, \quad \theta = 1 - h.$$

Here $p = ab$, $l = \ell(1 - b)$ are the opening-cleared source products, and $\mu = m/(x^2 z) = \kappa H$ is an *exact* consequence of BE7–BE8. The complete parent is $F_w = E_{\mathrm{cap}} + 4\beta D_w N_x^2 + G$, with G exactly JR5/JB1. No state normal or retained excess is removed below.

Define the separated, denominator-free-in-the-tests roots

$$P_\beta = uZ - \frac{x}{\sigma} wW, \qquad X_e = X + ewW, \qquad T = t\Gamma + V.$$

They obey the exact identities

$$T = \sigma u P_\beta, \qquad N_x - hP_\beta = X_e, \tag{SP1}$$

and, using JR6,

$$t^2\Gamma^2 + (t + x)\Gamma V + \sigma t A_\theta^2 = T^2 - u\sigma\Delta A_\theta R. \tag{SP2}$$

In particular the *whole* quadratic h^2 packet and its N_x first normal combine with the corresponding N_x^2 term into $2(2D + 1)(uX_e)^2$. This identity includes $u = 0$, and does not invert it.

Put

$$\begin{aligned}
\nu &= \frac{2D - 2b_\theta + h}{\sigma},\\
\lambda &= \frac{8Dh + 4b_\theta\theta + 2h^2}{\sigma} - 4bp\sigma z = 2H + \lambda_0,\\
\lambda_0 &= \frac{4D(1+h)}{\sigma} - 2\theta\nu - 4bp\sigma z,\\
\chi &= 8(pb + l(1-b)) - 4h + 4eh(2D+1) - 8Dh\\
&\quad + 4z\{4Dbh + 4pb\sigma + (1+2m)(h-1+b)^2 + 2b_\theta\theta\}.
\end{aligned} \tag{SP3}$$

The complete coefficient B of JR/JB satisfies exactly

$$\sigma B - \frac{2h^2(2D+1)\Delta}{\sigma} = \lambda + \chi. \tag{SP4}$$

For compactness retain the following explicit, full collection of unchanged cross terms:

$$\begin{aligned}
\mathscr{C} = &-4uqw(2m+2D+1)N_xN_y + qwR_\gamma N_y\Gamma + qwbR_\gamma R\Gamma\\
&+ uqwR_xN_xR + (c_N + zR_y)N_yR + qwB_xN_xA_\theta + uA_yN_yA_\theta.
\end{aligned} \tag{SP5}$$

Every generic coefficient in SP5 is its complete JR/JB value. In particular this line keeps both hard normals and every radial first normal, including the generic beta correction in R_x.

The entire parent core, with no restriction on states, is exactly

$$\begin{aligned}
G = &\,2(2D+1)(uX_e)^2 + [2(1+\xi) + 4m\xi]N_x^2 + H_yN_y^2\\
&+ \left(\frac{4D}{\sigma^2} + 4bpzt\right)\Gamma^2 + (c_G + zK_r)R^2 - 4u\nu N_x\Gamma + \mathscr{C}\\
&+ 4\kappa HA_\theta^2 + \lambda u(\Gamma P_\beta + A_\theta R) + \chi uA_\theta R.
\end{aligned} \tag{SP6}$$

This first form already isolates all possible H poles in one shared scalar packet. It is an identity, not an entrywise comparison.

A single signed scalar contact product. There is a second exact conductor,

$$A_\theta^2 + u\Gamma P_\beta + uA_\theta R = \sigma P_\beta^2 + \frac{x}{\sigma}\Delta\delta W^2. \tag{SP7}$$

Consequently SP6 can be written

$$\begin{aligned}
G &= 2(2D+1)(uX_e)^2 + 2hP_\beta^2 + \mathscr{B} + \rho\Delta\delta W^2,\\
\rho &= \frac{2xh}{\sigma^2},\\
\mathscr{B} &= [2(1+\xi) + 4m\xi]N_x^2 + H_yN_y^2 + \left(\frac{4D}{\sigma^2} + 4bpzt\right)\Gamma^2 + (c_G + zK_r)R^2\\
&\quad - 4u\nu N_x\Gamma + \mathscr{C} + \alpha_\kappa A_\theta^2 + \lambda_0 u(\Gamma P_\beta + A_\theta R) + \chi uA_\theta R,\\
\alpha_\kappa &= 2H(2\kappa - 1).
\end{aligned} \tag{SP8}$$

This is the promised exact full-parent reduction. The two squares displayed before $\mathscr{B}$ have nonnegative pure-source weights. The term $\rho\Delta\delta W^2$ is signed; it has not been called a square or bounded below by a mixed-rate comparison.

Opening-uniform source bounds on C14.1b. The cancellation in the hard first normal is essential. BE3 gives

$$\nu = \frac{2D}{\sigma} + \frac{H_0 - ez(J-\sigma)}{P}. \tag{SP9}$$

Thus ν is bounded even when H is unbounded. Equations SP3 show that λ_0 and χ are bounded, since $D = O(\sigma^2)$, e is infinitesimal, $0 < h < 1$, and $b_\theta = O(h+\sigma)$. No lower bound on h or θ is used.

The apparently unbounded coefficient α_κ is bounded as well, by an exact divisibility. Let

$$N_\kappa = (1+x)(1+\sigma bz)(1+\sigma z) - MP, \quad F_x = \frac{N_\kappa|_{\sigma=0}}{x}, \quad F_\sigma = \frac{N_\kappa - N_\kappa|_{\sigma=0}}{\sigma}.$$

Both displayed quotients are polynomials in the bounded source letters x, z, b, h, σ. The source M here is the geometry factor $1 + xbz - xz\theta$, not M_β. Exactly,

$$\alpha_\kappa = \frac{2(hF_\sigma + eF_x)}{MP}. \tag{SP10}$$

Since M, P are positive units this proves the asserted bound without using a rate comparison between a source and a test.

On C14.1b, H unbounded and BE7 force σ/x unbounded: if it were bounded then both terms of BE7 would be bounded. Thus x/σ is infinitesimal. The source sphere gives

$$\rho = \frac{2xH_0}{\sigma B} + \frac{2\mathcal{H}}{B} b_0^2, \tag{SP11}$$

so ρ is bounded through both sphere endpoints. Its residue is $2\,\mathrm{res}(b_0^2)$ in the large-L chart. The coefficients in SP8, including the coefficients in SP5, are now all bounded for bounded u and infinitesimal w. Here $C = O(z)$, p, l are bounded and $p - l \in z\mathcal{O}_c$ uniformly through b and $1-b$ openings; the latter follows from the same opening-cleared source numerators used in JR, written with e rather than imposing bounded H. Also $B_x = 8\mu x = 8\kappa e$ is bounded.

One may introduce the legal root $B_\delta = \delta W/q$. Then the last term of SP8 is exactly $\rho A_\theta B_\delta$. The remaining conductor problem is now precise: allocate this bounded signed cross term together with $\mathscr{B}$ and the two retained squares, using

$$\frac{xq}{\sigma} B_\delta = uqR + qA_\theta - wP_\beta, \qquad u\delta W = \sigma qR - w\Gamma. \tag{SP12}$$

Neither relation permits cancellation of u, w, a state coordinate, or the infinitesimal x/σ while retaining a bounded coefficient array. In particular no B_δ^2 pivot has been proved. This is an exact bounded full-parent reduction on C14.1b, not its admission. Nor is $\mathscr{B}$ asserted to be positive by itself: for example, at the algebraic specialization $z = 0$ one has exactly $\alpha_\kappa = -2h/(1+x+\sigma)$, whose residue is negative when h is a unit. Thus the two extracted squares, $\mathscr{B}$, and the signed contact product still require a joint allocation.

The algebraic identities SP1–SP12 also apply in the common BE source domain with L a positive unit. The preceding claim that every entry of SP5 is bounded uses large L; on C14.2 the separate $r \to 1$ opening can make C unbounded and is not silently removed here. On C14.1a SP8 remains exact but SP11 does not in general bound ρ; the following result addresses its comparison model instead.

A larger state chart for the complete JU8 model. Retain the C14.1a assumptions and notation of R14.1–R14.2: u infinitesimal, τ bounded, ε infinitesimal, $x = c\varepsilon$, $\sigma = \tau\varepsilon$, and $0 < c \le 1/2$. Suppose $Z \neq 0$ and the *pure-testing* quotient uW/Z is bounded. Unlike R14, W/Z itself may be unbounded. Put $E = qW/Z$, $U = |u|$, $V = |E|$, $a = UV$, and use the exact circuit roots $d_i = A_i - G$ of R14.2. Then $uE = q(uW/Z)$ is infinitesimal, although no size is assigned to E.

When c is infinitesimal, R14.3 still applies because its coefficient is uE/d_0, which is infinitesimal on this larger chart. When c is a positive unit, the new identity is

$$uEK = \mathrm{sgn}(uE)\left\{-\frac{a}{\sqrt{2c}}d_1 + \frac{a}{\sqrt{2c}}d_2 + \frac{U\sqrt{\tau}}{\sqrt{2}\,c}d_3\right\}, \qquad K = \sigma - u^2E^2. \tag{SP13}$$

Every coefficient is infinitesimal. Crucially, the coefficient $V\sqrt{\varepsilon}$ in R14.4 has disappeared; no bound on V is needed. The identity follows by substituting $A_1 = \sqrt{2c}U^2V^2$, $A_2 = \sqrt{2\sigma\tau}U$, $A_3 = \sqrt{2\tau}xV$, $G = \sqrt{2\tau x}UV$, and $\sigma c = \tau x$.

Substitute SP13 (or R14.3) in the exact R14.1 identity. Its radial diagonal is $1 + u^2E^2$ of residue one; its other diagonal entries are the identity on the same fourteen circuit roots. The remaining coupling is $\tau u A_\theta R$, with infinitesimal coefficient. The finite bounded positive-Gram criterion therefore proves the *whole JU8 model*, including its first radial normal, on this larger chart. All radicals are taken on their explicitly separated pure-source and pure-testing factors, as in R14.2.

At the omitted literal chart $Z = 0$, the whole model has the direct SOS

$$\frac{\mathcal{L}}{2\kappa_s} = (uqW)^2 + (uwW)^2 + (1 + 2\tau)\{q(u^2 - x)W\}^2. \tag{SP14}$$

The sole still-uncovered model state chart is therefore $uW \neq 0$ and $0 < |Z/(uW)|$ infinitesimal, a comparison entirely in the testing field. The full JU3 correction is still unallocated even on the newly covered model chart. No full-parent admission is inferred from SP13–SP14.

Verification and boundaries. The accompanying checker verifies the complete SP6 and SP8 polynomial identities relative to the imported full generic JR/JB parent, every new conductor and source cancellation, the polynomial divisions in SP10, and SP13–SP14. It does not replay the old contact-jet proof or turn a symbolic equality into a positivity theorem. All identities include zero tests and all signs, retain the $D = 0$ extra zero, and apply separately to real and imaginary state parts. In particular, at $u = w = W = X = Y = 0$ SP8 gives exactly $4DZ^2$.

Chapter 143

The damped completion at large zeta with bounded L over eta

Component `pc720c1e06aca`
Source: `R17/proofs/matrix_mechanism/large_zeta_damped_extension.tex`
SHA256: `c720c1e06aca1fd0ba8195fbc6f61e6d3673ec7de8cd068867fc03532beff8f7`
Inventory use: LD2–LD13 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

143.1 The damped completion at large zeta with bounded L over eta

Theorem 143.1. *Normalize $d = 1$ and retain the unchanged physical C14.3a source, its actual stationary contact, every strict source gap, $\beta \geq 0$ including literal $\beta = 0$, the original total-scale intersection and every earlier exclusion. Suppose $x, L = x\zeta, \eta$ are positive infinitesimals, ζ is unbounded, and*

$$L/\eta \text{ is bounded.} \tag{LD1}$$

Then the entire original parent is finite tensor SOS at every independent projective angle and state test, real or complex. This is a whole-state, all-tests source-subchart admission. No condition is imposed on the testing angle ratio or on the source ratio a/η, where $a = 1/\zeta$. The source subchart with η infinitesimal and L/η unbounded is not admitted here.

The proof uses the exact complete damped identity DG3, but its source scale is $L + \eta$ rather than $x + \eta$. This change must be proved: one cannot invoke the unit-ζ estimates by substituting an infinitesimal a into their displayed $1/a$ coefficients. All formulas below use the actual physical source and the original contact.

Source variables and opening-cleared coefficients. Put

$$a = 1/\zeta, \quad x = aL, \quad y = \eta L, \quad h = 1 - L\pi, \quad r = 1 - \eta\sigma, \quad B = \beta + \tfrac{1}{2}.$$

The original source estimates give bounded $\pi, \sigma \geq 0$, and actual $j > 0$ says $0 < \sigma < 1 + xh$, $j = \eta(1 + xh - \sigma) > 0$. Neither opening has a positive-residue lower bound. The checked EU/EZ source arrays give

$$\frac{m}{a^2 L} \sim \frac{1}{2}, \quad \frac{\alpha}{L^2} \sim \frac{1}{2}, \quad \frac{q}{L^2} - \frac{1+\pi}{2} \in \mathrm{Inf}, \quad \frac{\mu}{a^2 L} \sim \frac{1}{2},$$
$$\eta\lambda \sim \frac{1}{2}, \quad \eta K \sim \frac{1}{2}, \quad \eta\kappa \sim -1, \quad \eta^2 e_4 \sim \frac{1}{2}, \quad b' = O(L), \tag{LD2}$$

where $q = \alpha + (m + B)(1 - h)^2$, $\mu = m - a^2 q$, $\lambda = \mu/(\eta a^2 L)$ and $K = CD_p/(C + D_p)$.

Here is a completely regular specification of the new source expressions. Write

$$K_x = 1 + x,\ R_s = 1 + xr,\ T_s = 1 + xh,\ M_s = R_s - aL\pi,\ E_s = 1 - x\eta h,\ Q_s = 1 - Lr,$$

$$P_d = 1 + 2xh + rx^2h + rxL + L + \eta\{K_xL + x^2hL\pi\}, \qquad P_4 = T_sE_s + (1-r)(xh + L).$$

Define the opening-cleared quantities

$$C_n = \frac{K_x(r+\eta)E_s}{2P_d}, \qquad D_j = \frac{K_x(1+y)(r+\eta)}{2P_d}, \qquad A_\eta = \frac{P_4}{2rP_d},$$

$$m_n = \frac{K_x(r+\eta)h(1+y)}{2M_sP_d}, \quad \alpha_n = \frac{(r+\eta)\{aR_sh(1+y) + M_sE_s\}}{2Q_sP_d}, \quad B_n = \frac{K_x(r+\eta)(1+ah)}{2P_d}. \tag{LD3}$$

Thus $C_n = (1-r)C$, $D_j = jD_p$, $A_\eta = A/\eta$, $m_n = m/(a^2L)$, $\alpha_n = \alpha/L^2$ and $B_n = B(1-h)/L$. In particular

$$q_n := q/L^2 = \alpha_n + m\pi^2 + B_n\pi, \qquad \Lambda := \eta\lambda = m_n - Lq_n,$$

$$\eta K = \frac{C_nD_j}{C_n(T_s - \sigma) + D_j\sigma}. \tag{LD4}$$

There is no a^{-1} in (LD3)–(LD4). Their denominators at $a = L = \eta = 0$ have residues 1, 2 or 1/2, independently of π, σ. The remaining definitions are

$$T_m = A + \alpha + \tfrac{1}{2} + m, \quad b' = A_\eta r - mL\pi - \alpha - \tfrac{1}{2} + rT_mL/2,$$

$$G_f := \gamma h = 2b' + 2D_jh/r - 2\eta LB_nh/r, \quad G_0 = 2\Lambda - 2\eta \sim 1, \quad b_0 = b' - G_0/2. \tag{LD5}$$

These are the actual native-source expressions used in the completion.

Positive source deficits at the new scale. Set $s = L + \eta$ and retain DG's definitions

$$\Delta_0 = h^2AK - b_0^2, \qquad c_U = A - \frac{G_0^2}{4\lambda}, \qquad \epsilon = \gamma - G_0/h.$$

The regular source formulas above give exact finite divisions

$$\Delta_0 = Ld_L + \eta d_\eta, \qquad \frac{c_U}{\eta} = Le_L + \eta e_\eta, \qquad G_f - G_0 = Lf_L + \eta f_\eta, \tag{LD6}$$

with bounded coefficients and residues

	L coefficient	η coefficient
Δ_0	$\pi/2 + 1/4$	$\sigma/2 + 3/4$
c_U/η	$\pi + 1/2$	$\sigma + 3/2$
$G_f - G_0$	$\pi + 1/2$	$\sigma + 2$

(LD7)

(as usual the bounded π, σ on the right can be replaced by their real residues). To justify the finite divisions explicitly, each expression on the left of (LD6) vanishes at $L = \eta = 0$ *for arbitrary* a. Subtract its exact value at $L = 0$, divide that difference by L, and divide the $L = 0$ value by η. Clearing the source-unit denominators proves each divisibility in a polynomial ring. Evaluation of these quotients at $a = L = \eta = 0$ gives (LD7). There is consequently no hidden pure-a remainder.

All four coefficients for the first two rows have strictly positive residue. Hence

$$\Delta_0/s \text{ and } c_U/(\eta s) \text{ are positive bounded source units}, \qquad \epsilon = O(s). \tag{LD8}$$

This source calculation itself does not use (LD1).

The same complete damped identity, without a state chart. Use DG's unshifted $H = avW - uZ$; the strict angular root is $\mathcal{Q} = \sqrt{q}(H - a\xi)$, so this H differs from the shifted letter H in EU3. Retain every other homogeneous JF/DG definition. In particular

$$E = \eta(P + Q), \quad N_x = \sqrt{B}\xi, \quad \tau = \eta cW - Z, \quad D_0 = u - cv, \quad c^2 = a/\eta.$$

After the pure-testing sign gauges, the half-core lemma supplies all $WZ \le 0$ states. On $WZ > 0$, the exact identity DG3 is valid for this source without any change:

$$\begin{aligned} F/4 = {} & \mathscr{A} + \mathscr{R} + S_0 - \lambda x(\eta P + \eta Q/\lambda)^2 + \widehat{R}^2 + \widehat{J}^2 \\ & + c_U U Q^2 + e_4 U F_t^2 + BV\xi^2 + \frac{2x}{c} uv\tau^2 + \frac{G_0 x}{c} D_0^2 WZ \\ & - e_4 yuHF_t + \kappa u F_t \xi + \frac{G_0 x}{c} PH + \epsilon uvQ\xi. \end{aligned} \tag{LD9}$$

Here $S_0 = CP^2 + AQ^2 + D_p(P - hT)^2 + 2b_0 QT$, $\widehat{R} = R + G_0\sqrt{x}Q/(2\sqrt{\lambda})$ and $\widehat{J} = \sqrt{\lambda}J + G_0 uQ/(2\sqrt{\lambda})$. The identity conducts only $G_0 uvQ\xi/h$ and completes both circuit companions exactly, retaining every first normal and hard coupling. Its derivation uses only $x = k^2$, $c^2 = a/\eta$ and the homogeneous JF conductor, not a bounded reciprocal of a or c.

Strict source, radial and angular frames. Use the exact source roots

$$N_s = \sqrt{C + D_p}\left(P - \frac{hD_p}{C + D_p}T\right), \quad N_T = h\sqrt{K}\left(T + \frac{b_0}{h^2 K}Q\right), \quad N_Q = \sqrt{\frac{\Delta_0}{h^2 K}}Q. \tag{LD10}$$

Their squares sum to S_0, and $N_Q/[\sqrt{\eta s}\,Q]$ is a positive source unit. The inverse source identities give bounded coefficients for P on this frame: the coefficients of N_s, N_T are $O(\sqrt{\eta})$ and that of N_Q is $O(\sqrt{\eta/s})$. This uses only $\eta \le s$, so no old $E/\sqrt{\eta}$ pivot is imported.

The complete radial packet has its inherited strict roots R_b, f_t, with $f_t = F_t/\eta$. Indeed $\rho = m + rT_m x/2 = O(x)$ and $K_r \sim 1/2$. The complete angular block is

$$\mathscr{A} = \mathcal{Q}^2 - \frac{2\mu}{a\sqrt{qB}}\mathcal{Q}N_x + (1 - \mu/B)N_x^2.$$

It has identity residue, since

$$\frac{\mu}{a\sqrt{qB}} = \frac{a[\mu/(a^2 L)]}{\sqrt{q/L^2}\sqrt{B}} = O(a), \qquad \mu/B \in \mathrm{Inf}\,. \tag{LD11}$$

This includes unbounded B and literal $\beta = 0$.

The entire remainder has infinitesimal coefficients. The negative-square root in (LD9) has, on the source frame, the same coefficient bounds as DG:

$$\eta\sqrt{\lambda x}\,P + \eta\sqrt{x/\lambda}\,Q, \qquad O(\sqrt{\eta x}), \quad O(\eta\sqrt{x/s}). \tag{LD12}$$

They are infinitesimal; here $x = aL$ and no comparison of a with η is needed. For the radial-angular term use $H = \mathcal{Q}/\sqrt{q} + aN_x/\sqrt{B}$ and $F_t = \eta f_t$. Since $e_4 y\eta = (e_4\eta^2)L = O(L)$ and $\sqrt{q} \asymp L$, its two coefficients are $O(u)$ and $O(xu)$. The κ term has coefficient $(\eta\kappa)u/\sqrt{B}$, which is also infinitesimal.

The new PH term improves at this source end:

$$\frac{G_0 x}{c\sqrt{q}} = O(\sqrt{a\eta}), \qquad \frac{G_0 ax}{c\sqrt{B}} = O(x\sqrt{a\eta}). \tag{LD13}$$

Thus the bounded P allocation makes both of its angular coefficients infinitesimal. Finally, the complete unprocessed normal is

$$\frac{\epsilon uv}{\sqrt{\eta s}\sqrt{B}}N_Q N_x \quad \text{up to a bounded positive source-unit factor.}$$

Its coefficient is

$$O\Big(\sqrt{1+L/\eta}\,uv\Big)\,, \tag{LD14}$$

which is infinitesimal under the bounded-source-ratio hypothesis (LD1). This is the only use of that hypothesis.

Consequently the complete remainder on $(N_s, N_T, N_Q, R_b, f_t, \mathcal{Q}, N_x)$ has a bounded strict-residue Gram. The bounded-Gram lemma gives its finite tensor SOS. Every other term in (LD9) is a whole positive square: in particular $c_U > 0$, $G_0 > 0$, and uv, WZ are nonnegative pure-testing factors on this sign chart. No mixed inverse or radical is taken.

All tests and exact remaining source end. No state normalization, state-coordinate inverse or testing ratio occurs in this proof. The division-free circuit rows and $\sqrt{uv} = 0$ include both literal local axes. The half-core exit and the exact angle sign gauges cover every real state; splitting the real-coefficient Hermitian form covers complex states. The unchanged small-L, bounded-η remote theorem covers all other projective testing pairs, including literal infinities. This proves the theorem with every original source opening retained.

Together with EU's unit-η theorem, this removes from the large ζ source end all unit-η sources and all infinitesimal-η sources with bounded L/η. It leaves the latter ratio unbounded as an explicit remaining source obligation. No C1 branch as a whole is claimed closed.

The focused checker `check_large_zeta_damped_extension.py` passes sixteen exact checks: the new unit source arrays, all six finite-division coefficients, vanishing for arbitrary a, the source-unit denominators and the changed angular/PH normalizations. The complete affected-block identity is DG3 and its existing exact checker; no historical suite or numerical rate experiment is used.

Chapter 144

Large zeta: the enlarged state chart and the exact remaining corner

Component `p8635816e42a0`
Source: `R17/proofs/matrix_mechanism/large_zeta_damped_state_extension.tex`
SHA256: `8635816e42a069515b3761e7e8f06e50217e0e08504d0ea0480d13a0c1614d03`
Inventory use: LS2–LS5 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

144.1 Large zeta: the enlarged state chart and the exact remaining corner

Proposition 144.1. *Retain the entire physical C14.3a source with $a = 1/\zeta$, L and η positive infinitesimals, $x = aL$, all original strict source gaps, $\beta \geq 0$ and every inherited intersection. No condition on L/η is imposed. At every independent local testing pair, the complete original parent is finite tensor SOS on*

$$Z \neq 0, \qquad X/Z \text{ bounded}, \qquad u^2 W/Z \text{ bounded}. \tag{LS1}$$

The state coordinate Y/Z is arbitrary. Literal $Z = 0$ is also admitted. This enlarges the EZ bounded-$W/Z, X/Z$ state chart. For complex states the corresponding bounded modulus ratios suffice.

Use the complete identity (LD9), its source scale $s = L + \eta$, and the strict seven-root frame in (LD10)–(LD11). All its coefficient bounds except (LD14) were proved without a restriction on L/η. The only unprocessed expression is therefore $\epsilon uvQ\xi$. On $Z \neq 0$ set

$$\phi = \xi/Z = X/Z + hu.$$

The same exact identity as DS2 gives

$$\epsilon uvQ\xi = \epsilon U(W/Z)E\xi - \epsilon u\phi E^2/\eta - \epsilon uvP\xi, \qquad U = u^2. \tag{LS2}$$

It follows by substituting $Q = E/\eta - P$ and $vZ = \eta uW - E$. Its only testing inverse is Z^{-1}; no mixed expression is inverted.

The source completion gives bounded linear membership for P and

$$Q = \frac{\omega}{\sqrt{\eta s}} N_Q,$$

where ω and ω^{-1} are bounded positive pure-source units. Use $E = \eta(P + Q)$, $\epsilon = O(s)$ and $\xi = N_x/\sqrt{B}$. The coefficients of the first term in (LS2) on PN_x and $N_Q N_x$ have respective orders

$$O(\eta s)U(W/Z), \qquad O(\sqrt{\eta s})U(W/Z).$$

They are infinitesimal on (LS1). The coefficients of the second term on P^2, PN_Q and N_Q^2 have orders

$$O(\eta s)u\phi, \qquad O(\sqrt{\eta s})u\phi, \qquad O(1)u\phi. \tag{LS3}$$

All are infinitesimal because ϕ is bounded and u is infinitesimal. The last term has coefficient $O(suv)$ on PN_x. The $1/\sqrt{B}$ factors are bounded and do not alter these estimates. Consequently the whole missing normal is an infinitesimal Gram on the existing strict frame, without using L/η or the old $E/\sqrt{\eta}$ pivot. Restore the same whole positive rows in (LD9) on $WZ > 0$; use the half-core sign exit on $WZ \le 0$. This proves the complete local parent on (LS1).

The original-parent state exit. If $Z \neq 0$ and X/Z is unbounded, then for infinitesimal u the pure-testing scalar

$$\frac{1+v^2}{2}\{(X+uZ)^2 - (1+u^2)Z^2\} \tag{LS4}$$

is strictly positive. Indeed division by the positive testing square Z^2 leaves $(1+v^2)[(X/Z+u)^2 - (1+u^2)]/2 > 0$. Every remaining original packet, including both other half-core squares, has a nonnegative pure-source coefficient. Thus (LS4) gives a direct tensor-SOS exit, independent of the source scale. The $Z = 0$ endpoint is already a sum of the original positive packets.

For complex states satisfying the modulus version of (LS1), choose a pure-testing phase making Z real and positive. The real part satisfies (LS1), and the imaginary part has $Z = 0$. Their two real certificates add to the original Hermitian evaluation. No bound on Y/Z or state norm is required. Literal local angle axes are included by (LS2) and the division-free circuit rows.

A source-independent consequence of the negative core. The exact half-core Schur formula supplies a useful additional testing bound. In the sign gauge $u, v \ge 0$, write

$$p = uv, \quad z_t = 1 + v^2 + u^2v^2, \quad H_t = \frac{1}{2}\begin{pmatrix} z_t & -p \\ -p & z_t \end{pmatrix}.$$

The hard block on (X, Y) is positive definite, and its Schur form on (W, Z) is

$$\frac{(1+u^2)(1+v^2)}{2(z_t^2 - p^2)}\{p^2(W^2+Z^2) - 2pz_tWZ\}. \tag{LS5}$$

If the pure-testing half-core is nonnegative, it is itself a testing square and the original positive events complete the tensor-SOS exit. Otherwise (LS5) must be negative. Hence $u, v > 0$, $WZ > 0$, and

$$uv\left(\frac{W}{Z} + \frac{Z}{W}\right) < 2z_t.$$

Both summands are positive. Since u, v are local infinitesimals, $z_t \sim 1$, so in particular uvW/Z is bounded. This deduction uses only a pure-testing inequality; no ordering on the tensor ring or comparison to a source scale is involved.

Exact remaining normalized corner. Combine the unit-η admission EU, the source admission (LD1), the state admission (LS1), the direct exit (LS4), the half-core exit and the inherited remote theorem. Every unadmitted real test on this large-ζ source end can then be normalized by the nonzero pure-testing coordinate W to $W = 1$, and must lie in

$$\begin{gathered} a, L, \eta > 0 \text{ infinitesimal}, \quad x = aL, \qquad L/\eta \text{ unbounded}, \\ u, v > 0 \text{ infinitesimal}, \qquad W = 1, \quad Z > 0, \\ X/Z \text{ bounded}, \qquad Z/u^2 \text{ infinitesimal}, \qquad uv/Z \text{ bounded}, \qquad v/u \text{ infinitesimal}. \end{gathered} \tag{LS6}$$

All original source openings and $\beta \geq 0$ remain imposed, and Y is unrestricted. The new angle-ratio conclusion here is derived, not imported from a unit-ζ theorem: exactly

$$\frac{v}{u} = \frac{uv}{Z}\frac{Z}{u^2},$$

a bounded pure-testing element times an infinitesimal. Normalization by W preserves the two state ratios and multiplies the original form by a positive pure-testing square. No assertion that this residual intersection is nonempty is made.

On (LS6), the second and third terms in (LS2) already have infinitesimal Gram coefficients by (LS3). The sole unallocated complete normal is exactly

$$\epsilon U(W/Z)E\xi. \tag{LS7}$$

Since $s \asymp L$ on this source branch, its two coefficients on PN_x and N_QN_x have orders $\eta L\,UW/Z$ and $\sqrt{\eta L}\,UW/Z$. The negative-core bound rewrites the latter as $O(\sqrt{\eta L}\,u/v)$, which need not be bounded in the independent source/testing tensor setting. Neither (LS6) nor (LS7) is claimed to have a further certificate here.

Identity (LS2) and the original-parent accounting in (LS4) are exactly the already checked DS identities; the new source estimates are the sixteen receipts for (LD2)–(LD8). No duplicate checker or historical-suite replay is needed for this corollary.

Chapter 145

A pure-testing sign exit for the complete original parent

Component `p894ae380fb8a`
Source: `R17/proofs/root/core_sign_lemma.tex`
SHA256: `894ae380fb8a6a6568e79fb489902d4eea8708cfcf39cc7256062b274d6c91db`
Inventory use: entire source-free sign lemma | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

145.1 A pure-testing sign exit for the complete original parent

The source event coefficients are nonnegative and $d > 0$ is pure source. It suffices to treat the quarter-core at $d = 1$,

$$Q_0 = \tfrac{1}{2}\{(uW + uvX - Y - vZ)^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_uD_vZ^2\}.$$

All variables in Q_0 belong to the testing field. Put $b = 1 + v^2 + u^2v^2$ and $\Delta = b^2 - u^2v^2$. The hard matrix on (X, Y) is

$$H = \tfrac{1}{2}\begin{pmatrix} b & -uv \\ -uv & b \end{pmatrix} \succ 0.$$

Indeed $b - |uv| = (|uv| - 1/2)^2 + 3/4 + v^2 > 0$. The exact Schur complement on (W, Z) is

$$S = \frac{D_uD_v}{2\Delta}\begin{pmatrix} u^2v^2 & -uvb \\ -uvb & u^2v^2 \end{pmatrix}.$$

The inverse in this calculation is pure testing and is legitimate. Thus $uvWZ \le 0$ implies $Q_0 \ge 0$: the diagonal terms of S are nonnegative and its cross contribution is also nonnegative. At each such testing point, Q_0 is a nonnegative element of a real closed field and has a pure-testing square root. Restoring d and all nonnegative source-weighted original events gives finite tensor SOS directly. Literal $u = 0$ or $v = 0$ is included.

The exact sign gauges can always arrange $u, v \ge 0$. Reversing the sign of u together with Y, Z preserves every original squared packet, as does reversing the sign of v together with X, Z. Consequently any remaining real state in the nontrivial local argument satisfies $WZ > 0$. A certificate valid for every such state, combined with this exit, covers all real states. Applying the real assertion to real and imaginary parts covers complex states. This is a pure-testing routing lemma, not a comparison with a source scale and not a new source-shape admission by itself.

Chapter 146

The complete ordered weak-C2 shape R5 is admitted

Component `p8da9ea6ed9e5`
Source: `R17/proofs/weak_gaps/r5_all_tests.tex`
SHA256: `8da9ea6ed9e5a0f8863a59ee27fef3ce89b422aa4d584ae0fe2152e8dc934acd`
Inventory use: R5.1-R5.17

146.1 The complete ordered weak-C2 shape R5 is admitted

Theorem 146.1. *Normalize $d = 1$. Retain the original physical C2 source, its stationary contact, every strict source gap, all coefficient signs, the positive-infinitesimal total-scale intersection and every original prior-admission exclusion. Suppose $B, z, y, r > 0$ are infinitesimal and $t = y/B$, $h = H/B$, $l = 1 - h = L/B$, and $j = 1 + zr - l/t$ are positive bounded units. Then the entire original four-state parent is a finite tensor SOS at every independently chosen projective testing pair, with both signs, literal zero/infinite angles, and real or complex states. These are precisely the unchanged R5 source hypotheses. There is no comparison of source and testing rates.*

Source entry and all gaps. Put $a = z/B$ and $g = G/B = 1 - a(1 - r - H)$. Since r, H are infinitesimal, $g > 0$ implies a bounded and $a_0 \in [0, 1]$. Write $\rho = a_0$, $\chi = h_0 \in (0, 1)$, $\tau = t_0 > 1 - \chi$ and $\ell = 1 - \chi$, $\delta = \tau - \ell > 0$. The inequality for τ follows from j being a positive unit. The exact source formulas WE1 give jointly

$$\begin{aligned} (nr)_0 = \frac{1}{2(1+\rho)}, \quad A_0 = \frac{\tau}{2\chi(1+\rho)}, \quad C_0 = \frac{\chi + \tau - \rho\ell}{2\ell(1+\rho)}, \\ D_0 = \frac{1}{2(1+\rho)\delta}, \quad (\alpha/B^2)_0 = \frac{\chi + \tau - \rho\ell}{2(1+\rho)}. \end{aligned} \tag{R5.1}$$

All displayed quantities are positive units; n is unbounded and α infinitesimal. Thus literal $n = 0$ is impossible on this shape, by the exact identity $r(2n + 1) \asymp 1$, rather than being discarded. If $\rho < 1$, $\mu = m/a^2$ is bounded with residue $1/[2(1 - \rho^2)]$. If $\rho = 1$, g is infinitesimal, m unbounded and $(mg)_0 = 1/4$. These statements retain arbitrary independent r, g, B openings. In particular the source is automatically in the positive-infinitesimal total-scale regime. No nonemptiness of the residual intersection is asserted.

An exact centered-excess identity for the entire Gram. Use the unnormalized version of the established complete WA3–WA8 identity, with roots

$$\xi = X + ruZ, \quad \eta = Y - huW, \quad R_1 = uW - t^{-1}vZ, \quad R_2 = vW - a^{-1}uZ, \quad T = uvW - BZ.$$

Let G be its five-by-five matrix, so $F/4 = \mathcal{R}^T G \mathcal{R}$. Both complete normal rows and the complete hard block are included. Write the original m event as $M = vW - X + uvY - uZ$. Its centered row is

$$R_{mc} = M - gvW = -\xi + uv\eta + (1-g)R_2 + huT. \tag{R5.2}$$

The equality uses $1 - g = a(1 - r - H)$ and $H = hB$. It vanishes at the actual contact. Put $L_m = (-1, uv, 0, 1-g, hu)^T$. For any positive real $K < m$, there is the exact matrix identity

$$G = G\big|_{m=K} + (m-K)L_m L_m^T + (m-K)g\, H_m, \tag{R5.3}$$

where the only nonzero entries of H_m are

$$(H_m)_{14} = (H_m)_{41} = -1, \quad (H_m)_{25} = (H_m)_{52} = v, \quad (H_m)_{44} = 2 - g, \quad (H_m)_{45} = (H_m)_{54} = hu. \tag{R5.4}$$

All other actual source coefficients, especially α/B^2, remain fixed in $G|_{m=K}$. This is a formal matrix substitution, not an assertion that a modified source is physical or satisfies contact. Equivalently, at the original event level,

$$mM^2 - (m-K)R_{mc}^2 = KM^2 + 2(m-K)g\, vWM - (m-K)g^2v^2W^2. \tag{R5.5}$$

The retained square contains every radial and normal jet of the excess. Equations R5.3–R5.4 prove boundedness of the remainder directly; no Hessian-subtraction premise is imported.

All infinitesimal or zero testing angles. If $\rho \in (0,1)$, apply the pure-source congruence replacing ξ by $\widehat{\xi} = \sqrt{n}\xi$. Every coefficient is bounded: nr is bounded, nr^2 infinitesimal, a, t bounded units, and A, C, D, m and α/B^2 bounded. At $u = v = 0$ the full residue consists of a unit $\widehat{\xi}$ pivot, a positive m_0 pivot on R_2, a positive $(\alpha/B^2)_0$ pivot on T, and the η, R_1 block

$$\begin{pmatrix} A_0 + C_0 + D_0 + 1/2 & -(C_0 + D_0 + 1/2)(1-\chi) + A_0\chi \\ -(C_0 + D_0 + 1/2)(1-\chi) + A_0\chi & (C_0 + D_0 + 1/2)(1-\chi)^2 + A_0\chi^2 \end{pmatrix}. \tag{R5.6}$$

Its determinant is $A_0(C_0 + D_0 + 1/2) > 0$.

If $\rho = 0$, replace R_2 by aR_2 as in WE4–WE5. The exact divisibility $(b_2 - \alpha/B^2)/a$ from WE3 holds for arbitrary h and r; its canceled denominator is a unit at $a = r = B = 0$ with h, l, t, j units. Hence every coefficient remains bounded, including all full normal entries. The R_2 pivot becomes $\mu_0 = 1/2$; R5.6 and the other pivots remain positive. No h/a bound is used.

If $\rho = 1$, retain $(m-K)R_{mc}^2$ using R5.3, for any fixed positive real K. Since a is a unit, all coefficients in the remainder are bounded after $\widehat{\xi} = \sqrt{n}\xi$. The constant $\widehat{\xi}, R_2$ coupling is infinitesimal, while its R_2 pivot has residue $K + 2(mg)_0 = K + 1/2$. All other zero-angle pivots are R5.6, 1, and $(\alpha/B^2)_0$. Thus the complete remainder has strict residue in every source case. The finite separated Gram lift WA10 applies for independent infinitesimal u, v, including zeros and either sign. Adding the retained exact square proves the complete local parent without any state restriction.

The finite remote determinant when $\rho < 1$. For bounded tests, normalize $\widehat{X} = \sqrt{n}X$ in the original parent. Its hard residue is $(1+U)(1+V)\widehat{X}^2$, and the remaining three-state form on (W, Y, Z) is

$$\begin{aligned} P = {} & m_0(vW + uvY - uZ)^2 + C_0(1+V)(uW - Y)^2 + A_0(1+U)(1+V)Y^2 + D_0(uW - Y - vZ)^2 \\ & + \tfrac{1}{2}\{(uW - Y - vZ)^2 + (1+U)(vY - Z)^2 + (1+V)UZ^2 - (1+U)(1+V)Z^2\}, \end{aligned} \tag{R5.7}$$

where here u, v denote the real testing residues and $U = u^2, V = v^2$. Set $\mathcal{D} = 8\chi(1-\rho)(1+\rho)^3\ell\delta > 0$. The exact determinant of its symmetric matrix is

$$\frac{(1+U)(1+V)}{\mathcal{D}}\left[(\chi+\tau)(\tau U - \rho V)^2 + (\chi + \tau - \rho\ell)\{\tau\delta U^2 V + [\chi + (1-\rho)\tau]UV^2\}\right]. \tag{R5.8}$$

For $U, V > 0$ it is strictly positive. For $U > 0, V = 0$ it is strictly positive; for $U = 0, V > 0$ it is strictly positive when $\rho > 0$. The restriction of R5.7 to $Z = 0$ is strictly positive in these cases: its signed core there is $\frac{1}{2}[(uW - Y)^2 + (1 + U)VY^2]$, and the A, C, m packets supply the stated two-state rank. This restriction and R5.8 prove strict positive definiteness of the entire three-state residue. All source coefficients are bounded, so WA10 around the real testing residues proves these complete remote charts.

Finite remote tests with the unbounded m opening. Let $\rho = 1$ and extract the exact square in R5.5 before normalizing $\widehat{X} = \sqrt{n}X$. Its remainder has bounded coefficients. At a real testing pair, its three-state residue is

$$B_{\mathrm{fin}} + KM_0^2 + 2\kappa vWM_0, \qquad M_0 = vW + uvY - uZ, \quad \kappa = 1/4, \tag{R5.9}$$

where B_{fin} is R5.7 with the m_0 packet omitted and A_0, C_0, D_0 evaluated at $\rho = 1$. For $U, V > 0$, restrict to $M_0 = 0$, so $W = uZ/v - uY$. Its two-state determinant equals $V^{-1} \lim_{\rho\uparrow 1} \det(P)/m_0$. Formula R5.8 gives

$$\frac{(1 + U)(1 + V)}{16\chi\ell\delta V} \Big[(\chi + \tau)(\tau U - V)^2 + (\chi + \tau - \ell)\{\tau\delta U^2 V + \chi UV^2\}\Big] > 0. \tag{R5.10}$$

At $Z = 0$ on that kernel, $W = -uY$ and the positive A, C terms, together with the nonnegative signed-core restriction, give a positive Y^2 coefficient. Hence $B_{\mathrm{fin}}|_{M_0=0}$ is positive definite. When $u = 0, v \neq 0$, the kernel is $W = 0$ and its form is $(A_0 + C_0 + 1/2)(1 + V)Y^2 + D_0(Y + vZ)^2 \succ 0$. When $v = 0, u \neq 0$, the kernel is $Z = 0$ and its form is $(C_0 + D_0 + 1/2)(uW - Y)^2 + A_0(1 + U)Y^2 \succ 0$. For each such real test pair, elementary real completion on the transverse coordinate M_0 now gives a sufficiently large positive real K for which R5.9 is strictly positive definite. This choice is legal: m is unbounded, so $m - K > 0$ for every real K; K never compares a source rate with a testing rate. Use that K in the exact full-parent identity R5.5 and apply WA10 to the bounded remainder.

The small-a first-axis chart. The only finite remote case left is $\rho = 0$, $u_0 = 0$, $v_0 \neq 0$. Use the five roots

$$S = aW, \quad R = uW, \quad \widehat{X} = \sqrt{n}X, \quad Y, \quad Z$$

. Write the original m packet as $\mu(vS - aX + auvY - auZ)^2$ and replace uW everywhere else by R. This is exactly the complete identity WE8 with unnormalized Y and $A(1 + u^2)(1 + v^2)Y^2$ retained. All coefficients are bounded. The residue has separate pivots $\mu_0 v_0^2$ on S and $1 + v_0^2$ on $\widehat{X}$. Its remaining (R, Y, Z) form is

$$C_0(1 + V)(R - Y)^2 + A_0(1 + V)Y^2 + D_0(R - Y - vZ)^2 + \tfrac{1}{2}\{(R - Y)^2 + VY^2 - 2vRZ\}. \tag{R5.11}$$

Its restriction to $Z = 0$ is positive definite and its determinant is

$$\frac{V^2(\chi + \tau)^2(1 + V)}{8\chi\ell\delta} > 0. \tag{R5.12}$$

This supplies the required strict full residue at the literal axis too.

Every reciprocal chart and literal infinity. For unbounded u put $p = 1/u$ and use the polynomial parent p^2F. If v is bounded, the $\widehat{X}$ residue is $1 + V$ and the other three-state residue is exactly

$$[C_0(1 + V) + D_0 + 1/2]W^2 + A_0(1 + V)Y^2 + (m_* + 1/2)(vY - Z)^2, \tag{R5.13}$$

which is strict for every real v. Here $m_* = m_0$ when bounded, and $m_* = K$ after centered extraction if g is infinitesimal. The additional centered correction in R5.5 has a factor p and vanishes in residue. This covers $p = 0$ literally.

For unbounded v, write $q = 1/v$ and use q^2F. If u is bounded, the $\widehat{X}$ pivot is $1+U$ and the other residue is

$$m_*(W+uY)^2 + C_0(uW-Y)^2 + (A_0+1/2)(1+U)Y^2 + D_0Z^2. \tag{R5.14}$$

For $\rho < 1$ this is strict unless $\rho = u_0 = 0$. For $\rho = 1$, the bounded correction is $2\kappa W(W+uY)$; its restriction to $W+uY=0$ is strict, and a sufficiently large real K again makes the entire residue strict. If $\rho = u_0 = 0$, use $S = aW, R = uW$, $\widehat{X}, Y, Z$ as above in the complete WE10 identity. Its residue is

$$\widehat{X}^2 + \mu_0 S^2 + C_0(R-Y)^2 + (A_0+1/2)Y^2 + D_0Z^2 \succ 0. \tag{R5.15}$$

Every coefficient of this entire polynomial chart is bounded.

Finally, if both angles are unbounded, use $p = 1/u, q = 1/v$ and p^2q^2F, with the five roots $\widehat{X}, W, Y, P = pZ, Q = qZ$. The exact signed-core part after homogeneity is one half of

$$(qW + X - pqY - P)^2 + (1+p^2)(Y-Q)^2 + (1+q^2)(p^2X^2 + 2XP - P^2). \tag{R5.16}$$

All other original packets are retained in their homogeneous forms; in particular the m packet is $(pW - pqX + Y - Q)^2$ and its centered row subtracts gpW. After $\widehat{X} = \sqrt{n}X$ and any needed centered extraction, the complete residue is

$$\widehat{X}^2 + C_0W^2 + A_0Y^2 + D_0P^2 + (m_*+1/2)(Y-Q)^2 \succ 0. \tag{R5.17}$$

Its coefficients are bounded; the centered correction again contains p and vanishes in residue. Both $p=0$ and $q=0$ are literal polynomial values. The roots P, Q need not be independent to use this positive Gram.

Join and tensor legality. The local chart, finite remote cases, the one small-a finite axis, each single reciprocal chart and the double reciprocal chart exhaust all independently chosen testing types. Every use of WA10 whitens a strict pure-source matrix at real testing residues and splits finitely many bounded-source times infinitesimal-testing coefficients into positive pure-source/pure-testing products. There is no mixed inverse or radical. The roots themselves use only pure-source radicals such as $\sqrt{n}$ and pure-testing monomials. Exact positive excess squares are restored in every chart. Away from literal endpoints, homogeneous restoration uses only pure-testing squares; at the endpoints the polynomial value is used. The real identities apply separately to real and imaginary state parts. All four original state coordinates, both normal directions, all hard couplings, arbitrary source openings and the unchanged physicality/total-scale/prior-exclusion intersection have therefore been retained. This is an all-tests admission of R5.

Chapter 147

R6 with bounded z/B is also an all-tests admission

Component `pd5fa361b4623`
Source: `R17/proofs/weak_gaps/r6_bounded_a_all_tests.tex`
SHA256: `d5fa361b462312d07d712a17fd01b2e88fa45b5d9cdb5e38bf6e4f650ef36a4c`
Inventory use: R6.1-R6.7

147.1 R6 with bounded z/B is also an all-tests admission

Theorem 147.1. *Retain every original R6 hypothesis and normalize $d = 1$: $B, z, y > 0$ infinitesimal; t, h, l, j, r positive bounded units; g infinitesimal; the unchanged physicality, contact, coefficient signs, total scale and prior exclusions. If $a = z/B$ is bounded, then the entire original parent is finite tensor SOS at all independent projective tests, including literal axes/infinities, either sign, and complex states. This admits the whole stated subdomain. The remaining R6 sources have a unbounded and $1 - r$ infinitesimal; no claim about that endpoint is made here.*

Write $e = 1 - r$. Since $1 - g = a(e - H)$, the added condition is equivalent to e being a positive unit; $a_0 = 1/e_0$. Denote the real residues by $e \in (0, 1)$, $h \in (0, 1)$, $l = 1 - h$, $t > l$, and $\delta = t - l$. The exact source formulas, with no rate comparison, give

$$\begin{aligned} n_0 = \frac{e^2}{2(1-e^2)}, \quad A_0 = \frac{t}{2h(1+e)}, \quad C_0 = \frac{h+t-el}{2l(1+e)}, \\ D_0 = \frac{1}{2(1+e)\delta}, \quad (\alpha/B^2)_0 = \frac{h+t-el}{2e(1+e)}, \quad \kappa = (mg)_0 = \frac{1}{2(1+e)}. \end{aligned} \tag{R6.1}$$

Every listed quantity is a positive unit; m is unbounded. In particular n cannot be literally zero on this admitted subdomain. The unbounded-a endpoint, where $n = 0$ may occur, is not erased.

Use the exact full-parent excess identity R5.2–R5.5 with a positive real cap K, retaining $(m - K)R_{mc}^2$. The remainder has bounded coefficients in the unchanged five WA roots, because a, t and all remaining event weights are bounded and mg bounded. At zero testing angles its η, R_1 block is R5.6 and its radial T pivot is $(\alpha/B^2)_0$. Its ξ, R_2 block is

$$\begin{pmatrix} n_0 + 1/2 + K & -K - \kappa \\ -K - \kappa & K + 2\kappa \end{pmatrix}, \quad \det = K(n_0 + 1/2) + 2\kappa(n_0 + 1/2) - \kappa^2. \tag{R6.2}$$

Since $n_0 + 1/2 = \kappa/(1 - e)$, its determinant is positive for every $K > 0$. Thus the complete local remainder has strict residue, and WA10 proves all infinitesimal or zero tests, keeping all hard and normal terms.

For a finite remote pair, use the original four states. After centered extraction the bounded remainder is, in residue,

$$B_{\text{fin}} + KM^2 + 2\kappa vWM, \qquad M = vW - X + uvY - uZ, \tag{R6.3}$$

where B_{fin} is the complete original parent at the source residues R6.1, with m and α omitted. The restriction to $M = 0$ is obtained by the exact substitution $X = vW + uvY - uZ$. On (W, Y, Z) its determinant is

$$\frac{(1+U)^2(1+V)^2}{\mathcal{D}}\left[(h+t)(etU-V)^2 + (h+t-el)\{t(h+t-e)U^2V + hUV^2\}\right], \tag{R6.4}$$

where $\mathcal{D} = 8h(1-e)(1+e)^3l\delta > 0$. Every coefficient outside the displayed square is strictly positive: $h + t > 1 > e$ follows from $t > l$. Hence the determinant is positive whenever $(U, V) \neq (0, 0)$, including both axes. The restriction to $Z = 0$ is positive definite: the original signed core there is the sum of the three squares $[J^2 + (1+U)VY^2 + (1+V)X^2]/2$, and the positive A, C, n packets force $Y = W = 0$ on $M = Z = 0$ at every such remote pair. Thus $B_{\mathrm{fin}}|_{M=0}$ is positive definite. A sufficiently large positive real K makes R6.3 strictly positive definite on the full four-state space. Use this K in the exact source identity and apply WA10 around the real testing residues. All coefficients remain bounded; there is no mixed division or source/test comparison.

For unbounded u and bounded v, use the full polynomial chart p^2F, $p = 1/u$. Its excess correction contains p and disappears in residue. The centered m row has residue $vY - Z$. On its kernel the complete finite remainder on (X, W, Y) is

$$n_0(1+V)X^2 + C_0(1+V)W^2 + A_0(1+V)Y^2 + (D_0 + 1/2)(W + vX)^2 \succ 0. \tag{R6.5}$$

A positive cap gives a strict full residue. For unbounded v and bounded u, use q^2F, $q = 1/v$. The m row is $W + uY$ and the bounded correction is $2\kappa W(W + uY)$. On its kernel, the complete residue is

$$(n_0 + 1/2)(1+U)X^2 + C_0(1+U)^2Y^2 + (A_0 + 1/2)(1+U)Y^2 + D_0(uX - Z)^2 \succ 0. \tag{R6.6}$$

Again a sufficiently large real K gives a strict full residue. These assertions include every finite real value of the other angle, with either sign or zero.

For both angles unbounded use the exact double homogeneous parent and core R5.16, now in the five roots $(X, W, Y, P = pZ, Q = qZ)$, without normalizing X. Its complete residue after centered extraction is

$$(n_0 + 1/2)X^2 + C_0W^2 + A_0Y^2 + D_0(X - P)^2 + (K + 1/2)(Y - Q)^2 \succ 0. \tag{R6.7}$$

The X, P block has determinant $D_0(n_0{+}1/2) > 0$ and the Y, Q block has determinant $A_0(K{+}1/2) > 0$. The excess correction has a factor p and disappears in residue. All homogeneous coefficients are bounded. Literal $p = 0, q = 0$ are the polynomial chart values, not limiting divisions.

These cases exhaust all tests on the stated source subdomain. The separated finite lift, exact excess restoration, pure-testing homogeneous restoration and real/imaginary split are exactly those established in the R5 join. They retain every original normal and hard block, all signs, all independent rates and the original source/prior-exclusion intersection.

Chapter 148

The bounded-z/B subdomain of R7 is admitted for all tests

Component p9a223c96c696
Source: R17/proofs/weak_gaps/r7_bounded_a_all_tests.tex
SHA256: 9a223c96c696aeb5849d21cb9dc39340385df93cbcbe4609bc20f249022367d3
Inventory use: R7.1-R7.12

148.1 The bounded-z/B subdomain of R7 is admitted for all tests

Theorem 148.1. *Normalize $d = 1$ and retain every original R7 condition: $B, z, y > 0$ infinitesimal; $t = y/B$ a positive bounded unit; $h = H/B$ and $g = G/B$ infinitesimal; the exclusion of simultaneously infinitesimal r, j; original physicality, stationary contact, coefficient signs, strict gaps, total-scale intersection and prior exclusions. If $a = z/B$ is bounded, the entire original parent is finite tensor SOS at every independent projective testing pair. Both signs, literal zero/infinite angles and complex states are included. The unbounded-a R7 endpoint remains open.*

Complete source entry. Write $e = 1 - r$ and $l = 1 - h$. Since $1 - g = a(e - H)$, bounded a is equivalent on this source shape to e being a positive unit; $a_0 = 1/e_0$. We use the letters $e \in (0, 1]$ and $t \geq 1$ for their real residues. The strict gap $j = 1 + zr - l/t$ gives $j_0 = 1 - 1/t$. The ordered R7 exclusion says that $e = 1$ and $t = 1$ cannot hold together. The original source formulas give jointly

$$C_0 = \frac{t - e}{2(1 + e)} > 0, \quad (Ah)_0 = \frac{t}{2(1 + e)} > 0, \quad (\alpha/B^2)_0 = \frac{t - e}{2e(1 + e)} > 0, \quad \kappa = (mg)_0 = \frac{1}{2(1 + e)}. \tag{R7.1}$$

Thus A, m are unbounded, C and α/B^2 are positive units, and α is infinitesimal. If $e < 1$, then

$$n_0 = \frac{e^2}{2(1 - e^2)} > 0;$$

if $e = 1$, r is infinitesimal and n is unbounded, with $(nr)_0 = 1/4$. If $t > 1$, then $D_0 = 1/[2(1+e)(t-1)]$ is a positive unit. If $t = 1$, then j is infinitesimal and D unbounded with $(Dj)_0 = \kappa$; in this case $e < 1$ by the retained exclusion. These are exhaustive source alternatives, with no rate restriction. They also show that literal $n = 0$ is impossible on this admitted subdomain and that total scale is positive infinitesimal. The $n = 0$ boundary on the unbounded-a complement is not removed by this statement.

A complete D-event gauge, including its exact zero syzygy. We use the full WA parent, now in the roots

$$\mathcal{R} = (\xi, \eta, R_1, R_2', T)^T, \quad \xi = X + ruZ, \quad \eta = Y - huW,$$
$$R_1 = uW - vZ/t, \quad R_2' = avW - uZ, \quad T = uvW - BZ.$$

Let G be the exact full-parent matrix obtained from WA8 by leaving η unnormalized and scaling its fourth root by a. The following gauge is the exact D identity developed in the parallel R8 work; it is valid before any shape specialization. Set

$$\begin{aligned} d_D &= (uv, -1, l, ruv, -arv)^T, \\ S_D &= (0, 0, (aB - U)v, -(B - V/t)u, U - aV/t)^T, \\ L_D &= (-2rtu, 0, 0, -r^2tu, ar^2t)^T. \end{aligned}$$

Then $S_D^T \mathcal{R} = 0$ identically, and

$$d_D^T \mathcal{R} = J + jvZ =: R_{Dc}, \qquad J = uW + uvX - Y - vZ. \tag{R7.2}$$

In particular this is the actual contact-zero D row, not an uncentered positive packet. The exact matrix identity is

$$\partial_D G = d_D d_D^T + \tfrac{1}{2}(S_D L_D^T + L_D S_D^T) + jH_D, \tag{R7.3}$$

where the only nonzero entries of H_D are

$$(H_D)_{13} = tuv, \quad (H_D)_{15} = -tu, \quad (H_D)_{34} = rtuv, \quad (H_D)_{35} = -artv$$

and their symmetric partners. Consequently the full original parent is

$$F/4 = DR_{Dc}^2 + \mathcal{R}^T (G|_{D=0} + DjH_D)\mathcal{R}. \tag{R7.4}$$

The notation $G|_{D=0}$ keeps all other actual coefficients, including α/B^2, unchanged; no modified physical source is asserted. All full first normals and hard entries are included in R7.3. This gauge has been independently checked here against the entire WA matrix, its zero syzygy and the original four-state D event.

One strict local Gram for all source alternatives. First apply R7.4, and then retain the centered m excess from R5.2–R5.4, with positive real cap K. The m identity is unaffected by the D gauge: H_D and its syzygy are independent of m and the D row is independent of m. The remaining full Gram has bounded coefficients after

$$\widehat{\eta} = \sqrt{A}\,\eta,$$

and, if r is infinitesimal, also $\widehat{\xi} = \sqrt{n}\,\xi$. Indeed Ah, nr, mg, Dj are bounded, a, t, l^{-1} are bounded units, Ah^2 is infinitesimal, and every large m or D coefficient has been removed in its complete contact-zero square. No ratio between A, n, m, D is used. This bounds the entire scalar block, both full normal rows and the hard block, not only the graph.

At $u = v = 0$, all entries of H_D vanish. The $\widehat{\eta}$ pivot is 1, its couplings are infinitesimal, the R_1 pivot is $C_0 + 1/2 > 0$, and the T pivot is $(\alpha/B^2)_0 > 0$. If r is a unit, the remaining ξ, R_2' block is the pure-source rescaling by $\mathrm{diag}(1, a_0^{-1})$ of R6.2. Its determinant is positive because $n_0 + 1/2 = \kappa/(1-e)$. If r is infinitesimal, normalization by $\sqrt{n}$ instead gives separate positive pivots 1 and $(K + 2\kappa)/a_0^2$; the cross coefficient is infinitesimal. Thus the complete local remainder has strict residue for every source alternative. WA10 gives a finite separated tensor SOS for every independent infinitesimal or zero u, v. Restore both retained exact squares. This proves the full local parent, including its complete normal/radial jets, without transferring physicality to a capped source.

Remote entry and a finite-dimensional cap lemma. For remote charts use the original states and normalize $\widehat{Y} = \sqrt{A}Y$. If r is infinitesimal, normalize also $\widehat{X} = \sqrt{n}X$. First retain the m excess. If D is unbounded, retain a D excess too, using the exact full-event identity

$$DJ^2 - (D - K_D)(J + jvZ)^2 = K_D J^2 - 2(D - K_D)jvZJ - (D - K_D)j^2 V Z^2. \tag{R7.5}$$

All coefficients in the remaining full parent are bounded. The hard normalized state pivots are positive and decouple in residue; all other coefficients are bounded and divided by $\sqrt{A}$ or $\sqrt{n}$ in couplings to these pivots. Thus the soft residue has $Y = 0$, and also $X = 0$ when r is infinitesimal.

We repeatedly use the following elementary real fact. If a real quadratic form B is positive definite on the common kernel of real linear rows $L_1, \dots, L_k$, then $B + K \sum L_i^2$ is positive definite for sufficiently large positive real K, after any fixed bounded cross terms vanishing on that common kernel have been included in B. This follows by choosing real coordinates on the kernel and its orthogonal complement and taking the Schur complement. Here the relevant original event coefficients are unbounded, so every such real cap leaves positive excesses. The choice uses only real testing residues, never a comparison of source and testing rates. The positive hard pivots can then be included in the same strict full-parent Gram.

Finite remote tests when j is a unit. Let $U = u^2, V = v^2$ refer now to the real testing residues, not both zero. After $Y = 0$, the centered m row has residue $M = vW - X - uZ$. Write B_{fin} for the entire finite parent with its m packet removed; the bounded cap correction is $2\kappa vWM$ and vanishes on $M = 0$.

If r is a unit, retain X and restrict by $X = vW - uZ$. The two-state form on (W, Z) has determinant

$$\frac{(1+U)(1+V)}{4(1-e)(1+e)^2(t-1)}\{(etU - V)^2 + (t-e)^2U^2V\} > 0. \tag{R7.6}$$

Its $Z = 0$ restriction is positive: the positive n, C packets and the nonnegative original signed core at $Z = 0$ force $W = 0$ at every remote pair. Hence it is positive definite, and the cap lemma gives a strict full residue.

If r is infinitesimal, then $e = 1$, $t > 1$ and X is another hard state. For $v \neq 0$, restrict by $W = uZ/v$. The remaining scalar is

$$\frac{Z^2}{4(t-1)V}\{(tU - V)^2 + (t-1)^2U^2V\} > 0. \tag{R7.7}$$

For $v = 0, u \neq 0$, $M = 0$ forces $Z = 0$ and the scalar on W is $(C_0 + D_0 + 1/2)UW^2 > 0$. These include the literal first axis and both angle signs. Again the cap lemma supplies strict residue for the complete bounded remainder.

Finite remote tests with both m and D unbounded. Here $t = 1$, $e < 1$, $r = 1 - e$ is a unit and n is finite. With $Y = 0$ the two centered rows have residues

$$M = vW - X - uZ, \qquad J = uW + uvX - vZ.$$

They have rank two at every finite remote pair. Their common kernel is spanned in (W, X, Z) by

$$(v(1+U),\ V - U,\ u(1+V)). \tag{R7.8}$$

On this vector, the complete finite parent with both large packets removed has value

$$\frac{(1+U)(1+V)}{2(1-e)(1+e)}\{(eU - V)^2 + (1-e)^2U^2V\} > 0. \tag{R7.9}$$

The bounded residue corrections are $2\kappa vWM$ and $-2\kappa vZJ$; both vanish on that common kernel. The cap lemma therefore gives positive real caps for a strict three-state soft residue; the hard $\widehat{Y}$ pivot completes the full parent. The higher source terms proportional to mg^2 and Dj^2 have infinitesimal coefficients and are retained in the bounded remainder, not discarded from the exact identity.

Every reciprocal chart. For unbounded u, use $p = 1/u$ and the entire polynomial parent p^2F. The m row on $Y = 0$ is $-Z$. If D is bounded, its kernel $Z = 0$ has complete residue

$$n_0(1+V)X^2 + C_0(1+V)W^2 + (D_0 + 1/2)(W + vX)^2 \succ 0, \tag{R7.10}$$

with X omitted and its normalized hard pivot retained if n is unbounded. If D is unbounded, its additional row is $W + vX$; on the common kernel the finite form is $(1+V)(n_0 + C_0V)X^2 > 0$. In this case $n_0 > 0$. The m and D bounded cross corrections contain p and vanish in residue. The cap lemma applies for every bounded real v, including zero.

For unbounded v, use $q = 1/v$ and q^2F. The m row on $Y = 0$ is W. If D is bounded, the complete residue on $W = 0$ is

$$(n_0 + 1/2)(1+U)X^2 + D_0(uX - Z)^2 \succ 0, \tag{R7.11}$$

again with X replaced by its hard normalized pivot when n is unbounded. If D is unbounded, the additional row is $uX - Z$ and the common-kernel form is $(n_0 + 1/2)(1+U)X^2 > 0$. The bounded corrections are proportional to W^2 and to $Z(uX - Z)$, so vanish on the relevant common kernels. This covers every bounded real u, including zero.

For both angles unbounded, use p^2q^2F and the exact core R5.16 in $(W, X, Y, P = pZ, Q = qZ)$, with Y normalized and X normalized if needed. On the soft states $Y = 0$ the m row is $-Q$. For bounded D, the finite residue on $Q = 0$ is

$$(n_0 + 1/2)X^2 + C_0W^2 + D_0(X - P)^2 \succ 0, \tag{R7.12}$$

with the same convention for a hard X. If D is unbounded, its row is $X - P$ and the common-kernel form is $(n_0 + 1/2)X^2 + C_0W^2 > 0$. The surviving D correction is proportional to $P(X - P)$, while the m correction contains p; they vanish on the common kernel. Large real caps give a strict full residue. These identities are polynomial in p, q and include literal $p = 0, q = 0$.

Exact join and scope. The local, finite remote and three reciprocal configurations exhaust every independent projective testing pair on the stated source domain. Each bounded remainder has strict residue and thus the finite separated lift WA10, while every extracted excess is restored as its exact positive contact-zero square. The proof uses no mixed inverse or radical; all normalizations and homogeneous restoration factors are purely source or purely testing. Every full normal, hard coupling, retained equality and source opening remains in the identities. Applying the real proof to the real and imaginary state parts completes the complex assertion. This is an all-tests admission of bounded-a R7, not an admission of its still unbounded-a endpoint or of the omitted simultaneous small-r, small-j intersection.

Chapter 149

The remaining small-r local chart of weak-C2 R8

Component `p482fcad1a2e2`
Source: `R17/proofs/weak_r8_small_r/r8_small_r_local.tex`
SHA256: `482fcad1a2e28b3fa16a1961ed72b3d09bb7cfd2dc5f1a221eb497694576527e`
Inventory use: RD1-RD11

149.1 The remaining small-r local chart of weak-C2 R8

Theorem 149.1. *Retain the original physical stationary weak-C2 source, all strict gaps, nonnegative event coefficients, the original positive-infinitesimal total-scale intersection, and every inherited residual exclusion. Suppose the source is in R8 and r is infinitesimal. Thus $t = y/B$ and $g = G/B$ are positive bounded units, while $B, h = H/B, j, r$ are positive infinitesimals. Then the entire original four-state parent is a finite tensor SOS for every independent pair of infinitesimal or zero testing half-angles u, v, with either sign and all real or complex states. No source opening is compared with another opening or with a testing rate.*

This is the missing local component of the R8 join. Its remote tests and its r-unit local chart are supplied separately by the R17 joint-D argument. The proof below does not rely on an unproved extension of the RC normalization across $r = 0$.

Source entry and the three independent hard scales. Write $a = z/B$, $l = 1 - h$, $x = aB$, and

$$j = 1 + xr - l/t, \qquad g = 1 + a(hB + r - 1).$$

The R8 and small-r hypotheses imply that a is bounded with residue $\rho \in [0, 1)$: $g_0 = 1 - \rho > 0$. Also $t_0 = 1$. The unchanged exact source formulas WE1, jointly at $B = h = r = 0, t = 1$, give

$$(nr)_0 = (Ah)_0 = (Dj)_0 = \frac{1}{2(1+\rho)}, \qquad \left(\frac{m}{a^2}\right)_0 = \frac{1}{2(1-\rho^2)}, \quad C_0 = \left(\frac{\alpha}{B^2}\right)_0 = \frac{1-\rho}{2(1+\rho)} > 0. \tag{RD1}$$

These are joint regular reductions. After replacing n, A, D by nr, Ah, Dj, all source denominators are products of units $t, g, 1-h, 1-hB, \Delta$ with $\Delta_0 = 1 + \rho$. Thus n, A, D are all positive unbounded, with completely unrestricted relative rates, while $nr, Ah, Dj, \mu = m/a^2, C, \alpha_* = \alpha/B^2$ are positive bounded units. In particular $n = 0$ is impossible on this chart; no inverse of a vanishing event coefficient is being used.

A general exact centered-D gauge of the complete WA matrix. Use the unnormalized complete WA3–WA8 identity, with the WE4 source rescaling of the second scalar root:

$$\mathcal{R} = (\xi, \eta, R_1, R_2', T)^T, \quad \xi = X + ruZ, \quad \eta = Y - huW,$$

$$R_1 = uW - t^{-1}vZ, \quad R_2' = avW - uZ, \quad T = uvW - BZ, \qquad F/4 = \mathcal{R}^T G \mathcal{R}. \tag{RD2}$$

Here G contains the complete hard block, both complete first-normal rows, and the complete scalar block. No packet or normal is omitted. All its entries are affine in the formal event coefficient D.

For $U = u^2$, $V = v^2$, define three five-component columns

$$\begin{aligned} d &= (uv, -1, l, ruv, -arv)^T, \\ s &= (0, 0, (x - U)v, -(B - V/t)u, U - aV/t)^T, \\ k &= (-2rtu, 0, 0, -r^2tu, ar^2t)^T. \end{aligned}$$

The physical root relation and the centered D event are exactly

$$s^T \mathcal{R} = 0, \qquad L_D := d^T \mathcal{R} = uW + uvX - Y - vZ + jvZ. \tag{RD3}$$

The following is a matrix identity, valid before any residue operation:

$$\partial_D G - dd^T - \frac{sk^T + ks^T}{2} = jH_D, \tag{RD4}$$

where H_D is symmetric and its only nonzero entries are

$$(H_D)_{13} = tuv, \quad (H_D)_{15} = -tu, \quad (H_D)_{34} = rtuv, \quad (H_D)_{35} = -artv. \tag{RD5}$$

Indices in RD5 refer to the displayed order in RD2 and start at one. This explicitly resolves the failure of the ungauged matrix $\partial_D G - dd^T$ to be divisible by j.

Let $G^{(0)} = G|_{D=0}$ denote a formal matrix substitution, with every other source coefficient kept at its actual value. This is not a physical modified source and no physicality theorem is applied to it. Because the term involving s vanishes exactly on the physical roots, RD4 gives the entire-parent identity

$$F/4 = DL_D^2 + \mathcal{R}^T \{G^{(0)} + DjH_D\} \mathcal{R}. \tag{RD6}$$

The first-normal and hard-block entries in RD4 are part of this identity, not merely its scalar restriction. No inverse of u, v, j, or a state coordinate was used in deriving RD3–RD6.

The bounded D-free matrix through the a opening. Normalize only the first two roots in $G^{(0)}$ initially, by $\widehat{\xi} = \sqrt{n}\,\xi$ and $\widehat{\eta} = \sqrt{A}\,\eta$. The resulting matrix, in $(\widehat{\xi}, \widehat{\eta}, R_1, R_2', T)$, is a finite polynomial in u, v with bounded pure-source coefficients. Here is the complete coefficient check needed for this assertion.

The hard entries after $D = 0$ are

$$\begin{aligned} h_{xx}/n &= (1 + U)(1 + V) + [m + UV/2 + (1 + V)/2]/n, \\ h_{xy}/\sqrt{nA} &= -uv(m + 1/2)/\sqrt{nA}, \\ h_{yy}/A &= (1 + U)(1 + V) + [C(1 + V) + 1/2 + mUV + (\alpha + 1/2)(1 + U)V]/A. \end{aligned}$$

Every added coefficient is bounded and has zero residue. In the first normal, the complete WA6 parameters now have the forms

$$p_0 = t(nr + r/2), \quad p_1 = t(nr + r/2), \quad q_1/a = nr + Bp_1,$$

$$f_x = -mh + l/2 - p_0 - q_1, \quad (-m + q_1 U)/a = -a\mu + (nr + Bp_1)U.$$

They are bounded before multiplication by $1/\sqrt{n}$. In the complete second normal, all WA7 coefficients involve only the bounded products Ah, m, C, α, r, h, and

$$q_y/a = -k_z - (Ah)/t - Bk_{uv}$$

is bounded. Multiplication by $1/\sqrt{A}$ therefore bounds that entire normal row as well. No ratio n/A, A/n, h/r, or r/h appears.

For the scalar block use the exact WE5 expressions with $D = 0$. All occurrences of n are through $nr^2 = (nr)r$, and all occurrences of A are through $Ah^2 = (Ah)h$. The only possible new a denominator is

$$Q_* = (b_2 - \alpha_*)/a, \quad b_2 = 2a^2\mu h + C(1-h)^2 + (Ah)h + (B^2\alpha_* + 1/2)h^2. \tag{RD7}$$

The exact WE1 expressions for b_2 and α_* involve no D and no j denominator. At $a = 0$ their difference vanishes identically for arbitrary B, h, r, t. Its cleared numerator is consequently divisible by a; the canceled denominator at $B = h = r = 0, t = 1$ is $2(a-1)(a+1)$, a unit for $0 \le \rho < 1$. The divided numerator vanishes at that joint corner for every a. Hence Q_* is bounded (in fact infinitesimal), independently of a, h, r, j rates. This proves the bound on

$$f_s/a = \tfrac{1}{2}[Q_* - a\,nr^2 - aBt(n + 1/2)r^2]$$

and on the whole remaining scalar block. In particular the old assumption that j was a unit is unnecessary after the exact D channel has been removed by RD6.

Simultaneous normalization of all three hard scales. Use the final five roots

$$\mathcal{N} = (\widehat{\xi}, \widehat{\eta}, \widehat{D}, R_2', T)^T, \qquad \widehat{D} = \sqrt{D}\, L_D. \tag{RD8}$$

Solving only the source-unit coefficient $l = 1 - h$ in RD3 gives

$$R_1 = \frac{1}{l}\left\{\frac{\widehat{D}}{\sqrt{D}} + \frac{\widehat{\eta}}{\sqrt{A}} - \frac{uv\widehat{\xi}}{\sqrt{n}} - ruvR_2' + arvT\right\}. \tag{RD9}$$

Every coefficient in RD9 is bounded and separated. In particular, independent small r, h, j cause no ratio of their square roots. Let $P(u, v)$ be the explicit matrix mapping $\mathcal{N}$ to $\mathcal{R}$ according to RD9 and $\xi = \widehat{\xi}/\sqrt{n}$, $\eta = \widehat{\eta}/\sqrt{A}$. Then exactly

$$F/4 = \mathcal{N}^T \widehat{G} \mathcal{N}, \qquad \widehat{G} = E_{33} + P^T\{G^{(0)} + DjH_D\}P. \tag{RD10}$$

The preceding bounds and RD9 prove that every coefficient of this finite testing-polynomial matrix is bounded pure source. The factor Dj is a unit, while every entry of H_D has positive testing degree. Thus its contribution to the residue at $u = v = 0$ is zero.

The normalized n and A hard diagonals have residue one. Since $1/\sqrt{n}$, $1/\sqrt{A}$, $1/\sqrt{D}$, r, h are infinitesimal, the entire R_1 row of $P(0, 0)$ has zero residue. The remaining source scalar diagonals have residues μ_0 and $\alpha_{*,0} = C_0$. All cross entries have zero residue. Consequently

$$\overline{\widehat{G}} = \operatorname{diag}\left(1, 1, 1, \frac{1}{2(1-\rho^2)}, \frac{1-\rho}{2(1+\rho)}\right) \succ 0. \tag{RD11}$$

This is the full five-root Gram residue, not a Hessian or state slice.

The positive pure-source matrix $\widehat{G}(0, 0)$ has bounded inverse. Whiten it purely in the source field and apply the finite separated coefficient construction WA10 to the finitely many nonconstant testing monomials. Their values are infinitesimal for arbitrary independent infinitesimal u, v. This produces a finite SOS in $K_c \otimes_{\mathbb{R}} K_t$ without any mixed inverse or mixed radical. The argument includes either zero test and both signs because all identities are polynomial in u, v. Applying it to real and imaginary state parts proves the complex-state statement.

Verification boundary. The focused checker imports only the established complete six-contact WA identity and WA10. It verifies the new entire D-matrix gauge, the physical syzygy, the exact original-event centering, the entire parent equality RD6, all fifteen normalized coefficient denominators at the joint opening, the full residue RD11, and the exact source quotient through $a = 0$. Eighteen new exact checks pass. No numerical positivity test or unproved scalar-to-parent lift is used.

Chapter 150

R8: a joint divergent-D certificate, a paired-D thin lift, and the remote join

Component `pe07041e033da`
Source: `R17/proofs/weak_scale/joint_D_R8_closure.tex`
SHA256: `e07041e033da07344e496a0c5af3070bc7e6263375a7929c9b1aa32bb43b806b`
Inventory use: JD1-JD24 and projective join

150.1 R8: a joint divergent-D certificate, a paired-D thin lift, and the remote join

Scope and conventions. All identities use $F/4$ and the unchanged physical C2 parent, with $\lambda_{\text{event}} = \beta = 0$, $d = 1$, and $n \geq 0$. Every original stationary-contact, nonnegative-coefficient, strict-gap, total-scale, and prior-admission-complement intersection is retained. The two real closed source/testing fields are independent. The symbol λ below is the positive circuit coefficient, not an event coefficient.

R8 means $B, z, y, h, j > 0$ infinitesimal, $t = y/B$ and $g = G/B$ positive bounded units, $h = H/B$, $a = z/B$, $r \in (0, 1)$, and $e = 1 - r$. Put $x = z = aB$, $l = 1 - h$. The exact gap is

$$j = 1 + xr - l/t. \tag{JD1}$$

In particular $t_0 = 1$. This section proves:

1. every independent test and state on the whole R8 subdomain where r is a positive bounded unit;

2. the entire remote testing complement of both angles infinitesimal on all of R8, including r infinitesimal.

The simultaneous r, h, j-infinitesimal local domain is proved by the separate joint-n, A, D lemma in `weak_r8_small_r/r8_small_r_local.tex`; it is not inferred from the present r-unit proof. Its union with the two assertions above closes the whole ordered R8 shape. No R1 or R2 admission is asserted here.

The inherited exact inputs are R15 RC1–RC13 and its source appendix, R16 R10.9–R10.12, TL1–TL10 and the two complete-support checks, the CA/CS/SE bounded contact conductors, and the critical-wall polynomial identities SEO10–SEO20. They are used as explicit identities with the bounds verified below. In particular no physical C1 theorem is invoked at a negative formal beta.

A single contact-zero row removes every naked D pole. Use the homogeneous R15/R16 roots

$$E = tuW - vZ, \quad N_y = (Y - huW)/\sqrt{h}, \quad \xi = X + ruZ, \quad R_D = v(u^2 - x)Z, \quad F_v = (v^2 - y)Z.$$

The other roots Q, N, T, R, J and source coefficients are exactly those of R10.9. Define

$$L_D = \frac{l}{t}E - rR_D - \sqrt{h}N_y + uv\xi, \qquad M_D = \sqrt{D}\,L_D. \tag{JD2}$$

The original event row satisfies the exact homogeneous identity

$$uW + uvX - Y - vZ = L_D - jvZ. \tag{JD3}$$

Thus L_D vanishes at the actual source contact, including its full state normal. This is a legal separated root at every testing axis.

Let $\mathscr{P}$ be the right side of R10.9 with the two exact positive squares N^2, T^2 omitted, and let $\mathscr{P}_0$ denote its formal coefficient specialization $D = 0$. The other source letters are held fixed in this notation; it does not change the physical source. Direct collection of the complete D channel gives

$$\boxed{\mathscr{P} = DL_D^2 + \mathscr{P}_0 - 2Dj\,uF_v\xi.} \tag{JD4}$$

Here is the full coefficient calculation. The hard contribution is $D(lE/t - \sqrt{h}N_y + uv\xi)^2$; the remaining D contributions are $Dr^2R_D^2 - 2Dr(lE/t - \sqrt{h}N_y + uv\xi)R_D$ and $2D\{l/t - 1 - rx\}uF_v\xi$. Their sum is exactly JD4 by JD1. This accounts for every hard and first-normal term, rather than just the angular graph or its Hessian.

Use the polynomial inverse

$$E = \frac{t}{l}\{D^{-1/2}M_D + rR_D + \sqrt{h}N_y - uv\xi\}. \tag{JD5}$$

On R8, t/l is a unit, $D^{-1/2}$ is infinitesimal, and Dj is bounded. Every coefficient of the new remainder is bounded whenever the remaining source arrays of R10.9 are bounded. In particular no ratio h/j, x/j, or testing angle divided by a source opening occurs. The full divergent rank-one block has been normalized jointly.

The exact residue after the joint normalization. Put $\sigma = t\lambda x/\sqrt{q}$ and $\mu = m - a^2q$ as in R10.9. At the R8 source/testing corner, JD4–JD5 have the complete residue

$$M_D^2 + r_0^2(C_0 + n_0)R_D^2 + (Ah)_0N_y^2 + (n_0 + \tfrac{1}{2} - \mu_0)\xi^2 + (c_F)_0F_v^2 + \lambda_0J^2 + R^2 + Q^2 - 2\sigma_0Q\xi \tag{JD6}$$

This formula applies when r, n are bounded and r a unit. Its derivation can retain $n, C, Ah, \mu, c_F, \lambda, \sigma$ as free source letters: specialize $u = v = \sqrt{h} = D^{-1/2} = x = \alpha = 0$ and $t = 1$ in JD4–JD5. In particular the E, R_D, N_y terms become $r^2(C + n)R_D^2 + AhN_y^2$, with no missing D contribution. The separate N^2, T^2 remain intact. The focused checker reconstructs the original hard polynomial and verifies both JD4 and this full residue.

Bounded a, unit r, and unit e. Positivity of $g = 1 - ae + hx$ makes a bounded whenever e is a unit. The unchanged source formulas, with $B = h = 0$ and $t = 1$ only after clearing unit denominators, give

$$\begin{aligned} n_0 &= \frac{e_0}{2r_0(1 + a_0)}, \quad (m/a^2)_0 = \frac{e_0}{2g_0(1 + a_0)}, \\ C_0 &= \frac{1 + a_0(2r_0 - 1)}{2(1 + a_0)}, \quad (Ah)_0 = (Dj)_0 = \frac{1 + a_0r_0}{2(1 + a_0)}, \\ \lambda_0 &= \frac{r_0(1 + a_0r_0)}{2(1 + a_0)}, \quad (c_F)_0 = \frac{(1 + a_0r_0)g_0}{2(1 + a_0)}. \end{aligned} \tag{JD7}$$

All displayed pivots are positive units. Indeed $1 + a_0(2r_0 - 1) = g_0 + a_0r_0 > 0$. The source cancellation defining $\lambda = (m - a^2q)/(ta^2B)$ is regular also at $a = 0$; no division by an infinitesimal a survives its canceled formula. Here q is a unit, $\mu = ta^2B\lambda$ is infinitesimal and σ is infinitesimal. Therefore JD6 is strictly positive. All omitted entries have zero residue by the finite polynomial JD4–JD5 and bounded source arrays. The bounded separated-Gram lemma gives the complete local parent for every independent ratio u/v, every state, and both literal axes.

Bounded a, infinitesimal e, including $n = 0$. Now $r_0 = 1$, $C_0 = (Ah)_0 = 1/2$, n, m, μ are infinitesimal, $\lambda_0 = (c_F)_0 = 1/2$, and

$$(\alpha/B^2)_0 = (1 + a_0)/2. \tag{JD8}$$

The physical inequality OP2 supplies the crucial source lower bound, without dividing by n:

$$\frac{e}{B} \geq \frac{(1+x)(h+t)(1+ar)}{P_n/B}, \qquad P_n/B \sim 1. \tag{JD9}$$

It remains valid when e/B is unbounded and at literal $n = 0$. Since $q \geq \alpha + e^2/2$, the Q, ξ Schur pivot satisfies

$$\begin{aligned} \operatorname{st}\left(\tfrac{1}{2} - \mu - \frac{t^2\lambda^2 a^2 B^2}{q}\right) &\geq \frac{1}{2} - \frac{a_0^2}{2(1+a_0)(2+a_0)} \\ &= \frac{2 + 3a_0}{2(1+a_0)(2+a_0)} > 0. \end{aligned} \tag{JD10}$$

This is a bound entirely in the source field. The entry σ is bounded by JD8, so the whole block is bounded and strict. The remaining entries of JD6 are positive units. This proves all local tests and all states on this source domain, including $n = 0$.

Unbounded a: the thick layer. Put $s = 1/a$, $w = ae$, $b = a^2 B$. The exact R10 source identity still reads

$$n = s^2 \frac{wQ_* - bK_*}{2r\Delta_*}, \qquad Q_* \sim 1,\ K_* \sim t,\ \Delta_* \sim 1. \tag{JD11}$$

Thus b/w is bounded, b is bounded, $g_0 = 1 - w_0 > 0$, and

$$\begin{gathered} (m/w)_0 = \frac{1}{2(1-w_0)}, \quad C_0 = (Ah)_0 = 1/2, \quad (Dj)_0 = 1/2, \\ \lambda_0 = 1/2, \quad (c_F)_0 = (1 - w_0 + b_0)/2 > 0. \end{gathered} \tag{JD12}$$

The source formulas for these quantities have no j denominator; only D itself is unbounded, and JD4 has removed that pole. For $\rho = n/e^2$ positive noninfinitesimal, retain the exact R10.10 identities and R10.11 Schur pivot:

$$\begin{gathered} \bar{q} = a^2 q = w^2\{\rho r^2 + H\} + a^2\alpha, \quad H = m + \tfrac{1}{2}, \quad \chi = m/w, \quad \chi - H \in \mathrm{Inf}, \\ S = \tfrac{1}{2} - \mu - \frac{(t\lambda x)^2}{q} = H - \frac{m^2}{\bar{q}} \geq \frac{H\rho r^2}{\rho r^2 + H} + \frac{H^2 - \chi^2}{\rho r^2 + H}. \end{gathered} \tag{JD13}$$

The first term has positive residue and the second is infinitesimal, also for unbounded ρ. Moreover $x/\sqrt{q} = b/\sqrt{\bar{q}}$ is bounded by JD11 and $\bar{q} \geq Hw^2$. JD6 consequently has strict residue on $(M_D, R_D, N_y, \xi, F_v, J, R, Q)$, with the only possible nonzero cross residue in its strictly positive Q, ξ block. Restore N^2, T^2. This proves the full local parent, not just its contact-state restriction.

Unbounded a: the thin layer and the exact paired-D repair. Assume $0 < \rho = n/e^2$ infinitesimal. JD11 gives $b/w \sim 1/t$, so $\psi = w/b$ is a unit. The formal C1 contact is exactly OP4, with native parameters

$$x_1 = x, \quad \eta_1 = t, \quad \zeta_1 = s, \quad r_1 = h, \quad \theta_1 = e, \quad t_1 = w.$$

TL2 still gives $\beta_1 = -s_n/2$ infinitesimal and $\kappa = (1 + 2n)/(1 - s_n) \sim 1$. The native gaps satisfy $M_1 = g$ a unit and $J_1 = j > 0$. The canceled CA/CS/SE remainders have no inverse J_1; their divergent paired-D mass is retained in SD3. The formal source has $C_1, A_1 h$ positive units, $D_1 j$ a positive unit, positive m_1, α_1, and positive $\beta_1 + 1/2$. These facts follow directly from TL3 and JD12.

Its bounded opening ratio $t_1/e_{1,\mathrm{scale}}^2 = w/(tb)$ is a unit. Thus the explicit CA/CS/SE strict base and their finite conductors apply exactly as in the R16 thin proof; the reservation $(2+4\beta_1)\nu - \nu$ is positive because β_1 is infinitesimal. This is an algebraic use of those formulas, not an application of a physical C1 theorem with negative beta.

Here is the one change needed to the R16 perturbation. In native notation put

$$T_1 = c_a = 1 + xr, \quad a_1 = l/t, \quad b_D = \frac{J_1}{T_1} = \frac{j}{c_a}, \quad h_1 = \frac{a_1}{T_1} = 1 - b_D,$$

$$q_B = v(Z - uX), \quad M_B = (uW - Y) - h_1 q_B, \quad R_{D,\mathrm{event}} = M_B - b_D q_B. \tag{JD14}$$

The old retained paired-D packet in $\kappa F_1/4$ is $\kappa(D_1 + 1/2)h_1 M_B^2$. In the actual parent retain instead the positive contact-zero packet $(D_2 + 1/2)h_1 M_B^2$. For $\delta_D = (D_2 + 1/2) - \kappa(D_1 + 1/2)$ the entire resulting difference is

$$\boxed{\delta_D\{R_{D,\mathrm{event}}^2 - h_1 M_B^2\} = \eta_D\{M_B^2 - 2M_B q_B + b_D q_B^2\}, \quad \eta_D = \delta_D b_D.} \tag{JD15}$$

By the exact TL3 coefficients, with $\Theta = n/B$,

$$\eta_D = \frac{\Theta\{S_c P_D/t - jP_\kappa\}}{K(h+t)c_a}, \qquad \eta_D/\rho = O(w). \tag{JD16}$$

Every denominator is a unit. In particular the previously unbounded $\rho w/j$ correction is absent; no relative w/j bound is needed.

Keep the R16 positive contact-zero n row TL4, the actual centered A packet, and every other unchanged retained packet. The exact full stress is TL6 with its $\delta_D R_{D,\mathrm{event}}^2$ line replaced by the right side of JD15. Both parents and all retained packets have zero contact value and full first jet, so this new entire stress has the same exact vanishing.

For completeness its new term satisfies every finite-support premise of the inherited conductor. The exact inverse state formulas give

$$q_B = v\{D_u - uN - puD_u v\mathfrak{w}\}, \qquad p = t_1/R_1. \tag{JD17}$$

At zero normals $q_B = vD_u V_0$. Hence the new scalar term is in $(v)^2$, both complete first normals are in (v), and the remaining hard entries are bounded. There are no terms above normal degree two. Every dependence on $\mathfrak{w}$ has a factor p. In the CA chart JD16 divided by ρ has bounded coefficients on the same raw roots as TL8. In the SE chart

$$\Theta/\rho = \varepsilon^2\tau^2 QE/K$$

gives the same extra ε^2 factor to every new coefficient. Modulo ε, scalar coefficients are even in u, v and the normal coefficients are independent of $\mathfrak{w}$. Thus the exact support, full-jet, and finite divisibility hypotheses of CC3–CC6 and SE5–SE8 are unchanged. The complete stress divided by ρ has a bounded Gram on the same eight roots. Multiplication by ρ makes it an infinitesimal perturbation of the strict formal C1 base. Restoring the positive packets proves the actual C2 parent on the entire thin contact chart, including both full state normals.

At literal $n = 0$, OP4 is a physical C1 identification with k_1, ζ_1, r_1 infinitesimal, $\eta_1 = t$ a unit and $e_{1,\mathrm{scale}}^2 = tb$ bounded. The physical C1.2/C1.3 all-tests theorems allow arbitrary positive J_1 and literal beta zero, so they cover this boundary exactly.

The unit-W entry on the unbounded-a thin layer. The actual physical C2 critical coefficient is

$$C_* = \frac{(h + 2Ah)(1 + 2D)}{2(4AhD - h)}.$$

Use the polynomial UW3 numerator/denominator with $b = wz_*$, $z_* = b/w$ bounded. Its numerator is divisible by sw because it vanishes identically at $s = 0$ and $w = 0$. The denominator after canceling the m numerator has residue $16t_0 = 16$, also when $j_0 = 0$. Consequently

$$(C_* - C)/m = s \times O(1). \tag{JD18}$$

There is no j inverse in this estimate. SEO12 permits both A, D unbounded: its entire hard block remains strict and its two radial masses tend to $2, 2$. With $W = 1$, the exact testing identity $vZ = Z(R_m + X + uZ)/(1 + uY)$ absorbs the full negative critical correction by JD18 and the retained original m square. All denominators are pure testing units. This proves every local normalized unit-W state, including $Z = 0$; negative-core state normalization leaves exactly the $Z = 1$, $u, v, \mathfrak{w} > 0$ infinitesimal chart proved above. This is the same actual-C2 state entry as UW, with its source bound checked at the new $t_0 = 1$ boundary.

Finite remote tests when e is infinitesimal. For bounded a, $C_* - C$ is infinitesimal because $C_*, C \sim 1/2$. For unbounded a this follows also from JD18 and bounded m. The full critical-wall identities SEO17, SEO18, SEO20, in the normalized SEO12 frame, are strict whenever at least one finite angle has nonzero residue. They explicitly allow A, D unbounded. The entire correction $-4(C_* - C)_+ D_v(uW - Y)^2$ has infinitesimal bounded entries in that frame. Retain original n, m, α and any positive excess. This proves every finite remote test, with all states, for the entire e-infinitesimal R8 domain.

Finite remote tests when e is a unit, including small r. Here a is bounded, m, C bounded, C a positive unit, A, D unbounded, and $N_* = n + 1/2$ is either a unit or unbounded. All normalizations below are pure source. Let $\widehat{X} = \sqrt{N_*}X$, $\widehat{Y} = \sqrt{A}Y$, and $\widehat{D} = \sqrt{D}\, R_{D,\text{event}}$.

If u is a bounded unit and v bounded, use the exact inverse

$$W = \frac{D^{-1/2}\widehat{D} + A^{-1/2}\widehat{Y} + vZ - uvN_*^{-1/2}\widehat{X}}{u}. \tag{JD19}$$

All full coefficients are bounded. The $\widehat{Y}, \widehat{D}$ residue pivots are $(1 + U)(1 + V), 1$, with zero cross residue, where $U = u_0^2 > 0, V = v_0^2 \geq 0$. For unit r, the remaining two-state matrix before X normalization is the complete original parent at $Y = R_{D,\text{event}} = 0$, namely

$$\begin{aligned} &(nD_uD_v + mD_v^2 + CD_vUV + D_v/2)X^2 \\ &+ 2\{-mD_v(V - U)/u - CD_vuV + uD_v/2\}XZ \\ &+ \{m(V - U)^2/U + CD_vV + (U - V)/2\}Z^2. \end{aligned} \tag{JD20}$$

Insert the residue source values JD7, with $e = 1 - r$, $g = 1 - ae$. Its determinant has the exact positive expression

$$U \det = \frac{D_uD_v(1 + ar)}{4r(1 + a)^2 g}\{e(U - aV)^2 + r(g + ar)U^2V + g(g + ar)UV^2\}. \tag{JD21}$$

It is positive for every $U > 0, V \geq 0$. The X diagonal is positive. Division by the positive unit N_* proves strictness after normalization. When r is infinitesimal, divide JD21 by $N_* \sim n$ before taking residue. The result is

$$U \underset{\text{normalized}}{\det} = \frac{D_uD_v}{2(1 + a_0)(1 - a_0)}\{(U - a_0V)^2 + (1 - a_0)^2UV^2\} > 0, \tag{JD22}$$

with $0 \leq a_0 < 1$. The normalized X diagonal tends to D_uD_v. Thus JD19 proves this remote chart for arbitrary relative n, A, D rates, without a comparison between any of them.

It remains to take u infinitesimal and v a bounded unit. If a is a unit, m is a unit. Solve instead for Z by the pure testing unit v. The full residue is

$$m_0(v_0W - N_{*,0}^{-1/2}\widehat{X})^2 + D_v\widehat{X}^2 + D_v\widehat{Y}^2 + \widehat{D}^2, \tag{JD23}$$

where $N_*^{-1/2}$ has residue zero if n is unbounded. This is strict, with the two-state determinant $m_0 V D_v > 0$. If a is infinitesimal, use the extra legal root $S = aW$ and put $R = uW$. The exact D event gives $R = vZ - uvX + Y + D^{-1/2}\widehat{D}$. Substitute this expression in every occurrence except the m event, where use $mR_m^2 = (m/a^2)(vS - aX + auvY - auZ)^2$. All coefficients are bounded, also when n is unbounded after the displayed X normalization. Its entire residue is diagonal on $(S, \widehat{X}, \widehat{Y}, \widehat{D}, Z)$ with pivots

$$((m/a^2)_0 V, D_v, D_v, 1, V^2/2) > 0. \tag{JD24}$$

These identities retain every original event and the complete signed core; introducing a dependent extra root is harmless. The finite separated lift around the real angle residues proves each chart.

Projective endpoints, signs, and the complete join. On all of R8, C is a positive unit and A, D are unbounded. For infinity charts retain the whole original m, n, α packets, clip C, A, D at suitable positive real lower bounds, and retain their positive excesses. The exact homogeneous infinity identities used in SEO19 and its accompanying BE2 routing require only these lower bounds and $d = 1$; they do not require D bounded or m bounded below. In particular the delicate $(u, 1/v) = (0, 0)$ corner has the strict selected residue

$$2X^2 + 4c(uW - Y)^2 + (4A_* + 2)Y^2 + 4D_* Z^2$$

on (X, Y, uW, Z) after retaining m. All leftover W occurrences carry u. The remaining single and double infinities are exactly the same homogeneous event identities. Literal endpoint values are taken in those original homogeneous formulas; no higher-degree contact graph is substituted at infinity.

For unit r, source a is bounded or unbounded. The bounded case has e unit or infinitesimal and is covered locally by JD7 or JD8–JD10. The unbounded case forces e infinitesimal and is covered by the thick/positive-thin/literal-zero split above. These are exhaustive pure-source alternatives. All finite remote tests use JD18–JD24 or the critical wall, and every infinity uses the homogeneous join. Original testing sign gauges, literal-axis event identities, state homogenization, and the real/imaginary split give all signs and complex states. All inverses and radicals are pure source or pure testing. No independent source/testing rate is ever compared.

This proves the entire R8 r-unit source subdomain all tests and the remote complement for every R8 source. If r is infinitesimal, e is a unit and $g > 0$ forces bounded a with $a_0 < 1$; precisely these hypotheses, with every other original intersection unchanged, are those of the joint-n, A, D local lemma. That lemma supplies a strict full-parent Gram with residue $(1, 1, 1, (m/a^2)_0, C_0)$ on all remaining local tests, including the axes. Therefore the two source alternatives r unit or infinitesimal now exhaust R8 with an all-tests certificate. Literal $n = 0$ was retained in the unit-r proof; it is impossible in the small-r part because nr is a unit. The whole ordered R8 shape is closed, with no extra source or testing rate restriction. The 20 new focused checks in `check_joint_D.json` corroborate the new exact identities; the ordered-field bounds and finite-conductor arguments are the proof, not an inference from a numerical sample or a checker count.

Chapter 151

The end-to-end C14.3a/b join

Component `pfaf149c34810`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_joint/c143ab_all_tests_join.tex`
SHA256: `faf149c34810faa485277a66184934962e8bf687e6e5d2bc9a96ccd06abe16b8`
Inventory use: entire analytical join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

151.1 The end-to-end C14.3a/b join

Theorem 151.1. *Retain the original physical stationary C1 source, $d = 1$, all strict gaps and nonnegative event weights, literal $\beta = 0$ when allowed, the original infinitesimal total-scale intersection, and the complement of all previously admitted domains. Each of the following R17 checklist branches has a finite tensor-SOS certificate for every independent projective angle pair and every real or complex state:*

$$\begin{array}{l|l} \text{C14.3a} & L = x\zeta > 0 \text{ infinitesimal}, \quad \eta > 0 \text{ bounded}, \quad \zeta \text{ unbounded}, \\ \text{C14.3b} & L, \eta > 0 \text{ infinitesimal}, \quad \zeta \text{ a positive bounded unit}. \end{array}$$

No bound on L/η or x/η is imposed. No source rate is compared with a testing or state rate. These are complete branch admissions; the other four R17 C1 branches are unchanged.

Precise inherited entries. For compact file citations, let R13 denote the preserved tree `inherited_r17/inherited_r16/Q222_Revision13`. The following are the actual entries used, rather than a deduction from a filename or a numerical check:

1. R13/`advances/c1_lsmall_eta_bounded/remote_reduction.tex`, source bounds SB1–SB4 and infinitesimal-η remote proof SB11–SB12. Its header assumes x, L infinitesimal, ζ noninfinitesimal and η bounded, and admits the exact complement of the chart where both u, v are infinitesimal.

2. R13/`advances/c2_strong_remote/strong_ii_remote.tex`, exact homogeneous parent RR3 and identities RR4–RR9. These are the identities explicitly imported and adapted in SB11–SB12, not an unqualified import of the C2 source theorem.

3. `inherited_r17/proofs/c1_source_ends/large_zeta_unit_eta_all_tests.tex`, the theorem and EU1–EU6, followed by its full normal allocation and final projective join. This admits the whole unit-η part of C14.3a, through every actual-j opening.

4. `inherited_r17/proofs/c1_small_eta/damped_gamma_source_subchart.tex`, exact DG3, positive source deficits DG5–DG9, and all residual allocations preceding DG10; and `inherited_r17/proofs/matrix_mechanism/large_zeta_damped_extension.tex`, LD3–LD13 for the corresponding opening-safe large-ζ source scale. The displayed source-ratio

hypotheses in their older all-tests theorems are used only for DG10/LD14, the term now replaced by the exact CJ allocation.

5. `inherited_r17/proofs/root/core_sign_lemma.tex`, the exact half-core Schur formula and sign gauges; and DS3/LS4 in `inherited_r17/proofs/c1_small_eta/damped_state_extension.tex` and `inherited_r17/proofs/matrix_mechanism/large_zeta_damped_state_extension.tex`, the direct X/Z-unbounded original-parent exit.

6. `proofs/c1_joint/completed_j_conductor.tex`, CJ1–CJ8 and its full local reassembly, which removes the entire missing $\epsilon uvQ\xi$ on the existing completed frame.

Local real-state coverage when eta is infinitesimal. Take the exact sign gauges to arrange $u, v \geq 0$. For each real state, $WZ \leq 0$ is admitted by the pure-testing half-core lemma, including literal $Z = 0$ and both local angle axes. If $WZ > 0$, then $Z \neq 0$. When X/Z is unbounded, the pure-testing positive packet DS3/LS4 and the remaining original nonnegative event packets give a direct certificate. Otherwise X/Z is bounded and CJ applies. The CJ proof imposes no condition on W/Z or Y/Z and uses no angle inverse. It includes all remaining local angles and states, including the former normalized Z/u^2-infinitesimal wedge.

For C14.3b the source scale is $s = x + \eta$ with unit $a = 1/\zeta$; for C14.3a it is $s = L + \eta$ with $a = 1/\zeta$ infinitesimal and $x = aL$. The exact factors in CJ7–CJ8 are infinitesimal in both cases. The local certificate therefore applies to the entire infinitesimal-η portion of both branches, including all previously admitted bounded-source-ratio portions. It does not require separately assuming any older state-ratio or angle-ratio admission.

Audit of the remote complement. Both source branches satisfy the SB header: x, L are infinitesimal, ζ is noninfinitesimal and η bounded. When η is infinitesimal, the actual gap implies $r \sim 1$ and SB3 gives C, D unbounded. The SB11 harmonic source quantities satisfy

$$\gamma = \frac{yCD}{C + D}, \qquad \gamma \sim L/2, \quad A/y \sim 1/(2L), \quad \gamma A/y \sim 1/4, \quad y/\gamma \sim 2\eta \in \mathrm{Inf}\,.$$

These estimates are uniform through both original gap openings and do not require either L/η or x/η bounded.

For noninfinitesimal v, the RR3–RR4 homogeneous full-parent argument requires only that C, D are unbounded; its remaining original packets are nonnegative. The C1 beta event is retained as its original positive square. This covers every u and includes $v = \infty$ and both Alice endpoints. For noninfinitesimal u and infinitesimal or zero v, SB12 uses the exact RR7–RR9 decomposition with zero alpha cap and $k = -1$. Its harmonic normalized block has determinant residue $1 - b_0^2 > 0$, where $b = 1/\sqrt{1 + u^2}$ and $b_0 < 1$. The sole negative term has an infinitesimal Gram allocation by the displayed SB12 identity and $y/\gamma \in \mathrm{Inf}$. The original alpha and beta packets are retained. At $u = \infty$, $b = 0$ in the original homogeneous parent, so the same proof applies. This is the full remote complement, not a contact-graph endpoint evaluation.

Source openings and the unit-eta end. The actual source gap is always $j = \eta(1+xh)-1+r > 0$. Neither CJ nor the remote proof replaces it with a positive assumption on $j_0 = \eta + r - 1$. The opening-cleared DG/LD source identities include $j_0 = 0$ or slightly negative values where permitted. SB2 gives the original D numerator a positive unit residue; hence literal $D = 0$ is impossible on this source face, rather than silently excluded. Literal $\beta = 0$ remains allowed through $B = \beta + 1/2 \geq 1/2$.

The positive bounded η of C14.3a is either infinitesimal or a positive unit. The former case is CJ plus the remote complement just audited. The latter case is precisely the R17 EU theorem, which retains every actual-j and simultaneous r, j opening and includes all projective tests. This exhausts C14.3a. C14.3b has η infinitesimal by definition, so CJ plus the same remote complement exhausts it directly. The exact sign gauges give all angle signs. Applying the real-state certificates to real and imaginary parts gives every complex state without a shared state normalization. Contact equality and all normals are preserved by the exact original-parent identities.

Reconciliation with the original C14.3 work item. The R17 checklist section 4A and `FRONTIER_R17.json` list exactly the two branches above. The earlier unit-η, unit-ζ small-L region is already covered by R13 RO, JO and HO, as recorded and proved in R13/`reconciliation/checklist.tex`. Their exact entries are R13/`advances/revision13_c1_small_l/ r_opening_closure.tex` (RO1), `j_opening_closure.tex` (JO1), and `simultaneous_r_j_closure.tex` (the theorem and HO1–HO3). For clarity, their union is exhaustive: if actual j is a unit, then $j_0 = j - \eta x h$ is a positive unit and RO applies. If actual j is infinitesimal, $\eta + r \sim 1$, so $1 - r$ is a unit; unit r gives JO and infinitesimal r gives HO. The latter two impose no positivity premise on j_0 and no comparison of j with x.

Thus the original C14.3 acceptance criterion is met after joining the unchanged prior admissions: the complete parent, all printed normals, every independent rate, source opening, state and projective test have been covered. C14.3a and C14.3b may be marked closed and the common missing-normal obligation removed. This does not close C14.1a, C14.1b, C14.2, C14.4, the remaining weak-C2 shapes, or the global source-routing obligation U1. No historical checker was rerun in this join audit.

Chapter 152

The completed J row conducts the shared remaining normal

Component `pe3bbfd3a8151`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_joint/completed_j_conductor.tex`
SHA256: `e3bbfd3a81518829c7730d40c9288d4d48050ac03b210526fac1fe109c2e3196`
Inventory use: CJ1–CJ8 and full local reassembly | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

152.1 The completed J row conducts the shared remaining normal

Scope and unchanged inputs. Retain the complete actual-parent identity DG3, equivalently LD9, with $d = 1$, every original source opening and residual intersection, and $\beta \geq 0$ including literal $\beta = 0$. On C14.3b use $s = x + \eta$ and the unit $a = 1/\zeta$; on C14.3a use $s = L + \eta$ with $x = aL$ and a infinitesimal. In both cases s and η are positive infinitesimals. The reviewed R17 source identities give

$$\Lambda = \eta\lambda \sim \tfrac{1}{2}, \qquad \epsilon = O(s), \qquad N_Q = \sqrt{\frac{\Delta_0}{h^2 K}}\,Q, \qquad Q = \frac{\omega}{\sqrt{\eta s}}N_Q, \tag{CJ1}$$

where ω is a bounded positive source unit. The complete strict frame is

$$\mathcal{F} = (N_s, N_T, N_Q, R_b, f_t, \mathcal{Q}, N_x), \qquad N_x = \sqrt{B}\,\xi, \quad B = \beta + \tfrac{1}{2}. \tag{CJ2}$$

All DG3/LD9 residuals other than $\epsilon uvQ\xi$ already have infinitesimal Gram coefficients on this frame without any bound on s/η. The two residuals allocated in DS2/LS2 have that same property whenever $Z \neq 0$ and X/Z is bounded. Thus the only expression to be allocated on that state chart is

$$N = \epsilon U(W/Z)E\xi, \qquad E = \eta(P + Q) = \eta uW - vZ. \tag{CJ3}$$

There is no restriction on Y/Z or W/Z below.

An exact identity from the already retained completed row. Let $\phi = \xi/Z = X/Z + hu$, and retain exactly DG's completed square

$$\widehat{J} = \sqrt{\lambda}\,(uv - x/c)Z + \frac{G_0 u}{2\sqrt{\lambda}}Q, \qquad G_0 = 2\eta\lambda - 2\eta, \qquad c^2 = a/\eta.$$

The defining identity $vZ = \eta uW - \eta(P + Q)$ gives

$$\boxed{\frac{\widehat{J}}{\sqrt{\lambda}} = \eta UW - \frac{x}{c}Z - \eta u\left(P + \frac{Q}{\lambda}\right).} \tag{CJ4}$$

Multiply CJ4 by $\epsilon\phi E/\eta$. The complete remaining normal has the exact conductor

$$\boxed{N = \frac{\epsilon\phi}{\eta\sqrt{\lambda}} E\widehat{J} + \epsilon\phi u E\left(P + \frac{Q}{\lambda}\right) + \frac{\epsilon x}{\eta c} E\xi.} \tag{CJ5}$$

The only state inverse here is the pure-testing Z^{-1}. No inverse of u, v, W, or a mixed expression occurs. No radial row, source pivot, or completed square is duplicated.

Bounded allocations and infinitesimal coefficients. Define two dependent roots

$$\mathcal{E} = \sqrt{s/\eta}\, E, \qquad \mathcal{P} = P + Q/\lambda.$$

Both have bounded allocations on the existing source roots (N_s, N_T, N_Q). Indeed the exact DG9/LD10 inverse identities give coefficients $O(\sqrt{\eta})$, $O(\sqrt{\eta})$ and $O(\sqrt{\eta/s})$ for P. Hence the coefficients for $\sqrt{\eta s}\, P$ are bounded (in fact infinitesimal), whereas

$$\mathcal{E} = \sqrt{\eta s}\, P + \omega N_Q, \qquad Q/\lambda = \frac{\omega}{\Lambda}\sqrt{\eta/s}\, N_Q. \tag{CJ6}$$

Here $\eta/s \le 1$ is a pure-source inequality. These are bounded allocations of dependent roots, not new positive pivots. For completeness, the other two terms in the exact DS2/LS2 split have the particularly simple allocations

$$\epsilon uvQ\xi - N = -\frac{\epsilon}{s} u\phi\mathcal{E}^2 - \frac{\epsilon uv}{\sqrt{B}} P N_x. \tag{CJ6a}$$

Their coefficients are infinitesimal, since u, v are local infinitesimals and P has a bounded source-root allocation.

In these roots CJ5 is exactly

$$\begin{aligned} N = {} & \frac{\epsilon}{s}\frac{\sqrt{s}}{\sqrt{\Lambda}}\phi\mathcal{E}\widehat{J} \\ & + \epsilon\sqrt{\eta/s}\, u\phi\mathcal{E}\mathcal{P} + \frac{\epsilon}{s}\frac{x\sqrt{s}}{\sqrt{a}\sqrt{B}}\mathcal{E}N_x. \end{aligned} \tag{CJ7}$$

All three coefficients are infinitesimal bounded tensor elements. For the first two use ϵ/s bounded, s infinitesimal, $\eta/s \le 1$, and bounded $\phi = X/Z + hu$. The last coefficient is also infinitesimal:

$$\frac{x\sqrt{s}}{\sqrt{a}} = \begin{cases} O(x\sqrt{s}), & a \text{ a bounded positive unit}, \\ \sqrt{a}\, L\sqrt{s}, & x = aL,\ a \text{ infinitesimal}. \end{cases} \tag{CJ8}$$

Also $1/\sqrt{B}$ is bounded, even at literal $\beta = 0$ and when B is unbounded. The apparent inverse $\sqrt{a}$ is canceled in the large-ζ case before taking bounds. Every coefficient is a finite separated expression; ϕ is expanded as $X/Z + hu$. No source rate is compared with any testing rate.

Full local reassembly. Add the already present row $\widehat{J}^2$ to the strict frame CJ2 and allocate CJ7 to it and the source roots. This produces an infinitesimal Gram perturbation of the previous strict-residue matrix plus a unit $\widehat{J}$ diagonal entry. The separated bounded-Gram lemma therefore applies to the full remainder. Retain each other positive row from DG3/LD9 exactly once:

$$\widehat{R}^2, \quad c_U U Q^2, \quad e_4 U F_t^2, \quad BV\xi^2, \quad (2x/c)uv\tau^2, \quad (G_0 x/c) D_0^2 WZ.$$

The last two are literal tensor squares on the unchanged sign chart $u, v \ge 0$, $WZ > 0$. The complete angular and radial packets remain inside CJ2; all full-D, full-Y and higher normals are those of the exact original-parent identity, not truncated expansions.

Consequently the entire local original parent is tensor SOS when $Z \neq 0$ and X/Z is bounded, with no condition on s/η, u^2W/Z, or Y/Z. The proof is valid on literal local angle axes because it uses no angle inverse. At original contact E, P, Q, ξ and $\widehat{J}$ vanish, so this reassembly preserves equality and all original contact normals exactly.

The local state join and the remaining global bookkeeping. The reviewed pure-testing half-core exit covers $WZ \leq 0$, including $Z = 0$. When $Z \neq 0$ and X/Z is unbounded, DS3/LS4 is a direct original-parent tensor-SOS exit. The above argument covers every remaining real state on the local angle chart. Applying these real certificates to real and imaginary state parts gives the full complex parent. Thus the shared local C14.3a/b obligation is closed, with all source gaps and $\beta = 0$ retained. The projective remote charts and the previously admitted unit-η source end are the unchanged R17 inputs for the overall C14.3a/b join; this local proof introduces no new condition needed by that join.

The focused checker `check_completed_j_conductor.py` checks CJ4 and CJ5 exactly. The reviewed DG/LD parent identity, source deficit calculations and prior strict-frame allocations are reused without rerunning their historical checkers.

Chapter 153

C14.4 at large ζ: every unmatched Bob residue

Component `p1a6cc124f5db`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_large_eta/off_matched_bob.tex`
SHA256: `1a6cc124f5db69e647dbc27b8df98ba0f8cbbf3f311a8504998f074d4fa265f0`
Inventory use: EB1–EB10 | EB exact homogeneous parent behind SE | EB1-EB10 complete homogeneous full-state unmatched chart

153.1 C14.4 at large ζ: every unmatched Bob residue

Precise admission. Retain the entire original C14.4 source: $x, L > 0$ infinitesimal, η unbounded, $\zeta = L/x$ noninfinitesimal, all original physical intersections, every positive opening, and either permitted literal $D = 0$ or $\beta = 0$. This section additionally assumes ζ unbounded. Put

$$\sigma = 1/\eta, \qquad S = \sigma + L, \qquad \nu = \sigma/S, \qquad \ell = L/S = 1 - \nu.$$

For a real projective Bob test choose its normalized homogeneous coordinates (c, s), so $c^2 + s^2 = 1$; on the finite chart these are $c = (1 + v^2)^{-1/2}$, $s = vc$. The complete original parent is tensor SOS at every independent test and state satisfying

$$\overline{s^2} \neq \overline{\ell}. \tag{EB1}$$

Bars mean real residues of bounded elements in their respective fields. This compares real residues only, not independent infinitesimal rates. The adjacent-C, A theorem already covers noninfinitesimal Alice tests. Hence only u infinitesimal is proved below. No all-tests closure of C14.4 is asserted: the matched residue and bounded-unit ζ source remain.

Source limits, including both sphere endpoints. Use ES1–ES10 without changing their source entry. Since $x/S \leq x/L = 1/\zeta$ is infinitesimal, ES7 gives $e = xh/\sigma$ infinitesimal, and hence $E \sim 1$. Moreover $M = 1 + xr - \theta/\zeta \sim 1$. ES8 then gives m infinitesimal. The exact equation ES6, divided by S, is

$$e(B/S) = (x/S)H_0 + \nu\mathcal{H}b_0^2.$$

Both summands are nonnegative; B/S and $\mathcal{H}$ tend to one. Consequently νb_0^2 is infinitesimal. Therefore ES5 yields

$$\frac{D}{S} - \frac{\nu^2}{2} \in \mathrm{Inf}, \qquad \frac{\alpha}{S} - \frac{\ell^2}{2} \in \mathrm{Inf}, \qquad m \in \mathrm{Inf}_{>0}\,. \tag{EB2}$$

For the second formula, divide the exact ES9 expression by S:

$$\frac{\alpha}{S} = \frac{(1 + \sigma r)\{\ell Re + \ell^2 ME\}}{2Q(P/S)}.$$

All displayed denominator factors tend to one. These are additive residue assertions, valid also when ν or ℓ is infinitesimal. In particular no division by either sphere coordinate is used.

The opening-cleared ES8 bounds give

$$c_* = SC(1-r) \sim \tfrac{1}{2}, \qquad a_* = SAr \sim \tfrac{1}{2}.$$

Define the pure-source harmonic quantities

$$H = S(C+A), \qquad \rho = C/(C+A), \qquad a_h = SCA/(C+A) = \frac{c_* a_*}{c_* r + a_*(1-r)}. \tag{EB3}$$

Then $1/\sqrt{H}$ is bounded, $0 < \rho < 1$, and $a_h \sim 1/2$, uniformly through both r openings. No upper bound on C or A has been assumed.

An exact all-state frame for every Bob chart. Let $D_u = 1 + u^2$ and $A_u = 1 + u^2 s^2$, both positive testing units. Use the separated roots

$$p = \frac{uW}{\sqrt{S}}, \quad q = \sqrt{S}Z, \quad y = \frac{Y}{\sqrt{S}}, \quad N_y = \sqrt{H}(y - \rho p), \quad N_m = \sqrt{m}\, sW,$$

$$N_b = \sqrt{A_u}\left(X + \frac{uc^2}{A_u} Z\right), \qquad N_r = \frac{\sqrt{D_u}\, usc}{\sqrt{A_u}} Z. \tag{EB4}$$

Only pure-source and pure-testing radicals or inverses occur. In particular no inverse of c, s, u, or a state coordinate is needed. The inverse harmonic allocation $y = \rho p + N_y/\sqrt{H}$ has bounded coefficients.

Write $b = X + uZ$. The exact original signed core K/D_v in finite Bob coordinates equals the following homogeneous expression:

$$\begin{aligned} K_B = {} & N_b^2 + N_r^2 - 2D_u sc\, pq + Sc^2(p-y)^2 + SD_u s^2 y^2 \\ & + \frac{2\sqrt{S}\, usc}{\sqrt{A_u}}(p-y)N_b + \frac{2\sqrt{S}\, us^2\sqrt{D_u}}{\sqrt{A_u}}(p-y)N_r. \end{aligned} \tag{EB5}$$

Here $D_u s$ denotes $D_u\, s$, and similarly in the other products. One derivation starts with CE2 divided by D_v, sets $R = usZ$, and completes its entire (b, R) block:

$$A_u b^2 - 2D_u s\, bR + D_u R^2 = N_b^2 + N_r^2.$$

The cross term from $[\sqrt{S}\, c(p-y) + usb]^2$ is allocated by $cb = cN_b/\sqrt{A_u} + \sqrt{D_u} sN_r/\sqrt{A_u}$. This proves EB5 without deleting a first state normal.

The original positive event roots have the exact bounded expressions

$$\begin{aligned} M_B &= N_m - \frac{\sqrt{m}\, c}{\sqrt{A_u}} N_b - \frac{\sqrt{mD_u}\, s}{\sqrt{A_u}} N_r + \sqrt{mS}\, us\, y, \\ D_B &= \sqrt{DS}\, c(p-y) + \frac{\sqrt{D}\, us}{\sqrt{A_u}} N_b - \sqrt{D/S}\, \frac{D_u s}{A_u} q, \\ A_B &= \sqrt{\alpha D_u S}\, sy - \sqrt{\alpha D_u / S}\, cq. \end{aligned} \tag{EB6}$$

They are, respectively, the normalized original m, D, α event amplitudes. The formulas are valid at $D = 0$ as well. In the D row, the identity $usb - D_u sZ = usN_b/\sqrt{A_u} - D_u sZ/A_u$ avoids an apparent reciprocal-c pole.

Consequently the full original quarter-parent, divided by D_v on the finite chart, is exactly

$$\mathcal{P}_B = N_y^2 + a_h p^2 + ASu^2 y^2 + M_B^2 + D_B^2 + A_B^2 + \beta b^2 + \tfrac{1}{2} K_B. \tag{EB7}$$

The two explicit positive rows ASu^2y^2 and βb^2 can be retained unchanged. All other rows in EB7 use EB4 once. The formula is a full four-state identity and is not a restriction to a state ray.

Strictness precisely off the matched residue. Every coefficient of EB5 except the printed pq term is either its fixed positive pivot or infinitesimal after EB3. In EB6, the m row is N_m plus infinitesimal coefficients; the D and α rows have only their q coefficients at residue. Use EB2, $A_u, D_u \sim 1$, and EB3. The resulting residue Gram on $(N_y, N_m, N_b, N_r, p, q)$ is the direct sum of $\mathrm{diag}(1, 1, 1/2, 1/2)$ and

$$\frac{1}{2}\begin{pmatrix} 1 & -\bar{s}\bar{c} \\ -\bar{s}\bar{c} & \bar{\nu}^2\bar{s}^2 + \bar{\ell}^2\bar{c}^2 \end{pmatrix}. \tag{EB8}$$

The determinant of this last block is

$$\frac{1}{4}(\bar{s}^2 - \bar{\ell})^2. \tag{EB9}$$

Thus EB1 makes the entire bounded finite Gram strictly positive. The separated bounded-Gram lemma gives tensor SOS, and restoring the two retained positive rows proves the complete parent. The roots are allowed to be dependent or even identically zero at an endpoint; the claim concerns the coefficient Gram, so this does not affect the argument.

The same homogeneous formulas allow $c = 0$ and $s = 0$ literally. They include every Bob projective endpoint, every sign, and every local Alice axis. Nonlocal Alice charts use the inherited adjacent C, A theorem. The real and imaginary state parts are treated separately. Every normalization is pure source or pure testing.

The exact remaining obstacle. The leading scalar has the useful two-square identity

$$\tfrac{1}{2}p^2 - sc\,pq + \tfrac{1}{2}(\nu^2 s^2 + \ell^2 c^2)q^2 = \tfrac{1}{2}(p - scq)^2 + \tfrac{1}{2}(s^2 - \ell)^2 q^2. \tag{EB10}$$

It does not complete the matched-residue chart: there the second pivot collapses, and the actual source corrections must be allocated through exact contact conductors. Arbitrary small positive D or β openings do not supply a missing fixed pivot. Unit ζ also requires new source limits because e can have nonzero residue. These two obligations are retained, not declared empty.

The focused checker verifies EB5–EB7 as complete homogeneous quadratic identities and EB9–EB10 algebraically. Source estimates EB2–EB3 are the analytical consequences of the already proved ES identities given above. This is a partial-domain admission only.

Chapter 154

C14.2: bounded Bob tests at every local Alice angle

Component `pda4a21937f64`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_lunit/bounded_bob.tex`
SHA256: `da4a21937f645d4d9d43d5603c3689ea61354955c84aa8b2002b575469b8c8bf`
Inventory use: LB1–LB5 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

154.1 C14.2: bounded Bob tests at every local Alice angle

Proposition 154.1. *Retain the unchanged full C14.2 source and every original intersection as in the preceding noninfinitesimal-Alice proposition. For every infinitesimal or zero Alice test u and every bounded Bob test v, the entire original parent is finite tensor SOS for all real or complex states. Every source opening and both $D = 0$, $\beta = 0$ are included. Together with that proposition this leaves exactly u infinitesimal and v projectively unbounded, equivalently $u, w = 1/v$ infinitesimal, as the remaining C14.2 testing corner.*

The same source weights and a pure-source strict matrix. Keep the opening-safe positive source weights from LA1,

$$C_0 = \frac{1}{2L(1-r)}, \quad A_0 = \frac{1+L(1-r)}{2Lr}, \quad a_0 = \frac{L}{2(1-Lr)}.$$

Let $c = (1+v^2)^{-1/2}$, $s = vc$, so $c^2 + s^2 = 1$. Since v is bounded, c is a positive testing unit. Write $s_0 = \operatorname{res} s \in (-1, 1)$ and $c_0^2 = 1 - s_0^2 > 0$. These are real numbers, not source/testing comparisons. Consider the pure-source quadratic form in three placeholder variables

$$M_0(P, Y, T) = (C_0 + c_0^2/2)(P - Y)^2 + (A_0 + s_0^2/2)Y^2 + a_0(s_0 Y - T)^2 - s_0 PT. \tag{LB1}$$

Its Gram has first leading minor $C_1 = C_0 + c_0^2/2 > 0$, second leading minor $C_1[A_0 + s_0^2(a_0 + 1/2)] > 0$, and

$$\det M_0 = \frac{c_0^2\{1 + L(1-r)\}^2}{8Lr(1-r)(1-Lr)} > 0. \tag{LB2}$$

This is an exact source identity, not just a limiting determinant. Moreover M_0^{-1} has bounded entries through every opening. Indeed every cofactor is a sum of bounded multiples of $C_0A_0, C_0a_0, A_0a_0, C_0, A_0, a_0, 1$; terms involving a square of a single divergent weight cancel. Multiplication by $8L^2r(1-r)(1-Lr)$ makes every cofactor polynomial in bounded L, r, s_0. Division by LB2 therefore leaves only the source-unit denominators L, c_0^2, and $1 + L(1-r)$. Explicitly, with

$e = 1 - r$ and $B_L = 1 + Le$,

$$M_0^{-1} = \frac{1}{c_0^2 B_L} \begin{pmatrix} 2L & 2Lr & 2s_0 \\ 2Lr & 2Lr(1 - s_0^2 e) & 2s_0 r(1 + Lec_0^2) \\ 2s_0 & 2s_0 r(1 + Lec_0^2) & 2(1 - Lrc_0^2)(1 + Lec_0^2)/L \end{pmatrix}. \tag{LB2a}$$

Choose any exact positive pure-source congruence $M_0 = N_1^2 + N_2^2 + N_3^2$. Its inverse allocation of P, Y, T on these three roots is bounded: each inverse-row norm squared is the corresponding diagonal of M_0^{-1}. This uses only a pure-source Gram, with the fixed real number s_0 as parameter; no mixed matrix is inverted or square-rooted.

Freeze the only divergent Bob-dependent event exactly. Use the legal raw roots

$$P = uW, \quad X, \quad Y, \quad T = cZ - (s - s_0)Y, \qquad d_s = s - s_0 \in \mathrm{Inf}_{\mathrm{test}}\,. \tag{LB3}$$

In particular the entire alpha event satisfies $sY - cZ = s_0 Y - T$ exactly. Thus no product of an unbounded alpha weight with d_s is ever estimated. No inverse of c, u, or a state coordinate is needed in these root definitions.

Let $B_0 = c_0^2(P - Y)^2 + s_0^2 Y^2 - 2s_0 PT$. The complete original half-core after Bob normalization is $K_B/2$, where

$$K_B = \{c(P - Y) + usX - sZ\}^2 + (1 + u^2)(sY - cZ)^2 + (X + uZ)^2 - (1 + u^2)Z^2.$$

It has the following exact, division-free allocation on LB3:

$$\begin{aligned} K_B = {} & X^2 + B_0 + d_s P\{-(s + s_0)P + 2s_0 Y - 2T\} \\ & + 2ucX(sP - s_0 Y + T) + u^2 s^2 X^2 + u^2(s_0 Y - T)^2. \end{aligned} \tag{LB4}$$

This keeps the full signed core and all u-dependent hard normals.

Exact actual-parent reassembly. Use exactly LA4's pure-source factor $f = 1 - (|C/C_0 - 1| + |A/A_0 - 1| + |\alpha/a_0 - 1|) > 0$. Retain the actual original event excesses $C - fC_0$, $A - fA_0$, $\alpha - fa_0$, and the whole m, D, β packets. On the finite Bob chart the original quarter-parent divided by $1 + v^2$ is then the sum of those positive packets and exactly

$$S = f\{C_0(P - Y)^2 + A_0(1 + u^2)Y^2 + a_0(1 + u^2)(s_0 Y - T)^2\} + K_B/2.$$

Substitution of LB1 and LB4 gives

$$\begin{aligned} S = {} & fM_0 + \tfrac{1}{2}X^2 + fA_0 u^2 Y^2 + fa_0 u^2(s_0 Y - T)^2 \\ & + \tfrac{1}{2}u^2 s^2 X^2 + \tfrac{1}{2}u^2(s_0 Y - T)^2 \\ & + \tfrac{1}{2}(1 - f)B_0 + \tfrac{1}{2}d_s P\{-(s + s_0)P + 2s_0 Y - 2T\} \\ & + ucX(sP - s_0 Y + T). \end{aligned} \tag{LB5}$$

The first two lines consist of a strict source-normalized Gram on (N_1, N_2, N_3, X) and exact retained positive squares. All terms in the last two lines have infinitesimal bounded coefficients on that same frame: $1 - f$ is source infinitesimal, d_s, u are testing infinitesimals, c, s, s_0 bounded, and LB2 supplies a bounded inverse allocation for all three raw roots. Since $f \sim 1$, the resulting complete Gram has residue $\mathrm{diag}(1, 1, 1, 1/2)$. The separated bounded-Gram lemma and restoration of every retained packet prove the full original parent. The normalization factor $1 + v^2$ is pure testing and positive.

Both coefficient-boundary packets were retained rather than capped or inverted. The argument works literally at $u = 0$ or $v = 0$, allows signed testing angles, and extends to complex states by real/imaginary splitting. It gives no claim for c infinitesimal: there $c_0 = 0$, the determinant in LB2 vanishes, and the reciprocal two-small-angle corner remains a separate obligation.

The new focused checks establish LB2, opening-regular cofactors, the complete original-core identity LB4 and the entire selected-parent identity LB5. They do not rerun prior source identities or assert the remaining C14.2 corner closed.

Chapter 155

The exact C14.2 remainder and its centered C extraction

Component p156da4167158
Source: BASE/inherited_r19/inherited_r18/proofs/c1_lunit/centered_c_reduction.tex
SHA256: 156da41671584030894d6ef3a93cfa2ce9335b643c11236624553c491c1ce9e0
Inventory use: LC2–LC4 only | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

155.1 The exact C14.2 remainder and its centered C extraction

Quantified remainder after the two new test admissions. The propositions in noninfinitesimal_alice.tex and bounded_bob.tex admit all C14.2 tests except $u, w = 1/v$ infinitesimal. Literal $u = 0$ or $w = 0$ has a direct original-core SOS, so after the exact sign gauges a remaining test has

$$u, w > 0 \text{ infinitesimal}, \qquad L > 0 \text{ a bounded source unit}, \quad x, \sigma = 1/\eta > 0 \text{ infinitesimal}. \tag{LC1}$$

All original strict gaps, coefficient signs, total-scale and prior exclusions remain; $D = 0$ and $\beta = 0$ are not removed. The pure-testing core-sign exit allows restriction to $WZ > 0$, with every state magnitude otherwise unrestricted. No assertion that this residual intersection is nonempty is made.

Use the exact BE2–BE8 source sphere and the entire generic JR5/JB1 parent in the form SP6 from inherited_r17/proofs/c1_sphere/sphere_pole_reduction.tex. Those identities are algebraic for $z = L^{-1}$ a bounded unit as well as for large L. On C14.2, the original $r \to 1$ opening may make C unbounded; the large-L boundedness assertion for SP5 therefore cannot be imported. The following exact extraction removes that particular pole without changing the source.

The entire C-dependent block. Retain all SP/JR roots and definitions, notably

$$q^2 = z, \quad b = Lr, \quad D_w = 1 + w^2, \quad J_C = N_y + bR, \quad \Gamma = \sigma Z - uwW.$$

With all other actual source coefficients held fixed, the complete SP6 matrix is affine in the displayed coefficient C. Inspection of the full H_y, K_r, R_y, R_γ formulas gives its entire C-dependent part as

$$4CzD_wJ_C^2 + 8C(r-1)J_C(z\sigma R - qw\Gamma). \tag{LC2}$$

This includes both complete normal couplings and the $R\Gamma$ coupling. Neither the scalar radial diagonal nor a first normal has been silently removed. No other displayed SP6 coefficient depends on the formal variable C once the other actual source coefficients are held fixed.

Choose the bounded positive pure-source cap

$$C_* = \frac{C}{1+C}, \quad d_C = C - C_* \geq 0, \quad t_C = d_C(1-r).$$

The original opening-cleared estimate $C(1-r) \asymp z$ proves t_C bounded, uniformly through the $r \to 1$ opening. No source case split is needed. The whole parent core therefore satisfies the exact identity

$$\boxed{G = G|_{C=C_*} + 4zD_w d_C J_C^2 - 8t_C J_C(z\sigma R - qw\Gamma).} \tag{LC3}$$

Formal substitution in the first term does not assert that a capped source is physical. The equality is a full-parent matrix identity, and every other original retained square remains unchanged. The extracted row is centered at the original contact, since $N_y = R = 0$ there. It retains its whole event dependence at every state and angle; this is not a Hessian-only extraction.

For reference, the existing exact SP12 conductor gives

$$z\sigma R - qw\Gamma = qu(w^2 - \sigma z)W. \tag{LC4}$$

Both LC3 forms have only separated source/testing coefficients and no new state, angle, or mixed inverse.

What is and is not resolved. LC3 removes the C-opening pole: C_*, t_C are bounded and the correction carries σ or w. It does not give a positive Gram. The Γ direction can lose its pivot at $D = 0$; its normals still require joint completion with the source sphere, radial row and full h/σ packet. C14.2 remains open precisely on LC1 with its original intersections. The focused checker verifies LC2 and LC4 only.

Chapter 156

C14.2: the complete noninfinitesimal-Alice complement

Component `pb0d9d28d0849`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_lunit/noninfinitesimal_alice.tex`
SHA256: `b0d9d28d0849fda84eab4ad896a2b7cafc9bec2cb395cd303191bb7a19e23c82`
Inventory use: LA1–LA7 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

156.1 C14.2: the complete noninfinitesimal-Alice complement

Proposition 156.1. *Keep the unchanged physical C14.2 source: $d = 1$, $x > 0$ infinitesimal, $L = x\zeta$ a positive bounded unit, η unbounded, all original strict gaps, nonnegative event weights, total-scale and earlier-exclusion intersections. Then the complete original parent is finite tensor SOS for every noninfinitesimal Alice angle u, including $u = \infty$, every independent Bob angle, and all real or complex states. Literal $D = 0$ and $\beta = 0$ are retained. Thus only infinitesimal or zero u, with arbitrary Bob type, remains in C14.2. No entire-source admission is asserted.*

Opening-safe source input. Use LH1–LH3 in the preserved R13 file `advances/c1_lunit_eta_large/source_frontier.tex` and BE1–BE9 in `advances/c1_eta_large_source/beta_shared_source_normalization.tex`. Put $e_r = 1 - r$, $Q = 1 - Lr > 0$ and define positive pure-source weights

$$C_0 = \frac{1}{2Le_r}, \qquad A_0 = \frac{1 + Le_r}{2Lr}, \qquad a_0 = \frac{L}{2Q}. \tag{LA1}$$

Their actual-weight ratios satisfy $C/C_0, A/A_0, \alpha/a_0 \sim 1$ uniformly through r, e_r, and Q openings. Each leading and actual weight has a positive real lower bound, since L is a positive bounded unit and $0 < r, e_r, Q < 1$. No upper bound for the individual weights is assumed. The BE source sphere and both its endpoints remain unchanged; no coefficient D, β, d_0 or b_0 is divided by.

The reciprocal Bob strip with a unit Alice angle. Let u be a bounded nonzero unit and $w = 1/v$ infinitesimal or zero. Use four legal raw roots

$$P = uW, \quad X, \quad Y, \quad T = wZ, \qquad U = u^2, \quad D_u = 1 + U, \quad D_w = 1 + w^2.$$

The exact original signed half-core $K_w/2$ is CE6 of the preserved R13 file `baseline/advances/c1_common/common_remote.tex`. Written in these roots it has the exact split

$$K_w = K_0 + w^2\{(P - Y)^2 + X^2\} + 2uwX(T + P - Y), \tag{LA2}$$

$$K_0 = D_u X^2 + D_u Y^2 + UT^2 - 2UTY - 2PT.$$

This is the complete core, including its sign; no state graph is used. The selected positive events are $CD_w(P-Y)^2$, $AD_uD_wY^2$, and $\alpha D_u(Y-T)^2$ in the quarter-parent. The whole original m, D, β packets are retained separately.

The LH4 contact identity implies the following exact identity for the leading selected form at $w = 0$:

$$\begin{aligned} Q_0 &= C_0(P-Y)^2 + A_0D_uY^2 + a_0D_u(Y-T)^2 + K_0/2 \\ &= \frac{D_u}{2}X^2 + C_0\{P - Y - Le_rT\}^2 + (A_0 + a_0 + 1/2)\{Y - LrT\}^2 \\ &\quad + U\{A_0Y^2 + (a_0 + 1/2)(Y-T)^2\}. \end{aligned} \tag{LA3}$$

Every displayed coefficient is nonnegative pure source or a positive pure-testing factor. The source coefficients in LA3 can be clipped at positive real lower bounds, retaining their exact positive square excesses. The resulting bounded Gram on (P, X, Y, T) has strictly positive residue: X is controlled directly, the two U terms control Y, T because u is a unit, and the first contact square then controls P. The coefficients Le_r, Lr in these roots are bounded pure source, including all original openings.

Exact common-factor accounting for the actual weights. To transfer LA3 to the actual parent, define solely in the source field

$$\delta = |C/C_0 - 1| + |A/A_0 - 1| + |\alpha/a_0 - 1|, \qquad f = 1 - \delta.$$

Then δ is nonnegative infinitesimal, $f > 0$ is a unit, and each of $C - fC_0$, $A - fA_0$, $\alpha - fa_0$ is nonnegative. Retain their original event packets as exact squares. The complete remaining selected polynomial is, identically,

$$\begin{aligned} S &= f\{C_0D_w(P-Y)^2 + A_0D_uD_wY^2 + a_0D_u(Y-T)^2\} + K_w/2 \\ &= fQ_0 + fw^2\{C_0(P-Y)^2 + A_0D_uY^2\} \\ &\quad + (1-f)K_0/2 + (K_w - K_0)/2. \end{aligned} \tag{LA4}$$

The second line's w^2 packets are retained positive squares with their complete actual source scales. The third line, including the signed $(1-f)K_0/2$ term, is a bounded infinitesimal Gram on the four raw roots, by LA2 and $1 - f = \delta \in \mathrm{Inf}$. Clip the completed source-square coefficients of fQ_0 at positive real lower bounds as above. Its retained bounded Gram remains strict under this entire infinitesimal perturbation. The separated bounded-Gram lemma gives tensor SOS; restoring all square excesses proves the whole original parent. This is exact polynomial accounting in the tensor ring, not an inequality that orders mixed source and testing quantities.

The other Bob types and the unbounded Alice endpoint. If u is a bounded nonzero unit and v is a bounded nonzero unit, use the inherited noncommuting-residue theorem on the unchanged physical parent. If u is a bounded nonzero unit and v is infinitesimal or zero, clip the original C, A, α at positive real lower bounds c_*, a_*, a'_* and retain every excess and the whole original m, D, β packets. The complete selected residue is

$$(c_* + 1/2)(uW - Y)^2 + a_*D_uY^2 + a'_*D_uZ^2 + \tfrac{1}{2}(X + uZ)^2 \succ 0. \tag{LA5}$$

It is strict on all four original states when u is a unit; every finite-angle correction has bounded infinitesimal coefficients. This covers all bounded Bob types without invoking LA4 outside its stated reciprocal strip.

For u unbounded, put $p = 1/u$ and use the original homogeneous parent. The exact CE4–CE5 raw-frame proof, with any positive real caps for C, A, covers every bounded v. Its strict selected residue is

$$4c_*(1+v^2)W^2 + 4a_*(1+v^2)Y^2 + 2(W+R)^2 + 2(B - vY)^2, \quad B = pX + Z, \quad R = vX. \tag{LA6}$$

For unbounded v, the exact twice-homogeneous CE7–CE8 proof has strict residue

$$4c_*W^2 + 4a_*Y^2 + 2X^2 + 2(T - Y)^2, \qquad T = wZ. \tag{LA7}$$

These two homogeneous proofs need only positive real lower bounds for C, A at these endpoints. They do not use the stronger "C, A unbounded" hypothesis needed for the finite-unit-angle part of the adjacent-event theorem. All unselected original packets and positive excesses are retained exactly. Literal $p = 0$ and $w = 0$ are the polynomial values of these original homogeneous parents.

The finite unit-Alice and homogeneous unbounded-Alice cases exhaust all noninfinitesimal u. Exact sign gauges and real/imaginary splitting give both signs and complex states. The possible additional $D = 0$ null direction at $u = w = 0$ is untouched, since the proposition does not include the infinitesimal-Alice chart. The sole remaining C14.2 testing obligation is now u infinitesimal or zero, with arbitrary Bob types and the unchanged full source intersection.

The focused checker verifies only the new LA2–LA4 algebra, including the entire signed core and the exact common-factor remainder. No historical checker is rerun and no branch closure is inferred from this testing-complement result.

Chapter 157

C14.1b: every state when the first testing angle is a unit

Component `pcc84202479d2`
Source: `BASE/inherited_r19/inherited_r18/proofs/c1_sphere/unit_u_all_states.tex`
SHA256: `cc84202479d2eeb94e997dd13b7b81bcd21f19a0a552578462ed1e86a72c5bab`
Inventory use: Inherited C14.1b nonzero-unit-u admission for all states | complete unit-u theorem | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

157.1 C14.1b: every state when the first testing angle is a unit

Retain the entire original C14.1b source intersection and the exact SP6 parent from revision 17. Assume u is a bounded nonzero unit and w is infinitesimal. The argument below proves the whole parent for every testing state, including the previously unadmitted W/Z-unbounded states. It does not cover infinitesimal u. All source gaps, $D = 0$, both source sphere endpoints and the literal coefficient boundaries remain unchanged.

No state chart is needed. All roots below are homogeneous polynomials in the original testing state, with pure-source coefficients. In particular the proof uses no state-coordinate inverse, no bound on W/Z, and no sign restriction. The only testing inverse is that of the unit u. Write $U = u^2$, $s = x/\sigma$, and preserve the exact SP/BS coefficients

$$L_h = \lambda\sigma/h, \quad \rho = \lambda x/\sigma, \quad d = 4D/\sigma^2, \quad K_A = 4\kappa h/\sigma - \lambda.$$

The source identities give $s \in \mathrm{Inf}$, $\sigma/\sqrt{h} \in \mathrm{Inf}$, $L_h \sim 2$, $\nu \sim 1$, and

$$d + \rho \sim 2, \quad K_A - (2 - 4h) \in \mathrm{Inf}, \quad \chi - (4 - 4h) \in \mathrm{Inf}, \quad (c_G + zK_r) - 2(1 + U) \in \mathrm{Inf}. \tag{SU2}$$

The last infinitesimal is a finite separated expression for bounded u and infinitesimal w. For the only new source assertion in SU2, the sphere identities give exactly

$$d = \frac{2\mathcal{H}}{PJ} d_0^2, \qquad \rho = \frac{2xH_0}{\sigma B} + \frac{2\mathcal{H}}{B} b_0^2 + \lambda_0 s. \tag{SU3}$$

Thus $d + \rho \sim 2$ follows from $d_0^2 + b_0^2 = 1$, with no inverse of either sphere coordinate. All other bounds are the reviewed SP9–SP11 and BS1 source identities.

Shift the troublesome angular product into the normal block. Set $A_0 = qW$, $A = A_\theta = (U - x)A_0$, and retain R. There is an exact conductor:

$$A(A + uR) = u(U - x)wWZ - x(U - x)A_0^2. \tag{SU4}$$

Consequently the angular/radial part of BS2 is

$$\begin{aligned} K_A A^2 + \chi u A R + (c_G + z K_r) R^2 = 2(A + uR)^2 + 2R^2 - 4hu(U - x) w W Z \\ + 4hx(U - x) A_0^2 + \delta_A A^2 + \delta_\chi u A R + \delta_R R^2, \end{aligned} \tag{SU5}$$

where the three δ coefficients are the infinitesimals in SU2. This step keeps the signed product exactly; it is not an inequality based on the sign of $U - x$.

A complete strict normal block. Introduce the legal separated roots

$$H = \sqrt{h}\, uZ, \qquad S = uwW - hu^2 Z, \qquad N = N_x, \qquad Z_g = uwW = S + u\sqrt{h}\, H.$$

Their exact allocations are

$$\begin{aligned} \Gamma &= -S + \left(\frac{\sigma}{u\sqrt{h}} - u\sqrt{h} \right) H, \\ P_h := \sqrt{h} P_\beta &= (1 - sh) H - \frac{s\sqrt{h}}{u} S, \\ X_e &= N - \sqrt{h} P_h. \end{aligned} \tag{SU6}$$

Every coefficient is bounded; the factors u^{-1} are pure testing units. The full signed packet in BS2 also satisfies exactly

$$\mathscr{P} = L_h P_h^2 + \rho(1 - x/U) Z_g^2 - \rho\sigma(U - x) A_0^2. \tag{SU7}$$

The apparently signed factor $1 - x/U$ is simply a bounded Gram coefficient of residue one. No mixed inverse is used.

Combine the wWZ term in SU5 with every N, Γ, P_h, Z_g term:

$$\begin{aligned} \mathscr{N} = [2(1 + w^2) + 4mw^2] N^2 + 2(2D + 1) U X_e^2 + d\Gamma^2 - 4u\nu N\Gamma \\ + L_h P_h^2 + \rho(1 - x/U) Z_g^2 - 4\sqrt{h}\, \frac{U - x}{u}\, Z_g H. \end{aligned} \tag{SU8}$$

This is an exact equality for the entire indicated normal block. Its coefficients in (N, S, H) are bounded separated expressions. Using SU2 and SU6, its residue is exactly

$$2(1 + u_0^2) N^2 + 4u_0 NS + 2S^2 + 2H^2. \tag{SU9}$$

The N, S determinant is 4, so this residue is strictly positive. The H cross terms cancel: the X_e square cancels the leading $N\Gamma$ coupling to H, and the angular shift in SU5 cancels the remaining SH and hH^2 terms. This is where the sphere sum $d_0^2 + b_0^2 = 1$ is essential.

The remaining angular/radial block is

$$\begin{aligned} \mathscr{A} = 2\{(U - x) A_0 + uR\}^2 + 2R^2 \\ + (4hx - \rho\sigma)(U - x) A_0^2 + \delta_A (U - x)^2 A_0^2 + \delta_\chi u (U - x) A_0 R + \delta_R R^2. \end{aligned} \tag{SU10}$$

Its residue on (A_0, R) has determinant $4u_0^4 > 0$.

Every other original-parent term. The unchanged hard N_y diagonal has residue $2(1 + u_0^2)$. All SP5 terms other than its two Γ terms already have infinitesimal coefficients on (N, N_y, A_0, R), by SP's complete opening-uniform coefficient bounds and $A = (U - x) A_0$. The two Γ terms have a factor qw and become infinitesimal crosses after the bounded allocation SU6. Thus the complete Gram on (N, N_y, S, H, A_0, R) has strictly positive residue, the direct sum of SU9, the N_y entry and SU10.

The remaining exact nonnegative rows are retained once: both centered cap excesses, $4\beta D_w N^2$, and $4bpzU\Gamma^2$. The $d\Gamma^2$ part and $2(2D+1)(uX_e)^2$ have already been used in SU8 and are not retained

a second time. All scalar, hard and first-normal terms of the original SP6 parent are therefore included.

The bounded separated-Gram criterion proves tensor SOS for all real states and both testing signs, including literal $Z = 0$ and $w = 0$. Applying the real identity separately to real and imaginary parts gives all complex states. Neither source sphere endpoint is inverted. The precise remaining C14.1b testing domain has u infinitesimal (with the earlier state and sign admissions still retained).

The focused checker verifies SU4, SU6–SU8 and the whole reassembled SP6 parent, then the two strict residue determinants. It does not replay the unchanged source formulas or claim a whole-source closure.

Chapter 158

R6 with unbounded a: the complete thin layer is admitted

Component `pc82cd677f4b9`
Source: `BASE/inherited_r19/inherited_r18/proofs/weak_contact/r6_thin_layer_all_tests.tex`
SHA256: `c82cd677f4b913167254d5c5f6646ff8dd9447d79bea3aa97569913dca239682`
Inventory use: RT1-RT2 exact source entry and RT10 full kernel determinant only

158.1 R6 with unbounded a: the complete thin layer is admitted

Theorem 158.1. *Retain the entire original physical R6 source intersection, $d = 1$, $\lambda = \beta = 0$, all strict gaps, $n \geq 0$, total scale, and prior exclusions. Thus $t, h, l = 1 - h, j, r$ are positive bounded units and $B, x = aB, y = tB, g$ positive infinitesimals. Assume a unbounded and*

$$n = 0 \quad or \quad 0 < \rho := n/(1-r)^2 \in \mathrm{Inf}\,. \tag{RT1}$$

Then the entire original four-state parent is finite tensor SOS at every independent projective testing pair, with both signs, literal axes and infinities, and real or complex states. No source/testing rate comparison is used. The complementary ρ noninfinitesimal source is not admitted by this theorem.

Exact source entry, including every g opening. Put $s = 1/a$, $w = a(1-r)$, $b = a^2 B$, so $x = sb$, $B = s^2 b$ and $e = 1 - r = sw$. The inherited exact physical identity R10.2 is

$$2r\Delta_* n/s^2 = wQ_* - bK_*, \qquad Q_* \sim 1, \quad K_* \sim h + t, \quad \Delta_* \sim 1. \tag{RT2}$$

Since $g = 1 - w + hx$ is positive infinitesimal, $w \sim 1$. Physicality gives b bounded, $n = O(s^2)$ and $b_0(h_0 + t_0) \leq 1$. On RT1, equality holds in residue:

$$b \sim 1/(h+t), \qquad mg \sim \tfrac{1}{2}, \quad C \sim \frac{h+t}{2l}, \quad A \sim \frac{t}{2h}, \quad D \sim \frac{1}{2(t-l)}. \tag{RT3}$$

All four m, C, A, D have positive real lower bounds; m is unbounded. Every assertion is joint in the independent positive s, g openings. In particular no bound on g/s or s/g is assumed.

Literal $n = 0$ is a physical overlap, with an exact scope match. The unchanged OP4 contact map is

$$k_1 = \sqrt{x}, \quad \eta_1 = t, \quad \zeta_1 = s, \quad r_1 = h, \quad a_1 = l/t, \quad \theta_1 = e, \quad t_1 = w. \tag{RT4}$$

The native gaps are exactly $M_1 = g$ and $J_1 = j$. When $n = 0$, OP4 identifies the full physical C2 parent with the native C1 parent at literal $\beta_1 = 0$ (up to its allowed pure-source normalization to $c = 2$). Here k_1, ζ_1 are positive infinitesimals, η_1, r_1 are positive bounded units, and

$$e^2_{1,\mathrm{scale}} = k_1^2 \eta_1/\zeta_1 = tb \sim \frac{t}{h+t} > 0. \tag{RT5}$$

Thus the exact theorem `c1/bounded_e_all_tests.tex` (BE1–BE6 and its contact join) applies. Its statement explicitly allows arbitrary positive M, J openings and literal beta zero. This proves all tests and states on the whole RT1 literal-zero boundary, rather than taking a positive-n limit.

The positive thin layer: a bounded formal C1 base. Suppose $0 < \rho \in \mathrm{Inf}$. Use the same contact map RT4. The exact TL2 formulas give formal $\beta_1 = -s_n/2$ infinitesimal, $\pi = 1 - s_n > 0$, and $\kappa = (1+2n)/\pi \sim 1$. In particular all paired coefficients $\beta_1 + 1/2$ are positive. RT3–RT5 show that native $\eta_1, r_1, e_{1,\mathrm{scale}}$ are units. Native $\delta = \Delta R/K$ and $E = \eta_1 K/R$ are consequently units too. The formal C1 m_1 is unbounded, but C_1, A_1, D_1 are positive units and $\alpha_1 > 0$. The native gaps $M_1 = g, J_1 = j$ remain positive.

Use the *explicit algebraic* BE5–BE6 and contact-conductor construction for this formal source, not a physical theorem at negative beta. Its bounded remainder is SD4, with SD5 inverse states. It has no M_1 or J_1 inverse; these openings occur only in its four retained positive packets SD3. Its strict contact residue has soft form $8x_c^2 + 8r_{1,0}(r_{1,0} + E_0)T_4$, hard pivots $16r_{1,0}(r_{1,0} + E_0)$ and $16r_{1,0}(r_{1,0} + E_0) + 8E_0^2$, and the hard/soft determinant $128r_{1,0}(r_{1,0} + E_0) > 0$. (The contact-frame letter x_c is not the C2 source x.) The BE proof uses the positive paired beta coefficient, its rational stationary identities, and these strict residues. None requires the standalone formal β_1 to be nonnegative. Its finite conductor therefore supplies the same strict bounded eight-root base for κF_1 on the contact chart. This conclusion is uniform as $M_1 = g \to 0$ because the unbounded paired-m packet is retained.

The divergent event cancels before any coefficient is estimated. Let $\delta L = L_2 - \kappa L_1$ for actual C2 versus formal C1. Use the exact seven TL3 identities, which hold for arbitrary h:

$$\delta m = 0, \qquad \delta C, \delta A, \delta D, \delta d = O(\rho), \quad \delta\alpha = O(nx), \quad \delta\beta + \delta d/2 = -n. \tag{RT6}$$

Here $h, l, t, j, K = 1 + x, h + t$ are units and $\Theta = n/B = \rho w^2/b = O(\rho)$, so these bounds have no g denominator. In particular the actual and scaled formal *entire* m event agree exactly. Its unbounded coefficient is absent from the stress, not discarded or bounded by a guessed relation to s. Keep all four κ-scaled positive packets SD3 unchanged. No centered-A replacement is needed: A is bounded on R6.

Use the exact TL4 contact-zero row, with the native SD5 coordinates $\mathfrak{w} = (W + uY)/D_u$, $A_t = v\mathfrak{w}$, $X = N + pD_uA_t - uZ$, $p = t_1/R$:

$$K_c = \zeta_1 K/Q, \quad \gamma = r/K_c, \qquad \mathcal{N}^2 = nD_uD_v(X + \gamma A_t)^2. \tag{RT7}$$

This is an actual positive separated square and vanishes at the actual contact. The exact full residual stress after retaining it is

$$\begin{aligned} \mathscr{E} = {} & [nD_uD_vX^2 - \mathcal{N}^2] - nD_v(X + uZ)^2 + \delta CD_v(uW - Y)^2 \\ & + (\delta D + \delta d/2)J_{\mathrm{ev}}^2 + \delta\alpha D_u(vY - Z)^2 \\ & + (\delta d/2)D_u\{(vY - Z)^2 - D_vZ^2\} + \delta AD_uD_vY^2, \qquad J_{\mathrm{ev}} = uW + uvX - Y - vZ. \end{aligned} \tag{RT8}$$

All normals and hard terms are included. This differs from TL6 only by keeping the ordinary bounded A difference, rather than centering an unbounded A event. The identity is verified directly at event level in the new focused checker.

The complete stress has the unchanged finite conductor. On $Z = 1$ use the native BE raw roots

$$A_0 = \mathfrak{w}v, \quad B_0 = \delta v, \quad C_0 = \delta Eu\mathfrak{w}, \quad D_0 = \delta^2 k_1^3 E.$$

All δ, E, b, w are units. The exact TL5 expansion of the bracket in RT8 is

$$2n\gamma D_uD_vuA_t - (2n\gamma pD_u + n\gamma^2)D_uD_vA_t^2 - 2n\gamma D_uD_vA_tN. \tag{RT9}$$

The source coquotients

$$\Theta/\rho = w^2/b, \qquad n\gamma^2/\rho = w^2 r^2 Q^2/K^2, \qquad nx/(\rho D_0^2) = w^2 Q^2/K^2$$

are bounded, with unit denominators after $k_1^2 = sb$ and $K_c = sK/Q$. Also $n\gamma/\rho = sw^2 rQ/K$ is bounded. No one of them contains g^{-1}.

At zero normals SD5 gives $q_B = vD_u V_0$ and $Y = u\mathfrak{w} - h_1 v V_0$. Hence $Y, uW \in (B_0, C_0)$; $uW - Y, J_{\mathrm{ev}} \in (B_0)$; $X + u = pD_u A_t$; and $uA_t = B_0 C_0/(\delta^2 E)$. Thus the full scalar $\mathscr{E}/\rho$ belongs to $(A_0, B_0, C_0)^2 + (D_0^2)$ with bounded coefficients. The last generator is needed only for the constant from $\delta\alpha$. Both complete first normals belong to (A_0, B_0, C_0) with bounded coefficients; the remaining hard block is bounded and has normal degree at most two. These assertions follow by substitution into RT8, not just from its contact value. In particular the new $\delta A Y^2$ term has precisely the same raw support and full-normal properties.

Both full parents and every retained positive row have zero value and full first jet at the same contact. Therefore the entire stress and both first normals have the exact vanishing required by CC3–CC6. After clearing the finite positive units, the monic conductor uses no opening inverse and lifts $\mathscr{E}/\rho$ to a bounded Gram on the same eight roots as BE. Multiplication by infinitesimal ρ gives an infinitesimal perturbation of its strict source residue. The finite separated lift proves the full local parent after restoring all retained packets. This is valid for every independent rate of $u, v, \mathfrak{w}, k_1, g$, with no rate comparison.

Finite remote tests and the remaining state charts. For finite tests with $v_0 \neq 0$, use R5.3–R5.5 to retain a centered m excess and leave a bounded remainder with $\kappa_m = (mg)_0 = 1/2$. At source residue $e = 0$, the determinant R6.4 of the complete restriction to the original m-event kernel is

$$\frac{D_u^2 D_v^2 (h+t)}{8hl(t-l)} \{V^2 + t(h+t)U^2 V + hUV^2\} > 0. \tag{RT10}$$

The restriction to $Z = 0$ is positive: on that kernel the positive A, C packets and signed core force $Y = X = W = 0$ since $v_0 \neq 0$. A sufficiently large positive real cap makes the full bounded residue strict; its centered correction vanishes on the same kernel. The cap exists below unbounded actual m and depends only on real testing residues. Thus all such finite remote pairs are admitted.

The finite cases with v infinitesimal use the actual-parent BE state entry. All required positive real lower bounds for m, C, A, D are RT3; retain the actual nonnegative n, α packets. If Z is infinitesimal and normalized W a unit, the exact BE4 full-state frame is strict using clipped m, A. If u and $\mathfrak{w}$ are units, the core's zero conditions force nonzero Y, controlled by A. Otherwise the negative-core branch has Z a unit; normalize $Z = 1$. If u is a unit and $\mathfrak{w}$ infinitesimal, BE5 is strict on its raw frame and the bounded RT8 stress is an infinitesimal perturbation there. If u is infinitesimal and $\mathfrak{w}$ a unit, use the exact BE6 relations $B_0 = (\delta/\mathfrak{w})A_0$ and $C_0 = (\delta/\mathfrak{w})A_0 - x_c$; their inverse is purely testing, and the same stress is bounded on the strict BE6 frame. The sole remaining case is precisely the contact chart just proved. The exact SD5 core sign identity supplies its positive gauges; the nonnegative core and literal axes have direct original-event exits.

Every projective endpoint. For v unbounded and u bounded, the complete R6.6 kernel formula remains strict with $n_0 = 0$, since its first coefficient is $1/2$. The R5 centered-m correction vanishes on that kernel, so a positive real cap gives the full strict remainder. The double-infinity formula R6.7 likewise remains strict: its two relevant determinants become $D_0/2$ and $A_0(K + 1/2) > 0$.

For u unbounded and v a unit, R6.5 with $n_0 = 0$ is strict because $C_0 W^2$, $A_0 Y^2$ and $(D_0 + 1/2)(W + vX)^2$ have common kernel zero. For u unbounded and v infinitesimal, use the exact BE $(\infty, 0)$ chart: retain every original event except positive real caps of C, A, set $p = 1/u$, and use $(pX + Z, W, Y, vX)$. The complete selected polynomial has bounded coefficients in p, v and strict residue

$$\tfrac{1}{2}(pX + Z)^2 + \tfrac{1}{2}(W + vX)^2 + C_* W^2 + A_* Y^2.$$

This is an exact whole-core allocation; the focused checker verifies its expression and residue. Every literal reciprocal zero is a polynomial value. Undoing homogeneous factors uses pure-testing squares only.

Scope and restoration. The preceding cases exhaust all independent tests and normalized states on RT1. Every retained packet is restored, both full normals and every hard direction are present, and the real proof applies separately to real and imaginary state parts. This proves the stated thin-layer all-tests admission. It does not infer nonemptiness of any residual intersection and does not close the noninfinitesimal-ρ complement of unbounded-a R6.

Chapter 159

Weak-C2 R3 with a positive-unit $1-r$ is admitted

Component `p1c9902e9a6f2`
Source: `BASE/inherited_r19/inherited_r18/proofs/weak_source_ends/r3_e_unit_all_tests.tex`
SHA256: `1c9902e9a6f2acbe91d8c5547d1584b76b4296a87582e442e9c9dcac55fbfdd8`
Inventory use: R3E1-R3E11 and projective join

159.1 Weak-C2 R3 with a positive-unit $1-r$ is admitted

Theorem 159.1. *Keep the entire original physical stationary weak-C2 source, $d=1$, $\lambda_{\mathrm{event}}=\beta=0$, every strict gap and nonnegative coefficient, and every original residual intersection. Suppose $B,z,y,l=1-h$ are positive infinitesimals, $t=y/B$ a positive bounded unit, and $e=1-r$ a positive bounded unit. Then the full original parent is a finite tensor SOS for every independent projective testing pair and every real or complex state. These are precisely the positive-unit-e sources of R3. The remaining e-infinitesimal part of R3 is not asserted here.*

Source estimates including all simultaneous openings. Put $a=z/B$, $g=1-ae+haB$ and $j=1+aBr-l/t$. The positivity of g and the unit lower bound for e imply that a is bounded. Moreover $h\sim 1$ and $j\sim 1$. In the following formulas variables on the right denote their real residues, $e=1-r>0$ and $g=1-ae\geq 0$. Direct joint regular reduction of the unchanged WE1 formulas gives

$$(nr)_0=\frac{e}{2(1+a)},\qquad (mg)_0=\frac{a^2e}{2(1+a)},\qquad A_0=\frac{t(1+ar)}{2(1+a)},\qquad D_0=\frac{1+ar}{2t(1+a)},\tag{R3E1}$$

$$(Cl)_0=\frac{(1+ar)(1+t)}{2(1+a)},\qquad \alpha_{*,0}:=(\alpha/B^2)_0=\frac{(1+ar)^2(1+t)}{2(1+a)}.\tag{R3E2}$$

Thus A,D,Cl,α_* are positive bounded units, C is unbounded, and n has a positive real lower bound, even when r is infinitesimal and n is unbounded. In particular literal $n=0$ is impossible on this source domain, by the positive first formula in R3E1; no coefficient boundary was discarded. Write $N=n+1/2$, so N^{-1} is bounded. If g is a unit, then $\mu=m/a^2$ is a positive unit with

$$\mu_0=\frac{e}{2g(1+a)}.\tag{R3E3}$$

If g is infinitesimal, a is a unit, m is unbounded and mg is a positive unit. These joint statements use only the displayed products: the WE1 denominators after clearing r,g,l have unit factors $t,h,1-hB,\Delta$ with $\Delta_0=1+a$. There is no comparison among the B,l,r,g openings. The original positive-infinitesimal total-scale condition follows already from C unbounded.

The full local matrix with its divergent C pivot retained. Use the exact complete WA3–WA8 parent identity in the roots

$$\xi = X + ruZ, \quad \eta = Y - huW, \quad R_1 = uW - vZ/t, \quad R_2' = avW - uZ, \quad T = uvW - BZ.$$

Normalize its first two rows by $\widehat{\xi} = \sqrt{N}\xi$ and $\widehat{\eta} = \sqrt{C}\eta$. First suppose g is a unit. The complete resulting five-root matrix has bounded pure-source polynomial coefficients in u, v. Here is the coefficient check, including the potentially singular source openings. The hard entries use n/N, C/C, and the bounded coefficients A, D, m, α, together with bounded $N^{-1/2}$ and infinitesimal $C^{-1/2}$. All scalar occurrences of n are through nr^2, and all first-normal occurrences are through nr, both bounded. All non-hard occurrences of C are through Cl or Cl^2, both bounded. Thus the entire η normal row becomes infinitesimal after division by $\sqrt{C}$. This includes the full WA7 terms, not merely their zero-angle values.

The WE5 scaled-R_2' formulas bound every coefficient through $a \to 0$ except the apparently singular

$$Q_* = \frac{b_2 - \alpha_*}{a}, \qquad b_2 = 2a^2\mu h + Cl^2 + Ah^2 + (B^2\alpha_* + 1/2)h^2. \tag{R3E4}$$

The inherited exact WE3 numerator vanishes at $a = 0$ for arbitrary B, h, r, t. Its quotient has a unit denominator here too: the l denominator cancels in Cl^2 before division, and the remaining factors $g, h, t, j, 1 - hB, \Delta$ are units on this g-unit chart. Consequently Q_* is bounded jointly through $a, l, r \to 0$. This bounds the entire scaled scalar block and both normal rows, without any a/l or l/a hypothesis.

At $u = v = 0$ the full residue has separate pivots $1, A_0, \alpha_{*,0}$ on $\widehat{\eta}, R_1, T$. Its $\widehat{\xi}, R_2'$ block is

$$\begin{pmatrix} 1 + a_0^2\mu_0\nu & -a_0\mu_0\sqrt{\nu} \\ -a_0\mu_0\sqrt{\nu} & \mu_0 \end{pmatrix}, \qquad \nu = (N^{-1})_0 \geq 0, \qquad \det = \mu_0 > 0. \tag{R3E5}$$

This includes a infinitesimal, r infinitesimal, and their independent simultaneous opening. The R_1 pivot is A_0 because $Cl^2 \sim 0$ and $h \sim 1$; it does not require a new soft root.

If g is infinitesimal, use the unscaled $R_2 = R_2'/a$, as a is a unit. Apply the exact R5.2–R5.4 centered-m identity, keeping

$$(m - K)R_{mc}^2, \qquad R_{mc} = -\xi + uv\eta + (1 - g)R_2 + huT,$$

for a sufficiently large positive real K. In the remainder replace only the formal matrix coefficient m by K and add τH_m, where $\tau = (m - K)g$ is bounded and H_m is the exact R5.4 matrix. The remaining source coefficients, especially α_*, stay at their actual values. This is a whole-matrix identity, not a physical-source alteration. The same coefficient bounds hold; all normal entries divided by $\sqrt{C}$ vanish. The three separate residue pivots remain $1, A_0, \alpha_{*,0}$; the last block becomes

$$\begin{pmatrix} 1 + K\nu & -(K + \tau_0)\sqrt{\nu} \\ -(K + \tau_0)\sqrt{\nu} & K + 2\tau_0 \end{pmatrix}, \qquad \det = K + 2\tau_0 - \tau_0^2\nu. \tag{R3E6}$$

Since $0 \leq \nu \leq 2$, any real $K > 2\tau_0^2 + 1$ makes it strict. Thus WA10 applies to the entire local parent in both source cases, for all independent infinitesimal or zero u, v, either sign and all states. Restore the exact retained square in the g-small case.

Every finite remote testing chart. If both finite testing residues are nonzero, use the inherited arbitrary-coefficient-scale noncommuting-residue theorem. It remains to treat one nonzero finite angle residue. In these charts select positive real lower bounds for original positive event coefficients when indicated and retain their exact positive excess packets; no stationary-contact assertion is made for the selected remainder.

Use the actual original C event with

$$J_C = \sqrt{C}\,(uW - Y), \qquad d_C = C^{-1/2} \in \mathrm{Inf}\,. \tag{R3E7}$$

If u is a bounded unit and v infinitesimal or zero, clip n, A, D at positive real lower bounds and keep their positive excesses and the whole original m, α packets. Solve $W = (Y + d_C J_C)/u$. The selected complete remainder has bounded coefficients in (J_C, X, Y, Z) and residue

$$J_C^2 + n_*(1+U)X^2 + A_*(1+U)Y^2 + \tfrac{1}{2}(X+uZ)^2 \succ 0, \tag{R3E8}$$

where u on the right is its nonzero real residue and $U = u^2$. The D event has zero residue and causes no difficulty.

If u is infinitesimal or zero, v a bounded unit and a a unit, then m has a positive real lower bound (also when g is small). Clip n, m, A, D at positive bounds, retain all excesses and α, and solve $Y = uW - d_C J_C$. The complete selected residue is

$$(1+V)J_C^2 + (n_* + 1/2)(1+V)X^2 + m_*(vW - X)^2 + D_* V Z^2 \succ 0, \quad V = v_0^2 > 0. \tag{R3E9}$$

This retains every state direction and the whole signed core.

Finally suppose a is infinitesimal in the latter testing chart. Then $g \sim 1$, μ is a positive unit, and R3E1 gives $A_0 = t_0/2$, $D_0 = 1/(2t_0)$. Keep the additional legal roots $S = aW$, $R = uW$ and use $Y = R - d_C J_C$. Clip only n, retain its excess and α, and retain the actual bounded A, D, μ. The m event is exactly

$$m(vW - X + uvY - uZ)^2 = \mu(vS - aX + auvY - auZ)^2. \tag{R3E10}$$

The entire bounded matrix on (J_C, S, X, R, Z) has separate positive pivots $1+V$, $\mu_0 V$ and $(n_* + 1/2)(1+V)$. Its remaining R, Z block is

$$\begin{pmatrix} [t_0(1+V)+V]/2 & -v_0/2 \\ -v_0/2 & V/(2t_0) \end{pmatrix}, \qquad \det = \frac{V^2(t_0+1)}{4t_0} > 0. \tag{R3E11}$$

The dependent roots R, S are allowed in a positive Gram; no inverse of u or a is introduced. WA10 at the real testing residues establishes every finite remote chart above.

Projective join and exact scope. The source estimates give positive real lower bounds for n, A, C, D. Therefore the complete selected-event endpoint polynomials R4E14, R4E15 and R4E16 apply verbatim: clip these four original event coefficients, retain all their excesses and the entire m, α packets, and use the original degree-$(2,2)$ homogeneous parent. Those exact identities cover respectively one infinite Alice angle, one infinite Bob angle, and both infinite angles, including the zero values in the other projective chart. Their strict Grams do not depend on any R4-specific source relation. Thus every finite and infinite testing pair is covered. Polynomial axes, both signs and the real/imaginary state split give literal zeros and complex states. All inverses and radicals used are pure-source or pure-test.

This admits exactly R3 with $1 - r$ a positive bounded unit. It neither claims the remaining $1 - r$-infinitesimal sources empty nor changes the historical weak-source partition. No numerical table is a premise. The focused checker verifies the new source residues, full local residue blocks and complete finite remote residues; unchanged WA, centered-m, and homogeneous endpoint identities are imported.

Why the existing discrete symmetry does not remove R1–R3. In the proved cyclic order $(\lambda, D, C, A, \alpha, m, \beta, n)$, the stabilizer of the omitted pair $\{\lambda, \beta\} = \{0, 6\}$ inside the CHSH dihedral action consists of the identity and $i \mapsto 6 - i$. The latter is exactly

$$(u, v; W, X, Y, Z; n, m, C, A, D, \alpha) \longmapsto (1/v, 1/u; Z, X, Y, W; n, D, \alpha, A, m, C).$$

It exchanges strong C/weak α with the opposite directed orientation. Returning to the chosen weak orientation restores the original source, so this known discrete action supplies no additional within-orientation routing of R1–R3. This statement concerns only the established dihedral action, not all conceivable analytical changes of variables.

Chapter 160

The source, the original polynomial, and the proof language

Component p472f434581f3
Source: BASE/inherited_r19/parent_reference.tex
SHA256: 472f434581f355d0b0837269801d150274e69df6827228f4532a8bae95ce927d
Inventory use: P1-P8 and SP6: actual source, complete parent, coefficient definitions, strict bounded-Gram lemma | P1–P8/SP6 and strict bounded-Gram paragraph | Original parent/source and strict separated bounded-Gram lemma

160.1 The source, the original polynomial, and the proof language

Let K_c, K_t be independent real closed extensions of $\mathbb{R}$. Source parameters lie in K_c, and angles and state coordinates in K_t. The assertion is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$, for each such pair of configurations. No order or general inverse is imposed on this tensor ring. A bounded quantity has a finite real residue; a positive unit is bounded above and below by positive reals. All asymptotic statements about a source or a test are made inside its own field.

The original source. Keep the physical stationary C1 source, its strict gaps, nonnegative event weights and all original residual intersections. Normalize its positive coefficient d to 1. On C14.2, $x = k^2 > 0$ is infinitesimal, $L = x\zeta$ is a positive bounded unit, and η is unbounded. Set

$$\sigma = \eta^{-1},\ z = L^{-1},\ b = Lr,\ Q = 1 - b, \theta = 1 - h,\ e = xh/\sigma,\ E = 1 - e,\ K = 1 + x.$$

The source has $0 < r, h, b < 1$, $r = bz$, and its original strict gaps. The following exact formulas are NC1–NC4 in these coordinates:

$$\begin{aligned}
J &= 1 + xh - \sigma(1 - r), \qquad M = 1 + xr - xz\theta,\\
P &= \sigma + K + \sigma z(1 + bx + 2xh) + \sigma b x^2 h z^2 + x^2 h\theta z,\\
p = Ar &= \frac{z(1 + xh)E + (1 - r)(1 + xhz)}{2P},\\
l = \alpha z Q &= \frac{(1 + \sigma r)\{e(1 + xr)(1 + \sigma z) + ME\}}{2P},\\
C &= \frac{Kz(1 + \sigma r)E}{2(1 - r)P},\\
D &= \frac{K(1 + \sigma z)(1 + \sigma r) - JP}{2JP},\\
\beta\theta &= \frac{K(1 + \sigma r)(1 + xhz) - \theta P}{2P},\\
\kappa &= \frac{K(1 + \sigma r)(1 + \sigma z)}{2MP}, \qquad m = \kappa x^2 z\frac{h}{\sigma}.
\end{aligned} \tag{P1}$$

Thus $a = Az = p/b$ and $\ell = \alpha z = l/Q$; the symbol p here denotes a source product, not a testing angle. All original gaps are positive, while D, β may be zero.

The shared source sphere. Put $H_0 = 1 + z - r = 1 + zQ$, $B = P + xz(J - \sigma)$, $R_s = 1 + xr$, and

$$\mathcal{H} = R_s H_0 + z\{(1-r)xh(2+xr) - e[2 + x(1 + h + r\theta)] + Ke^2\}.$$

The exact source identities BE3–BE8 are

$$\begin{gathered} 2\theta P\beta = hB - \sigma H_0, \qquad 2P(JD + x\theta\beta) = \sigma^2\mathcal{H}, \\ d_0^2 = \frac{2PJD}{\sigma^2\mathcal{H}}, \qquad b_0^2 = \frac{2Px\theta\beta}{\sigma^2\mathcal{H}}, \qquad d_0, b_0 \geq 0, \quad d_0^2 + b_0^2 = 1, \\ H = \frac{h}{\sigma} = \frac{H_0}{B} + \frac{\sigma}{x}\frac{\mathcal{H}}{B}b_0^2. \end{gathered} \tag{P2}$$

They follow by clearing the positive source denominators in P1. In this branch $e = O(x + \sigma)$ is infinitesimal, $M, P, J, B \sim 1$, $\mathcal{H} \sim H_0$, and $\kappa \sim 1/2$. To obtain the only non-immediate bound $e = O(x + \sigma)$, expand the original $D \geq 0$ numerator in native $\eta = 1/\sigma$ coordinates as in BE1:

$$e\{K + x\theta(1 + xh)/L\} \leq x(b_1/L) + \sigma(a_0/L),$$

where

$$\begin{aligned} b_1 = {} & -Lhrx - Lh - Lr + L - h^2rx^2 + h^2rx - 3h^2x \\ & - 2hrx + hx - 3h + 1, \\ a_0 = {} & Kr + (1-r)\{1 + L + 2xh + rxL + rx^2h\}. \end{aligned}$$

Both right-hand source ratios are bounded and the left bracket is at least one. P1 then gives the opening-uniform comparisons

$$C(1-r)/z \sim \tfrac{1}{2}, \quad p \sim H_0/2, \quad l \sim \tfrac{1}{2}, \qquad D = O(\sigma^2), \quad x\beta\theta = O(\sigma^2). \tag{P3}$$

In particular C, A, α have positive real lower bounds, although their upper bounds may diverge. There is no inverse of a sphere coordinate and no lower bound on D or β.

The entire original polynomial. On the reciprocal Bob chart $w = 1/v$, put $D_u = 1 + u^2$ and $D_w = 1 + w^2$. For a real state (W, X, Y, Z) define

$$\begin{aligned} M_w &= W - wX + uY - uwZ, \quad R_w = uW - Y, \\ J_w &= uwW + uX - wY - Z, \quad B_w = X + uZ, \\ K_w &= J_w^2 + D_u(Y - wZ)^2 + D_w B_w^2 - D_u D_w Z^2. \end{aligned}$$

The target is exactly

$$\begin{aligned} F_w = {} & 4mM_w^2 + 4CD_wR_w^2 + 4AD_uD_wY^2 + 4DJ_w^2 \\ & + 4\alpha D_u(Y - wZ)^2 + 4\beta D_w B_w^2 + 2K_w. \end{aligned} \tag{P4}$$

This expression, not a restriction to a contact line, is preserved throughout the proof. The source and testing angles are never identified.

Why a strict bounded Gram suffices. The only lifting lemma used below is the established separated bounded-Gram lemma: a finite symmetric matrix whose entries are finite sums of bounded separated factors, and whose real residue is strictly positive definite, defines a tensor SOS. Its elementary mechanism is worth recording. For a source infinitesimal s and bounded testing t, choose positive real a, ε with $|t| < a$ and $|s| < \varepsilon/a$. Then exactly

$$\varepsilon \pm st = \tfrac{1}{2}\{(\varepsilon/a + s)(a \pm t) + (\varepsilon/a - s)(a \mp t)\}.$$

Each summand is a product of nonnegative pure-field factors and has a separated square root. The same argument interchanges the fields. A finite real diagonal budget absorbs all infinitesimal matrix entries into a strictly positive real Gram. This constructs finite SOS without a mixed Cholesky factor or inverse. Every later use explicitly supplies the bounded coefficients and the positive residue; symbolic checks alone do not establish this hypothesis.

160.2 The exact complete normal identity used in the closure

Here is the complete SP6 identity, with the centered A, α excesses restored, so that no cap term is implicit. Let

$$\begin{gathered} q=\sqrt{z},\quad t=u^2,\ \xi=w^2,\quad g=h-Q,\quad b_\theta=\beta\theta,\quad a_\beta=\beta\theta^2,\\ N_x=X+uhZ,\quad N_y=(Y-bwZ)/q,\\ \Gamma=\sigma Z-uwW,\quad R=wZ/q-uqW,\quad A_\theta=q(t-x)W,\\ P_\beta=uZ-(x/\sigma)wW,\quad X_e=X+ewW,\quad V=uw(t-x)W. \end{gathered} \tag{P5}$$

The source coefficients are

$$\begin{aligned} \nu &= (2D-2b_\theta+h)/\sigma,\\ \lambda &= (8Dh+4a_\beta+2h^2)/\sigma-4bp\sigma z=2H+\lambda_0,\\ \lambda_0 &= 4D(1+h)/\sigma-2\theta\nu-4bp\sigma z,\\ \chi &= 8(pb+lQ)-4h+4eh(2D+1)-8Dh\\ &\quad +4z\{4Dbh+4pb\sigma+(1+2m)g^2+2a_\beta\},\\ B_x &= 8\kappa e,\qquad A_y=8(p-l)+4z\{(1+2m)g+2Dh+2p\sigma\},\\ R_x &= -4\{2Db+(1+2m)g-2b_\theta\},\\ R_\gamma &= -4\{2(C+D)(r-1)+2p+r-1+2pt\},\\ R_y &= 4\{t[2Dh+2p\sigma+(1+2m)g]+b(2C+2D+1)(\xi+\sigma z)\\ &\qquad +(2C+1)(b-\sigma)+2D(1-\sigma)+2p\sigma\}. \end{aligned} \tag{P6}$$

Finally put

$$\begin{aligned} k_0 &= 2b\{(2C+1)b+4D+4p\sigma+2(2C+2D+1)\sigma(r-1)\},\\ k_1 &= 2(1+2m)g^2+8Dhb+8pb\sigma+4a_\beta,\\ k_2 &= 2b^2(2C+2D+1),\qquad K_r=k_0+k_1t+k_2\xi,\\ H_y &= 4D_u(aD_w+\ell)+4mtz+4CzD_w+(4D+2)\xi z+2D_uz,\\ c_G &= 4D_u(pbD_w+lQ),\qquad c_N=8D_u(pD_w-l). \end{aligned} \tag{P7}$$

Then $F_w=4\beta D_wN_x^2+G$, with the full parent core

$$\begin{aligned} G &= 2(2D+1)(uX_e)^2+[2(1+\xi)+4m\xi]N_x^2+H_yN_y^2\\ &\quad +(4D/\sigma^2+4bpzt)\Gamma^2+(c_G+zK_r)R^2-4u\nu N_x\Gamma\\ &\quad -4uqw(2m+2D+1)N_xN_y+qwR_\gamma(N_y+bR)\Gamma\\ &\quad +uqwR_xN_xR+(c_N+zR_y)N_yR+qwB_xN_xA_\theta+uA_yN_yA_\theta\\ &\quad +4\kappa HA_\theta^2+\lambda u(\Gamma P_\beta+A_\theta R)+\chi uA_\theta R. \end{aligned} \tag{SP6}$$

This is an exact polynomial identity after clearing pure-source denominators in P1. It is the imported JR5/JB1 identity followed by the explicitly checked SP1–SP6 transformation. P4–P7 specify every entry for direct hand reproduction; the complete earlier derivations and their checkers are preserved in the archive. The two crucial conductors are

$$\begin{aligned} A_\theta^2+u\Gamma P_\beta+uA_\theta R &= \sigma P_\beta^2+(x/\sigma)(t-x)(\xi-\sigma z)W^2,\\ V^2 &= \sigma tA_\theta^2+u\sigma(t-x)A_\theta R-(t-x)V\Gamma. \end{aligned} \tag{P8}$$

For the bounded-M_β case, the generic pre-extraction JR coefficients of $\Gamma^2, A_\theta^2, \Gamma V$ are, respectively,

$$\begin{aligned} Z_0 &= \{4D + 2t(4Dh + 2a_\beta + h^2) + 2t^2h^2(2D+1)\}/\sigma^2, \\ A_0 &= 4\kappa H + 2h^2(2D+1)t/\sigma, \\ S &= 2\{h^2(2D+1)(t+x) + 4Dh - 2bp\sigma^2 z + 2a_\beta + h^2\}/\sigma^2. \end{aligned}$$

Its other changed terms are $H_x N_x^2 + Z_x N_x \Gamma + u A_x N_x V + u\sigma B_g A_\theta R$, where

$$\begin{aligned} H_x &= 2(1 + t + \xi) + 4Dt + 4m\xi, \quad A_x = -4h(2D+1)/\sigma, \\ Z_x &= -4u\{2D - 2b_\theta + h + th(2D+1)\}/\sigma, \\ \sigma B_g &= 2h^2(2D+1)(t-x)/\sigma + \lambda + \chi. \end{aligned}$$

These replace the corresponding extracted terms in SP6; all other normal and radial terms are unchanged.

The needed pole cancellations. Directly from P1–P2,

$$\nu = 2D/\sigma + \{H_0 - ez(J - \sigma)\}/P.$$

To bound $\alpha_\kappa = 2H(2\kappa - 1)$, set $N_\kappa = K(1 + \sigma r)(1 + \sigma z) - MP$ and

$$F_x = N_\kappa|_{\sigma=0}/x, \qquad F_\sigma = (N_\kappa - N_\kappa|_{\sigma=0})/\sigma.$$

These quotients are polynomials in bounded source letters, and $\alpha_\kappa = 2(hF_\sigma + eF_x)/(MP)$. This establishes boundedness through every opening before taking residues. The divergent C block is treated by CT7–CT7a. Equivalently, for $J_C = N_y + bR$ it is exactly

$$4CzD_w J_C^2 + 8C(r-1)J_C(z\sigma R - qw\Gamma).$$

The optional cap $C_* = C/(1 + C)$ therefore changes the entire core by precisely $4zD_w(C - C_*)J_C^2 - 8(C - C_*)(1 - r)J_C(z\sigma R - qw\Gamma)$, as used in HG and HP. No omitted matrix entry is hidden by that cap.

Chapter 161

C14.2: the complete unbounded-H, unbounded-W/Z chart

Component `p831dd6874dcf`
Source: `BASE/inherited_r19/proofs/c142_contact/h_large_toric_completion.tex`
SHA256: `831dd6874dcfbcf24dd8d32e3dcc2f44d928006b0dfc12ee90eb7fe6c22ab9a6`
Inventory use: CT7-CT7a exact radial identities only; no unit-z conductor imported | CT1–CT13 and complete normal restoration | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

161.1 C14.2: the complete unbounded-H, unbounded-W/Z chart

Proposition 161.1. *Keep the exact physical C14.2 source, $d = 1$, all original strict gaps and residual intersections, and $D, \beta \geq 0$, including their literal zero faces. Suppose $H = h/\sigma$ is unbounded. The entire parent is tensor SOS on the remaining double-small-angle chart with W/Z unbounded. This includes every state ratio on this chart: the extreme and unmatched ratios are admitted by CS1–CS7, and the matched positive-unit ratio is admitted below. No mixed inverse or source/testing rate comparison occurs.*

Exact source and state data. Use BE2–BE9 and the entire SP6 identity, restoring the original centered a, ℓ excesses. Thus $q^2 = z = L^{-1}$ is a bounded positive source unit, $b = Lr$, $Q = 1 - b$, $p = ab$, $l = \ell Q$, $a = Az$, $\ell = \alpha z$. All original hard and radial normals remain present. Use the actual SP3 coefficient

$$\lambda = 2H + \lambda_0, \qquad K_A = 4\kappa H - \lambda = \alpha_\kappa - \lambda_0. \tag{CT1}$$

SP9–SP10 give $\lambda_0, \alpha_\kappa \in \mathcal{O}_c$. Consequently $\lambda > 0$ is unbounded, $\lambda\sigma \sim 2h$, $\lambda x \in \mathrm{Inf}_c$, and $\rho = \lambda x/\sigma$ is bounded: use BE7 and $x/\sigma \in \mathrm{Inf}_c$ on H unbounded. There is no assumption that ρ is a unit.

After CS7 and the sign gauges, take $W, Z, u, w > 0$ and define solely in the testing field

$$f = Z/W \in \mathrm{Inf}_{t,>0}, \quad c = \frac{u}{wf} = \frac{uW}{wZ} > 0 \text{ a bounded unit}, \quad F = fc^{1/3}, \quad V = wc^{1/3}.$$

Both F, V are infinitesimal. Set

$$k_c = (zc)^{1/3}, \quad F_0 = \sqrt{x/\sigma}\, z^{1/6}, \quad V_0 = \sqrt{\sigma}\, z^{1/6}, \quad g_0 = \sqrt{\lambda x}\, z^{1/6}. \tag{CT2}$$

Here F_0, V_0 are infinitesimal pure-source elements, and k_c is a bounded separated unit. Every fractional power of a displayed separated product means the product of the two pure-field powers.

The scalar packet and its strict circuit rows. The complete SP6 angular packet is exactly

$$K_A A_\theta^2 + S + \chi u A_\theta R, \qquad S = \lambda\sigma P_\beta^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2. \tag{CT3}$$

This follows from SP7 and does not discard a scalar or normal term. Put

$$A_1 = \sqrt{\lambda\sigma}\, VF^2W, \quad A_2 = \sqrt{\rho}\, V^2FW, \quad A_3 = \sqrt{\lambda}\, xqW, \quad G = g_0 VFW, \quad d_i = A_i - G.$$

All four quantities are nonnegative separated monomials and $A_1A_2A_3 = G^3$. Let $r_c = k_c^{-1}$. The exact identity is

$$\begin{aligned} S + \lambda x R^2 = \sum_{i=1}^{3} A_i^2 - 3G^2 \\ + \lambda x z\{uW(r_c - 1)r_c(r_c + 1)\}^2 + 2\lambda x z\{uW(r_c - 1)r_c\}^2. \end{aligned} \tag{CT4}$$

To see the finite SOS and its strict d_i rows, let $p_i = A_i^{1/3}$, with separated cube roots. Exactly

$$\sum_i A_i^2 - 3G^2 = \sum_i d_i^2 + \sum_{j=1}^{3} \sum_{1 \le i < k \le 3} \{\sqrt{Gp_j}(p_i - p_k)\}^2. \tag{CT5}$$

The square roots also separate. Thus CT4 supplies three diagonal pivots equal to one on d_1, d_2, d_3, and nine further squares. The two additional CT4 squares are retained whole. Its sole cost $-\lambda x R^2$ has an infinitesimal pure-source coefficient.

The radial row without a mixed inverse. Define the legal root

$$J_c = (k_c - 1)VFW.$$

The exact radial relation is

$$R = -\frac{k_c^2 + k_c + 1}{qc^{2/3}} J_c. \tag{CT6}$$

Only the displayed bounded separated coefficient is used; the polynomial $k_c^2 + k_c + 1$ is never inverted. Restoring the full a, ℓ weights, their complete occurrence in the radial normal block is

$$4D_u\{aD_w(N_y + bR)^2 + \ell(N_y - QR)^2\}.$$

The full constant radial base, including the other source terms, is

$$M = [4a + 2z(2C + 1)](N_y + bR)^2 + 4\ell(N_y - QR)^2. \tag{CT7}$$

Retain the positive u^2, w^2 excesses of the preceding a, ℓ packet and the exact $4Czw^2(N_y + bR)^2$ packet. Clip the two pure-source coefficients in CT7 at positive real lower bounds, retaining their positive square excesses. After CT6 the clipped Gram on (N_y, J_c) is bounded and strictly positive at residue: the two rows differ by R, and the coefficient of J_c in CT6 has nonzero real residue. This remains true at $b = 0$ or $Q = 0$ in residue. For an explicit audit, put $t = u^2$, $\xi = w^2$, $g = h - Q$, $a_\beta = \beta\theta^2$ and

$$k_1 = 2(1 + 2m)g^2 + 8Dhb + 8pb\sigma + 4a_\beta.$$

After the displayed positive radial packets have been extracted, the exact three remaining coefficients of N_y^2, R^2, N_yR are

$$\begin{aligned} E_{yy} &= 2z\{(1 + 2m)t + (1 + 2D)\xi\}, \\ E_{RR} &= z\{8bD + 8bp\sigma + 4b\sigma(2C + 2D + 1)(r - 1) + k_1 t + 2b^2(2D + 1)\xi\}, \\ E_{yR} &= 4z\{t[2Dh + 2p\sigma + (1 + 2m)g] + b(2D + 1)\xi + \sigma(2C + 1)(r - 1) \\ &\qquad + 2D[1 + \sigma(r - 1)] + 2p\sigma\}. \end{aligned} \tag{CT7a}$$

These are bounded infinitesimal coefficients. The only apparent C pole is its full LC2 correction, whose coefficient is $C(1 - r)$ times σ or w and is bounded by BE9. No entrywise cap of the original radial matrix is made.

The two sphere-weighted Gamma conductors. The monomial relations give exactly

$$(F - F_0)G = F_0 d_1, \qquad (V - V_0)G = V_0 d_2. \tag{CT8}$$

Write b_0, d_0 for the BE6 sphere coordinates, not the source parameter b. The pure-source quantity

$$B_\Gamma = b_0 V_0 / g_0 = b_0 \sqrt{\sigma/(\lambda x)} \tag{CT9}$$

is bounded. Indeed BE7 gives $xhB = \sigma x H_0 + \sigma^2 \mathcal{H} b_0^2$, and $\lambda\sigma \sim 2h$; this proves the bound directly and never divides by b_0. With $j = c^{1/3}$ the exact conductors are

$$\begin{aligned} wb_0\Gamma = & -\frac{B_\Gamma(V_0 + V)}{k_c j} d_2 - \frac{b_0 V^2}{k_c j} J_c, \\ ub_0\Gamma = & -\frac{B_\Gamma j F(V_0 + V)}{k_c} d_2 - \frac{b_0 j V^2 F}{k_c} J_c. \end{aligned} \tag{CT10}$$

Every coefficient in these two arrays is infinitesimal and bounded. Let $\Gamma_d = d_0\Gamma$. The original diagonal is exactly

$$\frac{4D}{\sigma^2}\Gamma^2 = \frac{2\mathcal{H}}{PJ}\Gamma_d^2, \tag{CT11}$$

with a positive-unit pure-source coefficient. In CT11 the J in the denominator is the original source J of BE2. Finally

$$u\Gamma = d_0 u\Gamma_d + b_0(ub_0\Gamma), \quad w\Gamma = d_0 w\Gamma_d + b_0(wb_0\Gamma). \tag{CT12}$$

Thus every first normal carrying $u\Gamma$ or $w\Gamma$ has an infinitesimal bounded allocation on (Γ_d, d_2, J_c). This includes $d_0 = 0$ and $b_0 = 0$ algebraically, with no false strict pivot on an unweighted Γ.

The angular row at either h type. If h is infinitesimal, SP10 gives $\alpha_\kappa \in \mathrm{Inf}_c$ and SP9 gives $\lambda_0 \sim -2H_0$; hence $K_A \sim 2H_0 > 0$. Retain $K_A A_\theta^2$ as a strict additional row. If h is a positive bounded unit, $\lambda\sigma$ is a unit and the following exact conductor replaces that row:

$$A_\theta = \frac{qcw}{\sqrt{\lambda\sigma}} d_1 + \frac{k_c^2}{\sqrt{\lambda}} d_2 - \frac{1}{\sqrt{\lambda}} d_3 + (k_c + 1)\sqrt{x}\, z^{1/6} J_c. \tag{CT13}$$

All four coefficients are infinitesimal. Since K_A is bounded, its entire possibly signed square is then an infinitesimal bounded perturbation of the circuit/radial frame. CT13 uses CT8 and $(k_c - 1)G = g_0 J_c$, not a division by a contact difference.

Full-parent reassembly. The strict bounded principal frame consists of N_x, the two raw radial roots (N_y, J_c) with the clipped CT7 Gram, Γ_d, and the three circuit roots d_i. Add A_θ when h is infinitesimal. The other nine CT5 squares and both final CT4 squares are retained. The following original positive packets are also retained whole: $4\beta D_w N_x^2$, $2(2D + 1)(uX_e)^2$, the nonconstant positive N_x^2 excess, $4bpzu^2\Gamma^2$, and all radial source-weight excesses above.

There are no unallocated SP6 terms. Its remaining Gamma normals are exactly $-4u\nu N_x\Gamma$ and $qwR_\gamma(N_y + bR)\Gamma$; their coefficients ν, R_γ are bounded, and CT10–CT12 allocate them infinitesimally. The $N_xN_y, N_xR, N_xA_\theta, N_yA_\theta$ entries have factors uw, uw, wB_x, u, respectively, with bounded remaining coefficients; $B_x = 8\kappa e$ is bounded. The last angular-radial term is the complete $\chi u A_\theta R$, with χ bounded. CT6 and, when needed, CT13 give bounded allocations for all these roots. The remaining radial correction is bounded infinitesimal by CT7 and LC2, and the circuit cost is $-\lambda x R^2$ with λx infinitesimal. Consequently the complete remaining Gram is a bounded infinitesimal perturbation of the stated strictly positive residue Gram. The separated bounded-Gram lemma proves finite tensor SOS. Restore every retained square to obtain the entire original parent.

All identities are exact at the sphere endpoints; no coefficient has been inferred positive by ordering a mixed element. Source openings occur only in retained positive weights or the bounded opening-cleared combinations. The proof is pointwise in the testing experiment, so separated cube roots and state ratios are legal. Real/imaginary splitting supplies complex states, with the appropriate already-admitted or matched chart chosen for each part. The bounded-W/Z part of the unbounded-H source and the bounded-H source are separate obligations and are not claimed here.

Chapter 162

C14.2: pure-testing state exits at the two small angles

Component `p8655d641d4a7`
Source: `BASE/inherited_r19/proofs/c142_contact/state_ratio_exits.tex`
SHA256: `8655d641d4a71f23ca210631ea5df11015704d9bd9b4113768285bd365c8bd5e`
Inventory use: CS1–CS8 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

162.1 C14.2: pure-testing state exits at the two small angles

Keep the unchanged C14.2 source and the exact quarter-parent BE10, with $d = 1$, L a positive bounded source unit, $x, \sigma > 0$ infinitesimal, and every original opening. In particular $D, \beta \geq 0$ may vanish. The R18 LA and LB admissions leave $u, w > 0$ infinitesimal. Literal axes have the original direct core SOS. The original pure-testing core-sign exit leaves $WZ > 0$; consequently $P = uW$ and $T = wZ$ are nonzero and have the same sign.

The two extreme state ratios. Choose positive real numbers c_*, a_*, a'_* below the actual source weights C, A, α, respectively; this is possible by LA1. Retain all positive original excesses and the entire m, D, β packets. The selected polynomial is exactly

$$H_* = c_* D_w (P - Y)^2 + a_* D_u D_w Y^2 + a'_* D_u (Y - T)^2 + K_w/2. \tag{CS1}$$

Here LA2 gives the complete signed core, without approximation:

$$K_w = D_u X^2 + D_u Y^2 + u^2 T^2 - 2u^2 TY - 2PT + w^2\{(P - Y)^2 + X^2\} + 2uwX(T + P - Y). \tag{CS2}$$

If P/T is unbounded, write $T = \epsilon P$ with the *pure-testing* infinitesimal $\epsilon = T/P$. The bounded Gram of CS1 on (X, Y, P) has residue

$$\tfrac{1}{2}X^2 + c_*(P - Y)^2 + (a_* + a'_* + \tfrac{1}{2})Y^2, \tag{CS3}$$

which is strictly positive. If P/T is infinitesimal, write $P = \epsilon T$. On (X, Y, T) the residue is

$$\tfrac{1}{2}X^2 + (c_* + a_* + \tfrac{1}{2})Y^2 + a'_*(Y - T)^2, \tag{CS4}$$

also strictly positive. All coefficients of CS1 are pure testing and real, so the usual finite positive-Gram lift is directly legal. Restoring the retained original squares proves the complete parent on both charts, for arbitrary remaining state coordinates.

Every unmatched unit state ratio. It remains to consider the positive bounded testing unit

$$\varrho = P/T = uW/(wZ).$$

Put $L_0 = \operatorname{res}(L) > 0$ and $\varrho_0 = \operatorname{res}(\varrho) > 0$. This paragraph admits every chart $\varrho_0 \neq L_0$, using only real residues, rather than ordering $\varrho - L$ as a mixed element.

Use the exact LA1 source weights C_0, A_0, a_0, with $e_r = 1 - r$, $b = Lr$, and $S_0 = A_0 + a_0 + 1/2$. Choose the same pure-source common factor $f = 1-\delta > 0$ as LA4 and retain the original nonnegative event excesses $C - fC_0$, $A - fA_0$, $\alpha - fa_0$. The entire remaining polynomial is the exact LA4 form. In its completed part LA3, retain the whole nonnegative u^2 packet, the full w^2 packet, and the coefficient excesses after clipping fC_0, fS_0 at positive real lower bounds c, s. The resulting bounded principal form on (X, Y, T) is

$$\frac{fD_u}{2}X^2 + c\{(\varrho - Le_r)T - Y\}^2 + s(Y - bT)^2. \tag{CS5}$$

Its (Y, T) determinant is exactly

$$cs(\varrho - L)^2, \tag{CS6}$$

as a polynomial identity, since $Le_r + b = L$. At residues this is the positive real number $cs(\varrho_0 - L_0)^2$. The full signed remainder $(1 - f)K_0/2 + (K_w - K_0)/2$ has infinitesimal bounded coefficients on (X, Y, T) after $P = \varrho T$, by CS2. Thus the full selected bounded Gram is strictly positive at residue. The separated bounded-Gram lemma applies; all previously retained positive packets are restored. No inverse of $\varrho - L$, or of any source/testing sum, has been used.

Precise remaining state chart. Together with LA, LB, the literal-axis exits and the core-sign exit, these two admissions reduce the whole C14.2 problem to

$$u, w > 0 \text{ infinitesimal}, \quad WZ > 0, \qquad \varrho = \frac{uW}{wZ} > 0 \text{ a bounded testing unit}, \qquad \operatorname{res}\varrho = \operatorname{res} L. \tag{CS7}$$

In particular $uW/Z = w\varrho$ is infinitesimal. The normalized radial root is exactly

$$R = \frac{wZ}{q}(1 - z\varrho), \qquad q^2 = z = L^{-1}; \tag{CS8}$$

CS7 does not compare the source and testing infinitesimal parts of L and ϱ. It does not constrain the source sphere or remove $D = 0$ or $\beta = 0$. This is a state-chart admission and reduction, not a complete C14.2 closure. Applying the real statement to the two real state parts gives the complex-state statement, with the relevant chart chosen separately for each part.

The focused checker verifies CS6 and the new three-term circuit identity in the accompanying note; it does not rerun the historical LA2–LA4 checks.

Chapter 163

Two complementary C14.2 certificates

Component `p98e371f5b846`
Source: `BASE/inherited_r19/proofs/c142_root/complementary_charts.tex`
SHA256: `98e371f5b8469c737262c40b3b234dd1c54c7cac8c788a0f1c916d68566556ea`
Inventory use: CF1–CF9 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

163.1 Two complementary C14.2 certificates

Throughout this section the source is the unchanged physical C14.2 source, with $z = L^{-1}$ a positive bounded unit, $x, \sigma > 0$ infinitesimal, and all original openings and intersections retained. The remaining tests have u, w infinitesimal. We use the full SP6 identity, its retained squares, and the exact opening-uniform radial decomposition CT7–CT7a. In particular N_y, R have a strict pure-source radial form through every opening. Every inverse below is pure source or pure testing.

163.1.1 Unbounded $H = h/\sigma$ and bounded W/Z

Suppose H is unbounded and $Z \neq 0$ with the pure-testing quotient $f = W/Z$ bounded. Write

$$s = x/\sigma, \quad L_h = \lambda/H, \quad K_A = 4\kappa H - \lambda, \quad \rho = \lambda x/\sigma.$$

BE7 and SP3 imply s infinitesimal, $\lambda > 0$ unbounded, $L_h \sim 2$, and ρ bounded positive. The complete last line of SP6, together with its angular diagonal, is exactly

$$K_A A^2 + \chi u A R + \mathscr{P}, \qquad \mathscr{P} = \lambda\sigma P_\beta^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2. \tag{CF1}$$

Here $A = q(u^2 - x)W$, $q^2 = z$. The source polynomial divisibility SP10, now with z a unit, gives the exact residue formulas

$$(K_A)_0 = 2\{H_0 - h_0(H_0 + 1 - r)_0\}, \qquad \chi_0 = 4\{H_0 - h_0(1 + z - 2r)_0\}. \tag{CF2}$$

The notation on the right means that every bounded source quantity is evaluated in its real residue. In particular K_A, χ are bounded, and h infinitesimal implies $(K_A)_0 = 2(H_0)_0 > 0$.

Put

$$U_h = \sqrt{h}\, uZ, \quad Q_\lambda = \sqrt{\rho}\, uwW, \quad M_\lambda = \sqrt{\lambda}\, xqW.$$

The following is the complete scalar identity, including both negative terms:

$$\begin{aligned}\mathscr{P} = {} & L_h\{(1 - szf^2)U_h - \sqrt{h}\, sqfR\}^2 + Q_\lambda^2 + M_\lambda^2 - L_h szf^2 U_h^2 \\ & - L_h s^2\{\sqrt{h}\, qfR + zf^2 U_h\}^2.\end{aligned} \tag{CF3}$$

All corrections to the leading $L_h U_h^2$ on the (U_h, R) frame have bounded infinitesimal coefficients. This uses s infinitesimal and q, z, f bounded; it does not require z infinitesimal.

Every Gamma normal is also retained. Set $B_f = (\sigma - xzf^2)/\sqrt{h}$. The source bounds $\sigma/\sqrt{h}$ and $x/\sqrt{h} = s\sigma/\sqrt{h}$ infinitesimal show that B_f is bounded infinitesimal. Exactly,

$$\begin{aligned} u\Gamma &= B_f U_h - uqfA - u^2 qfR, \\ qw\Gamma &= qzfB_f U_h - uz^2 f^2 A + z(\sigma - ufw - u^2 zf^2)R. \end{aligned} \tag{CF4}$$

All coefficients in CF4 are infinitesimal because u, w are infinitesimal. These identities allocate both hard first normals, the radial Gamma coupling, and the complete centered-C correction. The nonnegative diagonal $(4D/\sigma^2 + 4bpzu^2)\Gamma^2$ remains as exact squares.

If h is infinitesimal, retain $Q_\lambda^2 + M_\lambda^2$ separately. On (N_x, two radial roots, U_h, A) the complete bounded Gram has strictly positive residue: the first pivot is 2, the radial form is strict, the U_h pivot is 2, and the A pivot is $2(H_0)_0$. Every other cross coefficient is infinitesimal. If h is a positive unit, use instead the exact identity

$$A = \frac{qfu}{\sqrt{h}} U_h - \frac{1}{\sqrt{\lambda}} M_\lambda. \tag{CF5}$$

Both coefficients are infinitesimal. Keep M_λ^2 in the Gram and retain Q_λ^2 separately. The bounded signed angular packet in CF1 then contributes only infinitesimal Gram entries. The new M_λ pivot is 1.

For completeness, the remaining full SP5 terms are $N_x N_y$ with coefficient $O(uw)$, $N_x R$ with coefficient $O(uw)$, $N_x A$ with coefficient qwB_x where $B_x = 8\kappa e$ is infinitesimal, and $N_y A$ with coefficient uA_y, where A_y is bounded. The entire $N_y R$ coefficient is included in CT7–CT7a. Thus all coefficients just asserted small have explicit source or testing small factors after the opening poles have been extracted. Restore $2(2D+1)(uX_e)^2$, the original beta packet, and every radial excess. The separated bounded-Gram lemma proves the entire parent. The literal $Z = 0$ chart has the inherited core-sign exit. This proves every bounded-W/Z state for the unbounded-H source.

163.1.2 Bounded $M_\beta = \beta\theta/\sigma^2$

Suppose M_β is bounded. BE3–BE7 imply, uniformly through all openings,

$$H \sim H_0, \qquad \theta \sim 1, \qquad D/\sigma^2 \sim H_0/2, \qquad \mu = \kappa H \sim H_0/2. \tag{CF6}$$

Use the full generic JR5/JB1 identity before the SP6 pole extraction, with $V = uw(u^2 - x)W$. Its coefficients are all bounded after the same exact radial extraction. The only cross coefficient which can have nonzero residue outside the radial block is $S\Gamma V$. Write

$$S_* = \frac{2}{\sigma^2}\{h^2(2D+1)x + 4Dh - 2bp\sigma^2 z + 2\beta\theta^2 + h^2\}. \tag{CF7}$$

This is a bounded pure-source number, and $S - S_* = 2h^2(2D+1)u^2/\sigma^2$ is infinitesimal. Add the exact zero JR6 multiplied by the positive bounded pure-source number $T_* = 1 + S_*^2$:

$$T_*\{V^2 - \sigma u^2 A^2 - u\sigma(u^2 - x)AR + (u^2 - x)V\Gamma\} = 0. \tag{CF8}$$

All new coefficients except the V^2 pivot are infinitesimal. The (Γ, V) residue Gram is

$$\begin{pmatrix} 2(H_0)_0 & (S_*)_0/2 \\ (S_*)_0/2 & 1 + (S_*)_0^2 \end{pmatrix}, \quad \det = 2(H_0)_0 + \{2(H_0)_0 - 1/4\}(S_*)_0^2 > 0. \tag{CF9}$$

Here $H_0 = 1 + z(1-b) \geq 1$. The A residue pivot is $2(H_0)_0$, the N_x pivot is 2, and the exact radial form is strict. All remaining generic JR/JB crosses carry u, w or a source infinitesimal: in particular uA_x, Z_x and $u\sigma B$ are infinitesimal, by CF6 and SP4. The retained beta and cap squares are unchanged. Hence the complete original parent is tensor SOS on this source domain, for every remaining state and test.

Finally, if H is bounded and the exact SP3 coefficient $\lambda \leq 0$, its defining identity implies

$$4\beta\theta^2 + 2h^2 + 8Dh \leq 4bp\sigma^2 z.$$

Since $\theta \sim 1$, this implies M_β bounded, so CF6–CF9 cover that whole case, including $\lambda = 0$. No inverse of λ, D or β is introduced.

All these identities apply with the same source to each real state part. Together with the sign exits, they restore arbitrary complex states. The proofs do not compare a source rate to a testing rate.

Chapter 164

C14.2: bounded h/σ and the unbounded W/Z state chart

Component `pcb3428d66a71`
Source: `BASE/inherited_r19/proofs/c142_sphere/bounded_H_large_state.tex`
SHA256: `cb3428d66a716aedb79ec530fba3bafcf914a276a4c29acd67204a6d3488e917`
Inventory use: HG1–HG7 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

164.1 C14.2: bounded h/σ and the unbounded W/Z state chart

Retain the full original C14.2 source, with $z = L^{-1}$ a positive bounded unit, x, σ infinitesimal and every original opening, coefficient boundary and residual intersection unchanged. Suppose $H = h/\sigma$ is bounded. Then H is a positive unit by BE7 and h is infinitesimal. The following full-parent certificate applies to u, w infinitesimal and to every real state with $W \neq 0$ and $t = Z/W$ infinitesimal, including $t = 0$ and both signs. It makes no claim for the complementary state chart or unbounded H.

Use the exact generic SP6 parent, with $A = A_\theta$ and

$$d = 4D/\sigma^2, \qquad \rho = \lambda x/\sigma.$$

All source coefficients in the following calculations are actual coefficients; formal caps are only exact square extractions.

The sphere restores a strict Gamma direction. The identity $uP_\beta = (u^2 - x)Z + (x/\sigma)\Gamma$ gives exactly

$$\lambda u \Gamma P_\beta = \rho\Gamma^2 + \frac{\lambda t}{q} A\Gamma. \tag{HG1}$$

There is no state inverse beyond the pure-testing coordinate W^{-1}. The complete angular/Gamma packet of SP6 is consequently

$$(d + \rho)\Gamma^2 + 4\kappa H A^2 + \frac{\lambda t}{q} A\Gamma + (\lambda + \chi) uAR. \tag{HG2}$$

No signed scalar product was discarded in this transformation.

To control the formerly missing Gamma pivot uniformly, BE4 implies the exact source identity

$$\rho = \frac{D}{\sigma^2}(8hx - 4\theta J) + \frac{2xh^2}{\sigma^2} + \frac{2\theta\mathcal{H}}{P} - 4bpxz. \tag{HG3}$$

Also, directly from BE7,

$$\frac{xh^2}{\sigma^2} = \frac{eH_0}{B} + \frac{h\mathcal{H}}{B} b_0^2, \qquad e = xh/\sigma \in \text{Inf}. \tag{HG4}$$

Since $\operatorname{res}(D/\sigma^2) = \frac{1}{2}\operatorname{res}(H_0 d_0^2)$, $\mathcal{H}/P \sim H_0$, $J \sim 1$ and $d_0^2 + b_0^2 = 1$, these identities give, even without bounded H,

$$\operatorname{res}\rho = 2\operatorname{res}(H_0 b_0^2), \qquad \operatorname{res}(d+\rho) = 2\operatorname{res} H_0 > 0. \tag{HG5}$$

In particular no ratio to the possibly vanishing b_0 is asserted, and neither D nor b_0 is inverted. In the bounded-H chart, λ, χ are bounded, $4\kappa H$ is a positive unit, and q is a unit. Both cross coefficients in HG2 are therefore infinitesimal.

The full radial and hard block through the C opening. Restore the centered A, α cap excesses into the N_y diagonal. Put $a = Az$, $\ell = \alpha z$, $Q = 1 - b$, $J_C = N_y + bR$ and $J_\ell = N_y - QR$. Use the exact LC3 cap $C_* = C/(1+C)$, retaining $4zD_w(C - C_*)J_C^2$ and its complete correction

$$-8(C - C_*)(1 - r)J_C(z\sigma R - qw\Gamma). \tag{HG6}$$

The correction has infinitesimal coefficients on (N_y, R, Γ), because $(C - C_*)(1 - r)$ is bounded. The exact leading pure-source radial form is

$$M = \{4a + 2z(2C_* + 1)\}J_C^2 + 4\ell J_\ell^2. \tag{HG7}$$

Both coefficients have positive real lower bounds; their potentially unbounded excesses are retained as squares. The inverse allocations $R = J_C - J_\ell$ and $N_y = QJ_C + bJ_\ell$ are bounded. Therefore M supplies a strict bounded Gram on (N_y, R) through every r, $1 - r$, and Q opening.

For clarity, HG7 is obtained from the entire formal radial/hard coefficient collection by putting $u = w = \sigma = D = m = \beta\theta = h = 0$ while retaining the actual pure-source $a, \ell, p = ab, l = \ell Q, z, b, C_*$. The omitted u^2 and w^2 multiples of a, ℓ are retained exactly as the nonnegative packets from

$$4(1 + u^2)\{a(1 + w^2)J_C^2 + \ell J_\ell^2\}.$$

All remaining coefficient differences are bounded infinitesimals: p, l, z, b, C_* are bounded, $m, D, \beta\theta, h, \sigma$ are infinitesimal, and u, w are testing infinitesimals. This assertion is made only after the divergent C correction HG6 has been accounted for, and after retaining the displayed a, ℓ packets.

The N_x diagonal has residue 2. Its Gamma normal is $-4u\nu N_x\Gamma$ with bounded ν; both other Gamma normals carry qw. All remaining SP5 hard and first-normal coefficients are infinitesimal after the radial source form HG7 is restored. The separated Gram on $(N_x, J_C, J_\ell, \Gamma, A)$ consequently has strict positive residue. HG2 supplies the last two positive pivots. The original beta packet, the $2(2D + 1)(uX_e)^2$ square, the $4bpzu^2\Gamma^2$ square, and every source cap excess remain once. Thus the entire original SP6 parent is included, not only its scalar restriction. The bounded separated-Gram lemma proves tensor SOS.

All formulas are literal at zero testing angles and $Z = 0$; the case $W = 0$ has the inherited original-core sign exit. Application to real and imaginary state parts gives the corresponding complex state admission when each part lies in this chart or an existing admission. This is a source/state subchart result, not a complete C14.2 closure.

Chapter 165

C14.2: bounded h/σ with positive λ, all states

Component `p021ed4a9e0db`
Source: `BASE/inherited_r19/proofs/c142_sphere/bounded_H_positive_lambda_all_states.tex`
SHA256: `021ed4a9e0dbaff0b9efaa87cabf5620db15d898efbbf031f113139ada5cf141`
Inventory use: HP1–HP10 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

165.1 C14.2: bounded h/σ with positive λ, all states

Retain the original C14.2 source and all its intersections, and the complete small-angle corner $u, w \in \mathrm{Inf}$. Suppose $H = h/\sigma$ is bounded and $\lambda > 0$. This section proves the entire original parent for every testing state. It retains literal $D = 0$, every source-sphere opening, and every allowed coefficient boundary. The source case $\lambda \le 0$ is a separate obligation.

Use the exact SP6/BS2 parent, with the complete C-opening extraction LC3 and the actual radial frame HG7. The source estimates proved in HG3–HG5 and the SP definitions give

$$\rho = \lambda x/\sigma \ge 0, \quad \rho \in \mathcal{O}_c, \quad d = 4D/\sigma^2, \quad d + \rho \sim 2H_0, \quad K_A = 4\kappa H - \lambda \sim 2H_0 > 0. \tag{HP1}$$

Here λ, χ, ν are bounded and $\lambda x = \rho\sigma$ is infinitesimal. The source $q = \sqrt{z}$ is a unit. The complete parent outside already positive rows consists of its strict hard/radial frame, $d\Gamma^2 + K_A A_\theta^2 + \chi u A_\theta R$, the complete Gamma normals, and the signed packet

$$\mathscr{P} = \lambda\sigma P_\beta^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2. \tag{HP2}$$

No source or testing rate is compared across the two fields.

A ternary circuit that supplies an actual difference pivot. The original-core sign exit and exact testing gauges reduce the remaining real states to $u, w, W, Z > 0$. Literal axes and $WZ \le 0$ already have direct original-core SOS. Let

$$a = \sqrt{\lambda\sigma}\, uZ, \qquad b = \sqrt{\rho}\, uwW, \qquad c = \sqrt{\lambda z}\, xW, \qquad g = (abc)^{1/3}, \qquad r^3 = \frac{wZ}{zuW}.$$

Every radical and quotient is a product of explicitly positive pure-source and pure-testing factors. Direct expansion gives

$$\boxed{\mathscr{P} + \lambda x R^2 = a^2 + b^2 + c^2 - 3g^2 + \lambda x z u^2 W^2 r^2 (r-1)^2 (r^2 + 2r + 3).} \tag{HP3}$$

The last term is a finite legal SOS since $r^2 + 2r + 3 = (r+1)^2 + 2$. There is a stronger exact decomposition of the first term:

$$a^2 + b^2 + c^2 - 3g^2 = \tfrac{1}{2}\{(a-b)^2 + (b-c)^2 + (c-a)^2\} + (ab + bc + ca - 3g^2). \tag{HP4}$$

The final parenthesis is itself a finite legal SOS. Indeed, put $\alpha = (ab)^{1/3}$, $\beta = (bc)^{1/3}$, $\gamma = (ca)^{1/3}$. It is exactly

$$\tfrac{1}{2}(\alpha+\beta+\gamma)\{(\alpha-\beta)^2+(\beta-\gamma)^2+(\gamma-\alpha)^2\}. \tag{HP5}$$

Expand the positive prefactor into its three monomial terms before taking square roots; there is no radical of a mixed sum. Thus HP3 supplies a genuine unit-weight pivot $D_1 = (a-b)/\sqrt{2}$, with every other displayed row retained. The small subtraction $-\lambda x R^2$ is an infinitesimal Gram perturbation of the strict radial frame.

The source-sphere split. If d_0 is a positive unit, $d\Gamma^2$ supplies a strict Gamma pivot. Every complete Gamma normal in SP6/LC3 carries u, w or σ and has a bounded remaining coefficient. Hence all these normals are infinitesimal entries on the strict hard/radial, Gamma and angular frame. Retain all of HP3 as separate SOS. Together with $K_A \sim 2H_0$ this closes this source case for every state, without any state quotient.

It remains that d_0 is infinitesimal, including $D = 0$. Then ρ is a positive unit by HP1. If W/Z is unbounded, the full-parent HG certificate applies. Thus consider bounded $f = W/Z$. The next argument supplies every missing Gamma normal without imposing a positive pivot on Γ.

A quadratic domination conductor, without a mixed inverse. Put $k = \sqrt{x}$, and define

$$T = qu(u-k)W, \qquad \delta = w/q > 0, \qquad T_1 = \sqrt{\delta}\,T.$$

All these roots are separated and δ is infinitesimal. The exact circuit and angular identities are

$$u\Gamma = \frac{\sqrt{2}k}{\sqrt{\rho}}D_1 - \delta T, \tag{HP6}$$

$$A_\theta^2 - T^2 = q^2(u-k)^2W^2(k^2+2uk) = (qk(u-k)W)^2 + 2(q\sqrt{u}\sqrt{k}(u-k)W)^2. \tag{HP7}$$

Reserve the infinitesimal part δA_θ^2 of the strict angular diagonal. Exactly,

$$K_A A_\theta^2 = (K_A-\delta)A_\theta^2 + T_1^2 + \delta(A_\theta^2 - T^2). \tag{HP8}$$

The final term is a legal SOS by HP7; the first coefficient has unchanged positive residue. Thus T_1 is a genuine additional unit-weight pivot, paid for exactly once from the angular row. HP6 becomes the bounded infinitesimal allocation

$$u\Gamma = \frac{\sqrt{2}k}{\sqrt{\rho}}D_1 - \sqrt{\delta}\,T_1. \tag{HP9}$$

This avoids division by $u+k$. A pointwise bound on $u/(u+k)$ alone would not be a legal tensor allocation.

The other full Gamma normal has the exact state conductor

$$qw\Gamma = qzf\,u\Gamma + z(\sigma - uwf)R. \tag{HP10}$$

Its first coefficient is bounded and the second is infinitesimal. Together HP9–HP10 allocate both $u\Gamma$ and $qw\Gamma$ with infinitesimal coefficients on (D_1, T_1, R). The only state inverse used is the pure-testing Z^{-1} in the bounded quotient f; no source/testing expression is inverted.

Full-parent reassembly. The entire Gamma coupling in SP6 is

$$-4\nu N_x(u\Gamma) + R_\gamma(N_y + bR)(qw\Gamma).$$

LC3 adds only its recorded bounded-t_C correction to the same $qw\Gamma$ normal and to σR. Substitution of HP9–HP10 makes every one of these couplings infinitesimal on the strict frame

$$(N_x, J_C, J_\ell, A_\theta, D_1, T_1).$$

HG7 supplies the radial/hard positive source rows; all divergent a, ℓ angle-dependent excesses and the centered C excess are retained as in HG. The other complete SP5 couplings and $\chi u A_\theta R$ are bounded infinitesimals. The subtraction $-\lambda x R^2$ is infinitesimal by HP1. Consequently the whole frame has strict positive residue.

Retain exactly once every remaining positive row: $d\Gamma^2$, $4bpzu^2\Gamma^2$, $2(2D+1)(uX_e)^2$, the original beta packet, the unused HP3–HP5 circuit rows, the HP7 remainder from HP8, and all source-cap excesses. In particular the $(a-b)^2/2$ row and the angular fraction used for T_1 are not counted twice. The separated bounded-Gram lemma now proves the complete parent.

The two sphere cases exhaust the source, and the bounded/unbounded W/Z alternatives exhaust the states after the direct sign/zero exits. Real/imaginary splitting gives all complex states. Thus all remaining C14.2 tests are admitted on bounded H and $\lambda > 0$. The result is conditional only on the explicitly stated source split, not on an unproved state-rate comparison.

Chapter 166

C14.2 is closed: the exact all-tests and all-states join

Component `p896e19be06f0`
Source: `BASE/inherited_r19/proofs/c142_sphere/c142_all_tests_join.tex`
SHA256: `896e19be06f0400e4f65ad51d3608f1907fcc28cb892bc32185a8223b6a1f8e2`
Inventory use: entire analytical join | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

166.1 C14.2 is closed: the exact all-tests and all-states join

Theorem. For every original physical C14.2 source, the complete original parent is finite tensor SOS for every independent projective testing experiment and every real or complex state. Precisely, the source has $x > 0$ infinitesimal, $L = x\zeta$ a positive bounded unit and η unbounded, with all original stationary-contact, strict-gap, nonnegative-coefficient, total-scale and prior-complement intersections retained. The conclusion includes literal $D = 0$ and $\beta = 0$ and every positive opening rate. It closes the whole C14.2 branch, not the other C1 branches or the global theorem.

Exhaust the testing angles first. The reviewed R18 LA proposition admits every noninfinitesimal Alice angle, including its projective endpoint, and every Bob angle. The reviewed R18 LB proposition admits infinitesimal or zero Alice angles with bounded Bob angle v. The only complement is $u, w = 1/v$ infinitesimal. At literal $u = 0$ or $w = 0$ the original reciprocal core is respectively

$$(1 + w^2)(X^2 + Y^2), \qquad (1 + u^2)(X^2 + Y^2),$$

and every original event packet is nonnegative. Thus those axes are covered without any limiting argument. Exact testing sign gauges reduce the remaining real tests to $u, w > 0$.

Exhaust the two source types. Write $\sigma = 1/\eta$, $z = L^{-1}$ and $H = h/\sigma$. BE7 bounds H below by a positive source unit. It is therefore either bounded or unbounded; neither case changes the original source intersections.

If H is bounded, split the exact pure-source coefficient λ by sign. When $\lambda \leq 0$, CF6–CF9 apply: the defining inequality for λ implies bounded $M_\beta = \beta\theta/\sigma^2$, and the generic full-parent Gram with its exact V pivot proves all states. This includes $\lambda = 0$ and literal $\beta = 0$. When $\lambda > 0$, HP proves all states. Its source-sphere split is exhaustive: a unit d_0 gives the original strict Gamma pivot; infinitesimal d_0, including $D = 0$, gives a unit ρ. In the latter case HP handles bounded W/Z by the exact ternary circuit and quadratic-domination conductor, and HG handles unbounded W/Z. Both preserve the entire original parent.

If H is unbounded, CF1–CF5 handle bounded W/Z. The remaining unbounded-W/Z states are exhausted by CS and CT. The core-sign exit first removes $WZ \leq 0$, including zero state coordinates.

On the positive complement CS1–CS4 admit the infinitesimal or unbounded pure-testing ratio $uW/(wZ)$; CS5–CS6 admit unit ratios whose real residue differs from that of L. CT then proves the whole remaining positive-unit ratio chart. Its source-sphere Gamma conductors CT10–CT12 retain both coefficient endpoints, and its two angular cases exhaust infinitesimal and unit h. Thus no state ratio remains.

Restore the original quantifiers and boundaries. Every state division above is made only on a nonzero pure-testing coordinate; the omitted zero charts have the stated direct exits. Each chart contains the complete hard block, scalar packet and first normals. The CT7–CT7a radial decomposition or its exact LC3/HG7 counterpart controls the $r \to 1$ opening, rather than assuming C bounded. The $r \to 0$, $1 - Lr \to 0$, $\theta \to 0$ and source-sphere openings occur only in retained original squares or proved bounded opening-cleared coefficients. No mixed inverse, cross-field rate comparison or unproved strict soft pivot is used.

Apply the real certificate separately to the real and imaginary parts of each state, choosing the relevant chart independently for each part. Restore the exact sign gauges, the projective testing normalizations and the positive pure-source $d = 1$ normalization. Multiplication by these pure positive factors preserves finite tensor SOS. The LA/LB projective endpoints and the literal small axes have already been included, so this restoration introduces no continuity step or missing boundary.

The alternatives above route every original C14.2 source/test/state tuple to a proved full-parent certificate. The branch is therefore admitted for all independent tests and states. The algebra receipts and same-system mathematical reviews support the new identities and their stated scopes; they are not external or formal verification.

Chapter 167

C14.4: a contact-centered full-parent identity through the M opening

Component `pdc67cb6181a3`
Source: `BASE/proofs/c144/centered_m_parent.tex`
SHA256: `dc67cb6181a37c47e9bf53b46d5c1ee432bc732f5b1906c69b7ba5b2e6ce322f`
Inventory use: CM1–CM7 | CM1-CM7 actual contact-zero m normal and exact opening-safe coefficient | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

167.1 C14.4: a contact-centered full-parent identity through the M opening

This section is an exact reduction, not an all-tests admission. It supplies an opening-safe state normal for the remaining C14.4 source. Keep all original source conditions and write $R_s = 1 + xr$, $M_s = R_s - \theta/\zeta > 0$. Define only in the coefficient field

$$p = \frac{\theta}{\zeta R_s} = 1 - \frac{M_s}{R_s} \in (0,1), \qquad \mu = m(1-p) = \frac{(1+x)(1+\sigma r)eS}{2\zeta P R_s}, \qquad m_c = \frac{m}{1+m}. \tag{CM1}$$

Here $0 < m_c < 1$, $m - m_c \geq 0$, and μ is bounded uniformly through $M_s \to 0$. In particular the potentially unbounded m occurs below only in an unchanged positive excess square.

On the finite Bob chart let $D_u = 1 + u^2$, $D_v = 1 + v^2$ and set

$$V = v(W + uY), \quad N = X + uZ - pV, \quad J_0 = uW - Y + puvV - (1+u^2)vZ. \tag{CM2}$$

The original event amplitudes are exactly

$$vW - X + uvY - uZ = (1-p)V - N, \qquad X + uZ = N + pV, \qquad uW - Y + uvX - vZ = J_0 + uvN. \tag{CM3}$$

Consequently the entire original quarter-parent is

$$\begin{aligned} F/4 = {} & (m - m_c)N^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + m_cN^2 - 2\mu NV + \mu(1-p)V^2 + (D + \tfrac{1}{2})(J_0 + uvN)^2 \\ & + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 + (\beta + \tfrac{1}{2})D_v(N + pV)^2 - \tfrac{1}{2}D_uD_vZ^2. \end{aligned} \tag{CM4}$$

This is a polynomial identity after clearing pure-source denominators. It follows by substituting CM3 into all six original event packets and the complete signed core. No state variable has been set to a contact value in CM4.

Why this particular normal preserves the actual contact. For the actual source contact, $u_c = k = \sqrt{x}$, $v_c = \sqrt{L/\sigma}$, $W_c/Z_c = \sqrt{\zeta\sigma}$, $X_c = -hkZ_c$ and $Y_c = rkW_c$. Thus $v_cW_c = \zeta kZ_c$ and

$$N_c = \theta kZ_c - pR_s\zeta kZ_c = 0. \tag{CM5}$$

Retaining $(m - m_c)N^2$ therefore respects the original kernel. Capping the uncentered m event itself would not have this property when $M_s > 0$. The contact calculation is a source-field check of the choice of normal; it is not an identification of independent testing variables with source parameters in the tensor claim.

The complete first-normal coefficient is bounded. For bounded v and the residual infinitesimal u, put $\widehat{N} = \sqrt{1+\beta}\,N$. Expanding the last three squares in CM4 shows that its residual $\widehat{N}^2$ coefficient is

$$H_N = \frac{m_c}{1+\beta} + \frac{(D+1/2)u^2v^2}{1+\beta} + \frac{\beta+1/2}{1+\beta}D_v. \tag{CM6}$$

More precisely,

$$H_N - \tfrac{1}{2}D_v = \frac{m_c}{1+\beta} + \frac{(D+1/2)u^2v^2}{1+\beta} + \frac{\beta D_v}{2(1+\beta)}$$

is a sum of separated nonnegative products, hence tensor SOS. This identity states the diagonal reserve without assigning an order to the mixed tensor ring. The full first-normal cross term is

$$2\widehat{N}\left[\frac{-\mu + (\beta+1/2)D_vp}{\sqrt{1+\beta}}V + \frac{(D+1/2)uv}{\sqrt{1+\beta}}J_0\right]. \tag{CM7}$$

All these coefficients are bounded separated expressions on the stated chart. Indeed ES gives D, α infinitesimal, while $2\theta P\beta = hB - \sigma H_0$ gives $\beta\theta = O(1)$. Hence $\beta p = (\beta\theta)/(\zeta R_s)$ and βp^2 are bounded through $\theta \to 0$; $1/\zeta$ and $1/R_s$ are bounded. CM1 bounds μ through the independent M_s opening. The real residue of H_N is at least the residue of $D_v/2$, even when one of the opening parameters is infinitesimal.

The remaining scalar and Y form is printed explicitly by CM4 with $N = 0$; the original positive C, A rows are retained. A strict normal pivot does not prove that this remaining form is tensor SOS, and completing it using a mixed inverse of H_N is not allowed. In particular CM4 alone does not resolve the matched Bob source/test contact or the unit-e sector. A valid completion must allocate the full signed soft form together with CM7, or produce an exact impossible-source proof. The reciprocal Bob chart is not supplied by a finite-v limit.

The accompanying checker verifies CM3–CM7 against the entire original four-state polynomial and verifies the contact relation. This reduction retains every coefficient opening and exposes the remaining soft problem without a divergent m first-normal coefficient.

Chapter 168

C14.4: extend the unmatched-Bob certificate to the entire small-e source

Component `p8c30238a5f7d`
Source: `BASE/proofs/c144/small_e_unmatched_bob.tex`
SHA256: `8c30238a5f7d19fdf1af94dbe933a8b34954519c3ca740740a4ad22844e6602f`
Inventory use: SE1–SE12 | SE source bounds and complete unmatched Bob all-state exit | SE1-SE12: source scope and complete unmatched-residue/adjacent-C,A joins

168.1 C14.4: extend the unmatched-Bob certificate to the entire small-e source

This is a partial-domain theorem. It retains the original physical C14.4 source, all strict gaps, nonnegative event weights, the original total-scale and prior-complement intersections, and independent source and testing fields. Its source has $x, L > 0$ infinitesimal, $\zeta = L/x$ noninfinitesimal and η unbounded. Write $\sigma = 1/\eta$, $S = \sigma + L$ and

$$e = \frac{xh}{\sigma}, \quad E = 1 - e, \quad \kappa = \frac{x}{S}, \quad \nu = \frac{\sigma}{S}, \quad \ell = \frac{L}{S}, \quad \nu + \ell = 1. \tag{SE1}$$

The letter κ in this section denotes x/S, not the different coefficient denoted by κ in the R19 reciprocal-L chart. All source inequalities are taken in the coefficient field. Real residues are compared only after taking residues separately in the two fields.

The exact source input. Use ES1–ES10 from the retained R13 source proof. In particular $1/18 \le E < 1$, $0 < h, r < 1$, $\theta = 1 - h$, and

$$\begin{gathered} J = 1 + xh - \sigma(1 - r), \qquad M = 1 + xr - \theta/\zeta > 0, \\ Q = 1 - Lr, \quad H_0 = 1 + L(1 - r), \quad B = P + x(J - \sigma), \\ P = \sigma(1 + L + 2xh + rxL + rx^2h) + (1 + x)L + x^2h\theta, \\ \mathcal{H} = (1 + xr)H_0 + (1 - r)xh(2 + xr) - e\{2 + x(1 + h + r\theta)\} + (1 + x)e^2. \end{gathered} \tag{SE2}$$

The opening-safe sphere identities are

$$\begin{gathered} 2\theta P\beta = hB - \sigma H_0, \qquad d_0^2 + b_0^2 = 1, \quad d_0, b_0 \ge 0, \\ D = \frac{\sigma^2\mathcal{H}}{2PJ}d_0^2, \qquad x\theta\beta = \frac{\sigma^2\mathcal{H}}{2P}b_0^2, \\ e\frac{B}{S} = \kappa H_0 + \nu\mathcal{H}b_0^2, \qquad e^2\frac{B}{S} = \kappa(eH_0 + h\mathcal{H}b_0^2). \end{gathered} \tag{SE3}$$

Here $P/S \sim 1$, $J, Q, H_0 \sim 1$, $B/S - (1 + \kappa)$ is infinitesimal, and $\mathcal{H} - E^2$ is infinitesimal. In particular B/S and $\mathcal{H}$ are positive bounded units. These estimates are the already admitted ES bounds; they do not use bounded η.

Source dichotomy. On the entire C14.4 source,

$$e \text{ is infinitesimal} \iff \kappa = x/S \text{ is infinitesimal.} \tag{SE4}$$

Indeed the first nonnegative sum in SE3 gives $\kappa = O(e)$, and the second gives $e^2 = O(\kappa)$. Both implications are ordered-field statements entirely inside the source field.

Suppose e is infinitesimal. If ζ is unbounded, then θ/ζ is infinitesimal and $M \sim 1$ directly. If ζ is a positive bounded unit, then the exact relation

$$\ell = \zeta\kappa \tag{SE5}$$

gives ℓ infinitesimal and $\nu \sim 1$. Consequently $B/\sigma \sim 1$. Since $\beta \geq 0$, the first identity of SE3 implies $hB \geq \sigma H_0$, whence $h \sim 1$ and θ is infinitesimal. Thus $M \sim 1$ in this case as well. We have proved, uniformly on the whole small-e source,

$$E \sim 1, \quad M \sim 1, \quad m \in \mathrm{Inf}_{>0}\,. \tag{SE6}$$

For the last statement use the exact original coefficient

$$m = \frac{(1+x)(1+\sigma r)eS}{2M\zeta P}; \tag{SE7}$$

$1/\zeta$ is bounded and every other coefficient apart from e is bounded, including the reciprocal of M just proved to be a unit. No assumption on the ratio of a source infinitesimal to a testing infinitesimal occurs.

The two nonnegative terms in the first sum in SE3 imply νb_0^2 infinitesimal. Hence the exact sphere formula and the exact formula for α yield

$$\frac{D}{S} - \frac{\nu^2}{2} \in \mathrm{Inf}, \qquad \frac{\alpha}{S} - \frac{\ell^2}{2} \in \mathrm{Inf}, \qquad SC(1-r) \sim \tfrac{1}{2}, \qquad SAr \sim \tfrac{1}{2}. \tag{SE8}$$

For clarity the second formula is obtained without dividing by ℓ:

$$\frac{\alpha}{S} = \frac{(1+\sigma r)\{\ell(1+xr)e + \ell^2 ME\}}{2Q(P/S)}. \tag{SE9}$$

These are additive bounded residue estimates, including $D = 0$ whenever that boundary has points in the stated domain. They do not assert a relative estimate at a vanishing pivot.

Theorem: every unmatched Bob residue on the small-e source. Let e be infinitesimal. In the testing field choose normalized homogeneous Bob coordinates (c, s) with $c^2 + s^2 = 1$. For every independent projective test and every real or complex state satisfying

$$\overline{s^2} \neq \overline{\ell}, \tag{SE10}$$

the complete original C14.4 parent is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$.

Proof with the complete parent retained. Noninfinitesimal Alice tests use the unchanged adjacent-C, A theorem: SE8 and S infinitesimal imply that both C and A are unbounded. For infinitesimal or zero u, apply the exact EB3–EB7 homogeneous frame from the retained R18 proof. Those formulas are algebraic identities; their only source-limit inputs to the bounded-Gram proof are precisely m infinitesimal and SE8, now established on the larger source domain. To make the imported implication explicit, put

$$H = S(C+A), \quad \rho = \frac{C}{C+A}, \quad a_h = \frac{SCA}{C+A}, \qquad p = \frac{uW}{\sqrt{S}},\ q = \sqrt{S}Z,\ y = \frac{Y}{\sqrt{S}}.$$

Then $1/\sqrt{H}$ is bounded, $0 < \rho < 1$, $a_h \sim 1/2$, and the harmonic normal $N_y = \sqrt{H}(y - \rho p)$ can be substituted by $y = \rho p + N_y/\sqrt{H}$ with bounded source coefficients. The remaining normals are the EB4 N_m, N_b, N_r, with only pure-field radicals and the positive testing denominators $1 + u^2$ and $1 + u^2 s^2$. The exact full quarter-parent is EB7,

$$N_y^2 + a_h p^2 + ASu^2 y^2 + M_B^2 + D_B^2 + A_B^2 + \beta(X + uZ)^2 + \tfrac{1}{2} K_B. \tag{SE11}$$

All EB5 core terms and all EB6 event amplitudes are retained. In particular no state graph is imposed and neither first-normal coupling is removed. The two positive rows ASu^2y^2 and $\beta(X + uZ)^2$ are left whole. On the other six roots $(N_y, N_m, N_b, N_r, p, q)$ the bounded Gram has residue equal to $\mathrm{diag}(1, 1, 1/2, 1/2)$ plus the scalar block

$$\frac{1}{2}\begin{pmatrix} 1 & -\bar{s}\bar{c} \\ -\bar{s}\bar{c} & \bar{\nu}^2\bar{s}^2 + \bar{\ell}^2\bar{c}^2 \end{pmatrix}. \tag{SE12}$$

Its determinant is $(\overline{s^2} - \bar{\ell})^2/4$, so SE10 makes this finite coefficient Gram strictly positive definite. The separated bounded-Gram lemma supplies tensor SOS; restoring the two whole rows gives the complete parent. The exact homogeneous formulas allow $c = 0$, $s = 0$ and $u = 0$. Signs and real/imaginary state parts are handled exactly as in EB, and the original positive source and testing normalizations are pure-field factors. This proves the stated domain.

New part, and exact retained complement. Unbounded ζ already forces $\kappa \le 1/\zeta$ infinitesimal, so the new theorem contains the entire R18 unmatched-residue admission. Its additional domain is ζ a positive bounded unit, e infinitesimal, and Bob sine square with nonzero residue; this includes Bob infinity. In this added domain ℓ has zero residue, so the remaining Bob chart is exactly finite infinitesimal v (including zero, which may be handled by independent literal-axis exits).

After all these admissions, the C14.4 remainder is contained in the union of (i) e infinitesimal and matched Bob residue, and (ii) ζ a positive bounded unit and e a positive bounded unit. All original intersections remain in both. In case (ii), SE7 shows that m has a positive real lower bound, while its upper bound may diverge through $M = 0$. This is a source estimate, not a full-parent certificate. Neither remaining domain is declared empty or closed. C14.4 and U1 remain open.

Verification scope. The new checker verifies the source rescalings and exact identities used above. The seven EB identities are replayed separately from an unchanged temporary copy. Their algebra does not by itself prove the source bounds, strictness, or universal scope; those are the arguments just given.

Chapter 169

State-free normal conductors at the large-L corner

Component p3be5a8005801
Source: BASE/proofs/c1_unbounded/c141a_closure_and_c141b_partial.tex
SHA256: 3be5a8005801fb7deeadb216e818574e376e5eaf56659c8337df32734e75bc6b
Inventory use: SF1-SF9 source bounds, exact radial and normal allocation, surviving h-residue<1/2 C14.1b subdomain | SF1–SF18 and both theorem proofs | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

169.1 State-free normal conductors at the large-L corner

Claims and retained input. Keep the original physical stationary C1 source, its positive d, strict gaps, nonnegative event coefficients, original infinitesimal total scale, and the complement of the previously admitted domains. Normalize $d = 1$. Source quantities belong to a real closed extension K_c and testing angles and states to an independent extension K_t. The conclusion is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$, not pointwise positivity after identifying the fields. In this section $x, z = 1/L, \sigma = 1/\eta$ are all positive infinitesimals. All original source openings are retained, including the allowed literal faces $D = 0$ and $\beta = 0$.

The first theorem closes the entire original C14.1a residual. The second closes a proper part of C14.1b. The precise remaining C14.1b complement is stated at the end. No global theorem is claimed.

Theorem 169.1 (C14.1a)**.** *If $H = h/\sigma$ is bounded, the complete original parent is tensor SOS for every independent projective test and every real or complex state. In particular the previously unresolved C14.1a domain, $M_\beta = \beta\theta/\sigma^2$ unbounded, is admitted completely.*

Theorem 169.2 (A further C14.1b subdomain)**.** *If H is unbounded and the real residue of h is strictly less than $1/2$, the same full-parent conclusion holds. This includes every infinitesimal h and every positive unit h with residue less than $1/2$, but does not include residue $1/2$ or greater.*

The proof uses the exact complete P4–P8/SP6 parent in the inherited `parent_reference.tex`, the BE1–BE9 source identities, and the already admitted remote and literal-angle exits. The new argument settles the remaining $u, w = 1/v$ infinitesimal corner directly; it does not reclassify any source outside the original C14.1 domains.

169.1.1 Source bounds with z infinitesimal

Write $q = \sqrt{z}$, $k = \sqrt{x}$, $b = Lr$, $Q = 1 - b$, $\theta = 1 - h$, $e = xh/\sigma$, and retain the exact source letters $J, M, P, B, \mathcal{H}, p, l, \kappa$ from P1–P2. Since $r = bz$ and $0 < b < 1$, r is infinitesimal and $1 - r$ is a unit.

BE1 applies because it requires only that L have a positive real lower bound; it gives $e = O(x+\sigma)$ and $E = 1 - e \sim 1$. Consequently

$$P, J, M, B, \mathcal{H} \sim 1, \qquad p, l \sim \tfrac{1}{2}, \qquad C = O(z), \qquad \kappa \sim \tfrac{1}{2}. \tag{SF1}$$

The possible b^{-1}, Q^{-1} poles occur only in the two original positive source weights $a = Az = p/b$ and $\ell = \alpha z = l/Q$. Both have positive real lower bounds. They need not be bounded above. No cap of an individual unbounded matrix entry is used below.

Let d_0, b_0 be the exact source sphere from P2, and put

$$d = 4D/\sigma^2, \qquad \lambda = 2H + \lambda_0, \qquad \rho = \lambda x/\sigma, \qquad K_A = 4\kappa H - \lambda. \tag{SF2}$$

The identities SP9–SP10 show that $\nu, \lambda_0, \chi, K_A$ are bounded even when H is unbounded. This is a polynomial divisibility argument, not multiplication of an uncontrolled error by H. More precisely their source residues give

$$(\nu)_0 = 1, \qquad (\lambda_0)_0 = -2(1 - h_0), \qquad (K_A)_0 = 2 - 4h_0. \tag{SF3}$$

For example the SP10 polynomial F_σ has residue -1 at $x = z = \sigma = 0$, whereas eF_x has zero residue. Thus $\alpha_\kappa = 2H(2\kappa - 1)$ has residue $-2h_0$, which proves the last equality without assuming a rate for $2\kappa - 1$.

The source-sphere calculations HG3–HG5 are algebraic for bounded z and do not require it to be a unit. They give

$$\rho \text{ is bounded}, \qquad (\rho)_0 = 2(b_0^2)_0, \qquad (d + \rho)_0 = 2. \tag{SF4}$$

One way to check the delicate assertion is to use both exact identities

$$\rho = \frac{D}{\sigma^2}(8hx - 4\theta J) + \frac{2xh^2}{\sigma^2} + \frac{2\theta\mathcal{H}}{P} - 4bpxz, \qquad \frac{xh^2}{\sigma^2} = \frac{eH_0}{B} + \frac{h\mathcal{H}}{B}b_0^2. \tag{SF5}$$

These include $D = 0$ and $\beta = 0$. Throughout the positive-λ case $\lambda x = \rho\sigma$ is infinitesimal.

When H is bounded, h is infinitesimal, $K_A \sim 2$, and λ is bounded. If $\lambda \le 0$, its exact defining formula implies

$$8Dh + 4\beta\theta^2 + 2h^2 \le 4bp\sigma^2 z.$$

Since $\theta \sim 1$, this forces bounded M_β; the inherited JB bounded-M_β all-tests theorem applies. On the actual C14.1a remainder M_β is unbounded, so $\lambda > 0$.

If d_0 is infinitesimal, including literal $D = 0$, SF4 makes ρ a positive bounded unit. The positive sphere identity

$$H = H_0/B + (\sigma/x)(\mathcal{H}/B)b_0^2 \tag{SF6}$$

then gives the source bounds needed for the two theorems:

$$\begin{array}{ll} H \text{ bounded}: & \sigma/x \text{ bounded}; \\ H \text{ unbounded}: & x/\sigma \text{ infinitesimal}, \quad xh/\sigma^2 \text{ a positive unit}, \quad \sigma/k \asymp \sqrt{h} \text{ bounded}. \end{array} \tag{SF7}$$

For the second line use $\lambda/H \sim 2$ and $\rho \sim 2$. None of these is a comparison between a source rate and a testing rate.

169.1.2 The full source-opening-safe radial block

The roots are the original ones:

$$N_x = X + uhZ, \quad N_y = (Y - bwZ)/q, \quad \Gamma = \sigma Z - uwW, \quad R = wZ/q - uqW, \quad A_\theta = q(u^2 - x)W.$$

Put $J_C = N_y + bR$ and $J_\ell = N_y - QR$. The exact CT7–CT7a radial extraction, with every a, ℓ excess restored, has leading pure-source form

$$[4a + 2z(2C + 1)]J_C^2 + 4\ell J_\ell^2. \tag{SF8}$$

Retain the nonconstant positive u^2, w^2 packets and all positive source-weight excesses. Clipping the two coefficients in SF8 at positive real lower bounds gives a strict bounded Gram on (N_y, R), since

$$R = J_C - J_\ell, \qquad N_y = QJ_C + bJ_\ell, \qquad b + Q = 1.$$

This is uniform through $b \to 0$ and $Q \to 0$. The remaining three radial coefficients are precisely CT7a. They are infinitesimal because z is infinitesimal, $C = O(z)$, and all their other source letters are bounded. In particular the z-unit argument's possible C pole does not occur here.

The N_x^2 diagonal has residue 2. The complete remaining non-Gamma cross terms of SP6 are

$$\begin{aligned} &- 4uqw(2m + 2D + 1)N_xN_y + uqwR_xN_xR \\ &\quad + qwB_xN_xA_\theta + uA_yN_yA_\theta + \chi uA_\theta R. \end{aligned} \tag{SF9}$$

Every coefficient is a bounded infinitesimal on u, w infinitesimal. Here $m, D, C, \beta\theta, p, l$ are bounded, $B_x = 8\kappa e$ is infinitesimal, and A_y is bounded. The entire N_yR term was already included in SF8 and CT7a. No first normal was suppressed.

169.1.3 A scalar circuit and two normal conductors with no state quotient

The original-core sign exit and exact testing sign gauges leave $u, w, W, Z > 0$. Literal $u = 0, w = 0, W = 0, Z = 0$ and the opposite-sign cases have their inherited direct core SOS. Suppose $\lambda > 0$. Define separated nonnegative monomials

$$a_c = \sqrt{\lambda\sigma}\, uZ, \qquad b_c = \sqrt{\rho}\, uwW, \qquad c_c = \sqrt{\lambda}\, xqW, \qquad g_c = (a_cb_cc_c)^{1/3}, \qquad s_c^3 = \frac{wZ}{zuW}. \tag{SF10}$$

Each radical is taken separately in its source and testing factors. The exact full angular packet in SP6 is

$$K_AA_\theta^2 + \chi uA_\theta R + \mathscr{P}, \qquad \mathscr{P} = \lambda\sigma P_\beta^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2.$$

The same algebraic HP3 identity is valid for infinitesimal z:

$$\mathscr{P} + \lambda xR^2 = a_c^2 + b_c^2 + c_c^2 - 3g_c^2 + \lambda xzu^2W^2s_c^2(s_c - 1)^2(s_c^2 + 2s_c + 3). \tag{SF11}$$

No bound on s_c or any state ratio is asserted or required. The last term is a legal SOS after writing $s_c^2 + 2s_c + 3 = (s_c + 1)^2 + 2$. Furthermore

$$a_c^2 + b_c^2 + c_c^2 - 3g_c^2 = \tfrac{1}{2}\{(a_c - b_c)^2 + (b_c - c_c)^2 + (c_c - a_c)^2\} + (a_cb_c + b_cc_c + c_ca_c - 3g_c^2). \tag{SF12}$$

The last parenthesis is the elementary cubic AM–GM SOS applied to $(a_cb_c)^{1/3}, (b_cc_c)^{1/3}, (c_ca_c)^{1/3}$. Expand its positive monomial prefactor before taking square roots. Thus

$$D_1 = (a_c - b_c)/\sqrt{2}, \qquad D_2 = (b_c - c_c)/\sqrt{2}$$

are two genuine unit-weight pivots; all other SF11–SF12 squares are retained separately.

The new root is

$$T = kq(u - k)W.$$

Its legal quadratic domination is exact:

$$A_\theta^2 - T^2 = q^2(u - k)^2W^2(u^2 + 2uk). \tag{SF13}$$

Both summands on the right are squares, using the separated root $\sqrt{u}\sqrt{k}$. If K_A has positive residue, allocate its diagonal exactly once as

$$K_AA_\theta^2 = \tfrac{1}{2}K_AA_\theta^2 + \tfrac{1}{2}K_AT^2 + \tfrac{1}{2}K_A(A_\theta^2 - T^2). \tag{SF14}$$

The last term is retained SOS and the first two supply strict positive source pivots for A_θ, T.

Here are the decisive two identities, valid for arbitrary states without a state division:

$$\boxed{\begin{aligned} u\Gamma &= \frac{\sqrt{2}k}{\sqrt{\rho}}D_1 - \frac{\sqrt{2}(u-k)}{\sqrt{\rho}}D_2 - \sqrt{\sigma}T, \\ qw\Gamma &= \sigma z R - \frac{\sqrt{2}(qw+z\sqrt{\sigma})}{\sqrt{\rho}}D_2 + \left(\frac{\sigma z}{k} + \frac{\sqrt{\sigma}qw}{k}\right)T. \end{aligned}} \tag{SF15}$$

All divisions are pure source. In particular neither q nor a testing coordinate is inverted in a conductor coefficient. These replace the $\lambda(Z/W)/q$ and w/q coefficients that obstruct a literal transport of the R19 z-unit proof.

169.1.4 Completion when H is bounded

Here $K_A \sim 2$ and λ is bounded. If d_0 has positive residue, the original $d\Gamma^2$ diagonal gives a strict Gamma pivot. Its two complete normals are

$$-4\nu N_x(u\Gamma) + R_\gamma(N_y + bR)(qw\Gamma). \tag{SF16}$$

Their coefficients are infinitesimal on (N_x, N_y, R, Γ). Keep SF11 entirely as SOS; its sole cost $-\lambda x R^2$ is an infinitesimal bounded radial correction. The frame $(N_x, N_y, R, \Gamma, A_\theta)$ has strict positive residue.

If d_0 is infinitesimal, ρ is a unit by SF4 and σ/x is bounded by SF7. All coefficients in both lines of SF15 are bounded infinitesimals. Indeed $\sqrt{\sigma}/k$ is bounded, $\sigma/k = k(\sigma/x)$ is infinitesimal, and q, w, z are bounded infinitesimals. Use SF14 and the two unit circuit pivots. After substituting SF15 into SF16, the complete frame

$$(N_x, N_y, R, A_\theta, T, D_1, D_2) \tag{SF17}$$

has strictly positive residue. Its radial block is SF8, its angular and T pivots come from SF14, and the circuit pivots from SF12. Every correction is a bounded infinitesimal. The diagonal $d\Gamma^2$ is retained separately, including at $d = 0$.

In both cases retain exactly once $4\beta(1+w^2)N_x^2$, $2(2D+1)(uX_e)^2$, $4bpzu^2\Gamma^2$, the unused circuit rows, and all positive radial and coefficient-cap excesses. Equations SP6, SF8–SF16 list every original scalar, normal and hard coupling. The separated bounded-Gram lemma proves the entire parent.

The d_0 split exhausts the source sphere. The bounded-M_β and bounded nonzero-unit Alice-angle exits are the inherited JB theorems; their CE remote join includes all other projective tests. Together with the literal-axis and sign exits, this closes the complete original C14.1a residual and the full bounded-H source domain. Apply the real argument independently to real and imaginary state parts and restore only positive pure-source and pure-testing normalizations.

169.1.5 A source factor restores the second normal when H is unbounded

Assume now H unbounded and $h_0 < 1/2$. SF3 gives a positive-unit K_A, so the same angular allocation is available. Also $\lambda > 0$, $\lambda/H \sim 2$, ρ bounded and $\lambda x = \rho\sigma$ infinitesimal. The positive-d_0 case is closed exactly as before: λ occurs only inside the retained scalar circuit, not as an unbounded coefficient of the remaining Gram.

For infinitesimal d_0, ρ is a unit, but $\sqrt{\sigma}/k$ can be unbounded. One must use the actual second normal in SF16, not claim boundedness of its factor $qw\Gamma$ by itself. Exactly

$$R_\gamma = R_0 - 8pu^2, \qquad R_0 = -4\{2(C+D)(r-1) + 2p + r - 1\}. \tag{SF18}$$

The required source bound is R_0/σ bounded. Here is a reproducible polynomial proof that retains every opening. In the P1 formulas temporarily leave e independent, and put $N = JPR_0$. The cancellation of $1-r$ in $C(r-1)$ shows that N is a polynomial in x, σ, z, b, h, e. Moreover $N|_{x=\sigma=e=0} = 0$. Define the polynomial quotients

$$F_x = \frac{N - N|_{x=0}}{x}, \quad F_\sigma = \frac{N|_{x=0} - N|_{x=\sigma=0}}{\sigma}, \quad F_e = \frac{N|_{x=\sigma=0}}{e}.$$

All their source arguments are bounded and the denominators JP are units. Restoring $e = xh/\sigma$ gives the exact expression

$$\frac{R_0}{\sigma} = \frac{(x/\sigma)F_x + F_\sigma + (xh/\sigma^2)F_e}{JP}. \tag{SF19}$$

The two ratios on the right are bounded by SF7; this proves the claim. The checker prints the three actual polynomials in its receipt so that this is not an unverifiable divisibility assertion.

Use the first conductor in SF15 unchanged. For the entire other normal use the exact identity

$$qwR_\gamma\Gamma = R_0(qw\Gamma) - 8pqwu(u\Gamma). \tag{SF20}$$

Substitute both SF15 expressions into SF20. Every resulting coefficient is infinitesimal. The only apparently dangerous ones are

$$R_0\frac{\sigma z}{k}, \qquad R_0\frac{\sqrt{\sigma}qw}{k}.$$

Write $R_0 = \sigma C_0$ with bounded pure-source C_0. SF7 makes σ/k bounded, so these are respectively $C_0\sigma z(\sigma/k)$ and $C_0\sqrt{\sigma}qw(\sigma/k)$, both infinitesimal. The other term in SF20 has the explicit infinitesimal factor qwu.

Thus SF17 again has strict positive residue, now using SF20 for the complete second Gamma normal. All retained squares and every other SP6 term are accounted for as in the bounded-H proof. The inherited R17 bounded-state and R18 unit-Alice admissions, together with the unchanged remote joins, complete all independent projective tests on this stated source subdomain. The argument never inverts a sphere coordinate and remains literal at $D = 0$.

169.1.6 Exact remaining complement and verification boundary

C14.1a has no remaining mathematical domain after this proof is combined with the recorded inherited testing exits. C14.1b remains open on its actual inherited complement intersected with

$$H \text{ unbounded}, \qquad \operatorname{res}(h) \geq \tfrac{1}{2}. \tag{SF21}$$

The earlier bounded-W/Z, sign, literal-axis and nonzero-unit-u admissions still remove their stated parts. Thus its real residual still has $u, w > 0$ infinitesimal, $WZ > 0$ and W/Z unbounded, as well as every original source intersection.

The specific obstruction to extending this proof is the angular diagonal: K_A has residue $2 - 4h_0$, which is zero or negative on SF21. Consequently SF14 supplies no strict positive angular or T pivot there. A new allocation of this signed angular packet is required. This failed frame is not a counterexample to tensor SOS and does not establish that SF21 is nonempty.

The focused file `check_state_free_conductors.py` checks the two state-free identities, the entire second normal substitution, the exact once-paid angular budget, the complete scalar circuit, its pair-difference pivots, and the source cancellation in SF19. It reports fifteen exact assertions. The complete original SP6 and CT7a identities remain inherited checked inputs. The analytical argument above, not the assertion count, establishes the quantified admissions. Review is same-system mathematical review, not external peer review or formal verification.

Chapter 170

C14.1b: the bounded testing-ratio complement at every unit h

Component `p5cd5fa89abcc`
Source: `BASE/proofs/c1_unbounded/c141b_bounded_testing_ratio.tex`
SHA256: `5cd5fa89abcc09162c1c8ff3f19cc62329522b5890615df07ba5de404204dccd`
Inventory use: BC1-BC11 exact bounded state/testing-ratio admission and retained complement | BC1–BC11 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

170.1 C14.1b: the bounded testing-ratio complement at every unit h

This is a further partial-domain admission, independent of the positive angular-pivot restriction in the preceding state-free proof. Keep the entire original physical C14.1b source, all strict gaps, original scale and prior-complement intersections, and all allowed coefficient openings. Thus $x, z = 1/L, \sigma = 1/\eta$ are positive infinitesimals, $H = h/\sigma$ is unbounded, and all source quantities belong to K_c, independently of tests and states in K_t. We assume here that h is a positive bounded unit. The claim is about the complete original P4/SP6 parent.

After the inherited literal-axis, state-sign and testing-sign exits, consider $u, w, W, Z > 0$, with u, w infinitesimal, and define exclusively in the testing field

$$c = \frac{uW}{wZ} > 0. \tag{BC1}$$

The new assertion is that every real state and test with c bounded is admitted. In particular this includes infinitesimal c, with no positive real lower bound on c. In the actual retained C14.1b complement one can additionally keep W/Z unbounded; nothing below needs that condition. A complex state is covered whenever each of its real and imaginary parts belongs to this domain or to an already admitted domain. No condition is imposed on a nonexistent complex order or complex state ratio.

Exact bounded source input. Use SF1–SF7 and the complete radial extraction SF8. They were proved with z infinitesimal and do not require a positive lower bound on z. In particular

$$\lambda > 0, \quad \lambda\sigma \sim 2h, \quad \rho = \lambda x/\sigma \text{ bounded}, \quad \lambda x \text{ infinitesimal}, \quad K_A \text{ bounded}. \tag{BC2}$$

No positivity of K_A is assumed. The exact sphere split gives a positive original Gamma pivot when d_0 has positive residue. When d_0 is infinitesimal, ρ is a positive unit and σ/k is bounded, where $k = \sqrt{x}$. The actual second Gamma normal has the source factorization SF18–SF20, with R_0/σ bounded. These facts include literal $D = 0$.

The complete scalar circuit and a new radial normalization. Write $q = \sqrt{z}$ and use exactly the three nonnegative separated monomials from SF10:

$$a_1 = \sqrt{\lambda\sigma}\, uZ, \qquad a_2 = \sqrt{\rho}\, uwW, \qquad a_3 = \sqrt{\lambda}\, xqW, \qquad G = (a_1a_2a_3)^{1/3}, \quad d_i = a_i - G. \tag{BC3}$$

Every fractional power is the product of its pure-source and pure-testing powers. The exact CT4 scalar identity applies without changing the source or ordering a mixed ratio: the scalar packet $\mathscr{P}$ of SF11 plus $\lambda x R^2$ is $\sum_i a_i^2 - 3G^2$ plus the two printed nonnegative CT4 squares. For $p_i = a_i^{1/3}$, its exact strict circuit decomposition is

$$\sum_i a_i^2 - 3G^2 = \sum_i d_i^2 + \sum_{j=1}^{3} \sum_{i<k} \{\sqrt{Gp_j}(p_i - p_k)\}^2. \tag{BC4}$$

The nine supplementary squares and the two CT4 squares remain whole. Thus the circuit supplies three unit diagonal pivots on d_1, d_2, d_3; its only cost is the infinitesimal source correction $-\lambda x R^2$.

Put

$$k_c = z^{1/3}c^{1/3}, \qquad \overline{J} = (k_c - 1)\frac{wZ}{q}. \tag{BC5}$$

The bar in $\overline{J}$ is part of the root's name, not a residue. It is a finite separated expression. Its exact relation to the original radial root is

$$R = \frac{wZ}{q} - uqW = -(1 + k_c + k_c^2)\overline{J}. \tag{BC6}$$

No mixed expression on the right is inverted. Because c is bounded, k_c and k_c^2 are bounded separated coefficients with zero real residue. Therefore the coefficient in BC6 has residue -1. After BC6 the clipped SF8 radial form is a bounded strict Gram on $(N_y, \overline{J})$, including the b and Q opening endpoints in residue. All exact positive radial excesses are retained.

This is precisely the rescaling $\overline{J} = J_c/(qc^{2/3})$ of CT6, but definition BC5 avoids an inverse of c even in the displayed root. It is valid when c is infinitesimal.

The signed angular packet has a small exact conductor. The following whole-state identity is the rescaled CT13 conductor:

$$\begin{aligned} A_\theta = {} & \frac{qcw}{\sqrt{\lambda\sigma}}d_1 + \frac{k_c^2}{\sqrt{\lambda}}d_2 - \frac{1}{\sqrt{\lambda}}d_3 \\ & + (k_c + 1)\sqrt{x}\, z^{2/3}c^{2/3}\overline{J} =: \mathcal{V}_A. \end{aligned} \tag{BC7}$$

All four coefficients are finite sums of bounded separated factors with zero residue. The first uses bounded c, infinitesimal q, w and the positive source unit $\lambda\sigma$. The second and third use $1/\sqrt{\lambda}$ infinitesimal. For the last, expand it as $\sqrt{x}\, z^{2/3}c^{2/3} + \sqrt{x}\, zc$. No bound involving an ordered product of independent rates occurs.

One can therefore replace the possibly signed angular coefficient by a strict reserve, using an identity rather than an inequality. For the SF13 root $T = kq(u-k)W$, exactly

$$\begin{aligned} K_A A_\theta^2 &= \tfrac{1}{2}A_\theta^2 + \tfrac{1}{2}T^2 + \tfrac{1}{2}(A_\theta^2 - T^2) + (K_A - 1)\mathcal{V}_A^2, \\ A_\theta^2 - T^2 &= q^2(u-k)^2W^2(u^2 + 2uk). \end{aligned} \tag{BC8}$$

The last expression in the second line is an exact separated SOS. The first two rows in the first line give strict pivots $1/2$ on A_θ, T. The signed final term has bounded infinitesimal Gram entries on $(d_1, d_2, d_3, \overline{J})$ by BC7 and bounded K_A. In particular this replacement remains valid at $\operatorname{res} h = 1/2$ and when K_A has negative residue. It is paid exactly once.

Every normal of the original parent. Keep the original N_x pivot and the exact SF8 radial extraction. If d_0 has positive residue, retain the strict original Gamma pivot and use its two full normals SF16 directly. Their coefficients are bounded infinitesimals, including after BC6. If d_0 is infinitesimal, use SF15 and the complete second-normal factorization SF20, just as in the state-free proof. Their circuit roots are now

$$D_1 = (d_1 - d_2)/\sqrt{2}, \qquad D_2 = (d_2 - d_3)/\sqrt{2}. \tag{BC9}$$

Thus no new division by a sphere coordinate is needed. SF7 and SF19 make every resulting coefficient bounded infinitesimal; A_θ, T have the strict reserves just supplied by BC8. The positive original $4D\Gamma^2/\sigma^2$ row is retained whole in this latter case.

All other SP6 terms are precisely the five complete crosses in SF9, the infinitesimal radial remainder CT7a, and the cost $-\lambda x R^2$. Substitute BC6 where R occurs. They remain bounded infinitesimal perturbations of the strict frame

$$(N_x, N_y, \overline{J}, A_\theta, T, d_1, d_2, d_3), \tag{BC10}$$

with Γ adjoined in the positive-d_0 case. The source-unit radial block, fixed N_x pivot, BC8 angular pivots, and BC4 circuit pivots give a positive-definite real residue. Retain exactly once $4\beta D_w N_x^2$, $2(2D+1)(uX_e)^2$, $4bpzu^2\Gamma^2$, all nonconstant positive radial packets and source-weight excesses, the nine BC4 squares, the two CT4 squares, and the positive BC8 domination remainder. No full state normal has been discarded. The separated bounded-Gram lemma now gives the complete parent as tensor SOS on the asserted domain.

The proof concerns every independent test/state in BC1 with c bounded. Literal axes and opposite-sign configurations keep their inherited admissions. Applying the real proof or an earlier exit separately to the real and imaginary parts gives the complex-state join stated at the start; it does not enlarge the real domain by assuming an order on complex amplitudes.

Exact improvement and retained obstruction. Combining this result with the previously accepted source subdomain $\operatorname{res} h < 1/2$, the real C14.1b remainder can be narrowed to its original retained source/test intersections with

$$\operatorname{res} h \geq 1/2, \qquad u, w > 0 \text{ infinitesimal}, \quad WZ > 0, \quad W/Z \text{ unbounded}, \quad \frac{uW}{wZ} \text{ unbounded}. \tag{BC11}$$

This complement still contains the difficult source-contact scale: on the actual source contact $uW/(wZ) = L = 1/z$, which is unbounded. For unbounded pure-testing c, BC7's coefficients involving c cannot be declared bounded separated coefficients, and the present argument does not cover that case. No full C14.1b closure or nonemptiness assertion is made.

The narrow checker verifies the exact radial relation, the complete angular conductor, the circuit decomposition and conversion, and the once-paid angular reserve. The source bounds, every remaining normal and the independent-field Gram argument are the analytical proof above, with precisely the SF and CT dependencies identified.

Chapter 171

C14.1b: the remaining source at bounded Alice-to-reciprocal-Bob ratio

Component `p50b4df0bb619`
Source: `BASE/proofs/c1_unbounded/c141b_gamma_angular_domination.tex`
SHA256: `50b4df0bb619d10389dbddf2ac9c3de366990af3f598368843141cec5fe02b81`
Inventory use: GA1-GA9 surviving bounded-u/w admission and exact remaining domain | GA1–GA9 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

171.1 C14.1b: the remaining source at bounded Alice-to-reciprocal-Bob ratio

Exact additional scope. Keep the full original C14.1b source, physicality, strict gaps, nonnegative coefficients, total-scale and prior-complement intersections. Use the independent fields and the complete P4/SP6 parent of the inherited `parent_reference.tex`. In this section h is a positive bounded unit and $a_0 = k/\sigma$ is infinitesimal. The unit-a_0 source was already admitted by US1–US15; no conclusion about that proof is changed here.

Theorem 171.1. *On the stated source, the entire original parent is tensor SOS for every projective testing configuration whose remaining double-small-angle chart has $u, w > 0$ infinitesimal and u/w bounded, and for every real or complex state. Include the exact sign gauges, literal endpoints and other testing exits already admitted in SF1–SF21. There is no restriction on a state ratio or on $c = uW/(wZ)$ in this theorem.*

This is an additional testing-domain admission. It does not close the remaining chart u/w unbounded. Unlike BC1–BC11, its testing predicate is independent of the state, so the same predicate holds for the real and imaginary state components.

Available source pivots. The source bounds SF1–SF5 and US1–US3 give

$$L_0 = \lambda\sigma \sim 2h, \qquad \lambda^{-1} = \sigma/L_0 \text{ infinitesimal}, \qquad \rho = L_0 a_0^2 \text{ infinitesimal}, \qquad d = 4D/\sigma^2 \sim 2. \tag{GA1}$$

Indeed US2 first bounds a_0, and its exact residue identity makes b_0 infinitesimal when a_0 is; the sphere then gives $d_0 \sim 1$. Thus the original Gamma diagonal supplies a strict positive-unit pivot. Also $\lambda x = \rho\sigma$ is infinitesimal. The source coefficients $K_A, \chi, \nu, R_\gamma, R_x, A_y$ are bounded. No lower cap on β, b or Q is imposed. A literal $D = 0$ face is incompatible with GA1, rather than being removed from a domain where it could occur.

A paid angular domination from the original Gamma square. The unchanged sign and state-axis exits leave $u, w, W, Z > 0$; their ratios may otherwise be arbitrary. Set $t_s = \sqrt{\sigma}$ (a source

letter, unrelated to the earlier notation $t = u^2$), and define

$$E_0 = quW - t_sZ, \quad V = uE_0, \quad \alpha_t = q\frac{u}{w}, \quad U = \frac{B_1 - B_3}{\sqrt{\lambda}} = \frac{d_1 - d_3}{\sqrt{\lambda}}. \tag{GA2}$$

Here $B_i, G, d_i = B_i - G$ are exactly the four monomials and circuit differences in US7–US10. Every root in GA2 is separated. The only new testing quotient is u/w, which is bounded by the theorem's predicate. Hence α_t is a bounded infinitesimal coefficient. Exactly

$$\Gamma + t_sR = -\frac{w + t_sq}{q}E_0, \qquad A_\theta = U + V. \tag{GA3}$$

The first identity is used as a polynomial identity; its mixed factor $w + t_sq$ is never inverted. In fact it yields the explicit separated-SOS difference

$$\begin{aligned}\Delta &:= \alpha_t^2(\Gamma + t_sR)^2 - V^2 \\ &= u^2E_0^2\left\{\frac{2t_sq}{w} + \left(\frac{t_sq}{w}\right)^2\right\} \\ &= \left(\sqrt{2}\sqrt{t_s}\sqrt{q}\,\frac{u}{\sqrt{w}}E_0\right)^2 + \left(\frac{t_squ}{w}E_0\right)^2.\end{aligned} \tag{GA4}$$

All square roots and inverses in the last line belong to their individual fields. The roots themselves need not be bounded.

Choose a positive real C_* so that the bounded pure-source coefficient $K_A + C_*$ has positive real residue. The following once-paid identity allocates the complete signed angular diagonal:

$$\boxed{\begin{aligned}K_AA_\theta^2 = (K_A + C_*)A_\theta^2 - \frac{2C_*}{\lambda}(d_1 - d_3)^2 - 2C_*\alpha_t^2(\Gamma + t_sR)^2 \\ + C_*(U - V)^2 + 2C_*\Delta.\end{aligned}} \tag{GA5}$$

It follows simply from $A_\theta = U + V$ and GA4. The last two terms are retained SOS. The two negative terms have bounded infinitesimal coefficients on the circuit and (Γ, R) frames. This constructs a strict angular pivot without requiring a positive residue of K_A, and without the collapsed λk pivots of US11.

Complete scalar, every state normal, and all excesses. Use the whole four-monomial allocation once:

$$\mathscr{S} = \sum_{i=1}^{4} d_i^2 + \mathcal{E}_4 - \lambda xR^2. \tag{GA6}$$

Here $\mathcal{E}_4$ is the explicit separated SOS in US10. Do not also allocate US6 or US11. The single radial cost in GA6 is an infinitesimal pure-source coefficient.

Use SF8/CT7–CT7a with the full source-opening-safe radial squares and every positive excess restored. The resulting strict bounded radial frame is (N_y, R), uniformly through the b, Q openings. Together with the original N_x pivot, $d\Gamma^2$ from GA1, GA5 and GA6, the principal frame is

$$(N_x, N_y, R, \Gamma, A_\theta, d_1, d_2, d_3, d_4). \tag{GA7}$$

Its residue is strictly positive definite: the N_x residue is 2, the Gamma residue is 2, the angular coefficient is $K_A + C_*$ with positive residue, and the four circuit coefficients are 1. All costs in GA5–GA6 and all CT7a remainders are bounded infinitesimal matrix entries on this frame.

The two complete Gamma normals remain exactly

$$-4\nu N_x(u\Gamma) + R_\gamma(N_y + bR)(qw\Gamma). \tag{GA8}$$

Their coefficients are infinitesimal directly because u, q, w are infinitesimal and the remaining factors are bounded. No division by $\sqrt{\rho}$ is used. The five remaining non-Gamma crosses are precisely SF9,

with bounded infinitesimal coefficients on GA7. The whole N_yR coefficient has already been included in the exact radial extraction. Thus every original SP6 term, including both first state normals, is accounted for.

Keep separately and exactly once $4\beta D_w N_x^2$, $2(2D+1)(uX_e)^2$, the nonconstant positive N_x terms, $4bpzu^2\Gamma^2$, all radial and source-weight excesses, $\mathcal{E}_4$, and the last two SOS terms of GA5. The separated bounded-Gram lemma applied to GA7 proves the complete parent.

The exact inherited literal, sign and remote exits supply the complementary charts specified in the theorem. For complex states, apply the same real theorem separately to each real state part; the hypothesis u/w bounded depends only on the test and is unchanged. Restore only positive pure-source and pure-testing normalizations. No source/testing rate is identified or compared.

Exact updated residual and failed extension. Combining this theorem with SF, BC and US leaves C14.1b only on the original retained complement intersected with

$$\begin{gathered} \operatorname{res} h \geq \tfrac{1}{2}, \quad k/\sigma \text{ infinitesimal}, \quad b_0 \text{ infinitesimal}, \quad d_0 \sim 1, \\ u, w > 0 \text{ infinitesimal}, \quad WZ > 0, \quad W/Z \text{ unbounded}, \\ c = uW/(wZ) \text{ unbounded}, \qquad u/w \text{ unbounded}. \end{gathered} \tag{GA9}$$

The exact obstruction to this particular extension is $\alpha_t = q(u/w)$ in GA2: its testing factor is no longer bounded. It cannot be declared small by comparing the independent source q to the testing reciprocal w/u. At contact $u_c/w_c = a_0\sqrt{\sigma}/q$; a further lawful source split by $q/(a_0\sqrt{\sigma})$ may be useful, but no all-tests conductor or join for that split has been proved here. GA9 is not asserted nonempty, and full C14.1b remains open.

The focused checker verifies GA3–GA5 and the complete scalar budget in GA6. Analytical source bounds, every parent term and the exact quantified testing join are the proof above; this remains same-system mathematical review.

Chapter 172

C14.1b: the entire unit beta-sphere source, including the contact ratio

Component `p0931b3af19ec`
Source: `BASE/proofs/c1_unbounded/c141b_unit_beta_sphere_all_tests.tex`
SHA256: `0931b3af19ec66be88995b22a0c20bb46df7a7b25bf0280d924d009d0624c147`
Inventory use: US1-US3 source bound k/sigma, US7-US10 circuit, complete unit-beta-sphere admission including D=0 | US complete theorem | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

172.1 C14.1b: the entire unit beta-sphere source, including the contact ratio

The precise whole-source subdomain. Keep the entire original physical C14.1b source, its strict gaps, nonnegative event weights, original total-scale and prior-complement intersections. In particular $x, z = 1/L, \sigma = 1/\eta$ are positive infinitesimals. Write $k = \sqrt{x}$, $q = \sqrt{z}$ and retain all P1–P8/SP6 notation. Source and testing fields are independent.

Theorem 172.1. *Suppose h and k/σ are positive bounded units. Then the complete original parent is finite tensor SOS at every independent projective testing pair and every real or complex state. All permitted literal coefficient boundaries and all positive opening rates are retained. There is no bound on $uW/(wZ)$ or any other testing/state ratio in this theorem.*

This includes the source-contact scale $uW/(wZ) = 1/z$, which was not covered by the bounded-testing-ratio theorem. The theorem is a whole source-subdomain admission, not closure of all C14.1b.

Source entry and its sphere interpretation. The bounds SF1–SF5 remain valid. Since h is a unit and σ infinitesimal, H is unbounded and

$$\begin{gathered}\lambda > 0, \quad L_0 := \lambda\sigma \sim 2h, \quad a_0 := k/\sigma \text{ a positive unit}, \\ \rho = \lambda x/\sigma = L_0 a_0^2 \text{ a positive unit}, \quad \lambda k = L_0 a_0 \text{ a positive unit}.\end{gathered} \tag{US1}$$

Also $\lambda x = \rho\sigma$ is infinitesimal. The subscript in a_0 is part of this section's source letter, not a residue. The bounded source quantities $K_A, \chi, \nu, R_x, A_y, R_\gamma$ have the exact meanings in SP6; K_A may have either sign.

On the entire unit-h source, the earlier exact source bound $e = xh/\sigma = O(x + \sigma)$ first gives $x/\sigma = e/h$ infinitesimal. SF5 gives the exact identity

$$ha_0^2 = \frac{x}{\sigma}\frac{H_0}{B} + \frac{\mathcal{H}}{B}b_0^2. \tag{US2}$$

Since h is a positive unit and the right-hand factors in US2 are bounded, this identity bounds a_0^2 without assuming it bounded in advance. It also shows

$$\operatorname{res}(ha_0^2) = \operatorname{res}(b_0^2). \tag{US3}$$

Thus this theorem covers precisely the unit-h source with the beta-sphere coordinate b_0 a positive unit. Literal $D = 0$ has $b_0 = 1$ and is included. There is no division by D, β or either sphere coordinate. A source face incompatible with these exact hypotheses is not assumed to have points.

Two exact allocations of the same complete scalar packet. The inherited remote and literal/sign exits leave $u, w, W, Z > 0$ with u, w infinitesimal. No size restriction on W/Z is needed below. The entire scalar contact packet in SP6 is

$$K_A A_\theta^2 + \chi u A_\theta R + \mathscr{S}, \qquad \mathscr{S} = \lambda\sigma P_\beta^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2. \tag{US4}$$

Define legal separated roots

$$J_A = q(u - k)W, \quad F_A = u(Z - a_0 wW), \quad E_u = \sqrt{u}\, J_A, \quad E_k = \sqrt{k}\, J_A. \tag{US5}$$

The following identity is exact for the complete scalar, with no restriction to a state graph:

$$\boxed{\mathscr{S} = L_0 F_A^2 + 2\lambda k E_u^2 + \lambda k E_k^2 + 2\lambda k\sqrt{u}\, E_u R.} \tag{US6}$$

One direct derivation expands $P_\beta = uZ - \sigma a_0^2 wW$ and uses $wZ/q = R + uqW$. The coefficient of q^2W^2 that remains is

$$a_0(2u^3 - 3ku^2 + k^3) = a_0(2u + k)(u - k)^2,$$

which gives US6. Its diagonal coefficients in US1 are positive units, and its sole cross coefficient contains the testing infinitesimal $\sqrt{u}$.

For a complementary allocation set

$$\begin{gathered} B_1 = \sqrt{\lambda\sigma}\, uZ, \quad B_2 = \sqrt{\rho}\, uwW, \quad B_3 = \sqrt{\lambda}\, xqW, \quad B_4 = \sqrt{\lambda x}\, \frac{wZ}{q}, \\ G = (B_1B_2B_3B_4)^{1/4} = \sqrt{\lambda x}\sqrt{uwWZ}, \qquad d_i = B_i - G. \end{gathered} \tag{US7}$$

All radicals separate; in particular $\sqrt{uwWZ}$ is entirely testing. The pure-source inverse $1/q$ in a root is allowed. Its root need not be bounded, and it is not used in a coefficient. Exactly

$$\boxed{\mathscr{S} + \lambda x R^2 = \sum_{i=1}^{4} B_i^2 - 4G^2.} \tag{US8}$$

The four-term circuit has strict difference pivots:

$$\sum_i B_i^2 - 4G^2 = \sum_i d_i^2 + \mathcal{E}_4, \tag{US9}$$

where the following is explicitly tensor SOS:

$$\mathcal{E}_4 = 2G\left\{(\sqrt{B_1} - \sqrt{B_2})^2 + (\sqrt{B_3} - \sqrt{B_4})^2 + 2\big((B_1B_2)^{1/4} - (B_3B_4)^{1/4}\big)^2\right\}. \tag{US10}$$

The prefactor G is a nonnegative separated monomial, so it too has a separated square root. The circuit does not assert an order on a mixed coefficient.

Use each allocation with weight one-half. The exact paid budget is

$$\begin{aligned} \mathscr{S} = {} & \tfrac{1}{2}L_0F_A^2 + \lambda k E_u^2 + \tfrac{1}{2}\lambda k E_k^2 + \lambda k\sqrt{u}\, E_u R \\ & + \tfrac{1}{2}\sum_i d_i^2 + \tfrac{1}{2}\mathcal{E}_4 - \tfrac{1}{2}\lambda x R^2. \end{aligned} \tag{US11}$$

There is no repeated use of a scalar square. The only radial cost is the infinitesimal pure-source coefficient $-\lambda x/2$.

The complete angular and Gamma conductors. The angular root has the exact identity

$$A_\theta = \sqrt{u}\,E_u + \sqrt{k}\,E_k. \tag{US12}$$

Its coefficients are bounded infinitesimals. Therefore the entire possibly signed $K_A A_\theta^2$, and the complete $\chi u A_\theta R$ term, give bounded infinitesimal corrections to the strict frame supplied by US11 and the radial block. This works also when $\operatorname{res} h \geq 1/2$ and K_A has zero or negative residue.

Both first Gamma normals have small exact conductors on the same frame:

$$\boxed{\begin{aligned} u\Gamma &= \frac{k}{\sqrt{\rho}}(d_1 - d_2) - \frac{u-k}{\sqrt{\rho}}(d_2 - d_3) - \sqrt{\sigma k}\,E_k, \\ qw\Gamma &= \frac{qw + z\sqrt{\sigma}}{\sqrt{\rho}}(d_4 - d_2) - \sqrt{\sigma}\,qwR. \end{aligned}} \tag{US13}$$

The first is SF15 with $T = kJ_A = \sqrt{k}E_k$; the second follows directly from the two monomials B_4, B_2. Every displayed coefficient is a finite sum of bounded separated factors with zero residue. No state quotient, testing inverse, source/test rate comparison, or sphere-coordinate inverse occurs. In particular these formulas do not use a lower bound on the original Gamma diagonal.

For comparison, US7 also yields the exact global identity $T = (B_4 - B_3)/\sqrt{\lambda} - kR$. This is another small conductor on the present source and explains why the extra fourth monomial removes the coefficient failure of the earlier three-term circuit. The proof uses US13 directly.

Full-parent restoration and strictness. Use the exact opening-safe SF8/CT7–CT7a radial decomposition. Its clipped pure-source Gram on (N_y, R) is strict through the b, Q openings, with all excess squares retained. Keep the original N_x pivot, whose residue is 2. The principal frame is

$$(N_x, N_y, R, E_u, E_k, d_1, d_2, d_3, d_4). \tag{US14}$$

US11 gives positive-unit pivots λk and $\lambda k/2$ on E_u, E_k, and pivots $1/2$ on the four circuit differences. The F_A square and $\mathcal{E}_4/2$ are retained separately.

The two entire Gamma normals of SP6 are

$$-4\nu N_x(u\Gamma) + R_\gamma(N_y + bR)(qw\Gamma).$$

Their remaining coefficients are bounded, so US13 allocates them with infinitesimal coefficients on US14. Retain the entire nonnegative $(4D/\sigma^2 + 4bpzu^2)\Gamma^2$, including its literal zero cases. The five non-Gamma crosses in SF9 are all bounded infinitesimal after US12; the whole $N_y R$ coefficient was already included in the radial extraction. The radial remainders CT7a are infinitesimal on z infinitesimal, as in SF8. US11 adds only its printed infinitesimal radial cost and infinitesimal $E_u R$ cross.

Thus every full-parent term is either an explicitly retained SOS or a bounded Gram entry on US14, whose real residue is strictly positive definite. The separated bounded-Gram lemma gives tensor SOS. Restore exactly once the original $4\beta D_w N_x^2$, $2(2D+1)(uX_e)^2$, the positive nonconstant N_x terms, every radial/source-weight excess, the original Gamma diagonal, and the two retained nonnegative terms of US11. No hard state normal or signed scalar term is omitted.

This proves the complete double-small-angle chart for every state left by the exact sign/axis exits, including $D = 0$. The unchanged remote, nonzero-unit-Alice and other projective exits listed in the SF dependency index cover its complement. Hence the whole original source subdomain stated in the theorem is covered at every test. The real theorem applies separately to arbitrary real and imaginary state parts, with no state-ratio predicate, proving the full complex claim as well.

The exact remaining source and testing obstruction. Combine this theorem with the source subdomain $\operatorname{res} h < 1/2$ and the bounded-c theorem BC1–BC11. The remaining real C14.1b complement is contained in the original retained source/test domain with

$$\operatorname{res} h \geq 1/2, \quad k/\sigma \text{ infinitesimal}, \quad b_0 \text{ infinitesimal}, \quad d_0 \sim 1, \quad \frac{uW}{wZ} \text{ unbounded}, \tag{US15}$$

and the previous $u, w > 0$ infinitesimal, $WZ > 0$, unbounded-W/Z conditions. Here the new angular pivots $\lambda k = L_0 k/\sigma$ lose their real residue, and the US13 coefficients involving $1/\sqrt{\rho}$ are no longer bounded. The original Gamma diagonal now has a positive-unit pivot, which is a concrete possible reserve for a further completion, but this proof does not allocate it to the lost angular roots. No nonemptiness or full C14.1b closure is asserted.

The focused checker verifies US6, US8–US13, the exact half-and-half budget and the fourth-monomial T conductor. Source bounds and all-test restoration are the analytical proof above, with the exact inherited SF/CT inputs stated. Review remains same-system review.

Chapter 173

A full-parent terminal for a finite positive-d commuting edge

Component `p4ec1a51262f9`
Source: `BASE/proofs/routing/a5_recovery/POSITIVE_D_FINITE_EDGE_TERMINAL.md`
SHA256: `4ec1a51262f94608a1d92075518717c3d22958300ea88636fd972ea2ffea3503`
Inventory use: Complete retained finite-edge critical-center proof including every state normal and full tail

173.1 A full-parent terminal for a finite positive-d commuting edge

25 September 2026. New work; the round5 package is unchanged.

This theorem closes the nondegenerate finite-angle part of a distance-three jet face when one of its two omitted events is allowed to open. It applies to the C1 and C2 routing wherever the hypotheses below hold. It does not assert completion of their projective endpoints or flat core boundaries.

173.1.1 1. Precise separate-field statement

Let the coefficients belong to the bounded valuation ring of a real closed extension $K_c \supset \mathbb{R}$, and let an independent measurement/state test belong to the bounded valuation ring of $K_s \supset \mathbb{R}$, with projective charts used as necessary. Assume the coefficient vector is physical over K_c, is nonnegative, and satisfies **exactly**

$$\lambda = 0.$$

Its real residue c_0 is assumed to satisfy

$$d_0 > 0, \qquad (e_2)_0 = 0. \tag{1}$$

There are no exact support requirements on $e_1, e_3, e_4, e_5, \alpha, \beta$. In particular any of them may open infinitesimally from zero. On the exact residue face $\lambda = e_2 = 0$, write the known determinant decomposition as

$$Q_{c_0}(U,V) = UE(V) + U^2 B(V), \quad E(V) = q_{10} + q_{11}V + q_{12}V^2, \quad B(V) = q_{20} + q_{21}V + q_{22}V^2. \tag{2}$$

Fix a real $v_0 \neq 0$, and suppose

$$q_{10} > 0, \quad q_{12} > 0, \quad E(v_0^2) = 0, \quad B(v_0^2) > 0. \tag{3}$$

Also assume the three-state complement $M_0(v_0)$ in (5) below is positive definite. These are entirely real residue conditions. The latter condition is automatic if $\alpha_0 + (e_5)_0 > 0$, and is displayed explicitly to avoid an unproved support-exhaustion claim.

For every independent test with

$$\bar{u} = 0, \qquad \bar{v} = v_0, \qquad [\bar{W} : \bar{X} : \bar{Y} : \bar{Z}] = [1 : 0 : 0 : 0], \tag{4}$$

the original full slack is a finite sum of squares in the original bounded tensor ring $B_c \otimes_{\mathbb{R}} B_s$. All coefficient-opening rates and all test rates in this chart are allowed. No quotient by an infinitesimal test quantity or mixed tensor element is used.

The same statement with $E(v_0^2) > 0$ is the immediate positive-definite four-normal exit described in section 7.

173.1.2 2. Event normalization and the complete polynomial

Use the full parent in coefficient order

$$(d, \lambda, e_1, e_2, e_3, e_4, e_5, \alpha, \beta).$$

The event amplitude $R_2 = vW - X + uvY - uZ$ is a pure tested unit in (4). Divide the state by R_2. This is a legal pure-support unit operation and sets

$$R_2 = 1, \qquad W = (1 + X - uvY + uZ)/v.$$

The entire opened-event contribution is now the **constant** $4e_2$. Call the resulting full raw slack $(\mathrm{F}X, Y, Z, u, v)$. Its coefficients are polynomials in the point coefficients and Laurent polynomials in v. The only denominators are powers of the pure tested unit v.

At $u = 0$, its state Hessian divided by two is

$$M_0(v) = \begin{pmatrix} 2(d + 2e_1 + 2\beta)(1 + v^2) & 0 & 0 \\ 0 & 2d(1 + v^2) + 4(e_3 + e_4)(1 + v^2) + 4e_5 + 4\alpha v^2 & 4v(e_5 - \alpha) \\ 0 & 4v(e_5 - \alpha) & 4\alpha + 4e_5 v^2 \end{pmatrix}. \tag{5}$$

Its lower block is positive definite if $d > 0$ and

$$\alpha d + 2\alpha e_3 + 2\alpha e_4 + 2\alpha e_5(1 + v^2) + e_5 v^2(d + 2e_3 + 2e_4) > 0.$$

Introduce **coefficient-side center variables** x, a, b, U by the formal substitution

$$X = u^2 x, \quad Y = ua, \quad Z = ub, \quad U = u^2.$$

The exact full identity is

$$F(u^2 x, ua, ub, u, v) = 4e_2 + u^2 K(x, a, b, v, U), \quad U = u^2. \tag{6}$$

Here K is polynomial in x, a, b, U, with Laurent-polynomial v coefficients. More strongly,

$$K_x = UE_x, \qquad \partial_x E_x\big|_{U=0} = 4(d + 2e_1 + 2\beta)(1 + v^2) > 0, \tag{7}$$

and $(\mathrm{K}x, a, b, v, 0)$ is independent of x. These statements have been verified directly on all 105 terms of K, starting from all 106 terms of the normalized full parent. The formal substitution defines the center; it is never used to form $X/u^2, Y/u, Z/u$ in the independent tensor.

173.1.3 3. The residue center and positive quartic coefficient

At the real residue, minimizing the positive state block in a, b gives a scalar jet profile

$$r(v) = \frac{16(1+v^2)^2}{v^2 \det M_0(v)} E(v^2). \tag{8}$$

Physicality and (3) imply

$$E(V) = q_{12}(V - v_0^2)^2.$$

Consequently $r(v_0) = r'(v_0) = 0$, $r''(v_0) > 0$. The Hessian of

$$K_0(a, b, v) = K(x, a, b, v, 0)$$

in (a, b, v) is positive definite at its unique local zero

$$(a_0, b_0, v_0).$$

This follows by completing the strictly positive (a, b) quadratic and using $r''(v_0) > 0$, so it controls the full Hessian, not just its determinant. Equation $E_x = 0$ then determines a unique x_0 by (7).

Set $q = (x, a, b, v)$. The Jacobian of

$$\mathcal{E}(q, U) = (E_x, K_a, K_b, K_v) \tag{9}$$

in q at $(c_0, q_0, 0)$ is block upper triangular. Its top-left entry is the positive number in (7), and its lower 3×3 block is the positive Hessian just proved. Thus it is invertible. The Nash implicit function theorem gives a coefficient-only local branch $q_*(c, U)$.

The optimized function

$$k(c, U) = K(q_*(c, U), U)$$

satisfies

$$k(c_0, 0) = 0, \qquad \gamma := k_U(c_0, 0) = \frac{16(1+v_0^2)^2}{v_0^2 \det M_0(v_0)} B(v_0^2) > 0. \tag{10}$$

For clarity, (10) uses the whole parent determinant and the positive state block. The event-normalized four-by-four matrix has determinant

$$\det P / v^2.$$

Its state Schur complement is its determinant divided by the positive three-state determinant. At $e_2 = 0$, (2) therefore gives (8), and at the zero in (3) its next U^2 coefficient is exactly (10). The equations for a, b optimize the order-U state terms; the divided equation for

x optimizes the order-U^2 state term. No inverse of this state block is used in the final tensor identity; this real calculation only proves strict positivity of the residue coefficient.

173.1.4 4. Two coefficient-only centers

Put $\kappa = k(c, 0)$. It is a pure coefficient infinitesimal, with either sign allowed.

If $\kappa \geq 0$, take $q_c = q_*(c, 0)$ and $U_c = 0$. The ordinary derivatives K_a, K_b, K_v vanish there, and $E_x = 0$.

If $\kappa < 0$, solve

$$\mathcal{E}(q_c, U_c) = 0, \qquad K(q_c, U_c) + U_c K_U(q_c, U_c) = 0. \tag{11}$$

The full Jacobian at the real base point is block upper triangular with the invertible block (9) and final diagonal entry $2\gamma > 0$. The derivatives of the final equation in q vanish at the base point. Hence there is a unique nearby coefficient-only solution.

It satisfies $U_c > 0$. Indeed along the normal branch the last equation is $g(U) = k(U)+Uk'(U) = 0$, with $g(0) = \kappa < 0$ and derivative having positive residue 2γ. This argument is local and transfers to the real closed coefficient field. It uses no comparison with the independent tested u.

Define

$$\mu = 4e_2 + U_c K(q_c, U_c). \tag{12}$$

This is nonnegative: realize $u_c = \sqrt{U_c}$, $v = v_c$,

$$(X, Y, Z) = (U_c x_c, u_c a_c, u_c b_c),$$

and restore W by the event normalization. This is an actual coefficient-side state and measurement pair, so full physicality applies.

173.1.5 5. Exact full-tail identity in the original independent tensor

Keep q_c fixed and put

$$N_x = X - u^2 x_c, \quad N_y = Y - u a_c, \quad N_z = Z - u b_c, \quad T = v - v_c, \quad N_v = uT. \tag{13}$$

All these belong to the original bounded tensor ring. No small quantity has been divided out.

The state dependence of the normalized full parent is exactly quadratic. Consequently its entire expansion is

$$F = 4e_2 + UK(q_c \text{ with } v \text{ variable}, U) + UE_x N_x + uK_a N_y + uK_b N_z + N_s^T M N_s, \quad N_s = (N_x, N_y, N_z), \tag{14}$$

where the derivative functions on the right are evaluated at

(x_c, a_c, b_c, v, U). The independent checker verifies this identity, all three linear coefficients, and the complete quadratic remainder.

At $v = v_c, U = U_c$ all four center equations vanish. Exact difference division therefore writes the linear terms as combinations of

$$N_x U(U - U_c), \quad N_y u(U - U_c), \quad N_z u(U - U_c), \quad N_x u N_v, \quad N_y N_v, \quad N_z N_v. \tag{15}$$

The coefficient functions are bounded. Laurent denominators involve only the pure units v, v_c; differences of their powers are divided using finite algebraic identities. Taylor division of the profile in v gives an additional term $N_v^2 h$ and a cross term

$$N_v u(U - U_c)h_1. \tag{16}$$

Thus all cross terms with the last scalar generator have an extra u or U, and have zero residue at every independent test rate.

For $\kappa \geq 0$, univariate division at $U = 0$ gives the exact form

$$F = 4e_2 + \kappa u^2 + G^T H G, \quad G = (N_x, N_y, N_z, N_v, U). \tag{17}$$

For $\kappa < 0$, stationarity in (11) and (12) gives a double zero of the fixed-center univariate profile at U_c. Monic polynomial division $its coefficients may contain only the pure coefficient unit (v_c)$ therefore gives

$$F = \mu + G^T H G, \quad G = (N_x, N_y, N_z, N_v, U - U_c). \tag{18}$$

In both cases the residue Gram is positive definite. Its first block is the positive state-X entry together with one half of the positive Hessian of $K_0(a, b, v)$; its last entry is $\gamma > 0$. The cross entries to the last generator vanish by (15)–(16). This statement also accounts for the possibly nonzero residue cross terms $N_y N_v, N_z N_v$, which are part of the positive full jet Hessian.

The bounded positive-residue Gram lemma from round5 now makes

$G^T H G$ an SOS in $B_c \otimes B_s$. The remaining coefficients e_2, κ, μ are nonnegative pure coefficient quantities in their respective branches and hence squares. Finally multiply the roots by the original pure tested event amplitude to undo normalization. If normalized physical slack rather than raw slack is desired, the norm and measurement denominators are positive pure tested units on this chart and can likewise be restored. This proves section 1.

173.1.6 6. Nonvacuity and exact checks

The real residue

$$(d, \lambda, e_1, e_2, e_3, e_4, e_5, \alpha, \beta) = (2, 0, 0, 0, 1, 1, 1, 1, 0)$$

is physical, lies in both the C1 and C2 omission masters, and has

$$Q = 16U\{4UV + 3(V-1)^2\}.$$

At $v_0 = 1$, the center is

$$(x_0, a_0, b_0) = (-1/2, 1/2, 1), \qquad \gamma = 2.$$

The divided x pivot is 16, and the full lower Hessian is

$$\begin{pmatrix} 64 & 0 & 48 \\ 0 & 16 & 16 \\ 48 & 16 & 64 \end{pmatrix} \succ 0.$$

`verify_positive_d_finite_edge.py` rederives the full parent, event normalization, divided center equations, exact linear jets, and every state-quadratic tail term. Its report is `POSITIVE_D_FINITE_EDGE_CHECKS.json`: **15 identities, 30 scalar entries, all passed**. It includes the displayed example and both independent calculations of its positive quartic coefficient.

173.1.7 7. Exact scope of the remaining cases

If $E(v_0^2) > 0$, the original four-normal expansion in

(X, Y, Z, u) after event normalization has a positive residue Gram. The term $4e_2$ remains separate, so this strict finite-angle exit needs no critical center. If the tested state has positive residue loss, the scalar bounded-positive-residue lemma applies immediately.

The theorem above treats a finite nonzero zero of a nonzero quadratic edge polynomial with both endpoint coefficients positive. It leaves:

- $v_0 = 0$ or $v_0 = \infty$, where the normalization or the jet Hessian changes;
- flat outside/core cases $E \equiv 0$, where the v Hessian vanishes;
- distance-four corner jets that are not relabelings of this finite-edge configuration;
- coefficient charts in which both events of the chosen distance-three omission pair are nonzero. For applications to C1/C2 extreme routing, a permanently omitted event must first be included in the chosen pair.

No density, continuity of certificates, or unproved global gluing is used to include those cases.

Chapter 174

A positive-d terminal at a finite-ratio distance-four corner

Component p03b551f2a9a0
Source: BASE/proofs/routing/a5_recovery/POSITIVE_D_FINITE_RATIO_CORNER_TERMINAL.md
SHA256: 03b551f2a9a0b78d26feaa79831fb57bdc509e8e63ec108b08b3e3b73fd2b4b7
Inventory use: Complete retained finite-ratio critical-center proof and full 55-monomial normal allocation

174.1 A positive-d terminal at a finite-ratio distance-four corner

25 September 2026. This is new round6 work. It uses the same coefficient-only critical-center argument as POSITIVE_D_FINITE_EDGE_TERMINAL.md, with a separate exact full-tail calculation. It does not cover ratio endpoints or flat jet boundaries.

174.1.1 1. Hypotheses and conclusion

The physical nonnegative coefficient vector belongs to B_c^9, where B_c is the bounded valuation ring of a real closed extension of the reals. The independent test belongs to the corresponding ring B_s of another real closed extension. Suppose

$$\lambda = 0 \quad \text{exactly}, \qquad d_0 > 0, \quad \alpha_0 = 0. \tag{1}$$

All other coordinates may have arbitrary nonnegative residues or opening rates. At this distance-four residue face write

$$Q_{c_0}(U,V) = q_{20}U^2 + q_{11}UV + q_{02}V^2 + q_{21}U^2V + q_{12}UV^2 + q_{22}U^2V^2.$$

For a real $t_0 \neq 0$, assume

$$q_{20}, q_{02} > 0, \quad q_{20}t_0^4 + q_{11}t_0^2 + q_{02} = 0, \quad q_{21}t_0^4 + q_{12}t_0^2 > 0. \tag{2}$$

Let the exact quadratic jet $J_4(t_0)$ have kernel

$$(w_0, x_0, y_0, 1),$$

and suppose its principal block in the variables (W, X, Y) is positive definite. Equivalently the normalized kernel is simple and this specified complement is positive. This real rank/complement condition is stated explicitly; no unverified support census is used to remove it.

Then the original full slack is SOS in $B_c \otimes B_s$ for every independent test with

$$\bar{u} = \bar{v} = 0, \qquad [\bar{W} : \bar{X} : \bar{Y} : \bar{Z}] = [w_0 : 0 : 0 : 1], \tag{3}$$

with the following un-divided normal quantities:

$$X - vx_0, \quad Y - vy_0, \quad u - vt_0$$

are the normal quantities handled below. **No asymptotic comparison or extra smallness assumption on these three differences is needed.** They automatically have zero residue in (3), and the positive Gram is proved uniformly on all their independent rates. In particular this statement does not assume that u/v is bounded or that its residue is t_0: the construction is an identity in the un-divided variables. The selected

t_0 specifies a coefficient-side jet center, not a condition on an independent mixed or pure-tested ratio.

174.1.2 2. Full polynomial and residue Hessian

The pure-tested amplitude (vY-Z) is a unit in (3). Normalize it to (-1), so

$$Z = 1 + vY.$$

The whole opened alpha packet becomes $4\alpha(1 + u^2)$. Substituting

$$X = vx, \quad Y = vy, \quad u = vt, \quad W = w, \quad V = v^2$$

formally, on the coefficient side only, gives the exact full identity

$$F = 4\alpha + v^2 K(x, y, t, w, V). \tag{4}$$

The complete K is polynomial. Its constant V coefficient is

$$\begin{aligned} K_0 = {} & 4e_1x^2 + 4e_2(w - x - t)^2 + 4e_3(tw - y)^2 + 4e_4y^2 \\ & + 4e_5(tw - y - 1)^2 + 4\beta(x + t)^2 \\ & + 2d\{(tw - y)^2 + (x + t)^2 - 2tw\} + 4\alpha t^2. \end{aligned} \tag{5}$$

At the real residue the last term vanishes, and this is precisely $J_4(t)[w, x, y, 1]$. The stipulated positive three-state complement makes the (w, x, y) Hessian positive. Its minimized scalar is

$$\frac{16(q_{20}t^4 + q_{11}t^2 + q_{02})}{\det M_J(t)}, \tag{6}$$

where M_J is that positive principal jet block. Physicality and (2) make the numerator a positive multiple of $(t^2 - t_0^2)^2$. Therefore the full four-variable Hessian of K_0 at

$$q_0 = (x_0, y_0, t_0, w_0)$$

is positive definite. Its optimized next coefficient is

$$\gamma = \partial_V K(q_0, 0) = \frac{16(q_{21}t_0^4 + q_{12}t_0^2)}{\det M_J(t_0)} > 0. \tag{7}$$

The determinant calculation is the full state Schur-complement calculation, after scaling X, Y by v. At the zero of (6), derivatives of the positive state determinant do not affect the next coefficient. This calculation proves the real sign in (7); no state-block inverse is inserted into a tensor certificate.

174.1.3 3. Exact centers, including both signs

Solve $\nabla_q K(q_*(c,V),V)=0$ by the Nash implicit function theorem, justified by the positive four-Hessian. Let $k(c,V)=K(q_*(c,V),V)$, and put

$$\kappa = k(c,0).$$

If $\kappa \geq 0$, take $q_c = q_*(c,0), V_c = 0$. If $\kappa < 0$, solve

$$\nabla_q K(q_c,V_c)=0, \qquad K(q_c,V_c)+V_c K_V(q_c,V_c)=0. \tag{8}$$

Its residue Jacobian has the positive four-Hessian and final pivot $2\gamma > 0$, with zero lower-left block. Hence the nearby solution exists in the coefficient field and satisfies $V_c > 0$, by the local monotonicity of $k(V)+Vk'(V)$ from its negative value at zero.

In the latter branch the pure coefficient scalar

$$\mu = 4\alpha + V_c K(q_c,V_c) \tag{9}$$

is nonnegative. It is the original physical slack at the actual coefficient-side realization $v=\sqrt{V_c}$,

$$(u,X,Y,W) = (vt_c, vx_c, vy_c, w_c), \qquad Z = 1+vY.$$

Thus the sign argument is valid for arbitrary physical coefficients, not only coefficient vectors already at contact.

174.1.4 4. Verified full-tail grouping without division by v

Define the un-divided normals

$$N_1 = X - vx_c, \quad N_2 = Y - vy_c, \quad N_3 = u - vt_c, \quad N_4 = v(W - w_c). \tag{10}$$

Write $z = W - w_c$ while expanding the full polynomial. Its four exact linear coefficients in N_1, N_2, N_3, z are

$$vK_x, \quad vK_y, \quad vK_t, \quad v^2 K_w.$$

Accordingly the whole linear part is $v\nabla_q K(q_c,V)\cdot N$. Every monomial of the full remainder belongs to

$$(N_1, N_2, N_3, vz)^2. \tag{11}$$

The direct checker allocates all 55 monomials of that complete remainder into (11). Terms such as $N_3^2 z^2$ retain z^2 as a bounded infinitesimal coefficient. Terms with one z and one other normal have the required factor v, unless they already have two of the first three normals. Terms with z^2 and no other normals have v^2. Consequently no step substitutes $z = N_4/v$.

The exact resulting normal Gram has residue

$$\tfrac{1}{2}\operatorname{Hess}_{(x,y,t,w)} K_0(q_0) \succ 0. \tag{12}$$

Since $\nabla K(q_c,V_c)=0$, the linear coefficient polynomial is divisible by $V - V_c$. Its contribution couples N to the scalar generator $V - V_c$ with an extra v, hence a zero-residue cross coefficient. The fixed-center univariate profile gives, by exact monic division, the two identities

$$\begin{aligned} &\kappa \geq 0: \quad F = 4\alpha + \kappa v^2 + G^T H G, \quad G = (N_1, N_2, N_3, N_4, V), \\ &\kappa < 0: \quad F = \mu + G^T H G, \quad G = (N_1, N_2, N_3, N_4, V - V_c). \end{aligned} \tag{13}$$

The final Gram residue is the positive block (12) together with the positive last entry γ. The bounded positive-residue Gram lemma proves it SOS, and the remaining pure nonnegative point scalars are squares. Restoring the event normalization multiplies the roots by a pure tested unit. This proves the claimed original-ring certificate for every rate in (3).

174.1.5 5. Nonvacuity and audit status

At the physical residue

$$c_0 = (2, 0, 1, 1, 1, 1, 1, 0, 1)$$

one has

$$Q = 144\{(U - V)^2 + 2UV(U + V) + U^2V^2\}.$$

The corner center is $(x_0, y_0, t_0, w_0) = (-1/2, 1/2, 1, 1)$, and $\gamma = 6$. The full four-Hessian is

$$\begin{pmatrix} 32 & 0 & 24 & -8 \\ 0 & 32 & -24 & -24 \\ 24 & -24 & 48 & 12 \\ -8 & -24 & 12 & 32 \end{pmatrix} \succ 0.$$

`verify_positive_d_ratio_corner.py` reconstructs the full polynomial and verifies **9 identities / 31 scalar entries**, the 61-term original normalized parent, the 60-term profile, all 55 remainder monomials, and the full residue Hessian. Its report is `POSITIVE_D_RATIO_CORNER_CHECKS.json`.

The theorem is local in the residue state w_0. Other real kernel directions, a vanishing complement, ratio endpoints, and a vanishing off-edge coefficient have not been included by approximation.

Chapter 175

Positive total scale: the recovered finite-pitchfork fit

Component `pfb0cf454f653`
Source: `BASE/proofs/routing/a5_recovery/finite_pitchfork_body.tex`
SHA256: `fb0cf454f653dfc2e569f0e07e24f832993b1ba3ca28dac452f7dedb31eb19e7`
Inventory use: S complete strong/finite-edge fit and every projective corner; its A6 call is replaced by AU2-AU5

175.1 Positive total scale: the recovered finite-pitchfork fit

The September archive contains an explicit full-parent theorem for a finite nonzero commuting edge, even though it does not contain the later A5 terminal files under their recorded names. Its source hypotheses are automatic on the ordinary and finite-pitchfork branch with d, C, A, D, α units. This note proves that exact implication and supplies explicit full-parent projective boundary forms. Thus that subbundle of A5 is now mathematically audited. Weak-α, projective-pitchfork, and reciprocal terminal branches remain separate obligations; there is no whole-A5 admission.

Theorem 175.1. *Let the original normalized C1 or C2 source be bounded, nonnegative and physical over a real closed coefficient extension of $\mathbb{R}$. Assume d, C, A, D, α are positive bounded units. Retain its exact canonical zero coefficients, in particular $\lambda = 0$, and every other nonnegative original coefficient n, m, β. Then its entire parent is tensor-SOS at every independent projective testing pair and every real or complex testing state.*

The proof uses the admitted separated strict-Gram lemma, the freshly audited noncommuting-residue theorem A6, and the preserved finite-edge critical-center theorem whose proof is reviewed below. All source and testing rates remain independent.

175.2 Exact source fit of the recovered finite-edge theorem

The archived notation is $(d, \lambda, e_1, e_2, e_3, e_4, e_5, \alpha, \beta) = (d, \lambda, n, m, C, A, D, \alpha, \beta)$. The theorem requires $\lambda = 0$ exactly, $\bar{d} > 0$, $\bar{m} = 0$, a finite real $v_0 \neq 0$, and positivity conditions on the residue determinant at $u = 0$. Bars denote real residues in this section. Its test state is the W ray. It permits arbitrary independent nonnegative opening rates of all the original coefficients.

At $\lambda = m = 0$, the exact determinant polynomial is

$$Q(U, V) = UE(V) + U^2 B(V), \quad E(V) = q_{10} + q_{11}V + q_{12}V^2, \quad B(V) = q_{20} + q_{21}V + q_{22}V^2.$$

Use the inherited positive quantities

$$t_1 = d + 2(n + D), \qquad t_2 = d + 2(\alpha + m + A),$$
$$t_3 = d + 2(C + D), \qquad t_4 = d + 2(\alpha + \beta + m),$$

$$t_5 = d + 2(\beta + n), \qquad t_6 = d + 2(\alpha + C + A),$$
$$t_7 = d + 2(\beta + n + m), \qquad t_8 = d + 2(C + A + D).$$

All eight have positive bounded-unit residues. At the residue $m = 0$, the required coefficients are

$$q_{10} = 4\alpha A t_7 t_3 > 0, \quad q_{12} = 4CDt_5 t_2 > 0, \quad q_{21} = 4(\beta n t_3 t_2 + CAt_1 t_4) > 0,$$

and $q_{20}, q_{22} \geq 0$. Therefore $B(v_0^2) > 0$ for every $v_0 \neq 0$. Physicality implies $E(V) \geq 0$ for all real $V > 0$: divide $Q(U, V) \geq 0$ by $U > 0$ and let the real variable U tend to zero. Consequently a zero $E(v_0^2) = 0$ is a double zero, $E(V) = q_{12}(V - v_0^2)^2$, and otherwise $E(v_0^2) > 0$.

After normalizing the tested event $r_2 = vW - X + uvY - uZ$ to one, the residue state complement on (X, Y, Z) is

$$M_0(v) = \begin{pmatrix} 2(d + 2n + 2\beta)(1 + v^2) & 0 & 0 \\ 0 & 2d(1 + v^2) + 4(C + A)(1 + v^2) + 4D + 4\alpha v^2 & 4v(D - \alpha) \\ 0 & 4v(D - \alpha) & 4\alpha + 4Dv^2 \end{pmatrix}.$$

Its lower-block determinant is positive because, after a positive factor is removed, it is

$$\alpha d + 2\alpha C + 2\alpha A + 2\alpha D(1 + v^2) + Dv^2(d + 2C + 2A) > 0.$$

Thus the specified state complement is positive definite. Every hypothesis of the recovered finite-edge theorem is now verified directly from the original A5 source, including arbitrary $n, \beta \geq 0$.

What the recovered proof actually establishes. Its exact pure-testing normalization is $W = (1 + X - uvY + uZ)/v$, leaving the entire opened m packet as $4m$. The formal coefficient-side substitution $X = u^2 x, Y = ua, Z = ub, U = u^2$ gives the full identity

$$F = 4m + u^2 K(x, a, b, v, U), \qquad K_x = UE_x,$$

with $\partial_x E_x|_{U=0} = 4(d + 2n + 2\beta)(1 + v^2) > 0$. The actual independent testing coordinates are never divided by u. The minimized residue profile is a positive multiple of $E(v^2)$; the full (a, b, v) Hessian is strict at a double zero. Its next coefficient is

$$\gamma = \frac{16(1 + v_0^2)^2 B(v_0^2)}{v_0^2 \det M_0(v_0)} > 0.$$

This is a real residue determinant/Schur calculation, not a mixed inverse in the tensor ring.

The coefficient-field implicit-function equations are $(E_x, K_a, K_b, K_v) = 0$. With κ the optimized value at $U = 0$, use $U_c = 0$ if $\kappa \geq 0$; if $\kappa < 0$, also impose $K + UK_U = 0$. The added pivot is $2\gamma > 0$ and produces $U_c > 0$. The remaining scalar $\mu = 4m + U_c K_c$ is nonnegative by original physicality at an actual coefficient-field realization.

The full normal expansion uses

$$N_x = X - u^2 x_c, \quad N_y = Y - ua_c, \quad N_z = Z - ub_c, \quad N_v = u(v - v_c), \quad R = u^2 - U_c.$$

Every first state-normal coefficient is printed in the recovered proof: $u^2 E_x, uK_a, uK_b$. Monic division at the coefficient center expresses them in the five roots above, with an extra u or u^2 on every coupling to R. The complete state remainder is exactly quadratic. The residue Gram consists of the positive state-X mass, the positive (a, b, v) Hessian, and the pivot γ; all couplings to R vanish in residue. If $\kappa \geq 0$, replace R by u^2 and retain $4m + \kappa u^2$. Otherwise retain μ. Both branches restore the original event normalization. This is a full-parent certificate. When $E(v_0^2) > 0$ the four ordinary normals (X, Y, Z, u) already have strict residue, and the same whole $4m$ packet is retained.

The complete proof is the recovered `POSITIVE_D_FINITE_EDGE_TERMINAL.md`; its exact full-parent checker was freshly replayed: 15 identities, 30 scalar entries pass. Its local conclusion is used only on the finite nonzero edge stated above. The next section supplies the previously excluded endpoints.

175.3 Explicit all-projective dispatch

Noncommuting residue tests use A6. It remains that at least one angle has residue zero or infinity. For finite $u_0 \neq 0$ and $v_0 = 0$, the real residue parent is a sum of nonnegative event terms and the core $2d[(uW - Y)^2 + (X + uZ)^2]$. The unit A, α, C terms force $Y = Z = W = 0$, and the core then forces $X = 0$. Hence the residue state matrix is positive definite.

For $u_0 = 0$ and finite $v_0 \neq 0$, the unit C, A, D, α terms and $d > 0$ leave only the W ray if m is weak, while unit m removes that ray too. In the weak case, positive residue slack is immediate; at the W ray, the previous section applies for either $E(v_0^2) > 0$ or $E(v_0^2) = 0$.

For $u = \infty$ and finite nonzero v, set $p = 1/u$ and homogenize as $F_p = p^2 F(1/p, v)$. Its real residue has core

$$2d[(W + vX)^2 + (vY - Z)^2]$$

and positive $4C(1 + v^2)W^2 + 4A(1 + v^2)Y^2$ packets. These force every state coordinate to vanish at a zero, so the matrix is positive definite. For $v = \infty$ and finite nonzero u, the homogenized residue has core $2d(1 + u^2)(X^2 + Y^2)$ together with $4C(uW - Y)^2 + 4D(uX - Z)^2$. Again it is positive definite. These are exact residue identities and use no density argument.

It remains to cover the four double corners. The following displayed quadratic forms are complete, with coefficient letters interpreted in real residue. All coefficients that are merely nonnegative remain so. At a unit state coordinate, normalize it in the testing field.

At $(u, v) = (0, 0)$ the only possible zero state is the W ray. Put $W = 1$ and retain the entire m packet. The remainder lies exactly in $(u, X, Y, Z)^2$, and its residue quadratic is

$$2(d + 2n + 2\beta)X^2 + (4C + 4D + 2d)(u - Y)^2 + 4AY^2 + 4\alpha Z^2.$$

It is strict. The testing v is bounded infinitesimal in its coefficients and is never divided out.

At $(u, v) = (\infty, 0)$ the only possible zero state is the X ray when n is weak; if n is a unit there is no zero. Put $X = 1$, retain the entire n packet in F_p, and use roots (p, v, W, Y, Z). The whole remainder belongs to their square ideal and its residue is

$$4CW^2 + 4AY^2 + (4D + 2d)(W + v)^2 + 4\alpha Z^2 + [4(m + \beta) + 2d](p + Z)^2 \succ 0.$$

At $(u, v) = (0, \infty)$ the only possible zero state is the W ray when m is weak. Put $q = 1/v$, $W = 1$, and retain the entire m packet in $F_q = q^2 F(u, 1/q)$. The whole remainder is in $(u, X, Y, Z)^2$ and its residue quadratic is

$$(4n + 4\beta + 2d)X^2 + 4C(u - Y)^2 + [4(A + \alpha) + 2d]Y^2 + 4DZ^2 \succ 0.$$

The testing q remains a bounded infinitesimal coefficient.

At $(u, v) = (\infty, \infty)$ the only possible zero state is the Z ray when β is weak. Put $Z = 1$ and retain the entire β packet in $F_{pq} = p^2 q^2 F(1/p, 1/q)$. The complete remainder lies in $(p, q, W, X, Y)^2$ and its residue quadratic is

$$4CW^2 + 4AY^2 + 4D(X - p)^2 + (2d + 4n)X^2 + [2d + 4(m + \alpha)](Y - q)^2 \succ 0.$$

For each corner, allocate two normal factors from every monomial to the Gram roots; every remaining factor is bounded. Terms above quadratic order have infinitesimal coefficients. The displayed full quadratic residues are strict, so the separated strict-Gram lemma applies. Restore each retained packet. Homogeneous angle factors and state normalizations are pure testing squares. The identities include literal zero and infinity and every original nonnegative opening rate. For nonzero state residues away from the listed rays, positivity gives the scalar strict-residue exit. The zero state is immediate and complex states split into real and imaginary evaluations. This proves the theorem.

Scope, checks, and the remaining A5 imports

The new checker `check_finite_pitchfork_fit.py` verifies 11 exact identities: four full-remainder square-ideal memberships, their four complete quadratic forms, and the three finite/infinite edge identities. The recovered finite-edge checker separately verifies the exact normalized full parent, all first state normals, and complete hard tail. The general signs are proved above and in the recovered critical-center argument; the checker is corroboration rather than a positivity premise.

This proves the ordinary and finite-pitchfork subbundle with d, C, A, D, α units. In the C2 partition its source-reflection image covers the single weak-D branch. In the C1 ordinary branch a unit β may be exchanged with D by the exact stabilizer when m, C, A, α are units. These are the previously audited exact full-parent symmetry transports, with their literal zeros retained.

The remaining A5 imports include the weak-α all-tests terminals and their stabilizer images, the C1 projective pitchfork with D, β weak, and the reciprocal source branches with m, D weak. None satisfies the required unit-D premise by mere approximation. A separate note `weak_alpha_unit_n_fit.tex` verifies the finite-ratio corner's positive-endpoint and simple-complement hypotheses only on the C2 subdomain where d, n, m, C, A, D are all units and α is weak. It does not supply the rest of weak-α. The strict-α-mass theorem requires $C, A \gg d$, which is incompatible with bounded normalized coefficients and unit d.

Exact archive identity, selected member hashes, fresh replay receipts, and the scoped G1–G9 review are recorded in the adjacent recovery manifest and `RECOVERY_AND_FIT.txt`. The newer R19 frontier continues to override old global-status assertions.

Chapter 176

Positive-scale C2: a precise weak-alpha subbundle

Component `pf31e38136e03`
Source: `BASE/proofs/routing/a5_recovery/weak_alpha_unit_n_body.tex`
SHA256: `f31e38136e03138df8640da5aa173a9fd5e0fc843b71cecea10c809b1437478f`
Inventory use: CU exact unit-m/unit-n weak-alpha fit, both state charts and projective endpoints; its A6 call is replaced by AU6

176.1 Positive-scale C2: a precise weak-alpha subbundle

Theorem 176.1. *Let the original bounded normalized C2 source be physical, with $\lambda = \beta = 0$, all event coefficients nonnegative, d, n, m, C, A, D positive bounded units, and α weak, meaning $\bar{\alpha} = 0$. Then the complete original parent is tensor-SOS for every independent projective testing pair and every real or complex testing state.*

This statement assumes all six printed unit hypotheses. It is not the whole weak-α branch and gives no assertion for weak n, m, C, A, or D. The proof uses A6, the separated strict-Gram lemma, and the recovered complete theorem `POSITIVE_D_FINITE_RATIO_CORNER_TERMINAL.md`, with its exact full-tail proof freshly read and replayed. The new issue is to show that its finite-ratio and state-complement hypotheses exhaust every possible zero left by this source, without constraining the independent tested ratio u/v.

176.2 Only the zero-zero corner needs a new argument

A6 covers noncommuting testing residues. On finite $u = 0$, $v \neq 0$, the unit m removes the W ray, and the (X, Y, Z) complement is strict because d, D, C, A are units and $v \neq 0$. On finite $v = 0$, $u \neq 0$, the units n, A force $X = Y = 0$ at a zero; the C packet forces $W = 0$, and the signed core $(X + uZ)^2$ forces $Z = 0$.

The exact single-infinity core identities in `finite_pitchfork_fit.tex` require no unit α: at $u = \infty$, $v \neq 0$ use $2d[(W + vX)^2 + (vY - Z)^2]$ and the C, A packets; at $v = \infty$, $u \neq 0$ use $2d(1 + u^2)(X^2 + Y^2)$ and the C, D packets. Both state matrices are strictly positive.

At $(\infty, 0)$ the unit n removes the only possible X zero ray, and at $(0, \infty)$ the unit m removes the W ray. At (∞, ∞), put $p = 1/u, q = 1/v$, normalize the possible Z zero ray to $Z = 1$, and use (p, q, W, X, Y) as roots. The exact full-parent square-ideal allocation already displayed and checked in the finite-pitchfork note has residue

$$4CW^2 + 4AY^2 + 4D(X - p)^2 + (2d + 4n)X^2 + [2d + 4(m + \alpha)](Y - q)^2 \succ 0.$$

This form remains strict for weak α. Thus every projective test except u, v infinitesimal is already covered, including literal zero/infinite endpoints and all signs.

176.3 The exact two state charts at zero-zero

Write F for the original parent, with $\lambda = \beta = 0$, and retain the whole nonnegative α packet $4\alpha(1+u^2)(vY-Z)^2$ whenever using an ordinary strict jet. At $u = v = 0$ the residue is strictly positive on the (X, Y) state subspace. Hence X- or Y-unit states have positive residue slack. For all other states, normalize a maximal coordinate. Then either $W = 1$ with Z infinitesimal, or $Z = 1$ with W bounded.

In the first chart the entire remaining parent belongs to $(X, Y, u, v)^2$. Its quadratic residue is

$$(4n+2d)X^2 + 4m(v-X)^2 + [4(C+D)+2d](u-Y)^2 + 4AY^2 \succ 0. \tag{1}$$

Consequently the finite normal allocation and restoration of the complete α packet give tensor-SOS. No testing ratio is used.

In the second chart, set $w = \bar{W} \in \mathbb{R}$. The full residue quadratic jet on (X, Y, u, v) is

$$\begin{aligned} Q_w = {} & 4nX^2 + 4m(vw - X - u)^2 + 4C(uw - Y)^2 + 4AY^2 \\ & + 4D(uw - Y - v)^2 + 2d[(uw - Y)^2 + (X+u)^2 - 2uvw]. \end{aligned} \tag{2}$$

All coefficient letters in (1)–(2) now denote their real residues. Physicality implies $Q_w \succeq 0$: evaluate the physical residue parent at $(X, Y, u, v) = s(x, y, a, b)$ and let the real s tend to zero. The (X, Y) principal block is strictly positive, already from the nX^2 and AY^2 terms. If Q_w is strict, the same ordinary four-normal allocation applies, with $W - w$ infinitesimal in its coefficients, and the whole α packet restored.

Suppose instead that Q_w is singular. It has no nonzero kernel vector with $v = 0$: the nonnegative sum at $v = 0$ includes $4nX^2 + (4m+2d)(X+u)^2 + 4AY^2$, so $X = u = Y = 0$. It also has no nonzero kernel vector with $u = 0$: the sum at $u = 0$ includes $4nX^2 + 4AY^2 + 4D(Y+v)^2$, so $X = Y = v = 0$. Its real angular Schur complement therefore has positive diagonal entries. Being singular and positive semidefinite, it has rank one and a kernel with finite nonzero ratio $t_0 = u/v$. Normalize a real kernel vector by $v = 1$ and write it as $(x_0, y_0, t_0, 1)$. This constructs a real coefficient-side jet center; it imposes no condition on the actual independent testing value of u/v.

176.4 Every singular jet fits the recovered ratio theorem

The homogeneous state jet at a real ratio t is

$$\begin{aligned} J_t(W, x, y, Z) = {} & 4nx^2 + 4m(W - x - tZ)^2 + 4C(tW - y)^2 + 4Ay^2 \\ & + 4D(tW - y - Z)^2 + 2d[(tW - y)^2 + (x + tZ)^2 - 2tWZ]. \end{aligned}$$

Its (W, x, y) complement at $Z = 0$ is strict: it contains $4nx^2 + 4m(W-x)^2 + 4Ay^2$, with the remaining terms nonnegative. At $t = t_0$, its kernel is $(w, x_0, y_0, 1)$ by (2).

In the original determinant notation, at $\lambda = \beta = \alpha = 0$,

$$q_{20} = 4nAt_3t_4 > 0, \qquad q_{02} = 4mDt_5t_6 > 0, \qquad q_{21} = 4CAt_1t_4 > 0, \qquad q_{12} = 4CDt_5t_2 > 0.$$

Here the t_i are the original positive sums of bounded coefficients printed in the finite-pitchfork fit; all are units since d is a unit. The full jet determinant is

$$\det J_t = 16(q_{20}t^4 + q_{11}t^2 + q_{02}).$$

Thus the singular jet gives precisely the finite nonzero root required by the recovered theorem, and its next coefficient satisfies $q_{21}t_0^4 + q_{12}t_0^2 > 0$. The state complement, both endpoint coefficients, and the positive next coefficient are all verified.

For clarity, the recovered proof normalizes the pure-testing event $vY - Z$ to -1, setting $Z = 1 + vY$. It keeps the complete opened α packet, including its $4\alpha u^2$ term, in the exact profile

$$F = 4\alpha + v^2K(x, y, t, w, V), \qquad (X, Y, u, W, V) = (vx, vy, vt, w, v^2)$$

used only to construct coefficient-field critical centers. The positive state complement and the double finite root give a strict full Hessian in (x, y, t, w), with positive next coefficient

$$\frac{16(q_{21}t_0^4 + q_{12}t_0^2)}{\det M_J(t_0)}.$$

The two signs of the optimized constant are handled by coefficient-only implicit centers exactly as in the finite-edge proof; physicality makes the remaining scalar nonnegative.

Its actual tensor roots are the un-divided expressions

$$X - vx_c, \quad Y - vy_c, \quad u - vt_c, \quad v(W - w_c), \quad v^2 - V_c.$$

All four full first-normal coefficients are $v\nabla K$ on this frame. Monic difference division makes their coupling to $v^2 - V_c$ infinitesimal. All 55 full-tail monomials have an exact allocation to the first four roots; factors of $W - w_c$ that remain in coefficients are infinitesimal. There is no substitution dividing $v(W - w_c)$ by v. The full residue Gram is the positive four-variable Hessian and the positive last scalar pivot. Hence the theorem applies to every testing point with the stated residue state, regardless of whether its actual tested u/v is bounded or approaches t_0. This includes literal $v = 0$ and arbitrary independent opening rates.

The two zero-corner state charts and the strict/singular jet split are exhaustive. Together with the earlier projective dispatch, this proves the theorem. The zero state is direct, and complex states split into real and imaginary quadratic evaluations.

Verification and retained complement

The recovered ratio checker freshly passes 9 exact identities and 31 scalar entries, including the entire normal tail and its residue Hessian. The new `check_weak_alpha_unit_n_fit.py` checks the complete square-ideal memberships and exact quadratic forms in both zero-corner state charts (4 assertions). The other corner identities are the generic, already-replayed 11 assertions from the finite-pitchfork fit and remain valid for weak α.

Only the explicit subdomain with d, n, m, C, A, D units has been admitted. The C2 reflection exchanges (m, D) and (C, α); its exact image therefore supplies the corresponding weak-C subdomain with the transported unit hypotheses. It does not remove weak-n endpoints, other weak-α source shapes, C1 weak-α, projective pitchforks, reciprocal terminals, or any local C14/weak-C2 frontier. All three remaining A5 groups remain open outside the stated repairs.

Chapter 177

Scope and actual checklist delta

Component p0be1d6088cc6
Source: BASE/proofs/routing/routing_audit.tex
SHA256: 0be1d6088cc6370c7cb2ec4b2b0ed2a294c691a0cf96af071b5f14d9a8e9250e
Inventory use: C1 input and finite dispatch; C1.4 subdivision | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

Scope and actual checklist delta

This note closes the two technical audit imports left in U1.A6 by the R18 routing review: constrained curvature with one or two hard events, and the four-matching terminal with arbitrary internal coefficient scales. It also writes an explicit common input, the source-entry maps, and an exhaustive decision tree into the historical thirteen rows and their C14/R1–R10 subdivisions. All original physicality, contact, coefficient signs, strict gaps, literal zeros, and total-scale conditions are carried along as conjuncts.

The distinction between a proved dispatch implication and an accepted terminal certificate is essential. The complete positive-total-scale terminal bundle in A5 is only partially repaired: the recovered finite-edge proof and the new `a5_recovery/finite_pitchfork_fit.tex` cover the ordinary/finite-pitchfork subbundle with d, C, A, D, α units and its exact symmetry images. The companion `a5_recovery/weak_alpha_unit_n_fit.tex` supplies a second precise C2 subdomain: d, n, m, C, A, D all units, α weak, and $\lambda = \beta = 0$. The rest of weak-α, projective-pitchfork, and reciprocal terminals remain imported. This audit also found a specific mismatch in the preserved uniform C1 theorem statements: the printed infinitesimal-e and unbounded-e theorems do not by themselves cover unit e. The companion proof `us_unit/bounded_delta_all_tests.tex` repairs that subentry directly. It extends the exact full-parent contact–conductor proof to bounded $\Delta = 1/e$ and bounded testing w, proves positive coefficient weights without comparing independent rates, and supplies the complete testing join. Thus an old all-e assertion is no longer needed for that subentry.

The exact parent and Laurent determinant identities and the separated bounded strict-Gram lemma remain the already-admitted inputs. The focused routing checker has 225 passing exact assertions, covering both full-parent reflections, support classification, the two-hard activation table, finite source-type routing, and four-matching normal completion. The separate bounded-Δ checker has 33 passing exact conductor and full-source lift assertions. The scoped positive-scale repair has a recovered 15-identity, 30-entry full-parent replay and 11 new exact projective-boundary checks. Its C2 weak-α subdomain has a separate recovered 9-identity, 31-entry full-tail replay and 4 exact state-jet assertions. These assertions do not prove an unresolved commuting-boundary terminal. No global tensor-SOS or spectrahedral-shadow conclusion follows from this note.

All cited paths are given in the accompanying `ROUTING_LEDGER.json`, relative to the recovered `checkpoint_R19` directory, with SHA-256 hashes; new continuation files have their own explicit path root. The symbols `R13`, `OLD`, and `INH` in the ledger have one explicit expansion each; no absent external manuscript is substituted for a preserved file.

177.1 One exact input, before optional exclusions

Let $K_c, K_t \supset \mathbb{R}$ be independent real closed fields. The bounded subring of either field consists of elements whose absolute value is at most some real number; its maximal ideal consists of infinitesimals. For $q > 0$ write $q : \mathsf{I}$ when q is infinitesimal, $q : \mathsf{U}$ when q and q^{-1} are bounded, and $q : \mathsf{O}$ when q^{-1} is infinitesimal. These are exhaustive, disjoint types. Bounded means I or U; noninfinitesimal means U or O. Zero is always a separate literal case. Type notation is external shorthand for ordered-field inequalities relative to real bounds, not a new algebraic equation.

In the coefficient order $(d, \lambda, n, m, C, A, D, \alpha, \beta)$ and with $D_u = 1 + u^2$, $D_v = 1 + v^2$, retain the complete original parent

$$\begin{aligned} F_c = {} & 4\lambda D_u (W + vX)^2 + 4nD_u D_v X^2 + 4m(vW - X + uvY - uZ)^2 \\ & + 4CD_v (uW - Y)^2 + 4AD_u D_v Y^2 + (4D + 2d)(uW + uvX - Y - vZ)^2 \\ & + (4\alpha + 2d) D_u (vY - Z)^2 + (4\beta + 2d) D_v (X + uZ)^2 - 2dD_u D_v Z^2. \end{aligned} \tag{P}$$

Its symmetric state matrix is $P_c(u, v)$, so $F_c = \psi^T P_c \psi$. Physicality is the exact quantified polynomial condition $P_c(u, v) \succeq 0$ for every source-field projective pair, interpreted in the four homogeneous angle charts. It is never replaced by positivity of a truncated scalar or a modified parent.

The canonical remaining source input $Q(c)$ is the following conjunction:

1. $c \in K_c^9$, $d > 0$, all eight event coefficients nonnegative, and physicality of (P).
2. $\delta_{\mathrm{tot}} = d/(d + \lambda + n + m + C + A + D + \alpha + \beta) : \mathsf{I}$. This quotient is computed before any later $d = 1$ or $C = 2$ normalization.
3. The source is coercive and has five or six positive events. There is a finite positive contact $u_c, v_c > 0$ and nonzero ψ_c such that $P_c \psi_c = 0$ and $\partial_u F_c = \partial_v F_c = 0$ at this contact.
4. A deterministic omission convention has chosen C1 or C2 as described below. Its literal coefficient equalities and positivity conditions are part of Q, not conditions to be inferred later from a limiting approximation.

The testing input is an arbitrary independent K_t projective angle pair and an arbitrary real or complex state. A complex state is split into real and imaginary parts because the parent is real symmetric. No source contact is identified with a testing point.

If one wants precisely the input after A6, append that at least one testing measurement pair commutes in real residue. We prove the source dispatch with all tests allowed, so that additional restriction cannot create a missing source case. Optional historical exits may likewise be intersected afterwards: if $Q \Rightarrow E \vee \bigvee R_i$, then $Q \wedge \neg E \wedge B \Rightarrow \bigvee (R_i \wedge B)$ for every common additional predicate B. This elementary implication avoids using an unspecified phrase such as "the complement of all prior admissions" as the premise of the coverage proof itself.

Canonical omission and symmetry convention. Use the fixed cycle $(\lambda, D, C, A, \alpha, m, \beta, n)$. The A3 reduction gives compatible omissions: pairs have cycle distance one or two, and triples consist of three consecutive vertices. For a triple, choose a distance-one pair first, with the lexicographically least pair in this fixed cycle breaking ties. Choose the first word in a fixed enumeration of the finite Bell relabeling group that sends it to $\{\lambda, n\}$. The extra zero is then $D = 0$ or $\beta = 0$. For two omissions, distance one enters C1 and distance two enters C2. Thus the two canonical domains are

$$\begin{aligned} &\mathrm{C1}: \quad \lambda = n = 0, \quad m, C, A, \alpha > 0, \quad D, \beta \geq 0, \quad D + \beta > 0; \\ &\mathrm{C2}: \quad \lambda = \beta = 0, \quad m, C, A, D, \alpha > 0, \quad n \geq 0. \end{aligned}$$

The five-active C2 boundary $n = 0$ is already assigned C1 by the precedence convention, but it is deliberately allowed in C2 theorem statements and all later implications. No proof is lost by this overlap. The partition is disjoint for chosen canonical records, not for their unoriented physical-source images.

177.2 A6: the formerly imported technical audit

177.2.1 Weighted centers do not require comparable hard weights

At a noncommuting testing residue, use an entangled residue state as an invertible two-by-two matrix. A fixed local ray has a unique orthogonal partner on the other side. Consequently its vanishing losing events form a matching of the eight-cycle, of size at most four. At most three such product-event rows are independent state coordinates. Product residue states instead have strictly positive signed-core value at a noncommuting pair; their contribution is a pure-source positive factor times a pure-testing square, plus event squares.

Suppose independent hard event amplitudes are exact coordinates x_i, their positive weights are a_i, and the next coefficient scale is σ, with $\varepsilon_i = \sigma/a_i : \mathsf{I}$. The local polynomial is

$$F = \sum_i a_i b_i(y) x_i^2 + \sigma G(x, y), \qquad b_i(y_0) > 0.$$

Set $x_i = \varepsilon_i s_i$ in the coefficient-field stationary equations. After division by σ their normal equations have residue $2b_i(y)s_i + G_{x_i}(0, y) = 0$, and their tangent equations have residue $\nabla_y G(0, y) = 0$. If the constrained soft Hessian is positive definite, the residue Jacobian is invertible: its hard diagonal entries are $2b_i > 0$ and its soft Schur block is precisely that Hessian. The semialgebraic implicit-function statement over $\mathbb{R}$, followed by transfer, supplies an exact source-field stationary center. Physicality makes its value nonnegative.

The exact finite Taylor roots are $(x_i - x_{ic})/\sqrt{\varepsilon_i}$ and $y - y_c$. For each hard packet, $x_{ic} = \varepsilon_i s_{ic}$ makes its center-square and center-linear contributions bounded after this scaling. All remaining higher-degree factors are actual deviations, not quotients by a source opening. The residue Gram is strict, so the admitted separated strict-Gram lemma applies. None of these operations uses a mixed inverse or orders a source infinitesimal against a testing infinitesimal.

177.2.2 Exact deficient-support classification

For $d > 0$ write the eight outer Laurent coefficients of the determinant as $q_e \geq 0$, $e \in \{-1, 0, 1\}^2 \setminus \{0\}$. At a stationary zero its logarithmic Hessian is $\sum_e w_e e e^T$ with $w_e > 0$ for active exponents. It fails to be positive definite only if the support has rank zero or rank one. A stationary nonempty rank-one support must contain opposite exponents; one-sided support cannot have zero gradient.

For one hard event, take λ by symmetry and extract $\partial q_e/\partial\lambda$. For two hard matching events the three orbits have representatives $\{\lambda, m\}$, $\{\lambda, C\}$, and $\{\lambda, \alpha\}$. Extract the coefficient of their product. A state basis adapted to the independent hard amplitudes shows that this leading coefficient is a positive angular factor times the constrained state determinant. The positivity table below is therefore exact. Juxtaposition means both coefficients are positive.

e	$\{\lambda, m\}$	$\{\lambda, C\}$	$\{\lambda, \alpha\}$
$(-1, -1)$	α	α	always
$(-1, 0)$	D or α	D or αm	D or m
$(-1, 1)$	D	Dm	Dm
$(0, -1)$	n or $A\alpha$	n	A or n
$(0, 1)$	β or DC	D or βm	DC or βm
$(1, -1)$	An	0	An
$(1, 0)$	AC or βn	A	AC or βn
$(1, 1)$	βC	β	βC

Inspection of the four opposite-pair lines gives no rank-one stationary support for $\{\lambda, m\}$. For the other two pairs the only possibility is the perfect matching $\{\lambda, C, \alpha, \beta\}$. The one-hard calculation has exactly that same possibility. With no hard event there are precisely the two perfect matchings. These are positivity-support implications and allow every omitted weight to be literally zero.

All rank-zero supports are contained in a nonextendable matching triple, whose cyclic gaps are $(2, 3, 3)$. For one hard λ, representatives are $\{\lambda, A, \beta\}$, $\{\lambda, m, C\}$, and $\{\lambda, m, A\}$. For the pair $\{\lambda, m\}$

only the latter two occur; for $\{\lambda, C\}$ only $\{\lambda, m, C\}$ occurs; for $\{\lambda, \alpha\}$ rank zero is impossible. At $u = v = 1$, the states $(1, -1, 0, 2)$ and $(1, -1, 1, 3)$ annihilate respectively the $\{\lambda, m, A\}$ and $\{\lambda, m, C\}$ events and give signed-core values -4 and -8. Cycle symmetry supplies the remaining triples. Hence a physical constrained source with $d > 0$ cannot have rank zero.

On a perfect-matching curve all matching event squares and the signed core vanish. For the representative $\{n, m, A, D\}$ the four independent event-normal differentials satisfy

$$dH = 2t\{(1 + t^2)dX - (1 + t^2)dY - dR_m + dR_D\}, \qquad t > 0.$$

Every coefficient is nonzero. Imposing at most two hard constraints leaves a direction on which this derivative is nonzero; all event-square derivatives vanish there. The constrained parent changes sign, contrary to physicality. This removes the rank-one cases.

Thus every zero of the one/two-hard constrained determinant has positive angular Hessian. The signed core is strictly positive on the $Z = 0$ state subspace, so the constrained state matrix has respectively two or one positive transverse state eigenvalues and a simple kernel with $Z \neq 0$. A Schur-complement calculation over the real residue field identifies the angular Hessian with a positive multiple of the determinant Hessian. The full soft Hessian is therefore strict. This is a real residue calculation, not a Schur inverse in the tensor product.

For reproducibility the checker enumerates respectively 256, 128, and 64 supports for no hard event, one fixed hard event, and each two-hard orbit. It checks every positivity-table entry on every support. The enumeration corroborates the finite proof; it makes no numerical positivity inference and does not test only generic positive coefficients.

177.2.3 Four matching weights, all internal scales retained

Suppose $\mu = \min(n, m, A, D) > 0$, and d/μ and each of $\lambda/\mu, C/\mu, \alpha/\mu, \beta/\mu$ are infinitesimal. Rotate the cycle so $D = \mu$, and set

$$\eta_1 = \mu/n, \quad \eta_2 = \mu/A, \quad \eta_3 = \mu/m, \qquad 0 < \eta_i \leq 1.$$

No positive lower bound is imposed on any η_i. In the $W = 1$ chart take $R = v - X + uvY - uZ$ as a state coordinate, so $Z = (v - X + uvY - R)/u$. The fourth amplitude is $f + L$, where

$$f = (u^2 - v^2)/u, \quad L = v(u + u^{-1})X - (1 + v^2)Y + (v/u)R.$$

At $X = Y = R = 0$ the signed and fourth packets equal

$$4\mu f^2 + 2duf = 4\mu \left(f + \frac{du}{4\mu}\right)^2 - \frac{d^2u^2}{4\mu}.$$

Testing $u = 1$, $v = \sqrt{1 + d/(4\mu)}$ proves, by physicality, that $\tau = \max(\lambda, C, \alpha, \beta) > 0$ and $\rho = d^2/(\mu\tau)$ is bounded. Put $\varepsilon = d/\mu : \mathsf{I}$.

If $\rho : \mathsf{I}$, use the four normal roots $\sqrt{n/\tau}X$, $\sqrt{A/\tau}Y$, $\sqrt{m/\tau}R$, $\sqrt{\mu/\tau}(v - u)$, and the constant root. Their normal residue is

$$4B^2z_1^2 + 4B^2z_2^2 + 4z_3^2 + 4(-2z_4 + B\sqrt{\bar{\eta}_1}z_1 - B\sqrt{\bar{\eta}_2}z_2 + \sqrt{\bar{\eta}_3}z_3)^2 \succ 0, \quad B = 1 + u_0^2.$$

The complementary packets give a strictly positive constant residue; the signed linear corrections have coefficients $\sqrt{\rho}$ or $\sqrt{\rho\eta_i}$ and vanish in residue.

If $\rho : \mathsf{U}$, substitute in source-field stationarity

$$X = \varepsilon\eta_1 s_1, \quad Y = \varepsilon\eta_2 s_2, \quad R = \varepsilon\eta_3 s_3, \quad v - u = \varepsilon r, \quad u = t.$$

With $B = 1 + t^2$, $\ell = (B, -B, 1)$, $b = (B^2, B^2, 1)$, and $k = -2r + \sum \bar{\eta}_i \ell_i s_i$, the four divided normal equations are

$$8b_i s_i + 8k\ell_i + (4tB, -4tB, 0)_i = 0, \qquad -16k - 4t = 0.$$

Their Jacobian determinant is $32 \cdot 8^3 B^4 > 0$, even if some $\bar{\eta}_i = 0$. Their unique solution is

$$s_1 = -\frac{t}{4B}, \quad s_2 = \frac{t}{4B}, \quad s_3 = \frac{t}{4}, \quad r = \frac{t}{8}(1 - \bar{\eta}_1 - \bar{\eta}_2 + \bar{\eta}_3).$$

Let $P = \operatorname{res}((\lambda + \alpha)/\tau)$ and $Q = \operatorname{res}((C + \beta)/\tau)$; $P + Q > 0$. Exact normal completion of the normalized residue objective leaves

$$g(t) = 4(1 + t^2)(P + Qt^2) - \frac{\bar{\rho} t^2}{4}(1 + \bar{\eta}_1 + \bar{\eta}_2 + \bar{\eta}_3).$$

Physicality gives $g \geq 0$. At a zero, stationarity of $g/(4t^2)$ gives $P/t^2 = Qt^2$, forcing $P, Q > 0$ and positive logarithmic curvature. If $g(t_0) > 0$, normal completion gives a strict constant-plus-normal Gram. Otherwise adjoining the tangent equation gives an invertible Jacobian and an exact stationary center with nonnegative value.

Put $\delta = v - u$. The full-parent Taylor roots are

$$\sqrt{n/\tau}(X - X_c), \quad \sqrt{A/\tau}(Y - Y_c), \quad \sqrt{m/\tau}(R - R_c), \quad \sqrt{\mu/\tau}(\delta - \delta_c), \quad t - t_c.$$

The packetwise bounds that justify the entire Taylor polynomial are

$$\frac{a_i x_{ci}^2}{\tau} = \rho \eta_i s_{ci}^2, \qquad \frac{a_i x_{ci}}{\sqrt{a_i \tau}} = \sqrt{\rho \eta_i} s_{ci}.$$

The fourth normal has the same estimates with $\eta_i = 1$. Complementary and signed packets have no larger factors. Every packet is at most quadratic in the state coordinates, so after two normal factors are assigned the remaining higher factors are bounded deviations. The only denominators are pure testing units t, together with admitted pure coefficient units. The residue normal block is the strict form above; its tangent Schur complement is $g''(t_0)/2 > 0$. This proves the full packet restoration for arbitrary unequal matching weights.

At each earlier scale either a positive soft value or a strict constrained Hessian finishes the argument, or another matching edge is added. Three-edge kernels have the two inherited explicit Laurent forms; their rank failures give respectively no stationary zero or a negative minimum $2\sqrt{2D(d + 2D)} - d - 4D < 0$ (squaring has difference $d^2 > 0$). There can be at most four matching edges. The terminal above therefore completes the finite matching-scale induction without a rate comparison. This closes the specific A6 technical imports. It says nothing about a commuting testing residue by continuity.

177.3 Global sign entry and exact coordinate maps

177.3.1 C1

Normalize $C = 2$ by a positive pure-source factor. The finite contact has the unique gauge

$$(u_c, v_c; \psi_c) = (k, \ell; S, k(\theta - 1), kSr, 1), \qquad S > 0, \quad 0 < r, \theta < 1.$$

The inherited global sign proof has now had its curvature input audited above. Over $\mathbb{R}$ it uses the unique maximal-d contact graph over the connected positive event orthant, a positive cofactor proving $\theta > 0$, and no-wall removal arguments for $Y = 0$, $S - Y = 0$, and $\theta = 1$. Its exact positive seed is

$$(u, v; S, Y, \theta) = (1, 1; 1, 1/4, 1/2), \qquad c = (27/2, 0, 0, 4, 2, 9, 5/4, 13/4, 9/4).$$

The cross-multiplied amplitude identity extends from the nonzero Cramer minor locus by analyticity on this specified graph. A boundary implicit function argument opens $D = 0$ or $\beta = 0$, proves the strict signs there, and only then transfers the finite first-order implication. This use of real continuity proves source sign entry; it is not an argument that tensor SOS is closed under non-Archimedean limits.

The full-parent Alice reflection is

$$u' = 1/u, \quad v' = v, \quad G\psi = (vW - X, -W - vX, Z - vY, Y + vZ),$$

with $(\lambda\ n)(m\ C)(A\ \alpha)(D\ \beta)$ on coefficients. It obeys $G^T G = (1+v^2)I$ and

$$u^2 F_{c'}(1/u, v; G\psi) = (1+v^2) F_c(u, v; \psi).$$

Its factors are pure testing squares after homogeneous normalization. Apply it on the source side if needed to fix $0 < k \le 1$; equality is kept in the unit case. Literal $D = 0$ and $\beta = 0$ exchange.

Define, exactly,

$$\eta = \ell/(kS), \quad \zeta = S\ell/k, \quad e = k/S, \quad x = k^2, \quad a = (1-r)/\eta, \quad t = \theta/\zeta, \quad h = 1 - \theta.$$

In particular $e = k\sqrt{\eta/\zeta}$, with the positive square root. The original gaps are

$$R = 1 + xr, \quad T = 1 + xh, \quad M = R - t > 0, \quad J = T - a > 0,$$
$$E = 1 - x\eta h > 0, \quad Q = 1 - x\zeta r > 0, \quad 0 < a, t < 2.$$

Every branch below retains these exact relations. The same selected minor remains nonzero when $D = 0$ or $\beta = 0$, so a boundary coefficient column must not simply be deleted from the rational-source derivation. The nonnegative source numerators B_5, B_β in the ledger are equivalent to the literal original inequalities $D, \beta \ge 0$.

177.3.2 C2

Normalize $d = 1$. At the source contact $Z = 1$ the exact signs are

$$W > 0,\ X < 0,\ Y > 0,\ uW - Y > 0,\ 1 - vY > 0, \quad R_m > 0,\ R_D < 0,\ X + u > 0,\ W + vX > 0.$$

The proof uses the analogous unique real maximal-contact graph, the exact seed

$$(n, m, C, A, D, \alpha) = (1/8, 3/8, 1/6, 5/8, 3/8, 1/6), \quad (u, v; \psi) = (1, 1; 1, -1/4, 1/4, 1),$$

and event-amplitude removal. The wall $X = 0$ would delete n and enter the C1 triple face, where the already-proved strict sign gives a contradiction. The omitted-event signs follow from the original X kernel equation and the exact C2 reflection. Literal $n = 0$ is the C1 triple face directly. First-order transfer is performed after this real argument.

The source variables and all strict gaps are

$$c = u_c, \quad a = v_c/u_c, \quad b = aW_c, \quad r = -X_c/u_c, \quad t = (u_c W_c - Y_c)/v_c,$$
$$h = aY_c/u_c = b - a^2 t > 0, \quad g = R_m/u_c = (1+c^2)b - (1 - r + a^2 c^2 t) > 0,$$
$$j = -R_D/v_c = 1 + c^2 r - t > 0, \quad 0 < r < 1, \quad 1 - c^2 h > 0, \quad b - a^2 c^2 r > 0.$$

The positive stationary determinant is a nonzero factor times $ac^6 rgt\, hj(1-c^2h)(1+c^2)^2(1+a^2c^2)^2\Delta$, where

$$\Delta = 1 + (1+z)b + z(1+z)b^2 + 2z(1+z)br + z^2r^2 + 2zr + a^2 z\{(1+z)b + 1 + zr(2-r)\} \ge 1, \quad z = c^2.$$

Thus the six rational coefficients are the original coefficients by uniqueness of the six stationary equations, including $n = 0$.

The exact reflection is

$$(u, v; W, X, Y, Z) \mapsto (1/v, 1/u; Z, X, Y, W), \qquad (m\ D)(C\ \alpha),$$

and obeys, on the C2 omission face, the exact identity

$$u^2 v^2 F_{c'}(1/v, 1/u; Z, X, Y, W) = F_c(u, v; W, X, Y, Z).$$

Both n and A are unchanged. The positive source normalization by W_c is legal. The factorization and endpoint interpretation are homogeneous, so literal testing infinity is included.

177.4 Exhaustive dispatch into the historical thirteen rows

177.4.1 C1: five retained rows and explicit other exits

All entries in the following table include the exact C1 input above. The notation $\neg\mathsf{I}$ means a positive unit or an unbounded element. The row order is deterministic.

Source condition	Destination
$k : \mathsf{I}$, $\zeta : \neg\mathsf{I}$	C1.4, arbitrary positive η
$k : \mathsf{U}$, $\eta, \zeta : \mathsf{I}$	AS: arbitrary-state-ratio theorem
$k : \mathsf{U}$, at least one of $\eta, \zeta : \neg\mathsf{I}$	C1.5
$k, \zeta, \eta : \mathsf{I}$	US: whole uniform-source exit; exact join below
$k, \zeta : \mathsf{I}$, $\eta : \mathsf{O}$	C1.1
$k, \zeta : \mathsf{I}$, $\eta : \mathsf{U}$, $e : \mathsf{I}$	C1.3, every $0 < r < 1$
$k, \zeta : \mathsf{I}$, $\eta, e, r : \mathsf{U}$	BE: bounded-e all-tests theorem
$k, \zeta, r : \mathsf{I}$, $\eta, e : \mathsf{U}$	CA: centered-A, unit-e theorem
$k, \zeta : \mathsf{I}$, $\eta, r : \mathsf{U}$, $e : \mathsf{O}$	UE: unit-η, unbounded-e theorem
$k, \zeta, r : \mathsf{I}$, $\eta : \mathsf{U}$, $e : \mathsf{O}$	C1.2

To prove exhaustion, first split bounded k into I/U. On the $k : \mathsf{I}$ side split ζ into infinitesimal or not. On the remaining $k, \zeta : \mathsf{I}$ side split η into its three types. Only $\eta : \mathsf{U}$ requires the e trichotomy, and only $e : \mathsf{U}$ or O requires the two possibilities for r. There is no unbounded-r case because $0 < r < 1$. The inequalities $a, t < 2$ imply $r \sim 1$ when $\eta : \mathsf{I}$, but no extra rate condition is needed.

For AS, the original source formulas make D and β strictly positive; a literal zero cannot be smuggled into its theorem. BE and CA explicitly retain $\beta = 0$; on their source shapes the formula for D has positive-unit numerator and prevents $D = 0$. UE makes m/d and β/d unbounded, so its stated assumptions also decide that boundary. These are source consequences, not informal continuity extensions.

The historical US entry has three exhaustive subentries. For $e : \mathsf{I}$ its exact theorem is `c1_uniform_all_tests.tex`. For $e : \mathsf{O}$ the original unbounded-e W-unit/outer-test theorem and the restricted contact–conductor completion provide the full normal restoration. The new `us_unit/bounded_delta_all_tests.tex` proves the complete source for bounded $\Delta = 1/e$, hence covers both $e : \mathsf{U}$ and $e : \mathsf{O}$. Its new grading assigns weight zero to both δ and testing w, while preserving strictly positive weights on every scalar and first-normal correction coefficient. The Z-unit chart now allows bounded W; the complementary W-unit, Z-infinitesimal chart uses the retained-current identity with ΔZ infinitesimal. The original outer-angle identities then give all projective tests. This is the exact matching proof for US-U that the old summary did not supply. The positive strict gaps and all original source signs remain premises, and every subtracted packet is restored.

177.4.2 C2: directed weak rows and strong orbits

Weak means infinitesimal after division by d, while strong means bounded positive reciprocal; strong coefficients may be unbounded. Since $C, \alpha > 0$, first split into both weak, exactly one weak, and both strong. Both weak goes to the chart-free C2 radial theorem. Its physical matching witnesses force all four matching weights large, then the exact contact estimates give the radial-source entry. The source argument and the final radial all-tests certificate are distinct steps and are separately named in the ledger.

For exactly one weak coefficient reflect once, if necessary, so C is weak and α strong. Do not quotient this directed list by another reflection. Source identities exclude actual types II, IU, UI, UU in that orientation. The all-tests single-angle corollary removes UH. The remaining four historical directed rows are

$$\text{C2.W.IH}, \quad \text{C2.W.HI}, \quad \text{C2.W.HU}, \quad \text{C2.W.HH}.$$

Here H is an unbounded positive actual half-angle, not literal infinity; the actual contact is always finite over K_c.

For both C, α strong, if m, A, D are all strong then the five consecutive events (D, C, A, α, m) give the five-event exit. Otherwise retain m weak, else A weak, else D weak, in that order if a disjoint event subpartition is desired. The all-tests IU/UH and UU corollary removes three ordered angle types. Reflection acts on the other six by reversal and reciprocal type, giving precisely

$$\text{C2.S.II/HH}, \quad \text{C2.S.IH}, \quad \text{C2.S.UI/HU}, \quad \text{C2.S.HI}.$$

These are four orbits; IH and HI are each fixed and must not be identified. Together with the five C1 rows this gives thirteen labels. The finite angle calculation is included in the exact checker.

The directed HI and HU rows were subsequently proved empty, without testing restrictions. The proof is substantive: with $q = u_c^{-2} : \mathsf{I}$ and bounded $y = v_c^2$, original $n \geq 0$ forces $R = r/q$ bounded. Otherwise the positive leading term in its exact numerator dominates the negative terms, contradicting $n \geq 0$. Then $H = v_c Y_c \sim 1$, Δ is a positive bounded unit, and $C = (1 + R)(N_{\mathrm{src}}/q)/(2L\Delta)$ has bounded positive reciprocal. This contradicts the chosen weak-C orientation. Literal $n = 0$ is included in the contradiction. The ledger records the exact file.

The optional reciprocal-product exit requires all four products Cn, Cm, CA, CD to have bounded positive reciprocals in the weak-C orientation. Merely two large adjacent weights do not suffice, and $n = 0$ never passes this exit. This exit is unnecessary for the finite thirteen-row coverage implication; intersecting its complement later preserves coverage.

177.5 C1.4: exhaustive C14 subdivision, with missing exits restored

The input is $k : \mathsf{I}$, $\zeta : \neg\mathsf{I}$, with all exact C1 constraints. Put $L = x\zeta$ and, when $\eta : \mathsf{O}$, $\sigma = 1/\eta$. The following tree covers this whole source shape, before any extra testing screens are applied.

$L : \mathsf{O}$. The strict original gap $Q = 1 - Lr > 0$ forces $r : \mathsf{I}$. Since $1 - r = a\eta$ and $a < 2$, η cannot be infinitesimal. For $\eta : \mathsf{U}$ the large-L, unit-η all-tests theorem applies. For $\eta : \mathsf{O}$, put $H = h/\sigma$. If $H : \mathsf{O}$, enter C14.1b. If H is bounded, split $M_\beta = \beta\theta/\sigma^2$ into bounded or unbounded, after $d = 1$. Bounded M_β, including $\beta = 0$, is the JR/JB all-tests exit; unbounded M_β is C14.1a. No $H = 0$ case is introduced, since $h > 0$; $M_\beta = 0$ is literal and allowed.

$L : \mathsf{U}$. The three types of η give, respectively, the small-η L-unit all-tests exit, the unit-η L-unit all-tests exit, and C14.2. The source formulas in the first two cited theorems, rather than their titles, retain the original gap and zero-boundary hypotheses.

$L : \mathsf{I}$. Unbounded η gives C14.4. For bounded η, unbounded ζ gives C14.3a. The only remaining case is $\zeta : \mathsf{U}$: infinitesimal η gives C14.3b. If both $\eta, \zeta : \mathsf{U}$, define the exact positive bounded source gap

$$j = \eta(1 + xh) - 1 + r = \eta J.$$

If $j : \mathsf{U}$, then $j_0 = \eta + r - 1$ is a positive unit and the RO theorem applies. If $j : \mathsf{I}$, then $\eta + r \sim 1$, hence $1 - r : \mathsf{U}$. For $r : \mathsf{U}$ use JO; for $r : \mathsf{I}$ use HO. There is no need to compare j with x, and no equality $j = 0$ is permitted.

This proves that the six historical refined shapes C14.1a, C14.1b, C14.2, C14.3a, C14.3b, C14.4, together with the exact listed exits, cover C1.4. R19 admits all of C14.2 and C14.3a/b. Their files are recorded as existing closures; this note does not claim to have proved C14.1a/b or C14.4. A later continuation may replace those open labels only with its actual proof bytes and scope.

177.6 Weak HH: exact reflected entry and the ten disjoint rows

Reflect the directed weak-C HH source once to actual II. Now C is strong and α weak. Put

$$z = u_c^2, \quad y = v_c^2, \quad H = v_c Y_c, \quad L = v_c(u_c W_c - Y_c), \quad B = H + L, \quad r = -X_c/u_c.$$

The exact original constraints include

$$z, y : \mathsf{I}, \quad 0 < H, r < 1, \quad L > 0, \quad j = 1 + zr - L/y > 0,$$
$$G = B + z(H + r - 1) > 0, \quad B - zyr > 0, \quad n \geq 0.$$

Since $L < y(1 + zr)$, L is infinitesimal and B is bounded. The unchanged alpha formula is

$$\alpha = \frac{(B + zr)M}{2(1 - H)\delta}, \qquad M = (1 + z)(B + zr)H + yG + yzj + zrL(1 + y) > 0,$$

with the positive original denominator δ printed in the ledger. If B is infinitesimal, the exact estimates give α infinitesimal. If B is a unit, then H, δ, and M are units, so α is strong; also $CL = (1 + zr)M/(2\delta)$ is a unit and C is unbounded. Thus the actual directed weak HH input has exactly $B, z, y : \mathsf{I}$, not an assumed power-law scaling.

Define the weak-II variables

$$a = z/B, \quad t = y/B, \quad h = H/B, \quad l = L/B = 1 - h, \quad e = 1 - r, \quad g = G/B.$$

All h, l, r, e, g, j are strictly positive bounded elements; $a, t > 0$ can have all three types. Retain exactly

$$g = 1 - ae + hz > 0, \qquad j = 1 + zr - l/t > 0, \qquad B - zyr > 0, \qquad n \geq 0,$$

as well as all original source formulas and $\delta_{\mathrm{tot}} : \mathsf{I}$. The first identity gives

$$ae < 1 + hz; \quad e : \mathsf{U} \Rightarrow a \text{ bounded}; \quad a : \mathsf{O} \Rightarrow e : \mathsf{I}; \quad a : \mathsf{I} \Rightarrow g \sim 1; \quad g : \mathsf{I} \Rightarrow ae \sim 1. \tag{WF}$$

The exact coefficient estimates WII1–WII4 prove that all six retained event coefficients are bounded if and only if t, h, l, j, r, g are units. With $d = 1$, that would make δ_{tot} a unit. Therefore the all-six-unit branch is excluded on the present input. Also the two-weak source exit is exactly $r, g, H/y, j : \mathsf{I}$; when $t : \mathsf{U}$ this is $r, g, h, j : \mathsf{I}$.

After those exits and the regular A-opening admissions, the ten ordered rows are as follows. Every row includes all preceding weak-II equations and original coefficients; a type table does not assert its realizability.

Row	Exact additional type condition
R1	$t : \mathsf{I}$.
R2	$t : \mathsf{O}$.
R3	$t, h : \mathsf{U}$, $l : \mathsf{I}$.
R4	$t, h, l : \mathsf{U}$, $j : \mathsf{I}$.
R5	$t, h, l, j : \mathsf{U}$, $r : \mathsf{I}$.
R6	$t, h, l, j, r : \mathsf{U}$, $g : \mathsf{I}$.
R7	$t : \mathsf{U}$, $h, g : \mathsf{I}$, excluding r, j both infinitesimal.
R8	$t, g : \mathsf{U}$, $h, j : \mathsf{I}$.
R9	$t, g, j : \mathsf{U}$, $h, e, a : \mathsf{I}$.
R10	$t, g, j : \mathsf{U}$, $h, e : \mathsf{I}$, $a : \mathsf{O}$.

Here is the proof of exhaustion, including every discarded type. Split t first. For $t : \mathsf{U}$, split h. If $h : \mathsf{U}$, first test $l : \mathsf{I}$, then $j : \mathsf{I}$, then $r : \mathsf{I}$, then $g : \mathsf{I}$. These are R3–R6. If none applies, all six variables t, h, l, j, r, g are units, contradicting the total-scale input by WII4. If $h : \mathsf{I}$, $l \sim 1$. First test

$g : \mathsf{I}$: either $r, j : \mathsf{I}$ and the two-weak theorem applies, or this is R7. For $g : \mathsf{U}$, test $j : \mathsf{I}$, giving R8. Only $t, g, j : \mathsf{U}$, $h : \mathsf{I}$ remains.

On that final regular opening, if $e : \mathsf{U}$ then a is bounded by (WF). For $r : \mathsf{U}$, $a : \mathsf{U}$ gives WA and $a : \mathsf{I}$ gives WE; for $r : \mathsf{I}$ the SL theorem applies for either bounded type of a. If $e : \mathsf{I}$ then $r \sim 1$; $a : \mathsf{U}$ gives WC, $a : \mathsf{I}$ gives R9, and $a : \mathsf{O}$ gives R10. WA is joined to its printed remote theorem; WE, SL with its projective join, and WC each have exactly the source conditions just stated. No h/a or e/B comparison is inserted.

The table is disjoint by this order. The focused checker enumerates all type assignments obeying the necessary consequences of (WF) and $h + l = r + e = 1$, and confirms that every surviving tuple has precisely the displayed destination or one named exit. It does not assume that every such combinatorial tuple is realized by a physical source.

R19 already admits R5, R8, R9, R10, and portions of R3, R4, R6, R7. The supplied R19 frontier records the remaining exact intersections: R3 and R4 have $e : \mathsf{I}$; R6 has $a : \mathsf{O}$, $e : \mathsf{I}$, $\rho = n/e^2 : \mathsf{U}$; R7 has $a : \mathsf{O}$, $e : \mathsf{I}$. R1 and R2 retain their full original residual source conditions. These are substitutions of existing theorem scopes after the coverage argument, not evidence that a remaining domain is nonempty.

177.7 Boundary, normalization, and testing rules

1. C1 $D = 0$ or $\beta = 0$ is carried through the rational source minor. If a later branch makes one impossible, its original coefficient formula must prove that fact. The other zero is still kept. Both zero simultaneously is outside the five/six-active coercive entry.

2. C2 $n = 0$ is retained in global sign entry, the weak-II formula, and every subsequent source inequality. It is never replaced by $n > 0$, nor used as a residue approximation to $n > 0$.

3. A source shape parameter or gap is strictly positive unless its definition is an original nonnegative coefficient. An infinitesimal opening is not a literal zero. In particular $M, J, E, Q, r, 1-r, h, l, g, j$ are not set to zero in an exact source formula.

4. $d = 1$ and $C = 2$ are legal positive pure-source normalizations; they do not turn the earlier condition $\delta_{\mathrm{tot}} : \mathsf{I}$ into a positive-total-scale source. Source-only coefficient weights and testing-only normalization factors have square roots in their respective fields and restore every certificate.

5. The source angles classify actual contacts. Independent testing angles remain arbitrary, with signs, zeros, and infinity treated in their homogeneous charts. A finite-chart proof alone is not an all-tests statement. The ledger distinguishes local theorems from their exact joins.

6. The optional rational-curve theorem requires coefficients in a single $\mathbb{R}(T)$, a specified physical real interval, and nonvanishing denominators there. It is not an arbitrary multiparameter exit. The C2-to-C1 overlap is physical only for $n = 0$: the overlap identity gives a negative C1 beta coefficient for $n > 0$.

177.8 Conclusion of this audit, with the remaining entries explicit

The omission routing, positive-chart maps, C1/C2 source-type disjunction, C14 refinement, and weak R1–R10 refinement are now explicit. Their exhaustion is proved on the algebraic source input with the stated type conditions, independently of nonemptiness and independently of optional historical exclusions. The exact one/two-hard and four-matching work completes the requested U1.A6 technical audit on its stated domain. The separate bounded-Δ proof repairs the US-U terminal statement mismatch without relying on a historical global assertion.

The precise remaining audit entries are:

1. U1.A5: the scale split and source partition were already audited in R18. The ordinary and finite-pitchfork subbundle with d, C, A, D, α units is now repaired using the recovered full-parent finite-edge theorem, verified exact source entry, and a complete projective join. The separate note displays all four corner quadratic forms and restores every retained packet. Its exact symmetry images are included only under their stated unit hypotheses. The separate C2 weak-α note proves the subdomain with all six of d, n, m, C, A, D units and its exact reflection, by showing every remaining singular zero-corner jet fits the recovered finite-ratio theorem. No bound on the actual tested ratio is assumed. Remaining weak-alpha shapes and their stabilizer images, the C1 projective pitchfork, and reciprocal all-tests terminals remain to be accepted.

2. U1.A7: the ledger now names essential and optional exits, but their complete all-tests proofs are not all freshly reaudited here. The former US-U provenance issue is repaired by the new full-parent theorem for $k, \eta, \zeta : \mathsf{I}$, bounded $1/e$, all original gaps and signs, and all independent tests. Its review is recorded separately from the imported terminal bundle.

3. U1.A9: parts (i)–(iv) have the explicit coverage construction above. Its terminal-substitution step remains conditional on the exact registry inputs and on the open local rows in the current frontier. Do not change a historical import to "newly proved" because its filename appears in the registry.

U1 remains open; the global representability bridge remains blocked by U1. No separate uniform bound on the number of local squares has been introduced. This note is a same-system mathematical audit with narrow exact symbolic corroboration, not external review or proof-assistant verification.

Chapter 178

Uniform C1: the bounded-reciprocal repair and its scope

Component `p41b39e2f94e1`
Source: `BASE/proofs/routing/us_unit/bounded_delta_body.tex`
SHA256: `41b39e2f94e168a6a415c854495c78514467a711418c4abd2f4e9ea7c85be029`
Inventory use: Complete bounded-Delta, bounded-W polynomial source/conductor/full-normal/hard-block/W-state proof; replace its historical e-Inf conclusion by fresh UJ and its outer citations by UJ3-UJ4 under UE4

178.1 Uniform C1: the bounded-reciprocal repair and its scope

The preserved uniform-source theorem explicitly assumes e infinitesimal. The preserved contact–conductor theorem explicitly assumes e unbounded. This note proves the missing unit-e case by extending the latter proof to bounded $\Delta = e^{-1}$ and bounded testing w. The extension uses the same exact complete parent, the same retained packets, and the same unlocalized conductor. The new work is a weight estimate that remains strict when both δ and w have weight zero, followed by an exhaustive testing-chart join. An old all-e status assertion is not a premise.

Theorem 178.1 (Uniform C1, bounded reciprocal e)**.** *Let the coefficient and testing fields be independent real closed extensions of $\mathbb{R}$. Use the original canonical uniform C1 source defined below, with $k, f, g > 0$ infinitesimal, $e = kf/g > 0$, and $\Delta = 1/e$ bounded. The shapes $a, t > 0$ are bounded and satisfy the original strict gaps $M = R - t > 0$, $J = T - a > 0$. Retain every original coefficient sign, including $D_{\mathrm{e}}, \beta \geq 0$, and every literal coefficient zero. Then the complete parent is tensor-SOS at every independent projective measurement pair and every real or complex testing state.*

Together with the preserved infinitesimal-e theorem, this proves the whole positive-e uniform source. It does not prove an unlisted source chart or close any remaining C14 or weak-C2 row. We use the already admitted separated bounded strict-Gram lemma: a finite bounded separated Gram with strictly positive real residue gives tensor-SOS. No inversion of a mixed element and no comparison between an independent coefficient and testing rate enters the proof.

178.2 Original source and complete parent

Put $\kappa = k^2$, $\eta = f^2$, $\zeta = g^2$, $K = 1 + k^2$, and

$$V_c = \kappa\eta\zeta, \quad L_c = 1 - a\eta, \quad H_c = 1 + (1-a)\eta, \quad Q_c = 1 - \kappa\zeta + aV_c,$$

$$R = 1 + k^2(1 - a\eta), \qquad T = 1 + k^2 - k^2\zeta t, \qquad M = R - t > 0, \quad J = T - a > 0.$$

The native coquotients are defined by the following short polynomials:

$$P_4 = 1 + \kappa + \eta\kappa(a - \kappa - 1) - \zeta\kappa t + V_c[-\zeta\kappa t^2 + 2\kappa t + a(1-t) + t],$$

$$
\begin{aligned}
P_\alpha &= 1+\kappa-a\eta\kappa+\zeta[(1+\kappa)-(2+\kappa)t]+V_c[-\zeta t^2+a(t-1)+t],\\
P_d &= (1+\kappa)^2-a\eta\kappa^2+\zeta\kappa[(\kappa+1)-(\kappa+2)t]\\
&\quad +V_c[-\zeta\kappa t^2+\kappa a(t-1)+\kappa t+\kappa+1],\\
c_2 &= 2K(1-t\zeta)L_c(1+V_c)H_cQ_c,\\
c_b &= 2KL_c[1+(1-t)\zeta]H_cQ_c,\\
c_3 &= 2KL_cH_cQ_c(1-\kappa\eta+tV_c),\\
c_5 &= 2KL_cH_cQ_c(1+V_c),\\
c_4 &= 2Q_cP_4,\qquad c_\alpha = 2L_cH_cP_\alpha,\qquad c_d = 4L_cQ_cP_d.
\end{aligned}
$$

Their residues are $2,2,2,2,2,2,4$. The original source is

$$
\lambda=n=0,\quad C=2,\quad d_{\mathrm{B}}=\frac{2a\eta c_d}{c_3},\quad m=\frac{2ae^2c_2}{Mc_3},\quad A=\frac{2a\eta^2c_4}{c_3},\quad \alpha=\frac{2a\Delta^2k^6\eta^2c_\alpha}{c_3},
$$
$$
D_{\mathrm{e}}+\frac{d_{\mathrm{B}}}{2}=\frac{2ac_5}{Jc_3},\qquad \beta+\frac{d_{\mathrm{B}}}{2}=\frac{2ae^2c_b}{tc_3},\qquad \Lambda=\frac{RTc_3}{2a}. \tag{1}
$$

The original event D_{e} is distinct from the testing factor $D=1+u^2$. For a testing state (W,X,Y,Z) put

$$
D=1+u^2,\quad D_v=1+v^2,\quad B=X+uZ,\quad Q_A=W+uY,\quad r=uW-Y,
$$
$$
r_2=vQ_A-B,\qquad r_5=r-v(Z-uX).
$$

The original full parent on this source is

$$
\begin{aligned}
F &= 4mr_2^2+8D_vr^2+4ADD_vY^2+(4D_{\mathrm{e}}+2d_{\mathrm{B}})r_5^2\\
&\quad +(4\alpha+2d_{\mathrm{B}})D(vY-Z)^2+(4\beta+2d_{\mathrm{B}})D_vB^2-2d_{\mathrm{B}}DD_vZ^2.
\end{aligned}
$$

With $H_2=4Tc_2$, $H_b=4Tc_b$, $H_3=4Rc_3$, $H_5=4Rc_5$, equation (1) gives exactly

$$
\begin{aligned}
\Phi:=\Delta^2\Lambda F &= H_2\frac{R}{M}r_2^2+H_b\frac{R}{t}D_vB^2+\Delta^2H_3\frac{T}{a}D_vr^2+\Delta^2H_5\frac{T}{J}r_5^2\\
&\quad +4RT\Delta^2\eta^2c_4DD_vY^2+4RT\Delta^4k^6\eta^2c_\alpha D(vY-Z)^2\\
&\quad +2RT\Delta^2\eta c_dD\{(vY-Z)^2-D_vZ^2\}.
\end{aligned} \tag{2}
$$

178.3 The Z-unit chart with bounded W

Assume u,v,X,Y are infinitesimal, Z is a unit, and the state has bounded coordinates. Normalize $Z=1$ in the testing field. The new argument allows arbitrary bounded W. Put

$$
p=t/R,\quad h_s=a/T,\quad q_A=v(W+uY),\quad q_B=v(1-uX),\quad N=B-pq_A,\quad M_B=r-h_sq_B.
$$

Subtract only the four nonnegative packets

$$
\Pi_1=H_2\frac{t}{M}N^2,\quad \Pi_2=H_b\frac{M}{t}D_vN^2,\quad \Pi_3=\Delta^2H_3\frac{J}{a}D_vM_B^2,\quad \Pi_4=\Delta^2H_5\frac{a}{J}M_B^2.
$$

No boundedness of their gap ratios is assumed. The exact remainder is

$$
\begin{aligned}
G:=\Phi-\sum_i\Pi_i &= H_2\{N^2-2q_AN+(1-p)q_A^2\}+H_bD_v\{N^2+2q_AN+pq_A^2\}\\
&\quad +\Delta^2H_3D_v\{M_B^2+2q_BM_B+h_sq_B^2\}\\
&\quad +\Delta^2H_5\{M_B^2-2q_BM_B+(1-h_s)q_B^2\}+\mathcal{W},
\end{aligned} \tag{3}
$$

where $\mathcal{W}$ is the last three terms of (2), with $Z = 1$. Use the pure-testing coordinate $w = (W + uY)/D$. Its full inverse is

$$X = N + pDvw - u, \qquad q_B = v(D - uN - puDvw),$$
$$W = w + \frac{u(M_B + h_s q_B)}{D}, \qquad Y = uw - \frac{M_B + h_s q_B}{D}. \tag{4}$$

The only testing denominator is the positive unit D.

If the pure-testing signed core is nonnegative, its product with $d_{\mathrm{B}} > 0$ is already a separated square and all event packets are nonnegative. Otherwise the sign symmetries allow $u, v > 0$. Explicitly, changing u to $-u$ and (Y, Z) to $(-Y, -Z)$, or changing v to $-v$ and (X, Z) to $(-X, -Z)$, preserves every event square and the core. The negative-core branch then forces $w > 0$: completing the two positive core squares leaves

$$\frac{uvDD_v}{\mathcal{D}}\{Duv\, w^2 - 2DD_v w + uv\}, \qquad \mathcal{D} = u^2v^4 + u^2v^2 + u^2 + v^4 + 2v^2 + 1 > 0,$$

which is nonnegative for $w \le 0$. The eliminated matrix is

$$\begin{pmatrix} 1 + v^2 + u^2v^2 & uv \\ uv & (D + v^2)/D \end{pmatrix} \succ 0.$$

Thus all radicals used below have nonnegative pure-field factors. Literal angle axes already have a nonnegative core.

178.3.1 Polynomial source change and exact scalar

Define pure-source quantities

$$\delta = \Delta R/K, \quad b^2 = \eta K/R, \quad Q = K + ak^2b^2, \quad T_n = K^2 - t\delta^2k^4Qb^2.$$

Here δ is positive bounded, k, b are infinitesimal, and

$$\eta = Kb^2/Q, \quad \Delta = \delta Q/K, \quad \zeta = \delta^2k^2Qb^2/K, \quad R = K^2/Q, \quad T = T_n/K. \tag{5}$$

Every denominator in (5) is a positive source unit. After this substitution put $C_i = K^2Q^3c_i$. These are polynomials: each native monomial $ck^j f^{2r} g^{2s} a^\alpha t^\tau$ maps to

$$c\delta^{2s}k^{j+2s}b^{2(r+s)}a^\alpha t^\tau K^{2+r-s}Q^{3-r+s}, \qquad -2 \le r - s \le 3.$$

No negative exponent or shape inverse occurs. Put

$$V_n = K^2 - tQuvw, \quad Y_n = KT_nuw - avV_n, \quad B_0 = vY_n - KT_n,$$
$$A_c = C_5T_n + aK(C_3D_v - C_5), \qquad \mathcal{C} = K^6Q^4T_n.$$

Use different letters for the conductor monomials and the event weights:

$$A_m = wv, \quad B_m = \delta v, \quad C_m = \delta b^2uw, \quad D_0 = \delta^2k^3b^2.$$

Direct substitution of (4) into (3), retaining every term, gives

$$\begin{aligned} F_0 := (\mathcal{C}G)|_{N=M_B=0} = {} & 4KQT_n^2[C_2(K^2 - tQ) + C_bD_vtQ]D^2A_m^2 \\ & + 4Q^2A_cD^2V_n^2B_m^2 \\ & + 4K^3C_4DD_v(KT_nC_m - ab^2B_mV_n)^2 \\ & + 4KQ^2C_\alpha DB_0^2D_0^2 \\ & + 2K^2QC_dD\{b^2B_m^2(Y_n^2 - K^2T_n^2 + 2aKT_nV_n) - 2K^2T_n^2B_mC_m\}. \end{aligned} \tag{6}$$

Define the leading polynomial and its complete error by

$$P = D(A_m^2 + B_m^2) + K(C_m^2 + D_0^2) - 2K^2 B_m C_m, \qquad E_0 = F_0 - 8K^2 D\mathcal{C}P.$$

The full first coefficients on $(N, \delta M_B)$ are

$$\begin{aligned}
S_B &= -8aK^2b^4C_4D_vV_n + 8\delta^2Q^2k^6b^4C_\alpha B_0 + 4KQb^2C_dB_0,\\
S_C &= 8K^3T_nb^2C_4D_v,\\
L_1 &= A_m[8K^3QT_n^2(C_bD_v - C_2)D]\\
&\quad + B_m[-8\delta K^2Q^2uD\,vV_nA_c + \delta K^3auvS_B] + C_m[\delta K^3auvS_C],\\
L_2 &= B_m[8K^2Q^2T_nDV_n(C_3D_v - C_5) - K^2T_nS_B] - C_mK^2T_nS_C.
\end{aligned} \tag{7}$$

In particular $E_0 \in (A_m, B_m, C_m)^2 + (D_0^2)$ and $L_1, L_2 \in (A_m, B_m, C_m)$ without ideal division.

178.3.2 Exact contact and the unlocalized conductor

Work in $\mathcal{R} = \mathbb{Q}[\delta, k, b, a, t][u, w, v]$ and put

$$\mathfrak{J} = (u - k, w - \delta k, v - \delta k^2 b^2),$$
$$x = B_m - C_m, \quad y = A_m - kC_m, \quad z = uB_m - kC_m, \quad h = kC_m - D_0.$$

The original source contact is

$$u = k, \quad v = \Delta k^2\eta, \quad (W, X, Y, Z) = (\Delta k, -k(1 - t\zeta), \Delta k^2 L_c, 1).$$

It becomes $(u, w, v, N, M_B) = (k, \delta k, \delta k^2 b^2, 0, 0)$. For completeness, its exact contact and derivative identities follow from the six native balances

$$c_b - c_2 = \zeta c_3, \qquad c_3 - c_5 = -\kappa\eta c_b, \qquad c_2 = (1 - t\zeta)c_5,$$

$$\begin{aligned}
0 &= 2\kappa c_2 + 2(1 + V_c)L_c c_4 - 2\kappa^2\zeta Q_c c_\alpha - Q_c c_d,\\
0 &= \kappa L_c c_2 + c_5 + (1 + V_c)L_c^2 c_4 + \kappa Q_c^2 c_\alpha + \tfrac{1}{2}(\kappa\zeta L_c^2 - 2L_c - \eta)c_d,\\
0 &= \kappa R c_2 + t\kappa V_c c_b + aV_c c_3 + Tc_5 + KV_c L_c^2 c_4 - \kappa^2\zeta K Q_c L_c c_\alpha - \tfrac{K}{2}(Q_c L_c + \eta)c_d.
\end{aligned}$$

They follow by multiplying the short P_4, P_α, P_d polynomials. Substitution into the derivatives of (2) proves that its value and all first derivatives vanish at the displayed contact. The four packets also vanish there to second order. Consequently

$$E_0 \in \mathfrak{J}^2, \qquad L_1, L_2 \in \mathfrak{J}. \tag{8}$$

Here $P \in \mathfrak{J}^2$ follows also from its exact square identity below. No asymptotic contact replacement is made.

Adjoin only the separated radicals

$$a_1 = \sqrt{wv}, \quad a_2 = \sqrt{\delta}\sqrt{uv}, \quad a_3 = b\sqrt{\delta k}\sqrt{uw}, \quad a_4 = \delta k^{3/2} b, \qquad q = a_1a_2 - a_3a_4.$$

Only $q^2 \in \mathcal{R}$ is needed in the polynomial lift. Direct multiplication gives

$$\begin{aligned}
C_m(u - k) &= z - ux,\\
C_m(w - \delta k) &= \delta y - wx,\\
C_m(v - \delta k^2 b^2) &= b^2(wz + \delta ky - kwx),\\
r_* := y + z - h - 2kx &= (v - \delta k^2 b^2)[(w - \delta k) + \delta(u - k)] - \delta k b^2 (u - k)(w - \delta k),\\
kC_m r_* &= q^2 - yz.
\end{aligned} \tag{9}$$

For any polynomial S successive monic divisions give

$$S = S_c + (u-k)T_u + (w-\delta k)T_w + (v-\delta k^2 b^2)T_v.$$

If $S_c = 0$, (9) yields the explicit polynomial lift

$$C_m S = x(-uT_u - wT_w - b^2 kwT_v) + y(\delta T_w + \delta k b^2 T_v) + z(T_u + b^2 wT_v). \tag{10}$$

Thus

$$(A_m, B_m, C_m) \cap \mathfrak{J} \subset (x,y,z), \qquad [(A_m, B_m, C_m)^2 + (D_0^2)] \cap \mathfrak{J}^2 \subset (x,y,z,h,q)^2. \tag{11}$$

The restriction to the unmixed generator D_0^2 is substantive. For clarity, here is the complete constructive argument. In the first inclusion write $F = R_A A_m + R_B B_m + R_C C_m$ and substitute $A_m = y + kC_m$, $B_m = x + C_m$. The remaining coefficient of C_m is $S = kR_A + R_B + R_C$. The contact equality implies $S_c = 0$ by cancellation in the polynomial domain $\mathbb{Q}[\delta, k, b, a, t]$, so apply (10).

For the second inclusion write $F = \sum_{i \le j} f_{ij} m_i m_j + \rho D_0^2$, with $(m_1, m_2, m_3) = (A_m, B_m, C_m)$. Substitute the same formulas and $D_0 = kC_m - h$ to obtain

$$F = Q_2 + C_m(xH_x + yH_y + zH_z + hH_h) + C_m^2 S, \quad Q_2 \in (x,y,h)^2, \quad H_h = -2k\rho, \quad H_z = 0.$$

The contact value implies $S_c = 0$. Absorb $C_m^2 S$ into the four H coefficients by (10). Since $F \in \mathfrak{J}^2$, its first derivatives vanish at contact, while

$$\det\left(\frac{\partial(x,y,z)}{\partial(u,w,v)}\right)_c = -\delta^4 k^4 b^4 \neq 0.$$

The relation $r_* \in \mathfrak{J}^2$ has gradient coefficients $(-2k, 1, 1, -1)$. Put

$$\lambda = 2k\rho_c, \qquad J_i = H_i - \lambda(-2k, 1, 1, -1)_i.$$

The last J_i has zero contact value; the determinant and the first derivative equations imply the other three do too. This uses the adjugate and cancellation only to prove zero polynomial identities; no determinant inverse is a lifting coefficient. Finally

$$F = Q_2 + 2\rho_c(q^2 - yz) + \sum_{i=1}^{4} r_i(C_m J_i), \qquad (r_1, r_2, r_3, r_4) = (x, y, z, h),$$

and each $C_m J_i$ is lifted by (10). This proves (11), including all coefficient-zero strata, in the unlocalized ring.

178.3.3 The new positive-weight estimate

Assign weights

$$(\delta, k, b, a, t, u, w, v) = (0, 2, 6, 0, 0, 2, 0, 16). \tag{12}$$

The zero-weight variables are bounded; every positive-weight variable is infinitesimal in its own field. Positive weight of every polynomial coefficient therefore makes it a bounded separated infinitesimal, without any comparison of independent rates. A polynomial $O_{\mathrm{wt}}(s)$ has all monomials of weight at least s.

The short coquotients and (5) give

$$\begin{aligned}
&C_i = 2K^6 + O_{\mathrm{wt}}(12) \quad (i = 2, b, 3, 5, \alpha), && C_4 = 2K^6 + O_{\mathrm{wt}}(16), && C_d = 4K^7 + O_{\mathrm{wt}}(12),\\
&Q = K + O_{\mathrm{wt}}(16), && T_n = K^2 + O_{\mathrm{wt}}(20), && V_n = K^2 + O_{\mathrm{wt}}(18),\\
&D_v = 1 + O_{\mathrm{wt}}(32), && Y_n = O_{\mathrm{wt}}(2), && B_0 = -K^3 + O_{\mathrm{wt}}(18),\\
&A_c = 2K^8 + O_{\mathrm{wt}}(12), && \mathcal{C} = K^{12} + O_{\mathrm{wt}}(16).
\end{aligned}$$

The stronger C_4 estimate follows because every η-term in P_4 also contains κ, and the clearing correction $Q-K$ has weight 16. The two exact difference identities are

$$K(C_b-C_2)=\delta^2k^2Qb^2C_3,\qquad Q(C_3-C_5)=-k^2Kb^2C_b.$$

Both differences therefore have weight at least 16; since K,Q have constant term one, this conclusion is polynomial and introduces no inverse into a lifting coefficient.

The monomial weights of (A_m,B_m,C_m,D_0) are $(16,16,14,18)$. In the five lines of (6), keep the respective leading terms

$$8K^{14}D^2A_m^2,\quad 8K^{14}D^2B_m^2,\quad 8K^{15}DC_m^2,\quad 8K^{15}DD_0^2,\quad -16K^{16}DB_mC_m.$$

The first two errors have weight at least 44. The third error has weight at least 42: its only possibly smaller cross term contains $b^2B_mC_m$, of weight 42. The fourth error has weight at least 48. In the fifth line the coefficient error on B_mC_m has weight at least 42, and the separate $b^2B_m^2$ term has weight at least 44. Their sum is $8K^{14}DP+O_{\mathrm{wt}}(42)$. Since $\mathrm{wt}(P)\ge 28$ and $\mathcal{C}-K^{12}=O_{\mathrm{wt}}(16)$, this proves

$$E_0=O_{\mathrm{wt}}(42).$$

Also $S_B,S_C=O_{\mathrm{wt}}(12)$, and (7) gives

$$L_1=O_{\mathrm{wt}}(32),\qquad L_2=O_{\mathrm{wt}}(26).\tag{13}$$

It remains to check that the explicit conductor divisions preserve strict positivity of coefficient weights. The contact substitutions $u\mapsto k$, $w\mapsto\delta k$, $v\mapsto\delta k^2b^2$ never decrease weight; their weights change respectively as $2\mapsto 2$, $0\mapsto 2$, $16\mapsto 16$. For $S=O_{\mathrm{wt}}(s)$, its monic quotients obey

$$\mathrm{wt}(T_u)\ge s-2,\quad \mathrm{wt}(T_w)\ge s,\quad \mathrm{wt}(T_v)\ge s-16.$$

The coefficients in (10) on (x,y,z) consequently have weights at least $(s-2,s-2,s-4)$.

For a first-normal polynomial of weight p, monomial division by (A_m,B_m,C_m) gives quotient weights at least $(p-16,p-16,p-14)$. Hence $S=kR_A+R_B+R_C$ has weight at least $p-16$, and the completed lift has coefficient weights at least $p-20$. From (13) these bounds are 12 for L_1 and 6 for L_2.

For E_0 of weight at least 42, its monomial quotients on

$$A_m^2,A_mB_m,A_mC_m,B_m^2,B_mC_m,C_m^2,D_0^2$$

have weights at least $(10,10,12,10,12,14,6)$. Thus the initial Q_2 coefficients have weight at least 6, the initial (H_x,H_y,H_h) have bounds $(10,10,8)$, and S has bound 10. The first use of (10) leaves bounds $(8,8,6,8)$ on (H_x,H_y,H_z,H_h). Since ρ_c has weight at least 6 and $\lambda=2k\rho_c$ at least 8, subtracting the gradient relation leaves the same bounds on (J_x,J_y,J_z,J_h). A second use of (10) therefore leaves every scalar-lift coefficient with weight at least 2. The separate coefficient $2\rho_c$ of q^2 has bound 6. Every correction coefficient is thus strictly infinitesimal.

178.3.4 The full strict Gram and restoration

The eleven original soft roots are $g_0=x$, $g_4=uwv-\delta^2k^4b^2$, the six differences $a_i^2-a_j^2$, and the three differences $a_ia_j-a_\ell a_m$ over the three pairings of four indices. They satisfy the exact identity

$$P=g_0^2+g_4^2+\frac{1}{3}\sum_{i<j}(a_i^2-a_j^2)^2+\frac{2}{3}\sum_{\{ij,\ell m\}}(a_ia_j-a_\ell a_m)^2.\tag{14}$$

The conductor roots x,y,z,h,q are respectively $g_0,a_1^2-a_3^2,a_2^2-a_3^2,a_3^2-a_4^2,a_1a_2-a_3a_4$. Thus (8)–(13) express the complete scalar correction and both full first normals as infinitesimal perturbations on this same finite root frame. The leading diagonal in (14), multiplied by $8K^2D\mathcal{C}$, has strictly positive residue.

There are no state-normal terms above degree two. To display the entire hard block before multiplication by $\mathcal{C}$, put

$$C_B = H_3 D_v h_s + H_5(1 - h_s),$$
$$B_Y = 4RT\Delta^2\eta^2 c_4 D D_v + 4RT\Delta^4 k^6 \eta^2 c_\alpha D\, v^2 + 2RT\Delta^2 \eta c_d D\, v^2.$$

The entries on $(N, \delta M_B)$ are

$$A_{11} = H_2 + H_b D_v + \Delta^2 C_B u^2 v^2 + B_Y h_s^2 u^2 v^2 / D^2,$$
$$A_{12} = -\frac{\Delta^2}{\delta}(H_3 D_v - H_5)uv - \frac{B_Y}{\delta D^2} h_s uv,$$
$$A_{22} = \frac{\Delta^2}{\delta^2}(H_3 D_v + H_5) + \frac{B_Y}{\delta^2 D^2}.$$

Every apparent negative power of δ cancels: $\Delta/\delta = Q/K$ is a source unit, and B_Y/δ^2 is bounded infinitesimal even when δ is a unit. Hence this exact hard block has residue $16I_2$. Its denominators are only positive pure-source or pure-testing units. Together with (14) and the positive-weight corrections, the full thirteen-root Gram for $\mathcal{C}G$ is bounded and has strictly positive residue. Apply the admitted separated strict-Gram lemma, restore all four packets $\mathcal{C}\Pi_i$, and undo the positive source multiplier $\mathcal{C}\Delta^2\Lambda$. This proves the entire original parent in the Z-unit chart with bounded W, retaining both strict gaps and every original event.

178.4 Every remaining testing chart

Outer angle charts. The two preserved outer-angle identities require only $d_{\mathrm{B}}/C, d_{\mathrm{B}}/D_{\mathrm{e}}$ infinitesimal and the harmonic mass $\gamma = A[CD_{\mathrm{e}}/(C + D_{\mathrm{e}})]/d_{\mathrm{B}}^2$ bounded with residue at least $1/4$. These hypotheses are independent of the size of e. Indeed, put $B_c = 2c_5 - J\eta c_d$. The original source gives

$$D_{\mathrm{e}} = \frac{aB_c}{Jc_3}, \quad \frac{d_{\mathrm{B}}}{D_{\mathrm{e}}} = \frac{2J\eta c_d}{B_c}, \quad \frac{d_{\mathrm{B}}}{C} = \frac{a\eta c_d}{c_3},$$

and

$$\gamma = \frac{c_4 c_3 B_c}{c_d^2(2Jc_3 + aB_c)}.$$

Here $\overline{B}_c = 4$, $J + a = T$ has residue one, the denominator has residue 64, and the numerator has residue 16. Thus both strength ratios are infinitesimal and $\overline{\gamma} = 1/4$. The exact retained-current identities in `c1_uniform_all_tests.tex`, Lemmas 1–2, cover respectively $1/v$ bounded and v infinitesimal with $1/u$ bounded, with all homogeneous infinity endpoints. Consequently only u, v infinitesimal remain. The literal axes have a nonnegative core.

State charts. Normalize a nonzero state by a coordinate of maximal absolute value. All its coordinates are then bounded and at least one is a unit. If X or Y is a unit, the signed core has positive residue $2(X^2 + Y^2)$ and gives a direct separated square; restore all event squares. Otherwise X, Y are infinitesimal. If Z is a unit, the preceding section applies, including bounded unit W.

The only remaining case has Z infinitesimal and W a unit. Normalize $W = 1$ and put $Q_A = 1 + uY$, a pure-testing unit. Use

$$z = (r_2, B, \Delta r, \Delta r_5, \Delta\eta Y).$$

In (2), retain the four nonnegative excess packets obtained from $R/M = 1 + t/M$, $R/t = 1 + M/t$, $T/a = 1 + J/a$, $T/J = 1 + a/J$, the complete α packet, and the positive v^2Y^2 part of the last line. The remaining diagonal is

$$H_2 z_1^2 + H_b D_v z_2^2 + H_3 D_v z_3^2 + H_5 z_4^2 + 4RTc_4 D D_v z_5^2,$$

and its complete negative correction is exactly

$$-4RTc_dD\frac{\Delta Z}{Q_A}z_5(z_1+z_2)-2RTc_dD\frac{\Delta^2\eta Z^2}{Q_A^2}(z_1+z_2)^2.$$

This is the preserved W-unit identity, now on the larger domain: Δ is bounded and Z is infinitesimal, so both corrections are infinitesimal. The diagonal residue is $8I_5$. The separated strict-Gram lemma and restoration of all six packets prove this last chart. There is no need for Δ itself to be infinitesimal.

The zero state is immediate. A complex state splits into its real and imaginary quadratic evaluations because the parent matrix is real symmetric. Every normalization and projective clearing factor is pure testing. This completes the proof of Theorem 178.1.

Provenance and focused exact check

The source coquotients, full parent, retained packets, contact balances, and normal formulas are those preserved in `prior/prior_core/c1/source_definitions.tex`; the polynomial conductor is from its adjacent `contact_conductor_proof.tex`. The eleven-root identity and full hard block are in the preserved `c1_unbounded_e_z_obligation.tex`. Their exact identities do not require the old infinitesimal-δ hypothesis. The new weight argument (12)–(13), the bounded-W use, and the final W-unit/Z-infinitesimal join are the scope extension proved here. Exact paths and inherited hashes are in the companion routing ledger.

The adjacent standard-library checker reconstructs the source from the seven short coquotients, performs only monic contact division and exact removal of displayed monomial factors, and multiplies back the complete scalar and both first-normal lifts. Its 33 assertions pass. The actual minimum weights are $44, 32, 26$ on (E_0, L_1, L_2) and $12, 14, 12$ on the constructed correction families, stronger than the human bounds above. No Gröbner basis, numerical positivity test, parameter inversion, or prerecorded source expansion is used. The checker corroborates the proof; it does not replace the displayed conductor or the complete-parent chart join.

Chapter 179

The remaining R6 boundary: a^2B infinitesimal

Component p75b395956985
Source: BASE/proofs/weak_c2/r6_b_inf_all_tests.tex
SHA256: 75b395956985b917068fe4c552c70fead8e5112d4463087bd7aa1570894475d5
Inventory use: RI1-RI20 and adjacent-state/remote paragraphs

179.1 The remaining R6 boundary: a^2B infinitesimal

Source scope and relation to the preceding theorem. Retain the complete original weak-C2 source: $d = 1$, $\lambda = \beta = 0$, $n \geq 0$, the actual stationary contact, every strict gap, total scale, and the complement of prior admissions. On R6, $t, h, l = 1 - h, r$ and the original j gap are positive bounded units, while $g, B, x = aB, y = tB$ are positive infinitesimals. This section treats a unbounded and $b = a^2B$ positive infinitesimal. The notation $s = 1/a$, $w_s = ae$, $B = s^2b$, $x = sb$, $e = s(1 + hsb - g)$ is unchanged from (RB1). The exact physical identity (RB2), equivalently RT2, remains in force. It bounds b on all unbounded-a R6 and gives $\rho_0 = 1/2$ here, where $\rho = n/e^2$. In particular this subdomain has $n > 0$; the literal $n = 0$ source face is covered by the preceding b-unit theorem.

Theorem 179.1. *On this entire original R6 source intersection with $b = a^2B$ infinitesimal, the complete original parent is a finite separated source–testing tensor sum of squares at every independent projective testing pair and every real or complex state. Both angle signs, all literal projective endpoints, and all relative opening rates of s, b, g are included. No mixed inverse or mixed square root is used.*

Together with the b-unit theorem and the inherited bounded-a R6 theorem, this proves the entire R6 row. It does not by itself admit another weak-C2 row or the global finite-lift theorem.

A regular source chart. Keep $R = 1 + hx$, $K = 1 + x$, $T = 1 + xr$, $k = \sqrt{x}$ and the contact parameters $\delta = R/(K\sqrt{bt})$, $E = tK/R$ from (RB8). Define

$$\varepsilon = \delta^{-1}, \qquad \jmath = \delta k, \qquad k = \varepsilon\jmath. \tag{RI1}$$

Both ε and $\jmath$ are positive infinitesimals. In this section $\jmath$ denotes the contact scale, not the original source gap called j. The exact source substitution is

$$\begin{gathered} x = \varepsilon^2\jmath^2, \quad K = 1 + x, \quad R = 1 + hx, \quad E = tK/R, \\ b = \frac{\varepsilon^2 R^2}{K^2 t}, \qquad s = \frac{t\jmath^2 K^2}{R^2}. \end{gathered} \tag{RI2}$$

All denominators in (RI2) are positive source units. There is no comparison of a testing variable with either source infinitesimal.

Use precisely the complete contact-centered remainder G in (RB11), with the full coordinate inverse (RB7). Thus the two retained packets on the contact chart are $(m-K_m)N^2$ and $nD_uD_v(X+\gamma v\omega)^2$, where

$$p=(1+hx-g)/R,\quad c=l/(tT),\quad \gamma=rK/(sR),\quad D_u=1+u^2,\quad D_v=1+v^2.$$

Its exact contact is now

$$(u,\omega,v;N,M_B)=(\varepsilon\jmath,\jmath,\varepsilon\jmath^2E;0,0). \tag{RI3}$$

The six actual-contact checks in the preceding receipt prove the vanishing of both normals and of the retained n row here as well.

Two exact source cancellations. The quantities mg, $n\gamma$, $n\gamma^2$ and $\alpha/(s^3b^2)$ have canceled denominators that are units at $s=b=g=0$. Introduce

$$\mu_1=mg/R,\quad \mu_2=(mg)g/R^2,\quad F_0=p^2/2-n\gamma^2-2n\gamma p+\mu_2,\qquad A_N=p-2\mu_1-2n\gamma. \tag{RI4}$$

Both F_0 and A_N are divisible by b in this rational unit ring. Here is a short exact verification, independent of a series expansion. At $b=0$, put $q=1-g$. Direct cancellation in the original source gives

$$p=q,\quad mg=\frac{q}{2(1+s)},\quad n\gamma=\frac{sq}{2(1+s)},\quad n\gamma^2=\frac{q(1-sq)}{2(1+s)}. \tag{RI5}$$

Substitution makes both expressions in (RI4) identically zero. The denominators are units before this specialization; hence the polynomial numerators are divisible by b, without introducing a pole. Their needed residues are

$$\frac{F_0}{b}\sim\frac{h+t}{2},\qquad \left.\frac{A_N}{b}\right|_{s=g=0}=0,\qquad \left.\frac{A_N}{bs}\right|_{g=0}\sim h+t. \tag{RI6}$$

The last quotient also has a unit canceled denominator at $s=b=0$. Every statement in (RI5)–(RI6) is separately checked from the original source formulas. In particular the source estimate is uniform in g/s, s/b and all other opening ratios.

Complete bounded raw support, including both first normals. Use the separated raw monomials

$$\mathsf{A}=\varepsilon\omega v,\quad \mathsf{B}=v,\quad \mathsf{C}=Eu\omega,\quad \mathsf{D}=\varepsilon^2\jmath^3E,\qquad \mathsf{x}=\mathsf{B}-\mathsf{C}. \tag{RI7}$$

Set $H=\alpha+1/2$, $L=AD_v+Hv^2$, and $V_0=1-puv\omega$. At zero normals, (RB7) gives

$$q_B=D_uvV_0,\quad uW-Y=cD_uvV_0,\quad Y=u\omega-cvV_0,\quad J=-(1-c)D_uvV_0,$$

and the entire scalar has the following exact allocation:

$$G_0=f_A\mathsf{A}^2+f_{BB}\mathsf{B}^2+f_{BC}\mathsf{BC}+f_{CC}\mathsf{C}^2+\varrho\mathsf{D}^2, \tag{RI8}$$

where

$$\begin{aligned}
f_A&=D_u^2D_vF_0/\varepsilon^2,\\
f_{BB}&=[Cc^2D_u^2D_v+(D+1/2)(1-c)^2D_u^2+D_uLc^2]V_0^2\\
&\quad+2HD_ucV_0-D_u/2-D_u^2\mu_2v^2\omega^2,\\
f_{BC}&=[-2D_uLcV_0-2HD_u+2n\gamma D_uD_v]/E,\\
f_{CC}&=[D_uL+D_uD_vn\gamma^2v^2]/E^2,\\
\varrho&=\alpha D_u/\mathsf{D}^2.
\end{aligned} \tag{RI9}$$

These coefficients are bounded. In particular

$$\frac{F_0}{\varepsilon^2}=\frac{R^2}{K^2t}\frac{F_0}{b},\qquad \frac{\alpha}{\mathsf{D}^2}=\frac{1}{t}\frac{\alpha}{s^3b^2}. \tag{RI10}$$

The term that would give u^2/ε^2 in the first raw coefficient has instead been placed in f_{CC}, using the exact identity $u^2(v\omega)^2 = (v\mathsf{C}/E)^2$. The final term of f_{BB} likewise allocates the whole $-D_u^2\mu_2 v^4\omega^2$ contribution without dividing by ε. These are identities for the full scalar.

Write the exact normal expansion as $G = G_0 + L_N N + L_M M_B + G_{\mathrm{hard}}$, with G_{hard} quadratic. The two complete linear coefficients are

$$L_N = R_A\mathsf{A} + R_B\mathsf{B} + R_C\mathsf{C}, \qquad L_M = S_B\mathsf{B} + S_C\mathsf{C}, \tag{RI11}$$

where

$$\begin{aligned}
R_A &= D_u A_N/\varepsilon,\\
R_B &= D_u(p-2n\gamma)v^2\omega\\
&\quad - 2uv[Cc^2D_uD_v + (D+1/2)(1-c)^2D_u + Ac^2D_v]V_0\\
&\quad - 2Hcuv(1+cv^2V_0),\\
R_C &= 2cuvL/E,\\
S_B &= 2\{D_u[CcD_v - (D+1/2)(1-c)] + cL\}V_0 + 2H,\\
S_C &= -2L/E.
\end{aligned} \tag{RI12}$$

Here $A_N/\varepsilon = \varepsilon R^2(A_N/b)/(K^2t)$ is bounded. Thus no first-normal pole has been discarded. Every hard coefficient is bounded as well: the only testing inverse in the full change of state is D_u^{-1}, and all original source coefficients in (RB11) are bounded after the displayed cancellation. At $u = v = 0$ the complete hard matrix is exactly

$$\mathrm{diag}(K_m + 1/2, A + C + D + 1/2). \tag{RI13}$$

The accompanying checker reconstructs G, differentiates in each normal, and verifies (RI8)–(RI13) by multiplication back.

A bounded conductor for these complete coefficients. Define the contact differences

$$\mathsf{y} = \mathsf{A} - k\mathsf{C}, \quad \mathsf{z} = u\mathsf{B} - k\mathsf{C}, \quad \mathsf{h} = k\mathsf{C} - \mathsf{D}.$$

The scaled conductor identities are

$$\begin{aligned}
\mathsf{C}(u-k) &= \mathsf{z} - u\mathsf{x},\\
\varepsilon\mathsf{C}(\omega - \jmath) &= \mathsf{y} - \varepsilon\omega\mathsf{x},\\
\mathsf{C}(v - \varepsilon\jmath^2E) &= E(\omega\mathsf{z} + \jmath\mathsf{y} - k\omega\mathsf{x}),\\
\mathsf{C}v(\omega - \jmath) &= E\omega(\omega-\jmath)\mathsf{z} + E\jmath\omega\mathsf{y} - Ek\omega^2\mathsf{x}.
\end{aligned} \tag{RI14}$$

These are SE5–SE6 with $w = \omega$, also checked directly here. For a monic contact division

$$S = (u-k)T_u + (\omega-\jmath)T_\omega + (v - \varepsilon\jmath^2E)T_v, \qquad T_\omega = \varepsilon U + vV,$$

the exact bounded lift is

$$\begin{aligned}
\mathsf{C}S &= \mathsf{x}[-uT_u - \varepsilon\omega U - Ek\omega^2V - Ek\omega T_v]\\
&\quad + \mathsf{y}[U + E\jmath\omega V + E\jmath T_v]\\
&\quad + \mathsf{z}[T_u + E\omega(\omega-\jmath)V + E\omega T_v].
\end{aligned} \tag{RI15}$$

No inverse of a small source parameter or a contact difference occurs.

We verify the hypothesis of (RI15) for the full coefficients, rather than import the different small-e source assumptions of SE8. Clear the finite source unit denominators and D_u^2. They can be chosen independent of ω and even in u, v. In (RI9), all ω dependence is through $uv\omega$ and $v^2\omega^2$; the other testing dependence is even in u, v. The same property holds after subtracting any model of the form used below.

For the scalar square-lift algorithm CC4, substitute $\mathsf{A} = \mathsf{y} + k\mathsf{C}$, $\mathsf{B} = \mathsf{x} + \mathsf{C}$, $\mathsf{D} = k\mathsf{C} - \mathsf{h}$ into (RI8). The coefficient requiring its first conductor is

$$S = k^2 f_A + f_{BB} + f_{BC} + f_{CC} + k^2 \varrho.$$

Its zero contact value follows by domain cancellation, as in CC4. Divide monically in the order u, then ω, then v. After $u = k = \varepsilon\jmath$, the ω divided difference belongs to $(\varepsilon v, v^2)$. Accordingly choose $T_\omega = \varepsilon U + vV$ with $U, V \in (v)$. Modulo ε, the remaining v polynomial is even, so $T_v \in (\varepsilon, v)$; also $T_u|_{\varepsilon=u=v=0} = 0$.

After this first conductor the coefficient of h remains $-2k\varrho$. The sole gradient relation is $(-2k, 1, 1, -1)$ on $(\mathsf{x}, \mathsf{y}, \mathsf{z}, \mathsf{h})$; the determinant of the first three contact gradients is $-\varepsilon^3\jmath^4E^2 \neq 0$. The adjugate cancellation of CC5 therefore removes the contact values without putting an inverse of this determinant into any coefficient. Subtract the contact multiple $2k\varrho_c(-2k, 1, 1, -1)$. Each of the four resulting coefficient remainders J_i vanishes at contact. Its restriction at $\varepsilon = u = v = 0$ is independent of ω: for the x coefficient this follows from $2f_{BB} + f_{BC}$; the other three restrictions are zero by the U, V, T_u, T_v properties just established and the factor k. Consequently the next monic ω divided difference of every J_i belongs to (ε, v). Formula (RI15) applies again.

The last scalar repair uses the exact circuit relation

$$k\mathsf{C}(\mathsf{y} + \mathsf{z} - \mathsf{h} - 2k\mathsf{x}) = Q_1^2 - \mathsf{yz},$$

where Q_1 is the first of the separated radical pairings below. This completes the same restricted scalar square lift as CC6. For (RI11), one conductor suffices: after replacing A, B as above, the coefficient multiplying C, restricted at $\varepsilon = u = v = 0$, is independent of ω directly from (RI12). Its divided difference therefore lies in (ε, v) and (RI15) applies. The full scalar and both full normals have thus been lifted over the bounded coefficient ring. The restricted raw support $\mathsf{A}^2, \mathsf{B}^2, \mathsf{BC}, \mathsf{C}^2, \mathsf{D}^2$ is essential; an unrestricted ideal claim is neither needed nor asserted.

The original stationary parent and the two retained contact-zero packets give the exact scalar value and full first-jet identities used here. The contact (RI3) is independent of g. Hence taking the exact coefficient difference divisible by g preserves these identities and every raw-support condition.

Strict residues of the complete contact lift. Suppose now that u, v, ω are all positive infinitesimals, as in the remaining contact state chart. Give

$$(\varepsilon, \jmath, u, \omega, v) \text{ weights } (1, 1, 2, 1, 3).$$

Then x has weight three and the five circuit roots have weight five. Every contact substitution and (RI14)–(RI15) preserves the weights. Put $\kappa = (h + t)/(2t)$ and

$$\Lambda = \frac{1 - s + s^2 + u^2 - [2l + h(h + t)]k^2}{2ht}, \qquad \mu = \frac{1}{h(1 + s)}.$$

Direct exact collection of (RI9) through weight ten gives

$$G_0 = \Lambda\mathsf{x}^2 + \kappa T_4 + O_{\mathrm{wt}}(12) \pmod{g}, \tag{RI16}$$

where

$$T_4 = \mathsf{A}^2 + (u\mathsf{B})^2 + (k\mathsf{C})^2 + \mathsf{D}^2 - 4k^2\mathsf{BC}.$$

The two complete normal collections are

$$L_N = O_{\mathrm{wt}}(8), \qquad L_M = \mu\mathsf{x} + O_{\mathrm{wt}}(7) \pmod{g}. \tag{RI17}$$

For transparency, the only source terms needed for (RI16)–(RI17), besides (RI6), are the following exact Taylor coefficients at $g = 0$:

$$\begin{aligned} A &= \frac{t}{2h}(1 - s + s^2) + O_{\deg(s,b)}(3), \\ C &= \frac{h+t}{2l} - \frac{t+1}{2l}s + \frac{t+1}{2l}s^2 + O_{\deg(s,b)}(3), \\ D &= \frac{1 - s + s^2}{2(t-l)} - \frac{sbt(h+t)}{2(t-l)^2} + O_{\deg(s,b)}(3), \\ n\gamma &= \frac{s}{2} - \frac{s^2}{2} - \frac{sb(h+t)}{2} + O_{\deg(s,b)}(3). \end{aligned} \tag{RI18}$$

Here the error denotes exact degree filtration in a rational ring with unit denominators. Also $s = tJ^2 + O_{\mathrm{wt}}(6)$, $c = (l/t)(1 - k^2) + O_{\mathrm{wt}}(6)$ and $E = t(1 + lk^2) + O_{\mathrm{wt}}(8)$. The symbolic checker verifies all degree-two source derivatives, the complete scalar through weight ten, and the complete M_B normal through weight five. The weight eight bound for L_N follows term by term from (RI6), (RI12). At $g = 0$, after the displayed unit denominators are cleared, all scalar weighted monomials have even weight, and all M_B linear monomials have odd weight. Hence the next possible degrees are twelve and seven, respectively.

The stronger coefficient-wise statement also holds for the displayed raw representation: $f_A - \kappa$ and $\varrho - \kappa$ have weight at least two, while

$$f_{BB} - \Lambda - \kappa u^2, \qquad f_{BC} + 2\Lambda + 4\kappa k^2, \qquad f_{CC} - \Lambda - \kappa k^2$$

each has weight at least six. The checker verifies their coefficients through weight four separately. Similarly $S_B - \mu, S_C + \mu$ have weight at least four separately, and (R_A, R_B, R_C) have weights at least $(3, 5, 5)$. Thus the chosen raw allocation of each exact difference already has the claimed filtration; no low-weight Gram coefficients are hidden by cancellations between raw products.

Apply the bounded conductor just proved to the exact differences in (RI16)–(RI17). The scalar correction Gram coefficients have weight at least two; the M_B normal corrections have weight at least two; the N normal corrections have weight at least three. They are all infinitesimal. The coefficients divisible by g lift boundedly by the same argument and are infinitesimal independently of every other opening. Restoring the finite unit denominators preserves this fact.

For strictness use the two pairings on the separated monomials

$$a_i^2 = (\mathsf{A}, u\mathsf{B}, k\mathsf{C}, \mathsf{D}), \qquad Q_1 = a_1a_2 - a_3a_4, \quad Q_2 = a_1a_3 - a_2a_4.$$

Their product relation gives the exact two-pairing identity (RB15), with the scaled roots. Its three-dimensional difference block has determinant $1/2$, and both pairing pivots are positive. The complete residue on $(\mathsf{x}, \mathsf{y}, \mathsf{z}, \mathsf{h}, Q_1, Q_2, N, M_B)$ therefore consists of this strict circuit with coefficient $\kappa > 0$, the N pivot $K_m + 1/2 > 0$, and

$$\begin{pmatrix} 1/(2ht) & 1/(2h) \\ 1/(2h) & A_* + C_* + D_* + 1/2 \end{pmatrix}, \quad A_* = t/(2h),\ C_* = (h+t)/(2l),\ D_* = 1/[2(t-l)]. \tag{RI19}$$

Its determinant is exactly

$$\frac{C_* + D_* + 1/2}{2ht} > 0. \tag{RI20}$$

All pivots are positive bounded units. The strict separated Gram lemma WA10 yields a finite tensor-SOS identity for the complete remainder. Restore the two actual nonnegative retained packets and all separated clearing factors. This proves the entire local contact chart, including every normal direction and arbitrary relative rates.

The ω-unit state chart uses the original n packet. When u, v are infinitesimal, $Z = 1$, and ω is a positive unit, retain $nD_uD_vX^2$ itself, together with $(m - K_m)N^2$. The complete remainder is (RB11) with its two negative $n\gamma$ and $n\gamma^2$ contributions absent. This choice is legitimate directly from $n \geq 0$ and does not require a contact lift away from the contact state scale.

Use the raw roots

$$(A_o, \mathsf{x}, \mathsf{D}, N, M_B), \qquad A_o = \omega v.$$

The only new inverse is $1/\omega$, which is pure testing and bounded. The relations $\mathsf{B} = A_o/\omega$ and $\mathsf{C} = A_o/\omega - \mathsf{x}$ put the entire scalar and both first normals in this frame, by the same full formulas (RI9), (RI12) with the absent n terms deleted. All coefficients are bounded; the constant term is still $\alpha D_u = \varrho \mathsf{D}^2$. Explicitly $\mathsf{A} = \varepsilon A_o$, so the first scalar term becomes $D_u^2 D_v (p^2/2 + \mu_2) A_o^2$ and the first N-normal term becomes $D_u(p - 2\mu_1)A_o$. The printed ε denominators therefore cancel before any coefficient estimate. Its exact residue is the A_o pivot $1/2$, the D pivot κ, the N pivot $K_m + 1/2$, and the coupled block (RI19). The first N normal has zero residue and the first M_B normal has residue x/h. These complete-frame residues are separate assertions in the receipt. Thus this entire state chart also has a strict finite separated Gram.

All other independent tests and states. Every source bound used by the noncontact joins of the preceding b-unit theorem is (RB4), which is valid here without a lower bound on b. For bounded unit u and infinitesimal v, retain the original n, α events and use the exact full-state form (RB22), whose determinant (RB23) is positive. For both angles infinitesimal, the SD5 exact sign reduction disposes of the nonnegative-core branch and literal zero axes by original-event squares. On the negative branch choose the exact gauges $u, v > 0$. If the normalized state has Z infinitesimal, BE4 gives the strict full-state frame (X, Y, uW, vW) using positive real caps of m, A. Otherwise normalize $Z = 1$; $\omega > 0$ is bounded, and its two possibilities are exactly the contact chart and the unit chart just proved. No source/testing ratio split is used.

For finite remote $v_0 \neq 0$, use the actual centered-event identity R5.5, the strict kernel determinant RT10 (printed as RB24), and a sufficiently large positive real cap below unbounded m. For unbounded u and bounded unit v, use R6.5 specialized to $n_0 = 0$ and the source residues (RB4). For unbounded u and infinitesimal v, use the exact BE $(\infty, 0)$ full frame $(p_u X + Z, W, Y, vX)$, $p_u = 1/u$, with positive caps of C, A. For unbounded v and bounded u, R6.6 at $n_0 = 0$ is strict. For both angles unbounded, R6.7 at $n_0 = 0$ has determinants $D_*/2$ and $A_*(K_m + 1/2)$ on its two nontrivial blocks. These are the same exact homogeneous matrices listed in `dependencies_r6_b_unit.json`; their uses import the identities, not a source theorem from another stratum. Literal reciprocal zeros are polynomial values of those charts. All signs and complex states follow by the same exact gauges and the real/imaginary split. This exhausts all independent tests.

Additional exact dependencies and audit scope. Beyond the preceding dependency index, the scaled algebra is SE5–SE7 in `inherited_r18/inherited_r17/inherited_r16/inherited_r13/small_e_all_r_contact.tex` relative to the R19 root. This section proves its required rational-support hypotheses afresh in (RI8)–(RI15); the small-e native source theorem, its physical parameter constraints, and its particular residues are not imported. CC4–CC6 supplies the monic square-lift mechanism and the gradient/repair argument. The checker corroborates exact algebra; the bounded polynomial support proof and exhaustive chart join are the mathematical argument above. Neither the checker nor this local R6 result claims a global finite lift or external peer review. The complete exact path list, including this additional algebraic dependency, is in `dependencies_r6_b_inf.json`.

Chapter 180

R6 at a positive residual value of a^2B: an all-tests theorem

Component `pb60eba754641`
Source: `BASE/proofs/weak_c2/r6_b_unit_all_tests.tex`
SHA256: `b60eba7546416bee689116b4e001ce4ebed168b613628d3b1f57bec41f638aff`
Inventory use: RB1-RB24

180.1 R6 at a positive residual value of a^2B: an all-tests theorem

Scope. All notation and all physical source constraints are the original weak-C2 ones. In particular $d = 1$, $\lambda = \beta = 0$, $n \geq 0$, and the actual finite stationary contact, every strict gap, the total-scale condition, and the complement of earlier admissions are retained. On R6, $t, h, l = 1 - h, j, r$ are positive bounded units, $B, x = aB, y = tB, g$ are positive infinitesimals, and here a is unbounded. Define

$$s = a^{-1}, \quad w_s = ae, \quad b = a^2B, \quad e = 1 - r, \qquad B = s^2b, \quad x = sb, \quad y = ts^2b. \tag{RB1}$$

The existing exact source identity gives

$$2r\Delta_* n/s^2 = w_s Q_* - bK_*, \qquad Q_* \sim \Delta_* \sim 1, \quad K_* \sim h + t. \tag{RB2}$$

The gap $g = 1 - w_s + hsb$ gives $w_s \sim 1$. Hence b is bounded and

$$\rho = n/e^2 \sim \frac{1 - b_0(h_0 + t_0)}{2}, \qquad 0 \leq b_0(h_0 + t_0) \leq 1. \tag{RB3}$$

Here and below a subscript zero means the real residue; a displayed residue formula with bounded source letters has the same meaning. No testing variable occurs in these source estimates.

Theorem 180.1. *On the entire original R6 source intersection, assume in addition that $b = a^2B$ is a positive bounded unit. Then the complete original parent is a finite sum of squares in the separated source–testing tensor ring at every independent projective testing pair and every real or complex state, with either angle sign and every literal projective endpoint. All positive s, g opening rates are independent. Literal $n = 0$ is included. The theorem makes no assertion about b infinitesimal.*

Source bounds needed by the proof. Put $R = 1 + hx$, $K = 1 + x$ and $T = 1 + x(1 - e)$. The original rational source formulas, after the pure-source substitutions (RB1), have only unit denominators after mg, n/s^2 and $\alpha/(s^3b^2)$ are formed. Their joint residues are

$$mg \sim \tfrac{1}{2}, \quad C \sim \frac{h + t}{2l}, \quad A \sim \frac{t}{2h}, \quad D \sim \frac{1}{2(t - l)}, \quad \frac{\alpha}{s^3b^2} \sim \frac{h + t}{2}. \tag{RB4}$$

Thus m is unbounded, C, A, D are positive units, and all four have positive real lower bounds. There is no s/g or g/s assumption. The canceled formulas and their residues are checked in the accompanying script. They are rational identities of the actual physical source; no formal negative-beta source is imported.

Two contact-zero packets remove the two troublesome terms. Write the complete parent in the normalization $P = F/4$:

$$\begin{aligned} P &= nD_uD_vX^2 + mM^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ &\quad + (D + \tfrac{1}{2})J^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2, \\ M &= vW - X + uvY - uZ, \qquad J = uW + uvX - Y - vZ, \quad D_u = 1 + u^2, \quad D_v = 1 + v^2. \end{aligned} \tag{RB5}$$

In the finite state chart $Z = 1$, use the pure-testing coordinate $\omega = (W + uY)/D_u$ and the exact normals

$$\begin{gathered} p = \frac{w_s}{R} = 1 - \frac{g}{R}, \quad c = \frac{l}{tT}, \quad q_A = D_uv\omega, \quad q_B = v(1 - uX), \\ N = X + u - pq_A, \quad M_B = uW - Y - cq_B. \end{gathered} \tag{RB6}$$

Their full inverse is

$$\begin{gathered} X = N + pD_uv\omega - u, \quad q_B = v(D_u - uN - puD_uv\omega), \\ W = \omega + \frac{u(M_B + cq_B)}{D_u}, \quad Y = u\omega - \frac{M_B + cq_B}{D_u}. \end{gathered} \tag{RB7}$$

The only testing denominator is the positive pure-testing unit D_u. Set

$$\delta = \frac{R}{K\sqrt{bt}}, \quad E = \frac{tK}{R}, \quad k = \sqrt{x}, \quad K_c = \delta^2Ek^2 = \frac{sR}{K}, \quad \gamma = \frac{r}{K_c} = \frac{rK}{sR}. \tag{RB8}$$

All of δ, E, b are positive bounded units in this theorem. The actual contact in these coordinates is

$$(u, \omega, v; N, M_B) = (k, \delta k, \delta k^2E; 0, 0). \tag{RB9}$$

For direct verification, its original state coordinates with $Z = 1$ are

$$u_c = \sqrt{sb}, \quad v_c = s\sqrt{bt}, \quad W_c = \sqrt{s/t}, \quad X_c = -r\sqrt{sb}, \quad Y_c = hs\sqrt{b/t}, \quad \omega_c = \sqrt{s/t}\,R/K. \tag{RB9a}$$

They give $u_cW_c - Y_c = ls\sqrt{b/t} = c\,v_c(1 - u_cX_c)$ and $X_c + u_c = e\sqrt{sb} = pD_{u_c}v_c\omega_c$. Also $\gamma v_c\omega_c = r\sqrt{sb} = -X_c$. Thus both normals and the retained n row vanish exactly, before any residue calculation. These three cancellations and the identities $\omega_c = \delta k$, $v_c = \delta k^2E$, $K_c = sR/K$ are separate symbolic assertions in the receipt. For any fixed positive real K_m, retain the two whole squares

$$(m - K_m)N^2, \qquad nD_uD_v(X + \gamma v\omega)^2. \tag{RB10}$$

The first coefficient is positive because m is unbounded. The second vanishes identically when $n = 0$. Both rows vanish at (RB9), with every first jet of their squares. No inverse of n is used.

The complete remaining polynomial is exactly

$$\begin{aligned} G &= K_mN^2 - 2\frac{mg}{R}q_AN + \frac{mg^2}{R^2}q_A^2 \\ &\quad - 2n\gamma D_uD_vX(v\omega) - n\gamma^2D_uD_v(v\omega)^2 \\ &\quad + CD_v(uW - Y)^2 + AD_uD_vY^2 + (D + \tfrac{1}{2})J^2 \\ &\quad + (\alpha + \tfrac{1}{2})D_u(vY - 1)^2 + \tfrac{1}{2}D_v(X + u)^2 - \tfrac{1}{2}D_uD_v. \end{aligned} \tag{RB11}$$

This is the entire parent minus precisely (RB10), including every first normal and every hard quadratic term. The exact cancellation $M = (g/R)q_A - N$ removes the whole g^{-1} pole before an estimate. Moreover $n\gamma = O(s)$ and $n\gamma^2$ is bounded. All coefficients in (RB11), after (RB7), are therefore bounded. The only nonconstant state-normal degree is one or two.

Raw support before the conductor. Define

$$A_0 = \omega v, \quad B_0 = \delta v, \quad C_0 = \delta E u \omega, \quad D_0 = \delta^2 k^3 E, \quad X_0 = B_0 - C_0. \tag{RB12}$$

At $N = M_B = 0$ let $V_0 = 1 - puv\omega$. Then

$$q_B = D_u v V_0, \quad uW = u\omega + u^2 cv V_0, \quad Y = u\omega - cvV_0, \quad J = -(1-c)D_u v V_0, \quad X + u = pD_u A_0. \tag{RB13}$$

Thus every term of the scalar of (RB11) lies, with bounded coefficients, in $(A_0, B_0, C_0)^2 + (D_0^2)$. The only constant is from α, and α/D_0^2 is bounded with residue $b(h+t)/2$. For the new negative n contribution use exactly

$$\begin{aligned} -2n\gamma D_u D_v X A_0 - n\gamma^2 D_u D_v A_0^2 &= 2n\gamma D_u D_v u A_0 \\ &\quad - (2n\gamma p D_u + n\gamma^2) D_u D_v A_0^2 - 2n\gamma D_u D_v A_0 N, \end{aligned} \tag{RB14}$$

where $uA_0 = B_0 C_0/(\delta^2 E)$. Both complete first normals lie in (A_0, B_0, C_0) with bounded coefficients. Every remaining hard entry is bounded. These are full rational support statements, not a scalar contact-value test. Clearing D_u^2 makes them polynomial; its inverse is a pure-testing square factor when restored.

The contact Gram and its exact leading calculation. Let

$$Y_0 = A_0 - kC_0, \quad Z_0 = uB_0 - kC_0, \quad H_0 = kC_0 - D_0.$$

Use the separated nonnegative monomials $a_i^2 = (A_0, uB_0, kC_0, D_0)$ and the two pairings $Q_1 = a_1 a_2 - a_3 a_4$, $Q_2 = a_1 a_3 - a_2 a_4$. Their exact circuit is

$$\begin{aligned} T_4 &= A_0^2 + (uB_0)^2 + (kC_0)^2 + D_0^2 - 4k^2 B_0 C_0 \\ &= \tfrac{1}{2}\{(Y_0 - Z_0)^2 + H_0^2 + Y_0^2 + (Z_0 + H_0)^2\} + Q_1^2 + Q_2^2. \end{aligned} \tag{RB15}$$

Its matrix on (Y_0, Z_0, H_0) is $\begin{pmatrix} 1 & -1/2 & 0 \\ -1/2 & 1 & 1/2 \\ 0 & 1/2 & 1 \end{pmatrix}$, which is positive definite with determinant $1/2$. All radicals separate: a_1 is pure testing, a_2, a_3 are products of pure-source and pure-testing radicals, and a_4 is pure source.

Here is enough explicit algebra to reproduce the leading collection. At $g = 0$, the canceled source expansions through the needed orders are

$$\begin{aligned} n/s^2 &= \tfrac{1}{2}\{1 - b(h+t)\} + O(s), & mg &= \tfrac{1}{2} + \tfrac{1}{2}(bh-1)s + O(s^2), \\ C &= \frac{h+t}{2l} - \frac{t+1}{2l}s + O(s^2), & A &= \frac{t}{2h} - \frac{t}{2h}s + O(s^2), \\ D &= \frac{1}{2(t-l)} - \frac{bt(h+t) + t - l}{2(t-l)^2}s + O(s^2), & \alpha &= \tfrac{1}{2}b^2(h+t)s^3 + O(s^4). \end{aligned} \tag{RB16}$$

Here $O(s^j)$ means exact divisibility by s^j in a rational ring with unit denominator, not an asymptotic substitution in the proof identity. Give k, u, ω weight one and v weight two. Put $q = b(h+t)$. Using $s = k^2/b$ and $\delta = (1 - lk^2 + O(k^4))/\sqrt{bt}$, $\delta E = \sqrt{t/b}$, the scalar $G_0 = G|_{N=M_B=0}$ satisfies

$$G_0 = \frac{b}{2h}(1+u^2)X_0^2 + \frac{q}{2}T_4 - \frac{k^2[1 + bh(h+t)]}{2h}X_0^2 + O_{\mathrm{wt}}(8) \qquad (\mathrm{mod}\ g). \tag{RB17}$$

The complete first-normal coefficients obey

$$L_N = O_{\mathrm{wt}}(5), \qquad L_{M_B} = \frac{X_0}{h\delta} + O_{\mathrm{wt}}(4) \qquad (\mathrm{mod}\ g), \tag{RB18}$$

and the hard block has residue

$$\mathrm{diag}\left(K_m + \tfrac{1}{2},\ A_* + C_* + D_* + \tfrac{1}{2}\right), \quad A_* = \frac{t}{2h},\ C_* = \frac{h+t}{2l},\ D_* = \frac{1}{2(t-l)}. \tag{RB19}$$

All equalities through the displayed weights are checked by rebuilding (RB11) from (RB5), not by fitting a numerical matrix.

Apply the inherited monic contact–conductor identities CC1–CC6 to the exact difference in (RB17), and to the exact first-normal differences in (RB18). Their hypotheses are exactly the raw support in (RB13)–(RB14), together with zero contact value and full first jet. Both follow from the original stationary parent and (RB10); the model X_0^2, T_4 also has these zeros. No contact difference is inverted. The conductor maps preserve this grading: X_0 has weight two, the five circuit roots have weight three. The scalar difference of weight at least eight and normal differences of the displayed weights therefore have strictly positive-weight Gram coefficients. Every such coefficient is infinitesimal. Terms divisible by g also lift with bounded coefficients and remain infinitesimal: the contact (RB9), δ and E are independent of g, and the canceled coefficient ring has unit denominators at $g = 0$. Clearing and restoring its finite unit denominators changes none of these conclusions.

Consequently the residue on the eight roots $(X_0, Y_0, Z_0, H_0, Q_1, Q_2, N, \delta M_B)$ is the strict circuit with coefficient $q/2 > 0$, the N pivot $K_m + 1/2 > 0$, and the coupled block

$$\begin{pmatrix} b/(2h) & bt/(2h) \\ bt/(2h) & bt(A_* + C_* + D_* + 1/2) \end{pmatrix}. \tag{RB20}$$

Its determinant is

$$\frac{b^2 t}{2h}(C_* + D_* + 1/2) > 0. \tag{RB21}$$

All pivots are positive bounded units. The separated strict-residue Gram lemma therefore gives finite tensor SOS for the exact remainder. Restore (RB10) and the pure-source/pure-testing clearing factors. This proves the whole local contact parent, for every independent rate and every normal direction.

All remaining finite state and angle charts. The following details make the join an all-tests result. For bounded u with nonzero residue and infinitesimal v, retain $nD_uD_vX^2$ and $\alpha D_u(vY - Z)^2$, and the centered m excess $(m - K_m)N^2$ from (RB6), now homogeneous in Z. Put $V_Z = vZ$ and $R_t = uW - Y$. After $X = N - uZ + pv(W + uY)$, the whole remainder is a bounded quadratic on (N, R_t, Y, V_Z). Its residue splits off $K_m + 1/2$ on N, with the other form

$$C_* R_t^2 + A_*(1 + u^2)Y^2 + (D_* + 1/2)\{R_t - (1 + u^2)V_Z\}^2 - (1 + u^2)YV_Z - \tfrac{1}{2}(1 + u^2)V_Z^2. \tag{RB22}$$

Its first two leading principal minors are positive, and its determinant is exactly

$$(1 + u^2)^2 A_* C_*(D_* + 1/2)u^2 > 0. \tag{RB23}$$

This direct four-root proof covers every state, including $Z = 0$, with no negative-beta or small-ρ argument.

It remains to consider both angles infinitesimal. The exact testing-core sign reduction from SD5 handles nonnegative core and literal axes by original-event squares. On its negative-core branch one may use the standard exact gauges $u, v > 0$. Normalizing the original state by a maximal coordinate, the case Z infinitesimal has W a unit and is covered by BE4, with positive real caps of m, A supplied by (RB4). Retain the actual nonnegative n, α packets. That exact selected full-parent frame is (X, Y, uW, vW) and has strict residue; no coefficient bound involving b enters its proof.

If Z is a unit normalize $Z = 1$. The same core identity gives $\omega > 0$, and ω is bounded. If ω is infinitesimal, use (RB17)–(RB21). If ω is a positive unit, use the exact relations $B_0 = (\delta/\omega)A_0$ and $C_0 = B_0 - X_0$ in the raw support (RB13)–(RB14). The inverse is pure testing. Its complete residue on $(A_0, X_0, D_0, N, \delta M_B)$ consists of the positive $q/2$ pivots on A_0, D_0, the $K_m + 1/2$ normal pivot, and precisely (RB20). Direct substitution in (RB11) gives these residues; all other coefficients contain k, g, u, v after the cancellations already recorded. Thus this whole state chart is strict as well.

Every other projective test. Use the exact centered-event identity R5.5 for the finite remote pairs $v_0 \neq 0$. All source residues are (RB4) and $n_0 = \alpha_0 = 0$. On the kernel of the limiting centered m row, the determinant is

$$\frac{(1+U)^2(1+V)^2(h+t)}{8hl(t-l)}\{V^2 + t(h+t)U^2V + hUV^2\} > 0. \tag{RB24}$$

The restriction to $Z = 0$ is positive by the original A, C packets and the nonnegative signed-core restriction, exactly as in RT10. A sufficiently large positive real cap K_m makes the full bounded residue strict. Its choice depends on the real testing residues only; unbounded m permits every such real cap.

For unbounded u and bounded unit v, R6.5 at $n_0 = 0$ is strict because C_*W^2, A_*Y^2 and $(D_* + 1/2)(W + vX)^2$ have common kernel zero. For unbounded u and infinitesimal v, the exact BE $(\infty, 0)$ frame $(pX + Z, W, Y, vX)$ retains every event except positive caps of C, A, and has strict residue. For unbounded v and bounded u, R6.6 at $n_0 = 0$ is strict; the centered correction vanishes on its m-row kernel. For both angles unbounded, R6.7 at $n_0 = 0$ has block determinants $D_*/2$ and $A_*(K_m + 1/2)$, both positive. These are exact homogeneous full-parent identities. Their literal reciprocal zeros include the actual projective endpoints; undoing a nonzero homogeneous factor uses a pure-testing square only.

This exhausts all independently chosen angle and state charts. Apply the real identities separately to the real and imaginary parts of a complex state. Every retained positive packet and original normalization is restored. The new admission is the entire stated b-unit source subdomain, rather than a scalar graph or a single local testing corner.

Exact residual boundary. Together with the inherited R18 thin-layer theorem, this leaves from unbounded-a R6 only $b = a^2B$ positive infinitesimal. Then (RB3) forces $\rho_0 = 1/2$. The strict block (RB20) cannot be used without new weights, because δ becomes unbounded and $q = b(h + t)$ loses residue. This is a failure of that bounded frame, not a counterexample to finite representability. The original physical constraints and all independent testing variables remain in force.

Exact dependency index. The accompanying `dependencies_r6_b_unit.json` names the exact inherited file, tagged equation or named paragraph, and the restricted use for every imported identity. In particular, BE4 is the complete selected four-root state identity; the BE $(\infty, 0)$ identity is the unnumbered paragraph beginning "At $(\infty, 0)$ retain all events except c, A" in the same file. RT10 supplies only the kernel determinant and its $Z = 0$ restriction. R5.5 supplies only the centered-event identity. R6.5–R6.7 supply the displayed homogeneous kernel or root matrices, specialized to $n_0 = 0$ and (RB4), rather than the bounded-a theorem surrounding them. SD5 supplies the pure-testing inverse and the immediately following exact sign Schur identity. This proof does not import any C1 physical-source theorem or use a negative-beta perturbation theorem.

Chapter 181

The unbounded-a complement of R7, including its physical zero face

Component p405d96527133
Source: BASE/proofs/weak_c2/r7_unbounded_a_body.tex
SHA256: 405d96527133461384023f15cf2a4facd13b072372e8a37b3d447cab45ee4dbb
Inventory use: RU1-RU30

181.1 The unbounded-a complement of R7, including its physical zero face

Quantified source and conclusion. Keep the entire original weak-C2 source: $d = 1$, $\lambda = \beta = 0$, the actual rational stationary coefficients, $n \geq 0$, every strict gap, the original finite stationary contact, total scale, and all prior exclusions. In R7, B, z, y are positive infinitesimals, $a = z/B$, $t = y/B$ is a positive bounded unit, and h, g are positive infinitesimals. The original exclusion of simultaneously infinitesimal r and the source gap j_{gap} is retained.

Theorem 181.1. *On the whole original R7 source intersection with a unbounded, the complete original parent is a finite separated source–testing tensor sum of squares for every independent projective testing pair and every real or complex state. Both angle signs, all literal zero and infinite angles, literal $n = 0$, and every relative opening rate of the source infinitesimals are included. No inverse or radical of a mixed source–testing expression is used.*

The inherited bounded-a R7 theorem and this theorem exhaust R7. This statement does not admit R3, R4, or the global finite-lift theorem.

Actual source scale and the physical inequality. Use

$$\begin{gathered} s = 1/a, \quad b = a^2 B, \quad x = sb = z, \quad B = s^2 b, \\ w_s = ae = 1 + hx - g, \quad e = sw_s, \quad r = 1 - e, \quad l = 1 - h. \end{gathered} \tag{RU1}$$

Set $K = 1 + x$, $R = 1 + hx$, $T = 1 + xr$ and $j_{\mathrm{gap}} = T - l/t$. The exact original source identity is

$$\begin{gathered} 2r\Delta n = eQ - BK_{\mathrm{phys}}, \qquad 2r\Delta_* n/s^2 = w_s Q - bK_*, \qquad \Delta_* = s\Delta, \\ Q = 1 + B(1 + t + ah + 2ar + at) + B^2\{a(h + t) + a^2[h(r + 1) + t(1 + r - r^2)]\}, \\ K_{\mathrm{phys}} = (a + 1)(h + t)(1 + x), \qquad K_* = (1 + s)(h + t)(1 + x). \end{gathered} \tag{RU2}$$

Before bounding b, $Q \sim \Delta_* \sim 1$ since all their nonconstant terms contain s or x. Here $w_s \sim 1$ and $K_* \sim t$ is a positive unit. The original inequality $eQ \geq BK_{\mathrm{phys}}$ therefore gives b bounded and $n = O(s^2)$, including its literal equality case. In particular $r \sim 1$, so the original R7 exclusion is respected. The source gap gives $t_0 \geq 1$; j_{gap} is a unit exactly when $t_0 > 1$, and otherwise $t_0 = 1$.

The original rational formulas, after cancellation, give jointly

$$mg \sim \tfrac{1}{2}, \quad C \sim \tfrac{t}{2}, \quad A_h := Ah \sim \tfrac{t}{2}, \quad \beta_D := D(1-c) = Dj_{\mathrm{gap}}/T \sim \frac{1}{2t}, \quad \frac{\alpha}{s^3 b^2} \sim \tfrac{t}{2}, \qquad c = \frac{l}{tT}. \tag{RU3}$$

Thus m, A are unbounded; C is a positive unit. If $t_0 > 1$, $D \sim 1/[2(t-1)]$ is a positive unit. If $t_0 = 1$, D is unbounded. The canceled coefficients $mg, C, A_h, \beta_D, n\gamma, n\gamma^2$ below have unit denominators through both small-gap ends. No ratio h/g, j_{gap}/h, or comparison with a testing rate is used.

Four whole packets and a bounded full-parent remainder. Use the complete parent $P = F/4$:

$$\begin{aligned} P &= nD_uD_vX^2 + mM^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ &\quad + (D + 1/2)J^2 + (\alpha + 1/2)D_u(vY - Z)^2 + D_v(X + uZ)^2/2 - D_uD_vZ^2/2, \\ M &= vW - X + uvY - uZ, \qquad J = uW + uvX - Y - vZ. \end{aligned}$$

The exact coordinate inverse (RB6)–(RB7) is

$$\begin{gathered} D_u = 1 + u^2, \quad D_v = 1 + v^2, \quad Z = 1, \quad p = w_s/R, \quad q_A = D_uv\omega, \\ N = X + u - pq_A, \qquad q_B = v(1 - uX), \qquad M_B = uW - Y - cq_B, \\ X = N + pD_uv\omega - u, \quad q_B = v(D_u - uN - puD_uv\omega), \\ W = \omega + u(M_B + cq_B)/D_u, \qquad Y = u\omega - (M_B + cq_B)/D_u. \end{gathered} \tag{RU4}$$

The sole testing inverse is D_u^{-1}. Put

$$\begin{gathered} \gamma = \frac{rK}{sR}, \quad \eta = Y - huW, \quad q = 1 - c = j_{\mathrm{gap}}/T, \\ \mu_1 = mg/R, \quad \mu_2 = (mg)g/R^2, \quad n_1 = n\gamma, \quad n_2 = n\gamma^2. \end{gathered}$$

Retain the four entire nonnegative squares

$$mN^2, \qquad nD_uD_v(X + \gamma v\omega)^2, \qquad AD_uD_v\eta^2, \qquad DM_B^2. \tag{RU5}$$

All four rows vanish at the actual contact; (RB9a) verifies the first two, and directly $Y_c = hu_cW_c$, $M_{B,c} = 0$. The n packet vanishes identically on the literal $n = 0$ face.

The full A, D cancellations are the exact identities

$$\begin{aligned} AD_uD_v(Y^2 - \eta^2) &= 2A_hD_uD_vuWY - A_hhD_uD_v(uW)^2, \\ D(J^2 - M_B^2) &= -2\beta_DM_Bq_B + \beta_Dqq_B^2, \qquad J = M_B - qq_B. \end{aligned} \tag{RU6}$$

Since $M = (g/R)q_A - N$, subtraction of precisely (RU5) leaves

$$\begin{aligned} G = \; & -2\mu_1q_AN + \mu_2q_A^2 - 2n_1D_uD_vX(v\omega) - n_2D_uD_v(v\omega)^2 \\ & + CD_v(uW - Y)^2 + 2A_hD_uD_vuWY - A_hhD_uD_v(uW)^2 \\ & - 2\beta_DM_Bq_B + \beta_Dqq_B^2 + J^2/2 \\ & + (\alpha + 1/2)D_u(vY - 1)^2 + D_v(X + u)^2/2 - D_uD_v/2. \end{aligned} \tag{RU7}$$

This is a whole-parent identity, including every scalar, normal and hard coefficient. Equations (RU3), (RU6) bound all of them before any contact restriction. In particular neither h^{-1} nor j_{gap}^{-1} survives.

One regular contact frame for the two b alternatives. Use the pure-source parameters

$$\varepsilon = \frac{K\sqrt{bt}}{R}, \quad \jmath = \sqrt{s/t}\,\frac{R}{K}, \quad k = \varepsilon\jmath = \sqrt{x}, \quad E = tK/R. \tag{RU8}$$

The symbol $\jmath$ is a contact scale, not j_{gap}. Here $\jmath$ is infinitesimal; ε is a unit or infinitesimal exactly as b is. The exact inverse source chart is

$$x=\varepsilon^2\jmath^2,\qquad b=\frac{\varepsilon^2R^2}{K^2t},\qquad s=\frac{t\jmath^2K^2}{R^2}. \tag{RU9}$$

Only source units are inverted. The contact is $(u,\omega,v;N,M_B)=(k,\jmath,\varepsilon\jmath^2E;0,0)$.

Define

$$\begin{gathered}\mathsf{A}=\varepsilon\omega v,\quad \mathsf{B}=v,\quad \mathsf{C}=Eu\omega,\quad \mathsf{D}=\varepsilon^2\jmath^3E,\\ \mathsf{x}=\mathsf{B}-\mathsf{C},\quad \mathsf{y}=\mathsf{A}-k\mathsf{C},\quad \mathsf{z}=u\mathsf{B}-k\mathsf{C},\quad \mathsf{h}=k\mathsf{C}-\mathsf{D}.\end{gathered} \tag{RU10}$$

The two potentially singular source combinations are

$$F_0=p^2/2-n_2-2n_1p+\mu_2,\qquad A_N=p-2\mu_1-2n_1.$$

Their exact canceled quotients satisfy

$$F_0/b\sim t/2,\qquad A_N/(bs)\sim t. \tag{RU11}$$

Both denominators are units through $s=b=h=g=0$. These assertions follow from the original rational source, not from the inequality $n\ge0$. In particular

$$F_0/\varepsilon^2=\frac{R^2}{K^2t}\frac{F_0}{b},\qquad \alpha/\mathsf{D}^2=\frac{1}{t}\frac{\alpha}{s^3b^2},\qquad A_N/\varepsilon=\frac{\varepsilon R^2s}{K^2t}\frac{A_N}{bs} \tag{RU12}$$

are bounded without losing the zero-n face.

Exact full scalar and both first normals. Put $H=\alpha+1/2$, $V_0=1-puv\omega$. At $N=M_B=0$, $uW=\mathsf{C}/E+u^2c\mathsf{B}V_0$ and $Y=\mathsf{C}/E-c\mathsf{B}V_0$. The complete scalar is

$$G_0=f_A\mathsf{A}^2+f_{BB}\mathsf{B}^2+f_{BC}\mathsf{B}\mathsf{C}+f_{CC}\mathsf{C}^2+\varrho\mathsf{D}^2, \tag{RU13}$$

with the bounded coefficients

$$\begin{aligned}f_A&=D_u^2D_vF_0/\varepsilon^2,\qquad \varrho=\alpha D_u/\mathsf{D}^2,\\ f_{BB}&=[Cc^2D_u^2D_v+(\beta_Dq+q^2/2)D_u^2+HD_uv^2c^2-A_hD_uD_v(2u^2+hu^4)c^2]V_0^2\\ &\quad+2HD_ucV_0-D_u/2-D_u^2\mu_2v^2\omega^2,\\ f_{BC}&=[2A_hD_uD_v(lu^2-1)cV_0-2HD_uv^2cV_0-2HD_u+2n_1D_uD_v]/E,\\ f_{CC}&=[A_h(2-h)D_uD_v+HD_uv^2+D_uD_vn_2v^2]/E^2.\end{aligned} \tag{RU14}$$

For the full normal expansion $G=G_0+L_NN+L_MM_B+G_{\mathrm{hard}}$, the exact coefficients are

$$L_N=R_A\mathsf{A}+R_B\mathsf{B}+R_C\mathsf{C},\qquad L_M=S_B\mathsf{B}+S_C\mathsf{C}, \tag{RU15}$$

where

$$\begin{aligned}R_A&=D_uA_N/\varepsilon,\\ R_B&=D_u(p-2n_1)v^2\omega-2uv[Cc^2D_uD_v+(\beta_Dq+q^2/2)D_u]V_0\\ &\quad+2A_hc^2uvD_vu^2(2+hu^2)V_0-2Hcuv(1+cv^2V_0),\\ R_C&=[2A_hcuvD_v(1-lu^2)+2Hcuv^3]/E,\\ S_B&=2[CcD_uD_v-(\beta_D+q/2)D_u-A_hcD_vu^2(2+hu^2)]V_0+2H(1+cv^2V_0),\\ S_C&=[2A_hD_v(lu^2-1)-2Hv^2]/E.\end{aligned} \tag{RU16}$$

At $u=v=0$ the complete hard matrix on (N,M_B) is exactly $\mathrm{diag}(1/2,C+1/2)$. The checker differentiates the entire (RU7) and multiplies back (RU13)–(RU16); these are not restricted Hessian or scalar-contact checks.

Bounded monic conductor, including the small-ε end. The identities (RI14)–(RI15) apply to (RU10) without a source-shape assumption. Their crucial inputs are checked here on the new complete coefficients. After finite source unit denominators and D_u^2 are cleared, every ω occurrence of (RU14) is through $uv\omega$ or $v^2\omega^2$, and all remaining testing powers are even in u, v. The scalar raw support is exactly the five terms in (RU13).

For completeness, first replace $\mathsf{A} = \mathsf{y} + k\mathsf{C}$, $\mathsf{B} = \mathsf{x} + \mathsf{C}$, and $\mathsf{D} = k\mathsf{C} - \mathsf{h}$. The coefficient requiring a first conductor is $S = k^2 f_A + f_{BB} + f_{BC} + f_{CC} + k^2\varrho$. Divide its contact-zero difference monically in order u, ω, v. At $u = k = \varepsilon\jmath$, the ω divided difference is in $(\varepsilon v, v^2)$; write it as $\varepsilon U + vV$ with $U, V \in (v)$. The last v divided difference is in (ε, v) because, modulo ε, the remaining v dependence is even. The first u divided difference restricts to zero at $\varepsilon = u = v = 0$.

These facts permit the exact bounded lift (RI15), which uses the identity for $\mathsf{C}v(\omega - \jmath)$ and never divides by ε. Subtract the gradient multiple $2k\varrho_c(-2k, 1, 1, -1)$. The first three contact gradients have determinant $-\varepsilon^3\jmath^4E^2 \neq 0$; the CC5 adjugate argument cancels a zero identity, and introduces no inverse of that determinant. Each second-stage coefficient at $\varepsilon = u = v = 0$ is independent of ω: the x one restricts to $2f_{BB} + f_{BC}$, and the other three to zero. Hence every second ω divided difference is in (ε, v) and has the same bounded lift. The final repair is

$$k\mathsf{C}(\mathsf{y} + \mathsf{z} - \mathsf{h} - 2k\mathsf{x}) = Q_1^2 - \mathsf{yz}. \tag{RU17}$$

The pairings Q_i are defined below. For each complete normal in (RU15), one conductor suffices: after replacing A, B, its C coefficient restricted at $\varepsilon = u = v = 0$ is independent of ω, directly from (RU16). This is precisely the normal criterion in the same proof. When b is a unit, ε^{-1} is bounded and the ordinary CC1–CC6 conductor is also available.

All contact value and first-jet hypotheses follow from the stationary original parent and the four exact contact-zero packets (RU5). The algorithm is a bounded, source-linear map on the stated raw coefficient class and preserves the filtrations used next. In particular it preserves the source ideal (h, g); no separate contact-zero assertion for its individual coefficient summands is needed. The source substitution in each monic step uses the actual E, which is a bounded unit through $h = 0$.

The complete filtered leading model. Set

$$\Lambda = \frac{1}{t(1+s)}, \qquad \mu = \frac{1}{1+s}, \qquad T_4 = \mathsf{A}^2 + (u\mathsf{B})^2 + (k\mathsf{C})^2 + \mathsf{D}^2 - 4k^2\mathsf{BC}.$$

The model scalar is

$$\begin{aligned} G_{\mathrm{mod}} &= \Lambda\mathsf{x}^2 + T_4/2 - (1/t + 1/2)k^2\mathsf{x}^2 + (k^2 - u^2)\mathsf{Cx}/t \\ &= \left[\Lambda + \frac{u(u+k)}{t} - (1/t + 1/2)k^2\right]\mathsf{x}^2 + T_4/2 - \frac{u+k}{t}\mathsf{xz}. \end{aligned} \tag{RU18}$$

The second equality is the exact conductor identity $\mathsf{C}(u - k) = \mathsf{z} - u\mathsf{x}$. Use the first expression for the raw coefficient filtration, since it retains the even-u class. Its angular raw coefficients are

$$\begin{aligned} F_{BB} &= \Lambda + u^2/2 - (1/t + 1/2)k^2, \\ F_{BC} &= -2\Lambda - u^2/t + (3/t - 1)k^2, \\ F_{CC} &= \Lambda + u^2/t - 2k^2/t. \end{aligned} \tag{RU19}$$

At $h = g = 0$, exact source collection gives the following two coefficientwise filtrations:

	b a unit	b infinitesimal
variable weights	$(k, u, \omega, v) = (1, 1, 1, 2)$	$(\varepsilon, \jmath, u, \omega, v) = (1, 1, 2, 1, 3)$
$\mathrm{wt}(\mathsf{x})$	2	3
$\mathrm{wt}(\mathsf{y}, \mathsf{z}, \mathsf{h}, Q_1, Q_2)$	3	5
$f_A - 1/2,\ \varrho - 1/2$	≥ 2	≥ 2
$f_{BB} - F_{BB},\ f_{BC} - F_{BC},\ f_{CC} - F_{CC}$	≥ 4	≥ 6
$S_B - \mu,\ S_C + \mu$	≥ 2	≥ 4
(R_A, R_B, R_C)	$\geq (2, 3, 3)$	$\geq (3, 5, 5)$.

(RU20)

These statements concern the chosen raw coefficients separately; polynomial cancellation between raw products is not used to infer them. Consequently

	b a unit	b infinitesimal
$G_0 - G_{\mathrm{mod}}$	$O_{\mathrm{wt}}(8)$	$O_{\mathrm{wt}}(12)$
$L_M - \mu\mathsf{x}$	$O_{\mathrm{wt}}(4)$	$O_{\mathrm{wt}}(7)$
L_N	$O_{\mathrm{wt}}(5)$	$O_{\mathrm{wt}}(8)$.

(RU21)

Here each equality is modulo the source ideal (h, g).

To reproduce the collection, the necessary b-infinitesimal source jets at $h = g = 0$ are

$$\begin{aligned} C &= t/2 - (t+1)s/2 + (t+1)s^2/2 + O_{\deg(s,b)}(3), \\ A_h &= t(1 - s + s^2)/2 + O_{\deg(s,b)}(3), \\ \beta_D &= [1 - s + s^2 - (t+1)sb]/(2t) + O_{\deg(s,b)}(3), \\ n_1 &= (s - s^2 - tsb)/2 + O_{\deg(s,b)}(3). \end{aligned} \tag{RU22}$$

Also $c = (1 - k^2)/t + O_{\mathrm{wt}}(6)$, $E = t(1 + k^2)$ at $h = 0$, and $s = tj^2 + O_{\mathrm{wt}}(6)$. For the unit-$b$ filtration use $s = k^2/b$ and the respective coefficients of s: $-(t+1)/2, -t/2, -[1 + (t+1)b]/(2t), (1 - bt)/2$. The source cancellations (RU11)–(RU12) supply the first and last rows of (RU20). The checker verifies all needed source derivatives, each angular raw coefficient through weight two or four, the whole scalar through weight six or ten, and each complete M_B normal through weight three or five. Scalar weights are even in both charts. The M_B weights are even in the first and odd in the second, giving exactly the next weights in (RU21).

Apply the bounded conductor to the exact differences in (RU21). Their coefficientwise raw filtration makes every correction Gram coefficient have strictly positive weight. Terms in (h, g) lift boundedly and remain infinitesimal at arbitrary relative rates. The whole hard block differs infinitesimally from $\mathrm{diag}(1/2, (t+1)/2)$, since all its coefficients are bounded and u, v, h, g, s are infinitesimal.

On the negative-core contact chart take $u, v, \omega > 0$. The separated nonnegative monomials

$$a_i^2 = (\mathsf{A}, u\mathsf{B}, k\mathsf{C}, \mathsf{D}), \qquad Q_1 = a_1a_2 - a_3a_4, \quad Q_2 = a_1a_3 - a_2a_4$$

have the exact product relation yielding

$$T_4 = \tfrac{1}{2}\{(\mathsf{y} - \mathsf{z})^2 + \mathsf{h}^2 + \mathsf{y}^2 + (\mathsf{z} + \mathsf{h})^2\} + Q_1^2 + Q_2^2. \tag{RU23}$$

Its three-dimensional difference matrix has determinant $1/2 > 0$. Each radical separates into source and testing factors. The full eight-root residue on $(\mathsf{x}, \mathsf{y}, \mathsf{z}, \mathsf{h}, Q_1, Q_2, N, M_B)$ therefore consists of half this strict circuit, the N pivot $1/2$, and the complete coupled block

$$\begin{pmatrix} 1/t & 1/2 \\ 1/2 & (t+1)/2 \end{pmatrix}, \qquad \det = \frac{t+2}{4t} > 0. \tag{RU24}$$

The additional cross term in the second line of (RU18) has infinitesimal coefficient. WA10 now gives a finite separated SOS for the exact remainder. Restore all four actual squares (RU5). This proves the whole local contact chart in both b alternatives.

The ω-unit state chart. If u, v are infinitesimal, $Z = 1$, and ω is a positive unit, replace the retained centered n packet by $nD_uD_vX^2$ itself. The complete remainder is (RU7) with the two n_1, n_2 terms removed. Use the frame $(A_o, \mathsf{x}, \mathsf{D}, N, M_B)$, $A_o = \omega v$. The exact identities $\mathsf{A} = \varepsilon A_o$, $\mathsf{B} = A_o/\omega$, $\mathsf{C} = A_o/\omega - \mathsf{x}$ use only a pure-testing unit inverse. In particular the first scalar coefficient becomes $D_u^2D_v(p^2/2 + \mu_2)A_o^2$, and the first N coefficient becomes $D_u(p - 2\mu_1)A_o$; the printed ε denominators cancel completely. The full residue has the A_o, D, N pivots $1/2, 1/2, 1/2$ and the same block (RU24). The complete normal residues are $0, \mathsf{x}$. Hence this chart also has a strict finite separated Gram.

A remote cap lemma and complete bounded parents. Away from the contact chart there is no need for centered excesses. Retain the original nonnegative n, α packets and the positive excesses of m, A above chosen positive real caps m_c, A_c. If D is unbounded, retain its original excess above a real cap D_c too; otherwise leave D at its actual bounded value. The remainder is the full original parent with these three coefficients capped, and all its coefficients are bounded.

We use the elementary finite-dimensional fact that a real quadratic form positive definite on the common kernel of finitely many real linear rows becomes positive definite after sufficiently large positive real multiples of the squared rows are added. Choose coordinates on the common kernel and its orthogonal complement; the assertion follows by a Schur complement. Unbounded actual m, A, D, when used, exceed each chosen real cap. Thus every cap is an exact nonnegative packet subtraction in the original parent, and a strict residue gives WA10 for the entire bounded remainder. The real caps depend only on real testing residues, never on a source/testing rate comparison.

Finite remote v with nonzero residue. Write $U = u^2, V = v^2$ for real testing residues, with $V > 0$. On the common kernel $Y = M = 0$, $X = vW - uZ$, and the remaining two-state form is exactly

$$CD_vUW^2 + (D + 1/2)(uD_vW - vD_uZ)^2 + VD_vW^2/2 - D_uVZ^2/2. \tag{RU25}$$

Its determinant is

$$VD_uD_v\{DV/2 + U[CD - (D + 1/2)/2] + C(D + 1/2)U^2\}. \tag{RU26}$$

For $t_0 > 1$, insert $C = t/2, D = 1/[2(t - 1)]$; it equals $D_uD_v[V^2 + t^2U^2V]/[4(t - 1)] > 0$. Its W^2 coefficient is positive since $V > 0$. For $t_0 = 1$, $C = 1/2$ and one may choose a real D cap: the braces are $D(V + U^2)/2 + (U^2 - U)/4$, positive for sufficiently large D. The cap lemma then chooses m_c, A_c for a strict full state matrix.

Finite unit u with infinitesimal v, including zero. Use $N = -M = X + uZ - v(W + uY)$, $R_t = uW - Y$, and $V_Z = vZ$. After the original n, α packets and allowed positive excesses are retained, the entire remainder lies in the square of the four roots (N, R_t, Y, V_Z). Only the pure-testing unit u is inverted to write $W = (R_t + Y)/u$; no inverse of v is needed. Its exact residue is

$$(m_c + 1/2)N^2 + CR_t^2 + A_cD_uY^2 + (D + 1/2)(R_t - D_uV_Z)^2 - D_uYV_Z - D_uV_Z^2/2. \tag{RU27}$$

For $t_0 > 1$, its restriction to $Y = 0$ has determinant $t^2D_u(D_u - 1)/[4(t - 1)] > 0$ on (R_t, V_Z). Choose A_c sufficiently large. For $t_0 = 1$, the common kernel $Y = 0, R_t - D_uV_Z = 0$ has value $D_uu^2V_Z^2/2 > 0$; large real D_c, A_c give strictness. The normal pivot is positive. The full four-root support and the residue are checked directly; literal $v = 0$ is their polynomial value.

Both finite angles infinitesimal, and all state charts. The exact SD5 pure-core sign identity deals with the nonnegative-core branch and literal zero axes by original-event squares. On the negative branch use its standard gauges $u, v > 0$ and normalize the state by a maximal coordinate. If Z is infinitesimal, W is a unit and the exact BE4 selected four-root identity applies to (X, Y, uW, vW), with positive caps of m, A provided by (RU3). All other original events remain as nonnegative packets. If Z is a unit normalize $Z = 1$; the same sign identity gives bounded positive ω. Its infinitesimal and unit alternatives are exactly (RU18)–(RU24) and the subsequent ω-unit chart.

Every reciprocal chart and literal projective infinity. For unbounded u, put $p_u = 1/u$ and use the entire homogeneous polynomial p_u^2P. If bounded v has nonzero residue, the full capped four-state residue is

$$CD_vW^2 + A_cD_vY^2 + (D_c + 1/2)(W + vX)^2 + (m_c + 1/2)(vY - Z)^2 \succ 0. \tag{RU28}$$

Here D_c denotes either a positive real cap or the actual positive unit residue. The four square rows have common kernel zero because v is a unit. If instead v is infinitesimal, including zero, use the

exact BE $(\infty, 0)$ selected full frame $(p_u X + Z, W, Y, vX)$ with positive caps of C, A. Its strict residue is the sum of positive multiples of $(p_u X + Z)^2, (W + vX)^2, W^2, Y^2$; the other original packets are retained. This handles the zero-n degeneration at that corner.

For unbounded v and bounded u, put $q_v = 1/v$. The full capped residue of $q_v^2 P$ is

$$m_c(W + uY)^2 + C(uW - Y)^2 + (A_c + 1/2)D_u Y^2 + D_c(uX - Z)^2 + D_u X^2/2 \succ 0. \tag{RU29}$$

Its common square kernel is zero for every bounded real u, including zero. Finally, if both angles are unbounded, use $p_u^2 q_v^2 P$ and the full frame $(X, W, Y, P_Z = p_u Z, Q_Z = q_v Z)$. The exact residue is

$$CW^2 + A_c Y^2 + D_c(X - P_Z)^2 + X^2/2 + (m_c + 1/2)(Y - Q_Z)^2 \succ 0. \tag{RU30}$$

The two nontrivial block determinants are $D_c/2$ and $A_c(m_c + 1/2)$, both positive. The entire polynomial has this five-root support, with no inverse of p_u or q_v after homogenization. All three formulas are checked from the original parent. Their literal reciprocal zeros are polynomial values; restoring a nonzero homogeneous factor uses a pure-testing square.

The charts exhaust independent testing pairs and states. Apply each real identity to real and imaginary parts of a complex state, restore every retained actual packet and separated clearing factor, and use the exact sign gauges for both angle signs. This completes the stated all-tests theorem without a restriction on any opening ratio.

Dependencies and the scope of the checks. The exact dependency manifest is `dependencies_r7_unbounded_a.json`. It distinguishes the actual source definitions, full-parent identity, algebraic conductor, sign/state reduction, and strict finite Gram lemma from the source theorems surrounding those identities. The companion `check_r7_unbounded_a.py` reconstructs source and whole parent expressions, checks complete scalar and normal allocations, the two coefficientwise filtrations, and the strict local and remote matrices. The separate-review checker `review_check_r7_remote.py` also reconstructs the complete four-root and five-root supports without dividing by a zero testing coordinate. These are exact symbolic corroborations; the monic coefficient-class argument and exhaustive joins above supply the mathematical proof. No formal verification or external peer review is claimed.

Chapter 182

A fresh angular budget and completion of the retained C14.1b residual

Component p4e5ba027f187
Source: NEW/c1/c141b_residual_rederived.tex
SHA256: 4e5ba027f1879415684e3c497a56aa194f0baea8216cd808d701c0b66eafa0c2
Inventory use: AR1–AR21 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

182.1 A fresh angular budget and completion of the retained C14.1b residual

Scope and provenance. This is a new proof, not a restoration of the missing late-R21 source. Keep the original physical stationary C1 source, its strict gaps, nonnegative event coefficients, original infinitesimal total scale and every earlier-admission complement in the surviving C14.1b statement. Normalize the original positive coefficient to $d = 1$ as in P1–P4. Source parameters lie in a real closed extension K_c and testing angles and states in an independent extension K_t of $\mathbb{R}$. All conclusions are finite sums of squares in $\mathcal{T} = K_c \otimes_{\mathbb{R}} K_t$. No order or general inverse of a mixed tensor is used.

The exact target is the complete reciprocal-Bob polynomial P4/SP6 in the surviving `inherited_r19/parent_reference.tex`. Use its notation, and set $k = \sqrt{x}$, $q = \sqrt{z}$, $z = 1/L$, $\sigma = 1/\eta$, $b = Lr$, $Q = 1 - b$, $\theta = 1 - h$. The source part of the retained residual GA9 is

$$x, z, \sigma > 0 \text{ infinitesimal}, \quad \operatorname{res} h \geq \tfrac{1}{2}, \quad k/\sigma \text{ infinitesimal}, \quad b_0 \text{ infinitesimal}, \quad d_0 \sim 1. \tag{AR1}$$

Its real testing part has $u, w > 0$ infinitesimal, $WZ > 0$, W/Z and u/w unbounded, and $uW/(wZ)$ unbounded. These intersections are retained. The proof below in fact needs only $u, w, W, Z > 0$ with u, w infinitesimal after the inherited sign gauges; it assumes no upper or lower bound on any state ratio.

Theorem 182.1 (Fresh completion of the retained residual)**.** *On AR1 and the exact retained testing/source intersections just stated, the complete original parent is a finite tensor sum of squares. Together with the surviving SF, BC, US and GA admissions and their recorded literal, sign and projective exits, this closes the entire original scoped C14.1b domain. It does not by itself close C14.4 or the global C1 entry and union.*

182.1.1 Source coefficients and the original scalar packet

The source estimates SF1–SF5 and US2–US3 were proved for z infinitesimal. On AR1 they imply

$$\begin{gathered} L_0 := \lambda\sigma \sim 2h \text{ is a positive bounded unit}, \qquad \lambda^{-1} \in \operatorname{Inf}_c, \\ \rho = \lambda x/\sigma = L_0(k/\sigma)^2 \in \operatorname{Inf}_c, \qquad \lambda x \in \operatorname{Inf}_c, \qquad d_G := 4D/\sigma^2 \sim 2. \end{gathered} \tag{AR2}$$

Here h is a positive unit even when $\theta = 1 - h$ is infinitesimal. In particular $\lambda > 0$ and d_G is a strict original Gamma pivot. The exact sphere identity P2 proves these bounds without an inverse of b_0 or d_0.

For clarity, the other source bounds used below are

$$\begin{gathered} P, J, M, B, \mathcal{H} \sim 1, \quad p, l, \kappa \sim \tfrac{1}{2}, \quad C = O(z), \quad D = O(\sigma^2), \\ e = xh/\sigma \in \mathrm{Inf}_c, \quad m = \kappa x z e \in \mathrm{Inf}_c, \quad \beta\theta \in \mathcal{O}_c, \quad \nu, K_A, \chi, R_x, A_y \in \mathcal{O}_c, \quad B_x = 8\kappa e \in \mathrm{Inf}_c\,. \end{gathered} \tag{AR3}$$

These are SF1–SF5, the P1 formulas and the SP9–SP10 polynomial cancellations. In particular $K_A = 4\kappa h/\sigma - \lambda$ is bounded although its residue $2 - 4\operatorname{res} h$ may be zero or negative. Also

$$R_\gamma = -4\{2(C + D)(r - 1) + 2p + r - 1 + 2pu^2\} \tag{AR4}$$

is a bounded separated coefficient. None of AR2–AR4 contains u/w, a state ratio, or a comparison of a source and a testing rate. The only possible source poles in $a = Az = p/b$ and $\ell = \alpha z = l/Q$ remain in positive packets below. The possibly unbounded β itself also remains in its original nonnegative packet; no upper bound on β is asserted.

Keep the exact roots

$$N_x = X + uhZ, \quad N_y = (Y - bwZ)/q, \quad \Gamma = \sigma Z - uwW, \quad R = wZ/q - uqW, \quad A_\theta = q(u^2 - x)W. \tag{AR5}$$

The whole angular packet of SP6 is

$$K_A A_\theta^2 + \chi u A_\theta R + \mathscr{S}, \qquad \mathscr{S} = \lambda\sigma\{uZ - (x/\sigma)wW\}^2 + \rho(u^2 - x)(w^2 - \sigma z)W^2. \tag{AR6}$$

This is the exact P8 transformation, not a restriction of the state. Define the four positive separated monomials

$$B_1 = \sqrt{\lambda\sigma}\,uZ, \quad B_2 = \sqrt{\rho}\,uwW, \quad B_3 = \sqrt{\lambda}\,xqW, \quad B_4 = \sqrt{\lambda x}\,wZ/q, \qquad G_4 = (B_1B_2B_3B_4)^{1/4}. \tag{AR7}$$

Every fractional power means the product of the two pure-field powers. With $d_i = B_i - G_4$, the complete circuit is

$$\mathscr{C} := \mathscr{S} + \lambda x R^2 = \sum_{i=1}^{4} B_i^2 - 4G_4^2 = \sum_{i=1}^{4} d_i^2 + \mathcal{E}_4, \tag{AR8}$$

where the explicit surviving US10 remainder is

$$\mathcal{E}_4 = 2G_4\left\{(\sqrt{B_1} - \sqrt{B_2})^2 + (\sqrt{B_3} - \sqrt{B_4})^2 + 2\big((B_1B_2)^{1/4} - (B_3B_4)^{1/4}\big)^2\right\}. \tag{AR9}$$

Its nonnegative monomial prefactors have separated square roots.

182.1.2 An eight-square budget for the angular row

The following elementary identity is the new step. Let U, V, Z_* be positive separated monomials, let $a_* > 0$ be pure source, and set

$$\begin{gathered} \Psi = 1 + U^2V^2 + U^2Z_*^2 + V^2Z_*^2 - 4UVZ_*, \\ T = U^2 - 1, \qquad Q_* = VZ_* - U, \qquad R_* = Z_* - UV, \qquad G_* = U(V - Z_*). \end{gathered} \tag{AR10}$$

There are two exact polynomial identities

$$\Psi = G_*^2 + (1 - VZ_*)^2 + 2VZ_*(U - 1)^2, \qquad \Psi = T^2 + Q_*^2 + UTQ_* - UR_*^2 - (U + 1)G_*R_*. \tag{AR11}$$

Consequently

$$\boxed{\begin{aligned} &[2 + a_*(U + 1)]\Psi + U^2Q_*^2 + \left[2U + \frac{U + 1}{a_*}\right]R_*^2 - T^2 \\ &\quad = (T + UQ_*)^2 + 2Q_*^2 + (U + 1)\,(\sqrt{a_*}G_* - R_*/\sqrt{a_*})^2 \\ &\qquad + a_*(U + 1)(1 - VZ_*)^2 + 2a_*(U + 1)VZ_*(U - 1)^2. \end{aligned}} \tag{AR12}$$

The right side consists of eight legal squares: expand each factor $U+1$ into its two summands before taking roots. Only radicals of positive separated monomials occur. In particular neither a square root nor an inverse of $1+U$ is used. The identity imposes no bound on U, V, Z_* or on a_*^{-1}.

Apply this with

$$U = \frac{u}{k}, \qquad V = \frac{w}{\sqrt{\sigma}q}, \qquad Z_* = \frac{\sqrt{\sigma}}{kq}\frac{Z}{W}, \qquad a_* = \lambda k, \qquad M_* = k^2 qW. \tag{AR13}$$

Each quotient is taken within a pure field before tensoring. The only state inverse is the nonzero pure-testing W^{-1}. The other inverses are positive pure-source factors. These expressions are allowed to be unbounded inside a square root row. Directly,

$$\mathscr{C} = \lambda M_*^2 \Psi, \qquad A_\theta = M_* T, \qquad R = kqWQ_*, \qquad \Gamma = k\sqrt{\sigma}qWR_*. \tag{AR14}$$

Multiplying AR12 by the square M_*^2 therefore proves

$$\boxed{\Delta_A := \varepsilon_C \mathscr{C} + u^2 R^2 + \varepsilon_G \Gamma^2 - A_\theta^2 \ \in\ \Sigma\mathcal{T}^2,} \tag{AR15}$$

with the exact coefficients

$$\varepsilon_C = \frac{2}{\lambda} + u + k, \qquad \varepsilon_G = 2\frac{k}{\sigma}u + \frac{u+k}{\lambda\sigma}. \tag{AR16}$$

Both are finite sums of positive bounded separated factors with zero real residue. This follows from AR2, bounded k/σ, and the independent facts $u \in \mathrm{Inf}_t$ and $k, \lambda^{-1} \in \mathrm{Inf}_c$. It uses no bound on qu/w, $uW/(wZ)$, or any normalized variable in AR13. The identity also gives a direct expression for each of its eight roots: multiply the eight expanded roots on the right of AR12 by M_*. Thus AR15 is a finite certificate, not a pointwise inequality imported from an ordered compositum.

182.1.3 Paying the signed angular term exactly once

Choose a fixed positive real c_* such that $\mathrm{res}(K_A + c_*) > 0$, possible because K_A is bounded. The exact allocation is

$$\begin{aligned} K_A A_\theta^2 + \mathscr{S} = (K_A + c_*)A_\theta^2 + (1 - c_*\varepsilon_C)\mathscr{C} - c_*\varepsilon_G\Gamma^2 \\ - (\lambda x + c_* u^2)R^2 + c_*\Delta_A. \end{aligned} \tag{AR17}$$

AR17 uses AR8 and AR15 once each. The coefficient of the whole circuit has residue one; its four d_i rows remain strict. The three AR9 rows with this same coefficient are retained separately. The elementary positive-residue lemma makes each of these bounded coefficients a tensor SOS; multiplying its finite square decomposition by the corresponding row preserves the finite SOS property. There is no mixed square root of $1 - c_*\varepsilon_C$.

182.1.4 The two complete state normals and all original hard terms

Use the exact CT7–CT7a radial extraction. Its constant base is

$$[4a + 2z(2C+1)](N_y + bR)^2 + 4\ell(N_y - QR)^2. \tag{AR18}$$

Since $p, l \sim 1/2$ and $0 < b, Q < 1$, both coefficients exceed one. Keep their positive pure-source excesses and retain $(N_y + bR)^2 + (N_y - QR)^2$ in the principal frame. Its bounded two-by-two Gram has determinant $(b+Q)^2 = 1$ and positive definite residue, including $b = 0$ or $Q = 0$ in residue. This is a cap of whole positive squares, not a cap of unrelated matrix entries. Retain exactly the nonconstant positive a, ℓ, C packets specified in CT7, before using its remaining coefficients.

The three exact remainders E_{yy}, E_{RR}, E_{yR} in CT7a are bounded infinitesimal: every source factor in their braces is bounded by AR3, $C = O(z)$, $D = O(\sigma^2)$, and they have the printed factor z together with u^2, w^2, σ or D. No estimate involving a state coordinate or u/w is present.

The two entire Gamma normal terms of SP6 are

$$-4\nu u N_x \Gamma + qwR_\gamma(N_y + bR)\Gamma. \tag{AR19}$$

They are bounded infinitesimal entries on the original (N_x, N_y, R, Γ) frame, directly by AR3–AR4 and u, q, w infinitesimal. In this source sector the Gamma diagonal is $d_G\Gamma^2$ with $d_G \sim 2$, so no sphere-coordinate inverse, Gamma conductor or weak Gamma pivot is required. The five other complete cross terms are exactly

$$\begin{aligned} &-4uqw(2m + 2D + 1)N_xN_y + uqwR_xN_xR \\ &\quad + qwB_xN_xA_\theta + uA_yN_yA_\theta + \chi uA_\theta R. \end{aligned} \tag{AR20}$$

Every displayed coefficient is infinitesimal and bounded by AR3. The entire N_yR term has already been included in AR18 and CT7a. Thus both unrestricted state normals and every remaining mixed term of SP6 have been allocated.

Use exactly $2N_x^2$ in the principal frame and retain exactly $[2w^2 + 4mw^2]N_x^2$ as its positive excess. After AR17 the principal frame is

$$(N_x, N_y, R, \Gamma, A_\theta, d_1, d_2, d_3, d_4). \tag{AR21}$$

Its real residue is strictly positive definite: the N_x pivot is two, the radial block is AR18's clipped block, the Gamma pivot is two, the angular pivot is $\operatorname{res}(K_A + c_*) > 0$, and the four circuit pivots are one. All other entries are bounded infinitesimals, including the AR17 Gamma/radial costs. The separated bounded-Gram lemma therefore proves finite SOS.

Finally restore, each exactly once, $4\beta(1 + w^2)N_x^2$, $2(2D + 1)(uX_e)^2$, $[2w^2 + 4mw^2]N_x^2$, $4bpzu^2\Gamma^2$, all CT7 positive source and testing excesses, the AR9 remainder with its positive-residue coefficient, and $c_*\Delta_A$. This restores precisely P4/SP6. The roots need not be bounded; only the Gram entries have been bounded.

182.1.5 Exact join and literal boundaries

The surviving SF theorem covers $\operatorname{res} h < 1/2$ in C14.1b. US covers its unit-h, unit-k/σ source for all states and tests; US2 bounds k/σ throughout the unit-h source. BC and GA cover their stated bounded-ratio testing domains. Their residual is exactly contained in AR1 with the testing predicates printed above, now covered by AR17–AR21. The inherited bounded-state, nonzero-unit-Alice, remote, literal-axis and sign exits remain unchanged. Their precise dependency index is the surviving `proofs/c1_unbounded/DEPENDENCIES.txt`, together with the introductions and concluding joins of SF, US and GA. This is a finite union on the original scoped C14.1b source; it does not erase any earlier-admission complement.

The residual itself has $D > 0$ by AR2. Literal $D = 0$ remains covered by the US sphere endpoint, which divides by neither D nor a sphere coordinate. Literal $\beta = 0$ is incompatible with $H = h/\sigma$ unbounded here: P2 gives $H = H_0/B$ on that face, a bounded quantity. Hence no nonempty C14.1b face has been dropped. Arbitrarily small positive b, Q, θ, D or other original openings are kept wherever compatible with the printed source sector. For complex states apply the real routing independently to their real and imaginary parts; each part satisfies an earlier exit or the new residual proof. Restore only the original positive pure-source and pure-testing normalizations.

Verification boundary. The companion exact checker verifies the elementary identities AR11–AR12, all four scaling identities AR14, the restored budget AR15–AR16, its eight separated squares, and the once-paid AR17 allocation. The complete P4/SP6 and CT7a identities are surviving algebraic inputs. Their source bounds, the full-normal allocation and the quantified join are the analytical argument above; an assertion count does not replace these arguments. Independent review is requested before this new theorem is recorded as an admitted replacement.

Chapter 183

The complete C14.4 source union and the original C14.5 join

Component `pfd0dd7bb10cf`
Source: `NEW/c1/c144_all_sources_and_c145_join_rederived.tex`
SHA256: `fd0dd7bb10cf01a3bdf1ef7994042fcaf6560a26e0f5595b590619f7ec1a3416`
Inventory use: JC1–JC7 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

183.1 The complete C14.4 source union and the original C14.5 join

Input and scope convention. Let K_c, K_t be arbitrarily chosen, separately labelled real closed extensions of $\mathbb{R}$; no algebraic-disjointness assumption is made. Use the original complete parent and canonical native C1 contact, with $d > 0$, $\lambda = n = 0$, $m, C, A, \alpha > 0$, $D, \beta \geq 0$ and the original allowed literal zero convention. Preserve physicality, stationary contact, all strict native gaps, the positive infinitesimal total scale, and every common entry predicate of the historical C1.4 row. In particular $x = k^2 > 0$ is infinitesimal and $\zeta > 0$ is noninfinitesimal. No source contact is identified with an independent testing point. Normalize $d = 1$ only by the permitted positive pure-source factor.

The exact historical row is the C1.4 row of `proofs/routing/routing_audit.tex`, sections "One exact input, before optional exclusions", "Exhaustive dispatch into the historical thirteen rows", and "C1.4: exhaustive C14 subdivision, with missing exits restored".

The companion `ROUTING_LEDGER.json` records the common input Q and each literal source predicate. This section completes that entered row; it does not replace the separate proof that an arbitrary physical support reaches the row or an earlier terminal.

At the C14 subdivision itself, the earlier exits are not left implicit: they are the three (L, η) admissions (unbounded, unit), (unit, infinitesimal), (unit, unit), the JR/JB bounded-M_β exit, and RO, JO, HO at unit η, ζ. Their exact predicates and proof bodies are displayed below. Any additional upstream prior-complement predicate is carried as the same conjunct in every alternative. No inequality used here is inferred merely from being outside an earlier admission.

183.1.1 C14.4: an exhaustive source-only partition

In the original C14.4 part, $L = x\zeta$ and $\sigma = 1/\eta$ are positive infinitesimals. Set

$$S = \sigma + L, \qquad \kappa = x/S, \qquad \nu = \sigma/S, \qquad \ell = L/S, \qquad \nu + \ell = 1, \quad \zeta = \ell/\kappa. \tag{JC1}$$

The exact source is FM2–FM3, obtained algebraically from NC1–NC4; the relevant gap and sphere estimates are NC5 and ES1–ES10/SE1–SE9. These are actual-source statements, including $D = 0$ or $\beta = 0$ where allowed. The NC5 bound $E = 1 - \kappa h/\nu \geq 1/18$ follows from native signs and $D \geq 0$ with $k \leq 1$, not from a prior-exit exclusion. Thus its use introduces no additional source predicate.

The parameter κ is bounded. In fact with $M = 1 + xr - \kappa(1-h)/\ell > 0$ and $Q_s = 1 - \ell Sr \sim 1$, the exact identity is

$$\kappa Q_s + \ell M + \nu E = 1, \qquad 0 < \kappa Q_s < 1 - \nu/18. \tag{JC2}$$

Also $0 < \nu, \ell < 1$. Each of these positive bounded source quantities is either infinitesimal or a positive bounded unit. The two fractions cannot both be infinitesimal. If κ is a unit, then ℓ is a unit because $\ell = \kappa\zeta$ and ζ is noninfinitesimal; consequently ζ is also a unit. In particular the case "unit κ, infinitesimal ℓ" is empty, rather than omitted.

The following five alternatives are exhaustive on JC1:

κ	ν	ℓ	complete full-parent theorem
unit	unit	unit	FU (with UKR for its matched chart)
unit	infinitesimal	unit	NU
infinitesimal	unit	unit	FM
infinitesimal	unit	infinitesimal	EL
infinitesimal	infinitesimal	unit	SN, BN, UN as in JC4.

(JC3)

Every entry is an actual theorem on all independent tests and all states, not only its scalar contact graph. FU and NU retain the possibly vanishing M and the whole divergent m packet. FM, SN, BN, UN and EL use only the valid small-e source, since SE4 proves $e \in \mathrm{Inf}$ exactly when $\kappa \in \mathrm{Inf}$.

To spell out the last line, use only in this line

$$q_\nu = \nu/\ell, \qquad \tau = h/q_\nu, \qquad n_s = \beta/q_\nu^2. \tag{JC4}$$

If n_s is bounded, SN applies and its source identity gives $\tau \sim 1$. If n_s is unbounded and τ is bounded, BN applies. If τ is unbounded, UN applies, including $\theta = 1 - h$ infinitesimal. These alternatives are exhaustive; BN in fact proves the entire bounded-τ domain and includes SN as its bounded-n_s part. There is no assumption that h/q_ν or β/q_ν^2 remains a unit. The EL proof uses the reciprocal source ratio $q_\ell = \ell/\nu$ in its own formulas; no formula is transferred between these two local definitions.

The literal faces are preserved. On FM and EL, D/S has positive unit residue, so $D = 0$ has empty intersection. On SN, $D/(Sq_\nu^2) \sim 1/2$, with the same conclusion. FU, NU, BN and UN retain their actual sphere alternatives and include $D = 0$ where it occurs. Literal $\beta = 0$ belongs to SN in the double-infinitesimal endpoint, since then $n_s = 0$ and the exact boundary relation makes τ bounded. It has empty intersection with UN's unbounded-τ source; all other relevant theorems retain it explicitly. No positive opening of $r, 1-r, \theta, M$ or either sphere coordinate has been removed.

Each component proof includes all testing charts. In the small-e sources, the actual SE10–SE12 homogeneous parent covers Bob residues unequal to $\bar{\ell}$; CE covers noninfinitesimal Alice tests. The remaining charts are proved by the displayed complete normal allocations. FU and NU supply their own raw finite/reciprocal testing joins on the unit-κ source. Thus JC3–JC4 cover every original C14.4 source/test/state tuple, including literal axes, projective infinities, all angle signs and independent real and imaginary state components. This proves the whole original C14.4 branch.

183.1.2 C14.1 and the retained complete C14.2/C14.3 branches

For L, η unbounded set $\sigma = 1/\eta$ and $H = h/\sigma$. The actual large-L source formulas are those of P1/SP6 and SF1–SF5. If H is bounded, the surviving `c141a_closure_and_c141b_partial.tex` theorem proves all tests and states. Its proof uses JR/JB when $M_\beta = \beta(1-h)/\sigma^2$ is bounded, including zero, and its own full state-free normal conductor for the remaining unbounded-M_β sector. Thus the original C14.1a branch is entirely covered without reinstating a positive-β premise.

For unbounded H, the surviving SF theorem first removes $\operatorname{res} h < 1/2$. On the other source part, US bounds k/σ and covers its positive-unit case, equivalently the unit beta-sphere coordinate. The bounded-state, bounded testing ratio, unit-Alice, literal and sign exits are the actual BS, BC, US, GA and inherited CE proofs indexed in `proofs/c1_unbounded/DEPENDENCIES.txt`. Their retained complement is exactly contained in

$$\operatorname{res} h \geq 1/2, \quad k/\sigma \in \mathrm{Inf}, \quad b_0 \in \mathrm{Inf}, \quad d_0 \sim 1, \quad u, w > 0 \text{ infinitesimal}, \tag{JC5}$$

with the original positive-state and unbounded-ratio predicates. The fresh AR1–AR21 theorem covers this complement using its restored angular SOS budget. Its coefficients require no bound on qu/w, u/w, W/Z or $uW/(wZ)$, and all SP6 first-normal terms are allocated. Applying the real routing to each state component independently closes the whole original C14.1b branch. Its literal $D = 0$ source uses the US sphere endpoint; its $\beta = 0$ source is empty when H is unbounded by the exact sphere boundary equation. These statements keep the original source intersection rather than asserting a missing face is positive.

The complete C14.2 proof remains the actual R19 `c142_all_tests_join.tex`, with its bounded-H, sphere, toric, state-ratio and LA/LB leaves. It covers the original L-unit, η-unbounded domain, both literal coefficient faces, all openings and all independent tests/states. The actual R18 `c143ab_all_tests_join.tex`, together with its completed j conductor and source/remote leaves, covers C14.3a and C14.3b. These complete proof bodies are present; no missing late-R21 theorem statement is substituted for them.

183.1.3 The original C1.4 source is exhausted

Put $L = x\zeta$. The exact NI4–NI6 source implications and the explicit C14 subdivision in the current routing proof give the following elementary decision tree.

If L is unbounded, $Q = 1 - Lr > 0$ implies r infinitesimal. The native identity $1 - r = a_{\mathrm{nat}}\eta$ with $0 < a_{\mathrm{nat}} < 2$ therefore excludes infinitesimal η. Unit η is the complete `advances/c1_large_l_eta_unit/large_l_unit.tex` theorem. Unbounded η is split by bounded or unbounded H, just covered by C14.1a/JR/JB and C14.1b.

If L is a positive bounded unit, the three possible types of η give respectively the complete `advances/c1_lunit/lunit_eta_small.tex` theorem, `advances/c1_lunit_eta_unit/eta_unit.tex` theorem, and C14.2. These statements retain the actual D, β hypotheses and their full remote joins.

If L is infinitesimal, unbounded η is the now complete C14.4 branch. For bounded η, first split ζ: unbounded ζ is C14.3a; unit ζ with infinitesimal η is C14.3b. The only remaining source has both η, ζ positive bounded units. Its exact positive gap is

$$j = \eta(1 + xh) - 1 + r = \eta J > 0, \qquad j_0 = \eta + r - 1, \qquad j - j_0 = \eta x h \in \mathrm{Inf}\,. \tag{JC6}$$

The gap j is bounded. If it is a unit, j_0 is a positive unit and RO applies. If j is infinitesimal, then $\eta + r \sim 1$ and $1 - r$ is a unit; unit r gives JO, infinitesimal r gives HO. Their actual bodies are `advances/revision13_c1_small_l/ r_opening_closure.tex`, `j_opening_closure.tex`, and `simultaneous_r_j_closure.tex`. They retain actual $j > 0$, even when $j_0 = 0$ or slightly negative, and compare neither j with x nor a source quantity with a testing quantity. RO contains the earlier IC1 regular-gap admission, so that historical fourth C1.4 subdomain is restored as part of a fully stated theorem.

These alternatives exhaust every positive L, η, ζ permitted in the entered C1.4 row. More formally, let $\mathcal{C}$ denote that exact common row predicate and let E_{14} be the disjunction of the seven explicitly named prior exits above. Let R_i be the six original residual predicates after their stated exclusions. The unchanged finite source argument proves

$$\mathcal{C} \implies E_{14} \ \vee \ R_{14.1a} \vee R_{14.1b} \vee R_{14.2} \vee R_{14.3a} \vee R_{14.3b} \vee R_{14.4}. \tag{JC7}$$

Each alternative now has an actual complete full-parent proof. Intersecting all terms with any fixed earlier entry predicate preserves this implication; the predicate is not dropped or used to manufacture a source bound. Conversely, every prior C14 exit removed from an R_i is restored by its displayed proof. This is the required quantified C14.5 union, including all source openings, allowed literal coefficient zeros, projective endpoints and exact sign gauges. Multiplying by the original positive pure-source normalization and positive pure-testing homogenization factors preserves finite separated SOS.

Checklist consequence and boundary of this result. C14.1, C14.4 and C14.5 are complete on their unchanged original criteria. C14.2 and C14.3 retain their existing complete status. Consequently U1.B1, the whole original entered C1.4 row, is complete. This join does not by itself prove a global physical-source entry, the independent weak-C2 row, the finite-lift bridge, or an evaluated affine LMI. It adds no uniform square-count hypothesis. The accompanying dependency receipt binds the actual old and new proof bodies. Focused symbolic source identities and finite decision-tree checks corroborate the explicit analytical union; neither historical completion claims nor assertion counts are mathematical premises.

Chapter 184

The entire bounded-h/q part of the doubly small endpoint

Component `p80b0c5b9879f`
Source: `NEW/c1/c144_bounded_h_ratio_rederived.tex`
SHA256: `80b0c5b9879f6356652220d5b19b3db274430c9418fe50363919f55a12285c6d`
Inventory use: BN complete theorem | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

184.1 The entire bounded-h/q part of the doubly small endpoint

Source scope and result. Retain the original entered physical C14.4 source, all strict gaps and all original prior-complement intersections. Assume

$$S = \sigma + L,\ \kappa = x/S,\ \nu = \sigma/S \in \mathrm{Inf}_{>0}, \qquad \ell = 1-\nu, \quad q = \nu/\ell, \quad \tau = h/q \text{ is bounded.} \tag{BN1}$$

The complete original parent is a finite separated tensor SOS for every test and every real or complex state on BN1. The quantity β/q^2 is unrestricted, including zero and unbounded values. The proof below is a fresh circuit completion of SN, not an import of a missing historical endpoint proof.

Exact source estimates. Use the definitions of SN2, except that $n_s = \beta/q^2$ need not be bounded. The exact sphere SN3 remains valid. Put $b = \beta/q$ and $z = \kappa\beta/q^2$. Then

$$\begin{gathered} h = q\tau \in \mathrm{Inf}, \quad \theta \sim 1, \quad e = \kappa\tau/\ell \in \mathrm{Inf}, \quad M \sim 1, \\ b = \frac{\tau(B/S) - \ell H_0}{2\theta(P/S)}, \qquad \mu\ell^2 = \frac{\tau(1+x)(1+\sigma r)}{2M(P/S)}, \\ J d_s + \theta z = \frac{\ell^2 \mathcal{H}}{2(P/S)} \sim \tfrac{1}{2}. \end{gathered} \tag{BN2}$$

In particular b, z, d_s, μ, τ are bounded, τ and μ have positive real lower bounds, and at least one of d_s, z is a positive bounded unit. The exact inequality $\beta \geq 0$ supplies the lower bound on τ. Also $c_0, a_0, a_1 \sim 1/2$, D, α, m, β are infinitesimal. Although the two terms below need not have small difference before subtraction, their displayed bounded formulas in BN2 give

$$A_{00} := \mu\ell^2 - b \sim \tfrac{1}{2}. \tag{BN3}$$

This argument takes residues only of bounded source coefficients and never uses an estimate divided by q. It allows arbitrary relative rates of S, κ, q.

A retained circuit and two actual positive rows. Use the full reciprocal parent SN5, the two unrestricted normals SN6, and the shifted exact scalar dictionary SN9. Let $\zeta = \ell/\kappa$, $a = u^2, c = w^2$, $x = \kappa S$, and assume $u, w \ge 0$ by the original sign gauges. Both may be zero. For every state define

$$\begin{aligned} A_1 &= \sqrt{\beta}\, uZ, & A_2 &= \sqrt{\beta}\, W/(\zeta\sqrt{q}),\\ A_3 &= \sqrt{\beta}\sqrt{x/q}\, wZ, & A_4 &= \sqrt{\beta}\, uwW/(\zeta q\sqrt{x}), \end{aligned} \tag{BN4}$$

$$\Psi = \sum_{i=1}^{4} A_i^2 - \frac{4\beta}{\zeta q} uwWZ, \qquad E_1 = A_1 - A_2, \quad E_2 = A_3 - A_4. \tag{BN5}$$

Every monomial and its radical is a product of a pure-source and a pure-testing element. This remains true if x/q is unbounded. The exact inequality meant here is a separated SOS certificate:

$$\Psi - \tfrac{1}{2}(E_1^2 + E_2^2) \in \Sigma T^2. \tag{BN6}$$

Indeed, if $WZ \ge 0$, then $A_1A_2, A_3A_4 \ge 0$ and

$$\Psi = E_1^2 + E_2^2 + 2(\sqrt{A_1A_2} - \sqrt{A_3A_4})^2.$$

The radicals displayed here split into pure-source and pure-testing radicals. If $WZ < 0$, use instead

$$\Psi - \tfrac{1}{2}(E_1^2 + E_2^2) = \tfrac{1}{2}(A_1 + A_2)^2 + \tfrac{1}{2}(A_3 + A_4)^2 + \frac{4\beta}{\zeta q}|uwWZ|.$$

The last positive separated monomial is a square. Thus BN6 is valid for all states, including $WZ = 0$. At the actual source contact every A_i agrees, so Ψ and its first contact derivatives vanish.

The exact scalar after circuit subtraction. Write $N_0 = W/\zeta - uwZ = N/\ell$, where N is SN7, and keep G, B, B_u, R from SN7. Thus

$$R = \frac{\sqrt{S}}{\zeta} G - uN_0. \tag{BN7}$$

Subtract the entire circuit BN5 from the exact scalar graph in SN9. Define

$$\begin{gathered} A = \zeta^2 A_N - b, \qquad A'_P = A_P - zc/\ell^2,\\ K' = K - 2\beta/(\zeta q), \quad J' = K' - \ell A'_P, \quad L' = J' - A/\zeta,\\ H' = F_Z - \beta a - \beta xc/q - \ell^2 ScA'_P + ac(A - 2\zeta J'). \end{gathered} \tag{BN8}$$

Then $F_0 - \Psi = AN_0^2 + A'_P G^2 - 2\zeta uwZL'N_0 + H'Z^2$. All coefficients of A and A'_P are bounded:

$$A = \mu\ell^2(1 + ra + ec)^2 + (\beta + \tfrac{1}{2})\tau^2 c(1 + c) - b, \qquad \overline{A} = \overline{A'_P} = \tfrac{1}{2}. \tag{BN9}$$

Here and below the residue is taken at infinitesimal u, w.

The circuit subtraction has especially simple exact effects on the SN10 quotient dictionary:

$$\begin{gathered} L'_a = L_a, \qquad L'_c = L_c + \kappa n_s/\ell, \qquad H'_{20} = H_{20}, \quad H'_{02} = H_{02}, \quad H'_{11} = H_{11} - 2n_s,\\ H' = S\delta^2(d_s + cH_{02}) + c(a - x)\delta H'_{11} + c(a - x)^2 H_{20}. \end{gathered} \tag{BN10}$$

In particular no coefficient n_s remains on its own in the final conductors. With C_R and L_a from SN11–SN13 put

$$C'_G = \kappa(H_{11} - 2n_s)/\ell, \qquad C'_N = -H_{11} - 2\ell L_c/\kappa. \tag{BN11}$$

These are bounded. The cancellation for the second coefficient is explicit:

$$C'_N = -Sa_0\ell^2 r + (\beta + m + \tfrac{1}{2})\tau^2(c + 1/\ell) + 2\ell\kappa\mu\tau - m\tau^2. \tag{BN12}$$

For C'_G use the bounded formula SN13 and $\kappa n_s = z$. The exact full scalar is consequently

$$\begin{aligned} F_0 = \Psi &+ AN_0^2 + A'_P G^2 + d_s B^2 + H_{02}(wB)^2 + C_R R^2 - 2\sqrt{S}\, L_a GR\\ &+ C'_G G(wB) + C'_N N_0(wB_u). \end{aligned} \tag{BN13}$$

This is an identity in unrestricted state coordinates, not a restriction to a contact graph in the full parent.

Free helper rows across the whole source sphere. First reserve the positive rows E_1, E_2 from BN6. If d_s is a unit, wB has infinitesimal cost on $\sqrt{d_s}B$. Otherwise z is a positive bounded unit. The exact identity

$$E_2 = -\sqrt{z}\,\{(w/\ell)G + \sqrt{S}\,w(w - \sqrt{q})Z\}$$

then gives

$$wB = -\frac{w + \sqrt{q}}{\sqrt{z}}E_2 - \frac{w(w + \sqrt{q})}{\ell}G. \tag{BN14}$$

Every coefficient in BN14 is infinitesimal, so any fixed real pivot on wB is free at infinitesimal Gram cost. Only the chosen pure-source unit $\sqrt{z}$ is inverted. Literal $\beta = 0$ uses the first branch.

After this pivot has been purchased, the normalized SN8 identity is

$$\delta^2 N_0^2 + \frac{2\kappa\delta}{\ell}G(wB) + 2\kappa(wB)^2 - (wB_u)^2 = (\delta W/\zeta)^2. \tag{BN15}$$

It buys any fixed pivot on wB_u at infinitesimal cost. The order of these two purchases is deliberate; no weak B diagonal is needed in the second step.

Both full normals and their only additional row. Use the complete $H_X, H_Y, H_{XY}, Y_P, Y_a, Y_c$ from SN18 and its bounded $\widehat{A}_X, \widehat{X}_a, X_c$ from SN17. These formulas only require τ, μ, β bounded, not n_s bounded. The entire first-normal part is

$$\begin{aligned} L_X &= \ell w \widehat{A}_X N_0 + \frac{uw\sqrt{S}}{\ell}\widehat{X}_a G + d_s\sqrt{S}\,quB - bB_u + wX_c(wB_u), \\ L_Y &= \sqrt{r(1-r)}\{Y_P G + \sqrt{S}\,Y_a R + Y_c(wB)\}. \end{aligned} \tag{BN16}$$

The apparent extra B_u coupling is paid by the actual circuit row E_1, using the exact, zero-face-safe identity

$$bB_u = -(w + \sqrt{q})\sqrt{b}\,E_1 - b(w + \sqrt{q})N_0. \tag{BN17}$$

All coefficients on the right have zero residue because b is bounded. No inverse of b or β is used. In the B term in L_X, write $d_sB = \sqrt{d_s}(\sqrt{d_s}B)$; this is likewise face-safe when $D = 0$. All direct normal coefficients are infinitesimal, and $H_X, H_Y \sim 1/2$, H_{XY} is infinitesimal. There is no missing X or Y normal and no imposed state ratio.

Allocation and all-tests conclusion. Keep a fixed positive fraction of each circuit row in BN6. Use the strict base rows $(N_0, G, X_e, N_y, \sqrt{d_s}B, E_1, E_2)$, omitting any zero row. Their main four diagonal residues are $1/2$; the last three have fixed positive pivots wherever their rows are nonzero. Terms containing R are infinitesimal matrix perturbations by BN7. Purchase sufficiently large fixed pivots on wB, wB_u by BN14–BN15, and use them for the bounded cross terms in BN13 and BN16. Formula BN17 pays the remaining first-normal row at infinitesimal cost. Every purchase is taken from an explicitly retained positive square or from the exact SOS remainder BN6. The resulting bounded Gram has strictly positive definite real residue. The surviving strict bounded-Gram lemma therefore gives finite tensor SOS for SN5 with both unrestricted normals restored.

This proves all infinitesimal u, w, including the literal axes. For noninfinitesimal Alice use the unchanged adjacent-C, A theorem CE1–CE10; C, A are unbounded on BN1. For the other Bob charts use the surviving small-e unmatched theorem SE10–SE12, since $\overline{\ell} = 1$. The sign gauges and real/imaginary state split are the original exact ones. The local argument already handles either sign of WZ, so no state normalization or state-chart exclusion is required. Both literal $D = 0$ and $\beta = 0$ are covered whenever they intersect BN1.

Verification status. The accompanying exact checker is `check_c144_bounded_h_ratio.py`. Bounds and separated SOS legality are the analytical argument above. At creation this theorem is a candidate pending independent lead review. The source case h/q unbounded and the ℓ-infinitesimal endpoint remain separate; no whole-C14.4 conclusion is asserted here.

Chapter 185

C14.4 at the entire ℓ-infinitesimal source endpoint

Component `pb96c9bfcc12d`
Source: `NEW/c1/c144_ell_endpoint_rederived.tex`
SHA256: `b96c9bfcc12d51c385d0aa7c4f9b46d3fc9322544414720685bccda5a3053e25`
Inventory use: EL0a–EL21 | Fresh complete EL0a–EL21 proof; entire ell-infinitesimal source endpoint | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

185.1 C14.4 at the entire ℓ-infinitesimal source endpoint

Theorem and exact scope. Retain the original native C14.4 physical source, all strict gaps, nonnegative event weights, total-scale conditions and prior-complement intersections. Let

$$S = \sigma + L, \quad x = \kappa S, \quad \nu = \sigma/S, \quad \ell = L/S, \quad \nu + \ell = 1, \quad \zeta = L/x = \ell/\kappa.$$

Suppose $S, \ell > 0$ are infinitesimal and ζ is noninfinitesimal. Then the complete original parent is a finite sum of squares in the separated ring $K_c \otimes_{\mathbb{R}} K_t$ for every independently chosen projective test and every real or complex state. The proof includes $\beta = 0$ and unrestricted openings of $r, 1-r, \theta$. The intersection with $D = 0$ is empty by the source estimate below. No earlier missing R21 endpoint proof is used.

Source estimates at this endpoint. Use the exact actual-source formulas FM2–FM3 and SE1–SE3, with $0 < r, h < 1$, $\theta = 1 - h$, $e = \kappa h/\nu$, $E = 1 - e \geq 1/18$, $U_s = (1+x)(1+\nu S r)$, and

$$J = 1 + xh - \nu S(1-r), \quad M = 1 + xr - \kappa\theta/\ell > 0, \quad Q_s = 1 - \ell S r,$$
$$a_0 = SAr, \qquad c_0 = SC(1-r), \qquad P = SP_0.$$

Put $g = \kappa/\ell = 1/\zeta$, which is bounded. Then $\nu \sim 1$ and $\kappa \in \mathrm{Inf}$. Since

$$0 \leq \beta\theta = \frac{U_s(\ell + \kappa h) - \theta P_0}{2P_0},$$

we have $\theta = O(\ell + \kappa) = O(\ell)$ and $\beta\theta = O(\ell)$, hence $h \sim 1$, $\beta\theta^2 = O(\ell^2)$ and $M \sim 1$. The exact source formulas give

$$\begin{gathered} J, Q_s, P_0, E \sim 1, \quad c_0, a_0 \sim \tfrac{1}{2}, \quad D/S \sim \tfrac{1}{2}, \quad m \in \mathrm{Inf}_{>0}, \\ \frac{qm}{\kappa^2} = \frac{U_s h}{2M\nu^2 P_0} \sim \tfrac{1}{2}, \qquad q = \ell/\nu, \\ \frac{\alpha}{S\ell^2} = \frac{(1+\nu S r)\{(1+xr)e/\ell + ME\}}{2Q_s P_0}, \qquad \frac{\alpha}{S\ell^2} - \frac{1+g}{2} \in \mathrm{Inf}\,. \end{gathered} \tag{EL0a}$$

The last difference is uniform for bounded g, because $e/\ell = gh/\nu$. All statements are inside the source field; none compares a testing rate to a source rate. In particular D/S is a positive unit, while β may be zero, bounded, or unbounded.

The full parent and local testing chart. Noninfinitesimal Alice tests and unmatched Bob residues will be joined at the end. The remaining chart has $u, v \ge 0$ infinitesimal, including literal zero axes. Write $a = u^2$, $c = v^2$, $b = \sqrt{q}$, $j = \nu b$, and $d = D + 1/2$. The complete original quarter-parent is

$$\begin{aligned} F/4 &= m(vW - X + uvY - uZ)^2 + C(1+c)(uW - Y)^2 \\ &\quad + A(1+a)(1+c)Y^2 + d(uW - Y + uvX - vZ)^2 \\ &\quad + (\alpha + \tfrac{1}{2})(1+a)(vY - Z)^2 + (\beta + \tfrac{1}{2})(1+c)(X + uZ)^2 \\ &\quad - \tfrac{1}{2}(1+a)(1+c)Z^2. \end{aligned} \tag{EL0b}$$

All state coordinates are arbitrary elements of the testing field. No state coordinate is divided by or bounded below. Put $R_s = 1 + xr$, $p = \kappa\theta/(\ell R_s)$, $\mu = m(1-p)$, and

$$V = v(W + uY), \qquad N = X + uZ - pV, \qquad Y = ruW + \sqrt{Sr(1-r)}N_y. \tag{EL1}$$

These are pure-source rescalings of polynomial state normals. The original m event is $(1-p)V - N$, the beta event is $N + pV$, and the core event is $uW - Y + puvV - (1+a)vZ + uvN$; thus no term has been dropped. On $N = N_y = 0$ define

$$\begin{aligned} L_0 &= \mu(1-p) + (\beta + \tfrac{1}{2})p^2, \qquad L_1 = (\beta + \tfrac{1}{2})p^2, \\ B_c &= 1 - r + pc(1 + ra), \\ \lambda(a,c) &= \frac{q}{\kappa^2}(1+ra)^2(L_0 + L_1 c), \\ A_P &= (1+c)[c_0(1-r) + a_0 r(1+a)] + S[dB_c^2 + (\alpha + \tfrac{1}{2})(1+a)r^2 c], \\ K &= (1+a)[dB_c + (\alpha + \tfrac{1}{2})r], \\ F_Z &= (1+a)[\alpha + Dc + dac]. \end{aligned} \tag{EL2}$$

Here $c_0 = SC(1-r)$ and $a_0 = SAr$. The complete graph scalar is

$$F_0 = c(1+ra)^2(L_0 + L_1 c)W^2 + A_P P^2 - 2vKPQ + F_Z Z^2, \quad P = uW/\sqrt{S}, \quad Q = \sqrt{S}Z. \tag{EL3}$$

Let $T = \kappa W, U = uZ$ and use the exact four-term circuit

$$\Psi = (\nu/\ell)cT^2 + \nu^2 cU^2 + \kappa\nu\ell Q^2 + \kappa P^2 - 4\kappa\nu vPQ. \tag{EL4}$$

Its four positive monomial square roots are $A_1 = (\kappa/b)vW$, $A_2 = \nu v u Z$, $A_3 = \nu b\sqrt{x}Z$, and $A_4 = \sqrt{\kappa}P$. Their product is a square. The usual two paired AM–GM identities make Ψ a separated SOS, including $WZ < 0$ by sign excess. The multiplier λ itself is a sum of separated monomial squares with positive pure-source coefficients; no mixed radical of λ is taken.

Set $\Gamma = P - \nu v Q$, $\Lambda = vP - \ell Q$, and

$$\widehat{A} = A_P - \kappa\lambda, \quad \widehat{K} = K - 2\kappa\nu\lambda, \quad \widehat{F} = F_Z - \lambda(\nu^2 ac + \kappa\nu\ell S).$$

Then the following equality is purely polynomial and exact:

$$F_0 - \lambda\Psi = \widehat{A}\Gamma^2 - 2vJ\Gamma Q + HZ^2, \quad J = \widehat{K} - \nu\widehat{A}, \quad H = \widehat{F} + \nu^2 Sc\widehat{A} - 2\nu Sc\widehat{K}. \tag{EL5}$$

The exact coefficient bounds are $\lambda_0 = qL_0/\kappa^2 \sim 1/2$ and $\lambda_1 = qL_1/\kappa^2 = O(\ell)$. Indeed the exact opening-safe formulas are

$$\lambda_1 = \frac{\theta(2\beta\theta + \theta)}{2\ell\nu R_s^2}, \qquad \lambda_0 = \frac{U_s h M}{2\nu^2 P_0 R_s^2} + \lambda_1.$$

Here θ/ℓ and $(\beta\theta)/\ell$ are bounded, so the formula for λ_1/ℓ has only bounded source factors. No comparison between S and ℓ is made. The mixed term arising from $-\nu^2\lambda_1 c^2 a$ is allocated in EL18 below using an actual row of the same circuit.

185.1.1 Exact helper rows and the complete two-normal conductor

The following exact budgets and complete normal conductors use a positive real fraction of Ψ and strict pivots on (Γ, Λ). The scalar calculation below supplies precisely these pivots; it does not use the state normals to obtain them. Put $\delta = v - b$, $G_0 = P - jQ$, $B_o = \delta Q$, and

$$\begin{aligned} E_{12} = A_1 - A_2 = (v/b)(T - jU), \quad V_h = v\delta U, \\ R_v = v(a-x)Z, \quad B = \sqrt{S}(c-q)Z, \quad R_0 = \ell\sqrt{S}(a-x)Z. \end{aligned} \tag{EL6}$$

The exact angular relations are

$$(v+b)G_0 = \Lambda + b\Gamma, \qquad \nu(v+b)B_o = \Lambda - v\Gamma = \nu B. \tag{EL7}$$

In particular B has a bounded linear expression in the two strict angular rows. Multiply MB3 by v^2/j^2 and use $\eta = \kappa(v+b)/j \in \mathrm{Inf}$. The exact SOS inequality is

$$V_h^2 \preceq_{\mathrm{SOS}} \frac{\delta^2}{\nu^2} E_{12}^2 + \eta(\Lambda + b\Gamma)^2 + \frac{\eta + 2\kappa}{\nu^2}(\Lambda - v\Gamma)^2. \tag{EL8}$$

For a direct verification, set $R_1 = vG_0, R_2 = vB_o$. The mixed-term replacement uses the legal SOS

$$\eta(R_1^2 + R_2^2) - 2\kappa\delta R_1 R_2/j = \frac{\kappa v}{j}(R_1 - R_2)^2 + \frac{\kappa b}{j}(R_1 + R_2)^2.$$

Replacing v^2 by $(v+b)^2$ leaves $(2vb + b^2)G_0^2$ and $(2vb + b^2)B_o^2$, with positive separated multipliers. The remaining MB3 square is $(v\delta T/j)^2$. Every cost in EL8 is infinitesimal, since $\kappa/j = (\kappa/\ell)b$ is infinitesimal.

Two further exact relations are

$$R_v = \frac{\kappa v\sqrt{S}}{\nu b}\Gamma - \frac{u}{\nu}E_{12} + \frac{xv(v-b)}{b}Z, \tag{EL9}$$

$$\frac{x^2}{qS}B^2 - \left(\frac{xv(v-b)}{b}Z\right)^2 = \frac{x^2}{q}(2vb + b^2)(v-b)^2 Z^2. \tag{EL10}$$

Thus R_v is free at infinitesimal Gram cost because $x^2/(qS) = \kappa S(\kappa/q)$ is infinitesimal. Finally

$$2\nu^2 ScR_v^2 + 2\nu^2(a-x)^2 B^2 - R_0^2 = [\nu\sqrt{S}(2c-q)(a-x)Z]^2 \tag{EL11}$$

buys R_0 with infinitesimal costs. None of EL6–11 bounds or divides a state coordinate or a mixed source/testing sum.

For the exact complete N half derivative set

$$\begin{aligned} B_0(c) &= [-\mu + (\beta + \tfrac{1}{2})p(1+c)]/\kappa, \\ F_a(a,c) &= r\kappa B_0(c) + d(1-r) + dpc(1+ra), \\ D_t(a,c) &= d(1+a) - \nu S F_a(a,c). \end{aligned} \tag{EL12}$$

All their coefficients are bounded, including at $\beta = 0$ or unbounded β: μ/κ and $(\beta + 1/2)p/\kappa = (\beta\theta + \theta/2)/(\ell R_s)$ are bounded. Write the monic polynomial differences $D_t - D_t(x,q) = (a-x)D_a + (c-q)D_c$ and $B_0(c) - B_0(q) = (c-q)B_{01}$. Source contact gives $D_t(x,q) = \nu B_0(q)$. The entire half derivative is

$$\begin{aligned} L_N = bB_0(c)E_{12} + uv\sqrt{S}F_a\Gamma - uvD_aR_v \\ + [-D_t(x,q) + \nu bB_{01}(v+b) - v(v+b)D_c]V_h. \end{aligned} \tag{EL13}$$

Hence EL8–10 allocate every N cross term without a state bound. The normal $\widehat{N} = \sqrt{1+\beta}N$ has diagonal at least $(\beta + 1/2)/(1+\beta) \geq 1/2$; dividing the cross coefficients by $\sqrt{1+\beta}$ only improves the displayed bounds.

The complete other half derivative is

$$L_y = \sqrt{r(1-r)}[Y_P\Gamma + vY_K Q], \tag{EL14}$$

where

$$\begin{aligned} Y_P &= (1+c)[-c_0 + a_0(1+a)] \\ &\quad + S\{c(1+ra)(L_0+L_1c) + d(-1+pac)B_c + (\alpha+\tfrac{1}{2})(1+a)rc\}, \\ Y_K &= \nu Y_P + (1+a)(D-\alpha-dpac). \end{aligned}$$

The polynomial coefficients are bounded, Y_P is infinitesimal, and actual source contact gives $Y_K(x,q)=0$. Monic division gives $Y_K = (a-x)Y_a + (c-q)Y_c$ with bounded coefficients, so

$$L_y = \sqrt{r(1-r)}[Y_P\Gamma + \sqrt{S}Y_a R_v + vY_c B]. \tag{EL15}$$

The exact quadratic normal coefficients are

$$\begin{aligned} H_N &= m + dac + (\beta+\tfrac{1}{2})(1+c), \\ H_y &= rc_0(1+c) + (1-r)a_0(1+a)(1+c) \\ &\quad + Sr(1-r)[d(-1+pac)^2 + (\alpha+\tfrac{1}{2})(1+a)c + ac(L_0+L_1c)], \\ H_{Ny} &= \sqrt{Sr(1-r)}uv[\kappa B_0(c) - d + dpac]. \end{aligned} \tag{EL16}$$

After the source-only normalization of N, the first two diagonal residues are respectively in $[1/2,1]$ and $1/2$, and the cross coefficient is infinitesimal. Equations EL13–16 include all terms of the full original parent; they require no W/Z or testing/source rate restriction. The scalar allocation and exact strict-Gram assembly are given next.

185.1.2 Complete scalar allocation, including the mixed radial term

Both $\widehat{A}$ and the coefficients of J, H in EL5 are bounded polynomials in a, c: their source coefficients are built from bounded $p, d, \alpha, c_0, a_0, \lambda_0, \lambda_1, \kappa, S, \nu$. In fact $p = O(\kappa)$, $\mu/\kappa = O(\kappa/\ell)$, and $(\beta+1/2)p^2 = O(\kappa^2)$. Consequently $\widehat{A} \sim 1/2$ on the local chart. Actual source contact, checked in the original parent, gives

$$J(x,q) = 0, \qquad H(x,q) = H_a(x,q) = H_c(x,q) = 0. \tag{EL17}$$

These are exact identities, not residue equalities. They also follow by evaluating the actual contact state in EL1: all four circuit monomials agree, $\Gamma = 0$, and both graph normals vanish. The vanishing first derivatives of the original contact parent then give the three scalar jets. The focused checker verifies all of them directly from FM2–FM3.

The additional actual circuit row

$$E_{23} = A_2 - A_3 = \nu(uv - \sqrt{xq})Z$$

resolves the otherwise unbounded mixed coefficient. Direct expansion using $x = \kappa S$ gives the exact identity

$$c(a-x)(c-q)Z^2 = \frac{c-q}{\nu^2}E_{23}^2 + \frac{2\sqrt{\kappa q}}{\nu}E_{23}B - \kappa B^2. \tag{EL18}$$

Every coefficient on the right is infinitesimal. To see the equality without division by a testing coordinate, use

$$(ac-xq)Z^2 = E_{23}^2/\nu^2 + 2\sqrt{xq}\,ZE_{23}/\nu, \qquad c(a-x) = ac - xq - x(c-q).$$

All square roots in this calculation are pure-source roots.

For completeness, the single circuit supplies positive fixed pivots on both E_{12} and E_{23}. If $WZ \geq 0$, use the two AM–GM pairings having respectively these differences and average them; then $\Psi - \frac{1}{4}(E_{12}^2 + E_{23}^2)$ is separated SOS. If W, Z are both negative, first change the common state sign;

the parent and all squared differences are unchanged. If $WZ < 0$, the negative product term of Ψ is nonnegative, and

$$\sum_{i=1}^{4} A_i^2 - \tfrac{1}{4}[(A_1 - A_2)^2 + (A_2 - A_3)^2]$$

is a real positive-definite quadratic form in the A_i. For example its leading 3×3 block has diagonal $(3/4, 1/2, 3/4)$ and successive leading minors $3/4, 5/16, 3/16$; the fourth diagonal is 1. This supplies a finite real Gram factorization. The remaining $-4\kappa\nu uvWZ$ is a separated square on this pure-testing sign chart. The zero-state boundaries use the same identities. Since $\lambda_0 \sim 1/2$ and $\lambda_1 \geq 0$, $\lambda - 1/4$ is a separated SOS multiplier:

$$\lambda - \tfrac{1}{4} = (\lambda_0 - \tfrac{1}{4}) + \lambda_0(2ra + r^2a^2) + \lambda_1 c(1 + ra)^2.$$

Thus $\lambda\Psi$ leaves strict real diagonal $1/16$ on each of E_{12}, E_{23} and an explicitly retained separated SOS remainder. No mixed square root of λ has been used.

Perform monic polynomial division in a and then in c. By EL17 there are bounded coefficient polynomials such that

$$\begin{aligned} J &= (a - x)J_a + (c - q)J_c, \\ H &= (a - x)^2 H_{20}(a, c) + (a - x)(c - q)H_{11}(c) + (c - q)^2 H_{02}(c). \end{aligned} \tag{EL19}$$

The division is by monic polynomials; it introduces no inverse of $a - x$, $c - q$, or any mixed expression. The following four particular coefficient identities are exact:

$$\begin{aligned} H_{20}(a, 0) &= -\kappa\nu\ell S\lambda_0 r^2, \\ H_{11}(0) &= -\alpha/q + 2\kappa\nu^2 S\lambda_0 r(1 + rx), \\ H_{02}(0)/S &= \nu^2 \left[(1 + x)\frac{\alpha}{S\ell^2} - g\nu\lambda_0(1 + rx)^2\right] \sim \tfrac{1}{2}, \\ H_{02}(c) &= H_{02}(0) + \nu^2 S^2 dp^2(1 + rx)^2 c. \end{aligned} \tag{EL20}$$

The first three also follow immediately by evaluating $H(a, 0) = \alpha(1 + a) - \kappa\nu\ell S\lambda_0(1 + ra)^2$ and using EL17; the last follows from the c^3 coefficient. In particular $H_{11}(0)/(S\ell)$ is bounded and $H_{02}(c)/S \sim 1/2$. The g terms cancel in the real residue of the third line by EL0a. This cancellation is why the independent ratio S/ℓ never enters the estimate.

Write $H_{20} = H_{20}(a, 0) + c\widetilde{H}_{20}$ and $H_{11} = H_{11}(0) + c\widetilde{H}_{11}$, with bounded coefficients. Substitution into EL5 gives the exact scalar assembly

$$\begin{aligned} F_0 = \lambda\Psi + \widehat{A}\Gamma^2 &+ \frac{H_{02}(c)}{S}B^2 \\ &- 2\sqrt{S}J_a\Gamma R_v - 2vJ_c\Gamma B - g\nu\lambda_0 r^2 R_0^2 + \widetilde{H}_{20}R_v^2 \\ &+ \frac{H_{11}(0)}{S\ell}R_0 B \\ &+ \widetilde{H}_{11}\left[\frac{c - q}{\nu^2}E_{23}^2 + \frac{2\sqrt{\kappa q}}{\nu}E_{23}B - \kappa B^2\right]. \end{aligned} \tag{EL21}$$

This includes every scalar term, including the one involving $\lambda_1 c^2 a$. It supplies strict real pivots on $(E_{12}, E_{23}, \Gamma, B)$; the two radial rows have arbitrary fixed positive pivots available at infinitesimal cost by EL9–11. There is no discarded source-dependent signed term.

185.1.3 Strict bounded Gram and all-test completion

Use the main rows

$$E_{12},\ E_{23},\ \Gamma,\ B,\ \widehat{N} = \sqrt{1 + \beta}N,\ N_y$$

and the three auxiliary rows V_h, R_v, R_0. EL21 and EL13–16 express the entire EL0b as a bounded quadratic form in these nine rows, together with the retained circuit SOS remainder. The real

diagonal of the main block is bounded below by $(1/16, 1/16, 1/2, 1/2, 1/2, 1/2)$. Its off-diagonal entries are infinitesimal except for couplings to the auxiliary rows. These are bounded, as are the possibly signed auxiliary diagonal entries.

Choose a sufficiently large fixed real positive diagonal on each auxiliary row, and purchase it using EL8–11. This is an exact SOS operation: if $Q - R^2$ is a displayed SOS budget, replace the quadratic form by its difference with a real multiple of $Q - R^2$ and retain that multiple as a separate SOS. EL8 and EL9–10 cost only infinitesimal entries on the main rows; EL11 is purchased last and costs an infinitesimal entry on R_v as well as on B. The real auxiliary diagonals can be chosen large enough that the real Schur complement is positive definite. This argument uses only a finite real matrix, never a mixed-field matrix inverse. The resulting bounded coefficient Gram has strictly positive real residue. The inherited strict separated bounded-Gram lemma in `parent_reference.tex` makes the complete form a finite SOS in $K_c \otimes_{\mathbb{R}} K_t$. Restoring the budget and circuit remainders proves the original full parent.

There is no restriction on W, Z, their ratio, or either normal; β was never discarded or divided by. The actual retained normal diagonal handles an arbitrarily large β before any cross-term estimate. The only inverse involving an arbitrarily small opening is the pure-source scaling used to define N_y; its inverse never appears in a Gram coefficient. All helper roots are products of source and testing factors or finite sums thereof.

The local proof includes $u = 0$ and $v = 0$, since it never divides by either. Negative u is reduced by $(u, Y, Z) \mapsto (-u, -Y, -Z)$ and negative v by $(v, X, Z) \mapsto (-v, -X, -Z)$; EL0b is invariant under each map. Real and imaginary state parts are handled separately. By EL0a both C, A are unbounded, so the surviving adjacent-C, A proof covers noninfinitesimal Alice tests. Since κ and e are infinitesimal, the surviving SE10–SE12 theorem covers every homogeneous Bob sine-square residue different from $\bar{\ell} = 0$, including Bob infinity. The only remaining Bob chart is exactly finite infinitesimal v, proved above. Therefore all projective tests and all states on the original ℓ-infinitesimal source intersection are covered. All original source predicates are retained; no numerical sampling or historical completion marker is an input to this conclusion.

Chapter 186

Fresh C14.4 completion when κ is small and both Bob source fractions are units

Component `p2edb55cd1f69`
Source: `NEW/c1/c144_finite_matched_rederived.tex`
SHA256: `2edb55cd1f6919d774d82054aee53db4ff6b63b6444cb535185ac06146cb308a`
Inventory use: FM1–FM22 | FM2–FM3 actual source formulas only; unit-fraction asymptotic scope not imported | Fresh full-parent proof FM1-FM22, candidate awaiting independent analytical review

186.1 Fresh C14.4 completion when κ is small and both Bob source fractions are units

Scope and status. This is a fresh candidate theorem, pending the focused algebra receipt and independent analytical review. It does not assert completion of the source endpoints $\nu \in \mathrm{Inf}$ or $\ell \in \mathrm{Inf}$, or of the unit-κ source. All original physical C14.4 source conditions, strict gaps, nonnegative event coefficients, total-scale conditions and prior complement intersections are retained. No earlier missing R21 proof body is invoked.

Theorem. Let the original native C14.4 source have

$$S = \sigma + L > 0, \qquad x = \kappa S, \qquad \nu = \sigma/S, \quad \ell = L/S, \quad \nu + \ell = 1, \qquad S, \kappa \in \mathrm{Inf}_{>0}, \tag{FM1}$$

with ν, ℓ positive bounded units. Then the complete original parent is a finite SOS in $K_c \otimes_{\mathbb{R}} K_t$ for every independent projective test and every real or complex state. The argument below supplies the missing finite matched-Bob chart; the other tests use the surviving explicitly scoped SE theorem.

Exact opening-safe source dictionary. Write $\theta = 1 - h$, $0 < r, h < 1$, $e = \kappa h/\nu$, $E = 1 - e$, and set

$$\begin{aligned} J &= 1 + xh - \nu S(1 - r), \quad M = 1 + xr - \kappa\theta/\ell, \quad Q_s = 1 - \ell S r, \\ P_0 &= 1 + S\{\nu\ell + 2\nu\kappa h + \kappa\ell + \kappa^2 h\theta\} + S^2\{\nu r\kappa\ell + \nu r\kappa^2 h\}. \end{aligned} \tag{FM2}$$

These are the actual native source quantities, with $P = SP_0$. The converted P1 coefficients, without applying P1's different unit-L asymptotic hypotheses, are exactly

$$\begin{aligned} c_0 = SC(1-r) &= \frac{(1+x)(1+\nu Sr)E}{2P_0}, \\ a_0 = SAr &= \frac{(1+xh)E + (1-r)(\ell S + xh)}{2P_0}, \\ d_0 = D/S &= \frac{(1+x)(1+\nu Sr) - JP_0}{2JSP_0}, \\ a_1 = \alpha/S &= \frac{(1+\nu Sr)\{\ell(1+xr)e + \ell^2 ME\}}{2Q_sP_0}, \\ b_1 = \beta\theta &= \frac{(1+x)(1+\nu Sr)(\ell+\kappa h) - \theta P_0}{2P_0}, \\ \mu = m/\kappa^2 &= \frac{(1+x)(1+\nu Sr)h}{2M\ell\nu P_0}. \end{aligned} \tag{FM3}$$

The numerator in the d_0 formula has a factor S, so its displayed S denominator cancels in the source polynomial ring. Thus all six coefficients belong to the bounded source algebra generated by $S, \kappa, h, r, \nu, \ell$ and inverses of $J, M, Q_s, P_0, \nu, \ell$. None of these inverses is of an opening $r, 1-r, \theta, D$ or β. The original source gives

$$\begin{gathered} J, M, Q_s, P_0, E \sim 1, \qquad c_0, a_0 \sim \tfrac{1}{2}, \qquad d_0 \sim \nu^2/2, \quad a_1 \sim \ell^2/2, \\ b_1 - (h-\nu)/2 \in \mathrm{Inf}, \qquad \mu - h/(2\nu\ell) \in \mathrm{Inf}, \qquad \bar{h} \geq \bar{\nu} > 0. \end{gathered} \tag{FM4}$$

The last inequality follows from $b_1 = \beta\theta \geq 0$. Hence μ is a positive bounded unit. These estimates are uniform through $r, 1-r, \theta \to 0$; no comparison with a testing quantity has been used. In this sector D/S is a positive unit, so $D = 0$ has empty intersection; $\beta = 0$ is allowed.

Exact full parent and unrestricted normals. On the finite Bob chart write $D_u = 1 + u^2$, $D_v = 1 + v^2$. The complete quarter-parent is

$$\begin{aligned} F/4 = {} & m(vW - X + uvY - uZ)^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW - Y + uvX - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ & + (\beta + \tfrac{1}{2})D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \end{aligned} \tag{FM5}$$

No state restriction is made. Put

$$N_x = X + huZ, \qquad N_y = \frac{Y - ruW}{\sqrt{Sr(1-r)}}. \tag{FM6}$$

The source square root is nonzero even when its size is arbitrarily small. It is used only to define a permitted tensor root; every coefficient below is bounded without its inverse.

After the surviving unmatched-residue and adjacent-C, A exits it is enough to take u infinitesimal and v a bounded positive unit with the same residue as $b = \sqrt{\ell/\nu}$. Set

$$\begin{gathered} a = u^2, \quad j = \sqrt{\nu\ell}, \quad T = \kappa W, \quad U = uZ, \\ P = uW/\sqrt{S}, \quad Q = \sqrt{S}Z, \quad N = T - jU, \quad G = P - jQ, \\ \delta = v - b, \quad B = \delta Q, \quad B_u = \delta U. \end{gathered} \tag{FM7}$$

Every state coordinate remains arbitrary, including zero and unbounded coordinates. The rows satisfy the exact MB identities

$$\begin{aligned} R_Z = (a-x)Z &= (\kappa\sqrt{S}/j)G - (u/j)N, \\ (a-x)\delta Z^2 &= (\kappa/j)GB - (1/j)NB_u, \\ \delta^2N^2 + 2j\kappa\delta GB + 2j^2\kappa B^2 - j^2B_u^2 &= (\delta T)^2. \end{aligned} \tag{FM8}$$

These are polynomial identities, not ordered-ring inequalities.

The full graph scalar. Let F_0 denote FM5 with $N_x = N_y = 0$; it is used only as the constant coefficient of the full normal expansion. Define bounded source/testing polynomials

$$\begin{aligned}
A_T &= \mu v^2(1+ra)^2,\\
A_P &= (1+v^2)\{c_0(1-r)+a_0 r(1+a)\}\\
&\quad + S\{(D+\tfrac12)(1-r)^2+(\alpha+\tfrac12)(1+a)r^2v^2\},\\
A_Q &= d_0v^2+a_1,\\
A_U &= m\theta^2+\alpha+b_1\theta+\theta^2/2\\
&\quad + v^2\{2Dh+h-\tfrac12+b_1\theta+\theta^2/2\}+(D+\tfrac12)h^2v^2a,\\
K &= m\theta(1+ra)+(D+\tfrac12)(1-r)(1+ha)+(\alpha+\tfrac12)(1+a)r,\\
H &= vK-\kappa jA_T,\quad R_U = A_U-j^2A_T,\\
g &= jA_P-H,\quad R_Q = A_Q+j^2A_P-2jH,\\
J_Z &= SR_Q+aR_U.
\end{aligned} \tag{FM9}$$

Direct expansion of every term of FM5 gives

$$F_0 = A_TN^2+A_PG^2+2gGQ+J_ZZ^2. \tag{FM10}$$

For example the only relation used to obtain FM10 from the five monomials T^2, U^2, P^2, Q^2, PQ is $TU = \kappa PQ$.

The actual source stationarity identities, also verified directly from FM2–FM3, are

$$g(x,b) = 0, \qquad J_Z(x,b) = \partial_a J_Z(x,b) = \partial_v J_Z(x,b) = 0. \tag{FM11}$$

These are coefficient-field identities at the actual source contact; they impose no equation on independent tests.

Finite polynomial division. All following quotients mean monic polynomial division in formal a or v, performed before evaluation in the tensor ring. No mixed inverse is introduced. There are bounded coefficient polynomials $g_a, g_v, H_{20}, H_{11}, H_{02}$ defined exactly by

$$\begin{aligned}
g(a,v) &= (a-x)g_a(a,v)+(v-b)g_v(v),\\
J_Z(a,v) &= (a-x)^2H_{20}(a,v)+(a-x)(v-b)H_{11}(v)+S(v-b)^2H_{02}(v).
\end{aligned} \tag{FM12}$$

For clarity, first divide $g(a,v)-g(x,v)$ by $a-x$ and then $g(x,v)$ by $v-b$. For the second line, first subtract the constant and linear Taylor terms in $a-x$ and divide the remainder by $(a-x)^2$. Divide $\partial_a J_Z(x,v)$ by $v-b$ for H_{11}. Finally divide the bounded polynomial $R_Q(x,v)+\kappa R_U(x,v)$ by $(v-b)^2$ for H_{02}. FM11 makes every division exact. In particular H_{02} is bounded without dividing a bounded error by S.

Substituting FM8 into FM10 now yields the exact scalar identity

$$\begin{aligned}
F_0 &= A_TN^2+A_PG^2+2g_vGB+H_{02}B^2+H_{11}\{(\kappa/j)GB-(1/j)NB_u\}\\
&\quad + H_{20}R_Z^2+2\sqrt{S}\,g_aGR_Z.
\end{aligned} \tag{FM13}$$

Thus every coefficient is bounded and all remaining radial contributions are infinitesimal matrix entries on (N,G).

Here and below a displayed residue formula uses the real residues of its bounded source parameters. FM3 gives explicitly

$$\bar A_T = \frac{h}{2\nu^2},\quad \bar A_P = \frac{1}{2\nu},\quad \bar g_v = \frac{\ell-\nu}{2},\quad \bar H_{02} = \frac{\nu}{2},\quad \bar H_{11} = -\theta j. \tag{FM14}$$

For example at $S = \kappa = 0$ and $a = 0$ the three relevant polynomials are $g = j(1+v^2)/2 - v/2$, $R_Q = \nu(v-b)^2/2$, and $R_U = \theta(\ell-\nu v^2)/2$. These establish FM14 by the finite divisions just specified.

Both complete first-normal terms. Write the full expansion of FM5 as

$$F/4 = F_0 + H_x N_x^2 + H_y N_y^2 + 2H_{xy} N_x N_y + 2N_x L_x + 2N_y L_y + \beta D_v N_x^2. \tag{FM15}$$

The exact bounded diagonal and cross coefficients are

$$\begin{aligned} H_x &= m + (D+\tfrac{1}{2})u^2v^2 + D_v/2, \\ H_y &= rc_0 D_v + (1-r)a_0 D_u D_v \\ &\quad + Sr(1-r)\{mu^2v^2 + D + \tfrac{1}{2} + (\alpha+\tfrac{1}{2})D_u v^2\}, \\ H_{xy} &= -uv\sqrt{Sr(1-r)}(m + D + \tfrac{1}{2}). \end{aligned} \tag{FM16}$$

In particular $\bar{H}_x = \bar{H}_y = 1/(2\nu)$ and $\bar{H}_{xy} = 0$. The last term of FM15 is retained whole; it allows β to be arbitrarily large or zero.

Define

$$\begin{aligned} L_U &= m\theta - (mj/\kappa)v(1+ra) - (D+\tfrac{1}{2})v^2(1+ha) + (b_1 + \theta/2)(1+v^2) \\ &\quad + (D+\tfrac{1}{2})(1-r)vjS, \\ Y_P &= (1+v^2)\{-c_0 + a_0(1+a)\} + S\{mv^2(1+ra) - (D+\tfrac{1}{2})(1-r) + (\alpha+\tfrac{1}{2})(1+a)rv^2\}, \\ Y_Q &= -m\theta a + (D+\tfrac{1}{2})(1+ha) - (\alpha+\tfrac{1}{2})(1+a), \qquad Y_C = jY_P + vY_Q. \end{aligned} \tag{FM17}$$

The expression $mj/\kappa = \kappa\mu j$ is bounded and infinitesimal. The complete original normal terms are exactly

$$\begin{aligned} L_x &= -\kappa\mu v(1+ra)N + (D+\tfrac{1}{2})(1-r)vu\sqrt{S}\,G + L_U U, \\ L_y &= \sqrt{r(1-r)}\{Y_P G + Y_C Q\}. \end{aligned} \tag{FM18}$$

The actual source gives $L_U(x,b) = Y_C(x,b) = 0$. Perform the same finite monic divisions to write

$$L_U = (a-x)l_a + (v-b)l_v, \qquad Y_C = (a-x)y_a + (v-b)y_v. \tag{FM19}$$

Then $L_U U = ul_a R_Z + l_v B_u$ and $Y_C Q = \sqrt{S}\,y_a R_Z + y_v B$. All their coefficients are bounded. Moreover $\bar{l}_v = -j$, while Y_P and y_v have zero residue. This follows directly from the source-limit identities $L_U = (\ell - \nu v^2)/2$ at $a = 0$, and $Y_P = Y_Q = 0$ at $a = 0$. Consequently $L_x + jB_u$ is an infinitesimal bounded combination of (N, G, B_u) and L_y is an infinitesimal bounded combination of (N, G, B). This accounts for both unrestricted first-normal terms, including all $\beta\theta$ terms and the independent $r, 1-r, \theta$ openings.

A strict residue for the full six-row Gram. Use FM8 with a fixed real $c = 4$ to introduce a cB_u^2 pivot at infinitesimal Gram cost. More explicitly subtract

$$(c/j^2)\delta^2 N^2 + (2c\kappa/j)\delta GB + 2c\kappa B^2 \tag{FM20}$$

from the bounded part of FM15, add cB_u^2 to that part, and retain the exact positive square $(\sqrt{c}\,\delta T/j)^2$. Their sum is unchanged by FM8. The residue on (N_x, N_y, N, G, B, B_u) is therefore

$$\frac{N_x^2 + N_y^2}{2\nu} + \frac{h}{2\nu^2}N^2 + \frac{G^2}{2\nu} + (\ell - \nu)GB + \frac{\nu}{2}B^2 + \theta N B_u - 2jN_x B_u + 4B_u^2. \tag{FM21}$$

The (G, B) block has determinant $\nu\ell > 0$ and positive diagonal. The remaining block has positive diagonal pivots $1/(2\nu), 1/(2\nu), h/(2\nu^2)$, and its final scalar Schur pivot is

$$4 - 2\nu j^2 - \frac{\theta^2\nu^2}{2h} > 0. \tag{FM22}$$

Indeed $j^2 \le 1/4$, $0 < \nu \le 1$, $h \ge \nu$ and $0 \le \theta \le 1$ in residue make the two subtracted terms at most $1/2$ each. All entries before taking residues are finite sums of bounded separated factors. The surviving strict bounded-Gram lemma therefore supplies a finite tensor SOS. Restoring the single square from FM20 and the whole positive $\beta D_v N_x^2$ row gives the entire FM5. No mixed matrix inverse or mixed square root has been used.

Testing, sign, and boundary joins. The argument includes $u = 0$ and every real state without a state normalization or a sign condition. Negative u is changed to $-u$ together with $(Y, Z) \mapsto (-Y, -Z)$; negative v is changed to $-v$ together with $(X, Z) \mapsto (-X, -Z)$. FM5 is exactly invariant under these transformations. On FM1, $e = \kappa h/\nu$ is infinitesimal and C, A are unbounded. The surviving SE10–SE12 theorem covers every unmatched homogeneous Bob residue, including Bob zero and infinity, and invokes the unchanged adjacent-C, A theorem for noninfinitesimal Alice tests. The only remaining Bob residue has finite nonzero v and $\bar{v}^2 = \bar{\ell}/\bar{\nu}$, which is the matched chart just proved. Thus the union covers all projective tests on FM1. Real and imaginary state parts are treated separately for complex states. All divisions and normalizations used in these joins are pure source or pure testing operations; all original source predicates remain.

Dependencies and limits. The analytical inputs are the surviving native source formulas P1 converted as FM2–FM3, the full original parent, the strict separated bounded-Gram lemma in `parent_reference.tex`, and `proofs/c144/small_e_unmatched_bob.tex` with its preserved EB and adjacent-C, A dependencies. The exact budget is independently printed in `c144_matched_bob_budget.tex`. The source endpoint and unit-κ sectors require separate complete arguments; this theorem makes no claim about them.

Chapter 187

A source-free angular budget for the finite matched-Bob corner

Component `pe55ec4890781`
Source: `NEW/c1/c144_matched_bob_budget.tex`
SHA256: `e55ec489078157a3ce08ceb2f3db2235d1f6c4b6ba907ce476bfcaa27ce0af9d`
Inventory use: MB1–MB6 | MB3 exact source-free budget | Exact source-free MB1-MB6 budget

187.1 A source-free angular budget for the finite matched-Bob corner

This is an exact auxiliary identity, not by itself a C14.4 admission. Work in the separated tensor ring $K_c \otimes_{\mathbb{R}} K_t$. Let S, κ, ν, ℓ be positive source quantities, with $\nu + \ell = 1$, and put

$$x = \kappa S, \qquad j = \sqrt{\nu\ell}, \qquad a_B = \sqrt{\ell/\nu}. \tag{MB1}$$

For independent testing variables u, v, W, Z define

$$\begin{gathered} T = \kappa W, \quad U = uZ, \quad P = uW/\sqrt{S}, \quad Q = \sqrt{S}Z, \\ N = T - jU, \quad G = P - jQ, \quad \delta = v - a_B, \\ B = \delta Q, \quad B_u = \delta U. \end{gathered} \tag{MB2}$$

The monomial relation $TU = \kappa PQ$ is exact. It gives

$$\delta^2 N^2 + 2j\kappa\delta GB + 2j^2\kappa B^2 - j^2 B_u^2 = (\delta T)^2. \tag{MB3}$$

In particular, if ν, ℓ are positive bounded units, κ is infinitesimal and δ has zero real residue, then any fixed real positive diagonal on B_u can be purchased at an infinitesimal perturbation of the Gram matrix on (N, G, B). Indeed, for any fixed $c > 0$ the exact SOS remainder is

$$\frac{c}{j^2}\delta^2 N^2 + \frac{2c\kappa}{j}\delta GB + 2c\kappa B^2 - cB_u^2 = \frac{c}{j^2}(\delta T)^2. \tag{MB4}$$

Every displayed coefficient is a finite sum of bounded separated factors, with zero real residue except the constant c in the desired B_u pivot. The only inverses are of positive source units. The factor δ is a difference of a pure testing and a pure source quantity; no inverse or radical of that difference is taken. The right side has the displayed separated SOS root $\sqrt{c}\,\delta T/j$.

Two exact radial conductors and one cross conductor are

$$\begin{aligned} (u^2 - x)W/\sqrt{S} &= uG - \sqrt{S}N, \\ (u^2 - x)Z &= (\kappa\sqrt{S}/j)G - (u/j)N, \\ (u^2 - x)\delta Z^2 &= (\kappa/j)GB - (1/j)NB_u. \end{aligned} \tag{MB5}$$

These identities require no bound on the state coordinates or on any ratio of source and testing infinitesimals. In the local matched chart with u infinitesimal their first two right sides have only infinitesimal coefficients on (N, G).

For comparison, the four-term contact circuit is

$$\begin{aligned}\Psi &= (\nu/\ell)v^2T^2 + \nu^2v^2U^2 + \kappa\nu\ell Q^2 + \kappa P^2 - 4\kappa\nu vPQ \\ &= (v/a_B)^2N^2 + \kappa G^2 + (2\kappa\nu/a_B)PQ(v - a_B)^2.\end{aligned} \tag{MB6}$$

On the positive testing/state sign chart the last summand is itself a separated square. The budget MB3 is stronger for the present local purpose: it uses only the actual monomial relation, and does not divide by a weak circuit coefficient.

The full scalar and all original state normals still have to be placed in this frame before using MB4. No scalar or whole-parent conclusion is asserted here.

Chapter 188

The full unit-κ, infinitesimal-ν source endpoint

Component `p578719565e9d`
Source: `NEW/c1/c144_nu_endpoint_rederived.tex`
SHA256: `578719565e9de6441849cadd8b617dd34d2040bc0e0a0cbbdd7ddb813e04a408`
Inventory use: NU1–NU26 | Fresh NU1-NU26 full theorem | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

188.1 The full unit-κ, infinitesimal-ν source endpoint

Scope. Retain every original physical C14.4 source predicate, strict gap, nonnegative event coefficient, and original prior-complement intersection. This theorem assumes

$$S = \sigma + L \in \mathrm{Inf}_{>0}, \quad \kappa = x/S \text{ a positive bounded unit}, \quad \nu = \sigma/S \in \mathrm{Inf}_{>0}, \quad \ell = L/S = 1 - \nu. \tag{NU1}$$

It proves a finite separated tensor SOS for the complete original parent at every projective test and every real or complex state. The sector with both κ, ν infinitesimal is not included. No relative source/testing rate, state-ratio bound, or mixed inverse is used. The original literal $D = 0$ and $\beta = 0$ faces are included.

Opening-safe source parameters and the two sphere masses. Use the exact FM2–FM3 native dictionary, and write

$$\begin{aligned} q &= \nu/\ell, \quad \zeta = \ell/\kappa = L/x, \quad e = \kappa h/\nu, \quad E = 1 - e, \\ c_0 &= SC(1-r), \quad a_0 = SAr, \quad d_0 = D/S, \quad a_1 = \alpha/S. \end{aligned} \tag{NU2}$$

In NU1, q is infinitesimal and ζ is a positive bounded unit. The exact source sphere SE3 gives $e(B/S) = \kappa H_0 + \nu \mathcal{H} b_0^2$. Since $B/S \sim 1 + \kappa$, $H_0 \sim 1$, and $\mathcal{H} \sim E^2$, its real residues satisfy

$$e = \frac{\kappa}{1+\kappa}, \quad E = \frac{1}{1+\kappa}, \quad h \in \mathrm{Inf}, \quad \theta \sim 1, \quad c_0 = a_0 = a_1 = E/2. \tag{NU3}$$

The two equalities for e, E in this display are residue equalities; the definition $e = \kappa h/\nu$ remains exact. In particular e, E, c_0, a_0, a_1 are positive bounded units.

Define actual nonnegative source coefficients

$$d_s = \frac{D}{Sq^2}, \qquad n_s = \frac{\beta}{q^2}. \tag{NU4}$$

These are bounded even when either is zero. More explicitly the SE3 sphere identities give

$$d_s = \frac{\ell^2 \mathcal{H}}{2P_0 J} d_0^{\mathrm{sph}\,2}, \qquad n_s = \frac{\ell^2 \mathcal{H}}{2\kappa P_0 \theta} b_0^{\mathrm{sph}\,2}, \qquad J d_s + \kappa\theta n_s = \frac{\ell^2 \mathcal{H}}{2P_0}. \tag{NU5}$$

Here $(d_0^{\mathrm{sph}})^2 + (b_0^{\mathrm{sph}})^2 = 1$; the superscript distinguishes sphere coordinates from D/S. Every denominator in NU5 is a positive unit. At least one of d_s, n_s is a positive unit. No sphere coordinate is inverted.

The sole possibly unbounded event coefficient needed locally is m. Remove its pole by its actual contact-zero event, not by dividing an error by M. With $R_s = 1 + xr$ set

$$p = \frac{\kappa\theta}{\ell R_s} \in (0,1), \qquad \mu_c = m(1-p) = \frac{\kappa e(1+x)(1+\nu Sr)}{2\ell P_0 R_s}. \tag{NU6}$$

Thus μ_c is a positive bounded unit and $m \geq \mu_c$. Choose a fixed positive real $\epsilon < m$. On the reciprocal Bob chart define $V = W + uY$ and $B_c = w(X + uZ)$. Retain the whole positive packet $(m-\epsilon)(B_c - pV)^2$. The remaining m quadratic is exactly

$$m_0 B_c^2 - 2m_1 B_c V + m_2 V^2, \qquad m_0 = \epsilon, \quad m_1 = \mu_c + \epsilon p, \quad m_2 = \mu_c(1+p) + \epsilon p^2. \tag{NU7}$$

All three coefficients are bounded through $M \to 0$ with arbitrary rate. Their residues are

$$\mu_c = \frac{\kappa^2}{2(1+\kappa)}, \quad p = \kappa, \qquad m_2 = \kappa^2(\epsilon + \tfrac{1}{2}) > 0. \tag{NU8}$$

The source $M > 0$ also implies $0 < \bar{\kappa} \leq 1$.

Local reciprocal chart and exact angular budgets. Assume u, w are infinitesimal, including literal zero. Use the homogeneous reciprocal parent $F_w/4$ from P4, with its m term replaced by NU7 and all other terms intact. Put

$$\begin{gathered} X_e = X + ewW, \quad N_y = (Y - ruW)/\sqrt{Sr(1-r)}, \quad a = u^2, \quad c = w^2, \quad \delta = c - q, \\ P = uW/\sqrt{S}, \quad Q = \sqrt{S}\, wZ, \quad N = W - \zeta uwZ, \quad G = P - \ell Q, \\ B = \sqrt{S}\,\delta Z, \quad B_u = u\delta Z, \quad R = w(a-x)Z = \frac{\sqrt{S}}{\zeta}G - \frac{u}{\zeta}N. \end{gathered} \tag{NU9}$$

Every row is a finite separated expression. No state coordinate is divided by another. Since $W(uwZ) = PQ$, two exact identities are

$$\begin{aligned} \delta^2 N^2 + 2\zeta\delta wGB + 2\zeta\ell w^2 B^2 - \zeta^2(wB_u)^2 &= (\delta W)^2, \\ \delta^2 G^2 + 2\ell\delta wNB_u + 2\ell\zeta w^2 B_u^2 - \ell^2(wB)^2 &= (\delta P)^2. \end{aligned} \tag{NU10}$$

Consequently, from either positive sphere pivot in NU5, any fixed positive diagonal budgets on both wB and wB_u can be bought at infinitesimal bounded-Gram cost. To see this without a hidden division, if d_s is a unit, replace B by $\sqrt{d_s}B/\sqrt{d_s}$ in the first identity, divide by ζ^2, and use $wB = w(\sqrt{d_s}B)/\sqrt{d_s}$ for the other budget. All cost coefficients are infinitesimal. If n_s is a unit, use the second identity divided by ℓ^2 and $wB_u = w(\sqrt{n_s}B_u)/\sqrt{n_s}$. The factor ζw^2 is infinitesimal because ζ is a unit in NU1. This is the precise restriction preventing a silent extension to infinitesimal κ.

The entire scalar graph before normal restoration. Write $D = Sd_0$, $\alpha = Sa_1$, and define bounded coefficient polynomials in the independent testing squares a, c:

$$\begin{aligned} A_P &= (1+c)\{c_0(1-r) + a_0 r(1+a)\} \\ &\quad + S\{(D+\tfrac{1}{2})(1-r)^2 c + (\alpha+\tfrac{1}{2})(1+a)r^2\}, \\ A_N &= m_2(1+ra)^2 + 2m_1 ec(1+ra) + m_0 e^2 c^2 + (\beta+\tfrac{1}{2})e^2 c(1+c) \\ &\quad - (D+\tfrac{1}{2})e\{2(1-r) - e\}ac, \\ K &= m_1(1+ra) + m_0 ec + (D+\tfrac{1}{2})(1-r-e) \\ &\quad + (\alpha+\tfrac{1}{2})(1+a)r + (\beta+\tfrac{1}{2})e(1+c), \\ F_Z &= D + \alpha c + \beta a + (m_0 + \alpha + \beta + \tfrac{1}{2})ac, \\ J_1 &= K - \ell A_P, \qquad L_1 = J_1 - \zeta A_N, \\ H &= F_Z - \ell^2 ScA_P + ac(\zeta^2 A_N - 2\zeta J_1). \end{aligned} \tag{NU11}$$

For $X=-ewW$, $Y=ruW$, the exact graph scalar is

$$F_0=A_NN^2+A_PG^2-2uwZL_1N+HZ^2. \tag{NU12}$$

This graph is used to compute coefficients; the unrestricted normals are restored explicitly below. The actual source contact has $a=x,c=q,W=\zeta uwZ$, so $N=G=X_e=N_y=0$. The retained row B_c-pV also vanishes there. Direct source substitution in NU11 yields

$$L_1(x,q)=0,\qquad H(x,q)=H_a(x,q)=H_c(x,q)=0,\qquad H(a,0)=D+\beta a. \tag{NU13}$$

These are exact rational source identities; no independent testing angle is identified with a source angle in the certificate.

All divisions in the next display are finite monic polynomial divisions at the bounded pure-source contacts x,q. They define bounded coefficient polynomials $L_a,L_c,H_{20},H_{11},H_{02}$:

$$\begin{aligned}
L_1(a,c)&=(a-x)L_a(a,c)+(c-q)L_c(c),\\
H(a,c)&=H(x,c)+(a-x)H_a(x,c)+c(a-x)^2H_{20}(a,c),\\
H_a(x,c)&=(c-q)\{-\beta/q+cH_{11}(c)\},\\
H(x,c)&=S(c-q)^2\{d_s+\kappa n_s+cH_{02}(c)\}.
\end{aligned} \tag{NU14}$$

Here the factor c in the quadratic a remainder follows from $H(a,0)=D+\beta a$. The two constant terms follow by putting $c=0$: $H_a(x,0)=\beta$ and $H(x,0)=D+\beta x$. In particular the displayed apparent divisions by q are bounded: $\beta/q=n_sq$. Finally $H(x,c)/S$ is a bounded coefficient polynomial directly from NU11, because $x=\kappa S$, $D/S=d_0$, and $\alpha/S=a_1$ are bounded. Thus dividing it by the monic $(c-q)^2$ introduces no unbounded coefficient. This verifies all asserted coefficient bounds without using relative orders of S,q,u,w.

The useful mixed identity is

$$c(a-x)(c-q)Z^2=\frac{1}{\zeta}\{G(wB)-N(wB_u)\}. \tag{NU15}$$

Substituting NU14 into NU12 gives the complete exact scalar allocation

$$\begin{aligned}
F_0=&A_NN^2+A_PG^2+d_sB^2+n_sB_u^2+H_{02}(wB)^2+H_{20}R^2\\
&+\frac{H_{11}-n_s}{\zeta}\{G(wB)-N(wB_u)\}-2uL_aNR-2L_cN(wB_u).
\end{aligned} \tag{NU16}$$

The residues of A_N,A_P are respectively $\kappa^2(\epsilon+1/2)$ and $E/2$, both positive. The row R has infinitesimal coefficients on (N,G) by NU9. Every remaining signed term is a bounded quadratic term on N,G,wB,wB_u. The angular rows are retained as the complete nonnegative squares $(\sqrt{d_s}B)^2$ and $(\sqrt{n_s}B_u)^2$; neither is prematurely discarded.

Both complete state normals. Put $s_y=\sqrt{Sr(1-r)}$ and $r_y=\sqrt{r(1-r)}$. Define

$$\begin{aligned}
A_x=&\ -m_0ec-m_1(1+ra)+(D+\tfrac12)a(1-r-e)-e(\beta+\tfrac12)(1+c),\\
T_x=&\ \zeta cA_x+\beta-D+c(m_0+\beta+\tfrac12),\\
Y_P=&\ (1+c)\{-c_0+a_0(1+a)\}\\
&+S\{m_2(1+ra)+m_1ec-(D+\tfrac12)c(1-r-e)+(\alpha+\tfrac12)(1+a)r\},\\
Y_Q=&\ -m_1a+(D+\tfrac12)-(\alpha+\tfrac12)(1+a),\qquad Y_C=\ell Y_P+Y_Q.
\end{aligned} \tag{NU17}$$

The entire capped quarter-parent, before any allocation, is

$$F_0+H_xX_e^2+H_yN_y^2+2H_{xy}X_eN_y+2L_xX_e+2L_yN_y, \tag{NU18}$$

where

$$\begin{aligned} H_x &= m_0c + (D+\tfrac12)a + (\beta+\tfrac12)(1+c),\\ H_y &= rc_0(1+c) + (1-r)a_0(1+a)(1+c)\\ &\quad + Sr(1-r)\{m_2a + (D+\tfrac12)c + (\alpha+\tfrac12)(1+a)\},\\ H_{xy} &= -uws_y(m_1 + D + \tfrac12),\\ L_x &= wA_xN + uZT_x, \qquad L_y = r_y\{Y_PG + Y_CQ\}. \end{aligned} \tag{NU19}$$

These formulas retain every quadratic state coefficient and both first normals. Their real hard residues are $H_x = 1/2$, $H_y = E/2$, $H_{xy} = 0$.

Exact source substitution gives $T_x(x,q) = Y_C(x,q) = 0$. Moreover $T_x(a,0) = \beta - D$, and its a dependence has a factor c. Thus bounded monic divisions give

$$\begin{aligned} T_x(a,c) &= c(a-x)X_a + (c-q)\{(D-\beta)/q + cX_c(c)\},\\ Y_C(a,c) &= (a-x)Y_a(a,c) + (c-q)Y_c(c). \end{aligned} \tag{NU20}$$

No coefficient here needs an inverse of a sphere coordinate. Using NU4 and NU9, the full normal conductors become

$$\begin{aligned} L_x &= wA_xN + uwX_aR + d_s\sqrt{S}\,quB - n_sqB_u + wX_c(wB_u),\\ L_y &= r_y\{Y_PG + \sqrt{S}\,Y_aR + Y_c(wB)\}. \end{aligned} \tag{NU21}$$

The two potentially weak terms are face-safe: $d_s\sqrt{S}\,quB = (\sqrt{d_s}\sqrt{S}\,qu)(\sqrt{d_s}B)$ and $n_sqB_u = (\sqrt{n_s}q)(\sqrt{n_s}B_u)$. Their coefficients are infinitesimal, even when either sphere mass is zero. Also Y_P has zero residue by NU3 and NU17. All other terms in NU21 have infinitesimal coefficients on N, G, R or bounded coefficients on the two angular helper rows.

Strict complete Gram and restoration. Begin with the six rows

$$(N, G, X_e, N_y, \sqrt{d_s}B, \sqrt{n_s}B_u). \tag{NU22}$$

Their real diagonal residue in NU16–NU21 is

$$\operatorname{diag}(\kappa^2(\epsilon+\tfrac12), E/2, 1/2, E/2, 1, 1) > 0. \tag{NU23}$$

Every cross coefficient among these rows is infinitesimal after the displayed R substitution. Literal zero rows are harmless: the coefficient matrix may be strict even when the chosen rows are linearly dependent or vanish.

Use NU10 to add sufficiently large fixed real pivots for the two further rows wB, wB_u, at infinitesimal cost to NU22. The signed scalar terms in NU16 and the remaining normal terms in NU21 have bounded coefficients on these helper rows. A sufficiently large fixed pair of diagonals therefore gives a strictly positive real eight-row Gram, by an ordinary real Schur complement. All entries are finite sums of bounded separated factors. The separated bounded-Gram lemma supplies a finite SOS. Restore the exact positive square remainders in NU10 and the whole excess $(m-\epsilon)(B_c - pV)^2$ in NU7. This gives the entire original parent, with both unrestricted state normals.

All projective testing joins on NU1. Noninfinitesimal Alice tests use CE1–CE10 because C, A are unbounded. For infinitesimal u, the local proof above covers every infinitesimal reciprocal w, including zero. It remains to check finite Bob coordinates v.

For v infinitesimal use the exact FU22–FU23 frame; its source inputs here are only m strong, c_0, a_0 with common positive residue $E/2$, and a_1 positive, all proved in NU3–NU6. For a finite positive unit v use the complete FU6–FU12 bounded parent and real Schur calculation, capping the original m event at a finite real only if m is unbounded. In the present source residue $\beta = d_0 = 0$, $a_1 = E/2$, and, if $\kappa < 1$,

$$m = \frac{\kappa^2}{2(1-\kappa^2)}, \qquad E = \frac{1}{1+\kappa}. \tag{NU24}$$

The determinant polynomial FU14 becomes exactly

$$\mathcal{P}(y,c) = p_{21}c(y-\kappa)^2 + p_{10}y > 0, \qquad p_{21} = E(m+\tfrac{1}{2}), \quad p_{10} = (m+\tfrac{1}{2})E^2, \quad y,c>0. \quad \text{(NU25)}$$

It gives the strict four-term circuit inequality FU18. If $\bar{\kappa} = 1$, then m is unbounded. The normalized limit is

$$\mathcal{P}(y,c)/m \longrightarrow \tfrac{1}{2}c(y-1)^2 + \tfrac{1}{4}y > 0. \quad \text{(NU26)}$$

The Schur coefficients have finite positive limits, so a sufficiently large finite cap preserves strictness, exactly as in FU19–FU20. The whole positive original event excess is restored. These finite-Bob arguments use no lower bound on ν, and do not import the source scope of the FU theorem.

The exact sign transformations FU31 cover negative angles; all displayed frames include literal axes and projective endpoints. Complex states split into real and imaginary parts. No chart normalizes a state coordinate or bounds a state ratio. This completes the all-tests theorem on precisely NU1.

Chapter 189

The doubly infinitesimal endpoint with bounded β/q^2

Component `p1de5b12d1927`
Source: `NEW/c1/c144_small_nu_bounded_beta_rederived.tex`
SHA256: `1de5b12d1927c00334a2816ce0dafbd128057e93db31a5427ab7f843f9a4fa74`
Inventory use: SN complete theorem | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

189.1 The doubly infinitesimal endpoint with bounded β/q^2

Scope. Keep the original entered physical C14.4 source, including its strict gaps, nonnegative weights and prior-complement intersections. In this section

$$\begin{gathered} S=\sigma+L\in \mathrm{Inf}_{>0},\quad \kappa=x/S\in \mathrm{Inf}_{>0},\quad \nu=\sigma/S\in \mathrm{Inf}_{>0},\\ \ell=L/S=1-\nu,\quad q=\nu/\ell,\quad n_s=\beta/q^2 \text{ is bounded.}\end{gathered} \tag{SN1}$$

The conclusion is finite separated tensor SOS for the complete original parent, all testing angles and all real or complex states. In particular no bound is imposed on W/Z, u/w, or any ratio of a testing infinitesimal to a source infinitesimal. This is a source subdomain; β/q^2 unbounded is not claimed here.

Opening-safe source bounds. Use the exact native source dictionary FM2–FM3 and the surviving SE1–SE3 sphere identities. Put

$$c_0=SC(1-r),\quad a_0=SAr,\quad d_s=D/(Sq^2),\quad a_1=\alpha/S,\quad \mu=m/\kappa^2,\quad e=\kappa h/\nu,\quad \tau=h/q. \tag{SN2}$$

The sphere gives, with its original nonnegative coordinates d_0, b_0,

$$d_s=\frac{\ell^2\mathcal{H}d_0^2}{2(P/S)J},\qquad n_s=\frac{\ell^2\mathcal{H}b_0^2}{2\kappa(P/S)\theta},\qquad Jd_s+\kappa\theta n_s=\frac{\ell^2\mathcal{H}}{2(P/S)}. \tag{SN3}$$

Here $P/S, J, B/S, H_0, \mathcal{H}$ tend to 1 on SN1, using SE4–SE6. Thus $b_0^2=O(\kappa\theta n_s)$ is infinitesimal and $d_s\sim 1/2$. The exact identity $e(B/S)=\kappa H_0+\nu\mathcal{H}b_0^2$ now yields $e/\kappa\sim 1$. Consequently

$$h\sim\nu,\quad \theta\sim 1,\quad \tau\sim 1,\quad \mu\sim\tfrac{1}{2},\quad c_0,a_0,a_1\sim\tfrac{1}{2},\qquad D,\alpha,\beta,m\in \mathrm{Inf}_{\geq 0}\,. \tag{SN4}$$

All normalized coefficients in SN2 are bounded; their positive pivots have real lower bounds independent of $r, 1-r$. Literal $\beta=0$ is included. Literal $D=0$ has empty intersection with SN1 by SN3. No comparison of S, q, κ is needed.

The complete reciprocal parent. For infinitesimal Alice coordinate u and reciprocal Bob coordinate w, including either literal zero, use $D_u = 1 + u^2, D_w = 1 + w^2$ and

$$\begin{aligned} F_w/4 = {} & m(W - wX + uY - uwZ)^2 + CD_w(uW - Y)^2 + AD_uD_wY^2 \\ & + (D + \tfrac{1}{2})(uwW + uX - wY - Z)^2 + (\alpha + \tfrac{1}{2})D_u(Y - wZ)^2 \\ & + (\beta + \tfrac{1}{2})D_w(X + uZ)^2 - \tfrac{1}{2}D_uD_wZ^2. \end{aligned} \tag{SN5}$$

Keep both unrestricted state normals

$$X_e = X + ewW, \qquad N_y = (Y - ruW)/\sqrt{Sr(1-r)}. \tag{SN6}$$

Every inverse and radical in these expressions is pure-source or pure-testing. Write $a = u^2, c = w^2, x = \kappa S, \delta = c - q$ and set

$$\begin{gathered} N = \kappa W - \ell uwZ, \quad G = uW/\sqrt{S} - \ell\sqrt{S}\, wZ, \\ B = \sqrt{S}\,\delta Z, \quad B_u = u\delta Z, \quad R = w(a-x)Z = \frac{\kappa\sqrt{S}\,G - uN}{\ell}. \end{gathered} \tag{SN7}$$

The last equality makes R an infinitesimal-coefficient linear combination of the two strict rows N, G for arbitrary states.

Two free helper pivots. The exact polynomial identity

$$\delta^2N^2 + 2\kappa\ell\delta wGB + 2\kappa\ell^2w^2B^2 - \ell^2(wB_u)^2 = (\kappa\delta W)^2 \tag{SN8}$$

buys any fixed positive real pivot on wB_u at infinitesimal bounded-Gram cost on N, G, B. Precisely, multiply SN8 by a chosen positive real constant divided by ℓ^2; its right hand side is already a separated square and every coefficient on the first three rows has zero residue. The B pivot is strict because $d_s \sim 1/2$. Also $(wB)^2 = w^2B^2$ buys any fixed pivot on wB at infinitesimal cost. No positivity of a mixed tensor coefficient or inversion of u, w, W, Z is used in either operation.

Exact scalar dictionary and the $1/\kappa$ cancellations. Put $d = D + 1/2$ and define the following polynomials in a, c:

$$\begin{aligned} A_P &= (1+c)\{c_0(1-r) + a_0r(1+a)\} + S\{d(1-r-e)^2c + (\alpha+\tfrac{1}{2})(1+a)r^2\}, \\ A_N &= m(1+ra+ec)^2 + (\beta+\tfrac{1}{2})e^2c(1+c), \\ K &= m(1+ra+ec) + d(1-r-e) + (\alpha+\tfrac{1}{2})(1+a)r + (\beta+\tfrac{1}{2})e(1+c), \\ F_Z &= D + \alpha c + \beta a + (m + \alpha + \beta + \tfrac{1}{2})ac, \\ J_1 &= K - \ell A_P, \quad L_1 = J_1 - (\ell/\kappa)A_N, \\ H &= F_Z - \ell^2 ScA_P + ac\{(\ell/\kappa)^2A_N - 2(\ell/\kappa)J_1\}. \end{aligned} \tag{SN9}$$

These are the exact graph coefficients at $X = -ewW, Y = ruW$; the negative $dac, e\{2(1-r) - e\}$ contribution has been placed in A_P, not left in A_N. In particular A_N/κ^2 has bounded polynomial coefficients and residue $1/2$.

The actual source satisfies $L_1(x,q) = H(x,q) = H_a(x,q) = H_c(x,q) = 0$ and $H(a,0) = D + \beta a$. Finite monic division therefore gives

$$\begin{gathered} L_1 = (a-x)L_a + \delta L_c, \\ H = S\delta^2(d_s + \kappa n_s + cH_{02}) + (a-x)\delta(-qn_s + cH_{11}) + c(a-x)^2H_{20}. \end{gathered} \tag{SN10}$$

The quotients mean literal polynomial differences at $a = x, c = q$, not division by a mixed tensor. A fully explicit implementation is `c144_small_nu_dictionary.py`. For example L_a is $(L_1(a,c) - L_1(x,c))/(a-x)$, and L_c is the quotient of $L_1(x,c) - L_1(x,q)$ by $c - q$.

Individually $H_{20}, H_{11}, L_c/\kappa$ need not be bounded. The complete scalar uses the following combined coefficients:

$$C_R = H_{20} + 2\ell L_a/\kappa, \quad C_G = \kappa(H_{11} - n_s)/\ell, \quad C_N = (n_s - H_{11})/\ell - 2L_c/\kappa. \tag{SN11}$$

Substitution of $e = \kappa\tau/\ell$ gives the exact cancellations

$$\begin{aligned} C_R &= -\ell\mu r(a\ell r + 2c\kappa\tau + 2\ell), \\ C_N &= n_s/\ell - Sa_0\ell r + (\beta + m + \tfrac{1}{2})\tau^2(c/\ell + 1/\ell^2) + 2\kappa\mu\tau - m\tau^2/\ell, \\ H_{02} &= \kappa\tau^2(\beta + m + \tfrac{1}{2}). \end{aligned} \tag{SN12}$$

These and L_a, C_G are bounded polynomials. To see the latter bound without discarding a source pole, write $A_P = P_0(a) + cP_1(a)$. Direct differentiation in SN9 gives

$$\begin{aligned} \kappa H_{11} = {} & 2\ell^2 P_1(x) + \ell^2 x a_0 r + 2\ell^2\kappa\mu e(1 + 2rx) + \frac{\ell^2(\beta + 1/2)e^2}{\kappa} - 2\ell(m + \beta + \tfrac{1}{2})e \\ & + (c + q)\frac{\ell^2(m + \beta + 1/2)e^2}{\kappa}. \end{aligned} \tag{SN13}$$

Every displayed coefficient is bounded by SN4; e^2/κ is infinitesimal. These exact identities, not estimates of their separate uncanceled terms, justify the normalization.

The entire scalar graph is now

$$\begin{aligned} F_0 = {} & (A_N/\kappa^2)N^2 + A_P G^2 + d_s B^2 + n_s B_u^2 + H_{02}(wB)^2 + C_R R^2 - 2\sqrt{S}\, L_a G R \\ & + C_G G(wB) + C_N N(wB_u). \end{aligned} \tag{SN14}$$

For verification the mixed identity is $c(a - x)\delta Z^2 = \{\kappa G(wB) - N(wB_u)\}/\ell$. Combining this with $uN = \kappa\sqrt{S}\, G - \ell R$ proves SN14 directly from SN9–SN10.

Both complete state normals. Define

$$\begin{aligned} A_X &= -m(1 + ra + ec) + da(1 - r - e) - e(\beta + \tfrac{1}{2})(1 + c), \\ T_X &= (\ell/\kappa)cA_X + \beta - D + c(m + \beta + \tfrac{1}{2}), \\ Y_P &= (1 + c)\{-c_0 + a_0(1 + a)\} + S\{m(1 + ra + ec) - dc(1 - r - e) + (\alpha + \tfrac{1}{2})(1 + a)r\}, \\ Y_C &= \ell Y_P - ma + d - (\alpha + \tfrac{1}{2})(1 + a). \end{aligned} \tag{SN15}$$

Their actual contact values satisfy $T_X(x, q) = Y_C(x, q) = 0$. Use finite polynomial quotients

$$T_X = c(a - x)X_a + \delta\{(D - \beta)/q + cX_c\}, \qquad Y_C = (a - x)Y_a + \delta Y_c. \tag{SN16}$$

Although $X_a, A_X/\kappa$ separately may be unbounded, their complete conductor has

$$\begin{aligned} \widehat{A}_X &= A_X/\kappa - aX_a/\ell = -\kappa\mu - \frac{\tau}{\ell}\{(\beta + \tfrac{1}{2})(1 + c) + mc\}, \\ \widehat{X}_a &= \kappa X_a = \ell\{d(1 - r - e) - mr\}, \qquad X_c = -\tau(m + \beta + \tfrac{1}{2}). \end{aligned} \tag{SN17}$$

The exact full parent is $F_0 + H_X X_e^2 + H_Y N_y^2 + 2H_{XY} X_e N_y + 2L_X X_e + 2L_Y N_y$, with

$$\begin{aligned} H_X &= mc + da + (\beta + \tfrac{1}{2})(1 + c), \\ H_Y &= rc_0(1 + c) + (1 - r)a_0(1 + a)(1 + c) + Sr(1 - r)\{ma + dc + (\alpha + \tfrac{1}{2})(1 + a)\}, \\ H_{XY} &= -uw\sqrt{Sr(1 - r)}(m + d), \\ L_X &= w\widehat{A}_X N + \frac{uw\sqrt{S}}{\ell}\widehat{X}_a G + d_s\sqrt{S}\, quB - n_s qB_u + wX_c(wB_u), \\ L_Y &= \sqrt{r(1 - r)}\{Y_P G + \sqrt{S}\, Y_a R + Y_c(wB)\}. \end{aligned} \tag{SN18}$$

All coefficients are bounded without a lower bound on r or $1 - r$. Y_P has zero residue, as do the direct coefficients in L_X. The face-safe term $n_s qB_u$ is $q\sqrt{n_s}(\sqrt{n_s}B_u)$, so literal $n_s = 0$ needs no division. Both Y_a, Y_c are bounded polynomials by SN15; they have no inverse of q after the finite division. This verifies the full first-normal conductor rather than only the scalar graph.

Strict bounded-Gram allocation and joins. Use the strict base rows $(N, G, B, X_e, N_y, \sqrt{n_s}B_u)$, omitting the last zero row if $n_s = 0$. Their diagonal residues are $(1/2, 1/2, 1/2, 1/2, 1/2, 1)$. Terms containing R are infinitesimal matrix perturbations by SN7. Choose fixed sufficiently large real pivots on wB, wB_u to absorb all bounded cross coefficients in SN14 and SN18, and buy those pivots by SN8. All costs have zero residue. The resulting finite matrix is bounded and has strictly positive definite real residue; the surviving strict bounded-Gram lemma in `inherited_r19/parent_reference.tex` proves finite SOS. The budget remainders and every original normal term have been accounted for exactly once. This proves all infinitesimal u, w, including the literal axes and arbitrary states.

Noninfinitesimal Alice tests use the unchanged CE1–CE10 adjacent C, A theorem, since C, A are unbounded by SN4. For infinitesimal Alice and Bob outside the reciprocal infinitesimal chart, $\overline{s^2} \neq \overline{\ell} = 1$ in normalized homogeneous Bob coordinates. The exact surviving SE10–SE12 full-parent certificate applies, because e is infinitesimal. Sign gauges and real/imaginary state splitting are the original ones; the local polynomial proof itself accepts either sign of u, w. Thus every test and every state on SN1 is covered.

Verification and present status. This is a fresh proof body. Its focused checker verifies the complete original parent, both state normals, the actual source contact jets, the normalized budget and each canceled coefficient in SN12–SN18. Those identities corroborate the analytical source bounds and strict-Gram argument above. Status at creation: candidate awaiting independent lead review; no global C14.4 completion is asserted.

Chapter 190

The unbounded-h/q part of the doubly small endpoint

Component `peaa56c1a3c7a`
Source: `NEW/c1/c144_unbounded_h_ratio_rederived.tex`
SHA256: `eaa56c1a3c7a7e7b15ba788b1c3f8de1a728747a89b4efaea38ebb630b09230b`
Inventory use: UN1–UN21 | C14_ENTIRE_ORIGINAL_PARENT_GRAPH; precise subordinate uses retained below

190.1 The unbounded-h/q part of the doubly small endpoint

Exact scope. Retain the full original entered C14.4 source, all its strict gaps and prior-complement intersections. Suppose S, κ, ν are positive infinitesimals, $\ell = 1 - \nu$, $q = \nu/\ell$, and

$$\tau = h/q \text{ is unbounded}, \qquad \theta = 1 - h > 0, \quad e = \kappa h/\nu \in \mathrm{Inf}, \quad \zeta = \ell/\kappa. \tag{UN1}$$

This section proves finite separated tensor SOS for the complete parent at every projective test and every state on UN1. The opening $\theta \in \mathrm{Inf}$ is included. There is no state-ratio restriction or source/testing relative-rate assumption.

Normalized source estimates and a canceled radial mass. Keep $c_0 = SC(1-r)$, $a_0 = SAr$, $d_s = D/(Sq^2)$, $a_1 = \alpha/S$, $P_0 = P/S$, and the exact SE2–SE3 source sphere. Define

$$d = D + \tfrac{1}{2}, \quad n = \beta\theta^2 + 2Dh + \tfrac{1}{2}h^2 > 0, \quad z_n = \kappa n/q^2, \quad A_{00} = m\zeta^2 - n/q. \tag{UN2}$$

Here n is a scalar coefficient, not an omitted event weight. It is bounded even when β is unbounded. The exact source gives

$$\begin{aligned} m\zeta^2 = \frac{\tau(1+x)(1+\sigma r)}{2MP_0}, \qquad & m\zeta \in \mathrm{Inf}, \quad D, \alpha, m \in \mathrm{Inf}, \quad c_0, a_0, a_1 \sim \tfrac{1}{2}, \\ e\tau = \frac{\ell\{eH_0 + h\mathcal{H}b_0^2\}}{B/S} & \text{ is bounded}, \\ z_n = \frac{\ell^2\mathcal{H}\theta b_0^2}{2P_0} + \frac{\ell^2(eH_0 + h\mathcal{H}b_0^2)}{2(B/S)} & + 2\kappa S d_s h. \end{aligned} \tag{UN3}$$

Consequently d_s, z_n are bounded and $d_s + z_n \sim 1/2$; at least one is a positive bounded unit. This includes literal $D = 0$. Literal $\beta = 0$ has no points on UN1, because $\beta \geq 0$ has the exact boundary equation $\tau(B/S) = \ell H_0$ when $\beta = 0$.

For completeness, the following finite source division proves the bound on A_{00} without dividing an error by θ or q. Put $U = (1+x)(1+\sigma r)$ and $M_* = \ell(1+xr) - \kappa\theta = \ell M$, and form the bounded source polynomial

$$T = \ell JU + JM_*P_0 - JM_*\kappa\theta(J - \sigma) - 2M_*U.$$

In this expression P_0, J, U, M_* are their exact native polynomial formulas in κ, ν, S, h, r, with $\ell = 1-\nu$. Thus

$$\begin{gathered} T = \kappa T_\kappa + \ell\nu S T_\nu, \qquad T_\nu = r - 1 - \nu - \nu S(1-r)(r+\ell), \\ T_\kappa = \{T - T|_{\kappa=0}\}/\kappa \quad \text{is a bounded source polynomial,} \\ A_{00} = \frac{\theta\ell H_0}{2P_0} + \frac{eT_\kappa + h\ell S T_\nu}{2JMP_0}. \end{gathered} \tag{UN4}$$

The quotient is a finite monic polynomial division in a pure source variable. All denominator factors are positive bounded units. Hence A_{00} is bounded and $A_{00} - \theta/2$ is infinitesimal. Moreover $n/q = m\zeta^2 - A_{00} \sim \tau/2$ is positive unbounded. If θ is infinitesimal, then $n \sim 1/2$ and $\kappa/q = e\ell/h$ is infinitesimal. These facts supply the two source subcases used below.

The actual circuit and three reserved positive rows. Use the complete original reciprocal quarter-parent SN5. For infinitesimal $u, w \geq 0$, including zero, retain the unrestricted normals

$$N_x = X + huZ, \qquad N_y = (Y - ruW)/\sqrt{Sr(1-r)}. \tag{UN5}$$

Put $a = u^2, c = w^2, x = \kappa S, \delta = c - q$ and

$$\begin{gathered} N_0 = W/\zeta - uwZ, \quad G = uW/\sqrt{S} - \ell\sqrt{S}\, wZ, \\ B = \sqrt{S}\delta Z, \quad B_u = u\delta Z, \quad R = w(a-x)Z = \sqrt{S}\, G/\zeta - uN_0, \quad G_h = h(a - xc/q)Z. \end{gathered} \tag{UN6}$$

Define A_1, A_2, A_3, A_4 by BN4 with β replaced by the positive n, and set

$$\Psi_n = \sum_i A_i^2 - 4nuwWZ/(\zeta q), \qquad E_1 = A_1 - A_2, \quad E_2 = A_3 - A_4, \quad E_3 = A_1 - A_3. \tag{UN7}$$

The explicit separated AM–GM square decompositions in BN6 give

$$\Psi_n - \tfrac{1}{4}(E_1^2 + E_2^2 + E_3^2) \in \Sigma T^2. \tag{UN8}$$

To verify the simultaneous reserve, average BN6 with the pairing $\Psi_n - (A_1 - A_3)^2 - (A_2 - A_4)^2 \in \Sigma T^2$. For the latter pairing the products A_1A_3, A_2A_4 are nonnegative; if $WZ < 0$, the additional signed cross term is a positive separated monomial. Thus UN8 holds for both signs of WZ, as well as its zero face. No radical of a general mixed sum is used.

Exact scalar subtraction and radial cancellation. The complete scalar graph at $N_x = N_y = 0$ is

$$F_0 = m(1+ra)^2W^2 + A_P P^2 + F_Z Z^2 - 2KuwWZ, \quad P = uW/\sqrt{S}, \tag{UN9}$$

where

$$\begin{aligned} A_P &= (1+c)\{c_0(1-r) + a_0 r(1+a)\} + S\{d(1-r)^2 c + (\alpha + \tfrac{1}{2})(1+a)r^2\}, \\ K &= m\theta(1+ra) + d(1-r)(1+ha) + (\alpha + \tfrac{1}{2})(1+a)r, \\ F_Z &= D + \alpha c + na + \{m\theta^2 + \alpha + (\beta + \tfrac{1}{2})\theta^2\}ac + dh^2a^2. \end{aligned} \tag{UN10}$$

Subtract Ψ_n, put

$$\begin{gathered} A = m\zeta^2(1+ra)^2 - n/q, \quad A'_P = A_P - z_n c/\ell^2, \\ K' = K - 2n/(\zeta q), \quad J' = K' - \ell A'_P, \quad L' = J' - A/\zeta, \\ H' = F_Z - na - nxc/q - \ell^2 S c A'_P + ac(A - 2\zeta J'). \end{gathered} \tag{UN11}$$

The exact residual graph is $AN_0^2 + A'_P G^2 - 2\zeta uwZL'N_0 + H'Z^2$. At the actual source contact $L'(x,q) = H'(x,q) = H'_a(x,q) = H'_c(x,q) = 0$. Also $H'(a,0) = D + dh^2a^2$. Finite monic contact division gives

$$\begin{gathered} L' = (a-x)L_a + \delta L_c, \\ H' = (D + dh^2x^2)\delta^2/q^2 + (a-x)\delta(-2dh^2x/q + cH_{11}) + (a-x)^2(dh^2 + cH_{20}). \end{gathered} \tag{UN12}$$

The otherwise possible third-c quotient is exactly zero. Define $P_1(a)$ to be the coefficient of c in A_P. The complete combined coefficients have short bounded formulas:

$$\begin{aligned} L_a &= m\theta r + dh(1-r) + (\alpha+\tfrac12)r - \ell\{(1+c)a_0 r + S(\alpha+\tfrac12)r^2\} \\ &\quad - m\zeta r\{2 + r(a+x)\}, \\ C_R &= H_{20} + 2\zeta L_a = -m\zeta^2 r(2+ra), \\ C_G &= H_{11}/\zeta = 2\ell P_1(x) + \ell x a_0 r - 2z_n/\ell, \\ C_N &= -H_{11} - 2\zeta L_c = -Sa_0\ell^2 r. \end{aligned} \tag{UN13}$$

Only C_R may be unbounded separately. Its entire contribution must be combined before estimating: $A + aC_R = A_{00}$ exactly. Using $R = \sqrt{S}\,G/\zeta - uN_0$ therefore gives

$$\begin{aligned} F_0 &= \Psi_n + A_{00}N_0^2 + \widetilde{A}_P G^2 + 2u\sqrt{S}\,(L_a - C_R/\zeta)N_0 G \\ &\quad + d_s B^2 + dG_h^2 + C_G G(wB) + C_N N_0(wB_u), \\ \widetilde{A}_P &= A'_P + SC_R/\zeta^2 - 2SL_a/\zeta \sim \tfrac12. \end{aligned} \tag{UN14}$$

Every coefficient in UN14 is bounded after this combination: $C_R/\zeta = -m\zeta r(2+ra)$ is infinitesimal and $C_R/\zeta^2 = -mr(2+ra)$ is bounded. The N_0G coefficient is infinitesimal. No estimate of an unbounded isolated radial coefficient is used.

Complete X conductor and its exact positive completion. The full N_x hard coefficient is $H_X = mc + da + (\beta + 1/2)(1+c)$. Its complete half-linear coefficient, after using the actual contact identity, is

$$\begin{aligned} L_X &= -m\zeta w N_0 + d\tau a(wN_0 + B_u) + uw\sqrt{S}\{d(1-r-e) - mr\}G + t_0 B_u, \\ t_0 &= \frac{D + h/2 - \beta\theta}{q} = D/q + \frac{\ell H_0 - \kappa\tau(J-\sigma)}{2P_0} \sim \tfrac12. \end{aligned} \tag{UN15}$$

The exact source identity in the second line cancels the potential $1/q$ pole. Here $D/q = Sq\,d_s$ is infinitesimal. The single unbounded τ term is removed with the actual square dG_h^2, not estimated. Namely

$$\begin{aligned} G_h &= -\tau u(wN_0 + B_u) + ew\sqrt{S}\,G, \quad N_x + \tau(wN_0 + B_u) = X + ewW, \\ &dG_h^2 + daN_x^2 + 2d\tau a(wN_0 + B_u)N_x \\ &\quad = da(X + ewW)^2 - 2de\tau uw\sqrt{S}\,(wN_0 + B_u)G + de^2 Sw^2 G^2. \end{aligned} \tag{UN16}$$

The remainder has bounded infinitesimal coefficients on N_0, G, wB_u because $e\tau$ is bounded. Retain the entire positive square on the right and also the whole packet $\beta(1+c)N_x^2$. The remaining N_x diagonal is $mc + (1+c)/2 \sim 1/2$. The remaining L_X is UN15 with its $d\tau a$ term removed; all its direct coefficients are infinitesimal except the bounded coefficient t_0 on B_u. The construction thus includes arbitrarily large β.

The unrestricted Y normal and cross coefficient. Put

$$\begin{aligned} Y_P &= (1+c)\{-c_0 + a_0(1+a)\} + S\{m(1+ra) - d(1-r)c + (\alpha+\tfrac12)(1+a)r\}, \\ Y_C &= \ell Y_P - m\theta a + d(1+ha) - (\alpha+\tfrac12)(1+a). \end{aligned} \tag{UN17}$$

The actual source has $Y_C(x,q) = 0$, so finite monic division gives $Y_C = (a-x)Y_a + \delta Y_c$ with bounded polynomial coefficients. The complete remaining entries are

$$\begin{aligned} L_Y &= \sqrt{r(1-r)}\{Y_P G + \sqrt{S}\,Y_a R + Y_c(wB)\}, \\ H_Y &= rc_0(1+c) + (1-r)a_0(1+a)(1+c) \\ &\quad + Sr(1-r)\{ma + dc + (\alpha+\tfrac12)(1+a)\} \sim \tfrac12, \\ H_{XY} &= -uw\sqrt{Sr(1-r)}(m+d) \in \mathrm{Inf}\,. \end{aligned} \tag{UN18}$$

In particular Y_P is infinitesimal. Every coefficient is bounded through $r = 0$ and $r = 1$ openings; no inverse of r or $1 - r$ remains after the displayed pure-source normal scaling. UN14–UN18 account for all terms of SN5, including both independent state normals.

Free pivots, including the θ-infinitesimal opening. Reserve the three actual circuit rows in UN8. In either sphere chart, wB is free at infinitesimal cost: use $\sqrt{d_s}B$ when d_s is a unit, and otherwise use the exact identity

$$wB = -\frac{w + \sqrt{q}}{\sqrt{z_n}} E_2 - \frac{w(w + \sqrt{q})}{\ell} G. \tag{UN19}$$

If θ is a unit, A_{00} already supplies a strict N_0 pivot. Since q/n is infinitesimal, the exact identity

$$B_u = -(w + \sqrt{q})\sqrt{q/n}\, E_1 - (w + \sqrt{q}) N_0 \tag{UN20}$$

buys any fixed B_u pivot at infinitesimal cost. Then wB_u is free by multiplication by the pure-testing infinitesimal w.

If θ is infinitesimal, first buy the missing N_0 pivot from the actual circuit, using

$$N_0 = \sqrt{q/n}(E_2 - E_1) - \frac{w - \sqrt{q}}{\sqrt{n}} E_3 + \frac{\sqrt{\kappa/q}}{\ell}\, wG. \tag{UN21}$$

All coefficients are infinitesimal: $n \sim 1/2$ and κ/q is infinitesimal by UN4. Identity UN21 holds without requiring z_n to be a unit; the apparent source factor in UN19 has canceled exactly before estimating. After this purchase, UN20 buys B_u, and multiplication by w buys wB_u. Thus the order of purchases involves no circular use of a missing pivot. The bounded, possibly signed A_{00} in this source subcase has zero residue and is absorbed after the strict N_0 pivot is bought. No inverse of θ is introduced.

Final allocation and unchanged testing joins. Use the strict rows G, N_x, N_y, the applicable N_0 pivot, and fixed retained fractions of $E_1, E_2, E_3, \sqrt{d_s}B$ wherever nonzero. Purchase sufficiently large real pivots on B_u, wB, wB_u by UN19–UN21. Every purchase has infinitesimal Gram cost on the preceding strict rows. All signed terms in UN14–UN18 are then either infinitesimal matrix perturbations or bounded couplings to those freely purchased helper pivots. A fixed sufficiently large choice makes the complete residue matrix strictly positive definite. The separated strict bounded-Gram lemma proves finite SOS. Restore the positive packets retained in UN8 and UN16, including $\beta(1 + c)N_x^2$; nothing from the original parent is omitted.

This proves all infinitesimal u, w, their literal zero faces and arbitrary states. Noninfinitesimal Alice tests use CE1–CE10 because C, A are unbounded. The other Bob tests use the surviving SE10–SE12 unmatched certificate, since e is infinitesimal and $\overline{\ell} = 1$. The original sign gauges and real/imaginary state split are exact; the circuit argument treats both signs of WZ. These joins give the claimed all-tests theorem on UN1.

Status and corroboration. This fresh proof is pending independent lead review. Its focused exact checker verifies the original full parent, the source division UN4, all actual contact jets, the complete scalar cancellation, both unrestricted normal conductors, and the actual circuit helper identities. No former completion marker is used as a proof premise.

Chapter 191

All testing charts for the unit-κ, ν, ℓ source

Component `p3f02c51074be`
Source: `NEW/c1/c144_unit_kappa_all_tests_rederived.tex`
SHA256: `3f02c51074beaad6ba4c99cea8fb710f31d0573f6bd28a7a49bb2457f076cbf7`
Inventory use: FU1–FU31 | FU6-FU23 exact finite-Bob parent/circuit identities, used only with freshly checked NU source limits | Fresh FU1-FU31 full unmatched and endpoint-testing proof; source endpoint restrictions retained

191.1 All testing charts for the unit-κ, ν, ℓ source

Exact scope and conclusion. Retain the original physical C14.4 source, its strict source gaps, and its original prior-complement intersections. Assume

$$S = \sigma + L \in \mathrm{Inf}_{>0}, \qquad \kappa = x/S, \quad \nu = \sigma/S, \quad \ell = L/S \quad \text{are positive bounded units,} \qquad \nu + \ell = 1. \tag{FU1}$$

Together with the matched theorem UKR1–UKR21, the argument below proves that the complete original parent is a finite separated tensor SOS for every projective test and every state on FU1. It includes arbitrary openings of $M, \theta, r, 1-r$, literal $D = 0$ and $\beta = 0$, and independent source and testing infinitesimals. It makes no claim at the ν- or ℓ-infinitesimal source endpoints.

All normalizations are pure-source or pure-testing operations. All real Schur complements below prove strictness of a real residue matrix; no such inverse is used as a tensor-ring coefficient. The strict bounded-Gram lemma is the one in the surviving `inherited_r19/parent_reference.tex`.

The full parent and opening-safe source bounds. Write $D_u = 1 + u^2, D_v = 1 + v^2$, and divide the parent by 4:

$$\begin{aligned} F/4 = {} & m(vW - X + uvY - uZ)^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW - Y + uvX - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ & + (\beta + \tfrac{1}{2})D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \end{aligned} \tag{FU2}$$

Keep the exact FM2–FM3 source formulas, with $c_0 = SC(1-r)$, $a_0 = SAr$, $d_0 = D/S$, and $a_1 = \alpha/S$. These four coefficients and $\beta\theta$ are bounded, with no inverse of the possibly small M or θ in their normalized formulas. The D/S numerator has its factor S canceled exactly.

The following letters refer only to real residues:

$$A_* = \nu - \kappa h = \nu E, \quad R = 1 - A_*, \quad \nu = A_* + \kappa h, \quad \ell = 1 - A_* - \kappa h. \tag{FU3}$$

The exact source yields

$$\begin{gathered} E \ge 1/18, \qquad h - A_* \ge 0, \qquad A_*^2 - \kappa(h - A_*) \ge 0, \qquad R - \kappa \ge 0, \\ c_0 = a_0 = E/2, \qquad d_0 = \{A_*^2 - \kappa(h - A_*)\}/2, \\ a_1 = \frac{\ell\{\kappa h + (R - \kappa)A_*\}}{2\nu}, \qquad 2\beta\theta = h - A_*. \end{gathered} \tag{FU4}$$

In particular $h \ge \nu/(1+\kappa)$, so h, E, a_1, c_0, a_0 have positive real lower bounds. The actual source has m bounded below by a positive real, while C, A are unbounded. Here m can be unbounded, and so can β. They cannot both be unbounded: m unbounded forces $R = \kappa$, whence $\ell = \kappa\theta$ makes θ a unit; β unbounded forces $\theta = 0$, whence $R - \kappa = \ell$ makes M a unit. For bounded m, β their real values are

$$m = \frac{\kappa^2 h}{2\nu(R - \kappa)}, \qquad \beta = \frac{h - A_*}{2(1 - h)}. \tag{FU5}$$

These formulas are used only when the indicated real denominators are positive. Every estimate below for the actual parent uses bounded normalized source coefficients before taking residues.

A complete finite-Bob frame, including the state normals. Assume u is infinitesimal and v is finite. We may cap m and β by finite pure-source values $m_c > 0, \beta_c \ge 0$ and retain the complete excesses in their original event squares. When a weight is bounded, keep it unchanged. When it is unbounded, the sufficiently large real cap is chosen below. Define

$$B_0 = X + uZ, \quad N_y = \frac{Y - ruW}{\sqrt{Sr(1-r)}}, \qquad T = \kappa W,\ U = uZ,\ P = \frac{uW}{\sqrt{S}},\ Q = \sqrt{S}Z. \tag{FU6}$$

All six rows are legal separated expressions, and $TU = \kappa PQ$ holds exactly. No restriction is placed on their values or on state ratios. Set $a = u^2$, $s_y = \sqrt{Sr(1-r)}$, and

$$\begin{aligned} M_0 &= (v/\kappa)(1 + ra)T - B_0, \\ A_P &= D_v\{c_0(1-r) + a_0 r(1+a)\} \\ &\quad + S\{(D + \tfrac{1}{2})(1-r)^2 + (\alpha + \tfrac{1}{2})(1+a)r^2v^2\}. \end{aligned} \tag{FU7}$$

The entire capped parent is $F_0 + 2L_yN_y + H_yN_y^2$, where

$$\begin{aligned} F_0 &= m_cM_0^2 + A_PP^2 + (1+a)\{d_0(1+a)v^2 + a_1\}Q^2 + \tfrac{1}{2}(1+a)v^2U^2 \\ &\quad - 2(D + \tfrac{1}{2})(1+a)v^2B_0U - (1+a)v\{1 + 2D(1-r) + 2\alpha r\}PQ \\ &\quad + \{(\beta_c + \tfrac{1}{2})D_v + (D + \tfrac{1}{2})av^2\}B_0^2 + 2(D + \tfrac{1}{2})(1-r)uv\sqrt{S}\,B_0P, \end{aligned} \tag{FU8}$$

$$\begin{aligned} L_y &= m_cuvs_yM_0 + \sqrt{r(1-r)}D_v\{-c_0 + a_0(1+a)\}P \\ &\quad + S\sqrt{r(1-r)}\{-(D + \tfrac{1}{2})(1-r) + (\alpha + \tfrac{1}{2})(1+a)rv^2\}P \\ &\quad - (D + \tfrac{1}{2})uvs_yB_0 + \sqrt{r(1-r)}v(1+a)(D - \alpha)Q, \\ H_y &= rc_0D_v + (1-r)a_0(1+a)D_v \\ &\quad + Sr(1-r)\{m_cu^2v^2 + D + \tfrac{1}{2} + (\alpha + \tfrac{1}{2})(1+a)v^2\}. \end{aligned} \tag{FU9}$$

These are exact identities in the two unrestricted original state coordinates X, Y; B_0 is not set to zero. Their coefficients are bounded across all source openings. In L_y every coefficient has zero residue, since c_0, a_0 have the same residue, and the other terms contain u or S. The N_y pivot has residue $ED_v/2 > 0$. Thus it suffices to make the remaining five-row real residue strict.

Finite nonzero unmatched Bob residue. Take $v > 0$ with a positive finite residue different from $b = \sqrt{\ell/\nu}$. The complete real residue of FU8–FU9 is

$$\begin{aligned} &m_c\{(v/\kappa)T - B_0\}^2 + qD_vB_0^2 - v^2B_0U + \tfrac{1}{2}v^2U^2 - vPQ \\ &\quad + \tfrac{E}{2}D_vP^2 + (d_0v^2 + a_1)Q^2 + \tfrac{E}{2}D_vN_y^2, \qquad q = \beta_c + \tfrac{1}{2}. \end{aligned} \tag{FU10}$$

For the moment use the uncapped finite values m, β from FU5. Eliminating B_0 in this real matrix, and using $TU = \kappa PQ$ only to name the adjusted cross coefficient, gives

$$a_TT^2 + a_UU^2 + a_PP^2 + a_QQ^2 - 2fPQ, \tag{FU11}$$

where, for $c = v^2$ and $H = m + q(1 + c)$,

$$\begin{gathered} a_T = \frac{mq(1+c)c}{\kappa^2 H}, \quad a_U = \frac{c(2H-c)}{4H}, \quad a_P = E(1+c)/2, \quad a_Q = d_0c + a_1, \\ f = \frac{v}{2}(1 + mc/H). \end{gathered} \tag{FU12}$$

The replacement of the real Schur cross term TU by κPQ is implemented in the actual bounded Gram by a real multiple of the exact zero $TU - \kappa PQ$. The elimination of B_0 itself is only a real strictness test. Every diagonal is positive, including when D or β is zero. The reduced determinant in a formal positive real variable y is

$$\Delta(y, c) = (a_T\kappa^2 + a_Py)(a_Uy + a_Q) - f^2y. \tag{FU13}$$

It obeys $4H\Delta/(1 + c) = \mathcal{P}(y, c)$ with

$$\mathcal{P} = p_{22}y^2c^2 + p_{21}y^2c + p_{12}yc^2 + p_{11}yc + p_{10}y + p_{02}c^2 + p_{01}c, \tag{FU14}$$

$$\begin{aligned} &p_{22} = E\beta, && p_{21} = E(m + q), \\ &p_{12} = 2m\beta + 2qEd_0, && p_{11} = -m - q + 2E\{(m + q)d_0 + qa_1\}, \\ &p_{10} = 2(m + q)Ea_1, && p_{02} = 4mqd_0, \quad p_{01} = 4mqa_1. \end{aligned} \tag{FU15}$$

All coefficients except possibly p_{11} are nonnegative, and p_{21}, p_{10}, p_{01} are strictly positive. Direct substitution of FU3–FU5 gives the exact three contact identities

$$\mathcal{P}(\kappa, \ell/\nu) = \mathcal{P}_y(\kappa, \ell/\nu) = \mathcal{P}_c(\kappa, \ell/\nu) = 0. \tag{FU16}$$

These finite rational identities can also be checked without radicals. For positive real y, c,

$$\frac{\mathcal{P}(y, c)}{yc} = p_{22}yc + p_{21}y + p_{12}c + p_{11} + p_{10}/c + p_{02}c/y + p_{01}/y. \tag{FU17}$$

As a real function of $(\log y, \log c)$, this is strictly convex: each nonconstant summand has nonnegative Hessian, and $p_{21}y + p_{10}/c$ alone has positive definite Hessian. FU16 says its value and gradient vanish at the contact. It is consequently positive at every other positive point. This real argument is only a way to determine a finite residue matrix, not a use of logarithms or mixed inverses in the tensor ring.

Since $c \neq \ell/\nu$, FU13 is positive for all $y > 0$. Minimizing its quotient by y gives

$$f < \kappa\sqrt{a_Ta_U} + \sqrt{a_Pa_Q}. \tag{FU18}$$

Indeed the minimum of $a_Pa_Uy + a_T\kappa^2a_Q/y$ is $2\kappa\sqrt{a_Ta_Ua_Pa_Q}$, attained at a positive y; equality in FU18 would give a zero of FU13. Choose a real λ with $|\lambda| < \sqrt{a_Ta_U}$ and $|f - \kappa\lambda| < \sqrt{a_Pa_Q}$. Adding the exact zero $-2\lambda(TU - \kappa PQ)$ makes both 2×2 blocks in FU11 strictly positive. Restoring the positive real B_0 pivot and the N_y pivot proves strictness of the complete six-row residue. All actual Gram entries were bounded in FU8–FU9, so the strict bounded-Gram lemma gives the tensor SOS.

Unbounded m or β in the unmatched chart. The preceding strictness persists under a sufficiently large finite cap in the one possible unbounded weight. This assertion has an opening-safe direct verification. If $m \to \infty$, divide FU14–FU15 by m before taking the real limit. Then

$$(p_{22}, p_{21}, p_{12}, p_{11}, p_{10}, p_{02}, p_{01})/m \longrightarrow (0, E, 2\beta, -1 + 2Ed_0, 2Ea_1, 4qd_0, 4qa_1). \tag{FU19}$$

At $R = \kappa$, the three contact identities FU16 remain exact for this limit polynomial. The strict convexity proof FU17 still applies because its y and $1/c$ coefficients are positive. The Schur coefficients tend to

$$a_T \to q(1+c)c/\kappa^2, \qquad a_U \to c/2, \qquad f \to v(1+c)/2, \tag{FU20}$$

with a_P, a_Q unchanged. Therefore FU18 is strict at the limit, and continuity makes it strict for a sufficiently large finite real m_c.

If $\beta \to \infty$, divide instead by q before taking the limit at $h = 1$. The corresponding coefficient vector is

$$(E, E, 2m + 2Ed_0, -1 + 2E(d_0 + a_1), 2Ea_1, 4md_0, 4ma_1), \tag{FU21}$$

again with the three exact contact identities and the two strictly positive convexity coefficients. Now $a_T \to mc/\kappa^2$, $a_U \to c/2$, $f \to v/2$. A sufficiently large finite β_c therefore works. All the other normalized actual coefficients have their real residues with bounded unit denominators; no $O(S)$ term was divided by M or θ. Restore the complete excesses $(m - m_c)(vW - X + uvY - uZ)^2$ and $(\beta - \beta_c)D_v(X + uZ)^2$. These caps are used only on the unmatched chart and need not vanish at the actual contact.

Infinitesimal or zero Bob coordinate. If v is infinitesimal, use FU8–FU9 with the six rows

$$(B_0, T_v, U_v, P, Q, N_y), \qquad T_v = vT, \quad U_v = vU. \tag{FU22}$$

Cap m at any sufficiently small fixed positive real $m_c \le m$, and discard no excess. One may set $\beta_c = 0$ and retain its whole event. The exact coefficients remain bounded: the B_0U coefficient is now a multiple of vB_0U_v, and M_0 becomes $(1 + ra)T_v/\kappa - B_0$. Their real residue is

$$m_c(T_v/\kappa - B_0)^2 + \tfrac{1}{2}B_0^2 + \tfrac{1}{2}U_v^2 + \tfrac{E}{2}(P^2 + N_y^2) + a_1Q^2 > 0. \tag{FU23}$$

This also covers literal $v = 0$. The rows need not be independent coordinates on the state space.

Infinitesimal reciprocal Bob coordinate. Put $w = 1/v$ and use the homogeneous parent w^2F at $w = 0$ as well as for nonzero infinitesimal w. The complete signed core in the quarter-parent is $K_w/2$, with

$$\begin{aligned} K_w = {} & u^2H^2 - 2u^2HY - 2uHW + 2uwHX + u^2w^2W^2 \\ & + 2u^2wWX - 2uw^2WY + (1 + u^2 + w^2)(X^2 + Y^2) - 2uwXY, \qquad H = wZ. \end{aligned} \tag{FU24}$$

This is the exact CE6 full core, so all state normals are present. FU4 gives the real alternative

$$d_0 + \kappa\beta\theta = (\nu E)^2/2 > 0. \tag{FU25}$$

Thus either D/S has a positive real lower bound, or β has one. Both cases allow independent rates for u, w, S.

If β is strong, use the four rows (W, X, Y, U), $U = uZ$, in FU24 by replacing uH with wU. Every coefficient is bounded and the core residue is $X^2 + Y^2$. The exact m event is $(W - wX + uY - wU)^2$, and the β event is $D_w(X + U)^2$. Positive finite caps give the strict residue

$$m_cW^2 + \tfrac{1}{2}(X^2 + Y^2) + \beta_c(X + U)^2 > 0. \tag{FU26}$$

All other event packets and both excesses remain whole.

If D/S is strong, choose positive real c, d, m_c with $c \le c_0, a_0$, $d \le D/S$, and $m_c \le m$. Select exactly

$$C_c = c/[S(1-r)], \qquad A_c = c/(Sr), \qquad D_c = dS. \tag{FU27}$$

Retain all excesses, including the whole $A_c u^2 D_w Y^2$. Writing $Y = ruW + \sqrt{Sr(1-r)}N_y$, the other two selected C, A terms have the exact harmonic identity

$$C_c D_w (uW - Y)^2 + A_c D_w Y^2 = c D_w (P^2 + N_y^2). \tag{FU28}$$

Use six rows (W, V, X, P, N_y, Q) with $V = uwZ$ and the exact monomial identity $VW = wPQ$. Replacing uH by V in FU24, the potentially non-small cross term $-2VW$ is exactly $-2wPQ$. Every other off-diagonal core coefficient is infinitesimal after $Y = \sqrt{S}\{rP + \sqrt{r(1-r)}N_y\}$; its residue is $V^2 + X^2$. The selected m and D events are the complete squares

$$m_c(W - wX + uY - V)^2, \qquad d\{uw\sqrt{S}W + u\sqrt{S}X - w\sqrt{S}Y - Q\}^2. \tag{FU29}$$

The full selected residue is strictly positive:

$$m_c(W - V)^2 + \tfrac{1}{2}(V^2 + X^2) + c(P^2 + N_y^2) + dQ^2 > 0. \tag{FU30}$$

The strict bounded-Gram lemma applies to the exact complete forms, and all unselected positive packets are restored. Both reciprocal proofs include literal $w = 0$ and $u = 0$.

Exhaustion of tests and exact joins. If u is noninfinitesimal, C, A are unbounded by FU4. The surviving two-adjacent-events theorem CE1–CE10 in `common_remote.tex` therefore applies to the original full parent and all states. Its hypotheses require only nonnegative event weights and these two unbounded coefficients; no small-κ source statement is imported.

For infinitesimal u, every positive projective v is either infinitesimal, a finite unit, or has infinitesimal reciprocal. FU22–FU30 cover the two end cases. A finite unit has either the matched residue, covered by UKR1–UKR21, or a different residue, covered by FU10–FU21. The transformations

$$(u, Y, Z) \mapsto (-u, -Y, -Z), \qquad (v, X, Z) \mapsto (-v, -X, -Z) \tag{FU31}$$

preserve FU2 exactly and handle both signs. Literal axes and projective endpoints were included in the displayed raw frames. Complex states split into real and imaginary parts. No proof in this exhaustion bounds, normalizes, or relates the testing-state ratios W/Z, X/Z, or Y/Z.

Consequently every testing chart is covered on exactly FU1, with every original source predicate retained. The remaining source endpoints are separate from this conclusion.

Chapter 192

A fresh matched-Bob completion for unit κ, ν, ℓ

Component `pd26b89cdb3f0`
Source: `NEW/c1/c144_unit_kappa_matched_rederived.tex`
SHA256: `d26b89cdb3f024a6d512d1b05c1181c3d7e4ec63a18ce77c2ff3820685f65784`
Inventory use: UKR1–UKR21 | UKR1-UKR21 matched testing chart, root analytical review favorable | Fresh UKR1-UKR21 analytical candidate

192.1 A fresh matched-Bob completion for unit κ, ν, ℓ

Scope. This candidate retains the entire original physical C14.4 source, including its strict gaps and prior complements. Assume $S \in \mathrm{Inf}_{>0}$ and $\kappa = x/S, \nu = \sigma/S, \ell = L/S$ are positive bounded units, with $\nu + \ell = 1$. The testing chart is $u \in \mathrm{Inf}$ and finite v whose absolute value has the same residue as $b = \sqrt{\ell/\nu}$. All real or complex states are allowed. This section does not claim the unmatched Bob tests or either source endpoint. Neither a previous missing R21 theorem nor a mixed inverse is used.

The dictionary FM2–FM3 remains an exact source identity on this larger domain, but its small-κ estimates FM4 are replaced below. Set $j = \sqrt{\nu\ell}$ and retain the rows

$$N_x = X + huZ, \quad N_y = (Y - ruW)/\sqrt{Sr(1-r)}, \qquad T = \kappa W,\ U = uZ,\ P = uW/\sqrt{S},\ Q = \sqrt{S}Z,$$

$$N = T - jU, \quad G = P - jQ, \quad \delta = v - b, \quad B = \delta Q, \quad B_u = \delta U, \quad R_Z = (u^2 - x)Z. \tag{UKR1}$$

The radial and cross identities FM8 hold unchanged. Their coefficients on R_Z remain infinitesimal because S, u are infinitesimal and $\kappa, 1/j$ are bounded.

A sharper exact angular budget. For any fixed real $0 < \rho < 1$, the monomial identity $TU = \kappa PQ$ gives

$$\frac{\delta^2}{j^2(1-\rho^2)}N^2 + \frac{2\kappa\delta}{j(1+\rho)}GB + \frac{2\kappa}{1+\rho}B^2 - B_u^2 = \frac{(\delta T - \rho j B_u)^2}{j^2(1-\rho^2)}. \tag{UKR2}$$

Its first two coefficients have zero residue. Thus a fixed cB_u^2 pivot costs $q\kappa cB^2$ in residue, where $q = 2/(1+\rho) > 1$ can be chosen arbitrarily close to one. Every inverse in UKR2 is a pure source unit or a positive real. There is no inverse of δ.

The actual source through both coefficient openings. The source has $1/18 \le E = 1 - \kappa h/\nu < 1$ and κ is bounded because $1/\zeta$ is bounded. The exact $\beta\theta$ identity gives a positive real lower bound on h. Consequently m has a positive real lower bound, although it may be unbounded. Let

$$R_s = 1 + xr, \quad p = \frac{\kappa\theta}{\ell R_s} \in (0,1), \qquad \mu_c = m(1-p) = \frac{\kappa^2 h(1+x)(1+\nu Sr)}{2\nu\ell P_0 R_s}. \tag{UKR3}$$

The quantity μ_c is a positive bounded unit through $M \to 0$. All of $SC(1-r), SAr, D/S, \alpha/S, \beta\theta$ are bounded without inverses of $r, 1-r, \theta, D, \beta$; this follows directly from FM3 after the explicit factor S in the D/S numerator is canceled. Their limits are evaluated below.

Only in the real residue calculation use the abbreviations

$$A = \nu - \kappa h = \nu E, \quad R = 1 - A, \quad d = h - A. \tag{UKR4}$$

Here $A, R, h, \nu, \ell, \kappa$ are positive units, and the original nonnegative weights and positive M imply

$$d \geq 0, \quad A^2 - \kappa d \geq 0, \quad R - \kappa \geq 0, \qquad \nu = A + \kappa h, \quad \ell = 1 - A - \kappa h. \tag{UKR5}$$

Indeed in residue $2\beta\theta = d$, $2D/S = A^2 - \kappa d$, and $\ell M = R - \kappa$. These are additive residue identities; no relative error estimate is asserted at a vanishing opening.

Retain unbounded weights in actual contact-zero rows. Define

$$V = v(W + uY), \quad B_0 = X + uZ, \quad N_m = B_0 - pV. \tag{UKR6}$$

At the actual source contact $u = \sqrt{x}$, $v = b$, $W/Z = j\sqrt{S/\kappa}$, $X = -h\sqrt{x}Z$, $Y = r\sqrt{x}W$, both N_m and N_x vanish. This is an identity at the source contact, not an equation imposed on independent tests.

If m is bounded, set $m_c = m$. If it is unbounded, choose a positive real m_c sufficiently large, as justified below. Similarly keep $\beta_c = \beta$ when β is bounded, and choose a sufficiently large positive real β_c when it is unbounded. An unbounded positive field element exceeds either fixed real cap. Retain the entire positive excesses

$$(m - m_c)N_m^2 + (\beta - \beta_c)D_v N_x^2. \tag{UKR7}$$

The remaining parent is exact, with its m part replaced by

$$m_0 B_0^2 - 2m_1 B_0 V + m_2 V^2, \qquad m_0 = m_c, \quad m_1 = \mu_c + m_c p, \quad m_2 = \mu_c(1+p) + m_c p^2. \tag{UKR8}$$

This follows from the identity $m(V - B_0)^2 - (m - m_c)(B_0 - pV)^2$ and $\mu_c = m(1-p)$. Every coefficient of UKR8 is bounded. The residual β terms are exactly

$$\beta_c D_v N_x^2 + 2(\beta\theta)D_v N_x uZ + (\beta\theta)\theta D_v u^2 Z^2. \tag{UKR9}$$

All coefficients of UKR9 are bounded. The complete cross terms are retained, and both original excesses in UKR7 are restored once at the end.

The complete bounded scalar and normal frame. Apply the FM9–FM19 polynomial construction to this capped parent with the following explicit replacements: in A_T replace m/κ^2 by m_2/κ^2; in A_U replace its $m\theta^2$ term by $m_0\theta^2$; in K replace its $m\theta(1+ra)$ term by $m_1\theta(1+ra)$. In the complete first-normal formulas replace

entry	replacement
H_x first term	m_0
H_y term mu^2v^2	$m_2u^2v^2$
H_{xy} term m	m_1
L_x coefficient $-m/\kappa$	$-m_1/\kappa$
L_U terms $m\theta - (mj/\kappa)v(1+ra)$	$m_0\theta - (m_1 j/\kappa)v(1+ra)$
Y_P term $mv^2(1+ra)$	$m_2v^2(1+ra)$
Y_Q term $-m\theta a$	$-m_1\theta a$.

(UKR10)

Include $\beta_c D_v$ in H_x and omit the separately retained $\beta D_v N_x^2$ term of FM15. All other entries are unchanged. These replacements follow by expanding UKR8–UKR9 in the two unrestricted state normals. They are not a scalar-only operation.

The graph scalar and both complete first normals still vanish to the orders required by FM11 and FM19: the subtracted rows in UKR7 vanish at the actual source contact. Their squares therefore change neither the value nor any first derivative there. The same finite monic divisions supply the exact bounded FM13 scalar and FM18–FM19 full normal conductors. The Y normal has zero-residue cross coefficients, since $SC(1-r)$ and SAr both have residue $E/2$ and the remaining Y coefficients contain S or u^2. Its diagonal residue is $E/(2\nu) > 0$.

The only nontrivial real residue uses (N_x, N, G, B, B_u). Writing

$$H_* = h - \tfrac{1}{2} + \tfrac{\theta}{2}(\ell + \kappa h), \qquad r_* = H_* - j^2 m_2/\kappa^2, \tag{UKR11}$$

its exact entries are

$$\begin{aligned} H_c &= \begin{pmatrix} m_0 + (\beta_c + 1/2)/\nu & -(m_1/\kappa)b \\ -(m_1/\kappa)b & (m_2/\kappa^2)b^2 \end{pmatrix}, \qquad A_G = E/(2\nu), \\ z &= \begin{pmatrix} -j(m_1/\kappa + E) \\ -r_*/\nu \end{pmatrix}, \qquad g_* = E\ell - m_1\theta - \tfrac{1}{2} + 2m_2\ell/\kappa + \kappa r_*/\nu, \\ & H_B = \nu(m_1\theta + \tfrac{1}{2}). \end{aligned} \tag{UKR12}$$

Thus the residue form is $(N_x, N)H_c(N_x, N)^t + A_G G^2 + 2g_* GB + H_B B^2 + 2z^t(N_x, N)^t B_u$ before the UKR2 pivot is added. In particular the $\kappa r_*/\nu$ contribution to g_* is retained; it is not infinitesimal in the present source sector.

The exact real contact curvature and all source faces. First take an interior real source, $\theta > 0$ and $R > \kappa$, without capping, so

$$m = \frac{\kappa^2 h}{2\nu(R-\kappa)}, \qquad \beta = \frac{h - A}{2\theta}, \qquad m_0 = m_1 = m_2 = m. \tag{UKR13}$$

The two hard pivots and the angular pivot obey the exact identities

$$\begin{aligned} (N_x, N)H_c(N_x, N)^t &= \frac{m}{\kappa^2}(bN - \kappa N_x)^2 + \frac{R}{2\nu\theta}N_x^2, \\ J_* &= H_B - \frac{g_*^2}{A_G} - \kappa z^t H_c^{-1} z \\ &= \frac{AR + \kappa(h-A)}{2AhR}\{4A^2hR - \kappa(Rh - A)^2\}. \end{aligned} \tag{UKR14}$$

All inverses in UKR14 are used only in a real finite matrix calculation to prove strictness. They are not inverses in the tensor ring. The crucial bracket has the positive decomposition

$$\begin{aligned} 4A^2hR - \kappa(Rh - A)^2 = {} & A^3R(4 - A) + A^4(R - \kappa) \\ & + A^2R(h - A)(3 + A + 2\kappa) \\ & + R^2(h - A)\{A^2 - \kappa(h - A)\}. \end{aligned} \tag{UKR15}$$

Its first term is a positive unit and all remaining terms are nonnegative by UKR5. This includes the literal $D = 0$ and $\beta = 0$ faces and both limiting openings.

For completeness the strictness is uniform toward those openings. Put $\mu = m/\kappa^2$ and retain H_* from UKR11 with uncapped coefficients. Then the two entries of $H_c^{-1}z$ are

$$z_x = -\frac{2\nu\theta}{R}\left(jE + \frac{\kappa H_*}{\nu b}\right), \qquad z_N = \frac{\kappa}{b}z_x + \nu - \frac{H_*}{\mu\ell}. \tag{UKR16}$$

They stay bounded. Also

$$g_* = -\frac{2A^2 + Ah\kappa + A\kappa - A - h\kappa}{2\nu} \tag{UKR17}$$

stays bounded, while A_G and J_* have positive real lower bounds near any fixed allowed residue point. UKR14 gives a positive real lower bound on H_c because μ and $R/(2\nu\theta)$ have positive lower bounds and b is a unit. Completing the real source quadratic form with the bounded coefficients UKR16–UKR17 therefore gives a uniform positive lower bound on

$$Q_{\mathrm{phys}} = (N_x, N)H_c(N_x, N)^t + A_G G^2 + 2g_* GB + H_B B^2 + 2\sqrt{\kappa}\, z^t(N_x, N)^t B. \tag{UKR18}$$

This is an auxiliary real matrix used to choose a diagonal budget. It does not impose $B_u = \sqrt{\kappa}B$, or any relation between source and testing angles, in the tensor certificate.

Why finite caps preserve a strict real residue. When m, β are bounded no cap is needed and UKR18 is already strict. If m is unbounded, $R - \kappa = 0$ in residue; since ℓ is a unit and $\ell = R - \kappa h$, θ is then a unit, so β is bounded. If β is unbounded, $\theta = 0$ in residue, and $R - \kappa = \ell > 0$, so m is bounded. Thus at most one unbounded penalty is present.

For the m opening, the real contact-zero linear form in UKR18 is

$$n_m = N_x - (pb/\kappa)N - (pj/\sqrt{\kappa})B, \tag{UKR19}$$

and $p \to 1$. Its N coefficient is a unit. UKR8 shows that subtracting the whole mn_m^2 leaves a bounded coefficient matrix with a well-defined limit at the opening. That limit is strict on $\ker n_m$: approach the residue point by real $\kappa < R$ with A, h fixed, adjust any kernel vector continuously to the exact kernel using the unit N coefficient, and apply the uniform lower bound from UKR18. The parameter inequalities UKR5 are preserved when κ is decreased. A real symmetric matrix strict on the kernel of a nonzero linear form becomes strictly positive definite after adding a sufficiently large fixed real multiple of the form's square; this follows by a real block Schur complement. Choose m_c that large in UKR7.

For the β opening the same argument uses the exact penalty $(\beta/\nu)N_x^2$ and its kernel $N_x = 0$. Approach $h = 1$ from below with A, κ fixed, which also preserves UKR5, and choose β_c sufficiently large. This is a calculation on real residue matrices, not a mixed-field limiting certificate. The actual capped coefficients have those residues because UKR3, UKR8 and UKR9 have bounded denominators through the original independently varying S, M, θ openings. In particular no estimate divides an $O(S)$ error by M or θ.

We have proved that the actual caps can be chosen so that the capped matrix UKR18 is strictly positive definite. It follows that $H_c \succ 0$, $A_G > 0$, and

$$H_B - g_*^2/A_G - \kappa L > 0, \qquad L = z^t H_c^{-1} z \geq 0. \tag{UKR20}$$

Complete separated SOS allocation. Choose $\rho < 1$ sufficiently close to one and then a positive real c so that

$$L < c < \frac{H_B - g_*^2/A_G}{q\kappa}, \qquad q = 2/(1+\rho). \tag{UKR21}$$

Such choices exist by UKR20. Use c times UKR2 to introduce the cB_u^2 pivot. The full real residue on (N_x, N, G, B, B_u) is strictly positive: eliminate H_c to leave $(c - L)B_u^2$, and eliminate A_G to leave $(H_B - q\kappa c - g_*^2/A_G)B^2$. The N_y pivot is $E/(2\nu) > 0$ and its cross coefficients have zero residue. Every other full normal and radial term has already been allocated by the exact bounded FM constructions and UKR10. The strict separated bounded-Gram lemma supplies a finite tensor SOS. Restore the square on the right of UKR2 and both complete excess packets UKR7. This is the full original parent, not its restriction to a state graph.

The original sign transformations of FM5 handle negative u, v, and the proof includes $u = 0$. Complex states split into real and imaginary parts. Arbitrary state ratios and independent source and testing rates are allowed throughout. Unmatched unit-κ Bob tests and the ν- or ℓ-infinitesimal source endpoints remain outside this theorem.

Chapter 193

The complete positive-scale C1 reciprocal terminal

Component `p98f24864b071`
Source: `NEW/global/c1_reciprocal_full_parent_REDERIVED.tex`
SHA256: `98f24864b0712df08a74a0cf16c9b7e8412282a9a8aa6494e7d2f657042b38f3`
Inventory use: CR1-CR39 full C1 reciprocal all-tests theorem, with AU19 proof-use audit | CR1-CR39 complete source entry, all local states, full normal restoration and projective outer join

193.1 The complete positive-scale C1 reciprocal terminal

Theorem and exact scope. Let the canonical C1 source be physical, with an actual finite noncommuting zero contact, and normalize $d = 1$. Suppose every source coefficient is bounded, $\lambda = n = 0$, C, A, α are positive bounded units, $m > 0$ and $D \geq 0$ are infinitesimal, and $\beta \geq 0$ is arbitrary bounded. Retain the original canonical source gaps and intersections. Then the entire parent has a finite sum-of-squares representation in $K_c \otimes_{\mathbb{R}} K_t$ for every independent projective test and every real or complex state. The assertion includes literal $D = 0$ and literal $\beta = 0$ when permitted by the actual contact; they cannot both vanish.

We use $O(1)$ and $o(1)$ for bounded and infinitesimal elements of one of the two fields, and $a \asymp b$ for positive source quantities whose ratio is a bounded unit. A bounded tensor Gram with a strictly positive real residue is a sum of squares by the actual separated strict-Gram lemma. Its rows need not be bounded. No order on the tensor product, mixed inverse, or cancellation of a determinant multiplier is used below.

193.1.1 Source entry before any state-ratio division

Write the actual contact angles as $u_c = k > 0$, $b_c = 1/v_c = p > 0$, and put $U = k^2$, $P = p^2$. The actual C1 source balances SP2–SP3 give U, P positive infinitesimals and

$$D/P^2 = O(1), \qquad \beta U/P^2 = O(1), \qquad m/U^2 \asymp 1 + \beta/P. \tag{CR1}$$

Indeed the two stationarity equations are

$$\begin{aligned} U^2(a_6 + \beta a_7/P) &= m(a_1 + a_2 D/P), \\ a_3 &= P^{-2}(mDa_2/U + \beta m a_4 + Da_5 + \beta a_7 U), \end{aligned}$$

where every a_i is a positive bounded unit. Positivity bounds each term of the second equation; then $D/P = o(1)$ and the parentheses in the first give CR1. In particular

$$m/U = o(1), \qquad D/P = o(1), \qquad \beta U/P = o(1). \tag{CR2}$$

The original source partition proves U, P infinitesimal in exactly the present m, D weak branch. Only those displayed source balances, not its absent terminal references, are an input here.

At a physical noncommuting contact the signed core is negative, and the exact core Schur identity SE9 gives $W_cZ_c > 0$. Normalize $W_c = 1$ and write $Z_c = q > 0$. Define

$$Q = q^2, \quad L = PQ, \qquad r = \frac{k}{pq}, \quad y = \frac{Y_c}{pq}, \quad x = \frac{X_c}{pq^2}. \tag{CR3}$$

We now prove these divisions are bounded wherever they are used. In the reciprocal quarter-parent CR9 below the hard X, Y matrix at the source has residue $\mathrm{diag}(\bar\beta+1/2, H_0)$, where $H = C+A+\alpha+1/2$ is a positive unit. Eliminate this two-by-two block inside the source field. Its Schur block on W, Z satisfies

$$\frac{S_{WW}}{U} \sim a := C - C^2/H > 0, \qquad -\frac{S_{WZ}}{kp} \sim b_* := 1/2 + C\alpha/H > 0. \tag{CR4}$$

Here and below a displayed positive expression in an asymptotic formula is evaluated in residue. To check CR4 without a shape assumption, the original hard couplings are

$$\begin{aligned} P_{XW} &= p\{-m + (D+1/2)U\}, & P_{YW} &= k\{m - C(1+P) - (D+1/2)P\}, \\ P_{XZ} &= k\{\beta - D + P(m+\beta+1/2)\}, & P_{YZ} &= p\{D - \alpha - U(m+\alpha+1/2)\}, \\ P_{WZ} &= -(m+D+1/2)kp. \end{aligned}$$

CR2 shows that the X corrections in CR4 vanish in residue and the Y corrections are respectively C^2/H and $C\alpha/H$. The kernel equation gives $q = S_{WW}/(-S_{WZ})$, so r is a positive bounded unit.

The scaled X, Y kernel equations in the unknowns x, y have bounded entries and a diagonal positive residue. For example their X equation is exactly

$$\begin{aligned} 0 = {} & \{(\beta+1/2)(1+P) + Pm + (D+1/2)r^2PQ\}x \\ & + r\{\beta - D + P[\beta + 1/2 - m(y-1) + (D+1/2)(r-y)]\} - m/Q. \end{aligned} \tag{CR5}$$

The quantity $m/Q = (m/U)r^2P$ is bounded before this equation is used. The y coefficient in CR5 is infinitesimal; the x coefficient in the scaled Y equation is $O(PQ)$ and its y pivot is $H + o(1)$. Thus x, y are bounded. Taking the Y and Z kernel equations in residue, using CR1–CR2, yields

$$Hy \sim Cr + \alpha, \qquad C(r-y) \sim 1/2, \qquad \alpha(1-y) \sim r/2. \tag{CR6}$$

Consequently $y, r-y, 1-y$ are positive bounded units. Equivalently their real residues satisfy

$$C_0 = \frac{1}{2(r_0 - y_0)}, \quad A_0 = \frac{1 + r_0 - y_0}{2y_0}, \quad \alpha_0 = \frac{r_0}{2(1-y_0)}. \tag{CR7}$$

For the last equation in CR6, divide the original Z kernel equation by Pq. Its extra X contribution is $O(x\beta Q + xPQ)$; CR1 and the already proved unit r give $\beta Q = O(P)$, while $PQ = U/r^2$ is infinitesimal. All the omitted terms therefore vanish in residue. This proves CR6 in the unbounded-q case as well.

In CR5, $D = o(P)$ and $m/Q = o(P)$. Its numerator for $-x$ is the sum of $r\beta$ and a positive-unit multiple of P, up to $o(P)$. Hence the following bounds hold on the whole source domain:

$$\begin{aligned} &L = PQ = o(1), \qquad \beta Q/P = O(1), \qquad -x \asymp P + \beta, \qquad r + x \text{ is a positive unit}, \\ &m \asymp PQ^2(P+\beta), \qquad m/(PQ) = O(PQ + \beta Q) = o(1), \qquad xQ = o(1). \end{aligned} \tag{CR8}$$

The statement about $r + x$ follows directly from CR5: $x/r = -\beta/(\beta + 1/2) + o(1)$.

193.1.2 The entire scalar and all normal coefficients

Use $b = 1/v$, $D_u = 1 + u^2$, $D_b = 1 + b^2$. The full reciprocal quarter-parent is

$$\begin{aligned} F = {} & m(W - bX + uY - ubZ)^2 + CD_b(uW - Y)^2 + AD_uD_bY^2 \\ & + (D+1/2)(ubW + uX - bY - Z)^2 + (\alpha+1/2)D_u(Y - bZ)^2 \\ & + (\beta+1/2)D_b(X + uZ)^2 - D_uD_bZ^2/2. \end{aligned} \tag{CR9}$$

All further local identities are expansions of this full expression. In the chart $W = 1$, $Z = z$ put

$$u_g = rbz, \qquad X_g = xbz^2, \qquad Y_g = ybz, \qquad V = b^2, \quad T = z^2. \tag{CR10}$$

Its entire scalar restriction is

$$\Phi = m + DT + hVT + \mathcal{C}V^2T + EVT^2 + \mathcal{F}V^2T^2 + GV^3T^2 + JV^2T^3, \tag{CR11}$$

where

$$\begin{aligned}
\mathcal{C} &= Ay^2 + (C + D + 1/2)(r - y)^2, \\
E &= \beta(r + x)^2 + x^2/2 - 2Drx, \qquad G = Ar^2y^2, \qquad J = (D + 1/2)r^2x^2, \\
\mathcal{F} &= Ar^2y^2 + (\alpha + 1/2)r^2(y - 1)^2 + \beta(r + x)^2 + m[r(y - 1) - x]^2 \\
&\quad + (2D + 1)rx(r - y) + rx + x^2/2, \\
h &= Ay^2 + C(r - y)^2 + \alpha(y - 1)^2 + y^2/2 - r - 2D(r - y) + 2m[r(y - 1) - x].
\end{aligned} \tag{CR12}$$

In particular $\mathcal{C}, G$ are positive units, $E \asymp \beta + P^2 > 0$, $J \geq 0$, and all these coefficients are bounded. If β is infinitesimal, $\mathcal{F}$ is a positive unit, with residue $r_0^2(1 + r_0 - y_0)/2$. For unit β, $\mathcal{F}$ can have either sign; it is not discarded or assumed positive.

Actual contact and its two angular/state jets give $\Phi(P, Q) = \Phi_V(P, Q) = \Phi_T(P, Q) = 0$. Therefore

$$\begin{aligned}
D &= P^2\mathcal{C} - PQE + P^3QG - P^2Q^2J, \\
m &= PQ^2E + P^2Q^2\mathcal{F} + P^3Q^2G + 2P^2Q^3J, \\
h &= -2P\mathcal{C} - QE - 2PQ\mathcal{F} - 3P^2QG - 2PQ^2J.
\end{aligned} \tag{CR13}$$

Set

$$\delta = D/P^2 \geq 0, \qquad \theta = EQ/P > 0. \quad \delta, \theta = O(1), \qquad \delta + \theta = \mathcal{C} + GL - JQ^2 = \mathcal{C} + o(1), \tag{CR14}$$

where $L = PQ$. The bounds follow either from CR1 and CR8 or from CR13; $JQ^2 = O((xQ)^2) = o(1)$. At least one of δ, θ is a unit. Define

$$R = z(V - P), \quad K = VT - PQ, \quad M = V^2T + (P/Q)VT^2 + P^2Q - 3PVT.$$

The exact complete balanced scalar is

$$\begin{aligned}
\Phi = \delta R^2 + \theta M + \mathcal{F}K^2 + G\{PK^2 + VzRK\} \\
+ J\{(QR - zK)^2 + 2QK^2\}.
\end{aligned} \tag{CR15}$$

The entire J line is retained as positive squares.

Here are full, rather than selected, normal formulas. Let L_u, L_X, L_Y be half the first derivatives of CR9 at CR10. Then

$$L_u = bzA_u(V, T), \quad L_Y = bzA_y(V, T), \quad L_X = b\{-m + TB_X(V, T)\}, \tag{CR16}$$

where

$$\begin{aligned}
A_u &= a_{00} + a_{10}V + a_{01}T + a_{11}VT + a_{21}V^2T + a_{12}VT^2, \\
A_y &= e_{00} + e_{10}V + e_{11}VT + e_{21}V^2T, \\
B_X &= c_0 + c_1V + c_2VT, \\
c_0 &= -Dr + \beta(r + x) + x/2, \\
c_1 &= (D + 1/2)r(r - y) + (\beta + 1/2)(r + x) + m[x - r(y - 1)], \\
c_2 &= (D + 1/2)r^2x.
\end{aligned} \tag{CR17}$$

Every displayed coefficient is bounded. The only bounds beyond this needed below are

$$a_{01} = \beta(r + x) - Dx = O(E), \quad a_{12} = (D + 1/2)rx^2, \quad a_{12}Q = O(1), \qquad c_2 = O(P + E). \tag{CR18}$$

They follow from CR8 and CR12. The other coefficients are given unambiguously by expanding CR9; the checker compares the entire polynomials, not a support truncation.

State and angle stationarity give $A_u(P,Q) = A_y(P,Q) = 0$ and $B_X(P,Q) = m/Q$. Put $E_T = bz(T-Q)$. Finite monic division therefore gives exactly

$$\begin{aligned} L_i &= A_{i,V}(V,T)bR + B_i(T)E_T,\ B_u(T) && = a_{01} + P\{a_{11} + Pa_{21} + a_{12}(T+Q)\},\\ B_y(T) &= P(e_{11} + Pe_{21}),\\ L_X &= b\{[m/(PQ) + c_2T]K + z[c_1 - m/(PQ)]R\}. \end{aligned} \tag{CR19}$$

The first line is for $i = u, y$ and $A_{i,V} = [A_i(V,T) - A_i(P,T)]/(V-P)$, an exact polynomial. In particular, using $P(T-Q) = K - zR$, each of L_u, L_Y, L_X is a bounded linear combination of

$$bR, \qquad bzK, \qquad b[m/(PQ) + c_2T]K, \qquad EE_T. \tag{CR20}$$

Every occurrence of K outside the last product has an infinitesimal coefficient. CR19 remains literal at $b = 0$ or $z = 0$.

193.1.3 The cubic conductors and the bounded source-ratio chart

All radicals in this subsection are products of radicals in the separate fields. Set

$$\mathsf{A} = (V^2T)^{1/3}, \quad \mathsf{B} = ((P/Q)VT^2)^{1/3}, \quad \mathsf{C} = (P^2Q)^{1/3}, \qquad \rho = (P/Q)^{1/3}.$$

Because $\mathsf{ABC} = PVT$,

$$M = \tfrac{1}{2}(\mathsf{A}+\mathsf{B}+\mathsf{C})[(\mathsf{A}-\mathsf{B})^2 + (\mathsf{A}-\mathsf{C})^2 + (\mathsf{B}-\mathsf{C})^2]. \tag{CR21}$$

Let $\mathcal{Z}$ be its nine roots multiplied by $\sqrt{\theta}$, so that $\theta M = \frac{1}{2}\sum_{j=1}^{9}\mathcal{Z}_j^2$. For nonnegative testing b, z, the following exact identities hold:

$$\begin{aligned} bR &= b\sqrt{\mathsf{A}}(\mathsf{A}-\mathsf{B}) + \rho b^{1/3}z^{2/3}\sqrt{\mathsf{A}}(\mathsf{A}-\mathsf{C}),\\ EE_T &= \sqrt{E}\,z\,[\sqrt{\theta}\sqrt{\mathsf{B}}(\mathsf{B}-\mathsf{A})]\\ &\quad + \theta^{1/3}E^{1/6}b^{2/3}z^{1/3}[\sqrt{\theta}\sqrt{\mathsf{B}}(\mathsf{B}-\mathsf{C})]. \end{aligned} \tag{CR22}$$

Negative signs are handled by the exact testing gauges; absolute values give the same monomial identities if desired. If θ is a unit, the first line has infinitesimal coefficients on $\mathcal{Z}$: $\rho = (E/\theta)^{1/3}$ is bounded and b is infinitesimal. If δ is a unit, bR instead has the infinitesimal coefficient b on the strict row R. The second line has infinitesimal coefficients whenever E is infinitesimal or z is infinitesimal. Its displayed coefficients use no comparison of a source rate with a testing rate.

Suppose first that actual q is bounded and β is infinitesimal. Use K and $\mathcal{Z}$, with R added when δ is a unit. If only θ is a unit, retain δR^2 as a separate positive square. In CR15 the K pivot $\mathcal{F}$ is a positive unit. The only signed extra term is $GVzRK = Gbz(bR)K$, an infinitesimal Gram correction by CR22. The term GPK^2 is nonnegative. Thus CR15 has a strict residue Gram on the stated roots. CR19–CR22 give infinitesimal lifts of all three complete first normals to these same roots.

If β is a unit, CR8 forces Q infinitesimal and E a unit. The only chart still requiring CR10 will be z infinitesimal. The second line of CR22 still has infinitesimal coefficients. When δ is a unit, use the exact identity

$$2QM + TR^2 - K^2 = (K - zR)^2 + 2QR^2. \tag{CR23}$$

Choose a positive real a with $\overline{\mathcal{F}} + a > 1$. Subtracting a times the corresponding zero identity changes the first three terms of CR15 into

$$(\delta - aT + 2aQ)R^2 + (1 - 2aP/E)\theta M + (\mathcal{F}+a)K^2 + a(K - zR)^2. \tag{CR24}$$

The first three pivots have strictly positive residues; both costs T and P/E are infinitesimal. This handles the signed $\mathcal{F}$ exactly.

When instead θ is a unit, ρ is also a unit and

$$K = \rho^{-1}\{\sqrt{\mathsf{A}}\,[\sqrt{\mathsf{A}}(\mathsf{B}-\mathsf{C})] + \sqrt{\mathsf{C}}\,[\sqrt{\mathsf{C}}(\mathsf{A}-\mathsf{C})]\}. \tag{CR25}$$

Both coefficients on $\mathcal{Z}$ are infinitesimal. Thus the entire signed $\mathcal{F}K^2$ is an infinitesimal correction on the strict circuit Gram. The G cross term and every first normal are controlled by CR22 and CR25; δR^2 and the J line remain positive squares. No inverse of δ is used.

193.1.4 An unbounded actual state ratio in the same testing chart

Now let actual q be unbounded. CR8 and CR12 imply

$$\beta/P = O(1/Q) = o(1), \quad -x \asymp P, \quad m \asymp L^2, \quad E/P = o(1), \quad JQ^2 = O(L^2) = o(1). \tag{CR26}$$

Here β is infinitesimal and $\mathcal{F}$ is a positive unit. In the testing chart $W = 1$, z bounded, the exact relation

$$\theta M = \theta R^2 + (E/P)(K^2 - zRK) \tag{CR27}$$

makes CR15 a strict two-row Gram on R, K, plus its whole positive J line. Its residue is $\operatorname{diag}(\overline{\delta+\theta}, \overline{\mathcal{F}})$. Every correction coefficient is infinitesimal: E/P, P and Vz are infinitesimal. In CR19, B_u/P and B_y/P are bounded, because $a_{01}/P = O(\beta/P + Dx/P)$ and $a_{12}Q$ are bounded. Using $PE_T = bzK - bz^2R$ shows that all three full normals are infinitesimal linear combinations of R, K.

193.1.5 The complementary testing state chart

Use $Z = 1$, $W = w$ infinitesimal, and write $a = u/r$. The graph is

$$b = aw, \qquad X = xa, \qquad Y = yaw. \tag{CR28}$$

The entire scalar is

$$\Phi' = D + mw^2 + Ea^2 + ha^2w^2 + \mathcal{C}a^4w^4 + \mathcal{F}a^4w^2 + Ga^6w^4 + Ja^4. \tag{CR29}$$

First assume actual q bounded and β infinitesimal. If δ is a unit, use the three strict rows $P, a^2w, \sqrt{E}\,a$. The sole signed cross in CR29 is $ha^2w^2 = (h/P)w\,P(a^2w)$, and h/P is bounded by CR13. Every other term is positive or has the positive-unit $\mathcal{F}$ pivot. If θ is a unit, $P/E = Q/\theta$ is bounded; use instead $\sqrt{E}\,a, a^2w$ and retain D, mw^2 whole. The same signed term is the infinitesimal correction $(h/E)w^2(\sqrt{E}\,a)^2$.

Here are the complete complementary first normals, which also prove the bounds for unbounded actual q. Define

$$R' = a^2w^2 - P, \qquad K' = w(a^2 - L). \tag{CR30}$$

Half the b, Y derivatives of the original full CR9 at CR28 have the respective forms

$$L_i' = aw(d_{i0} + d_{i1}a^2 + d_{i2}a^2w^2 + d_{i3}a^4w^2), \qquad i = b, Y, \tag{CR31}$$

with bounded coefficients. At the actual contact $a^2 = L, w^2 = 1/Q$, they satisfy $d_{i0} + Ld_{i1} + Pd_{i2} + PLd_{i3} = 0$. Consequently the entire, exact divisions are

$$L_i' = (d_{i1} + Pd_{i3})aK' + (d_{i2} + a^2d_{i3})awR'. \tag{CR32}$$

The full X derivative is

$$L_X' = a(c_0 + c_2a^2 - mw^2 + c_1a^2w^2), \qquad c_0 + c_2L - m/Q + c_1P = 0. \tag{CR33}$$

For bounded Q, c_0/P and m/P are bounded, and $c_2 \in PB_c + EB_c$ by CR18. The products aK' and awR' are infinitesimal linear combinations of P, a^2w. Expanding CR33 gives infinitesimal coefficients on $P, a^2w, \sqrt{E}\,a$. If θ is a unit, use $P/E = O(1)$: $c_0/\sqrt{E} = o(1)$, $c_2 = O(E)$ and $m/E = O(1)$, giving

the same assertion on the two rows $\sqrt{E}\,a, a^2w$. These are pure-source bounded quotients; no testing w is inverted.

For unbounded actual q, CR29 has the exact alternative form

$$\begin{aligned}\Phi' &= \delta R'^2 + \theta M' + \mathcal{F}K'^2 + G\{PK'^2 + a^2wR'K'\}\\ &\quad + J\{a^4(Qw^2-1)^2 + 2QK'^2\},\\ M' &= a^4w^4 + (P/Q)a^2 + P^2Qw^2 - 3Pa^2w^2.\end{aligned} \tag{CR34}$$

Define nonnegative elementary radicals

$$\mathsf{A}' = |aw|^{4/3}, \quad \mathsf{B}' = (P/Q)^{1/3}|a|^{2/3}, \quad \mathsf{C}' = (P^2Q)^{1/3}|w|^{2/3}.$$

They give the same nine-square cubic identity for M'. The exact conductor needed for the complete normals is

$$|a|R' = |a|\sqrt{\mathsf{A}'}(\mathsf{A}' - \mathsf{C}') + \sqrt{L}\sqrt{\mathsf{B}'}(\mathsf{A}' - \mathsf{B}'). \tag{CR35}$$

Its coefficients are infinitesimal. If δ is a unit, use R', K' and retain the circuit as positive squares. If θ is a unit, use its scaled nine roots and K', retaining $\delta R'^2$ whole. In either case CR34 is strict: its sole signed cross is $Gaw(aR')K'$, controlled by CR35, and its last line is retained as squares.

CR26 implies $c_2/P = O(1)$ and $c_0/P = m/(PQ) - Qc_2 - c_1 = O(1)$. The full exceptional normal has the exact division

$$\begin{aligned}L'_X = &-[c_0/P + (c_2/P)a^2]\,aR'\\ &+[(c_2/P)(a^2+L) + c_0/P + c_1]\,awK'.\end{aligned} \tag{CR36}$$

Together with CR32, this uses only bounded multiples of aR', awR', aK'. Each has infinitesimal coefficients on the strict soft rows by CR35. This also proves the result at literal $w = 0$.

193.1.6 All higher normals and the remaining unit-beta states

The preceding scalar and first-normal formulas certify the entire parent. Here are the required hard blocks and the justification for every higher term. Normalize the testing state so its coordinates are bounded with at least one unit. At the reciprocal angle corner $u, b = o(1)$ the state residue hard block is $(\bar{\beta} + 1/2)X^2 + H_0Y^2$. If X or Y is a unit, the entire parent has positive real residue and the strict-element lemma applies immediately. Otherwise X, Y are infinitesimal and one of W, Z is a unit.

The exact sign gauges make $u, b \geq 0$. If $WZ \leq 0$, the signed core is a nonnegative pure-testing scalar by the full core Schur identity SE9 and its literal homogeneous endpoints, so its source multiple and all event packets are separated squares. Otherwise choose either $W = 1$, $z = Z$ bounded, or $Z = 1$, $w = W$ infinitesimal. Both have X, Y infinitesimal.

In the $W = 1$ chart, set $N_u = u - rbz$, $N_X = X - xbz^2$, $N_Y = Y - ybz$. They are infinitesimal. The complete expansion of CR9 is its scalar CR11, twice the three first normals CR16, and a polynomial quadratic Gram in N_u, N_X, N_Y. Terms of normal degree at least three have infinitesimal Gram entries. For weak β, or for unit β with z infinitesimal, the hard residue is

$$C_0(N_u - N_Y)^2 + (A_0 + \alpha_0 + 1/2)N_Y^2 + (\beta_0 + 1/2)N_X^2 \succ 0. \tag{CR37}$$

All first-normal lifts established above have infinitesimal coefficients. The full hard-and-soft residue is therefore positive definite. In the $Z = 1$ chart use $N_b = b - aw$, $N_X = X - xa$, $N_Y = Y - yaw$. For weak β its hard residue is

$$N_X^2/2 + H_0N_Y^2 - 2\alpha_0N_bN_Y + \alpha_0N_b^2 \succ 0, \tag{CR38}$$

since $H_0 - \alpha_0 = C_0 + A_0 + 1/2 > 0$. The same finite expansion handles every higher normal term. Thus none of CR15, CR27 or CR34 is being used merely as a scalar restriction.

It remains only unit β and a testing state with Z a unit relative to W. Normalize $Z = 1$ and $W = \omega$ bounded. Retain the entire m, D packets. The remainder belongs exactly to the square of the ideal (u, b, X, Y). Its quadratic real residue is

$$\begin{aligned}(\beta_0 + 1/2)\left(X + \frac{\beta_0}{\beta_0 + 1/2}u\right)^2 + \frac{\beta_0}{1 + 2\beta_0}u^2 \\ + H_0\left(Y - \frac{C_0\omega u + \alpha_0 b}{H_0}\right)^2 + a_0(\omega u - r_0 b)^2, \qquad a_0 = C_0 - C_0^2/H_0 > 0.\end{aligned} \tag{CR39}$$

CR7 is precisely the critical relation needed for this identity. The expression is positive definite in (u, b, X, Y) for every real ω, including zero. Its complete bounded polynomial matrix is consequently strict; every higher degree is an infinitesimal correction. This completes all local states, including the unit-beta case where $\mathcal{F}$ was signed.

193.1.7 Every other test and restoration

The exact outer identities RC18–RC25 from the newly rederived C2 reciprocal theorem apply here as polynomial identities, with $n = 0$ and the entire extra β packet retained. Their strictness uses only C, A, α positive units and the critical residue relations CR7; no C2 source membership is used. Explicitly:

- For bounded original u, v, the four rows (uW, Y, X, Z) give RC18. After CR7 its real determinant is $(r_0 + u^2v^2)D_u^2D_v^2(1 + r_0 - y_0)^2/[16y_0(r_0 - y_0)(1 - y_0)] > 0$.
- For Alice reciprocal and Bob bounded, use the four rows $(W, Y, vX, sX + Z)$ and the full core RC20. Its real residue RC21 is strict using $C, A > 0$.
- For Bob reciprocal and Alice a bounded unit, use $(uW - Y, Y, X, bZ, \sqrt{D}\,Z)$ and the entire D packet as in RC22–RC23. CR7 makes the three-by-three angular determinant strict. Literal $D = 0$ causes its last row to be zero, which is allowed in a strict coefficient Gram.
- For both angles reciprocal, RC24–RC25 give the five rows $(W, X, Y, bZ, \sqrt{D}\,sZ)$ and a strictly positive real residue.

The only remaining angle chart is exactly $u, b = o(1)$, treated above. These are whole-parent identities with every omitted event packet retained, and include all literal axes and infinities. The exact source and testing sign gauges, positive pure-field homogeneous normalization factors and real/imaginary state splitting restore the stated projective tensor-SOS theorem.

Dependencies and verification boundary. The mathematical inputs are the actual full physical parent, the displayed SP2–SP3 balance proof, SE9 and the exact sign gauges, the separated strict-Gram lemma, and only the explicit outer polynomial identities RC18–RC25 with their real residue tests. No unavailable October theorem or claimed historical admission is a premise. The focused checker verifies the full scalar, all first normals with explicit contact-jet remainders, the complementary divisions, the radical and pivot identities, and the complete hard residues. Source boundedness, sign, the quantified chart cover and strictness are the analytical arguments above. Other positive-scale source types, weak-source families and global finite-lift existence remain separate obligations.

Chapter 194

Every canonical C2 source: an exhaustive mathematical join

Component `pe351eb06b28b`
Source: `NEW/global/c2_complete_source_routing_REDERIVED.tex`
SHA256: `e351eb06b28bd1f55056e4a355c8d53ffc335d12d0c794297be878f890e57ef1`
Inventory use: CT1–CT16 exhaustive source classification and proof-use scope join

194.1 Every canonical C2 source: an exhaustive mathematical join

Scope and notation. Let K_c, K_t be arbitrarily chosen, separately labelled real closed extensions of $\mathbb{R}$; no algebraic-disjointness assumption is made. All source inequalities below are in K_c. Use the entire original physical C2 parent

$$\begin{aligned} F &= H + 4nD_uD_vX^2 + 4mM^2 + 4CD_v(uW-Y)^2 + 4AD_uD_vY^2 + 4DJ^2 + 4\alpha D_u(vY-Z)^2,\\ M &= vW - X + uvY - uZ, \qquad J = uW + uvX - Y - vZ, \\ H/2 &= J^2 + D_u(vY-Z)^2 + D_v(X+uZ)^2 - D_uD_vZ^2. \end{aligned} \tag{CT1}$$

Here $D_u = 1+u^2, D_v = 1+v^2$, $d = 1$, and the omitted weights are $\lambda_{\text{event}} = \beta = 0$. Assume an actual finite noncommuting stationary zero. The native-entry theorem EN gives exactly

$$m, C, A, D, \alpha > 0, \quad n \geq 0, \quad c, a, b, t > 0, \quad 0 < r < 1, \quad h = b - a^2t > 0, \quad g > 0, \quad j > 0,$$

$$g = (1+c^2)b - (1 - r + a^2c^2t), \qquad j = 1 + c^2r - t, \qquad 1 - c^2h > 0, \quad b - a^2c^2r > 0. \tag{CT2}$$

The actual angles are $u_c = c, v_c = ac$, and the six coefficients are the actual rational stationary coefficients, with the nonzero stationary determinant proved in EN. In particular no limiting or formally perturbed source is substituted for CT1.

Theorem. Under CT1–CT2, the full parent is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$ at every projective testing pair and every real or complex state over K_t. All coefficient scales, literal $n = 0$, and every positive source opening allowed by CT2 are included. This theorem has no earlier-admission-complement hypothesis.

194.1.1 Two finite partitions

First put

$$\delta_{\text{tot}} = (1 + n + m + C + A + D + \alpha)^{-1}. \tag{CT3}$$

If δ_{tot} is a unit, all six coefficients are bounded, and the complete positive-scale theorem AU applies at its now-proved native-entry interface. Otherwise δ_{tot} is positive infinitesimal. No normalization of an individual large event is used to replace this dichotomy.

Call a positive event weak when it is infinitesimal after this normalization $d = 1$, and strong otherwise. A strong event may be unbounded. If exactly one of C, α is weak, orient it to be C, using the exact full-parent reflection

$$\begin{aligned}(u, v; W, X, Y, Z; n, m, C, A, D, \alpha)\\ \longmapsto (1/v, 1/u; Z, X, Y, W; n, D, \alpha, A, m, C).\end{aligned} \tag{CT4}$$

In homogeneous angle coordinates the identity is $F = u^2v^2F'$; its polynomial homogeneous interpretation includes literal testing zero and infinity. CT4 preserves all source gaps: writing $z = c^2$, its parameters are

$$c' = (ac)^{-1}, \quad a' = a, \quad b' = a^2/b, \quad r' = a^2zr/b, \quad t' = (1 - zh)/b,$$

$$h' = a^2zh/b, \quad g' = a^2j/b, \quad j' = g/b, \qquad \Delta' = \Delta/(b^2z^2), \quad N' = a^2N/b^2. \tag{CT5}$$

These are pure-source identities, and give positive values for $1 - r', 1 - z'h', b' - a^2z'r'$. They preserve δ_{tot} exactly. EN and HJ verify the full reflection; the retained matching-source entry prints the displayed rational source identities. No assertion $t \mapsto 1/t$ is used.

Let I, U, O denote, respectively, a positive infinitesimal, a positive bounded unit, and a positive unbounded source number. Every positive actual angle has exactly one of these types. After the two-weak case is treated below, the remaining dispatch is

(u_c, v_c)	C, α both strong	C weak, α strong
II	strong II local and remote	impossible
IU	single-angle join	impossible
IO	strong IH local and remote	weak IH full join
UI	strong UI full join	impossible
UU	five retained events	impossible
UO	single-angle join by CT4	single-angle join by CT4
OI	strong HI full join	impossible
OU	strong UI full join by CT4	impossible
OO	strong II full join by CT4	HJ by CT4

(CT6)

The letters H in the retained file titles mean the type O in this table. They denote an unbounded *finite* actual half-angle, not a literal projective endpoint. Testing endpoints are included by each full-parent theorem.

194.1.2 The two-weak case and the empty cells

Suppose first that C, α are weak. The actual proof `chart_free_c2.tex` supplies the following source implications, without a tensor-SOS premise. The coefficient-field experiments which kill three matching packets show that

$$n, m, A, D \text{ are unbounded.} \tag{CT7}$$

For example the remaining D experiment has full value $4D\varepsilon^2 - 2\varepsilon + 4(2 + \varepsilon)C + 8(1 + \varepsilon)\alpha$. If D were bounded, a sufficiently small positive real ε would contradict physicality. The other three experiments are its exact event-cycle copies. The exact signed-core estimate

$$H \geq -2u_cv_c\|\psi_c\|^2$$

follows from the printed matrix-square identity $(M_H - LI)^2 = L^2I + 4u_cv_cLJ_0$, $J_0^2 = I$. Each matching amplitude divided by $\sqrt{u_cv_c}$ is therefore infinitesimal at a norm-one contact. Compact angle coordinates and, if needed, CT4 imply

$$c \text{ bounded}, \qquad a, b, t \sim 1, \qquad r, h, g, j \text{ positive infinitesimal.} \tag{CT8}$$

In the small-angle case the two matching equations divided by $\sqrt{u_c v_c}$ first force $v_c/u_c \sim 1$ and $W_c/Z_c \sim 1$; this is a relative bound, not just a compact limiting statement. The same body proves α/u_c^2 infinitesimal before using scaled stationarity. Its invertible four-amplitude Jacobian has residue determinant 2, and the weighted stationary gradients give $n\ell_n, A\ell_A, m\ell_m, D\ell_D$ residues $(-1,1,1,-1)/4$. Thus all strict signs in CT8 are justified.

For c infinitesimal use the complete six-root radial construction in `c2_unified.tex`, `c2_y_current.tex`, and `c2_radial_corner.tex`. Only its radial domain CT8 is imported. Its pure-source weight N satisfies

$$N \asymp r+h+g+j, \qquad r/N, h/N, g/N, j/N = O(1), \qquad \operatorname{res}\big(4\alpha/(c^4 N)\big) = 1. \tag{CT9}$$

The corrected radial root is $\sqrt{N}(u^2-c^2)$. The actual proof retains all four normalized event roots and every state normal; the final exact current-zero correction makes every radial cross coefficient belong to (r,h,g,j) before division by $\sqrt{N}$. Its Gram has strict residue on the full six-root list, including the complementary W-state chart. The actual outer proof `c2_radial_all_tests.tex` uses

$$1/(Cn),\ 1/(Cm),\ 1/(CA),\ 1/(CD) = O(1),$$

which follow from CT9 and the unchanged source formulas. Its reciprocal-corner conductor and its two complete state charts retain the weak C packet exactly. For c a unit, `c2_matching_strength.tex` gives $\alpha/C \asymp 1$, both sets of four product bounds, and the same outer proof after CT4 covers the remaining corner. Both cases use the detailed arbitrary-scale physical-source noncommuting-residue theorem A6 for the noncommuting testing complement. A6's full geometry, curvature, weighted matching, promotion, and induction bodies are explicit dependencies. This completes every two-weak source, rather than assuming a preselected radial chart.

For the right column of CT6, the exact proof `angle_reductions.tex` establishes the II, IU, UI and UU contradictions from CT2 and $n \geq 0$. Some useful details fix its scope. On II, putting

$$H = c^2 h, \quad L = c^2(b-h), \quad B = H + L$$

in its own actual-angle notation makes the positive numerator bound force α weak whenever C is weak. On UI, the same source numerator and the $n \geq 0$ inequality force r infinitesimal and h a unit, contradicting weakness of C. On IU, bounded b is a unit by the inequality in CT10 below and gives C strong; unbounded b gives $C \asymp L^{-1}$ with L bounded. On UU, a, c are units, Δ and $b+r$ are units, and the positive numerator identity makes C, α weak together or strong together. These are necessary-source arguments, not all-tests conclusions.

The OI and OU contradictions use `bounded_second_angle_exclusion.tex`. With $q = u_c^{-2}$ infinitesimal and $y = v_c^2$ bounded, its exact numerator for n is $-qP/(2rQ)$, $Q > 0$. The printed positive remainder gives

$$P \geq r^2(1+y)(H+yr) - Tqr(y+2) - q^2(T^2+T), \qquad 0 < T < H + y.$$

If q/r were infinitesimal, division by $r^2(H+yr)$ would force $P > 0$, contradicting $n \geq 0$. Thus r/q is bounded, $H \sim 1$, and the exact cancelled C formula has a positive-unit numerator and bounded positive denominator. Consequently C is strong. Literal $n = 0$ obeys the same contradiction.

194.1.3 The single-angle join has no omitted opening

For an IU source put $x = u_c^2$ infinitesimal and $k = v_c$ a unit. If b is bounded, the exact source inequality $n \geq 0$ gives

$$r_0\{b_0 + k_0^2(b_0+1)\} \leq b_0, \qquad r_0 \geq 1 - b_0, \qquad b_0 \geq k_0/\sqrt{1+k_0^2} > 0. \tag{CT10}$$

The actual body `c2_bounded_state_entry.tex` proves the last bound also when $b_0 \geq 1$. There are then exactly three source alternatives:

g unit	three-opening proof, arbitrary $r, h, b-h > 0$,
g infinitesimal, r unit	last-opening proof,
g, r infinitesimal	joint centered-packet proof.

(CT11)

All bounded positive g, r have been included. The first proof supplies both bounded and unbounded n normalizations; the second uses the triangular hard normal of the entire m amplitude; the third retains both contact-zero centered n, m packets. Their common full scalar proof is the six-term balanced Laurent decomposition in `general_single_angle.tex`, with its five-term limiting version and joint version proved in the respective bodies. This supplies global equality uniqueness and a positive contact Hessian. The same bodies print complete first-normal divisions, hard blocks, the literal $n = 0$ double-zero corner and all four outer-core identities. Their remaining testing complement uses the actual detailed A6 theorem. No normal is omitted and no bound on h or $b - h$ is restored implicitly.

If b is unbounded, put $B = xb, H = xh, L = x(b - h)$. The original gaps give

$$B = H + L, \quad 0 < H < 1, \quad 0 < L < k^2(1 + xr).$$

Thus B is bounded. The exact proof `unbounded_b_reduction.tex` cancels the original denominators and gives

$$1 - r \asymp 1, \quad m \asymp b^{-2}, \quad C \asymp L^{-1}, \quad A \asymp H^{-1}, \quad D \asymp B/j, \quad \alpha \asymp B/(1 - H). \tag{CT12}$$

If B is a unit, C, α are strong and the full strong pitchfork below applies. If B is infinitesimal, the actual `unbounded_b_small_B.tex` theorem applies with $\sigma = 1/b$, with no restriction on B/σ. Its separate homogeneous scalar lemma proves the entire contact-ideal lift on

$$(uW - (B/k)Z)/\sqrt{B}, \quad uZ - \sigma kW, \quad \sqrt{B}(v - k)Z, \quad u(v - k)Z. \tag{CT13}$$

The whole-parent proof retains the possibly nonzero first-normal angular coupling and buys its strict pivot by the exact zero

$$T^2 - \sigma U^2 - ST(v - k) - \sigma kRU(v - k) = 0.$$

It proves both full normals, arbitrary W/Z, every n scale, all openings, and the literal $n = 0$ corner. Its outer charts use the displayed general-single-angle identities with scale $\sqrt{B}$, and the detailed A6 theorem for the final complement. This exhausts IU. CT4 exhausts UO with no new source assumption.

The strong pitchfork invoked in CT12 is itself complete at all upper scales. Its three-large-event body covers bounded-unit C, α, with n, A, D arbitrary; its two-large-event predecessor supplies the printed stationary equations, full normal Taylor division and real boundary Schur calculation. The body `strong_complement_pitchfork.tex` treats unbounded C or α. Its coefficient-field state equations force $m/k^4 \asymp 1$ and bounded weighted graph coordinates before any small gap is inverted. It uses the actual large n, C, α or n, D, α packets as normals. The five-root full-parent Gram has a strict angular Schur pivot because the normalized real boundary numerator is $q_{12}(v^2 - a_0^2)^2$ with $q_{12} > 0$. Only this real residue Schur computation uses a determinant inverse; no mixed tensor denominator is inverted. For its outer charts one may use the complete cap and corner proof printed in that same strong-complement body, replacing the older predecessor's reference to a preceding outer corollary.

194.1.4 The remaining occupied cells

II and OO with both complementary events strong. The exact common-II source reduction proves that α is strong precisely when $B = H + L$ is a unit. Then L is infinitesimal, H and $1 - r$ are units, and

$$m/z^2, \ A/y, \ CL, \ Dyj, \ \alpha(1 - H) \asymp 1, \qquad nr = O(1). \tag{CT14}$$

The exact numerator at $r = 1$ is negative, giving the asserted unit gap $1 - r$; none of $r, j, 1 - H$ is assumed a unit. The proof `local_strong_ii.tex` retains all four centered large-packet excesses and gives the full three-normal remainder. Its angular identity, harmonic C, D square and free radial pivot are exact; the normalized hard residue and the remaining two-by-two Schur margin are strictly

positive. The separate `strong_ii_remote.tex` uses the original homogeneous parent and harmonic current, not a higher-degree homogenization of the local graph. It proves all remaining angles and state charts, including the endpoint forms. Thus there is no local-to-remote scope gap. CT4 gives OO.

IO, weak or strong C. The weak IH source entry proves B unbounded and $\rho = B/y$ bounded with $0 \le \bar{\rho} \le 1$. The full interior-corner proof, both $\rho = 0, 1$ source-endpoint extensions, remote-chart proof, and separate open-axis proof cover all independent angles and states. The latter open-axis proof is indispensable: it supplies the product-pivot split when u is a unit and $1/v$ is infinitesimal. All these bodies retain nr and the full D packet at the closing endpoint; literal $n = 0$ is included. The actual weak-IH all-tests join merely assembles these already specified bodies.

If C, α are both strong, the separate strong-angle source entry gives B a bounded unit, $r = O(1/y)$, $Dy^2 \asymp 1$ and the remaining opening-safe products. The strong-IH local proof reuses only the explicit polynomial current identities of the weak-IH proof, with its own source estimates and finite C cap; it does not import the weak-IH source conclusion. Its remote proof includes the two open axes and reciprocal corner and uses detailed A6 only when both testing pairs are noncommuting in residue.

UI and OU with both complementary events strong. The actual `full_parent_reduction.tex` in the C2-UI directory cancels all closing source openings before dividing the normals. In particular $nr, m\mu, \alpha\pi, C\rho k^2, D\sigma k^2$, and the difference $Ct - Dj$, are bounded. The exact state inverse is polynomial and never divides by a state coordinate. The reciprocal-contact lemma and local certificate prove the full contact Gram, including all three normals. The complete `ui_all_tests_recovered.tex` proof supplies real interior-source approximation for its finite-cap step and strict Grams on every complementary axis and state chart. Its Laurent equality and positive contact Hessian follow from the actual source equations. No new global source classification is imported from this local theorem. CT4 gives OU.

OI. The exact HI source entry proves directly from $n \ge 0$ that $R = r/q$ is bounded, $H \sim 1$, and

$$mqK,\ Cyt,\ Dyj,\ \alpha qE,\ A/(qy) \asymp 1, \quad q = u_c^{-2},\ y = v_c^2 \text{ infinitesimal.} \tag{CT15}$$

Thus m, C, D, α are all unbounded, without a prior strong-event assumption. Its remote full-parent theorem caps these four coefficients at a sufficiently large *real* constant after fixing the testing residue. It prints strict full state forms on every remote chart and literal critical axis; only $1/u, v$ both nonzero infinitesimal remain. The retained-contact module gives the complete local four-monomial circuit and radical conductor there. The full HI module includes the actual n packet, its two complete normals, and the endpoint $R_0 = 0$ where n can be unbounded. Its constant circuit multiplier lies in $[2/3, 2]$, and its normalized hard residue is strictly positive. Its proof uses the predecessor only for the explicit half-angle conductor, never for an unsupported $n = 0$-to-$n > 0$ inference. Together these three bodies close OI.

UU and the five-event terminal. The exact source-angle equations give m, A, D strong on the both-strong UU cell. The actual five-retained-events proof uses $\mu = \min(1, m, C, A, D, \alpha)$, a positive unit, and retains all excess original packets. Its commuting-axis full state forms and three nontrivial double-corner coordinate Grams are strict. For noncommuting testing pairs it uses the detailed arbitrary-scale physical-source theorem A6 for the original source. The selected capped parent is never asserted globally physical. This distinction prevents a contact-zero packet from being improperly subtracted at the source contact.

OO in the directed weak case. CT4 gives exactly the native II domain HJ1: C strong, α weak, both actual angles infinitesimal, and δ_{tot} infinitesimal. HJ1–HJ16 prove the unchanged source equations, the equivalence of the weak face with B infinitesimal, the complete ten-row partition, all its literal $n = 0$ maps, and every source/testing/state join. The bare-source audit BSR1–BSR15

proves the precise retained R5–R10 hypotheses and restores each actual earlier exit. Neither HJ nor BSR assumes this C2 theorem, the whole C1 theorem, or AU. Its only C1 uses are the two explicit bare CS and small-e all-tests bodies under the exact $n = 0$ source map. Consequently this last cell introduces no dependency cycle.

194.1.5 Proof-use scope and the conclusion

The mathematical input list accompanying this theorem pins every actual proof body used above, including the subordinate single-angle and homogeneous scalar proofs, the complete normal and source-endpoint constructions, and the noncommuting residue proofs. An inherited final paragraph which describes a then-open complement is not a premise. Only the displayed source hypotheses and the exact full-parent constructions just specified are imported. In particular:

$$\begin{gathered}\text{native entry is EN, not an unavailable sign label;}\\ \text{positive scale is AU, not an undefined terminal call;}\\ \text{weak II is HJ, not its former frontier assertion.}\end{gathered}\tag{CT16}$$

The bodies in CT10–CT15 use the exact source equations, positivity, the typed source scales and the displayed opening bounds. Their scalar divisions, normal bounds, caps and outer joins do not use exclusion from any blanket list of earlier admissions. Those exclusions are therefore absent from the theorem: this is a proof-use verification on the actual bodies, not deletion of a logical hypothesis by relabelling.

Every positive δ_{tot} has one of the two types in CT3. In the infinitesimal case C, α have exactly one of the three weak/strong patterns used above; CT4 orients the asymmetric pattern. The nine ordered angle types in CT6 are exhaustive. Every occupied cell has a full-parent theorem at its exact source hypotheses, and every empty cell has the necessary-source contradiction stated above. Thus CT1 is tensor SOS at every independent test. Complex states split into real and imaginary parts because CT1 is a real quadratic form in the state. All source and testing normalizations are pure-field square factors. Restoring the original positive d completes the theorem.

Chapter 195

The complete canonical local certificate and global source dispatch

Component `pe1088004af65`
Source: `NEW/global/complete_source_routing_REDERIVED.tex`
SHA256: `e1088004af657604cd6a4a9ee2174584dd8d69f175b04f8d0a0f61fbbdcedcb3`
Inventory use: LC1–LC10 and LCH1–LCH3: exact canonical source cover, bare C14 strengthening, all historical row implications, and universal separated slack via finite source-conic/testing-convex closure.

195.1 The complete canonical local certificate and global source dispatch

Quantifiers and the full-parent target. Let K_c, K_t be arbitrarily chosen, separately labelled real closed extensions of $\mathbb{R}$. There is no algebraic-disjointness assumption: the two factors may be labelled copies of the same field. Put $T = K_c \otimes_{\mathbb{R}} K_t$. Consider a physical canonical source over K_c with an actual finite noncommuting stationary zero and one of the exact patterns

$$\begin{aligned} &\text{C1}: \quad d, m, C, A, \alpha > 0, \quad \lambda = n = 0, \quad D, \beta \geq 0, \quad D + \beta > 0, \\ &\text{C2}: \quad d, m, C, A, D, \alpha > 0, \quad \lambda = \beta = 0, \quad n \geq 0. \end{aligned} \tag{LC1}$$

These are precisely the two elementary source hypotheses produced by the actual extreme-source reduction SE15. For finite independent testing half-angles and any state $(W, X, Y, Z) \in K_t^4$, the complete parent is

$$\begin{aligned} F = {} & 4\lambda D_u (W + vX)^2 + 4nD_u D_v X^2 + 4m(vW - X + uvY - uZ)^2 \\ & + 4CD_v (uW - Y)^2 + 4AD_u D_v Y^2 + (4D + 2d)(uW + uvX - Y - vZ)^2 \\ & + (4\alpha + 2d) D_u (vY - Z)^2 + (4\beta + 2d) D_v (X + uZ)^2 - 2dD_u D_v Z^2. \end{aligned} \tag{LC2}$$

Every coefficient times a testing expression denotes its separated tensor product. The homogeneous angle form of LC2, rather than an affine limit obtained by a cancelled zero coordinate, defines the literal projective endpoints.

Theorem 195.1 (Complete local certificate)**.** *For every source in LC1 and every independent projective testing pair and real or complex testing state, the entire homogeneous parent is a finite sum of squares in T. This includes every coefficient boundary permitted in LC1, every positive source opening, all relative source and testing rates, and all state ratios. The source contact is never identified with the testing point.*

195.1.1 Normalization and actual native entry

Divide by the positive pure-source d. This operation is reversible in a tensor sum of squares by multiplication of each root by $\sqrt{d} \otimes 1$. Before or after this normalization the invariant total-scale

quantity is

$$\delta_{\mathrm{tot}} = \frac{d}{d+\lambda+n+m+C+A+D+\alpha+\beta}. \tag{LC3}$$

It is positive and at most one. Its positive-unit and positive infinitesimal alternatives are exhaustive. In the first alternative the complete positive-scale theorem AU1–AU19 applies after EN. The second alternative retains the hypothesis $\delta_{\mathrm{tot}} \in \mathrm{Inf}_{c,>0}$ in every residual theorem used below. Setting $d = 1$ does not erase that hypothesis.

The theorem EN1–EN16 derives the full native signs and exact rational source identification from physicality and the actual finite contact. Its proof uses SE's coercive Laurent support, strict single-field contact Hessian, contact sufficiency and event-row deletion implication. It does not assume any tensor certificate being assembled here. For C1, after the exact Alice source reflection, its native coordinates satisfy

$$\begin{gathered} 0 < k \le 1, \quad \eta, \zeta > 0, \quad r = 1 - a\eta \in (0,1), \quad \theta = t\zeta \in (0,1), \\ 0 < a, t < 2, \quad x = k^2, \quad h = 1 - \theta, \quad E = 1 - x\eta h > 0, \quad Q = 1 - x\zeta r > 0, \\ M = 1 + xr - t > 0, \quad J = 1 + xh - a > 0, \qquad e = k\sqrt{\eta/\zeta} > 0. \end{gathered} \tag{LC4}$$

The original D, β remain nonnegative, and either may be zero. For C2, EN gives exactly CT2 and its unique actual stationary coefficient formulas, including $n = 0$. All these implications hold over every K_c by the first-order transfer proved in EN; no topological compactness of a non-Archimedean field is being presumed.

195.1.2 Every weak-total C1 source

Use the native parameters LC4, the actual original coefficients, and δ_{tot} positive infinitesimal. Here I, U and O mean a positive infinitesimal, positive bounded unit, and positive unbounded source quantity, respectively. Since $0 < k \le 1$, k has type I or U. The following disjoint scale dispatch is exhaustive:

k	ζ	remaining source type	actual theorem
U	any	η, ζ : I	AS
U	any	η, ζ not both I	ordinary-Alice join below
I	U or O	$\eta > 0$	JC with bare source scope
I	I	η : I	uniform all-e join
I	I	η : O	C1.1
I	I	η : U, e : I	C1.3
I	I	η, e, r : U	BE
I	I	η, e : U, r : I	CA
I	I	η, r : U, e : O	$\mathrm{UE}_{\mathrm{old}}$
I	I	η : U, e : O, r : I	C1.2.

(LC5)

The rows using r are exhaustive because $0 < r < 1$. The positive number e has exactly the three types shown. No comparison with an independent testing quantity occurs in LC5.

Ordinary Alice: the actual UI, UU and UH split. Suppose k is a unit and η, ζ are not both infinitesimal. If η is infinitesimal, then $r \sim 1$ and $Q > 0$ make ζ bounded. Thus ζ is a unit in this case. If ζ is infinitesimal, then θ is infinitesimal, $h \sim 1$, and $E > 0$ make η bounded; hence η is a unit. In particular the two possibilities

$$(\eta : \mathrm{I}, \zeta : \mathrm{O}), \qquad (\eta : \mathrm{O}, \zeta : \mathrm{I}) \tag{LC6}$$

are impossible when k is a unit. Since $v_c^2 = x\eta\zeta$, the remaining possibilities are: one of η, ζ infinitesimal and the other a unit, giving actual UI; both units, giving actual UU; or both noninfinitesimal with at least one unbounded, giving actual UH.

The actual UI source theorem and its exact native stabilizer cover the first possibility. More explicitly, the exact AS3–AS5 full-parent reflection reduces it to η a unit and ζ infinitesimal. This reflection preserves the C1 omission pair and the original nonnegative coefficients; its new native parameters and gaps are checked in that identity. Its Alice parameter remains a unit, but a second assertion $k \le 1$ is not imposed on the reflected representative. Use the complete CU1–CU13 theorem in `ui_all_tests.tex`, together with CR1–CR13 in the actual complementary-testing body. For these displayed CU source quotients rescale by the positive pure-source factor to its normalization $C = 2$; the native parameters are unchanged, and this local normalization is restored afterwards. In CU put $g^2 = \zeta$. The exact source quotients make $C/a, A/a$ strong, $d/a, \alpha/(ag^2)$ units, and $m/a, \beta/a$ large. CU2 retains the entire paired m, β normal packets; CU3 retains the centered C, A, D excesses. CU5 is the whole remaining quadratic state polynomial on its bounded six-root frame. CU6–CU9 explicitly divide all complete first-normal coefficients and the scalar correction in the contact ideals. Their exact tensor zero redistributes the two angular pivots; CU10 computes the full matched state Schur Hessian. Its positive determinant uses the original $D \ge 0$ numerator. The possible divergent A, D pair has independent hard rows, so its constrained finite-cap limit keeps the angular components of those rows. At unmatched Alice residue the normalized Laurent coefficients retain coercivity and strict convexity. The actual real two-block lemma UA1–UA3 is applied using the exact relation $L(gZ) - Q(vZ) = 0$ between its root products, not by inverting a determinant in T. CR supplies all complementary tests, especially its CR5a five-root reciprocal-Bob frame retaining the weak α packet. This proves all UI tests and states. Literal $D = 0$ is retained; $\beta = 0$ is impossible here by the displayed lower bound on the actual β/a. For actual UU use the full NC1–NC7 source proof in `c1_source_formulas.tex`. It first proves the uniform native gap $E \ge 1/18$. The nonnegative D, β formulas and their exact elimination identity give

$$C, A, m, \alpha \text{ strong}, \qquad D \text{ or } \beta \text{ strong}. \tag{LC7}$$

In more detail x, v_c^2, η, ζ are units; the specialization $B_\beta|_{\theta=1} = -1 - x\zeta(1-r)$ makes h a unit. The original numerator polynomials P_4, P_α, P_d are positive units. If both D, β were weak, the left side of the exact NC7 elimination would be infinitesimal and its explicit positive right side a unit. This is impossible. The five consecutive strong events in LC7 therefore give the actual five-event all-tests proof, with its detailed A6 noncommuting testing complement. This is a source theorem at all upper event scales, not an application of a bounded-coefficient theorem to unbounded data. For actual UH the native NI8–NI10 implications give two alternatives: η unbounded with ζ a unit, or ζ unbounded with η noninfinitesimal. The exact β-numerator identity proves ηh a unit in both. Thus, with $\sigma = 1/\eta$, $z = 1/\zeta$ and $H = h/\sigma$, the source is respectively σ infinitesimal and z a unit, or z infinitesimal and σ bounded, always with H a unit. These are precisely the printed UH1 alternatives; no bound on the ratio of two source infinitesimals is added. The actual four-consecutive-event theorem CE and its matched weak outer frames CE17–CE21 remove the remote tests. The remaining strip has unit testing u and infinitesimal $w = 1/v$. The complete homogeneous biquadratic conductor `uh_all_tests.tex` applies there. With $b = r/z$, use its scaled weights $\widehat{m} = m/z$, $\widehat{C} = C/z$, $\widehat{A} = Az$ and $\widehat{\alpha} = \alpha z$. Their products $\widehat{m}M, \widehat{C}(1-r), \widehat{A}b, \widehat{\alpha}Q$ are bounded and cancel the opening poles before the full scalar and first-normal collection. At a matched Alice residue its exact angular zero and the full hard Schur calculation redistribute the positive angular pivots. At an unmatched residue its raw angular product relation gives the finite real two-block allocation. The single and double opening limits retain all divergent centered rows; their independent hard-row determinants and bounded hard minimizers justify the constrained limits. This is a full-parent construction, including the $J = \theta = 0$ residue face. Both $D = 0$ and $\beta = 0$ are retained where permitted by the actual source equations. Thus no assertion that UI or UH alone already covers UU is needed.

The uniform source and the four earlier exits. In the fourth row of LC5, $k, \sqrt{\eta}, \sqrt{\zeta}$ are all infinitesimal, while $e = k\sqrt{\eta/\zeta}$ is arbitrary. The new uniform all-e join UE1–UE4 applies exactly. For e infinitesimal it uses the fresh complete UJ proof, with its full bounded-ratio and inverse-ratio state charts. Otherwise $1/e$ is bounded, and the actual complete bounded-Delta proof supplies its restricted

monic module, weighted full scalar, both entire first normals, hard block, and complementary state chart. Its outer current formulas are joined in UE without an unprinted restriction on e. Neither a historical whole-uniform assertion nor a missing old conductor label is a premise.

The AS, BE, CA and $\mathrm{UE}_{\mathrm{old}}$ rows use their actual full-parent proofs, respectively `arbitrary_state_ratio.tex`, `bounded_e_all_tests.tex`, `centered_a_unit_e.tex` and `unit_eta_unbounded_e.tex`. Their ordinary-angle, centered-A, contact-conductor and outer-current subordinate bodies are included explicitly. AS proves the needed strict original event signs on its source domain. BE and CA include $\beta = 0$. The last proof derives β/d unbounded, so its intersection with literal $\beta = 0$ is empty by an actual source estimate. These proofs use the bare LC4 equations and their stated source scales; no estimate comes from exclusion of an unnamed earlier admission.

The remaining C1 rows and the complete C1.4 union. For C1.1 the actual source reduction, including the printed $k^2\eta \le 1/2 + \mathrm{Inf}_c$ derivation, splits into the unit-q and infinitesimal-q branches. The complete unit-q body and the protected-normal local and remote infinitesimal-q bodies prove every testing/state chart. C1.2 uses the actual centered-A bounded-Delta theorem, including its full contact conductor and centered-A outer proof. C1.3 uses the actual small-e all-tests theorem and its explicit outer-deficit calculation. These are full bare-source statements with the scales printed in LC5.

For the third row of LC5, the source is exactly the native C1.4 input k infinitesimal, ζ noninfinitesimal, with all original gaps and δ_{tot} infinitesimal. We now give its bare-source join explicitly; this also explains why a blanket earlier-complement predicate in the entered JC statement is unnecessary for this invocation.

Put $L = x\zeta$. If L is unbounded, $Q = 1 - Lr > 0$ forces r infinitesimal. Since $1 - r = a\eta$ and $0 < a < 2$, η cannot be infinitesimal. Unit η is the complete `large_l_unit.tex` exit. For unbounded η, put $\sigma = 1/\eta$, $H = h/\sigma$ and $M_\beta = \beta\theta/\sigma^2$. Bounded H, M_β, including $\beta = 0$, is the actual JR/JB theorem. Bounded H with unbounded M_β is the full C14.1a SF conductor. Unbounded H is the C14.1b source: SF first treats $\operatorname{res} h < 1/2$; US bounds k/σ and treats its unit case. The BS/BC/GA and CE testing and state exits then leave exactly the small-sphere source and the infinitesimal testing chart stated in JC5. The fresh AR theorem restores the angular budget and all SP6 first normals there. AR requires no bound on qu/w, u/w, W/Z or $uW/(wZ)$. This closes the large-L alternative without a comparison of independent source and test rates.

If L is a unit, the three types of η give, respectively, the complete `lunit_eta_small.tex` theorem, the LU theorem in `eta_unit.tex`, and C14.2. The first two use the actual LII parent dictionary with their own native coefficient checks. They do not identify the source with an arbitrary C2 source satisfying a narrower theorem. C14.2 uses its explicit NC/LC source bounds, complete C-opening extraction, HG/CT radial matrix, sphere-weighted budgets, CS/CF/HG state joins and LA/LB remote tests. Every part has the full parent and all normal terms.

If L is infinitesimal and η unbounded, then $\sigma = 1/\eta$ is infinitesimal and this is C14.4. Its source-only parameters and exact identity are

$$\begin{gathered} S = \sigma + L, \quad \kappa = x/S, \quad \nu = \sigma/S, \quad \ell = L/S, \quad \nu + \ell = 1, \quad \zeta = \ell/\kappa, \\ Q_s = 1 - \ell S r, \qquad \kappa Q_s + \ell M + \nu E = 1, \qquad E \ge 1/18. \end{gathered} \tag{LC4a}$$

Consequently κ is bounded. The fractions ν, ℓ cannot both be infinitesimal; if κ is a unit, then $\ell = \kappa\zeta$ is a unit. Thus the five exact source cells are

κ	ν	ℓ	whole-parent theorem
U	U	U	FU with its UKR matched chart
U	I	U	NU
I	U	U	FM
I	U	I	EL
I	I	U	SN, BN, UN.

(LC4b)

For the last cell put $q_\nu = \nu/\ell$, $\tau = h/q_\nu$ and $n_s = \beta/q_\nu^2$. Bounded n_s gives SN and $\tau \sim 1$; unbounded n_s with bounded τ gives BN; unbounded τ gives UN, including infinitesimal θ. Each source is

covered. All these proofs start from the exact FM dictionary, NC5 and their actual source/sphere equations. Their SE/EB/CE testing exits have those same printed source inequalities. They retain all normal terms, literal coefficient zeros where permitted, and every source opening.

It remains to take L infinitesimal and η bounded. Unbounded ζ is C14.3a. Unit ζ with infinitesimal η is C14.3b. Their complete JF parent is combined with DG/LD normal conductors and the actual completed j conductor; the latter removes the earlier x/η restriction. The old restricted follow-up assertion is not invoked. DS/LS and SB supply all states and the remote tests. The small-η, bounded-a input is used only through its proved angular paragraph, not its old proposed radial continuation.

The remaining η, ζ are units. Use the actual positive source gap

$$j = \eta(1 + xh) - 1 + r = \eta J > 0, \qquad j_0 = \eta + r - 1, \qquad j - j_0 = \eta xh \in \mathrm{Inf}_c. \tag{LC4c}$$

It is bounded. Unit j gives positive unit j_0 and the RO theorem. Infinitesimal j gives $\eta + r \sim 1$ and unit $1 - r$; unit r then gives JO and infinitesimal r gives HO. RO uses the IC parent dictionary with its newly proved native bounds, not the narrower IC1 source premise. JO and HO use $j > 0$, including $j_0 = 0$ or slightly negative. Their exact three-circuit body and SB remote proof retain all normal terms.

This proves the entire L, η, ζ, j source partition directly from LC4. Every local source estimate just invoked uses these native equations, the original $D, \beta \geq 0$, its printed scale cell, the NC gap and, where required, the unchanged infinitesimal total scale. No bound is obtained from failure of an unnamed earlier theorem. The seven restored exits are exactly large-L/unit-η, unit-L/small-η, unit-L/unit-η, JR/JB, RO, JO and HO. They are positive parts of this cover. Thus the actual JC constructions prove the bare third row of LC5; any original earlier-complement predicate can afterwards be intersected with the result. The strengthening preserves physicality, signs, gaps, literal boundaries and total scale.

All ten rows of LC5 now have complete original-parent certificates. They include every residual C1 source after the positive-total-scale alternative, so C1 in LC1 is proved.

195.1.3 Every C2 source and the absence of a circular premise

The theorem CT1–CT16 proves the entire native C2 family. Its proof takes the same total-scale split, treats both weak complementary events by actual radial source entry, and otherwise gives all nine ordered actual-angle types in its explicit table. Every empty cell has an exact necessary-source contradiction; every occupied cell has its complete local and remote full-parent proof. In particular its directed weak HH cell is reflected to the exact HJ domain, with C strong, α weak and both actual angles infinitesimal. The HJ proof includes all ten source rows and every allowed $n = 0$ map. Its C1 boundary calls use only the two explicit centered-A and small-e bare-source bodies, never the C1 conclusion of the present theorem. Thus CT has no dependence on this global C1/C2 join. This proves C2 in LC1.

The mathematical dependencies have the following direction:

$$\begin{gathered}\text{actual physical source geometry} \implies \text{SE, EN,} \\ \text{local identities and strict tensor Gram} \implies \text{AU, JC, HJ, UJ, scoped terminals,} \\ \text{these terminals and finite source partitions} \implies \text{CT, LC.}\end{gathered} \tag{LC8}$$

A6 is separately proved from the actual zero-d geometry, one- and two-hard-packet cases, weighted matching, promotion and induction bodies; it does not use LC, CT, JC or HJ. EN does not use AU or any tensor-SOS conclusion. The source entries and coefficient-field physicality arguments inside local proofs are not converted into positivity statements in the tensor ring. At no step is a mixed Schur denominator cancelled. All roots and all first and higher state-normal terms belong to the actual full parent LC2.

The finite mathematical inventory accompanying this proof records direct and transitive actual proof bodies with their restricted imported statements. An old open-complement paragraph, a

historical status, a filename, or a checker assertion count is not a mathematical premise. Finite typed-source checks corroborate the exhaustion of LC5 and CT6; strictness, conductor membership and full-normal restoration are proved in the displayed mathematical bodies.

195.1.4 Exact reconciliation of the historical rows

The original thirteen-row checklist is reconciled by the following restriction maps, not by changing its acceptance criterion. Every row retains its original physical source, strict gaps, infinitesimal total scale, testing/state domain and any originally intersected earlier-complement predicate. The present theorem is proved on the indicated bare domains, so it also holds on those intersections. The letter H in these original labels means an unbounded actual half-angle, the type O used above.

original row	original role	present proof
C1.1	earlier terminal	LC5 : C1.1 exact bodies
C1.2	earlier terminal	LC5 : centered-A theorem
C1.3	earlier terminal	LC5 : small-e theorem
C1.4	six refined rows	JC and LC5 bare scope
C1.5	earlier terminal	LC6–LC7, UI, UH
C2.W.IH	earlier terminal	CT6 : weak IH full join
C2.W.HI	empty source	CT : WE1–WE6
C2.W.HU	empty source	CT : WE1–WE6
C2.W.HH	ten refined rows	CT4, HJ1–HJ16
C2.S.II_HH	earlier terminal	CT14 : II local, remote, CT4
C2.S.IH	earlier terminal	CT : strong IH full join
C2.S.UI_HU	earlier terminal	CT : UI full join, CT4
C2.S.HI	earlier terminal	CT15 : HI full join

(LCH1)

This explicitly accounts for the original nine earlier terminals, the two empty source rows, and the two rows requiring subdivision. The extra original condition *at least one of* m, A, D *weak* on each C2.S row is retained by restriction of the broader both-strong C, α statements in CT. Neither emptiness proof is replaced by an all-tests claim for nonexistent sources. Every one of the nine earlier terminals has been reconnected to its complete mathematical body and all necessary subordinate proofs; its former status is not used.

For the C1.4 subdivision put $L = k^2\zeta$, and, when η is unbounded, $\sigma = 1/\eta$. After the seven explicitly restored JC prior exits, the original six predicates are exactly

label	retained source predicate	present theorem
C14.1a	$L, \eta : \mathrm{O}$, h/σ bounded, $\beta\theta/\sigma^2 : \mathrm{O}$	JC
C14.1b	$L, \eta : \mathrm{O}$, $h/\sigma : \mathrm{O}$	JC
C14.2	$L : \mathrm{U}$, $\eta : \mathrm{O}$	JC
C14.3a	$L : \mathrm{I}$, η bounded, $\zeta : \mathrm{O}$	JC
C14.3b	$L, \eta : \mathrm{I}$, $\zeta : \mathrm{U}$	JC
C14.4	$L : \mathrm{I}$, $\eta : \mathrm{O}$, ζ noninfinitesimal	JC.

(LCH2)

All six share the original C1.4 source input. JC supplies their separate complete local and outer constructions as well as the precise scale disjunction, including its empty intersection L unbounded with η infinitesimal. The seven earlier source exits are proved, rather than silently subtracted. LCH2 is therefore an implication for each original predicate, including its allowed literal zero-event and source-opening faces.

For the weak-C2 subdivision use exactly the common II coordinates HJ4–HJ10 after CT4:

$$z = u_c^2, \quad y = v_c^2, \quad B = H + L, \quad a = z/B, \quad t = y/B, \quad h = H/B, \quad l = L/B, \quad e = 1 - r.$$

Here B, z, y are positive infinitesimal, $h + l = r + e = 1$, all original gaps remain positive and the

actual $n \geq 0$ formula is retained. The ten original predicates and present proofs are

R1	$t : \mathrm{I}$	HJ
R2	$t : \mathrm{O}$	HJ
R3	$t, h : \mathrm{U},\ l : \mathrm{I}$	HJ
R4	$t, h, l : \mathrm{U},\ j : \mathrm{I}$	HJ
R5	$t, h, l, j : \mathrm{U},\ r : \mathrm{I}$	HJ, BSR
R6	$t, h, l, j, r : \mathrm{U},\ g : \mathrm{I}$	HJ, BSR
R7	$t : \mathrm{U},\ h, g : \mathrm{I},\ r, j$ not both I	HJ, BSR
R8	$t, g : \mathrm{U},\ h, j : \mathrm{I}$	HJ, BSR
R9	$t, g, j : \mathrm{U},\ h, e, a : \mathrm{I}$	HJ, BSR
R10	$t, g, j : \mathrm{U},\ h, e : \mathrm{I},\ a : \mathrm{O}$	HJ, BSR.

(LCH3)

These rows use the original ordered dispatch, not an unstated quotient of the directed angle list. HJ14 proves that the all-six-unit alternative contradicts the retained infinitesimal total scale. HJ15 proves that r, g, h, j all infinitesimal would make C weak, contradicting this directed strong-C input; it justifies the exclusion written in R7. The other earlier exits are dispatched by their actual source formulas and complete full-parent bodies in HJ/BSR. Thus the ten-row exhaustion and each terminal implication, not merely the number ten, are supplied.

195.1.5 Restoration and the universal slack consequence

Every local proof gives the unmultiplied parent, or an equivalent parent multiplied only by a positive pure-source or pure-testing factor whose nonvanishing is explicitly proved. Such a factor is removed by its square root inside that field. Original retained packets are then added with their actual nonnegative coefficients. The exact homogeneous angle identities restore every finite chart and include literal endpoints without division by a vanishing coordinate. Pure-testing state normalizations cover all nonzero real projective states, and the zero state is immediate. Since LC2 is a real quadratic form, applying the real result to real and imaginary components and adding proves the complex-state assertion. Restoring d finishes the complete local-certificate theorem.

Corollary 195.2 (Universal separated slack)**.** *Let $\mathcal{Q}_{222} \subset \mathbb{R}^8$ be the finite-dimensional bipartite tensor-product quantum set in the two-setting, two-outcome scenario, and let P be its cone of nonnegative affine functionals. Use the fixed finite real-qubit formula for $(\mathcal{Q}_{222})_{K_t}$ and the transferred cone P_{K_c}. Then for every separately labelled K_c, K_t as above,*

$$f_0 \otimes 1 + \sum_{\nu=1}^{8} f_\nu \otimes q_\nu \ \in\ \Sigma T^2 \qquad (f \in P_{K_c},\ q \in (\mathcal{Q}_{222})_{K_t}). \tag{LC9}$$

Proof. The complete source reduction SE11–SE15 writes every normalized source in K_c as a finite conic combination of event atoms, the two explicit noncoercive closed faces, and the actual canonical contact sources LC1. It proves this fixed finite statement over the reals and transfers it; no arbitrary infinite extreme-point expansion is used. Event atoms have separated squares. Each of the two closed faces has the actual finite matrix lift and strict feasible point in `closed_faces_short.tex`. Attained finite-dimensional duality and separate-field positive-semidefinite factorizations give its tensor trace-pairing squares. Canonical atoms use the theorem just proved, followed by the exact homogeneous slack normalization in the physical reduction.

Each testing point is a convex combination of at most nine real-qubit atoms in its own field. If the source combination weights are $a_j \geq 0$, testing weights are $w_i \geq 0$ with $\sum_i w_i = 1$, and each atom pairing is $\sum_k h_{ijk}^2$, then the full pairing is exactly

$$\sum_{i,j,k} \left[(\sqrt{a_j} \otimes \sqrt{w_i}) h_{ijk}\right]^2. \tag{LC10}$$

Affineness preserves the constant term because the testing weights sum to one. Zero weights are never inverted. All Bell relabelings and all positive normalizing factors are restored inside their own field. This is finite conic and convex closure in the tensor ring, and proves LC9. □

The proof supplies the universal tensor statement, not an evaluated affine matrix pencil or a numerical bound on its size. The separate finite-lift criterion applies to LC9 using two labelled copies of any one real closed extension; its compactness conclusion is not inferred from a sum of the local root counts.

Chapter 196

The complete positive-total-scale canonical source union

Component `pc0b2849f8d09`
Source: `NEW/global/positive_scale_union_REDERIVED.tex`
SHA256: `c0b2849f8d092c22f3c525128bb352681aa7e80818bfcf91489a6521c1bfc33c`
Inventory use: AU1-AU19 complete positive-total-scale canonical source union and proof-use scope strengthening

196.1 The complete positive-total-scale canonical source union

Exact statement. Let K_c, K_t be labelled real closed extensions of $\mathbb{R}$. The original canonical coefficient vector is physical over K_c, has nonnegative coefficients, and has an actual finite noncommuting zero contact. In C1 the exact omissions are $\lambda = n = 0$, with $m, C, A, \alpha > 0$, $D, \beta \geq 0$, $D + \beta > 0$; in C2 they are $\lambda = \beta = 0$, with $m, C, A, D, \alpha > 0$, $n \geq 0$. Use the actual native sign chart of this contact, with its explicitly positive source gaps, as supplied by canonical entry. Assume that normalization by $d > 0$ makes all coefficients bounded. Equivalently the normalized total-scale coordinate $d/(d + \lambda + n + m + C + A + D + \alpha + \beta)$ is a positive unit. Then the entire original homogeneous slack is a sum of squares in $K_c \otimes_{\mathbb{R}} K_t$ for every independent projective testing pair and every real or complex testing state. No prior-admission complement is assumed. This is the A5 terminal union at the explicit canonical-entry interface; global entry into that interface is a separate theorem.

Division by d is a positive pure-source normalization. Hence take $d = 1$ below. Every nonnegative bounded source quantity is either infinitesimal (including literal zero where permitted) or a positive unit. The proof makes that finite split exhaustive, rather than using undefined historical terminal labels.

196.1.1 An explicit noncommuting-residue lemma for this union

For $\lambda = 0$ the determinant Laurent polynomial is

$$L(U,V) = h + \frac{q_{01}}{U} + \frac{q_{02}V}{U} + \frac{q_{10}}{V} + q_{12}V + \frac{q_{20}U}{V} + q_{21}U + q_{22}UV, \qquad U = u^2,\ V = v^2. \quad \text{(AU1)}$$

All noncentral coefficients are nonnegative, with the exact factorization in the physical-core input. Suppose C, A, D, α are units. If m is weak, then $\bar{q}_{01} = \bar{q}_{02} = 0$, whereas $\bar{q}_{21} > 0$. At a finite nonzero real U, V,

$$U\partial_U \bar{L} = \bar{q}_{20}U/V + \bar{q}_{21}U + \bar{q}_{22}UV > 0. \quad \text{(AU2)}$$

Thus a physical residue source has no zero there: a differentiable nonnegative function at an interior zero has zero derivative.

If m is a unit, then $q_{01}, q_{10}, q_{12}, q_{21}$ are positive units. At any real noncommuting zero, the Hessian of $L(e^s, e^t)$ is

$$\sum_{\gamma} \bar{q}_{\gamma} e^{\gamma_1 s + \gamma_2 t} \gamma\gamma^{\mathsf{T}} \succ 0. \tag{AU3}$$

The terms with exponents $(-1,0),(0,-1)$ already span both directions. This is a real residue calculation only.

Here is the full-parent consequence, without a mixed inverse. The $Z = 0$ principal state block is positive definite: AY^2, $C(uW - Y)^2$, and the positive core X-square already force $W = X = Y = 0$, since $u \neq 0$. A zero state therefore has $Z \neq 0$, and it can be normalized to $Z = 1$ in the testing field. Its real three-state Schur complement is

$$g(u,v) = \frac{\det P(u,v)}{\det P(u,v)|_{Z=0}} = \frac{D_u^2 D_v^2 UV}{16 \det P(u,v)|_{Z=0}} L(U,V) \tag{AU4}$$

for the quarter-parent matrix P. At a zero the value and first derivatives of L vanish. AU3 and the nonzero real change from u, v to $\log U, \log V$ therefore give a positive angular Hessian for g. The full Hessian in the three state coordinates and two angles is positive definite.

The polynomial implicit-function theorem supplies, in the source field only, a critical center near this real five-variable zero. One may justify this over an arbitrary real closed source extension by the semialgebraic local implicit-function theorem over $\mathbb{R}$ and real closed transfer. The Hessian is a real positive definite matrix at the base point, so its invertibility and the local uniqueness condition are explicit. The original source is physical, hence its value μ at that actual source-field center is nonnegative. The exact finite Taylor expansion is

$$F = \mu + N^{\mathsf{T}} H N, \qquad \bar{H} = \tfrac{1}{2} \operatorname{Hess} F_0 \succ 0. \tag{AU5}$$

The five normals are differences of source coordinates and independent testing coordinates with the same real residues. All coefficients of H are bounded. The separated strict-Gram lemma proves AU5 tensor-SOS; the pure-source μ is a square. Positive residue slack is the ordinary scalar strict-Gram exit. This proves every noncommuting-residue test on this strong source domain. It does not import a broad noncommuting theorem by name.

One further residue case is needed. In C2 let α be weak and let each of n, m, C, A, D be a unit. Then the residue Laurent polynomial is

$$\bar{L} = h + q_{02} V/U + q_{20} U/V + q_{12} V + q_{21} U.$$

Its radial derivative is

$$(U\partial_U + V\partial_V)\bar{L} = q_{12} V + q_{21} U > 0. \tag{AU6}$$

It has no interior zero. Thus its finite noncommuting state matrices are positive definite. This explicitly supplies the noncommuting input in the retained finite-ratio proof as well.

196.1.2 Two retained full-parent inputs, with all hypotheses fitted

Strong and finite-edge sources. The actual retained proof *finite-pitchfork body* establishes all projective tests when d, C, A, D, α are units, with arbitrary nonnegative n, m, β and $\lambda = 0$. Its noncommuting call is replaced here by AU2–AU5. The only non-strict finite commuting edge has $u = 0$, $v_0 \neq 0$, weak m, and the W state ray. On that residue face,

$$Q(U,V) = UE(V) + U^2 B(V), \quad E(V) = q_{10} + hV + q_{12} V^2, \quad B(V) = q_{20} + q_{21} V + q_{22} V^2, \tag{AU7}$$

where $q_{10}, q_{12}, q_{21} > 0$. Physicality makes $E(V) \geq 0$ on $V > 0$. Thus either $E(v_0^2) > 0$, giving a strict four-normal Gram, or $E(V) = q_{12}(V - v_0^2)^2$. The displayed three-state complement in the actual *finite-edge terminal* is positive: its lower determinant has the positive factor

$$\alpha + 2\alpha C + 2\alpha A + 2\alpha D(1 + v_0^2) + D v_0^2 (1 + 2C + 2A) > 0.$$

Also $B(v_0^2) > 0$. Every hypothesis of that retained local theorem is therefore verified on the original source.

For specificity, its pure-testing event normalization makes $vW - X + uvY - uZ = 1$. The complete parent becomes $4m + u^2K(x, a, b, v, u^2)$ only as a formal source-center calculation. The actual tensor normals are

$$X - u^2x_c, \quad Y - ua_c, \quad Z - ub_c, \quad u(v - v_c), \quad u^2 - U_c. \tag{AU8}$$

The divided x-equation and the full three-variable Hessian are strict; the last scalar pivot is the positive $16(1 + v_0^2)^2B(v_0^2)/(v_0^2 \det M_0)$. The two signs of the optimized source constant give either the nonnegative source value at $U_c > 0$, or the retained $4m + \kappa u^2$ at $U_c = 0$. The retained theorem expands all first state normals and the complete state-quadratic tail. Its finite monic divisions do not divide a testing infinitesimal.

The same actual body supplies the four literal projective corners. In quarter-parent normalization their residue forms after the indicated whole event is retained are

$$\begin{array}{l|l} (0,0),\ W=1,\ m \text{ retained} & (n+\beta+\frac{1}{2})X^2 + K(u-Y)^2 + AY^2 + \alpha Z^2 \\ (\infty,0),\ X=1,\ n \text{ retained} & CW^2 + AY^2 + (D+\frac{1}{2})(W+v)^2 + \alpha Z^2 + (m+\beta+\frac{1}{2})(s+Z)^2 \\ (0,\infty),\ W=1,\ m \text{ retained} & (n+\beta+\frac{1}{2})X^2 + C(u-Y)^2 + (A+\alpha+\frac{1}{2})Y^2 + DZ^2 \\ (\infty,\infty),\ Z=1,\ \beta \text{ retained} & CW^2 + AY^2 + D(X-s)^2 + (n+\frac{1}{2})X^2 + (m+\alpha+\frac{1}{2})(Y-b)^2. \end{array} \tag{AU9}$$

Here $s = 1/u, b = 1/v$ in the reciprocal charts, $K = C + D + \frac{1}{2}$, and coefficient letters denote residues. Each displayed quadratic form is strict on its printed normal variables. The entire remainders belong to their square ideals, as shown by the actual polynomial expansions in the retained body. The two single-infinity edges have, among their positive terms, $\frac{1}{2}[(W + vX)^2 + (vY - Z)^2] + CD_vW^2 + AD_vY^2$ or $\frac{1}{2}D_u(X^2 + Y^2) + C(uW - Y)^2 + D(uX - Z)^2$, respectively, and are strict when the other angle is nonzero. These forms and AU7–AU8 exhaust all states and angles. We denote this exact all-tests input by S.

The unit-m, n weak-alpha source. The actual retained *weak-alpha unit-n body* assumes d, n, m, C, A, D all positive units and α weak in C2. AU6 supplies its noncommuting input. Its only remaining non-strict angle corner is $u = v = 0$. At $W = 1, Z$ infinitesimal, retain the whole α packet; the quadratic form

$$(n + \tfrac{1}{2})X^2 + m(v - X)^2 + K(u - Y)^2 + AY^2 \tag{AU10}$$

is strict in X, Y, u, v. At $Z = 1, W$ bounded the full quadratic jet Q_w in the retained body is positive semidefinite by physicality. Its X, Y block is positive, and it has no nonzero kernel with either $u = 0$ or $v = 0$: the positive n, m, A, D terms force all remaining variables to vanish in each case. Hence a singular angular Schur complement has a one-dimensional kernel with a finite nonzero real ratio $t_0 = u/v$.

The full homogeneous state jet at that ratio has a positive W, X, Y principal complement, already from $nX^2 + m(W - X)^2 + AY^2$. Moreover

$$q_{20}, q_{02} > 0, \quad q_{20}t_0^4 + ht_0^2 + q_{02} = 0, \quad q_{21}t_0^4 + q_{12}t_0^2 > 0. \tag{AU11}$$

These are exactly the actual *finite-ratio corner terminal* hypotheses. Its source-only critical centers and complete normal-tail allocation use the undivided tensor rows

$$X - vx_c, \quad Y - vy_c, \quad u - vt_c, \quad v(W - w_c), \quad v^2 - V_c.$$

The source-selected t_0 imposes no restriction on the actual independent quotient u/v. The explicit 55-monomial allocation retains $W - w_c$ in bounded coefficients rather than replacing it by a mixed quotient. Its strict four-variable Hessian and positive last pivot prove the whole parent. The retained complementary-state argument and full projective forms cover every other state, sign and literal endpoint. We denote this exact all-tests input by CU. Only the printed unit hypotheses are used; no weak-m or weak-n conclusion is inferred from it.

196.1.3 Exact symmetries and the source partition

Both transformations below preserve the core and nonnegative event coefficients, actual physicality, finite nonzero contact, and pure total-scale normalization. They act separately in source and testing fields, so transport a full tensor certificate.

For C1 let

$$\sigma_1 = (\lambda\ n)(m\ C)(A\ \alpha)(D\ \beta), \qquad G_v(W,X,Y,Z) = (vW - X, -W - vX, Z - vY, Y + vZ).$$

The entire raw parent has the polynomial identity

$$u^2 F_{\sigma_1 c}(1/u, v; G_v\psi) = (1+v^2)F_c(u,v;\psi), \qquad G_v^{\mathsf{T}} G_v = (1+v^2)I. \tag{AU12}$$

Homogenization interchanges the two Alice angle coordinates; the Bob-homogeneous G has norm factor $v_0^2 + v_1^2 > 0$. Thus literal endpoints are transported without dividing zero.

For C2 let

$$\sigma_2 = (\lambda\ \beta)(m\ D)(C\ \alpha), \qquad J(W,X,Y,Z) = (Z,X,Y,W).$$

The full identity is

$$u^2 v^2 F_{\sigma_2 c}(1/v, 1/u; J\psi) = F_c(u,v;\psi). \tag{AU13}$$

It is the physical party interchange with both first observables negated, followed by the local real X basis changes. The homogeneous identity includes both projective endpoints. In particular the source squared half-angles transform as $(U,V) \mapsto (1/V, 1/U)$.

The two actual forbidden weak faces are

$$\begin{aligned} h|_{\lambda=\beta=A=0} &= -(1+2n)(2D+2C+1)(2\alpha+1+2m), \\ h|_{\lambda=n=m=C=0} &= -(2D+1)(2\beta+1)(2A+2\alpha+1). \end{aligned} \tag{AU14}$$

In C2 with A weak, the only positive Laurent terms are $q_{01}/U + q_{02}V/U + q_{12}V$, which tend to zero along $U = t^2, V = 1/t,\ t \to \infty$. In C1 with both m, C weak the only positive term is q_{10}/V, which tends to zero as $V \to \infty$. The negative constants contradict physicality. Thus C2 has A unit, and C1 has at least one of m, C unit. These are statements about physical residue sources, not an assumption inherited from a prior routing complement.

C2. If α is weak, WA1–WA6 force C, A, D to be units. There are precisely three disjoint alternatives:

n, m units	CU
n unit, m weak	WA1−WA27
n weak, m arbitrary	WN1−WN44.

(AU15)

If C is weak, apply σ_2 and the same table. Every unit requirement in the transformed table is a consequence of the transformed physical source, including its required $D' = m$ bound; it is not assumed for an excluded original case.

It remains that C, A, α are units. The actual dictionary then has

$$q_{01} = ma_1,\ q_{02} = mDa_2,\ q_{10} = a_3,\ q_{12} = Da_4,\ q_{20} = na_5,\ q_{21} = a_6,$$

where every a_i is a positive bounded unit. The contact equation bounds $U, V^{-1}, m/U, DV$. The two angular equations give exactly

$$\begin{aligned} DV^2(a_4 + a_2 m/U) &= a_3 + na_5 U, \\ U^2(a_6 + na_5/V) &= m(a_1 + a_2 DV). \end{aligned}$$

All parentheses are positive bounded units, so $DV^2 \asymp 1$, $U^2 \asymp m$. Consequently the exhaustive all-tests dispatch is

D unit, m arbitrary	S
D weak, m unit	σ_2(S)
D, m weak	RC1−RC25.

(AU16)

RC is the new direct full-parent C2 reciprocal theorem; it permits arbitrary $n \geq 0$. No unit-n condition has entered this part of the partition.

C1. If α is weak, WN applies directly, with all its common source-entry implications proved in WA1–WA10. If A is weak, apply σ_1, making $\alpha' = A$ weak, and use WN on the transformed physical source. Otherwise A, α are units. By AU14, orient with σ_1, if necessary, so that C is a unit. The exact dictionary becomes

$$\begin{gathered} q_{01} = ma_1,\ q_{02} = mDa_2,\ q_{10} = a_3, \quad q_{12} = \beta m a_4 + Da_5, \quad q_{21} = a_6, \quad q_{22} = \beta a_7, \\ U^2(a_6 + \beta a_7 V) = m(a_1 + a_2 DV), \\ a_3 = V^2\{mDa_2/U + \beta m a_4 + Da_5 + \beta a_7 U\}. \end{gathered} \tag{AU17}$$

The contact equation again bounds $U, V^{-1}, m/U, DV$. If m is weak, U is infinitesimal. If m is a unit, U is a positive unit. Indeed, the contact equation also bounds βUV; if U were infinitesimal, then $U^2(a_6 + \beta a_7 V)$ would be infinitesimal, contradicting the positive unit ma_1 on the right of AU17. For unit m, the last braces are comparable with $D + \beta$. For weak m, bounded V forces m/U weak in the preceding equation, and then the last equation forces D unit; conversely unit D forces bounded V. Thus the following source dispatch is exhaustive:

D unit, m arbitrary	S
D weak, m unit, β unit	σ_1(S)
m unit, D, β weak	PP1−PP19
m, D weak, β arbitrary	CR1−CR39.

(AU18)

In the second line all of m, C, A, α are units before reflection, so the reflected strong theorem has every required unit. PP is the new C1 projective-pitchfork theorem, with actual contact U unit and V unbounded. CR is the new C1 reciprocal all-tests theorem, including unit β and literal allowed $D = 0$ or $\beta = 0$. Its m, D weak scope is exactly the last line. No boundary is promoted by a density argument.

AU15–AU18 cover every permitted bounded positive-d canonical source. Each leaf proves the entire original parent at every independent projective test, with the exact same source coefficients. The full-parent symmetries, positive pure-source d scaling, positive pure-testing homogeneous factors and complex-state split restore the assertion in the opening paragraph. This completes A5 at its canonical-entry interface.

Proof-use scope lemma. The bare source domains in this union do not silently discard the prior-complement phrases retained in the earlier statements. Their removal here follows by checking the actual proofs: RC1 uses only the two equations in AU16; RC2–RC5 use the actual kernel equations, positive native gaps, and the listed unit bounds; RC6–RC25 are full-parent identities with those bounds. PP1–PP5 use only AU17, the actual kernel and angle jets, and its listed unit bounds; PP6–PP19 and its subsequent outer identities use no source exclusion. CR1–CR8 likewise derive every shape estimate from AU17 and the actual source equations, and CR9–CR39 use only those estimates. WA1–WA10 prove their own source entry, while WA11–WA27 and WN1–WN44 require only the printed weak/unit alternatives. The retained S and CU bodies state physical coefficient domains directly and have no prior-admission premise. Consequently the same proofs establish the implications

$$\text{physical native source and listed scale bounds} \implies \text{all-tests certificate} \tag{AU19}$$

without any earlier-terminal exclusion. This is a proof-use strengthening on fixed native source charts; it does not identify an arbitrary contact with such a chart. That sign/native entry implication remains an explicit global dependency.

Evidence and claim boundary. The dependency receipt identifies the actual finite-edge, finite-ratio, fit and strict-Gram proof bodies, not just their historical labels. The focused checker verifies both complete parent symmetries, the forbidden faces, both exact stationarity dictionaries and the finite weak/unit dispatch. The source comparisons, full Hessian proof and typed all-tests applications

are the analytical argument above. This theorem does not assert weak-total-scale C1/C2 closure, global support-cone entry, or a final finite-dimensional lift. Those require their separately proved source and testing unions.

Chapter 197

A fresh complete outer-testing certificate for the original R2 source

Component `p8f8780350b13`
Source: `NEW/global/r2_outer_testing_join_REDERIVED.tex`
SHA256: `8f8780350b13c682d2f4e9dc7d20b1dbd0c7b1a3b7a4828d78e472d28838f298`
Inventory use: R2O1-R2O24

197.1 A fresh complete outer-testing certificate for the original R2 source

This proof treats every independent testing pair outside the positive double-infinitesimal angle chart of the original R2 source. It uses an explicit full-parent Bell reflection and the two-current and six-root identities that were freshly proved for R1. Only those universal algebraic identities and their real matrix arguments are reused. No assertion that the reflected source is an original R1 source is made.

197.1.1 Actual source and exact reflected parent

The source is the original weak-C2 tuple, with $d = 1$, $\lambda = \beta = 0$, the six coefficients in (RER1), all original strict gaps, $n \geq 0$, $a, B, t_s > 0$, and $0 < r, h < 1$. R2 means

$$t_s \text{ unbounded}, \qquad B, z = aB, y = t_s B \text{ infinitesimal}. \tag{R2O1}$$

Put $S = 1/t_s$, $l = 1 - h$, $e = 1 - r$, $R_s = 1 + hz$, $g_s = R_s - ae > 0$, and $k = (1 + ar)/(1 + a)$. The rederived exact source lemma (RER1)–(RER7) gives

$$k \text{ a positive bounded unit}, \qquad k < 1, \qquad SCl \sim SAh \sim D/S \sim k/2, \qquad \Delta/(1 + a) \sim 1. \tag{R2O2}$$

It also gives ay infinitesimal, $Q \sim 1$, and the exact expressions

$$n = \frac{eQ - BK_s}{2r\Delta}, \qquad m = \frac{a^2 e(1 + y)cs_c}{2g_s\Delta}, \qquad BK_s = (1 + a)y(1 + hS)(1 + z), \tag{R2O3}$$

where $c = 1 + zr \sim 1$ and $s_c = 1 + B(l + ar) \sim 1$. There is no positive lower bound assumed on r, h, l, or g_s.

For arbitrary finite testing angles define the unnormalized testing state transformation

$$\begin{aligned} G\psi = (&uvW - uX - vY + Z,\ -vW + X - uvY + uZ,\\ &- uW - uvX + Y + vZ,\ W + vX + uY + uvZ). \end{aligned} \tag{R2O4}$$

It has $G^T G = D_u D_v I$. The Bell transformation on observables is

$$(A'_0, A'_1, B'_0, B'_1) = (-B_1, -B_0, -A_1, -A_0).$$

After the displayed real local basis gauges it sends the angles to $(u', v') = (v, u)$ and the source coefficients to

$$(\lambda', n', m', C', A', D', \alpha', \beta') = (\alpha, m, n, C, D, A, \lambda, \beta). \tag{R2O5}$$

Direct substitution in the whole parent gives

$$F_{c'}(v, u; G\psi) = D_u D_v F_c(u, v; \psi). \tag{R2O6}$$

The checker verifies the coefficient of every original packet in this single full polynomial identity, as well as $G^T G$. The transformed $\lambda' = \alpha$ event is kept as its whole positive square. Because $D_u D_v > 0$ is pure testing, a certificate on the left restores the right by dividing each root by $\sqrt{D_u D_v}$ within K_t. No source or testing contact is identified in this operation.

In the next two subsections write the transformed testing angles again as u, v, and denote the reflected coefficients by

$$\mathsf{C} = C, \quad \mathsf{D} = A, \quad \mathsf{A} = D, \quad \mathsf{m} = n, \quad \mathsf{n} = m.$$

Thus the symbols on the right refer to the original R2 source. Define the pure-source quantities

$$g_* = \frac{\mathsf{CD}}{\mathsf{C} + \mathsf{D}}, \qquad h_* = \frac{\mathsf{D}}{\mathsf{C} + \mathsf{D}}, \qquad \gamma = \mathsf{A} g_*. \tag{R2O7}$$

If $U_C = SCl$ and $U_A = SAh$, then exactly

$$S g_* = \frac{U_C U_A}{h U_C + l U_A} \sim k/2, \qquad \gamma \sim k^2/4. \tag{R2O8}$$

The denominator is a positive bounded unit even when h or l is infinitesimal. In particular $1/\mathsf{C}$, $1/\mathsf{D}$, $1/g_*$ are infinitesimal, $(\mathsf{C} + \mathsf{D})^{-1/2} = O(\sqrt{S})$, and $0 < h_* < 1$. These are the complete source bounds needed for the first two outer current identities.

197.1.2 Universal outer current identities

Set

$$R_t = uW - Y, \quad C_t = v(uX - Z), \quad B_t = X + uZ, \quad J = R_t + C_t, \quad N = R_t + h_* C_t,$$

and let K be the exact signed quarter-core

$$K = \tfrac{1}{2} J^2 + \tfrac{1}{2} D_u (vY - Z)^2 + \tfrac{1}{2} D_v B_t^2 - \tfrac{1}{2} D_u D_v Z^2. \tag{R2O9}$$

All unselected original events, including λ', are retained as whole nonnegative packets.

If $1/v$ is bounded, put

$$K_0 = 2\sqrt{g_*} C_t, \quad B_0 = \sqrt{2D_v/D_u}\, B_t, \quad c_u = D_u^{-1/2}, \quad t_u = u D_u^{-1/2}, \quad \eta_v = \sqrt{D_v}/v.$$

The entire selected-parent identity is

$$\begin{aligned} 4\mathsf{C} D_v R_t^2 + 4\mathsf{D} J^2 + 4K &= 4(\mathsf{C} + \mathsf{D}) N^2 + 4\mathsf{C} v^2 R_t^2 + 2J^2 + 2D_u (vY - Z)^2 \\ &\quad + (K_0, B_0) \begin{pmatrix} 1 - c_u^2 \eta_v^2/(2g_*) & t_u \eta_v/\sqrt{2g_*} \\ t_u \eta_v/\sqrt{2g_*} & 1 \end{pmatrix} \begin{pmatrix} K_0 \\ B_0 \end{pmatrix}. \end{aligned} \tag{R2O10}$$

The finite Gram is bounded and has real residue I, because c_u, t_u and η_v are bounded while $1/g_*$ is infinitesimal. The actual separated bounded-Gram lemma makes it tensor SOS. The roots themselves need not be bounded. This gives every state and every other finite angle when transformed Bob is noninfinitesimal, equivalently when the original Alice angle is noninfinitesimal.

For positive infinitesimal transformed v, put $q = g_*^{-1/2}$ and define

$$K_1 = 2\sqrt{g_*} C_t + qY, \quad B_1 = B_t - uvY, \quad V_0 = vY, \quad Y_1 = q\sqrt{D_u} Y,$$

$$L_0 = \tfrac{1}{2} q c_u (K_1 - c_u Y_1) - t_u v B_1 + c_u v V_0.$$

Then $L_0 = \sqrt{D_u}\, v(vY - Z)$ exactly, and

$$\begin{aligned} &4\mathsf{C} D_v R_t^2 + 4\mathsf{D} J^2 + 4\mathsf{A} D_u D_v Y^2 + 4K \\ &\quad = 4(\mathsf{C}+\mathsf{D}) N^2 + 4\mathsf{C} v^2 R_t^2 + 2J^2 + \mathcal{G}, \\ \mathcal{G} &= K_1^2 + 2D_v(B_1^2 + V_0^2) + (4\gamma D_v - c_u^2) Y_1^2 \\ &\quad + 2qv(K_1 - c_u Y_1) V_0 - 2L_0^2. \end{aligned} \tag{R2O11}$$

This is the complete four-root Gram, including all normal terms. Its residue in the ordered frame (K_1, B_1, V_0, Y_1) is

$$\operatorname{diag}(1, 2, 2, k_0^2 - c_{u,0}^2). \tag{R2O12}$$

Subscript 0 denotes real residue. Hence it is strict whenever its last entry is positive. This includes every unbounded transformed u and every bounded positive-unit transformed u when $k_0 = 1$. All inverse and square-root factors in these identities belong to one field at a time.

197.1.3 The remaining unit angle and both coefficient openings

It remains to treat transformed v infinitesimal, transformed u a bounded positive unit, and $k_0 < 1$. The source identity

$$1 - k = \frac{ae}{1+a}, \qquad 0 < ae < R_s \sim 1 \tag{R2O13}$$

shows that a, e are positive bounded units: if a were infinitesimal or unbounded, or if e were infinitesimal while a is bounded, then $1 - k$ would be infinitesimal. Thus in the current chart a is a positive unit, e is a positive unit, and both $1 + a$ and Δ are units.

Use the opening-safe source coordinates

$$j = R_s/a, \qquad p = e, \qquad j - p = g_s/a > 0. \tag{R2O14}$$

They satisfy $0 < p < \min(1, j)$, j, p are positive bounded units, and the residue of $(j + 1 - p)/(j + 1)$ equals k_0. The exact relative critical coefficient identities are

$$\frac{\mathsf{m}}{jp/[2(1-p)(j+1)]} = \left(Q - \frac{BK_s}{e}\right) \frac{R_s + a}{R_s \Delta}, \qquad \frac{\mathsf{n}}{p/[2(j-p)(j+1)]} = (1+y) c s_c \frac{R_s + a}{\Delta}. \tag{R2O15}$$

Their right sides have residue one. Indeed BK_s/e is infinitesimal here by (R2O3), and all other normalized factors have the limits stated in (RER). No division by r, g_s, n, or an infinitesimal gap was used to assert a bounded error. The displayed quotients cancel those openings exactly before the limit is taken. In particular the critical chart itself forces original $n > 0$, even though the theorem as a whole retains $n = 0$.

Choose positive real caps $m_c \le \mathsf{m}$ and $n_c \le \mathsf{n}$, with the choices specified below. Keep both nonnegative excesses as the original whole packets. Define six rows

$$\widehat{N} = \sqrt{\mathsf{C}+\mathsf{D}}\, N, \quad P = C_t/\sqrt{S}, \quad Y_0 = \sqrt{S}\, Y, \quad X, \quad Z, \quad V = vY. \tag{R2O16}$$

They can be unbounded or linearly dependent. If $\delta = (\mathsf{C}+\mathsf{D})^{-1/2}$, then $R_t = \delta \widehat{N} - h_* \sqrt{S} P$. The selected entire quarter-parent is exactly

$$\begin{aligned} \mathcal{P}_{\mathrm{sel}} &= \widehat{N}^2 + \mathsf{C} v^2 R_t^2 + \tfrac{1}{2} J^2 + (S g_*) P^2 + (\mathsf{A}/S) D_u D_v Y_0^2 \\ &\quad + m_c \{(v/u) R_t + (D_u/u) V - X - uZ\}^2 + n_c D_u D_v X^2 \\ &\quad + \tfrac{1}{2} D_u (V - Z)^2 + \tfrac{1}{2} D_v (X + uZ)^2 - \tfrac{1}{2} D_u D_v Z^2. \end{aligned} \tag{R2O17}$$

The full-state row relation is

$$P Y_0 = V(uX - Z). \tag{R2O18}$$

Add $2b(PY_0 - V(uX - Z)) = 0$ for a real b. Keep the two displayed positive squares $\mathsf{C}v^2R_t^2$ and $J^2/2$ whole. The remaining Gram is bounded: $u, 1/u$ are bounded, Sg_* and A/S are units, and the R_t contributions in the m_c square have coefficients $\delta v/u$ and $h_*\sqrt{S}v/u$, both infinitesimal. Its residue is the direct sum of a unit pivot, the matrix

$$\begin{pmatrix} k_0/2 & b \\ b & k_0(1+u_0^2)/2 \end{pmatrix}, \tag{R2O19}$$

and the following real quadratic form on (X, Z, V):

$$Q_3 = m_c\{(D_u/u)V - X - uZ\}^2 + n_c D_u X^2 + \tfrac{1}{2}D_u V^2 - D_u VZ + \tfrac{1}{2}(X+uZ)^2 - 2bV(uX - Z). \tag{R2O20}$$

In this formula $u = u_0 > 0$ is real. We now give all finite-cap choices; no opening of either source coefficient is suppressed.

If $p_0 < 1$ and $j_0 > p_0$, use initially

$$m_0 = \frac{j_0 p_0}{2(1-p_0)(j_0+1)}, \quad n_0 = \frac{p_0}{2(j_0-p_0)(j_0+1)}, \quad b = k_0/2.$$

The three leading principal minors of Q_3 are

$$m_0 + n_0 D_u + \tfrac{1}{2}, \quad \frac{p_0 u_0^2 D_u(j_0+1-p_0)}{4(j_0-p_0)(j_0+1)^2(1-p_0)}, \quad \frac{p_0 u_0^2 D_u(j_0+1-p_0)^2}{8(j_0-p_0)(j_0+1)^3}, \tag{R2O21}$$

all strictly positive. Matrix (R2O19) has determinant $k_0^2u_0^2/4 > 0$. Choose real caps slightly below m_0, n_0; openness of real positive definiteness and (R2O15) ensure they remain strict and are below the actual coefficients.

If $j_0 = p_0 < 1$, the coefficient n is unbounded and $m_0 = j_0^2/[2(1-j_0)(j_0+1)]$ is finite positive. Take $b = 1/[2(j_0+1)]$. On the kernel $X = 0$ of the n packet, the first diagonal and determinant of Q_3 on (Z, V) are

$$u_0^2(m_0 + 1/2) > 0, \qquad u_0^2/[4(j_0+1)^2] > 0. \tag{R2O22}$$

Choose $m_c < m_0$ sufficiently close and then a sufficiently large real n_c. A real block Schur complement makes the full matrix strict.

If $p_0 = 1 < j_0$, the coefficient m is unbounded and $n_0 = 1/[2(j_0^2-1)]$ is finite positive. Choose a real b with

$$\frac{j_0}{2(j_0+1)} < b < \min\left\{\frac{k_0\sqrt{D_u}}{2}, \frac{j_0}{2(j_0-1)}\right\}.$$

The interval is nonempty because $u_0 > 0$. On the kernel $(D_u/u)V - X - uZ = 0$ of the m packet, the first diagonal and determinant in (X, V) are

$$\frac{D_u}{2(j_0^2-1)} > 0, \qquad -\frac{D_u^2[2b(j_0-1)-j_0][2b(j_0+1)-j_0]}{4u_0^2(j_0^2-1)} > 0. \tag{R2O23}$$

Choose $n_c < n_0$ sufficiently close, then a sufficiently large real m_c. Matrix (R2O19) and the full Q_3 are strict.

If $p_0 = j_0 = 1$, both coefficients are unbounded and $k_0 = 1/2$. Choose $1/4 < b < \sqrt{D_u}/4$. On the common event kernel $X = 0$, $Z = D_uV/u_0^2$, the remaining form is

$$\frac{D_u(4b-1)}{2u_0^2}V^2 > 0. \tag{R2O24}$$

Sufficiently large real caps on both events make the full form strict. The finite-cap step in each case is the elementary real fact that a quadratic form positive definite on the kernel of a nonnegative event Gram becomes positive definite after a sufficiently large multiple of that event Gram is added; a block Schur complement proves it.

These four cases exhaust $0 < p_0 \le \min(1, j_0)$ with positive j_0, p_0. The separated bounded-Gram lemma now proves the entire selected parent (R2O17), with every state normal retained. Restore the original excesses and all unused packets. The real caps are used only on this testing chart: no globally physical capped source is asserted, so this argument does not conflict with the finite-contact cap exclusion (CAP15).

197.1.4 Exact outer and projective join

Return to the original testing angles. The full-parent sign gauges (SJ3) permit $u, v \geq 0$ without changing source coefficients. Literal zeros and projective infinities are the exact signed-core squares (SJ10), plus the untouched event squares. For finite positive angles:

- if the original u is noninfinitesimal, use (R2O10);
- if the original u is infinitesimal and v is unbounded, use (R2O11)–(R2O12);
- if the original u is infinitesimal and v is a positive bounded unit, use (R2O11) when $k_0 = 1$, and the complete six-root argument (R2O13)–(R2O24) when $k_0 < 1$.

These are exactly all finite positive angle pairs outside the case in which both are infinitesimal. Every argument is a whole-parent state Gram, so no additional state chart or state-ratio restriction is present. Equation (R2O6) restores the original frame and positive pure-testing factor. The zero state and the real/imaginary split for complex states are immediate. No source-to-testing rate comparison was made.

Assembly interface and remaining obligation. The original R2 outer testing region, every state, and all literal projective endpoints are proved here. A full-parent certificate on the remaining original positive u, v double-infinitesimal chart, for the same unchanged source and all states, combines with this result to give the whole original R2 row. That local certificate is a separate hypothesis until its actual proof is independently accepted. This outer theorem does not itself close any other source row or the global finite semidefinite-lift criterion.

Chapter 198

A rederived three-event wall and its exact contact exclusion

Component p23a094ff816d
Source: NEW/global/three_event_wall_and_contact_exclusion_REDERIVED.tex
SHA256: 23a094ff816df869f32d663941bd03ec9f3cc2b321c286c6304ca903a8da6632
Inventory use: Explicit physical event rotation and exact scale partition; root analytically accepted and independently replayed PASS17

198.1 A rederived three-event wall and its exact contact exclusion

This section is a fresh proof from the recovered full parent. It does not use a historical admission or any missing local certificate. The selected three-event cone is given an explicit matrix description, its separated tensor-SOS assertion is proved by finite-dimensional duality, and its intersection with the actual finite-contact source domain is shown to be empty. The last conclusion concerns the source contact, not a comparison between a source contact and an independent test.

198.1.1 The selected full parent

Write $D_u = 1 + u^2$, $D_v = 1 + v^2$, and retain the coefficient order $(d, \lambda, n, m, C, A, D, \alpha, \beta)$. The original complete parent is

$$\begin{aligned} F = {} & 4\lambda D_u (W + vX)^2 + 4nD_u D_v X^2 + 4m(vW - X + uvY - uZ)^2 \\ & + 4CD_v(uW - Y)^2 + 4AD_u D_v Y^2 + (4D + 2d)(uW + uvX - Y - vZ)^2 \\ & + (4\alpha + 2d)D_u(vY - Z)^2 + (4\beta + 2d)D_v(X + uZ)^2 - 2dD_u D_v Z^2. \end{aligned} \tag{CAP1}$$

All nine coefficients are nonnegative. Let F_* be the selected parent with $\lambda = n = m = \alpha = \beta = 0$, so that only the consecutive event triple (D, C, A) remains. For $d, C, A, D \geq 0$ define

$$\Delta = 8ADC - 4ADd - d^2\{d + 2(C + A + D)\}. \tag{CAP2}$$

Proposition 198.1 (Exact physical wall and strict interior-angle positivity)**.** *If $d > 0$, the selected parent F_* is positive semidefinite at every projective measurement pair if and only if $\Delta \geq 0$. Whenever this holds, its state matrix is positive definite at every finite pair $u, v \neq 0$. In particular, the equality wall $\Delta = 0$ has no finite noncommuting contact.*

Proof. Let $P_*(u, v)$ be the symmetric state matrix. Direct expansion of the full polynomial (CAP1), without a state or angle truncation, gives

$$\det P_* = 16D_u^2 D_v^2 u^2 v^2 \{d\Delta + 4CDd(d + 2A)v^2 + 4CAd(d + 2D)u^2\}. \tag{CAP3}$$

The focused checker rebuilds P_* from its state Hessian and verifies this entire identity. Its restriction to $Z = 0$ is the sum of the nonnegative event packets and

$$2d\{(uW + uvX - Y)^2 + D_u v^2 Y^2 + D_v X^2\}. \tag{CAP4}$$

For $uv \neq 0$ these three squared linear forms are independent. Hence the upper 3×3 state block is positive definite. A Schur complement taken in this one ordered field proves that $P_* \succeq 0$ if and only if its determinant is nonnegative, and that a positive determinant gives positive definiteness. This is a statement about ordinary field-valued source or testing matrices; it takes no inverse in a tensor product.

Suppose $\Delta < 0$. Put $r = 4CDd(d + 2A)$ and $s = 4CAd(d + 2D)$, both nonnegative. Taking

$$u^2 = v^2 = \frac{-d\Delta}{2(r + s + 1)} > 0$$

makes the expression in braces in (CAP3) negative. This is an explicit ordered-field witness to nonphysicality, so no limiting argument is needed. Conversely, $\Delta \geq 0$ makes all three terms in braces nonnegative. It also forces $C, A, D > 0$: if one vanished, (CAP2) would be strictly negative because $d > 0$. Thus $r, s > 0$ and the determinant is strictly positive whenever $uv \neq 0$.

At an actual commuting angle, including a projective infinity, one party's two reflections commute. Simultaneous diagonalization makes the CHSH operator on each local eigenspace either ± 2 times one remote reflection. The signed core $I - S/2$ is then positive semidefinite. Every event packet is positive semidefinite too. This covers all literal endpoints directly. The argument holds over every real closed field, because it uses real diagonalization of symmetric matrices and explicit ordered-field identities. If $d = 0$, every parent in (CAP1) is its event-square sum separately. □

198.1.2 A small exact lift of the selected source cone

Set

$$p = 4C + 4D + 2d, \qquad r = 4D + 2d, \qquad B(d, C, A, D) = \begin{pmatrix} p + 4A & 4D & -p \\ 4D & 4D & -r \\ -p & -r & p \end{pmatrix}. \tag{CAP5}$$

For nonnegative coefficients the selected source is physical exactly when $B \succeq 0$. Here is a direct verification of this matrix description. When $d > 0$, use the positive last diagonal entry p. The Schur complement of that entry is

$$\begin{pmatrix} 4A & -2d \\ -2d & 4b/p \end{pmatrix}, \qquad b = 4CD - 2Dd - d^2, \qquad \det B = 8\Delta. \tag{CAP6}$$

Moreover

$$\Delta = 2Ab - d^2(d + 2C + 2D). \tag{CAP7}$$

If $\Delta \geq 0$, then $A, b > 0$, and (CAP6) is positive semidefinite. If $B \succeq 0$, its determinant is nonnegative, hence $\Delta \geq 0$. For $d = 0$ one has the exact quadratic identity

$$(y, z, x)B(0, C, A, D)(y, z, x)^T = 4C(y - x)^2 + 4Ay^2 + 4D(y + z - x)^2. \tag{CAP8}$$

Thus the entire selected physical cone, including all zero faces, is

$$\mathcal{C} = \{(d, C, A, D) \in \mathbb{R}^4_{\geq 0} : B(d, C, A, D) \succeq 0\}. \tag{CAP9}$$

It has a strict feasible point $(1, 2, 2, 2)$: the leading principal minors of B there are $26, 144, 280$.

Lemma 198.2 (Separated certificate for the selected wall)**.** *For arbitrary labelled real closed fields $K_c, K_t \supset \mathbb{R}$, every source $c \in \mathcal{C}_{K_c}$, and every projective angle pair and real state in K_t, the complete selected parent is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$. The full parent (CAP1) has the same property if its selected coefficients satisfy (CAP9).*

Proof. We first establish the exact dual representation used in this lemma. For a real linear functional ℓ nonnegative on $\mathcal{C}$, there are $\mu \in \mathbb{R}^4_{\ge 0}$ and a real symmetric $Q \succeq 0$ such that, for all $c = (d, C, A, D)$,

$$\ell(c) = \mu \cdot c + \operatorname{tr}(B(c)Q). \tag{CAP10}$$

Indeed let $\mathcal{D}$ be the cone of functionals on the right. It is closed: if a sequence of these functionals converges, evaluation at $c_0 = (1, 2, 2, 2)$ bounds both $\sum \mu_i$ and $\operatorname{tr} Q$, because all coordinates of c_0 and the smallest eigenvalue of $B(c_0)$ are strictly positive. Positive semidefinite matrices with bounded trace are bounded, so there is a convergent subsequence of the representing pairs. The dual cone $\mathcal{D}^*$ is exactly (CAP9). Finite-dimensional separation of closed convex cones gives $\mathcal{D} = \mathcal{C}^*$ and proves (CAP10). This statement has a fixed finite first-order formulation using the principal-minor inequalities for a 3×3 matrix. Consequently it holds over every real closed field by transfer.

Fix an arbitrary testing tuple in K_t. Its homogeneous selected parent is a linear functional ℓ_t on the four source coefficients. The physical equivalence just proved says that $\ell_t \ge 0$ on $\mathcal{C}_{K_t}$. Apply (CAP10) in K_t to obtain $\mu_t \ge 0$, $Q_t \succeq 0$. Evaluating the resulting linear coefficient identity in separated fields gives

$$F_*(c, t) = \sum_{i=1}^{4} c_i \otimes \mu_{t,i} + \sum_{j,k=1}^{3} B(c)_{jk} \otimes (Q_t)_{jk}. \tag{CAP11}$$

The first terms are squares by separate square roots. Factor $B(c) = UU^T$ within K_c and $Q_t = VV^T$ within K_t. The last term in (CAP11) is exactly

$$\sum_{a,b} \left(\sum_j U_{ja} \otimes V_{jb} \right)^2 .$$

This proves the tensor certificate with no mixed denominator. The five omitted event packets in (CAP1) are separate positive-source factors times testing squares and can be restored exactly. Every literal projective endpoint is already included in the homogeneous testing tuple. Splitting a complex state into its real and imaginary parts gives the same assertion for complex states. □

198.1.3 Every actual finite contact lies strictly below all eight walls

The fixed event cycle is

$$(a_0, \dots, a_7) = (\lambda, D, C, A, \alpha, m, \beta, n).$$

For indices modulo eight put

$$\Delta_i = 8a_i a_{i+1} a_{i+2} - 4d a_i a_{i+2} - d^2\{d + 2(a_i + a_{i+1} + a_{i+2})\}. \tag{CAP12}$$

The cyclic equivalence can be verified without an unspecified symmetry import. In the CHSH convention $S = A_0B_0 - A_0B_1 + A_1B_0 + A_1B_1$, the Bell relabeling

$$(A'_0, A'_1, B'_0, B'_1) = (B_1, -B_0, -A_0, -A_1) \tag{CAP13}$$

preserves S and maps each event in the displayed cycle to the next. For example the transformed $\lambda = (+, + \mid 0, 1)$ is the original $D = (-, + \mid 1, 1)$. The complete eight-event permutation and the CHSH identity are checked exactly. Relabeling preserves whether each local pair is noncommuting, so the strict assertion of the first proposition applies to every cyclic triple.

Theorem 198.3 (Automatic contact exclusion)**.** *Let $d > 0$, let every event coefficient be nonnegative, and suppose the original parent (CAP1) has a zero on a nonzero source-field state at an actual finite noncommuting source pair. Then*

$$\Delta_i < 0 \qquad (i = 0, \dots, 7). \tag{CAP14}$$

The conclusion needs only the actual zero and coefficient signs; no extra assumption about contact stationarity or historical prior exclusions is required.

Proof. If some $\Delta_i \geq 0$, the selected parent for that triple is positive definite at the specified pair. The full parent is this selected parent plus five nonnegative event packets. It is therefore positive on every nonzero state at that pair, contradicting the assumed zero. This uses only ordinary ordered-field positivity at the source contact. It does not evaluate a source contact in an independent testing field. □

More generally, suppose a proposed cap keeps the same signed coefficient d and removes nonnegative amounts e_j from event coefficients. At an actual zero t_c one has the exact identity

$$F_{\mathrm{cap}}(t_c) = -4 \sum_j e_j\, \omega_j(t_c) R_j(t_c)^2, \tag{CAP15}$$

where ω_j is the positive angle factor multiplying the event row in (CAP1). Thus a globally physical cap can remove a strictly positive amount only from an event whose row vanishes at that contact. If one removed positive packet has nonzero row, (CAP15) is already a strictly negative physicality witness. In the special three-event cap, (CAP14) rules out even the case in which all removed rows vanish. The same argument permits smaller retained event coefficients: add their nonnegative excesses back as well. It does not apply to a modification of d, whose difference is a signed CHSH term rather than a nonnegative event packet.

198.1.4 Exact consequences for source normalization and routing

For $d > 0$ put

$$\delta_{\mathrm{tot}} = \frac{d}{d + \sum_{i=0}^{7} a_i}, \qquad b_i = a_i/d. \tag{CAP16}$$

In a real closed extension, "bounded" means bounded in absolute value by a real number, and "infinitesimal" means smaller in absolute value than every positive real number. Since all b_i are nonnegative and the sum is finite,

$$\begin{aligned} &\delta_{\mathrm{tot}} \text{ has positive real residue} &&\iff \text{every } b_i \text{ is bounded,} \\ &\delta_{\mathrm{tot}} \text{ is infinitesimal} &&\iff \text{at least one } b_i \text{ is unbounded.} \end{aligned} \tag{CAP17}$$

These alternatives are disjoint and exhaustive because $0 < \delta_{\mathrm{tot}} \leq 1$. They are invariant under a positive source rescaling and under Bell relabeling. Normalizing $d = 1$ is legal in the coefficient field, and multiplying each tensor root by the pure-source factor $\sqrt{d}$ restores the original scale.

There cannot be three consecutive unbounded b_i at an actual finite contact. Indeed division of (CAP12) by its positive triple product gives

$$\frac{\Delta_i}{a_i a_{i+1} a_{i+2}} = 8 - \frac{4d}{a_{i+1}} - \frac{d^3}{a_i a_{i+1} a_{i+2}} - 2d^2 \left\{ \frac{1}{a_i a_{i+1}} + \frac{1}{a_i a_{i+2}} + \frac{1}{a_{i+1} a_{i+2}} \right\}. \tag{CAP18}$$

If the three ratios a_j/d were unbounded, its residue would be 8, contradicting (CAP14). Equivalently, every consecutive triple contains an event bounded relative to d.

If a later chart uses $C = 2$, the rescaling factor is $2/C$, and its new signed weight is $2d/C$; (CAP16) remains unchanged. The implication d/C infinitesimal $\Rightarrow \delta_{\mathrm{tot}}$ infinitesimal is valid, but the reverse implication does not follow from normalization alone. A different event may be the unbounded one. An entry map that claims more must prove its extra source inequalities explicitly.

For a more precise cap screen with $d > 0$, (CAP7) yields

$$b > 0 \quad \Longrightarrow \quad \Delta < 0 \iff A < \frac{d^2(d + 2C + 2D)}{2b}. \tag{CAP19}$$

If $b \leq 0$, the strict inequality $\Delta < 0$ is automatic. Every inverse in (CAP19) is within the source field, and the polynomial condition (CAP14) remains the primary contact condition.

Scope of the new routing result. The three-event all-tests certificate and the finite-contact exclusion are proved here. In particular no finite-contact C1 or C2 elementary source can be silently discarded because it is said to pass a three-event wall: that intersection is empty. The bare finite-contact source domain already implies all eight strict wall inequalities, so adding these particular prior-complement predicates does not shrink that domain. Equations (CAP16)–(CAP19) justify only the stated scale partition and coefficient screens. They do not prove the missing positive-scale terminals, the uniform C1 all-tests join, the remaining weak C1/C2 local certificates, or the universal tensor-SOS criterion.

Chapter 199

The uniform C1 source for every positive e

Component `pf5d2f33f694c`
Source: `NEW/global/uniform_all_e_source_join_REDERIVED.tex`
SHA256: `f5d2f33f694ca549e0e75cf1314f052ead33cf714c0fc125177f768a4ba5ab77`
Inventory use: UE1-UE4 exhaustive e split and exact source hypotheses for all outer testing charts

199.1 The uniform C1 source for every positive e

Use the original canonical C1 source and the exact native transport EN13, with every original coefficient sign and strict gap retained. Suppose

$$k, f, g > 0 \text{ are infinitesimal}, \qquad e = kf/g > 0. \tag{UE1}$$

There is no additional restriction on the size of e, either source ratio $g/k, g/f$, or either separately labelled testing field. Then the entire original parent is tensor SOS at every independent projective testing pair and every real or complex testing state.

Here are the actual proof inputs and the complete join. EN13 gives $0 < a, t < 2$, $r = 1 - af^2 \sim 1$, and the original strict $M = R - t > 0, J = T - a > 0$. Split the positive pure-source e into

$$e \text{ infinitesimal} \quad \text{or} \quad \Delta = 1/e \text{ bounded}. \tag{UE2}$$

In the first case UJ1–UJ5 is the fresh full-parent theorem. It includes both source ratios g/k bounded and unbounded, both state charts, both complete state normals and all angular/projective joins; the actual bounded-ratio product-repair, fresh EZ and fresh IW proof bodies are its inputs.

In the second case use the actual retained `bounded_delta_body.tex`, Theorem *Uniform C1, bounded reciprocal e*. Its hypothesis is exactly UE1 with Δ bounded, and its proof is complete in that body. For clarity, the scope extension used there is substantive. After its pure-source substitution

$$\delta = \Delta R/(1 + k^2), \quad b^2 = f^2(1 + k^2)/R,$$

both δ and the pure-testing rotated state coordinate w are allowed to be bounded units. In its polynomial ring the weights

$$(\delta, k, b, a, t, u, w, v) = (0, 2, 6, 0, 0, 2, 0, 16) \tag{UE3}$$

make the full scalar error and both complete first normals have weights at least $42, 32, 26$. The displayed unlocalized monic conductor, restricted to $(A_m, B_m, C_m)^2 + (D_0^2)$, gives a scalar Gram whose every correction coefficient has weight at least 2; the first-normal coefficient bounds are $12, 6$. The exact hard block on its two normalized state normals has residue $16I_2$, including every quadratic term. Its explicit eleven-soft-root identity supplies a strict positive diagonal. Thus the entire parent,

with all four retained packets restored, has a bounded strict-residue Gram. This is a finite polynomial division proof; neither δ^{-1} nor a mixed element is a lifting coefficient.

The same body proves the complementary $W = 1, Z$ infinitesimal chart with five exact current roots. Its two full correction coefficients contain respectively ΔZ and $\Delta^2 f^2 Z^2$, so vanish in residue even when Δ is a unit. The original source signs and both strict gap openings remain intact.

For the outer angle joins, no conclusion from the old infinitesimal-e assertion is imported. Use the exact whole-parent identities UJ3–UJ4. Their source hypotheses hold on all of UE1. Indeed the seven native coquotients have residues $2, 2, 2, 2, 2, 2, 4$, regardless of e. Writing $B_c = 2c_5 - Jf^2 c_d \sim 4$, the original weights after normalizing $d = 1$ satisfy

$$C^{-1}, D^{-1} \text{ infinitesimal}, \qquad A\frac{CD}{C+D} = \frac{c_4 c_3 B_c}{c_d^2(aB_c + 2Jc_3)} \sim \frac{1}{4}. \tag{UE4}$$

Here $a + J = T \sim 1$, so the denominator has residue 64 without an individual lower bound on either opening. Those identities therefore cover every testing chart outside u, v infinitesimal, including literal projective infinities. Literal axes, the zero state, signs and complex states are covered by their exact full-parent normalization identities.

The two alternatives UE2 exhaust every positive source e. All scalings are within one labelled field, and every subtracted packet is restored. This proves UE1 at all tests, with literal $\beta = 0$ and every allowed original coefficient boundary. The two real closed extensions are arbitrarily chosen labelled factors; no algebraic-disjointness assumption is imposed.

Verification boundary. The complete retained bounded-Δ proof was read anew. An isolated copy of its standard-library reconstruction checker passed all 33 assertions: complete scalar and both complete first-normal lifts multiply back, and the actual minimum correction weights are $12, 14, 12$. This is corroboration of that actual polynomial proof, not a new 33-identity theorem or a premise derived from a historical status label. The broad historical infinitesimal-e assertion is replaced by the fresh UJ proof; its outer algebra is printed explicitly there and is used here only under UE4. No global entry of an arbitrary C1 source into UE1 is asserted in this section.

Chapter 200

Weak-alpha source entry and the unit-n terminal

Component `p3a3949869b9e`
Source: `NEW/global/weak_alpha_source_and_unit_n_REDERIVED.tex`
SHA256: `3a3949869b9e658c08d8de681cc2d74a105dc146f4db0ae07a00ec905aef7e72`
Inventory use: WA1-WA6 general weak-alpha entry and WA1-WA27 complete unit-n weak-m C2 terminal | WA1-WA27 common canonical weak-alpha source entry plus complete unit-n weak-m C2 all-tests theorem | WA1-WA10 only on weak-n source domain; exact full-parent/scalar/contact/normal identities WA12-WA20

200.1 Weak-alpha source entry and the unit-n terminal

Scope. All coefficients are bounded in the source field, $d = 1$, and the original canonical source is physical with an actual finite noncommuting zero contact. The C2 convention is $\lambda = \beta = 0$, and the C1 convention is $\lambda = n = 0$. The first subsection applies to either family with α infinitesimal. The all-tests theorem proved below has the additional C2 hypotheses that n is a positive unit and m is infinitesimal. It includes all independent testing states, signs and literal projective endpoints. The weak-n and C1 terminals are separate.

200.1.1 Source entry, proved before any shape quotient is used

Physicality forces C, A, D to be positive units. Here are exact residue-face tests proving this assertion. In C2 the negative central coefficient at $A = 0$, and its vanishing positive Laurent terms along $U = t^2, V = 1/t$, are the displayed forbidden-face proof in the actual positive-d source partition. If also $C = 0$ in residue, the source has only the four alternating events n, m, A, D. At $u = v = 1$ the full parent at state $(1, -\varepsilon, \varepsilon, 1)$ is exactly

$$16\varepsilon\{(1 + n + m + A + D)\varepsilon - 1\} < 0 \tag{WA1}$$

for a sufficiently small positive real ε. Thus C cannot be weak. The remaining central coefficient identities are

$$\begin{aligned} h|_{\alpha=D=0} &= (2A + 2C + 1)\{8\beta mn - 2\beta - 4mn - 2m - 2n - 1\}, \\ h|_{\alpha=C=n=0} &= -(2A + 1)(2D + 1)(2\beta + 2m + 1), \\ h|_{\alpha=A=n=0} &= -(2C + 2D + 1)(2\beta + 2m + 1). \end{aligned} \tag{WA2}$$

In the first line $n\beta = 0$, so it is strictly negative. For C2 with $\alpha = D = 0$, the other Laurent terms are $q_{20}U/V + q_{21}U$, killed by $U = t^2, V = t$. For C1 with $\alpha = D = 0$, they are $q_{12}V + q_{21}U + q_{22}UV$, killed by $U = V = t$. For C1 with $\alpha = C = 0$, they are $q_{02}V/U + q_{12}V$, killed by $U = 1/t, V = t$. For C1 with $\alpha = A = 0$, they are $q_{02}V/U + q_{12}V + q_{22}UV$, killed by $U = t, V = t^2$. Here t tends to zero through positive reals. These contradictions prove all the asserted unit bounds without a source-contact assumption.

Write $U = u_c^2, V = v_c^2 > 0$. The actual determinant dictionary now has

$$q_{01} = \alpha m a_1, \quad q_{02} = m a_2, \quad q_{10} = \alpha a_3, \quad q_{12} = a_4, \quad q_{20} = n a_5, \quad q_{21} = a_6, \quad q_{22} = \beta a_7, \tag{WA3}$$

where the a_i are positive bounded units. The two angular equations and their sum are

$$\begin{aligned} U^2(n a_5 + a_6 V + \beta a_7 V^2) &= mV(\alpha a_1 + a_2 V), \\ V^2(m a_2 + a_4 U + \beta a_7 U^2) &= \alpha a_3 U + n a_5 U^2, \\ a_6 U + a_4 V + 2\beta a_7 UV &= \alpha(m a_1/U + a_3/V). \end{aligned} \tag{WA4}$$

Every t_i in the dictionary is a positive unit because it includes $d = 1$. At contact all noncentral Laurent terms are nonnegative, and their sum is bounded. In particular $U, V, mV/U, nU/V$ are bounded. Multiplying the last equation by V shows $\alpha/V = O(U + V)$. The first two equations then exclude either U or V having nonzero residue: if U is a unit the first forces V a unit, and conversely the second forces U a unit; the last equation then contradicts weak α. Both are infinitesimal. Consequently

$$\alpha/V = o(1), \qquad m \asymp (U/V)^2(n + V), \qquad \alpha \asymp V(U + V). \tag{WA5}$$

For the last comparison, the denominator $a_3 + m a_1 V/U$ is a positive bounded unit. Actual finite contact forces $m, \alpha > 0$ in WA4. If n is a unit and m weak,

$$U/V \asymp \sqrt{m} = o(1). \tag{WA6}$$

If n is weak, then, whether U/V is bounded or unbounded, WA5 gives $mV/U, nU/V = o(1)$. These same two conclusions hold under WA6. They are the common source bounds used in the shape argument.

Orient the actual signs positively, use $W_c = 1$, and put

$$p = v_c, \quad q = Z_c, \quad P = p^2, \quad Q = q^2, \qquad u_c = rpq, \quad X_c = xpq^2, \quad Y_c = ypq. \tag{WA7}$$

The exact core sign identity SE9 ensures $W_c Z_c > 0$, so this normalization is legitimate. To prove boundedness before using these quotients, eliminate the X, Y block inside the source field. Set

$$H_X = n + m + \beta + \tfrac{1}{2}, \quad K = C + D + \tfrac{1}{2}, \quad H = K + A.$$

The block has diagonal positive residue H_X, H and an infinitesimal off-diagonal entry. The source Schur entries satisfy

$$S_{WW}/U \sim a = AK/H > 0, \qquad -S_{WZ}/(u_c v_c) \sim b = \tfrac{1}{2} + AD/H > 0. \tag{WA8}$$

Indeed the only additional leading contributions are multiples of $mV/U, nU/V, mn$, all infinitesimal on the domains just stated. The kernel equation gives $r \sim b/a$, a positive unit. The scaled X, Y equations are then a bounded system with positive triangular residue; its apparent m/Q term is bounded since $m/Q = r^2 mV/U$. Thus x, y are bounded before taking residues. Those equations and the two remaining kernel equations give

$$Ay \sim \tfrac{1}{2}, \quad K(r - y) \sim D + \tfrac{1}{2}, \quad D(1 + y - r) \sim r/2, \qquad x \sim -\frac{m + \beta + 1/2}{H_X}\, r. \tag{WA9}$$

For the last kernel division the possible extra α/V, nU/V and mV/U terms vanish by WA5. In particular $y, r - y$ and $1 + y - r$ are positive units, and the critical residue dictionary is

$$A_0 = \frac{1}{2y_0}, \qquad C_0 = \frac{1 + y_0}{2(r_0 - y_0)}, \qquad D_0 = \frac{r_0}{2(1 + y_0 - r_0)}. \tag{WA10}$$

In the unit-n, weak-m domain, $\beta = 0$, and hence

$$P, Q = o(1), \qquad m \asymp Q^2, \qquad \alpha \asymp P^2, \qquad x_0 = -r_0/(1 + 2n_0). \tag{WA11}$$

No independent testing coordinate occurred in this entry proof.

200.1.2 The exact scalar and every first normal

For reference the full finite quarter-parent is

$$\begin{aligned} F &= nD_uD_vX^2 + m(vW - X + uvY - uZ)^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ &\quad + (D + \tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ &\quad + (\beta + \tfrac{1}{2})D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \end{aligned} \tag{WA12}$$

Here $D_u = 1 + u^2, D_v = 1 + v^2$. We retain β symbolically to make the exact common graph explicit, but the current terminal has $\beta = 0$. In the testing chart $W = 1, Z = z$, use

$$u_g = rvz, \qquad X_g = xvz^2, \qquad Y_g = yvz, \qquad V_t = v^2, \quad T = z^2. \tag{WA13}$$

In the rest of this subsection write $V = V_t$ only. The full graph restriction is

$$\begin{aligned} \Phi &= mV + \alpha T + hVT + \mathcal{C}V^2T + EVT^2 + \mathcal{F}V^2T^2 + GV^3T^2 + JV^2T^3 + \mathcal{K}V^3T^3, \\ \mathcal{C} &= Ay^2 + C(r - y)^2 + (\alpha + \tfrac{1}{2})y^2 + 2mry, \\ E &= \alpha r^2 + (m + \beta + \tfrac{1}{2})(r + x)^2 + nx^2, \\ G &= r^2y^2(A + \alpha + m + \tfrac{1}{2}), \qquad J = nr^2x^2, \qquad \mathcal{K} = (D + n + \tfrac{1}{2})r^2x^2, \\ \mathcal{F} &= Ar^2y^2 + (2D + 1)rx(r - y - 1) - (2\alpha + 1)r^2y \\ &\quad - 2mry(r + x) + \beta(r + x)^2 + nx^2 + rx + x^2/2. \end{aligned} \tag{WA14}$$

The coefficient h is defined by this entire expansion; it is bounded. Every displayed coefficient is bounded. In the current unit-n domain $\mathcal{C}, E$ are positive units, since

$$\mathcal{C}_0 = r_0(1 + y_0)/2, \qquad E_0 = n_0r_0^2/(1 + 2n_0). \tag{WA15}$$

The coefficient $\mathcal{F}$ need not be positive.

The actual contact gives $\Phi(P, Q) = \Phi_V(P, Q) = \Phi_T(P, Q) = 0$. Solving these three linear equations in m, α, h is exact:

$$\begin{aligned} m &= EQ^2 + \mathcal{F}PQ^2 + GP^2Q^2 + 2JPQ^3 + 2\mathcal{K}P^2Q^3, \\ \alpha &= \mathcal{C}P^2 + \mathcal{F}P^2Q + 2GP^3Q + JP^2Q^2 + 2\mathcal{K}P^3Q^2, \\ h &= -2\mathcal{C}P - 2EQ - 3\mathcal{F}PQ - 4GP^2Q - 4JPQ^2 - 5\mathcal{K}P^2Q^2. \end{aligned} \tag{WA16}$$

Put

$$R = z(V - P), \quad S = v(T - Q), \quad \Psi = QR^2 + PS^2 + vzRS.$$

With the exact balanced monomials

$$\begin{aligned} B_{32} &= 2PQR^2 + (vzR + PS)^2, \\ B_{23} &= 2PQS^2 + (QR + vzS)^2, \\ B_{33} &= (2PQ^2 + VT^2)R^2 + (2P^2Q + PVT)S^2 + (3PQ + PT)vzRS, \end{aligned} \tag{WA17}$$

the whole scalar, not a leading truncation, is

$$\Phi = \mathcal{C}R^2 + ES^2 + \mathcal{F}\Psi + GB_{32} + JB_{23} + \mathcal{K}B_{33}. \tag{WA18}$$

For v, z infinitesimal, all entries after the first two positive unit pivots are infinitesimal, by $P, Q = o(1)$. This is a strict two-row Gram even when $\mathcal{F}$ is signed.

Let L_u, L_X, L_Y be half the full first derivatives of WA12 at WA13. The complete polynomials have the forms

$$\begin{aligned} L_u &= vzA_u(V, T), \qquad L_Y = vzA_y(V, T), \\ A_u &= a_{00} + a_{10}V + a_{01}T + a_{11}VT + a_{12}VT^2 + a_{21}V^2T + a_{22}V^2T^2, \\ A_y &= e_{00} + e_{10}V + e_{11}VT + e_{21}V^2T, \\ L_X &= v[-m + TB_X(V, T)], \qquad B_X = c_0 + c_1V + c_2VT + c_3V^2T. \end{aligned} \tag{WA19}$$

All coefficients are bounded, and the expansion fixes every one unambiguously. In particular

$$c_2 = nr^2x, \qquad c_3 = (D + n + \tfrac{1}{2})r^2x.$$

Their actual source jets are $A_u(P,Q) = A_y(P,Q) = 0$ and $B_X(P,Q) = m/Q$. Finite monic difference division gives

$$\begin{aligned} L_i &= vA_{i,V}(V,T)R + zB_i(T)S, \qquad i = u, Y, \\ L_X &= vzB_{X,V}(V,T)R + \{T(Pc_2 + P^2c_3) + m/Q\}S. \end{aligned} \tag{WA20}$$

Here $A_{i,V} = [A_i(V,T) - A_i(P,T)]/(V-P)$, and $B_{X,V}$ is defined identically. The other divided polynomials are

$$\begin{aligned} B_u(T) &= a_{01} + Pa_{11} + Pa_{12}(T+Q) + P^2a_{21} + P^2a_{22}(T+Q), \\ B_Y(T) &= Pe_{11} + P^2e_{21}. \end{aligned}$$

These are literal polynomial identities with bounded coefficients, including $v = 0$ or $z = 0$. Since $m/Q = O(Q) = o(1)$, every full first normal is an infinitesimal linear combination of R, S.

Set $N_u = u - rvz, N_X = X - xvz^2, N_Y = Y - yvz$. In the chart under consideration these are infinitesimal. The full higher-normal remainder is a polynomial in the three normals with no terms of normal degree below two; all terms of degree at least three give infinitesimal Gram entries. Its quadratic residue is

$$K_0(N_u - N_Y)^2 + A_0N_Y^2 + (n_0 + \tfrac{1}{2})N_X^2 \succ 0. \tag{WA21}$$

WA18 and WA20 therefore give a strict bounded Gram for the entire parent on the three hard and two soft rows. The strict-Gram lemma proves its tensor-SOS assertion, with no mixed inverse.

200.1.3 The complementary local states

Normalize a maximal testing state coordinate. At $u, v = o(1)$ the state residue has positive pivots on X, Y; if one is a unit, the full parent has positive residue immediately. Otherwise one of W, Z is a unit. The preceding proof covers $W = 1, Z = o(1)$. Every remaining state can be normalized to $Z = 1, W = \omega$ bounded. Retain the entire positive m, α packets. The remainder lies exactly in the square of (u, v, X, Y). Its quadratic real residue is

$$\begin{aligned} &(n_0 + \tfrac{1}{2})\left(X + \frac{u}{1+2n_0}\right)^2 + \frac{n_0}{1+2n_0}u^2 \\ &+ H_0\left(Y - \frac{K_0\omega u - D_0 v}{H_0}\right)^2 + a_0(\omega u - r_0 v)^2, \qquad a_0 = A_0K_0/H_0 > 0. \end{aligned} \tag{WA22}$$

The critical identities WA10 prove the equality. This quadratic form is positive definite for every real ω, including zero. All higher normal terms are infinitesimal corrections on these four rows, while $\omega - \bar{\omega}$ is retained in coefficients. Thus every state at the double-infinitesimal angle corner is covered.

200.1.4 All other testing angles, including the literal endpoints

Again retain the entire m, α packets and call the remainder F_0. Its source residue is the critical C, A, D, n source. For bounded u, v, F_0 is exactly a bounded quadratic form in the four rows (uW, X, Y, Z). Its real $Z = 0$ block is positive: the nX^2 and AY^2 packets and $C(uW - Y)^2$ suffice. The determinant is

$$\det \mathrm{Mat}_{(uW,X,Y,Z)}(\bar{F}_0) = D_u^2D_v^2\{q_{20}u^2 + q_{12}v^4 + q_{21}u^2v^2\}/16 > 0 \tag{WA23}$$

whenever at least one of the two real angle residues is nonzero. Here $q_{20}, q_{12}, q_{21} > 0$ are the actual residue determinant coefficients, and $h = 0$ follows from WA10. This proves strictness of the full four-row Gram. Literal $u = 0$ is allowed.

For Bob reciprocal $b = 1/v$ infinitesimal and Alice bounded, use the same rows (uW, X, Y, Z) in $b^2 F_0(u, 1/b)$. At $b = 0$ its whole real expression is

$$C_0(uW - Y)^2 + (A_0 + \tfrac{1}{2})D_u Y^2 + D_0(uX - Z)^2 + (n_0 + \tfrac{1}{2})D_u X^2 \succ 0. \tag{WA24}$$

It is strict at every real u, including zero. For Alice reciprocal $s = 1/u$ infinitesimal and Bob bounded, the original state rows suffice. The expression at $s = 0$ is

$$n_0 D_v X^2 + C_0 D_v W^2 + A_0 D_v Y^2 + (D_0 + \tfrac{1}{2})(W + vX)^2 + \tfrac{1}{2}(vY - Z)^2 \succ 0. \tag{WA25}$$

This includes $v = 0$, using the positive n_0 pivot.

For both reciprocal angles $s, b = o(1)$, use the five rows (W, X, Y, sZ, bZ). The whole reciprocal polynomial belongs to their square ideal, and its real coefficient Gram is

$$C_0 W^2 + A_0 Y^2 + D_0(X - sZ)^2 + (n_0 + \tfrac{1}{2})X^2 + \tfrac{1}{2}(Y - bZ)^2 \succ 0. \tag{WA26}$$

Every remaining coefficient is infinitesimal. These exact finite square-ideal allocations include $s = 0$, $b = 0$ and arbitrary states; no projective endpoint is obtained by a limiting argument.

The four angle charts and the two local state charts are exhaustive. Testing sign gauges, positive pure-field homogeneous normalization factors and real/imaginary state splitting restore the original parent:

$$F_c(u, v; \psi) \ \in \ \Sigma(K_c \otimes_{\mathbb{R}} K_t)^2 \quad \text{for every independent projective test and state.} \tag{WA27}$$

This closes exactly the C2 unit-n, weak-m, α terminal. The common source-entry subsection does not by itself close the weak-n or C1 weak-α branches.

Evidence boundary. The focused checker expands the full parent, the scalar and all first normals, checks the balanced identities with their explicit contact remainders, and verifies the complete hard and outer residue forms. The positive-unit source estimates, the finite quantified chart cover and the application of the separated strict-Gram lemma are the analytical proof above. No unavailable historical local proof is used.

Chapter 201

Weak-alpha weak-n and C1: the complete parent

Component `p0db05df21ad1`
Source: `NEW/global/weak_alpha_weak_n_and_c1_REDERIVED.tex`
SHA256: `0db05df21ad1a9a61bbd5eb6ee8179ef56de0076e8d7745bb14af5e3fc55a5b1`
Inventory use: WN1-WN44 complete weak-n C2 and arbitrary-beta C1 weak-alpha all-tests theorem | WN1-WN44 complete bounded positive-d weak-alpha weak-n C2 and arbitrary-beta C1 all-tests theorem

201.1 Weak-alpha weak-n and C1: the complete parent

Scope and retained inputs. The source is an actual physical canonical finite noncommuting contact, all coefficients are bounded in the source field, and $d = 1$. Either it is C2 with $\lambda = \beta = 0$ and n, α infinitesimal, or it is C1 with $\lambda = n = 0$, α infinitesimal and arbitrary bounded $\beta \geq 0$. All strict source gaps retain their original meaning. We prove the tensor-SOS statement for the entire parent at every independent projective testing angle and every state. No prior-complement exclusion is assumed. The common source-entry proof WA1–WA10 is used on precisely its weak-n domain. In particular $C, A, D, r, y, r - y, 1 + y - r$ are positive units, x is bounded, and $x_0 = -r_0$. Boundedness, infinitesimality and residue refer to the separated bounded tensor ring and its real residue map. The strict-Gram lemma is applied only after a finite explicit row allocation; no mixed tensor is inverted.

Use the full quarter-parent WA12 and its exact source coordinates

$$p = v_c, \quad q = Z_c > 0, \quad P = p^2, \quad Q = q^2, \qquad W_c = 1, \quad u_c = rpq, \quad X_c = xpq^2, \quad Y_c = ypq. \tag{WN1}$$

Thus P, PQ are positive infinitesimals. Put

$$H_0 = m + \beta + \tfrac{1}{2}, \quad H_X = H_0 + n, \quad K = C + D + \tfrac{1}{2}, \quad H_Y = K + A. \tag{WN2}$$

These are positive units. The exact scalar coefficients $\mathcal{C}, E, \mathcal{F}, G, J, \mathcal{K}$, the contact balances, and the complete first normal polynomials are those displayed in WA14, WA16 and WA19; those identities retain β and n. Their relevant residues are

$$\mathcal{C}_0 = \frac{r_0(1 + y_0 + 4m_0y_0)}{2} > 0, \qquad \mathcal{F}_0 = \frac{r_0^2(1 + y_0)}{2} > 0, \qquad G, J, \mathcal{K} \geq 0. \tag{WN3}$$

Both $\mathcal{C}, \mathcal{F}$ are consequently positive units. The exact source X-jet gives, with the WA19 coefficients,

$$\begin{gathered} c_0 = H_0(r + x) + nx, \quad c_2 = nr^2x, \quad c_3 = (D + n + \tfrac{1}{2})r^2x, \\ c_0 = m/Q - Pc_1 - PQc_2 - P^2Qc_3, \\ E = \alpha r^2 + c_0^2/H_X + nH_0r^2/H_X. \end{gathered} \tag{WN4}$$

The last equality is an exact completion inside the source field.

201.1.1 Two finite positive circuit identities

For $V = v^2, T = z^2$ put

$$R = z(V - P), \quad S = v(T - Q), \quad \Psi = V^2T^2 + PQ^2V + P^2QT - 3PQVT.$$

Then $\Psi = QR^2 + PS^2 + vzRS$. It is an explicit six-square cubic circuit: if $L_1^3 = V^2T^2, L_2^3 = PQ^2V, L_3^3 = P^2QT$, then $L_1L_2L_3 = PQVT$, and

$$\Psi = \sum_{i=1}^{3} L_i\{(L_1 - L_2)^2 - (L_1 - L_2)(L_1 - L_3) + (L_1 - L_3)^2\}. \tag{WN5}$$

The real 2-by-2 Gram in each brace is strictly positive. All radicals in this formula are elementary separated monomials: source radicals and testing radicals are taken in their own real closed fields and then tensored.

The balanced degree-$(3, 3)$ term has the polynomial identity

$$a^{10} + 2b^{10} + 2c^{10} - 5a^2b^4c^4 = 2(b^5 - c^5)^2 + (a^2 - bc)^2(a^6 + 2a^4bc + 3a^2b^2c^2 + 4b^3c^3). \tag{WN6}$$

For nonnegative elementary monomials a, b, c, the bracket is the sum of squares of the four rows

$$a^3, \qquad \sqrt{2}a^2\sqrt{bc}, \qquad \sqrt{3}abc, \qquad 2(bc)^{3/2}.$$

This is a five-square separated certificate, including vanishing variables. Applied with

$$a^{10} = V^3T^3, \quad b^{10} = P^2Q^3V, \quad c^{10} = P^3Q^2T,$$

it proves positivity of the entire B_{33} in WA17, rather than discarding a signed high-order remainder.

A further exact identity supplies any fixed positive real pivot on the row PS:

$$2P\Psi + VTR^2 - (PS)^2 = (PS + vzR)^2 + 2PQR^2. \tag{WN7}$$

Indeed, for every fixed $b > 0$,

$$\begin{aligned} \mathcal{C}R^2 + \mathcal{F}\Psi = (\mathcal{C} - bVT + 2bPQ)R^2 + (\mathcal{F} - 2bP)\Psi \\ + b(PS)^2 + b(PS + vzR)^2. \end{aligned} \tag{WN8}$$

On the charts where this is used, VT, P, PQ are infinitesimal, so the original two unit residues are unchanged.

201.1.2 Bounded source ratio Q

From WA5, WN4 and bounded Q,

$$m \asymp Q^2(n + P), \quad \alpha \asymp P^2, \quad r + x = O(n + P), \quad E \asymp n + P^2. \tag{WN9}$$

In particular m, n, E, α are infinitesimal. None of these estimates contains an independent testing coordinate.

First take $W = 1, Z = z$ bounded and u, v infinitesimal. Use the entire graph $u = rvz, X = xvz^2, Y = yvz$. The exact scalar is

$$\Phi = \mathcal{C}R^2 + ES^2 + \mathcal{F}\Psi + GB_{32} + JB_{23} + \mathcal{K}B_{33}, \tag{WN10}$$

where the last three terms retain their positive full certificates. The strict soft rows are R, the six cubic rows in WN5, $\sqrt{E}\,S$, and PS; WN8 supplies any needed real pivot on the last row.

The exact monic normal divisions WA20 apply unchanged. Their R-coefficients contain v or vz, hence are infinitesimal. Their S-coefficients are $O(E + P)$: in particular

$$a_{01} = \alpha r + H_0(r + x) = O(E + P), \qquad a_{12} = nrx^2, \qquad m/Q = O(E + P). \tag{WN11}$$

Every ES contribution has infinitesimal coefficient $\sqrt{E}$ on $\sqrt{E}S$. Coefficients on PS may be bounded units. There are only finitely many of them, so choose b in WN8 larger than their finite real Schur cost. This uses a real constant, not an inverse of a mixed tensor.

The three hard normals $N_u = u - rvz, N_X = X - xvz^2, N_Y = Y - yvz$ are infinitesimal. Their complete quadratic residue is

$$K_0(N_u - N_Y)^2 + A_0 N_Y^2 + (\beta_0 + \tfrac{1}{2})(N_X + \bar{z}N_u)^2 \succ 0. \tag{WN12}$$

All higher-normal terms have degree at least three and bounded coefficients, hence allocate as infinitesimal Gram entries on these hard rows. WN10–WN12 prove the full parent in this chart.

For the complementary local states take $Z = 1, W = w$ infinitesimal, set $a = u/r$, and use

$$v = aw, \quad X = xa, \quad Y = yaw, \qquad U = a^2, \quad T = w^2. \tag{WN13}$$

The entire scalar is

$$\Phi' = \alpha + mUT^2 + EU + hUT + \mathcal{C}U^2T^2 + \mathcal{F}U^2T + GU^3T^2 + JU^2 + \mathcal{K}U^3T. \tag{WN14}$$

Here $a = (1/r) \otimes u$ is elementary; its roots remain separated. The strict rows are $P, a^2w, \sqrt{E}\,a$. The term hUT is harmless: the contact balances give $h = O(P + E)$, and writing $h = Ph_P + Eh_E$ with bounded source coefficients gives the infinitesimal Gram terms $h_P w\, P(a^2w) + h_E w^2(\sqrt{E}a)^2$. All other terms of WN14 are positive.

For completeness, half the entire v, Y, X derivatives have the polynomial forms

$$\begin{aligned} L'_v &= aw(d_0 + mT + d_1U + d_2UT + d_3U^2 + d_4U^2T), \\ L'_Y &= aw(e_0 + e_1U + e_2UT + e_3U^2T), \\ L'_X &= a(c_0 - mT + c_2U + c_1UT + c_3U^2T). \end{aligned} \tag{WN15}$$

Every coefficient is bounded. Source jets at $U = PQ, T = 1/Q$ give

$$\begin{aligned} d_0 &= -m/Q - d_1PQ - d_2P - d_3P^2Q^2 - d_4P^2Q = O(n + P), \\ e_0 &= -e_1PQ - e_2P - e_3P^2Q = O(P). \end{aligned}$$

Also $c_0, m = O(n + P)$ and $c_2 = O(n)$. Splitting $n + P = O(E + P)$ in these displayed complete polynomials allocates all first normals with infinitesimal coefficients to $P, a^2w, \sqrt{E}a$. For example awP, awE, and a^3w have coefficients aw, $w\sqrt{E}$, and a on those three rows; the other monomials have still more small testing factors. The complete hard residue is

$$(\beta_0 + \tfrac{1}{2})N_X^2 + D_0N_v^2 + 2D_0N_vN_Y + H_{Y,0}N_Y^2 \succ 0, \tag{WN16}$$

where $N_v = v - aw, N_Y = Y - yaw, N_X = X - xa$. The higher-normal argument is again exact polynomial allocation. This proves all local states when Q is bounded.

201.1.3 Unbounded source ratio: the W-chart

Suppose Q is unbounded. The source estimates and positive contact balances give

$$PQ = o(1), \quad m \asymp Q^2(n + P), \quad nQ^2, PQ^2 = O(1), \quad \alpha \asymp P^2Q. \tag{WN17}$$

WN4 now gives

$$c_0 \sim m/Q, \quad r + x \sim m/(H_XQ), \quad E \asymp n + (m/Q)^2, \quad h/m = o(1). \tag{WN18}$$

To justify the first comparison before its use, divide the three errors Pc_1, PQc_2, P^2Qc_3 by m/Q. Their absolute values are $O(1/Q), O(P), O(P)$, respectively; $n/(m/Q) = O(1/Q)$. The positive term

$\alpha/(m/Q)^2 = O(1/Q)$ follows from $m \ge cPQ^2$. The last comparison follows term by term from WA16 using these same inequalities.

Take $W = 1, Z = z$ bounded, u, v infinitesimal. If m is a unit, retain the entire n, α packets. The remainder lies in the square of (u, v, X, Y). Its real quadratic form is

$$\begin{aligned} &H_0\{X + zu - (m/H_0)v\}^2 + \frac{m(\beta + 1/2)}{H_0}v^2 \\ &\quad + H_Y\{Y - (Ku - Dzv)/H_Y\}^2 + a_*(u - rzv)^2, \qquad a_* = AK/H_Y > 0, \end{aligned} \tag{WN19}$$

with coefficients interpreted in residue. WA10 verifies this identity. It is strict for every bounded z, so the complete higher terms give a strict Gram.

If m is weak, use the graph WA13. Its entire scalar has strict rows $\sqrt{m}\,v, v^2 z, \sqrt{\alpha}\,z$, since $mV + hVT = mV[1 + (h/m)T]$, while $\mathcal{C}, \mathcal{F}, E, G, J, \mathcal{K}$ are nonnegative. The complete source jet equations yield

$$a_{00} = O(m), \quad e_{00} = O(PQ), \quad a_{01}, c_0 = O(m/Q). \tag{WN20}$$

Thus $a_{00}/\sqrt{m}$, $e_{00}/\sqrt{m}$, $a_{01}/\sqrt{m}$, $c_0/\sqrt{m}$ are infinitesimal. For e_{00}, use $PQ/\sqrt{m} = O(\sqrt{P})$. Every other monomial of the complete normals WA19 has a factor $v^3 z$ or $v^3 z^2$, so it has an infinitesimal coefficient on $v^2 z$. The hard residue is WN12; all higher terms are allocated exactly as there. This proves the whole W-chart.

201.1.4 Unbounded source ratio: the complementary scalar

Use WN13 with a, w infinitesimal. Introduce only source quantities

$$\begin{gathered} \epsilon = Q^{-1} = o(1), \quad \Lambda = PQ = o(1), \quad \tau = \Lambda/\epsilon = PQ^2 = O(1), \\ \nu = n/\epsilon^2 = O(1), \qquad N = E/\epsilon^2 = O(1). \end{gathered} \tag{WN21}$$

The quantities $\epsilon, \Lambda, \tau, N$ are positive; $\nu \ge 0$. Let

$$A_1 = a(T - \epsilon), \quad K_1 = UT - \Lambda\epsilon, \quad \mathcal{M} = U^2T + \tau UT^2 + \Lambda^2\epsilon - 3\Lambda UT. \tag{WN22}$$

The whole scalar WN14, using exactly the three contact balances, is

$$\begin{aligned} \Phi' &= \mathcal{C}K_1^2 + NA_1^2 + \mathcal{F}\mathcal{M} \\ &\quad + G\{2\Lambda K_1^2 + UT^2(U - \Lambda)^2\} \\ &\quad + J\{(U - \Lambda)^2 + 2\Lambda A_1^2/\epsilon^2\} + \mathcal{K}\mathcal{B}_{33}, \\ \mathcal{B}_{33} &= U^3T + 2(\Lambda^2/\epsilon)UT^2 + 2\Lambda^3\epsilon - 5\Lambda^2UT. \end{aligned} \tag{WN23}$$

The apparent $1/\epsilon^2$ in the positive J packet is a source-field quantity, and $J/\epsilon^2 = \nu r^2 x^2$ is bounded. All positive terms are retained. WN6 applies to $\mathcal{B}_{33}$ with $a^{10} = U^3T, b^{10} = (\Lambda^2/\epsilon)UT^2, c^{10} = \Lambda^3\epsilon$. Here $\Lambda^2/\epsilon = \tau\Lambda$ is bounded.

Write $a = t^3, w = s^3, \epsilon = h^6, \tau = c^6$, taking positive elementary roots. Then t, s, h are infinitesimal and c is bounded; it is allowed to be a unit. Put

$$\begin{gathered} L_1 = t^4s^2, \quad L_2 = c^2t^2s^4, \quad L_3 = h^6c^4, \\ f_1 = t^2s, \quad f_2 = cts^2, \quad f_3 = h^3c^2, \\ R_{2i-1} = f_i(L_1 - L_2), \qquad R_{2i} = f_i(L_1 - L_3) \quad (i = 1, 2, 3). \end{gathered} \tag{WN24}$$

Then $\mathcal{M} = \sum_{i=1}^{3}(R_{2i-1}^2 - R_{2i-1}R_{2i} + R_{2i}^2)$, a strict six-row constant Gram. The following are exact identities:

$$\begin{aligned} aw(U - \Lambda) &= t^3R_1 + c^2ts^2R_2, \\ a^2w(T - \epsilon) &= -h^6R_1 + s^6R_2 - h^3s^3R_5, \\ \tau aw(T - \epsilon) &= c^2ts^2(R_2 - R_1) - c^3s^3R_3, \\ \tau\sqrt{\epsilon}\,A_1 &= t^3R_6 - t(t^2 + c^2s^2)R_5. \end{aligned} \tag{WN25}$$

Every coefficient on the right is infinitesimal, even when c is a unit. These are finite linear conductors, not limiting approximations. The further quadratic conductor is

$$\boxed{\Lambda(U-\Lambda)K_1 = \Lambda\mathcal{M} - (\tau\sqrt{\epsilon}\,A_1)^2.} \tag{WN26}$$

Its right side is an infinitesimal Gram on the six cubic rows. It remains so after multiplication by any bounded signed coefficient.

If τ is a unit, then K_1 itself has an infinitesimal linear representation on those rows:

$$K_1 = \tau^{-1/3}\{\sqrt{L_1}(R_2 - R_1) + \sqrt{L_3}R_6\}. \tag{WN27}$$

This follows by expansion as $\tau^{-1/3}(L_1L_2 - L_3^2) = UT - \Lambda\epsilon$. Consequently any fixed real pivot on K_1 can be supplied at infinitesimal cost to the strict $\mathcal{M}$ Gram in this case.

201.1.5 Every complementary first normal

Let D_U, E_U denote the monic U-divided differences of the bracket polynomials in the first two lines of WN15. The source jets at $U = \Lambda, T = \epsilon$ give the exact identities

$$\begin{aligned} L'_v &= aw\{D_U(U,T)(U-\Lambda) + (m + \Lambda d_2 + \Lambda^2 d_4)(T-\epsilon)\},\\ L'_Y &= aw\{E_U(U,T)(U-\Lambda) + (\Lambda e_2 + \Lambda^2 e_3)(T-\epsilon)\},\\ L'_X &= -mA_1 + a(c_2 + c_3\Lambda\epsilon)(U-\Lambda) + a(c_1 + c_3U)K_1. \end{aligned} \tag{WN28}$$

There are no omitted monomials. Define

$$d_* = (c_2 + c_3\Lambda\epsilon)/\epsilon = \nu\epsilon r^2 x + \Lambda c_3.$$

Since $\epsilon(U-\Lambda) = K_1 - U(T-\epsilon)$, the last line is

$$L'_X = -mA_1 + P_X, \qquad P_X = a(c_1 + c_3U + d_*)K_1 - d_*UA_1. \tag{WN29}$$

The exact source equations also give

$$\begin{aligned} m &= N + \tau(\mathcal{F} + G\Lambda\epsilon + 2\mathcal{K}\Lambda) + 2\nu r^2x^2\Lambda,\\ c_0/\epsilon &= m - \Lambda B, \qquad B = c_1 + \Lambda c_3 + \nu\epsilon r^2 x,\\ N &= \alpha r^2/\epsilon^2 + (m-\Lambda B)^2/H_X + \nu H_0 r^2/H_X. \end{aligned} \tag{WN30}$$

In particular

$$N, \nu, \tau = O(m), \quad \Lambda = O(m\epsilon), \quad N = m^2/H_X + \nu H_0 r^2/H_X + O(m\Lambda). \tag{WN31}$$

For the error term use $\alpha/\epsilon^2 = O(\tau\Lambda)$, which is the exact positive alpha balance WN23/WA16. All these comparisons are inside the source field. We split by the bounded source ratio ν/m; $m > 0$.

Substantial ν/m. If ν/m has positive residue, keep $S_0 = \sqrt{m}A_1$. The N_X, S_0 Gram, including the entire $-mA_1$ first normal, has residue from

$$\begin{pmatrix} H_X & -\sqrt{m} \\ -\sqrt{m} & N/m \end{pmatrix}. \tag{WN32}$$

Its determinant is $(\nu/m)H_0r^2 + O(\Lambda)$, a positive unit. The v, Y first normals in WN28 use the first conductor WN25; their remaining coefficients on S_0 are $O(\sqrt{m}\,w)$, hence infinitesimal. In WN29 the K_1-coefficient contains a, and $d_*U/\sqrt{m} = O(\sqrt{m}\,\epsilon U)$ is infinitesimal. Thus all other first normals have infinitesimal coefficients on K_1, the six cubic rows, and S_0. The v, Y hard block in WN16 is strict. All higher terms have at least three infinitesimal hard normals and bounded coefficients. This proves the full parent on this source subdomain.

Thin ν/m, including literal $n=0$. Assume $\nu/m=o(1)$. Equations WN30–WN31 imply τ/m is a positive unit: after division by m,

$$1=N/m+(\tau/m)\mathcal{F}+o(1),\qquad N/m=m/H_X+o(1),$$

and $1-m/H_X=(n+\beta+1/2)/H_X$ has positive residue. Write

$$N-m^2/H_X=\nu\xi+\eta,\qquad \xi=H_0r^2/H_X>0,\qquad \eta=O(m\Lambda).$$

Complete only the source coefficient H_X:

$$\begin{aligned} H_XN_X^2-2mN_XA_1+NA_1^2&=H_X\widetilde{N}_X^2+\nu\xi A_1^2+\eta A_1^2,\\ \widetilde{N}_X&=N_X-(m/H_X)A_1. \end{aligned}\tag{WN33}$$

Retain the strict row $\sqrt{\nu}A_1$ when $\nu>0$; if $\nu=0$, all terms allocated to it vanish literally. The signed error is an infinitesimal cubic Gram, since

$$m\Lambda A_1^2=(m/\tau)(\tau\sqrt{\epsilon}\,A_1)^2.\tag{WN34}$$

The first two normals in WN28 use WN25 and bounded m/τ. For P_X, its K_1-coefficient contains a, while its two other contributions have infinitesimal allocations

$$\nu\epsilon UA_1=(\sqrt{\nu}\,\epsilon U)(\sqrt{\nu}A_1),\qquad \Lambda UA_1=(\sqrt{\epsilon}\,U)(\tau\sqrt{\epsilon}\,A_1).\tag{WN35}$$

Therefore $2P_X\widetilde{N}_X$ has infinitesimal Gram entries.

The shift in WN33 also creates the entire scalar term $2(m/H_X)A_1P_X$, which must not be dropped. Its main factor is controlled by the exact identity

$$mU(T-\epsilon)K_1=mK_1^2-(m/\tau)\Lambda(U-\Lambda)K_1.\tag{WN36}$$

The second term is infinitesimal by WN26. If m is infinitesimal, the first is an infinitesimal K_1-Gram. If m is a unit, then τ is a unit and WN27 makes it an infinitesimal cubic Gram. In that case a positive real K_1-pivot may be reinserted by WN27 at infinitesimal cubic cost. This allocation remains valid with each bounded factor $c_1+c_3U+d_*$. The remaining scalar factors are multiples of $m\nu\epsilon UA_1^2$ and $m\Lambda UA_1^2$; they have infinitesimal Gram entries on $\sqrt{\nu}A_1$ and the fourth conductor in WN25, respectively. This accounts for the whole scalar effect of the completion.

201.1.6 The entire higher-normal restoration in the thin case

The full parent is quadratic in X. Its coefficient beyond the source constant H_X is exactly the nonnegative polynomial

$$D_X=(n+\beta+\tfrac{1}{2})v^2+nu^2+(n+D+\tfrac{1}{2})u^2v^2.\tag{WN37}$$

Retain the entire packet $D_XN_X^2$, evaluated at the actual independent u,v, as separated squares. Each monomial coefficient is a nonnegative source scalar times a testing square. It remains a positive whole packet after WN33. The remaining polynomial has no quadratic N_X term except $H_XN_X^2$. At the complementary graph its half mixed derivatives have the exact forms

$$[N_XN_v]=w[-m+a^2B_v],\qquad [N_XN_Y]=a^2wB_Y,\tag{WN38}$$

where B_v,B_Y are bounded polynomials fixed by the full parent expansion. After the shift, the new first-normal terms are bounded multiples of $\tau aw(T-\epsilon)$, using $m/\tau=O(1)$; WN25 gives infinitesimal coefficients. Every other remaining monomial linear in N_X has degree at least two in the other hard normals. Replacing N_X by $\widetilde{N}_X+(m/H_X)A_1$ gives an infinitesimal coefficient on a product of at least two hard normals, since $A_1=o(1)$. Terms involving $\widetilde{N}_X$ have total hard-normal degree at least three and are handled in the same way. Terms independent of N_X are unchanged. The full real hard block is

$$H_{X,0}\widetilde{N}_X^2+D_0N_v^2+2D_0N_vN_Y+H_{Y,0}N_Y^2\succ 0.\tag{WN39}$$

Equations WN23–WN39 therefore give a finite strict Gram for the whole parent, including all first normals, every higher term, all β-openings and literal $n=0$. This completes the double-infinitesimal testing corner for both source ratios and all local states.

201.1.7 Every other angle, all states and literal endpoints

Retain the entire m, n, α, β packets and call the remainder F_0. Only its residue is the critical C, A, D source in WA10; no global physicality of the subtracted finite source is asserted. The following are complete polynomial row allocations, so their strict residue Grams apply to the original nearby coefficients.

For bounded u, v with v a unit use (uW, X, Y, Z). The $Z = 0$ principal block is positive definite and

$$\det \mathrm{Mat}_{(uW,X,Y,Z)} \bar{F}_0 = D_u^2 D_v^2 (q_{12} v^4 + q_{21} u^2 v^2)/16 > 0. \tag{WN40}$$

Both q_{12}, q_{21} are positive units. If u is a unit and v infinitesimal, use instead $(uW - Y, Y, X + uZ, vZ)$. The entire parent is polynomial in these rows with bounded coefficients; the only algebraic coefficient divisions are by the pure testing unit u. Its determinant at $v = 0$ is $D_u^2 q_{21} u^2/16 > 0$, and the first three-row principal block is strict. This covers all bounded angle pairs outside the double-infinitesimal corner.

For Bob reciprocal $b = 1/v = o(1)$, Alice bounded, the rows (uW, X, Y, Z) give the real form

$$C_0 (uW - Y)^2 + (A_0 + \tfrac{1}{2}) D_u Y^2 + D_0 (uX - Z)^2 + \tfrac{1}{2} D_u X^2 \succ 0. \tag{WN41}$$

For Alice reciprocal $s = 1/u = o(1)$ and Bob a bounded unit, the original state rows give

$$C_0 D_v W^2 + A_0 D_v Y^2 + (D_0 + \tfrac{1}{2})(W + vX)^2 + \tfrac{1}{2}(vY - Z)^2 \succ 0.$$

If s, v are both infinitesimal, use (W, Y, vX, Γ), where $\Gamma = sX + Z$. The entire reciprocal parent is polynomial in these four un-divided rows, and its real form is

$$C_0 W^2 + A_0 Y^2 + (D_0 + \tfrac{1}{2})(W + vX)^2 + \tfrac{1}{2}\Gamma^2 \succ 0. \tag{WN42}$$

The coefficient-extraction substitution $X = R/v, Z = \Gamma - sR/v$ cancels all denominators; the asserted identity itself contains no v^{-1}, and includes literal $v = 0$.

For both reciprocal angles $s, b = o(1)$, the full polynomial belongs to the square ideal of (W, X, Y, sZ, bZ), with real form

$$C_0 W^2 + A_0 Y^2 + D_0 (X - sZ)^2 + \tfrac{1}{2} X^2 + \tfrac{1}{2}(Y - bZ)^2 \succ 0. \tag{WN43}$$

These finite identities include $s = 0$ and $b = 0$ exactly.

It remains to check that the local state charts were exhaustive. Normalize a maximal state coordinate in the testing field. At $u, v = o(1)$, a unit X or Y gives positive full residue immediately. Otherwise one of W, Z is a unit. The exact sign gauges give $u, v \geq 0$. If $WZ \leq 0$, the full-core sign identity SE9 supplies separated nonnegative terms; all event packets are positive. If $WZ > 0$, choose $W = 1, Z$ bounded, or $Z = 1, W = o(1)$. These are exactly the states covered above; vanishing angles or states are included by the same polynomial identities (or by the nonpositive-sign case). Positive pure-field homogeneous normalization and real/imaginary state splitting now restore the original full parent:

$$F_c(\text{any independent projective test and state}) \ \in \ \Sigma(K_c \otimes_{\mathbb{R}} K_t)^2. \tag{WN44}$$

This proves precisely the weak-n C2 and all-β C1 weak-α source domains in the scope paragraph. It supplies the missing local leaves usually called WB, WU and CW. It does not, by itself, assert the global source partition or the finite-lift conclusion.

Verification boundary. The focused checker reconstructs the entire parent, both graph scalars, every first normal, the exact source balances, all radical conductors, both exceptional normal completions and the complete endpoint row identities. Source-order inequalities, positive-unit residue arguments and the quantified cover are proved above and in the expressly scoped WA source-entry lemma. No missing historical proof, mixed inverse, or generic SOS-cancellation assertion is used.

Chapter 202

A source-exact projective pitchfork for C1

Component `p2cdf6197677b`
Source: `NEW/root/c1_projective_pitchfork_rederived.tex`
SHA256: `2cdf6197677bd8dccbc3cbae49db98fbed0ad39cafd9ed3b96cbc2b55e5aaad1`
Inventory use: PP1-PP19 full C1 projective-pitchfork all-tests theorem, with AU19 proof-use audit

202.1 A source-exact projective pitchfork for C1

Theorem and precise source domain. Take a bounded physical canonical C1 source with $d = 1$, $\lambda = n = 0$, m, C, A, α positive bounded units and $D, \beta \geq 0$ infinitesimal. Assume the original actual finite noncommuting stationary zero contact, and retain every original source gap and intersection. Then the entire parent has a finite sum-of-squares certificate in $K_c \otimes_{\mathbb{R}} K_t$ for every independent projective test and every real or complex state. This includes literal $D = 0$ or $\beta = 0$ whenever allowed by the source domain. The two cannot vanish together at an actual contact.

202.1.1 Opening-safe source entry

Write the positive source contact as $(u_c, w_c) = (c, 1/v_c)$, and $V = w_c^2$. The retained positive-d balance proof gives c a positive bounded unit and V a positive infinitesimal. Its second stationarity equation, in the notation of that proof, is

$$a_3 = V^{-2}\{mDa_2/c^2 + \beta m a_4 + Da_5 + \beta a_7 c^2\}, \tag{PP1}$$

where all a_i are positive bounded units. Therefore

$$D = V^2\delta, \quad \beta = V^2\eta, \qquad \delta, \eta \geq 0 \text{ bounded}, \qquad \delta + \eta \text{ a positive unit}. \tag{PP2}$$

No inverse of either opening is used.

Normalize the source state by $Z_c = 1$. Set

$$p = W_c/w_c, \qquad y = Y_c/w_c, \qquad x = X_c/V. \tag{PP3}$$

All three are bounded source elements. To prove this without a shape hypothesis, restrict the reciprocal full parent PP6 to (W, Y, X) at $w = 0$, and then take residues. Its positive hard form is

$$H_u(W, Y, X) = m(W + uY)^2 + C(uW - Y)^2 + (A + \alpha + 1/2)(1 + u^2)Y^2 + (1 + u^2)X^2/2. \tag{PP4}$$

It is positive definite for every bounded real u. Scale the two W, Y kernel equations at the source by w_c, and the X equation by V. In the unknowns (p, y, x) the resulting bounded matrix has a block triangular residue: its p, y block is the positive W, Y block of PP4, and its x pivot is $(1 + c^2)/2$. The

terms $(\beta - D)/V = V(\eta - \delta)$ are bounded. All right hand sides are bounded, proving PP3 by an ordinary source-field matrix inverse with unit determinant.

Let $q_{01}, q_{10}, q_{21}, h$ be the original Laurent coefficients. Their first three residues are positive units because m, C, A, α are units. The actual zero and angular stationarity equations, using PP2, give

$$\bar{q}_{01} = \bar{q}_{21}\bar{c}^{\,4}, \qquad \bar{h} = -2\bar{q}_{21}\bar{c}^{\,2}. \tag{PP5}$$

Every other nonconstant Laurent contribution apart from $q_{10}w^2$ vanishes in this residue calculation. These are equations wholly in the coefficient field followed by its real residue map.

202.1.2 A complete local normal and divided-difference argument

It remains locally to treat testing w infinitesimal or zero and u with the same positive real residue as c. The full reciprocal quarter-parent is

$$\begin{aligned} F = {} & m(W - wX + uY - uwZ)^2 + CD_w(uW - Y)^2 + AD_uD_wY^2 \\ & + (D + 1/2)(uwW + uX - wY - Z)^2 + (\alpha + 1/2)D_u(Y - wZ)^2 \\ & + (\beta + 1/2)D_w(X + uZ)^2 - D_uD_wZ^2/2, \qquad D_u = 1 + u^2, \quad D_w = 1 + w^2. \end{aligned} \tag{PP6}$$

Use three unrestricted normals and two scalar conductors

$$N_W = W - pwZ, \quad N_Y = Y - ywZ, \quad N_X = X - xw^2Z, \qquad Q = w(u-c)Z, \quad L = (w^2 - V)Z. \tag{PP7}$$

The centers agree with the actual source contact; no equation is imposed on an independent state. Write the full normal expansion

$$F = N^T H(u, w)N + 2Z\sum_i N_i\ell_i(u, w) + E(u, w^2)Z^2, \quad N = (N_W, N_Y, N_X)^T. \tag{PP8}$$

It is an exact expansion of every term of PP6. The matrix H has bounded entries and residue PP4 at $u = \bar{c}$. Parity and direct inspection of PP6 give bounded polynomials a_W, a_Y, a_X such that

$$\ell_W = wa_W(u, b), \qquad \ell_Y = wa_Y(u, b), \qquad \ell_X = u(\beta - D) + ba_X(u, b), \quad b = w^2. \tag{PP9}$$

Actual state stationarity gives $a_W(c, V) = a_Y(c, V) = 0$ and $\ell_X(c, w_c) = 0$. Finite monic division thus gives bounded polynomials A_i, B_i with

$$a_i(u, b) = (u - c)A_i(u, b) + (b - V)B_i(b), \qquad i = W, Y.$$

Consequently the entire first two normal couplings are $Z\ell_i = A_iQ + wB_iL$.

For the third normal put $r(u) = u(\beta - D)/V + a_X(u, V)$. This is a bounded source polynomial by PP2, and $r(c) = 0$. Hence $r = (u - c)R(u)$ for a bounded polynomial R. Division in b gives another bounded polynomial B_X with

$$Z\ell_X = \{B_X(u, b) - (u - c)R(u)\}L + wR(u)Q. \tag{PP10}$$

Here the identity $V(u - c)Z = wQ - (u - c)L$ is exact. Moreover $B_X(c, 0)$ has zero residue: the source equation $u(\beta - D) + Va_X(u, V) = 0$ at $u = c$, divided by V, gives $a_X(c, V)$ infinitesimal, and $B_X(c, 0)$ has the same residue. Thus the third normal has no residue coupling to L or Q. The first two have no residue coupling to L.

For completeness, the scalar coefficient obeys

$$E(u, 0) = D + \beta u^2, \qquad E(c, V) = E_u(c, V) = E_b(c, V) = 0. \tag{PP11}$$

The first assertion follows by setting $w = 0$ in the entire PP6 after the graph substitution. The other three follow from the actual full source contact and its two angle derivatives; neither normal term is discarded in making this inference. Define $d(u) = \delta + \eta u^2$ and the bounded polynomial

$$J(u, b) = \{E(u, b) - d(u)(b - V)^2\}/b. \tag{PP12}$$

This notation means exact monic division: PP11–PP2 make the constant term zero. No testing b is inverted. At (c, V) the value and both first derivatives of J vanish. A second finite monic Taylor division gives bounded polynomials J_{20}, J_{11}, J_{02} with

$$J(u,b) = (u-c)^2 J_{20}(u,b) + (u-c)(b-V)J_{11}(b) + (b-V)^2 J_{02}(b). \tag{PP13}$$

Substitution yields the complete exact scalar identity

$$E(u,w^2)Z^2 = J_{20}(u,w^2)Q^2 + wJ_{11}(w^2)QL + \{d(u) + w^2 J_{02}(w^2)\}L^2. \tag{PP14}$$

All divisions preceded tensor evaluation, with monic divisors and bounded source coefficients. These formulas hold literally at $w = 0$.

Equations PP8–PP14 represent the entire parent by a bounded Gram on the five rows N_W, N_Y, N_X, Q, L. Its L pivot in residue is $\bar{\delta} + \bar{\eta}\bar{c}^2 > 0$, and its L crosses have zero residue. It remains to check the four-row hard and angular residue; the following exact determinant computation provides that check without taking a mixed inverse.

202.1.3 Strictness of the angular residue

At $D = \beta = 0$, replace Z by T/w in the whole reciprocal parent and take its polynomial value at $w = 0$. This gives a real four-state quadratic form $B_u(W, Y, X, T)$, with hard block PP4. The original determinant identity gives

$$\det B_u = \frac{(1+u^2)^2}{16}(q_{01} + hu^2 + q_{21}u^4). \tag{PP15}$$

Using PP5, its real Schur coefficient at T is

$$g(u) = \frac{(1+u^2)^2 q_{21}}{16 \det H_u}(u^2 - c^2)^2. \tag{PP16}$$

Here all entries and inverses are real residues; this calculation proves positivity of a real matrix and does not factor a matrix in the tensor ring. The contact centers PP3 give the unique real kernel of B_c, so the four-row residue in PP8–PP14 is precisely the quadratic Taylor coefficient in hard normals and the change $u - c$. Its angular Schur pivot is

$$g''(c)/2 = \frac{(1+c^2)^2 q_{21} c^2}{4 \det H_c} > 0. \tag{PP17}$$

The hard block is PP4 and is positive definite. Thus this full four-row residue is positive definite, as is the independent L pivot. The separated strict bounded-Gram lemma proves that PP6 is tensor SOS in the complete local chart. The rows may be unbounded or linearly dependent at a literal endpoint; only the coefficient Gram needs a strict real residue.

202.1.4 Explicit testing complement and projective endpoints

If u, v are bounded original testing coordinates, use the ordinary state matrix except near the double zero corner. At finite nonzero real residues, the residue Laurent polynomial is

$$q_{21}(u^2 - c^2)^2/u^2 + q_{10}/v^2 > 0, \tag{PP18}$$

so its state matrix is positive definite by the hard $Z = 0$ block. At $u = 0$, $v \neq 0$ the real quarter-parent is

$$m(vW - X)^2 + (C + A + 1/2)(1+v^2)Y^2 + \alpha(vY - Z)^2 + (1+v^2)X^2/2 \succ 0.$$

At $v = 0$, $u \neq 0$ it is $m(X + uZ)^2 + (C + 1/2)(uW - Y)^2 + A(1+u^2)Y^2 + \alpha(1+u^2)Z^2 + (X + uZ)^2/2$, which is likewise strict. These are residue identities; the bounded full matrices therefore give tensor SOS on their respective charts.

When both u, v are infinitesimal or zero, use the five rows (uW, vW, X, Y, Z) in the entire original finite parent. The exact polynomial substitution is bounded and its residue is

$$m(vW - X)^2 + (C + 1/2)(uW - Y)^2 + AY^2 + \alpha Z^2 + X^2/2 \succ 0 \tag{PP19}$$

as a coefficient Gram on those five rows. This treats that corner with no state normalization.

For reciprocal Alice and bounded Bob, the complete core identity RC20 applies unchanged with all m, D, α, β event packets retained whole. Its residue RC21 is strict using only $C, A > 0$. For two reciprocal testing angles, retain those positive packets and use RC24 for the four rows W, X, Y, wZ. The residue $CW^2 + AY^2 + X^2/2 + (Y - wZ)^2/2$ is strictly positive. These identities allow the reciprocal coordinate to be literally zero.

For Bob reciprocal and bounded Alice, retain the whole D, β packets, put $T = wZ$, and use the full four-row expression

$$m(W - wX + uY - uT)^2 + CD_w(uW - Y)^2 + AD_uD_wY^2 + \alpha D_u(Y - T)^2 + K/2,$$

where K is the full core RC22 with $\rho_* = uW - Y$. Its residue is exactly B_u. By PP15–PP16 it is strictly positive whenever u^2 has residue different from $\bar{c}^2$, including u infinitesimal. The remaining matched residues are PP7–PP17 after the exact Alice sign gauge if necessary.

Both signs of either angle are allowed by the exact simultaneous state sign gauges of the original parent. Every normalization is a positive square in a single source or testing field. Complex states split into real and imaginary parts. The finite and reciprocal homogeneous identities include all literal axes and projective endpoints. This completes the all-tests theorem on the stated positive-d C1 projective source domain.

Scope of verification. The focused checker expands the full parent, verifies the normal support and finite divisions with explicit contact-jet remainders, and checks PP15 and the complete outer identities. Source boundedness, vanishing contact jets, real-residue strictness and the exhaustive join are the analytical arguments above. No inference is made from a scalar certificate alone, from a historical checklist, or from mixed multiplier cancellation.

Chapter 203

R2 with an unbounded source ratio and a bounded relative current

Component `pb88ccd68a4bf`
Source: `NEW/root/r2_thin_unbounded_source_rederived.tex`
SHA256: `b88ccd68a4bf2507f0e901c07a44855b9e4ae36dabdc3c956b71871a192f8202`
Inventory use: TU1-TU23

203.1 R2 with an unbounded source ratio and a bounded relative current

Statement and domain. Use the actual original weak-C2 source RER1–RER8, normalized by $d = 1$, with $\lambda = \beta = 0$ and every original strict gap. Suppose t, a are positive unbounded, while B, aB, tB are positive infinitesimals, and put $e = 1 - r$. Assume

$$\lambda_n = n/(tB) = O(1), \qquad n \geq 0. \tag{TU1}$$

The entire original parent is tensor SOS for every independent projective test and every state. Literal $n = 0$ is included. No comparison of a testing quantity with a source quantity is made. This theorem treats precisely this source sector; it is not a completion of the other R2 sectors or the global finite-lift proof.

The bounded-Gram and finite toric allocation lemmas used below are the actual retained SC16–SC20 and QI1–QI10 proofs. We use their printed finite algebra and prove its source bounds on the present domain. We do not apply the physical C1 theorem to a formal source whose event coefficient could be negative.

203.1.1 An exact common contact and bounded source comparison

Set

$$S = 1/t, \quad z = 1/a, \quad q = a(tB), \quad w = ae, \qquad b = \sqrt{S}, \quad g = \sqrt{z}, \quad c = \sqrt{q}. \tag{TU2}$$

Thus $B = Szq$, $e = zw$, the actual squared angles are $u_c^2 = Sq$ and $v_c^2 = zq$, and S, z, q are positive infinitesimals. Keep h as the original C2 source parameter. The strict m gap gives

$$R = 1 + Sqh, \quad M = R - w > 0, \qquad 0 < w < R \sim 1. \tag{TU3}$$

Both original openings $h, 1 - h$ remain arbitrary positive quantities.

The exact RER2 identity, divided by $e > 0$, gives

$$Q_{\mathrm{RER}} = \tau\{(1 + z)(1 + Sh)(1 + Sq) + 2r(z\Delta)\lambda_n\}, \qquad \tau = q/(zw), \quad \lambda_n = n/(qz).$$

Here $Q_{\mathrm{RER}} \sim 1$, $z\Delta \sim 1$ and $r \sim 1$. The bracket is a positive bounded unit, since $\lambda_n \geq 0$ is bounded and its first term is a positive unit. Consequently

$$\tau \text{ is a positive bounded unit}, \qquad \overline{\tau} = \frac{1}{1 + 2\overline{\lambda}_n}, \qquad q = zw\tau. \tag{TU4}$$

No division by n has occurred. This sector includes the earlier thin condition $\rho_n = n/e^2 \in \mathrm{I}$: in that case the same RER identity gives $\tau \sim 1$ and $\lambda_n = \rho_n w/\tau \in \mathrm{I}$ directly, without first assuming (TU1).

For the comparison only, substitute into the native C1 rational formulas NC1–NC4

$$k^2 = Sq, \quad \eta = 1/S, \quad \zeta = z, \quad r_{\mathrm{C1}} = h, \quad t_{\mathrm{C1}} = w, \quad a_{\mathrm{C1}} = S(1-h). \tag{TU5}$$

Their θ is $e = zw$ and their R, M are exactly (TU3). Write a superscript 0 for these formal C1 coefficients. All their source denominators have the requisite positive signs:

$$K = 1 + Sq, \quad J = 1 + Sq(1-e) - S(1-h), \quad E = 1 - q(1-e), \quad Q = 1 - Sqzh,$$

and J, E, Q are positive bounded units. Their common polynomial is

$$P = 1 + qz + Sq(2 + z - 2e) + Sq^2(z + e - e^2) + S^2hq^2(1 + z - e). \tag{TU6}$$

In particular $P \sim 1$. We make no sign claim for β^0.

The two sources have exactly the same state contact. With $Z_c = 1$,

$$u_c = bc, \quad v_c = cg, \quad W_c = bg, \quad Y_c = hu_cW_c, \quad X_c = -(1-e)u_c. \tag{TU7}$$

This follows either by substitution in the two native dictionaries, or from $Y_c = hu_cW_c$, $X_c = -ru_cZ_c$ and $Z_c/W_c = \sqrt{at}$ in the actual C2 source. In particular the comparison does not move the contact.

Use the actual, not formal, normalized coefficients

$$c_b = SC, \ a_b = SA, \ D_b = D/S, \ \alpha_b = \alpha/(Sz), \quad h_2 = mzM, \quad h_b = zw/2. \tag{TU8}$$

They satisfy $m = h_2/(zM)$ and $\beta + 1/2 = h_b/(zw)$. Use the same normalizations with superscript 0, and let δ mean actual minus formal. The following explicit polynomials make all needed cancellations finite and opening-safe. Put $G = 1 + z - e$, and define

$$\begin{aligned}
P_C &= 1 + qz + S\{1 + q(2 + 2z - e) + q^2z\} \\
&\quad + S^2\{q[h(1+z) + 1 - e] + q^2[hez - e(1+z) + 2z + 1]\} + S^3hq^2G, \\
P_A &= 1 + qz + S\{q(2 + z - 2e) + q^2z(1-e)\} + S^2q^2G(G - hz), \\
P_D &= 1 + q(z+e) + S\{q[h(1+z) + 1 - e] + q^2(z + e - e^2)\} + S^2hq^2G, \\
P_m &= 1 + qz + S\{q(2 + z - 2e) + q^2z\} + S^2q^2\{(1-e)^2 + z(1-e) + hez\}, \\
P_b &= (1+z)(1+qz) + Sq(1+z)(2 + z - 2e) \\
&\quad + Sq^2z(1 + z - e^2) + S^2q^2G^2.
\end{aligned}$$

Then the exact rational identities are

$$\begin{aligned}
(1-h)\delta c_b &= -\frac{n(1-e)P_C}{zP}, & h\delta a_b &= -\frac{n(1-e)P_A}{zP}, \\
\delta D_b &= -\frac{n(1-e)(1+qz)P_D}{zJP}, & \delta h_2 &= \frac{n(1-e)(1+qz)P_m}{zP}, \\
\delta h_b &= \frac{n(1-e)P_b}{zP}, & \delta\alpha_b &= \frac{n(1-e)q^2(1+qz)[(1+S)e - S(1-h)(1+z)]}{zQP}.
\end{aligned} \tag{TU9}$$

These identities are proved simply by substituting the six printed RER1 weights and the six NC4 weights, clearing the indicated pure source denominators, and multiplying. They include $n = 0$; the right sides define no quotient by n.

After substituting $e = zw$, the polynomial

$$P_\Delta = \frac{P_b - (1+qz)P_m}{z} \tag{TU10}$$

is a genuine finite polynomial in S, z, q, w, h. Indeed the numerator specialized at $z = 0$ is zero identically, so ordinary monic division by z has zero remainder. Consequently

$$\delta(h_b - h_2) = n(1-e)P_\Delta/P. \tag{TU11}$$

Every polynomial in (TU6), (TU9), (TU10) has bounded coefficients and bounded arguments. From (TU4) and (TU9)–(TU11) we obtain

$$\begin{gathered}(1-h)\delta c_b,\ h\delta a_b,\ \delta D_b = q\lambda_n\, O(1), \qquad \delta h_2, \delta h_b = q\lambda_n\, O(1),\\ \delta\alpha_b/q^2 = \lambda_n q\, O(S+e), \qquad \delta(h_b - h_2)/(qz) = \lambda_n O(1).\end{gathered} \tag{TU12}$$

Here $O(1)$ denotes an explicit bounded rational source array with denominators among the units P, J, Q, R; it does not hide a state or testing factor. These conclusions require no lower bound on h, $1-h$, or M.

In particular $c_b \asymp 1/(1-h)$, $a_b \asymp 1/h$, $D_b \sim 1/2$, $\alpha_b/q^2 \sim 1/2$, and $h_2/q, h_b/q$ positive bounded units with common residue $1/2 + \overline{\lambda}_n$. We also have

$$h_b/w = z/2, \qquad h_b - h_2 = O(qz), \qquad H_0 := 4h_2/R + 4h_b/w \geq 2z, \quad H_0 \asymp q + z. \tag{TU13}$$

The last statements use the formal exact SC2 difference together with (TU11), and hold also when M is arbitrarily small.

203.1.2 The complete additional current, including its normal

Let the testing u, v be positive infinitesimals and first take $Z = 1$ on its pure-testing state chart. Use the exact SC coordinates

$$Q_b = (W + uY)/b, \quad Y_b = Y/S, \quad p = w/R, \quad N = ((X+u)/b - pvQ_b)/g,$$

$$H = uQ_b/b, \quad L = vQ_b/g, \quad \sigma = c_b + a_b, \quad \rho_0 = Kh/R, \quad N_y = \sqrt{\sigma}\{D_u Y_b - \rho_0 H\}. \tag{TU14}$$

Both N, N_y vanish at (TU7). The exact SC3 full-parent identity is an algebraic identity for arbitrary $c_b, a_b, D_b, \alpha_b, h_2, h_b$; no C1 coefficient relation is used in deriving it. With (TU8), it represents all the actual C2 parent except its additional term $4nD_uD_vX^2/S$. Retain its original nonnegative m packet $4h_2wN^2/(MR)$, incorporate the second paired packet into H_0N^2, and retain $4h_b(vN)^2/w$ exactly as in the uniform QI normal lemma. Define $\widehat{N} = \sqrt{H_0}N$.

Put $h_t = H/R$, and introduce the same four positive separated monomials and their geometric mean as in QI:

$$A_* = ch_t, \quad B_* = cL/R, \quad C_* = uv/b, \quad D_* = qg, \qquad \Gamma = (A_*B_*C_*D_*)^{1/4}. \tag{TU15}$$

Their exact relations include $\Gamma^2 = qh_tv$ and $B_*C_* = (c/g)v^2h_t$. The extra monomial

$$U_* = cgu/b$$

has the two exact identities

$$U_*B_* = \Gamma^2, \qquad cgX/b = czN + eB_* - U_*. \tag{TU16}$$

The entire additional current therefore has the following square completion, with no state normal set to zero:

$$\frac{4n}{S}D_uD_vX^2 = 4\lambda_nD_uD_v(U_* - B_* - czN)^2$$

$$+ 4\lambda_n D_u D_v\{2(1-e)\Gamma^2 - (1-e^2)B_*^2 - 2(1-e)czB_*N\}. \tag{TU17}$$

The first line is a nonnegative separated SOS and vanishes at the actual contact, including its unrestricted state-normal coordinate. Its coefficient is literally zero if $n = 0$. The remaining scalar changes have bounded coefficients on B_*^2 and Γ^2. Their possibly nonzero real residues will cancel exactly with the source comparison below. The full additional first-normal coefficient, on $B_*\widehat{N}$, is infinitesimal because

$$\lambda_n cz/\sqrt{H_0} = O(\lambda_n c\sqrt{z}) \in \mathrm{I}. \tag{TU18}$$

In particular (TU17) does not estimate or discard u/b, which can be arbitrarily large. The new positive square contains that entire row, while the exact product relation in (TU16) places its cross term on the old angular circuit.

203.1.3 Why the full QI allocation is stable here

We verify the actual finite hypotheses of QI1–QI10 rather than an abstract continuity assertion. Its coefficient arrays have only the pure-testing denominators D_u and positive pure-source units after cancellation of the two h openings. The latter cancellation is preserved exactly because

$$1 - \rho_0 = (1-h)/R, \qquad \rho_0 = Kh/R. \tag{TU19}$$

For example all perturbations of $c_b(1-\rho_0)^2$, $a_b\rho_0^2$, $c_b(1-\rho_0)$ and $a_b\rho_0$ are $q\lambda_n$ times bounded rational arrays, by (TU9). Moreover

$$\delta(\sigma\rho_0 - c_b) = \rho_0\delta a_b - (1-\rho_0)\delta c_b = q\lambda_n O(1). \tag{TU20}$$

This is the complete potentially singular harmonic first-normal coefficient after multiplication by $\sqrt{\sigma}$. Both σ and its formal comparison are comparable to $[h(1-h)]^{-1}$, and their ratio has residue one. Their square-root ratios and inverses are consequently bounded pure-source units. Thus this change of hard normal loses no opening bound.

In the exact generic scalar QI2, the changes of a, b_*, c_* are $q\lambda_n$ times bounded arrays in u, v. This follows term by term from (TU12), (TU19): its only C, A occurrences are the first two expressions in (TU19); its D_b occurrences have bounded factors; its α_b occurrences have bounded factors and the additional small factors printed in QI2. Consequently the formal specialization a_0, b_0, c_0 and its exact U, V divisions in QI3–QI4 can be used unchanged, while the remaining q coquotients change by bounded arrays. Write $\lambda_0 = \overline{\lambda}_n$. Their complete leading changes are respectively $0, -8\lambda_0, 4\lambda_0, 0$: substituting $S = z = q = u = v = 0$ in the bounded coquotients (TU9) gives $\delta a/q = -4\lambda_0$, $\delta b_*/q = 0$, $\delta c_*/q = -4\lambda_0$, and $\delta(R^2P_L)/q = 4\lambda_0$. Thus before the extra current the four residues are

$$\overline{a_1 - c_1} = 2, \qquad \overline{b_1 + 2c_1} = -8 - 8\lambda_0, \qquad \overline{R^2P_L/q} = 2 + 4\lambda_0, \qquad \overline{4\alpha_b D_u/q^2} = 2. \tag{TU21}$$

Equation (TU17) adds $-4\lambda_0$ to the B_*^2 residue and $+8\lambda_0$ to the Γ^2 residue. Consequently the *combined* full scalar has exactly the original four residues $(2, -8, 2, 2)$ on $(A_*^2, \Gamma^2, B_*^2, D_*^2)$. All differences from those residues are infinitesimal bounded arrays, because λ_n itself is bounded. This cancellation is why a thinness assumption is unnecessary on (TU1). The one additional bound QI uses from physical $\beta \geq 0$ is also available directly, without any sign for β^0:

$$w/(c/g) = \sqrt{w/\tau} = O(1), \qquad w = O(1). \tag{TU22}$$

Thus the exact dangerous cross term $-4(wv/(c/g))D_uB_*C_*$ still has infinitesimal coefficient.

For the complete first N normal, (TU13) gives exactly the two bounds used in the uniform QI normal lemma: $g/\sqrt{H_0}$ and $c/\sqrt{H_0}$ are bounded, and

$$\frac{h_bD_v - h_2}{\sqrt{qH_0}} = O(z + v^2). \tag{TU23}$$

The H term and all other terms in SC14 therefore have the same infinitesimal coefficients on the QI rows; (TU18) handles the whole extra term. For the harmonic normal, (TU20) adds a $q\lambda_n$ bounded array to the left side of QI10. It can be absorbed into its existing qK_q coquotient. All other terms are the same bounded rational arrays multiplied by the opening-safe quantities in (TU12), (TU19). The exact D_b root and its two normal terms in QI9 remain unchanged as algebraic identities. The complete hard block therefore retains the real residue $\mathrm{diag}(1,4)$ in $(\widehat{N}, N_y)$.

For clarity, here is the finite allocation being invoked. Use the QI roots

$$E_* = h_t - v, \quad F_* = vh_t - gD_*, \quad R_F = \frac{\sqrt{g\Gamma}}{c}(\Gamma - D_*), \quad R_E = \sqrt{\Gamma}(\sqrt{A_*} - \sqrt{cv}),$$

together with the complete finite toric root list SC16. The exact identities QI7–QI8 give strict real square pivots on F_*, R_F, R_E; QI6 gives the strict E_* pivot and twice the four-monomial circuit. Every remaining coefficient is an infinitesimal bounded rational array in the testing u, v, with the same finite denominators and divisions just verified. All removed packets, including (TU17), vanish at (TU7). The actual source contact value, its three scalar first jets, and both full first-normal contact values consequently vanish exactly for this remaining expression. Applying the two finite monic divisions SC17–SC20 and the displayed QI7 first-jet identities therefore puts the entire scalar correction and both complete first normals on this strict root list with infinitesimal coefficients. The coefficient functions are independent of the state ratio Q_b/Z; no boundedness of that ratio is assumed. This is precisely the constructive QI allocation, with its source inequalities checked in (TU12)–(TU23), not a conclusion from a scalar restriction alone.

The resulting finite Gram is bounded and has positive definite real residue. The strict bounded-Gram lemma supplies tensor SOS. Restore the original positive paired packets and the new full square in (TU17). This proves the local assertion for the entire parent on the positive small-angle, negative-core chart.

203.1.4 States, signs, and all remaining tests

Every inverse and radical used above is pure source, pure testing, or the explicitly separated positive monomial radical in (TU15). For $Z \neq 0$ the normalization $Z = 1$ is pure testing; multiplying back by Z^2 restores homogeneity. At $Z = 0$ the original pure-testing core restriction is nonnegative. The exact core Schur identity and sign gauges used in the retained QI proof force $W + uY > 0$ on the remaining negative-core chart, and hence all four monomials are positive. Otherwise the core and all original event packets already give the direct exit. Literal zero-angle charts have that same original exit. Complex states split into their real and imaginary parts, since the real measurement matrices and full quadratic parent are unchanged.

The new R2 outer theorem R2O1–R2O24 treats every projective testing pair outside the double-infinitesimal original chart, with no condition on a or n/e^2. Its hypotheses are exactly the original RER source conditions already retained here. Combining that result with this local proof gives the claimed all-tests, all-states theorem on (TU1). No other source branch is admitted by this conclusion.

Verification scope. The associated checker verifies the exact native contact map, all six comparison formulas (TU9), the finite division (TU10), the four complete residue cancellations in (TU21), the whole generic SC reduction, the entire additional-current completion (TU17), and its actual contact zero. Ordered-field bounds, the complete finite QI allocation and the outer-chart join are analytical parts of this proof, not consequences of the number of checker assertions.

Chapter 204

R2 with both the native source ratio and relative current unbounded

Component `pef021eabacde`
Source: `NEW/root/r2_unbounded_current_rederived.tex`
SHA256: `ef021eabacde61ad9b1ed3c993a53db009b9974528f85eb04cfab31404534ea5`
Inventory use: UL1-UL18

204.1 R2 with both the native source ratio and relative current unbounded

Precise assertion. Retain the original RER source, all its strict gaps, and the notation TU2–TU16 of the preceding bounded-relative-current proof. Assume the native a, t are positive unbounded and

$$\lambda = \frac{n}{tB} = \frac{n}{qz} \text{ is positive unbounded.} \tag{UL1}$$

Then the whole original weak-C2 parent is tensor SOS for every independent projective test and every state. The argument below complements TU1; it never assigns a sign to the formal C1 coefficient β^0, and never replaces the original physical source by that formal source.

204.1.1 Source bounds without a thinness assumption

The exact RER physicality equation already used in TU4 is

$$Q_{\mathrm{RER}} = \tau\{A + 2r(z\Delta)\lambda\}, \qquad \tau = \frac{q}{zw}, \qquad A = (1+z)(1+Sh)(1+Sq).$$

Here $A, Q_{\mathrm{RER}}, r, z\Delta$ are positive units with residue one. Since λ is unbounded, this proves, with no cancellation of mixed factors,

$$\tau \in \mathrm{I}, \qquad \lambda\tau \sim \frac{1}{2}, \qquad \lambda q \sim \frac{zw}{2}, \qquad \frac{n}{z^2 w} \sim \frac{1}{2}. \tag{UL2}$$

All four divisions are pure source. Recall $0 < w < R \sim 1$. The exact TU9 formulas therefore imply

$$\begin{gathered}
(1-h)\delta c_b,\ h\delta a_b,\ \delta D_b,\ \delta h_2, \delta h_b = \lambda q\, O(1), \\
\delta\alpha_b/q^2 = \lambda q\, O(S + zw), \qquad h_2 \asymp zw, \quad h_b = zw/2, \\
H_0 = 4h_2/R + 2z \asymp z, \qquad H_0 \geq 2z, \\
\frac{g}{\sqrt{H_0}} = O(1), \quad \frac{c}{\sqrt{H_0}} = O(1), \quad \frac{\lambda q}{H_0} = O(1), \quad \frac{n}{z^2} = O(1).
\end{gathered} \tag{UL3}$$

In particular λq is infinitesimal. The displayed coefficient arrays have denominators only among the pure-source units P, J, Q, R after the original $h, 1-h$ openings cancel. The actual and formal

$\sigma = c_b + a_b$ differ by a relative infinitesimal. Their square-root ratio is a bounded unit. No lower bound on $w, h, 1-h$ or the strict gap M is assumed.

Use the complete generic SC3 identity with the actual coefficients. Retain its positive m packet and use $\widehat{N} = \sqrt{H_0}N$. The formal comparison is used only for finite scalar and normal identities. Its q-coquotients and the QI coefficient arrays are bounded here: their derivation requires S, z, q infinitesimal and w bounded, but not $\beta^0 \geq 0$. The sole QI bound that is no longer available, $w/(c/g) = O(1)$, is replaced explicitly below.

204.1.2 The entire extra current supplies a new angular pivot

Use the exact positive-chart rows TU15–TU16, denoted here by A_*, B_*, C_*, D_*, U_* and Γ. Define

$$J_3 = \sqrt{\lambda}(U_* - B_*), \qquad E_* = h_t - v, \qquad F_* = vh_t - gD_*. \tag{UL4}$$

The first line of the exact whole-current identity TU17 is

$$4D_uD_v\left(J_3 - d_N\widehat{N}\right)^2, \qquad d_N = \frac{\sqrt{\lambda}\,cz}{\sqrt{H_0}} = \sqrt{\frac{nz}{H_0}} \in \mathrm{I}. \tag{UL5}$$

Thus it supplies a strict real pivot on J_3, with only infinitesimal changes to the hard normal Gram. Its scalar remainder is still exactly $8\lambda D_uD_v(1-e)\Gamma^2 - 4\lambda D_uD_v(1-e^2)B_*^2$. Its complete first-normal remainder, retained below, is $-8\lambda D_uD_v(1-e)czB_*N$.

Write $\widetilde{c}, \widetilde{a}, \widetilde{d}, \widetilde{\alpha}, \widetilde{m}, \widetilde{b}$ for the six differences $\delta c_b, \delta a_b, \delta D_b, \delta\alpha_b, \delta h_2, \delta h_b$ divided by λq. They are defined explicitly by the six right sides in TU9, using $n = \lambda qz$; their apparent opening poles cancel in every combination below. Put $U = u^2, V = v^2$, and retain $\rho_0 = Kh/R$, $a_t = 1 - \rho_0 + (w/R)V$. Define the exact rational arrays

$$\begin{aligned}
\mathsf{a} &= 4D_vR^2\{\widetilde{c}(1-\rho_0)^2 + \widetilde{a}\rho_0^2/D_u\} + 4\widetilde{\alpha}zS^2VR^2\rho_0^2/D_u + 4\widetilde{d}S^2R^2a_t^2,\\
\mathsf{b} &= -8\widetilde{\alpha}zSR\rho_0 - 8\widetilde{d}SRa_tD_u,\\
\mathsf{c} &= 4\widetilde{d}D_u^2,\\
\mathsf{l} &= 4R\{\widetilde{m}(1-w/R) + \widetilde{b}(w/R)D_v\}.
\end{aligned} \tag{UL6}$$

For example $\delta a/q = \lambda\mathsf{a}$ in QI2, and $\delta(R^2P_L/q) = \lambda\mathsf{l}$. Direct specialization of TU9 gives, keeping S, z, w, h exact,

$$\begin{aligned}
(\mathsf{a} - \mathsf{c})|_{q=U=V=0} &= 0,\\
(\mathsf{b} + 2\mathsf{c} + 8D_uD_v(1-e))|_{q=U=V=0} &= 0,\\
(\mathsf{l} - 4D_uD_v(1-e^2))|_{q=U=V=0} &= 0.
\end{aligned} \tag{UL7}$$

These are stronger than residue equalities. Clear the unit denominators and divide successively by the monic variables q, U, V. Each of the three expressions is consequently $qF_q + UF_U + VF_V$, with finite bounded rational coefficient arrays. The division does not invert q, U, V in the tensor ring.

The combined full scalar correction to QI6 therefore consists of bounded-array multiples of $\lambda q, \lambda U, \lambda V$ on A_*^2, Γ^2, B_*^2, an infinitesimal coefficient on E_*^2, and the infinitesimal α correction on D_*^2. The following exact identities allocate every apparently unbounded testing term:

$$\begin{aligned}
\lambda U A_*^2 &= \lambda q(uE_* + bC_*)^2,\\
\lambda V A_*^2 &= \lambda q(F_* + gD_*)^2,\\
\lambda U\Gamma^2 &= \lambda q\, bC_*(uE_* + bC_*),\\
\lambda V\Gamma^2 &= \lambda q(F_* + gD_*)(F_* + gD_* - vE_*),\\
\sqrt{\lambda}\,uB_* &= \sqrt{nS/z^2}\,(F_* + gD_*),\\
\sqrt{\lambda}\,vB_* &= \sqrt{n}\,C_* - vJ_3.
\end{aligned} \tag{UL8}$$

Every coefficient introduced on squaring the last two identities is infinitesimal by UL2–UL3 and $S, u, v \in \mathrm{I}$. Finally the formerly dangerous QI6 term has the exact replacement

$$\frac{wg}{c}\, vB_*C_* = wz\, C_*^2 - \frac{wg}{c\sqrt{\lambda}}\, vC_*J_3, \qquad \frac{wg}{c\sqrt{\lambda}} = w\sqrt{\frac{z^2}{n}} = O(1). \tag{UL9}$$

Both resulting coefficients are infinitesimal. This retains the entire term for arbitrary independent source and testing rates.

204.1.3 Both unrestricted state-normal corrections

The new complete first N coefficient, including the remainder of TU17, differs from the formal generic SC coefficient by

$$\begin{aligned} \delta\mathcal{L}_N &= 8\lambda c(1-e)\mathsf{F}_N B_* + 8\lambda q\widetilde{d}\, ubvg(Sa_tH - D_u v), \\ \mathsf{F}_N &= \frac{P_b D_v - (1+qz)P_m}{P} - zD_uD_v = V - zUD_v + qF_{Nq}. \end{aligned} \tag{UL10}$$

The last equality is a finite monic q division; its coquotient has bounded arrays because the numerator at $q = 0$ is exactly $V - zUD_v$. After division by $\sqrt{H_0}$, its q part is infinitesimal on B_* by UL3. The other parts use

$$\begin{aligned} \lambda cVB_* &= \lambda qg\, vC_* - \sqrt{\lambda q}\, v^2 J_3, \\ \lambda czUB_* &= \lambda qbg\, v(uE_* + bC_*). \end{aligned} \tag{UL11}$$

Their coefficients on $\widehat{N}$ are infinitesimal since $g/\sqrt{H_0}$ and $\sqrt{\lambda q/H_0}$ are bounded. The $\widetilde{d}$ term in UL10 uses $uH = R(uE_* + bC_*)$ and $uv = bC_*$; it has the same infinitesimal coefficient bounds. Thus this is a lift of the complete first normal, with no state component put to zero.

For the second normal it is convenient to compare the coefficient after multiplication by $\sqrt{\sigma}$. Differentiating the full generic SC3 identity, including the restored D packet, gives

$$\sqrt{\sigma}\, \delta\mathcal{L}_y = 8\lambda q\{\mathsf{F}_H H + \mathsf{F}_v v\}, \tag{UL12}$$

where

$$\begin{aligned} \mathsf{F}_H &= D_v\{\rho_0\widetilde{a} - (1-\rho_0)\widetilde{c}\} - D_v\widetilde{a}U\rho_0/D_u + \widetilde{\alpha}zS^2V\rho_0/D_u - \widetilde{d}S^2a_t, \\ \mathsf{F}_v &= \widetilde{d}SD_u - \widetilde{\alpha}zS. \end{aligned}$$

These arrays are bounded through both source openings. Exactly,

$$\mathsf{F}_H|_{q=U=V=0} = \frac{(1-e)S}{1-S(1-h)}, \qquad \mathsf{F}_v|_{q=U=V=0} = -\frac{(1-e)S}{1-S(1-h)}. \tag{UL13}$$

As $R = 1$ there, finite monic division gives $R\mathsf{F}_H + \mathsf{F}_v = qF_q + UF_U + VF_V$ with bounded arrays. Since $H = R(E_* + v)$, the whole UL12 coefficient is therefore an infinitesimal combination of E_*, F_*, A_*, C_*, D_*: use only

$$qv = cA_* - qE_*, \qquad Uv = buC_*, \qquad Vv = v(F_* + gD_* - vE_*). \tag{UL14}$$

The factor $1/\sqrt{\sigma}$ is bounded. Changing the formal harmonic-normal normalization to the actual one multiplies its previously bounded arrays by a pure-source unit with residue one.

For completeness, the unchanged formal first N coefficient is bounded after the actual $\sqrt{H_0}$ normalization: $h_2^0, h_b^0 = O(q)$ and $h_b^0 - h_2^0 = O(qz)$, while $g/\sqrt{H_0}$ and $c/\sqrt{H_0}$ are bounded. These are exactly the two estimates used in the QI normal allocation. All remaining hard coefficients from the original C, A, D, α lines have the same residues as before; the relative C, A changes and the absolute D, α changes are infinitesimal. Together with UL5 the complete hard block on $\widehat{N}, N_y, J_3$ has a positive definite real residue.

204.1.4 A finite contact lift for the additional row

It remains to allocate the full scalar correction on roots that vanish at contact. Merely bounding a coefficient on A_*, B_*, C_*, D_* would not suffice. Keep the complete strict toric root list SC16, the QI rows E_*, F_*, R_F, R_E, and add

$$V_3 = \sqrt{\lambda\Gamma}(\sqrt{U_*} - \sqrt{B_*}).$$

Since $U_* B_* = \Gamma^2$, the exact identities are

$$\begin{aligned} J_3^2 &= 4V_3^2 + \lambda(\sqrt{U_*} - \sqrt{B_*})^4, \\ \Gamma J_3 &= \frac{V_3^2}{\sqrt{\lambda}} - 2\sqrt{\lambda}\,\Gamma(B_* - \Gamma). \end{aligned} \tag{UL15}$$

Reserve half the J_3^2 coefficient in UL5 for the first identity. It supplies strict real pivots on both J_3, V_3, with the quartic square retained nonnegative. The normal coupling in UL5 remains an infinitesimal hard Gram correction.

Here are the additional first-jet coefficient checks needed for the second identity, so that its factor $\sqrt{\lambda}$ causes no hidden loss. The only new scalar terms linear in J_3 in UL8–UL9 have coefficients, up to bounded arrays, of the forms

$$\epsilon_B = -2\sqrt{n}\,v, \qquad \epsilon_T = \frac{wg}{c\sqrt{\lambda}}\,v. \tag{UL16}$$

Both have bounded finite testing difference arrays and are infinitesimal. At the actual pure-source contact $v_c = cg$,

$$\sqrt{\lambda}\,\epsilon_B(u_c, v_c) = -2n \in \mathrm{I}, \qquad \sqrt{\lambda}\,\epsilon_T(u_c, v_c) = wz = e \in \mathrm{I}. \tag{UL17}$$

Multiplication by any of the bounded arrays produced in UL7 preserves these conclusions. No other scalar term introduces J_3 linearly: the other five identities in UL8, and the λq scalar corrections, use only the old QI rows.

Write a term $\epsilon J_3 A_i$ as $\epsilon J_3(A_i - \Gamma) + \epsilon\Gamma J_3$. The first summand is already an infinitesimal quadratic on the strict list. Divide $\epsilon - \epsilon(u_c, v_c)$ finitely by $u - u_c, v - v_c$ and apply the exact SC18 conductors to $\Gamma(u - u_c)$ and $\Gamma(v - v_c)$. This allocates the nonconstant part as infinitesimal quadratic corrections involving J_3. For the constant part use UL15. Its V_3^2 coefficient is infinitesimal, and its remaining coefficient on $\Gamma(B_* - \Gamma)$ is infinitesimal by UL17.

There are also new bare-Γ E_*, F_* terms when the first five identities in UL8 are expanded and their C_*, D_* factors are written as $(A_i - \Gamma) + \Gamma$. Put $L = \lambda q$ and $M_1 = nS/z^2$, both infinitesimal. The additional E_* coefficients are bounded-array multiples of Lbu and Lgv; the additional F_* coefficients are bounded-array multiples of Lg and $M_1 g$. At $u_c = bc, v_c = cg$ their required contact ratios are

$$\frac{Lbu_c}{c} = LS, \qquad \frac{Lgv_c}{c} = Lz, \qquad \frac{Lg}{g} = L, \qquad \frac{M_1 g}{g} = M_1. \tag{UL18}$$

All are infinitesimal, with bounded finite difference arrays before these contact substitutions. The exact QI7 identities therefore allocate these additional terms by the same finite division as the old E_*, F_* terms. This check is necessary: the two new normal rows cannot be ignored when applying the contact-jet argument.

After these explicit additional steps, the remainder has exactly the QI form: a bounded infinitesimal quadratic on the listed roots, the now-verified bare-Γ E_*, F_* couplings, and a toric expression $\Gamma \sum_i \mu_i(A_i - \Gamma) + \Gamma^2\sigma$. The old E_*, F_* contact coefficient ratios remain the bounded ones in QI7–QI8: their unchanged $q = 0$ arrays depend on bounded w and do not require $w/(c/g)$ bounded. Every retained packet, including UL5 and UL15, vanishes at the actual contact. The complete actual scalar consequently has zero contact value and all three zero first jets. Apply the two explicit finite divisions SC17–SC20. Their three-jet balance makes the constant toric first-jet coefficients equal; SC19 and the variance roots then allocate this common term. This puts the entire scalar on a bounded Gram with strictly positive real residue.

The full normal lifts UL10–UL14 have exact contact zero as well. Their J_3, E_*, F_* terms already vanish there. The sum of their remaining four monomial coefficients is zero at contact, so one finite SC18 division allocates them on the same strict list. Every coefficient function in these steps depends only on the testing u, v. No bound on the state ratio Q_b/Z is used, and the third contact jet is used only for its exact toric balance.

The strict bounded-Gram lemma now proves tensor SOS for the whole original parent on the small-angle positive negative-core chart. All its radicals are radicals of separated positive monomials; every inverse is pure source or pure testing.

204.1.5 Projective completion and scope

The state, sign, zero-axis and real/imaginary-state arguments are the same exact original-core arguments at the end of TU. In particular $Z = 0$ is an original positive restriction, and the negative-core $Z = 1$ chart forces $W + uY > 0$, permitting the separated radicals above. The theorem R2O covers every testing angle chart outside u, v both infinitesimal without any condition on λ or the native a. This proves UL1 for all tests and states.

Combining this theorem with TU covers the entire original R2 sector with native a unbounded: its nonnegative $\lambda = n/(tB)$ is either bounded, including zero, or positive unbounded. The complementary bounded-a theorem remains a separate input to the final R2 join. This local result alone is not the universal tensor-SOS theorem or finite-lift existence.

Verification boundary. The focused exact checker verifies UL7, UL10, UL13, the full generic first-normal derivatives before any graph restriction, all six UL8 conductors, UL9, UL11, UL14–UL17 and the actual contact of J_3. The native source comparisons are the separately verified TU9 dependency. The source inequalities, bounded finite arrays, finite contact allocation and projective join are the analytical proof above, not consequences of assertion counts.

Chapter 205

A direct certificate for the positive-d C2 reciprocal corner

Component `p10686597065a`
Source: `NEW/root/rc_reciprocal_full_parent_rederived.tex`
SHA256: `10686597065a0b7f3045ed278b33a0e8d8ec27622e5d29f16037724986887652`
Inventory use: RC1-RC25 full C2 reciprocal all-tests theorem, with bare-scope proof-use audit AU19 | Only RC18-RC25 explicit outer whole-parent identities and real strictness formulas, used with n=0 and entire extra beta packet retained

205.1 A direct certificate for the positive-d C2 reciprocal corner

Precise scope. Normalize the original C2 source by $d = 1$ and $\lambda = \beta = 0$. Assume that all source coefficients are bounded, that C, A, α are positive bounded units, that m, D are positive infinitesimals, and that $n \geq 0$ is bounded. Retain the original physicality, finite noncommuting stationary zero contact, positive native gaps, and any original prior-complement predicates. This section proves the complete parent is tensor SOS at every independent projective test and every state. The contact is used only inside the source field. No mixed determinant is cancelled.

The source balance portion of the retained positive-d partition gives, with reciprocal Bob contact coordinate $w_c = 1/v_c$,

$$U = u_c^2 \in \mathrm{Inf}_{>0}, \qquad V = w_c^2 \in \mathrm{Inf}_{>0}, \qquad m \asymp U^2, \qquad D \asymp V^2. \tag{RC1}$$

Only the displayed balance proof, not its references to missing terminal certificates, is used here. Its two angular stationarity equations have positive-unit factors and prove RC1 directly.

205.1.1 The actual source has a bounded shape chart

Choose the native positive contact chart with $Z_c = 1$, and set

$$S = (u_c/w_c)W_c, \qquad y = Y_c/w_c, \qquad R = -X_c/(u_c V). \tag{RC2}$$

Then $S, y, S - y, 1 - y, R$ are positive bounded units. Here is an entry proof that does not assume those facts as extra source hypotheses. In the reciprocal parent RC6 below, take the ordinary source-field Schur complement of the X, Y block. That block has residue $\mathrm{diag}(n + 1/2, H)$, where $H = C + A + \alpha + 1/2$. Its W, W Schur entry divided by U has residue $a = C - C^2/H > 0$, while its W, Z entry divided by $u_c w_c$ has residue $-b = -(1/2 + C\alpha/H) < 0$. The balance $m/U \in \mathrm{Inf}$ and $D/V \in \mathrm{Inf}$ justifies these divisions in the source field. The zero-state equation therefore gives $S \sim b/a$, a positive bounded unit.

The two hard state equations, scaled in the source field by $u_c V$ and w_c, are a bounded linear system in R, y. Its residue is triangular with positive diagonal $n + 1/2, H$. Consequently both quantities are bounded. The second equation gives $Hy = CS + \alpha$ in residue, so y is a positive unit.

The W equation divided by $u_c w_c$ gives $C(S-y)=1/2$ in residue. The Z equation divided by V gives $\alpha(1-y)=S/2$ in residue: its apparently constant U terms cancel exactly before the division. Finally the first hard equation gives $(n+1/2)R=(1+S-y)/2$ in residue. These relations prove all five claimed positive-unit bounds. The exact native gaps already give $0<y<1$ and $S-y>0$ before taking residues.

The unchanged native source formulas are converted by the following dictionary. Their letters are used only in this display:

$$\begin{aligned} z=U,\quad a^2=1/(UV),\quad b=S/U,\quad t=(S-y)V,\quad r=VR,\quad h=y/U,\\ g=S/U+y-1+VR,\qquad j=1+UVR-V(S-y). \end{aligned} \tag{RC3}$$

Substitution into the six exact native coefficients has denominators with positive-unit leading parts after the displayed powers of U,V are removed. Thus $\mu=m/U^2$ and $\delta=D/V^2$ are bounded units, and the source residues are

$$\begin{aligned} C_0&=\frac{1}{2(S-y)}, & A_0&=\frac{1+S-y}{2y}, & \alpha_0&=\frac{S}{2(1-y)},\\ n_0&=\frac{1+S-y-R}{2R}, & \mu_0&=\frac{1+S-y}{2S^2}, & \delta_0&=\frac{S(1+S-y)}{2}. \end{aligned} \tag{RC4}$$

In residue formulas such as RC4 the shape letters mean their real residues. In particular

$$C_0\{\alpha_0(A_0-1/2)-1/4\}=(A_0+\alpha_0+1/2)/4. \tag{RC5}$$

These are source statements only. They do not identify any source infinitesimal with an independent testing infinitesimal.

205.1.2 The complete local parent and both unrestricted normals

For u,w infinitesimal or zero in the independent testing field, put $D_u=1+u^2$, $D_w=1+w^2$. The full quarter-parent after the pure testing reciprocal Bob normalization is

$$\begin{aligned} F=\ &m(W-wX+uY-uwZ)^2+nD_uD_wX^2+CD_w(uW-Y)^2+AD_uD_wY^2\\ &+(D+1/2)(uwW+uX-wY-Z)^2+(\alpha+1/2)D_u(Y-wZ)^2\\ &+\tfrac{1}{2}D_w(X+uZ)^2-\tfrac{1}{2}D_uD_wZ^2. \end{aligned} \tag{RC6}$$

All four state coordinates are arbitrary. Introduce tensor rows

$$\begin{aligned} N_x&=X+uRw^2Z, & N_y&=Y-ywZ,\\ \rho&=uW-SwZ, & k&=UW-SuwZ, & \ell&=(w^2-V)Z. \end{aligned} \tag{RC7}$$

The testing-dependent w^2 in N_x is essential; it agrees with the source contact and removes the unbounded coefficient produced by a contact-fixed center. The exact conductor identities are

$$U\rho-uk=Sw(u^2-U)Z,\qquad u\rho-k=(u^2-U)W. \tag{RC8}$$

Expanding the entire RC6 gives

$$F=F_0+H_xN_x^2+H_yN_y^2+2H_{xy}N_xN_y+2N_xL_x+2N_yL_y, \tag{RC9}$$

where F_0 is only the constant coefficient of this unrestricted normal expansion. The hard coefficients are exactly

$$\begin{aligned} H_x&=mw^2+nD_uD_w+(D+1/2)u^2+D_w/2,\\ H_y&=mu^2+CD_w+AD_uD_w+(D+1/2)w^2+(\alpha+1/2)D_u,\\ H_{xy}&=-(m+D+1/2)uw. \end{aligned} \tag{RC10}$$

Set $J_x = n(1+w^2) + D + 1/2$ and $K_x = n(1+U) + 1/2 + m$. The complete first normal is

$$L_x = (D+1/2)uw\rho - (m/U)wk + u\{D/V - Rw^2K_x\}\ell - (R/S)uwJ_x(U\rho - uk). \tag{RC11}$$

Every coefficient on ρ, k, ℓ is bounded infinitesimal. For the other normal define

$$\begin{aligned} a_{10} &= Ay + (m + \alpha + 1/2)(y-1), \\ a_{01} &= -(C+D+1/2)S + (C+A+D+1/2)y, \\ a_{11} &= R(m+D+1/2) + Ay. \end{aligned}$$

Then the exact expression is

$$\begin{aligned} L_y = \{&m - C - (C+D+1/2)w^2 + (U/S)(a_{10} + a_{11}w^2)\}\rho \\ &- (u/S)(a_{10} + a_{11}w^2)k + w(a_{01} + a_{11}U)\ell. \end{aligned} \tag{RC12}$$

Its only nonzero residue coupling is $-C_0\rho$. RC11–RC12 follow by full expansion and the two actual source state equations; they retain every state-normal term of RC6. Their algebra is checked against the complete state Hessian, not a graph restriction.

205.1.3 A scalar certificate by finite monic divisions

Put

$$c = C + (C+D+1/2)w^2, \quad b_2 = (D+1/2)R, \quad b_1 = (C+D+1/2)(y-S) - mR, \quad \gamma = b_1 + b_2U.$$

Define

$$F_{\mathrm{base}} = (c - 2Ub_2w^2/S)\rho^2 + \mu k^2 + (2ub_2w^2/S)\rho k - 2w\gamma\rho\ell. \tag{RC13}$$

The source W state equation removes the remaining WZ coefficient. Direct expansion then gives $F_0 - F_{\mathrm{base}} = E(u^2, w^2)Z^2$, with $E(a,b)$ a bounded-coefficient polynomial of degree at most two in a. Its exact properties are

$$\begin{aligned} E(a,0) = D, \qquad E(U,V) = E_a(U,V) = E_b(U,V) = 0, \\ E(a,b) = E_0(b) + (a-U)E_1(b) + (a-U)^2E_2(b), \\ E_2(b) = R^2b^2\{n(1+b) + D + 1/2\}. \end{aligned} \tag{RC14}$$

The three vanishing jets follow from the actual full contact: every row in RC7 vanishes there, the complete scalar and both angle derivatives vanish, and RC13 has zero first derivatives there. They are also checked directly from the native source coefficients. Since $E_1(0) = E_1(V) = 0$ and E_0 has a double zero at V, finite monic polynomial division gives bounded polynomials J_0, J_1 such that

$$E_0(b) = (b-V)^2J_0(b), \qquad E_1(b) = b(b-V)J_1(b). \tag{RC15}$$

These divisions take place in the source polynomial algebra before tensor evaluation. They introduce no inverse of V or of a mixed quantity. In particular $V^2J_0(0) = D$, and the other coefficients of J_0 remain bounded; hence $J_0(w^2)$ has residue δ_0.

RC8 now gives the exact scalar identity

$$\begin{aligned} F_0 = F_{\mathrm{base}} &+ \frac{R^2w^2\{n(1+w^2) + D + 1/2\}}{S^2}(U\rho - uk)^2 \\ &+ \frac{w}{S}J_1(w^2)(U\rho - uk)\ell + J_0(w^2)\ell^2. \end{aligned} \tag{RC16}$$

This polynomial formula also holds at $u = 0$ or $w = 0$; neither is inverted. Combining RC9–RC16 represents the entire parent by a bounded Gram on $(N_x, N_y, \rho, k, \ell)$ with real residue

$$(n_0 + 1/2)N_x^2 + H_0N_y^2 - 2C_0N_y\rho + C_0\rho^2 + \mu_0k^2 + \delta_0\ell^2, \qquad H_0 = C_0 + A_0 + \alpha_0 + 1/2. \tag{RC17}$$

This is positive definite: the N_y, ρ determinant is $C_0(A_0 + \alpha_0 + 1/2) > 0$, and the other three pivots are strictly positive. The separated strict bounded-Gram lemma therefore gives a finite tensor SOS of the complete RC6. It requires bounded matrix entries, not bounded rows or states.

205.1.4 All other testing charts

We give complete parent identities for the outer charts so the local certificate is not mistaken for an all-tests conclusion. Every nonnegative event packet described as retained below is kept whole, as a separated source square times testing squares.

Both original testing coordinates bounded. Put $p = uW$. Retain the whole m, n, D packets and use the remaining Gram on (p, Y, X, Z):

$$\begin{aligned} CD_v(p-Y)^2 + AD_uD_vY^2 + \tfrac{1}{2}(p+uvX-Y-vZ)^2 + (\alpha+1/2)D_u(vY-Z)^2 \\ + \tfrac{1}{2}D_v(X+uZ)^2 - \tfrac{1}{2}D_uD_vZ^2. \end{aligned} \tag{RC18}$$

The $Z = 0$ hard block is positive definite by its displayed C, A, X squares. After RC4 the determinant of the real residue is

$$\frac{(S+u^2v^2)(1+u^2)^2(1+v^2)^2(1+S-y)^2}{16y(S-y)(1-y)} > 0. \tag{RC19}$$

Here also u, v mean real testing residues. Thus RC18 is a strict bounded Gram, even when u or v has zero residue.

Alice reciprocal, Bob bounded. Let $s = 1/u$ be infinitesimal or zero, set $D_s = 1 + s^2$, and use rows $(W, Y, V_x, R_x) = (W, Y, vX, sX + Z)$. Retain the entire m, n, D, α packets. The remaining expression is $CD_v(W - sY)^2 + AD_sD_vY^2 + K/2$, where exactly

$$\begin{aligned} K = \{W - sY + D_sV_x - svR_x\}^2 + D_vR_x^2 \\ + D_s\{v^2Y^2 - 2Y(vR_x - sV_x) - (vR_x - sV_x)^2\}. \end{aligned} \tag{RC20}$$

Its residue is

$$C_0D_vW^2 + A_0D_vY^2 + \tfrac{1}{2}(W+V_x)^2 + \tfrac{1}{2}(vY - R_x)^2 \succ 0. \tag{RC21}$$

This single formula includes all bounded v, including literal zero and arbitrary competing small s, v. There is no s/v coefficient.

Bob reciprocal, Alice a bounded unit. Let w be infinitesimal or zero, and u a bounded unit. Use $\rho_* = uW - Y$, $T = wZ$, $S_D = \sqrt{D}\,Z$, and the five rows (ρ_*, Y, X, T, S_D). Retain m, n whole. The D packet is exactly $\{\sqrt{D}(uX + w\rho_*) - S_D\}^2$. The other terms are

$$CD_w\rho_*^2 + AD_uD_wY^2 + \alpha D_u(Y-T)^2 + K/2,$$

where the complete core is

$$K = (D_u + w^2)X^2 + 2uwXT + w^2\rho_*^2 + 2uw\rho_*X - 2\rho_*T - T^2 + D_u(Y-T)^2. \tag{RC22}$$

The X, S_D residue pivots are $D_u/2, 1$. On (ρ_*, Y, T) the residue matrix is

$$\begin{pmatrix} C_0 & 0 & -1/2 \\ 0 & GD_u & -(\alpha_0+1/2)D_u \\ -1/2 & -(\alpha_0+1/2)D_u & \alpha_0D_u + u^2/2 \end{pmatrix}, \qquad G = A_0 + \alpha_0 + 1/2. \tag{RC23}$$

Its first two leading minors are C_0, C_0GD_u. By RC5 its determinant is $GD_uu^2(1+2C_0)/4 > 0$. This proves strictness. The alternative u infinitesimal or zero is precisely RC7–RC17.

Both testing coordinates reciprocal. Let s, w be infinitesimal or zero. Set $T = wZ$ and $S_D = \sqrt{D}\, sZ$. Use the five rows W, X, Y, T, S_D, and retain the entire m, n, α packets. The D packet is $\{\sqrt{D}(wW + X - swY) - S_D\}^2$. The other terms are $CD_w(W - sY)^2 + AD_sD_wY^2 + K/2$, where exactly

$$\begin{aligned} K = (wW + X - swY)^2 + D_s(Y - T)^2 + s^2D_wX^2 \\ - 2sWT + 2s^2YT + 2swXT - s^2T^2. \end{aligned} \tag{RC24}$$

The full five-row residue is

$$C_0W^2 + A_0Y^2 + X^2/2 + (Y - T)^2/2 + S_D^2 \succ 0. \tag{RC25}$$

This gives the final strict bounded Gram.

Every ordered-field testing coordinate is bounded or has infinitesimal reciprocal. RC18–RC25 and the local proof therefore cover both projective lines. The identities are polynomial in the chosen homogeneous coordinates, so literal infinity is included by $s = 0$ or $w = 0$. Both signs are allowed throughout; no state ratio or state normalization is used. Positive pure testing square normalizations restore the original homogeneous parent, and the original source scale is restored by a pure source square root. Real and imaginary parts of a complex state are handled separately.

Verification and limits. The local checker verifies the six native source limits, both full normal identities, every scalar monic-division identity, all contact jets and the critical residue relation. A separate outer checker verifies RC18–RC25 as complete parent/core identities. These exact checks corroborate the algebra; the source entry, boundedness, strictness and quantified joins are the analytical proof above. This result has exactly the positive-d C2 reciprocal scope stated at the start. It does not assert the remaining positive-d cases, the weak C1/C2 terminals or the final universal theorem.

Chapter 206

Both uniform C1 source charts on the entire bounded $Z = 1$ state chart

Component `pa4f6d1e57f34`
Source: `NEW/root/uniform_c1_z_charts_rederived.tex`
SHA256: `a4f6d1e57f340a918497f38bffc46735a5d8d0be05ef6da8c8f6b37cea682186`
Inventory use: fresh complete EZ1–19 Z-state charts

206.1 Both uniform C1 source charts on the entire bounded $Z = 1$ state chart

Domain and conclusion. Use the original physical uniform C1 source with $d = 1$, $\lambda = n = 0$, $k, f, g, e = kf/g$ positive infinitesimals, and all original nonnegative event weights. Retain

$$r = 1 - af^2, \quad R = 1 + k^2 r, \quad T = 1 + k^2 - tk^2 g^2, \quad M = R - t > 0, \quad J = T - a > 0, \qquad a, t > 0. \tag{EZ1}$$

In particular a, t are bounded, r, R, T are positive units, and M, J may approach zero at arbitrary rates. For every independent testing u, v, X, Y infinitesimal and $Z = 1$, W arbitrary bounded of either sign, the entire original parent is a separated tensor SOS. This includes both bounded and unbounded $b = g/k$, and every permitted $\beta = 0$ face. It does not claim the complementary $W = 1, Z$-infinitesimal chart.

206.1.1 The exact source and a common set of bounded coefficients

Use the seven short native coquotients $c_2, c_\beta, c_3, c_5, c_4, c_\alpha, c_d$ in the retained `endpoint_source.tex`, now with $\eta = f^2, \zeta = g^2$. Their residues are $2, 2, 2, 2, 2, 2, 4$. In the present $d = 1$ normalization the exact weights are

$$C = \frac{c_3}{af^2 c_d}, \quad A = \frac{f^2 c_4}{c_d}, \quad m = \frac{k^2 c_2}{g^2 M c_d}, \quad \alpha = \frac{k^4 g^2 c_\alpha}{c_d},$$
$$D + \tfrac{1}{2} = \frac{c_5}{Jf^2 c_d}, \qquad \beta + \tfrac{1}{2} = \frac{k^2 c_\beta}{g^2 t c_d}. \tag{EZ2}$$

Every use below is of these actual weights. In particular $\beta \geq 0$ is retained, not replaced by a formal source condition.

The following two pure-source coordinate choices are exhaustive:

	s	ω	P	G	$\kappa = P/G$
b bounded	e	b^2	k^2	k^2	1
b unbounded	f	1	k^2	g^2	$(k/g)^2$

(EZ3)

In both rows $f^2 = s^2\omega$, $g^2 = \omega G$, $s, \sqrt{P}, \sqrt{G}$ are infinitesimal, and ω, κ are positive bounded. In the second row κ is infinitesimal. Here $\beta \geq 0$ gives exactly $t \leq 2\kappa c_\beta / c_d$, so $t = \kappa\tau$ with $\tau > 0$ bounded; consequently M is a unit in this row.

Define bounded source coefficients

$$c_* = \frac{c_3}{Tc_d}, \quad d_* = \frac{c_5}{Tc_d}, \quad a_* = \frac{c_4}{c_d}, \quad \gamma_* = \frac{c_\alpha}{c_d}, \qquad B_* = \omega\beta t = \frac{\kappa c_\beta}{c_d} - \frac{\omega t}{2}. \tag{EZ4}$$

The first four have residue $1/2$; $B_* \geq 0$ may vanish. Use the further table

	μ	χ	ψ
b bounded	$c_2/(Rc_d)$	t	tM
b unbounded	$\kappa c_2/(Mc_d)$	0	$0.$

(EZ5)

All entries are bounded. In the first row μ has residue $1/2$; in the second μ is infinitesimal.

206.1.2 Exact removal of the source poles

The original quarter-parent is

$$\begin{aligned} F = {} & mr_2^2 + CD_v r_3^2 + AD_u D_v Y^2 + (D + \tfrac{1}{2})r_5^2 + (\alpha + \tfrac{1}{2})D_u(vY - 1)^2 \\ & + (\beta + \tfrac{1}{2})D_v(X + u)^2 - \tfrac{1}{2}D_u D_v, \end{aligned} \tag{EZ6}$$

where $r_2 = vW - X + uvY - u$, $r_3 = uW - Y$, $r_5 = uW + uvX - Y - v$, $D_u = 1 + u^2$, $D_v = 1 + v^2$. Put

$$z = sW, \qquad q_0 = f^2 uW^2 = \omega u z^2, \qquad z_0 = f^2 uW = s\omega uz.$$

Retain the following complete nonnegative packets of the original polynomial:

$$\frac{CJ}{T}D_v(r_3 - az_0)^2, \qquad \frac{(D + 1/2)a}{T}(r_5 + Jz_0)^2, \qquad \beta D_v(X + u - tq_0)^2. \tag{EZ7}$$

For bounded b, also retain

$$\frac{mt}{R}(r_2 - Mq_0)^2. \tag{EZ8}$$

Each vanishes at the actual source contact. They are literal positive squares, even when $\beta = 0$ or a source opening is small. The first two removals use $a + J = T$ and the exact identity

$$\frac{a}{T}[(L + Jq)^2 - TJq^2] + \frac{J}{T}(L - aq)^2 = L^2.$$

The last removal uses $r_2^2 - (t/R)(r_2 - Mq_0)^2 = (M/R)(r_2^2 + 2tq_0 r_2 - tMq_0^2)$. There is no mixed denominator.

Use all three unrestricted normal coordinates

$$N_x = X + u - tq_0, \qquad N_y = (Y - ruW)/s, \qquad N_v = (v - f^2 uW)/s. \tag{EZ9}$$

Only N_x is necessarily bounded. The exact bounds needed later are $N_x, sN_y, sN_v \in \mathrm{I}$; they follow from the stated testing chart and bounded W. No boundedness of N_y, N_v is assumed. Define

$$X_* = -u + tq_0 + N_x, \qquad y_* = ruz + s^2 N_y = sY, \qquad v_* = \omega uz + N_v = v/s,$$

$$L_2 = zv_* + uv_* y_* - X_* - u, \quad L_3 = T\omega uz - N_y, \quad L_5 = -N_y + v_*(uX_* - 1).$$

After multiplying the original quarter-parent by the positive pure-source ω and removing precisely the scaled packets EZ7–EZ8, the *entire* remaining polynomial is

$$\begin{aligned} \mathcal{F} = {} & \mu(L_2^2 + 2\chi q_0 L_2 - \psi q_0^2) + c_* D_v(L_3^2 - TJ\omega^2 u^2 z^2) \\ & + \omega^2 a_* D_u D_v y_*^2 + d_*(L_5^2 - aT\omega^2 u^2 z^2) \\ & + (\omega^2 P^2 G\gamma_* + \omega/2)D_u(v_* y_* - 1)^2 + \frac{\omega}{2}D_v(X_* + u)^2 \\ & + B_* D_v\{2q_0(X_* + u) - tq_0^2\} - \frac{\omega}{2}D_u D_v, \qquad D_v = 1 + s^2 v_*^2. \end{aligned} \tag{EZ10}$$

This is a short full-parent definition of every scalar and normal coefficient used below. In particular no normal was set to zero in obtaining it. All its coefficient arrays are bounded through a, J, M, t; the unbounded original weights occur only in the retained complete squares.

206.1.3 The entire scalar is a cubic circuit with finite corrections

Set $U = u^2, V = z^2$. The polynomial obtained from EZ10 at $N_x = N_y = N_v = 0$ is exactly

$$\mathcal{F}_0 = \omega^2 \Phi(U, V), \qquad \Phi = \gamma_* P^2 G(1 + U) + UV\, H(U, V), \tag{EZ11}$$

where H has bounded finite polynomial coefficient arrays. The apparent divisions by ω^2 and UV in this definition are coefficientwise polynomial extractions: substitution in EZ10 shows those factors before any evaluation. No testing coordinate is inverted.

At the actual source contact $u^2 = P, z^2 = G$ all three normal coordinates and all retained packets vanish. The original contact value and stationary equations give

$$\Phi(P, G) = \Phi_U(P, G) = \Phi_V(P, G) = 0. \tag{EZ12}$$

The exact native coquotients in EZ2 independently verify these identities. Substituting EZ11, and cancelling nonzero pure-source factors in the source field, gives

$$H(P, G) = -\gamma_* P(1 + P), \quad H_U(P, G) = \gamma_*, \quad H_V(P, G) = \gamma_* \kappa (1 + P).$$

Write $\Delta = U - P$, $\Gamma = V - G$. Finite monic division, first in $U - P$ and then in $V - G$, therefore gives the complete scalar identity

$$\begin{aligned} \Phi &= \gamma_* \{\mathcal{M} + P\kappa U \Gamma^2\} \\ &\quad + UV\{\Delta^2 H_{20} + \Delta\Gamma H_{11} + \Gamma^2 H_{02}\}, \\ \mathcal{M} &= U^2 V + \kappa U V^2 + P^2 G - 3PUV. \end{aligned} \tag{EZ13}$$

All quotients are finite bounded coefficient polynomials; no inverse of P, G is introduced by their monic construction. The extra term $\gamma_* P\kappa U\Gamma^2$ is retained as a nonnegative square.

In the inverse-b row one further coefficient factor is needed:

$$H_{02} = \kappa \widetilde{H}_{02}, \qquad \widetilde{H}_{02} \text{ has bounded arrays.} \tag{EZ14}$$

Here is a finite source-division certificate, rather than an estimate involving independent testing rates. Write $P = \kappa G$, $t = \kappa\tau$ before substituting the native coquotients into EZ10. Every denominator is among the pure-source units M, T, c_d and their factors. At formal $\kappa = 0$,

$$P = t = \mu = B_* = 0, \quad c_* = d_* = \frac{1 + (1 - a)s^2}{2}, \quad a_* = \frac{1}{2(1 - as^2)}.$$

The formula gives $H(0, V) = 0$ identically in V. After clearing unit denominators, ordinary polynomial division by κ therefore has zero remainder. Apply the two monic $V - G$ divisions to this quotient. This proves EZ14 with bounded arrays. In the bounded-b row $\kappa = 1$, so it needs no such factor.

206.1.4 Six explicit strict roots and their conductors

The following identities apply to both rows of EZ3. Put

$$\mathsf{h} = G^{1/6}, \quad \mathsf{c} = \kappa^{1/6}, \quad \mathsf{r} = \sqrt[3]{u}, \quad \mathsf{s} = \sqrt[3]{z},$$

$$(L_1, L_2, L_3) = (\mathsf{r}^4\mathsf{s}^2,\ \mathsf{c}^2\mathsf{r}^2\mathsf{s}^4,\ \mathsf{h}^6\mathsf{c}^4), \qquad (f_1, f_2, f_3) = (\mathsf{r}^2\mathsf{s},\ \mathsf{crs}^2,\ \mathsf{h}^3\mathsf{c}^2).$$

Define $R_{2i-1} = f_i(L_1 - L_2)$ and $R_{2i} = f_i(L_1 - L_3)$ for $i = 1, 2, 3$. Then, exactly,

$$\mathcal{M} = \sum_{i=1}^{3} (R_{2i-1}^2 - R_{2i-1}R_{2i} + R_{2i}^2). \tag{EZ15}$$

Each pair has the fixed strictly positive Gram $\left(\begin{smallmatrix} 1 & -1/2 \\ -1/2 & 1 \end{smallmatrix}\right)$. Real cube roots of signed testing quantities are permitted: $z = sW$ has the separated cube root $s^{1/3}\sqrt[3]{W}$. The $L_i = f_i^2$ identity is valid for either sign of u, z. No mixed radical occurs.

For explicit finite allocation define

$$J_1 = \mathsf{r}^3 R_1 + \mathsf{c}^2\mathsf{r}\mathsf{s}^2 R_2, \qquad J_2 = -\mathsf{h}^6 R_1 + \mathsf{s}^6 R_2 - \mathsf{h}^3\mathsf{s}^3 R_5.$$

Direct multiplication gives

$$\begin{aligned} uz\Delta &= J_1, & u^2 z\Gamma &= J_2, \\ \kappa uz\Gamma &= \mathsf{c}^2\mathsf{r}\mathsf{s}^2(R_2 - R_1) - \mathsf{c}^3\mathsf{s}^3 R_3, \\ UV\Delta^2 &= J_1^2, \\ UV\Delta\Gamma &= \mathsf{s}^6 R_2^2 - \mathsf{h}^6 R_1^2 - \mathsf{h}^3\mathsf{s}^3 R_1(R_5 + R_6), \\ \kappa UV\Gamma^2 &= \mathsf{s}^6(2R_1^2 - 3R_1R_2 + R_2^2 + R_3^2) + \mathsf{h}^3\mathsf{s}^3 R_1(R_5 - R_6). \end{aligned} \tag{EZ16}$$

All coefficients on the right are infinitesimal: $\mathsf{r}, \mathsf{s}, \mathsf{h}$ are infinitesimal and c is bounded. It is harmless that $\mathsf{c} = 1$ in the first row. Equations EZ13–EZ16 therefore put the entire scalar, apart from its retained positive square, on the six rows ωR_i with a bounded Gram whose real residue is $\frac{1}{2}$ times the strict matrix in EZ15.

206.1.5 Every complete first normal and all higher terms

Define L_x, L_y, L_v as half the three derivatives of the whole polynomial EZ10 at zero normals. Expanding its short factored form gives exactly

$$L_x = \omega u z^2 T_x(U, V), \qquad L_y = \omega uz T_y(U, V), \qquad L_v = \omega uz T_v(U, V), \tag{EZ17}$$

where each T_i is a polynomial with bounded coefficient arrays. This supplies a finite construction of every coefficient: differentiate EZ10, set the three normals to zero, and extract the displayed monomial. No derivative is dropped. The exact actual-source contact equations give $T_i(P, G) = 0$. Consequently the monic divisions are

$$T_i = (U - P)A_i(U, V) + (V - G)B_i(V). \tag{EZ18}$$

For inverse b, each $B_i = \kappa\widetilde{B}_i$ with bounded arrays. Indeed the same source substitution used in EZ14 gives $T_i(0, V) = 0$ identically at $\kappa = 0$. Divide the cleared source polynomial by κ and then by the monic $V - G$. This is an exact coefficient construction; it uses neither $1/u$, $1/z$, nor an evaluated mixed quotient. For bounded b, $\kappa = 1$ again suffices.

Combining EZ16–EZ18 proves that both complete state normals and the complete testing-angle normal have infinitesimal coefficients on the strict six-row list ωR_i. The additional factor z in L_x is bounded and infinitesimal. This argument uses the full first-normal polynomial, not just its value or linear approximation at contact.

The full normal expansion of EZ10 has a bounded quadratic Gram whose real residue, in order (N_x, N_y, N_v), is

$$\begin{pmatrix} \bar{\mu} + \bar{\omega}/2 & 0 & 0 \\ 0 & 1 & 1/2 \\ 0 & 1/2 & 1/2 \end{pmatrix} \succ 0. \tag{EZ19}$$

For bounded b, $\bar{\mu} = 1/2$; for inverse b, $\omega = 1$ and $\bar{\mu} = 0$. Both give a strictly positive first pivot. The other block has pivots $1, 1/4$.

Here is also a finite justification for all higher terms, since the scaled normals N_y, N_v may be unbounded. Expanding EZ10 in the three normals gives degree at most four. A monomial of normal degree at least three containing N_x has total N_y, N_v degree at most two. Assign those factors, and any needed extra N_x factor, to two Gram rows; the remaining factor contains $N_x \in \mathrm{I}$. For a higher monomial without N_x, each of the N_y, N_v exponents is at most two, and its coefficient has a factor s^2. If its degree is three, the unassigned coefficient is a bounded multiple of $s(sN_y)$ or $s(sN_v)$. If its degree is four, it is a bounded multiple of $(sN_y)(sN_v)$ (or of one of their squares). All are infinitesimal by EZ9. These support and divisibility statements follow directly from the factored quadratics in EZ10 and are checked on the entire polynomial. Thus no unbounded scaled normal is hidden in a Gram coefficient.

The scalar, all first normals, and the complete hard block now give a finite bounded Gram with strictly positive real residue. The separated strict-Gram lemma supplies tensor SOS. Restore the whole retained packets EZ7–EZ8 and the positive square in EZ13, and divide by the pure-source positive ω. This proves the stated original-parent theorem on both source rows, without a comparison of source and testing rates.

State and boundary scope. The signs of W and of the testing angles were not restricted in EZ10–EZ19. The polynomial and cube-root identities include literal zero testing coordinates. If a bounded normalized $Z = 1$ state has X or Y a unit, the original pure-testing core has positive residue at the double-small angle corner; its positive source multiple is already a separated square, and all original event packets are nonnegative squares. Thus this proof supplies the entire Z-unit state chart. Complex states split into their real and imaginary quadratic evaluations. The two strong-current outer testing identities remain separate retained inputs to a future uniform all-tests join, as does the complementary W-unit chart.

Verification boundary. The accompanying checker verifies the full packet restorations, all source contacts and first jets in both native substitutions, the full first-normal factorizations, the finite source-factor divisions, the exact cubic conductors, and the whole higher-normal support and hard residues. The proof of boundedness, strictness and the stated source/state scope is the analytical argument above. This result does not by itself establish the global finite semidefinite-lift theorem.

Chapter 207

The inverse-b uniform C1 source: a complete local conductor

Component `pabbcd0905dcd`
Source: `NEW/weak_c2/c1_inverse_b_full_local.tex`
SHA256: `abbcd0905dcdf27b0b1e63c23704b4e5b12a9731bd3242c8a542a08e881f1263`
Inventory use: fresh complete IW1–24 inverse-b all-state local proof

207.1 The inverse-b uniform C1 source: a complete local conductor

Statement and certificate ring. Let K_c, K_t be independent real closed extensions of $\mathbb{R}$, with bounded rings B_c, B_t and infinitesimal ideals $\mathfrak{m}_c, \mathfrak{m}_t$. The certificate ring is $\mathcal{T} = K_c \otimes_{\mathbb{R}} K_t$, with bounded coefficient ring $\mathcal{B} = B_c \otimes_{\mathbb{R}} B_t$ and zero-residue ideal $\mathfrak{n} = \mathfrak{m}_c \otimes B_t + B_c \otimes \mathfrak{m}_t$. We use the finite strict-residue Gram lemma in this ring. All radicals and inverses below are pure source, pure testing, or products of separately specified such factors.

Use the actual uniform native C1 stationary source, with $\lambda = n = 0$, $d > 0$, and the original nonnegative event coefficients. Normalize $d = 1$ by a positive pure-source scale. Suppose

$$k, f, g > 0 \text{ are infinitesimal}, \qquad k/g \in \mathfrak{m}_c, \qquad r = 1 - af^2, \quad h = 1 - tg^2,$$

$$R = 1 + k^2 r, \quad T_0 = 1 + k^2 h, \quad 0 < a < T_0, \quad 0 < t < R, \quad \beta \geq 0. \tag{IW1}$$

All original strict gaps remain imposed. Then the complete original parent is tensor-SOS whenever the independent positive testing angles u, v are infinitesimal, for every real or complex testing state. The proof below works in the rotated chart $w = 1, z > 0$ with z arbitrary; in particular it includes the previously missing original W-unit, Z-infinitesimal chart. It uses no bound on g/f and no comparison between any testing and source infinitesimals. The exact sign, zero-axis and state reductions are given at the end.

207.1.1 The actual C1 source and bounded source arrays

For clarity, the six weights used here come from the following unchanged native formulas. Put $x_n = k^2$, $\eta = f^2$, $\zeta = g^2$, $y_n = x_n \eta \zeta$, $K = 1 + x_n$, $G = h + \zeta$, and

$$M = R - t, \quad J = T_0 - a, \quad E_n = 1 - x_n \eta h, \quad Q_n = 1 - x_n \zeta r, \quad \theta = t\zeta.$$

Define

$$P_d = 1 + 2x_n h + r x_n^2 G + x_n \zeta + \eta x_n (K\zeta + x_n h\theta),$$

$$P_4 = T_0 E_n + x_n(1 - r)G, \qquad P_\alpha = Rh(1 + y_n) + \zeta M E_n.$$

The actual relative coefficients, now with $d = 1$, are

$$\begin{aligned} m &= \frac{Kx_n(r+\eta)h(1+y_n)}{2M\zeta P_d}, & C &= \frac{K(r+\eta)E_n}{2(1-r)P_d}, \\ A &= \frac{\eta P_4}{2rP_d}, & D+\tfrac{1}{2} &= \frac{K(1+y_n)(r+\eta)}{2J\eta P_d}, \\ \beta+\tfrac{1}{2} &= \frac{Kx_n(r+\eta)G}{2\theta P_d}, & \alpha &= \frac{x_n^2\zeta(r+\eta)P_\alpha}{2Q_nP_d}. \end{aligned} \tag{IW2}$$

No auxiliary C2 stationary source is used. Since a, t are bounded and k, f, g are infinitesimal, r, h, P_d, R, T_0 have residue one. The inequality $\beta + \frac{1}{2} \geq \frac{1}{2}$ in (IW2) gives $t = O(k^2/g^2)$. Thus t and $p = t/R$ are infinitesimal, and $M = R(1-p)$ is a positive unit. The two independent openings a and $J = T_0 - a$ are not capped.

Introduce the exact compact coordinates

$$j = \frac{T_0}{g^2R}, \quad B = \frac{k^2g^2R}{T_0}, \quad T = \frac{f^2T_0}{R}, \quad p = \frac{t}{R}, \quad H = \frac{a}{T_0}, \qquad x = jB, \quad y = TB, \quad q_z = jT. \tag{IW3}$$

Here j is unbounded, $0 < H < 1$, and

$$T, x, B, p, \sigma := 1/j, \delta := j^2B \in \mathfrak{m}_c, \qquad \delta \sim k^2/g^2, \qquad p = O(\delta). \tag{IW4}$$

The quantity q_z can have any positive size. The exact inverse is

$$\begin{aligned} R &= \frac{1+x}{1+HTx}, \quad T_0 = \frac{1+x}{1+Bp}, \quad r = \frac{1-HT}{1+HTx}, \quad h = \frac{j-p}{j(1+Bp)}, \\ f^2 &= \frac{T(1+Bp)}{1+HTx}, \quad g^2 = \frac{1+HTx}{j(1+Bp)}, \quad a = HT_0, \quad t = pR. \end{aligned} \tag{IW5}$$

In particular both strict gaps are retained exactly.

Here is a short rational dictionary sufficient to check every source estimate. Write $D_+ = D + \frac{1}{2}$, $b_+ = \beta + \frac{1}{2}$, $\mu = m(1-p)$, and set

$$\begin{aligned} P &= 1 + B(j+1-p) + BT(1+Hj) \\ &\quad + B^2T(Hj + jp - p^2 + p) + B^2HT^2j(1+Bp), \\ L &= 1 + T(1-H+Bp), \qquad E = 1 + B(HT + p - 1), \\ F_\alpha &= j + 1 - 2p + BHTj(1-p) + BTp(j-p). \end{aligned}$$

Substitution of (IW5) in (IW2) gives exactly

$$\begin{aligned} \mu &= \frac{Bj(j-p)(1+y)(1+Bp)L}{2(1+x)P}, \\ C &= \frac{(1+Bp)L[1+BTp-BTj(1-H)]}{2HT(1+x)P}, \\ A &= \frac{T(1+Bp)[1+BT(2Hj-Hp+H-j+p)+B^2H^2T^2j]}{2(1-HT)P}, \\ D_+ &= \frac{(1+y)(1+Bp)(1+HTx)L}{2T(1-H)(1+x)P}, \\ b_+ &= \frac{Bj(1+Bp)L(j+1-p+HTx)}{2p(1+x)P}, \\ \alpha &= \frac{B^2LF_\alpha}{2EP}. \end{aligned} \tag{IW6}$$

The denominators $P, E, 1+x, 1-HT$ are positive bounded units. After $j = 1/\sigma$, $B = \delta\sigma^2$, each of the following normalized arrays is a rational function with polynomial numerator and denominator in

the bounded variables σ,δ,T,p,H; its denominator has a nonzero real constant at $\sigma=\delta=T=p=0$, independently of H:

$$\frac{2\mu}{\delta},\quad \frac{2b_+p}{\delta},\quad 2TCH,\quad 2TD_+(1-H),\quad \frac{2A}{T},\quad \frac{2\alpha}{B^2j}\quad\text{all have residue 1.}\tag{IW7}$$

This finite polynomial criterion proves boundedness and the stated residues at every relative rate and at both H openings. It also shows $\beta p\le b_+p=O(\delta)$, including $\beta=0$. A useful exact cancellation is

$$K_d:=CH-D_+(1-H)=-\frac{b_+p}{j}.\tag{IW8}$$

207.1.2 The entire parent and both state normals

Put $D_u=1+u^2$, $D_v=1+v^2$. The quarter-parent is

$$\begin{aligned}\mathcal{P}&=m(vW-X+uvY-uZ)^2+CD_v(uW-Y)^2+AD_uD_vY^2\\&\quad+D_+(uW+uvX-Y-vZ)^2+(\alpha+\tfrac12)D_u(vY-Z)^2\\&\quad+b_+D_v(X+uZ)^2-\tfrac12D_uD_vZ^2.\end{aligned}\tag{IW9}$$

On the rotated chart $w=(W+uY)/D_u=1$, $z=(Z-uX)/D_u>0$, use the two unrestricted normals

$$N=\frac{X+uZ-pD_uv}{D_u},\qquad M_0=\frac{uW-Y-HD_uvz}{D_u}.$$

Their exact inverse is

$$X=N+pv-uz,\quad Z=z+uN+puv,\quad W=1+u(M_0+Hvz),\quad Y=u-M_0-Hvz.\tag{IW10}$$

Only the pure-testing unit D_u is inverted. Retain the positive packet $mpD_u^2N^2$, and put $G=(\mathcal{P}-mpD_u^2N^2)/D_u$. Direct substitution, before any state specialization, gives

$$\begin{aligned}G&=\mu D_u[N^2-2vN+(1-p)v^2]+CD_uD_v(M_0+Hvz)^2+AD_v(u-M_0-Hvz)^2\\&\quad+D_+D_u(M_0-(1-H)vz)^2\\&\quad+(\alpha+\tfrac12)[u(1-p)v-(1+Hv^2)z-uN-vM_0]^2\\&\quad+b_+D_uD_v(N+pv)^2-\tfrac12D_v(z+uN+puv)^2.\end{aligned}\tag{IW11}$$

In particular G is exactly quadratic in the two normals:

$$G=F+2L_NN+2L_MM_0+H_NN^2+H_MM_0^2+2(\alpha+\tfrac12)uvNM_0,\tag{IW12}$$

where $F=G|_{N=M_0=0}$ and

$$\begin{aligned}h_0&=\mu+b_+\ge\tfrac12,\\H_N&=h_0D_u+b_+v^2+\beta u^2v^2+\alpha u^2,\\H_M&=D_u(CD_v+D_+)+AD_v+(\alpha+\tfrac12)v^2.\end{aligned}\tag{IW13}$$

The potentially unbounded β is fully retained in (IW11)–(IW13). There is no omitted higher-normal polynomial.

207.1.3 An exact separated scalar circuit

Define the rows

$$U=u/\sqrt{x},\quad V=v/\sqrt{y},\quad Z_*=z/\sqrt{q_z},\quad Q_*=U-VZ_*,\quad B_*=V-UZ_*,\quad R_*=Z_*-UV,$$

$$\Phi=U^2+V^2+Z_*^2+U^2V^2Z_*^2-4UVZ_*.$$

None of U, V, Z_* is assumed bounded. On the positive chart,

$$\Phi = (U-V)^2 + Z_*^2(1-UV)^2 + 2UV(1-Z_*)^2 \tag{IW14}$$

is a legal separated SOS; for example $\sqrt{UV} = (xy)^{-1/4}\sqrt{uv}$.

For an explicit finite definition of the actual masses, put $K_0 = CH^2 + D_+(1-H)^2$ and

$$\begin{aligned}
&M_a = Ax, && M_b = [\mu(1-p) + b_+p^2]y,\\
&M_c = \alpha q_z, && M_{bc} = [K_0 + AH^2 + 2H\alpha + H - \tfrac{1}{2}]yq_z,\\
&M_{b^2} = b_+p^2y^2, && M_{b^2c} = H^2(A + C + \alpha + \tfrac{1}{2})y^2q_z,\\
&M_{ab^2c} = CH^2xy^2q_z, && M_{ab^2} = \beta p^2xy^2.
\end{aligned}$$

Set

$$f_c = M_b, \quad h_c = M_{ab^2}, \quad k_c = (M_a - M_b + M_{ab^2c} - M_{bc} + h_c)/2,$$
$$r_c = M_c - M_b - k_c, \quad q_c = M_a - f_c, \quad g_c = M_{ab^2c}, \quad d_c = M_{b^2c} - k_c, \quad e_c = b_+p^2y^2 - g_c.$$

The complete scalar from (IW11) satisfies the exact identity

$$\begin{aligned}
F = {} & q_cQ_*^2 + r_cR_*^2 + d_cV^2Q_*^2 + e_cV^2B_*^2\\
& + f_c\Phi + g_cV^2\Phi + k_cZ_*^2(V^2-1)^2 + h_cV^2R_*^2.
\end{aligned} \tag{IW15}$$

The last term is the extra nonnegative square supplied by the actual β event. The checker verifies all scalar monomials of the original parent and the four remaining source coefficient identities; (IW15) is not merely an identity at the contact.

The same finite bounded-array criterion as (IW7), with the displayed mass definitions, gives

$$\frac{2f_c}{Tx^2} \sim 1, \quad \frac{2q_c}{Tx} \sim 1, \quad \frac{2r_c}{TjB^2} \sim 1, \quad \frac{k_c}{T^2B^2j} \in \mathfrak{m}_c, \quad \frac{g_c}{f_c} \in \mathfrak{m}_c. \tag{IW16}$$

Here ~ 1 means residue one. In particular the three principal masses f_c, q_c, r_c are positive, and $|k_c|/r_c$, $|k_c|/(q_cy)$ are infinitesimal. All normalized mass denominators remain units after $j = 1/\sigma$, $B = \delta\sigma^2$; the companion finite-array checks test their residues independently of H.

For the signed terms use the exact identities

$$\begin{aligned}
(1+U^2)\Phi &= R_*^2 + B_*^2 + U^2Q_*^2 + U^2(1-UVZ_*)^2,\\
(1+V^2)\Phi &= Q_*^2 + R_*^2 + V^2B_*^2 + V^2(1-UVZ_*)^2,\\
Z_*(V^2-1) &= -VQ_* - R_*.
\end{aligned} \tag{IW17}$$

The e_c, g_c combination in (IW15) is exactly

$$b_+p^2y^2V^2B_*^2 - g_c\Phi + g_c[Q_*^2 + R_*^2 + V^2(1-UVZ_*)^2].$$

A k_c of either sign costs at most $3|k_c|V^2Q_*^2 + 2|k_c|R_*^2$, including its occurrence in d_c. The corresponding relative costs are $3|k_c|v^2/(q_cy)$ and $2|k_c|/r_c$, both in $\mathfrak{n}$. Every inequality here abbreviates a finite SOS identity, using $2A^2 + 2B^2 - (A+B)^2 = (A-B)^2$ and nonnegative pure-source factors. Consequently the scalar supplies strict fixed real fractions of

$$q_cQ_*^2, \qquad r_cR_*^2, \qquad f_c\Phi. \tag{IW18}$$

All other nonnegative packets remain available as whole squares.

207.1.4 Finite budgets for the complete normal rows

Call a row free if any fixed positive real multiple of its square can be subtracted at infinitesimal cost from (IW18), by a finite separated SOS identity. This definition allows a bounded normal coupling to be absorbed by a sufficiently large fixed real helper pivot. Put

$$q = vz - Tu, \quad b = v - uz/j, \quad R_z = z - uv/B, \quad R_u = v(u^2 - x), \quad R_v = u(v^2 - y),$$

$$J_u = u(u^2 - x), \qquad J_v = v(v^2 - y), \qquad \widehat{q} = q/\sqrt{T}.$$

Since $Q_* = -q/(T\sqrt{x})$, the first mass in (IW18) has a positive unit coefficient on $\widehat{q}^2$.

First, $R_u = \sqrt{y}\,x(-B_* - UR_*)$. Applying the first identity of (IW17) to B_*^2, and the actual r_c pivot to $U^2R_*^2$, shows that R_u is free. The three relative costs are bounded by constant multiples of

$$B, \qquad u^2/j, \qquad u^2, \tag{IW19}$$

all infinitesimal. Second, $R_v = \sqrt{x}\,y(-Q_* - VR_*)$, so $\sqrt{T}R_v$ is free; its two costs are $O(y^2)$ and $O(Tv^2)$.

The same-angle rows have exact decompositions

$$\begin{aligned} J_v &= \sqrt{y}\,v(v + \sqrt{y/x}\,u)(V - U) + (y/x)R_u, \\ J_u &= \sqrt{x}\,u(u + \sqrt{x/y}\,v)(U - V) + (x/y)R_v. \end{aligned} \tag{IW20}$$

If $\rho = O(\delta)$ is a nonnegative source factor, then ρJ_v is free. Indeed $\rho\sqrt{y/f_c} = O(\sqrt{\delta})$, $\sqrt{y/x} = \sqrt{T/j}$ is bounded, and $\rho y/x$ is infinitesimal. Thus every coefficient after division by the strict $\sqrt{f_c}(V - U)$ pivot is infinitesimal. Also $\sqrt{T}\,TxJ_u$ is free: the two coefficients of its first summand relative to $\sqrt{f_c}(U - V)$ are bounded multiples of

$$T\sqrt{x}\,u^2, \qquad \sqrt{T\delta}\,uv,$$

and its second summand is exactly $\delta(\sqrt{T}R_v)$, because $Tx^2/y = j^2B = \delta$. This last use of $\delta \in \mathfrak{m}_c$ is the additional source fact specific to the inverse-b sector.

Finally,

$$j\alpha b = -(\alpha/B)R_u - \alpha u R_z. \tag{IW21}$$

Here $\alpha/B = O(x)$, and the square of $\alpha u R_z$ costs $O(x^2u^2)$ relative to $r_cR_*^2$. Hence the entire left side of (IW21) is free. These identities involve no bound on z.

The unrestricted first normal. Define the bounded source quantities

$$K_z = H(\alpha + \tfrac{1}{2}) - \tfrac{1}{2}, \qquad K_R = -\mu - \alpha(1 - p) - \tfrac{1}{2} + b_+p + TK_z,$$

$$\rho = b_+p + \beta px \geq 0.$$

Direct differentiation of the entire (IW11), followed by the actual source contact identity, gives

$$L_N = -j\alpha b + (K_R + \beta pv^2)R_u + \rho J_v + K_z uvq. \tag{IW22}$$

For completeness, before imposing the source, the remainder is vS_N, with

$$S_N = j\alpha - \mu + b_+p + K_Rx + b_+py + \beta pxy = 0$$

identically under (IW6). Thus no constant or first-normal term is silently dropped. Equations (IW7) imply $K_R \sim -\frac{1}{2}$, $\rho = O(\delta)$ and $\beta p = O(\delta)$. The first three terms in (IW22) are free by (IW19)–(IW21); the last has coefficient $K_z\sqrt{T}\,uv \in \mathfrak{n}$ on $\widehat{q}$. Dividing by $\sqrt{h_0}$ preserves all these bounds because $h_0 \geq \frac{1}{2}$.

The unrestricted second normal. Use the exact K_d from (IW8), and put

$$C_q = K_d D_u + AH + \alpha + \tfrac{1}{2} + v^2[CHD_u + AH + (\alpha + \tfrac{1}{2})H],$$
$$c_2 = T[CH + AH + (\alpha + \tfrac{1}{2})H] - A - (\alpha + \tfrac{1}{2})(1 - p).$$

The complete second normal has the exact division

$$L_M = C_q q + (c_2 + TCHu^2)R_v + T(K_d + CHy)J_u. \tag{IW23}$$

Before substituting the actual source its only remainder is uS_M, where

$$c_0 = T[K_d + AH + \alpha + \tfrac{1}{2}] - A,$$
$$S_M = c_0 + TK_d x + yc_2 + TCHxy = 0.$$

The last equality is an exact identity from (IW6). Furthermore c_2 is bounded (in fact infinitesimal), TCH is a unit, and $(K_d + CHy)/x$ is bounded with residue $-\frac{1}{2}$. It follows from the two free rows in (IW20) that

$$\sqrt{TH(1-H)}[(c_2 + TCHu^2)R_v + T(K_d + CHy)J_u]$$

is free with bounded coefficients. For the other term,

$$\sqrt{TH(1-H)}C_q q = T\sqrt{H(1-H)}C_q \widehat{q}.$$

Its coefficient is in $\mathfrak{n}$: multiplying C_q by T clears the only CH pole, and every term then contains either T or v^2. This statement is made on the full expression in (IW23), before any relative source/testing rate is considered.

207.1.5 Strict full Gram and state restoration

Normalize the hard rows by pure-source factors,

$$\widehat{N} = \sqrt{h_0}\, N, \qquad M_0 = \sqrt{TH(1-H)}\, \widehat{M}.$$

Their full hard matrix is bounded: $b_+/h_0 \le 1$, $\beta/h_0 \le 1$, $\alpha/h_0 = O(\alpha)$, and the two source poles of H_M are exactly cleared by $TH(1-H)$. Equations (IW7), (IW13) give its real residue

$$\operatorname{diag}(1, \tfrac{1}{2}). \tag{IW24}$$

Its off-diagonal coefficient has the factor $uv\sqrt{TH(1-H)}/\sqrt{h_0}$ and is infinitesimal. The $\widehat{q}$ couplings in (IW22)–(IW23) are infinitesimal. All remaining complete-normal couplings are bounded coefficients of free rows. Buy sufficiently large fixed real pivots on the finitely many free rows, use the elementary real Young square completion, and apply the strict-residue Gram lemma to the remaining finite matrix. The strict scalar pivots (IW18) lose only infinitesimal amounts. This proves $G \in \Sigma\mathcal{T}^2$. No state normal has been set to zero in this conclusion. Restoring $mpD_u^2N^2$, the pure-testing factor D_u, and the positive source scale d proves the assertion for (IW9).

The proof has used $u, v > 0$ and $w = 1, z > 0$, with arbitrary z. The exact sign/state lemma SJ1–SJ10 applies to the same original core: sign gauges make u, v nonnegative while preserving every packet; a negative core at positive angles forces the rotated w, z to have the same nonzero sign. Dividing all state coordinates by the nonzero pure-testing w gives precisely this chart. When the core is nonnegative it is itself a pure-testing square, and every event packet is already positive. Zero angles have the direct core SOS exits. The zero state is immediate, and a complex state splits into its real and imaginary quadratic evaluations. Thus (IW1) is proved for all states at the double-infinitesimal angles. The existing two-current outer-angle identities give the separate all-tests join if needed; they are not required as a premise of the local conductor proved here.

Verification and scope. The new module `iw_actual_source.py` prints the finite actual C1 source dictionary. The independent-formula checks in `check_c1_inverse_b_local.py` compare it with (IW2), verify the entire original parent and both normals, all scalar monomials and the exact circuit, the elementary helper identities, and the finite normalized source arrays. The ordered-field estimates and finite Gram allocation are the proof above; a checker receipt alone does not establish them. No lost historical IW coefficient witness, R21 admission label, formal negative-β source, mixed inverse, or additional source-rate restriction is used.

Chapter 208

A new complete R1 theorem with no source-ratio split

Component `pbaab204a8ab1`
Source: `NEW/weak_c2/r1_all_sources_all_tests.tex`
SHA256: `baab204a8ab1db9336f65126c06bd3c57110f45d4ea11df2ab591f61d25a4ac1`
Inventory use: R1N1-R1N24 and all-testing restoration | Only R1N12-24 universal full-current/finite-cap algebra is reused; no original R1 source membership or local theorem is imported

208.1 A new complete R1 theorem with no source-ratio split

Statement and exact source. Use the original common-II source, with $d = 1$, $\lambda = \beta = 0$, the six native coefficients of the recovered `weak_c2_source_formulas.tex`, every original strict gap, and $n \geq 0$. In the original R1 sector,

$$t_s, \quad B_s, \quad a_s B_s, \quad t_s B_s > 0 \quad \text{are infinitesimal.} \tag{R1N1}$$

There is no bound on a_s, on any original positive opening, or on the source ratio previously called q_{old}. The conclusion is: the entire original parent is a finite separated tensor sum of squares at every independent real projective measurement pair and every real or complex state. This proof treats R1 only.

Source and testing quantities belong to separate real closed fields $K_c, K_t \supset \mathbb{R}$. All SOS assertions are in $K_c \otimes_{\mathbb{R}} K_t$. We use the actual recovered strict bounded-Gram lemma: a bounded finite Gram with positive definite real residue gives tensor SOS, with no boundedness requirement on its row entries. No order or mixed inverse is imposed on the tensor ring.

Put $c_s = 1 + a_s B_s r$, $R_s = 1 + h a_s B_s$, and define

$$\begin{aligned} j &= \frac{a_s c_s}{R_s}, \quad B = \frac{B_s R_s}{c_s}, \quad T = \frac{t_s c_s}{R_s}, \quad p = \frac{a_s(1-r)}{R_s}, \\ H &= \frac{1-h}{t_s c_s}, \qquad x = jB, \quad y = TB, \quad q_z = jT. \end{aligned} \tag{R1N2}$$

The native strict gaps give $0 < p < \min(1, j)$ and $0 < H < 1$. They also give $1 - h < t_s c_s$, so $h \sim 1$. Consequently B, T, x, y are infinitesimal, while q_z *is an arbitrary positive source quantity.* In fact $x = a_s B_s$ and $y = t_s B_s$ exactly. The inverse source substitution is

$$r = \frac{j-p}{j+px}, \quad h = \frac{1-HT}{1+HTx}, \quad a_s = \frac{j+px}{1+HTx}, \quad B_s = \frac{B(1+HTx)}{1+pB}, \quad t_s = \frac{T(1+pB)}{1+HTx}. \tag{R1N3}$$

All arguments below therefore concern the actual original source.

208.1.1 Source bounds without a bound on jT

The exact dictionary RSR4–RSR6 in the freshly reconstructed actual-source and scalar proof remains valid under (R1N2)–(R1N3), because it is an exact rational substitution. For completeness the two polynomials controlling its denominators and current masses are

$$\begin{aligned}\mathcal{P} &= j+1+B\{(j+1)^2-p\}\\ &\quad +TB\{Hj+j+1+B[Hj(2j+2-p)+p(2j+1-p)]\}\\ &\quad +HjT^2B^2(1+Bp+BHj),\\ \mathcal{L} &= K+T\{(1-H)j+1-p-H\}\\ &\quad +BT[pK+Hj(1-p)]+BHT^2\{(1-H)j-p\}, \qquad K=j+1-p.\end{aligned}$$

The estimates

$$\mathcal{P}\sim j+1, \qquad \mathcal{L}\sim K, \qquad \frac{1}{2}<k:=\frac{K}{j+1}<1 \tag{R1N4}$$

use only $T, B, Bj = x$ infinitesimal and $p, H < 1$. Indeed dividing each correction in $\mathcal{P}$ by $j+1$ bounds it by a finite sum of $B+x$, $TB(1+B+x)$ and $T^2B^2(1+B+x)$; dividing $\mathcal{L}-K$ by K bounds it by a finite sum of T, TB and BT^2. In particular no product jT is bounded in this argument.

The exact nonnegative numerator of n has the regrouping

$$\mathcal{N}=p\{1+B(3j+1-p)+BT(j+1)\}-Bj(j+1)\{1+(1-H)T\}+\mathcal{R}, \quad |\mathcal{R}|\le C_0BT\,Bj(j+1). \tag{R1N5}$$

This is the finite polynomial regrouping RSR8; its coefficient bound uses $p<\min(1,j)$ only. Since $n=\mathcal{N}/[2(j-p)\mathcal{P}]\ge 0$, it follows that

$$\eta_*:=\frac{Bj(j+1)}{p}=O(1), \qquad \eta_*\in \mathrm{I}\Longrightarrow n\sim\frac{p}{2(j-p)(j+1)}>0. \tag{R1N6}$$

Thus literal $n=0$ is retained. The same dictionary gives, with $\mu=m(1-p)$,

$$\begin{aligned}&\mu=O(p), \quad \mu+\tfrac{1}{2}\asymp 1, \quad \alpha\asymp B^2(j+1), \quad \frac{n(j-p)}{j}=O(1),\\ &\frac{A}{T}\sim\frac{k}{2}, \quad TCH\sim\frac{k}{2}, \quad TD(1-H)\sim\frac{k}{2}.\end{aligned} \tag{R1N7}$$

All statements are uniform at the independent openings H, $1-H$, $j-p$, and $1-p$.

Put

$$g_*=\frac{CD}{C+D}, \qquad h_*=\frac{D}{C+D}.$$

Writing $U_c=TCH$ and $U_D=TD(1-H)$ gives the exact source identity

$$Tg_*=\frac{U_cU_D}{(1-H)U_c+HU_D}\sim\frac{k}{2}. \tag{R1N8}$$

The denominator is a bounded positive unit even when either opening is infinitesimal. In particular $1/C, 1/D, 1/g_*$ are infinitesimal and $Ag_*\sim k^2/4$. Also $(C+D)^{-1/2}=O(\sqrt{T})$.

208.1.2 The complete small-angle theorem extends to all R1 sources

The reconstructed proof `fn_reconstructed.tex`, RFN4–RFN24, is a full-parent proof, not just a restriction to a scalar graph. We now justify its application under the larger source domain (R1N1), rather than importing its former domain label.

First, all of its exact identities hold for the rational dictionary (R1N3), without any size assumption on q_z. The parent, its two normals, their quadratic block, the scalar circuit, the radial Gram and

the three source cancellation identities were all proved as identities of this dictionary. Second, every uniform source bound used in that proof is exactly (R1N4)–(R1N7) or the following scalar bound:

$$f_c \asymp Tx^2, \quad q_c \asymp Tx, \quad r_c \asymp TjB^2(1-p), \quad g_c/f_c = O(Hy), \quad |k_c|/(q_c y) = O(p+B). \tag{R1N9}$$

To check the latter bounds without $jT \in \mathrm{I}$, use the explicit finite polynomials RSR15 and RSR18: each term of $\mathcal{F}_1/K, \mathcal{F}_2/K$ is bounded using $p/j < 1$ and $Bj = x$, and $\mathcal{K}/K = O(p+B)$ by the same inequalities. Thus $\mathcal{F} \sim K$ and all statements (R1N9) follow from RSR16–RSR20. No jT factor occurs in these estimates.

The signed scalar allocation retains the effective mass $M_e = nh_0 j^2/(n+h_0)$, $h_0 = \mu + 1/2$, after the complete correlated n-square completion. Its three source cases use $1-p$ a unit, or $p \sim 1$ with η_* a unit, or $p \sim 1$ with η_* infinitesimal. The proofs of these cases use only (R1N4)–(R1N9). In the last case M_e is a unit; in the second, $j^2 B$ is a unit. Literal $n = 0$ occurs only in the first two cases.

For clarity, the following are all potentially nonuniform costs in the complete normal lift, after cancellation of the source factors. The symbols u, v denote independent infinitesimal tests.

purchased row or source case	bounded diagonal costs
$v(u^2-x)$, $1-p$ unit	$O(B)$, $O(Bu^2)$, $O(u^2/(1-p))$
$v(u^2-x)$, $p \sim 1$, η_* unit	$O(B + u^2/j + u^4/(j^2 B))$, $O(Bu^2)$
$v(u^2-x)$, $p \sim 1$, $\eta_* \in \mathrm{I}$	$O(u^2/j)$, $O(u^4/j)$, $[8u^4 + 2(u^2-x)^2]/M_e$
$pv^2(v - uz/j)$, $\eta_* \in \mathrm{I}$	$(p^2/M_e)v^4$
$pv^2(v - uz/j)$, η_* unit	$(p^2 y/f_c)(yv^2 + v^4)$

(R1N10)

Every cost is infinitesimal for its stated case. The source quotients p^2/M_e or $p^2 y/f_c$ used in the last two lines are bounded by (R1N6) and $p < \min(1, j)$.

The full opposite normal row has squared multiplier

$$a_R = Tq_z v^2(\alpha - BTCHu^2)^2, \qquad J = B(j+1) + u^2.$$

The occurrence of the possibly unbounded q_z cancels exactly in its three cost ratios:

$$a_R/f_c = O((T/j)v^2 J^2), \qquad a_R/(q_c y) = O(v^2 J^2), \qquad a_R/(M_e yx) = O\Big(T(1+1/n)v^2 J^2/j^2\Big). \tag{R1N11}$$

The last ratio is used only for $p \sim 1, \eta_* \in \mathrm{I}$, where $(1+1/n)/j^2$ is bounded. In the other two cases the direct opposite-current or the $(1+U^2)\Psi$ budget applies exactly as in RFN24. The remaining normal coefficients are bounded multiples of $uv\sqrt{T}$, $T + v^2$ or the rows in (R1N10); their estimates require only B, T, x, y infinitesimal. This checks all complete first-normal and hard-normal coefficients. The variable z remains inside retained rows, with no bound on it.

Consequently RFN4–RFN24 proves the following enlarged local statement: for every original R1 source (R1N1), the entire parent is tensor SOS when $u, v > 0$ are infinitesimal, $w = (W + uY)/(1 + u^2) = 1$, and $z = (Z - uX)/(1+u^2) > 0$ is arbitrary. The complete source-only n-completion includes $n = 0$. The freshly proved SJ1–SJ10 state/sign reduction therefore gives every state for these small angles, as well as every literal zero-angle and projective endpoint.

208.1.3 Outer testing angles: two exact current identities

We include the outer mechanism and its precise positivity condition. Let

$$R = uW - Y, \quad C_t = v(uX - Z), \quad B_t = X + uZ, \quad N = R + h_* C_t, \quad D_u = 1 + u^2, \quad D_v = 1 + v^2,$$

and let K be the signed quarter-core

$$K = \tfrac{1}{2}(R + C_t)^2 + \tfrac{1}{2} D_u (vY - Z)^2 + \tfrac{1}{2} D_v B_t^2 - \tfrac{1}{2} D_u D_v Z^2. \tag{R1N12}$$

All unselected original nonnegative events are kept as whole squares.

If $1/v$ is bounded, use

$$K_0 = 2\sqrt{g_*}C_t, \quad B_0 = \sqrt{2D_v/D_u}\,B_t, \quad c_u = D_u^{-1/2}, \quad t_u = uD_u^{-1/2}, \quad \eta_v = \sqrt{D_v}/v.$$

The exact full selected-parent identity is

$$\begin{aligned} 4CD_vR^2 + 4D(R+C_t)^2 + 4K &= 4(C+D)N^2 + 4Cv^2R^2 + 2(R+C_t)^2 \\ &\quad + 2D_u(vY-Z)^2 + (K_0,B_0)Q(K_0,B_0)^T, \\ Q &= \begin{pmatrix} 1 - c_u^2\eta_v^2/(2g_*) & t_u\eta_v/\sqrt{2g_*} \\ t_u\eta_v/\sqrt{2g_*} & 1 \end{pmatrix}. \end{aligned} \tag{R1N13}$$

Since $1/g_*$ is infinitesimal and all the testing factors are bounded, $\overline{Q} = I$. This treats every other angle and every state, including Bob infinity by homogeneous clearing.

For v infinitesimal put $\epsilon = 1/g_*$, $q = \sqrt{\epsilon}$, $\gamma = Ag_*$, and

$$K_1 = 2\sqrt{g_*}C_t + \sqrt{\epsilon}Y, \quad B_1 = B_t - uvY, \quad V_0 = vY, \quad Y_1 = \sqrt{\epsilon D_u}\,Y,$$

$$L_0 = \tfrac{1}{2}qc_u(K_1 - c_uY_1) - t_uvB_1 + c_uvV_0.$$

Then $L_0 = \sqrt{D_u}\,v(vY - Z)$ exactly, and

$$\begin{aligned} &4CD_vR^2 + 4D(R+C_t)^2 + 4AD_uD_vY^2 + 4K \\ &\quad = 4(C+D)N^2 + 4Cv^2R^2 + 2(R+C_t)^2 + \mathcal{G}, \\ \mathcal{G} &= K_1^2 + 2D_v(B_1^2 + V_0^2) + (4\gamma D_v - c_u^2)Y_1^2 \\ &\quad + 2qv(K_1 - c_uY_1)V_0 - 2L_0^2. \end{aligned} \tag{R1N14}$$

The complete four-root Gram has residue

$$\operatorname{diag}(1, 2, 2, \overline{k}^{\,2} - \overline{c_u}^{\,2}). \tag{R1N15}$$

Thus it is strict whenever the last entry is positive. In particular it covers every unbounded u, and covers every noninfinitesimal u whenever $\overline{k} = 1$. All coefficients in the identity are bounded functions of the displayed normalized homogeneous Alice coordinates, so it includes the literal Alice-infinite endpoint.

208.1.4 The remaining bounded nonzero Alice chart

The only outer chart left by (R1N13)–(R1N15) has v infinitesimal, u a bounded positive unit, and $\overline{k} < 1$. Since $1 - k = p/(j+1)$, this implies that p, j are positive units and j is bounded. Therefore $\eta_* \in \mathrm{I}$ and (R1N6) gives the exact relative critical limits

$$m \sim \frac{jp}{2(1-p)(j+1)}, \qquad n \sim \frac{p}{2(j-p)(j+1)}. \tag{R1N16}$$

These statements allow either or both coefficients to be unbounded.

Choose bounded positive real caps $m_c \le m$, $n_c \le n$ as specified below and keep their nonnegative excesses. Retain all of α as its original square. Define six rows

$$\widehat{N} = \sqrt{C+D}(R + h_*C_t), \quad P = C_t/\sqrt{T}, \quad Y_0 = \sqrt{T}\,Y, \quad X, \quad Z, \quad V = vY. \tag{R1N17}$$

They need not be independent or bounded. Put $\delta = (C+D)^{-1/2} = O(\sqrt{T})$, so $R = \delta\widehat{N} - h_*\sqrt{T}\,P$. The selected *entire* parent is exactly

$$\begin{aligned} \mathcal{P}_{\mathrm{sel}} &= \widehat{N}^2 + Cv^2R^2 + \tfrac{1}{2}(R+C_t)^2 + (Tg_*)P^2 + (A/T)D_uD_vY_0^2 \\ &\quad + m_c\{(v/u)R + (D_u/u)V - X - uZ\}^2 + n_cD_uD_vX^2 \end{aligned}$$

$$+\tfrac{1}{2}D_u(V-Z)^2+\tfrac{1}{2}D_v(X+uZ)^2-\tfrac{1}{2}D_uD_vZ^2. \tag{R1N18}$$

This is obtained by the exact completion $CD_vR^2+D(R+C_t)^2=(C+D)N^2+g_*C_t^2+Cv^2R^2$. It keeps every state-normal term.

The row relation

$$PY_0=V(uX-Z) \tag{R1N19}$$

allows an arbitrary real zero shift $2b(PY_0-V(uX-Z))$. Keep the two explicitly nonnegative squares Cv^2R^2 and $(R+C_t)^2/2$ whole. The remaining complete Gram on (R1N17) is bounded: $u, 1/u$ are bounded, $Tg_*, A/T$ are units, and the R contribution in the m_c square has coefficients $\delta v/u$ and $h_*\sqrt{T}v/u$, both infinitesimal. Its residue is the direct sum of a unit pivot and the two blocks

$$\begin{pmatrix} k_0/2 & b \\ b & k_0(1+u_0^2)/2 \end{pmatrix}, \qquad Q_3(m_c,n_c;b), \tag{R1N20}$$

where subscripts denote real residues and Q_3 is the quadratic form in (X,Z,V)

$$\begin{aligned} Q_3 &= m_c\{(D_u/u)V-X-uZ\}^2+n_cD_uX^2 \\ &\quad+\tfrac{1}{2}D_uV^2-D_uVZ+\tfrac{1}{2}(X+uZ)^2-2bV(uX-Z). \end{aligned} \tag{R1N21}$$

In (R1N21), $u=u_0>0$ is real. The choices of b, m_c, n_c that make both blocks strict are now explicit.

Interior openings. Suppose $p_0<1$ and $j_0>p_0$. Use initially $m_0=j_0p_0/[2(1-p_0)(j_0+1)]$ and $n_0=p_0/[2(j_0-p_0)(j_0+1)]$, and take $b=k_0/2$. The first leading principal minor of Q_3 is $m_0+n_0D_u+1/2>0$; the other two are

$$\frac{p_0u_0^2D_uK_0}{4(j_0-p_0)(j_0+1)^2(1-p_0)}, \qquad \frac{p_0u_0^2D_uK_0^2}{8(j_0-p_0)(j_0+1)^3}, \tag{R1N22}$$

where $K_0=j_0+1-p_0$. They are positive. The first block in (R1N20) has determinant $k_0^2u_0^2/4>0$. Positive definiteness is open, so fixed real caps slightly below m_0, n_0 preserve strictness and are smaller than the actual source coefficients.

Only the n opening is infinite. If $j_0=p_0<1$, then n is unbounded and $m_0=j_0^2/[2(1-j_0)(j_0+1)]$ is finite positive. At $b=k_0/2=1/[2(j_0+1)]$, restrict (R1N21) to the kernel $X=0$ of the n packet. Its first diagonal on (Z,V) is $u_0^2(m_0+1/2)>0$, and its determinant is $u_0^2k_0^2/4>0$. Choose $m_c<m_0$ sufficiently close and then a sufficiently large fixed real n_c. The whole block is strict, with both caps below the original weights.

Only the m opening is infinite. If $p_0=1<j_0$, then m is unbounded and $n_0=1/[2(j_0^2-1)]$ is finite positive. Take a real b just above $k_0/2=j_0/[2(j_0+1)]$, with

$$b<\min\{k_0\sqrt{D_u}/2,\ j_0/[2(j_0-1)]\}.$$

This interval is nonempty. The first block in (R1N20) is strict. On the kernel $(D_u/u)V-X-uZ=0$ of the m packet, use (X,V) coordinates. The first diagonal of (R1N21) is $D_u/[2(j_0^2-1)]>0$, and its determinant is

$$-\frac{D_u^2[\,2b(j_0-1)-j_0\,][\,2b(j_0+1)-j_0\,]}{4u_0^2(j_0^2-1)}>0. \tag{R1N23}$$

Choose $n_c<n_0$ sufficiently close and then a sufficiently large real m_c. This gives the required full strict block.

Both openings are infinite. If $p_0 = j_0 = 1$, both m, n are unbounded and $k_0 = 1/2$. Choose $1/4 < b < \sqrt{D_u}/4$. The common kernel of the two event packets is $X = 0$, $Z = D_u V/u_0^2$. On it the rest of (R1N21) is exactly

$$\frac{D_u(4b-1)}{2u_0^2}V^2 > 0. \tag{R1N24}$$

A sufficiently large real cap on both packets therefore gives a strict block.

The elementary finite-cap fact used in these cases is as follows: if a real positive semidefinite event Gram is positive definite on the complement of its kernel, and the remaining real quadratic form is positive definite on that kernel, a sufficiently large real multiple of the event Gram makes their sum positive definite. It follows by a block Schur complement, and works for the sum of two event Grams as well. Thus the choices above prove strictness of the *complete* bounded Gram (R1N18), without restricting any normal or state coordinate.

208.1.5 Exhaustive testing and state join

The sign gauges SJ3 preserve every original source packet and the whole core, so affine u, v may be taken nonnegative. If v is not infinitesimal, (R1N13) applies. If v is infinitesimal and u is not infinitesimal, either (R1N15) applies or the complete six-root argument (R1N16)–(R1N24) applies. These cases exhaust the outer angles. If $u, v > 0$ are both infinitesimal, the enlarged local FN theorem proved above applies after SJ's full-state reduction to $w = 1, z > 0$; SJ gives the other state signs and the axes directly. Literal homogeneous measurement endpoints are given by SJ10 or the homogeneous current identities, without a limiting argument. The zero state is immediate. For a complex state, the real parent matrix splits its value into its real and imaginary quadratic evaluations, to which the same proof applies.

Restore every capped excess, the original n, m, α packets not used in a particular chart, and all cleared positive pure testing homogeneous factors. All normalizations, radicals and inverses are pure source or pure testing. This proves the complete original R1 source row at all independent tests and states, without a source-ratio split or a comparison of source and testing rates.

Verification scope. The companion checker verifies the exact source-coordinate map, the harmonic source identity, both full outer-current identities, the new full six-root parent representation, the zero relation, all positive-minor and kernel formulas, and their original source coefficients. The source bounds and exhaustive chart argument above are analytical. This theorem uses actual complete source and FN proof texts, not a historical status receipt. It makes no assertion about the still-separate R2–R4 rows.

Chapter 209

The entire original R2 source, with all independent tests and states

Component `p913a09bcdcda`
Source: `NEW/weak_c2/r2_all_original_sources_all_tests.tex`
SHA256: `913a09bcdcda2a6cc79cc099b32c4bb05596180f80eef27fe2e49f001419aa9e`
Inventory use: R2A1-R2A4

209.1 The entire original R2 source, with all independent tests and states

Theorem and unchanged source. Let K_c, K_t be independent real closed extensions of $\mathbb{R}$. Use the original common-II stationary source with $d = 1$, $\lambda = \beta = 0$, the six actual rational coefficients n, m, C, A, D, α in RER1, and all original source gaps and coefficient nonnegativity conditions. In particular $n \geq 0$; the strict m, D denominator gaps are

$$g_s = 1 + a(hB + r - 1) > 0, \qquad j_D = 1 + aBr - (1 - h)/t > 0.$$

Assume the original R2 size conditions

$$a, B, t > 0, \quad 0 < r, h < 1, \qquad t \text{ unbounded}, \qquad B, aB, tB \in \mathrm{I}_c. \tag{R2A1}$$

Then the complete original parent is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$ at every independent projective testing pair and every real or complex state. All allowed source openings and literal $n = 0$ are included.

For specificity, in finite testing half-angle coordinates the quarter-parent being certified is exactly

$$\begin{aligned} \mathcal{P} = {} & m(vW - X + uvY - uZ)^2 + nD_uD_vX^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 \\ & + (D + \tfrac{1}{2})(uW + uvX - Y - vZ)^2 + (\alpha + \tfrac{1}{2})D_u(vY - Z)^2 \\ & + \tfrac{1}{2}D_v(X + uZ)^2 - \tfrac{1}{2}D_uD_vZ^2, \qquad D_u = 1 + u^2, \quad D_v = 1 + v^2. \end{aligned} \tag{R2A2}$$

The statement is about this entire quadratic in the four state coordinates. It does not specialize either state normal to zero and does not use a scalar-only positivity assertion.

209.1.1 Exhaustion of the original source

The exact source-entry lemma RER1–RER8 derives its bounds from the original inequality

$$(1 - r)Q \geq BK,$$

including equality when $n = 0$. In particular it proves $a(tB) \in \mathrm{I}_c$ before any comparison source is used. No earlier-source exclusion is added to (R2A1).

Every positive source a lies in precisely one of the following two size classes.

Bounded a. Apply the whole-parent theorem RL1–RL24, with the exact circuit and complete-normal identities SH1–SH8. Its first source alternative is exactly bounded native a, with no restriction on $e = 1 - r$, n/e^2, or their ratios. Its correlated n-completion is literal at $n = 0$. The proof retains both complete normals and every positive rotated state ratio.

Unbounded a. Set

$$S = 1/t, \quad z = 1/a, \quad q = a(tB), \quad w = a(1 - r), \quad \Lambda = \frac{n}{tB} = \frac{n}{qz}. \tag{R2A3}$$

These are the exact original-source coordinates TU2. The additional nonnegative source quantity Λ is either bounded, including zero, or positive unbounded. If it is bounded, apply TU1–TU23. If it is unbounded, apply UL1–UL18. The formal C1 coefficients in those two proofs are only a finite algebraic comparison. Neither proof assumes their physicality or imports a C1 theorem at a negative event coefficient. The entire actual n-packet and both complete normal corrections are retained.

These alternatives exhaust all sources in (R2A1). There is no intermediate size class between bounded and positive unbounded in a real closed extension of $\mathbb{R}$. In particular $n = 0$ belongs to the bounded-Λ alternative whenever a is unbounded; it is not removed by a division by n.

209.1.2 Exhaustion of tests and states

The fresh exact sign and projective lemma SJ1–SJ10 permits nonnegative finite testing half-angles by real state gauges that preserve every event packet. Literal zero angles and homogeneous infinite endpoints have direct core SOS identities. On the remaining positive finite angles, R2O1–R2O24 proves the entire original parent whenever at least one angle is not infinitesimal. Its physical reflection, all four finite-cap opening cases, and the original positive excesses are part of that proof; no reflected R1 source hypothesis is imported.

It remains only to consider $u, v > 0$ both infinitesimal. The source alternatives above supply a complete local certificate for every such pair and every state. RL uses the nonzero pure-testing coordinate $w = (W + uY)/D_u$ to normalize $w = 1$, and permits the resulting $z > 0$ to be arbitrary. TU and UL use the pure-testing $Z = 1$ chart, with no bound on the other state ratio. Their respective original-core sign identities justify these normalizations and give direct SOS on the complementary state charts. Thus no state chart is lost when the two exact frames are used in different source sectors.

Every normalization is reversed by multiplying the roots by a positive pure-source or pure-testing factor. The zero state is immediate. Since the measurement matrices are real, the quadratic parent on a complex state is the sum of its values on the real and imaginary parts. This proves the stated real and complex all-state conclusion.

209.1.3 Exact proof dependencies and scope

The substantive new mathematical leaves are:

source or testing region	complete proof
$a = O(1)$	RL1−RL24, SH1−SH8
a unbounded, $\Lambda = O(1)$	TU1−TU23
a, Λ unbounded	UL1−UL18
outer testing angles and all endpoints	R2O1−R2O24, SJ1−SJ10.

(R2A4)

They use the actual source-entry identities RER1–RER8, the reconstructed exact compact source and scalar identities RSR, the complete parent RFN4–RFN16, and the printed finite QI/SC circuit and contact-division lemmas where explicitly invoked. These are proof bodies and exact finite identities; historical status files and assertion counts are not premises.

This theorem completes the whole original weak-C2 row R2. It does not by itself assert the global universal tensor-SOS theorem, produce an explicit finite LMI, bound an LMI size, or assert external proof verification. Those are separate assembly and deliverable questions.

Chapter 210

R2 with bounded a, or an infinitesimal paired-event parameter

Component `p393d3faa269a`
Source: `NEW/weak_c2/r2_bounded_a_or_small_p_local.tex`
SHA256: `393d3faa269a80f3be895a951ff29f03eb13c1ceb1e122cc5123d1978eb0a2ee`
Inventory use: RL1-RL24

210.1 R2 with bounded a, or an infinitesimal paired-event parameter

Exact statement. Use the actual original weak-C2 source and all its original strict gaps, with $d = 1$, $\lambda = \beta = 0$, and $n \geq 0$. Suppose t_s is positive unbounded and B_s, a_sB_s, t_sB_s are positive infinitesimals. Put $e = 1 - r$, $R_s = 1 + ha_sB_s$. On the source subset

$$a_s = O(1) \quad \text{or} \quad p = a_se/R_s \in \mathrm{I}_c, \tag{RL1}$$

the entire parent is separated tensor SOS for every independent projective testing pair and every real or complex state. Literal $n = 0$ and every opening of $r, h, 1 - h, g$ are retained. For the all-testing conclusion, the local proof below is joined explicitly to R2O1–R2O24.

The two fields and the bounded-Gram convention are those of EN1–EN4. In particular no mixed inverse is used. Every coefficient bound below refers to a finite array in $B_c \otimes B_t$; the rows need not be bounded.

210.1.1 Actual source entry and bounds

Use the exact compact source dictionary RSR2–RSR8, with

$$j = a_sc_s/R_s, \quad B = B_sR_s/c_s, \quad T = t_sc_s/R_s, \quad p = a_se/R_s, \quad H = (1-h)/(t_sc_s), \quad c_s = 1 + a_sB_sr.$$

Set

$$S = 1/T, \quad \ell = HT, \quad \bar{h} = 1 - \ell, \quad x = jB, \quad y = TB, \quad B = Sy, \quad q_z = j/S. \tag{RL2}$$

Thus S, x, y are positive infinitesimals, $0 < p < \min(1, j)$, $0 < \ell < 1$, and $j \asymp a_s$. Both ℓ and $\bar{h}$ may be arbitrarily small. The original physicality inequality RER2–RER7 gives $jy \in \mathrm{I}_c$. In terms of the actual finite polynomials $\mathcal{P}, \mathcal{L}, \mathcal{N}, \mathcal{F}, \mathcal{K}$ of RSR, put

$$K = j + 1 - p, \quad k = K/(j+1), \quad F_b = j + 1 + p(1-p)/j, \quad E = 1 + B(\ell + p - 1). \tag{RL3}$$

Here $1/2 < k < 1$ and $1 \leq F_b/(j+1) < 2$. Direct substitution $B = Sy, T = 1/S, H = S\ell$ in those finite polynomials gives

$$\mathcal{P}/(j+1) \sim 1, \qquad S\mathcal{L}/K \sim 1, \qquad S\mathcal{F}/F_b \sim 1. \tag{RL4}$$

For clarity, the terms after division contain only bounded $p/j, j/(j+1), K/(j+1)$, multiplied by S, y, or jy. No inverse of ℓ, $\bar{h}$, or $j-p$ is involved.

The same substitution in the exact numerator of n yields

$$\mathcal{N} = p(1+\epsilon_p) - yj(j+1)(1+\epsilon_y), \qquad \epsilon_p, \epsilon_y \in \mathrm{I}_c. \tag{RL5}$$

For an explicit finite division certificate, the zero-p term is

$$\mathcal{N}|_{p=0} = -yj(j+1)\{1+S(1-\ell)\}\{1+S\ell yj/(j+1)\}.$$

The quotient $(\mathcal{N} - \mathcal{N}|_{p=0})/p$ is a finite polynomial. This grouping can be read termwise from RSR5: the first coefficient is $1 + y(j+1) + O(Sy(j+1) + Sy^2(j+1))$; the second coefficient is $1 + O(S + y + Syj)$. Using $\mathcal{N} \geq 0$, therefore,

$$\eta_2 = \frac{yj(j+1)}{p} = O(1), \qquad \eta_2 \in \mathrm{I}_c \implies n \sim \frac{p}{2(j-p)(j+1)} > 0. \tag{RL6}$$

The estimate includes the face $\mathcal{N} = 0$, which can occur only in the positive-unit η_2 case.

Let $\mu = m(1-p)$ and $h_0 = \mu + 1/2$. The complete source formulas now imply

$$\begin{gathered} \mu \sim \frac{jp}{2(j+1)}, \quad h_0 \asymp 1, \quad CH \sim k/2, \quad SA\bar{h} \sim k/2, \quad D/S \sim k/2, \\ \alpha \sim \frac{Sy^2K^2}{2(j+1)}, \quad f_c \sim \frac{j^2y^2F_b}{2(j+1)}, \quad r_c \sim \frac{jy^2(1-p)K}{2(j+1)}. \end{gathered} \tag{RL7}$$

These use the exact masses in RFN12 and the closed RSR formulas. All displayed positive masses are strictly positive. Write $\kappa = -k_c > 0$. The exact positive-surplus formula RSR19 proves this sign. Its substituted finite polynomial gives

$$\frac{\kappa}{r_c} = O(p + Sy) \quad (p \in \mathrm{I}_c), \qquad \frac{\kappa}{jy^2} = O(1). \tag{RL8}$$

For the remaining case in (RL1), p and j are positive bounded units, and

$$\frac{\kappa}{f_c} \sim \frac{pK}{jF_b} \leq \frac{1}{2}, \qquad jF_b - 2pK = (j-p)K \geq 0. \tag{RL9}$$

The inequality is a bound on real residues when combined with the preceding asymptotic equality. On this case η_2 is infinitesimal, and the effective current defined below is a positive bounded unit.

210.1.2 The entire parent and its harmonic shift

Retain the complete original quarter-parent RFN4. Its exact inverse state change at $w = (W + uY)/(1+u^2) = 1$ is

$$X = N_0 + pv - uz, \quad Z = z + uN_0 + puv, \quad W = 1 + u(M_0 + Hvz), \quad Y = u - M_0 - Hvz. \tag{RL10}$$

Here $z > 0$ is arbitrary. Retain $mpD_u^2N_0^2$ and divide the remaining whole parent by $D_u = 1+u^2$. The exact expression is RFN7, with both complete state normals, hard entries RFN9, and scalar RFN12.

Use the additional exact change

$$q = vz - Tu, \qquad L = M_0 + Hq, \qquad M_0 = L - Hq, \qquad L = \sqrt{S\ell\bar{h}}\,\widehat{L}. \tag{RL11}$$

All divisions are pure source or pure testing. In particular this change does not invert the harmonic mixed unit $1 + \rho u^2$. The complete diagonal of $\widehat{L}$ is

$$S\ell\bar{h}\{D_u(CD_v + D + \tfrac{1}{2}) + AD_v + (\alpha + \tfrac{1}{2})v^2\}, \tag{RL12}$$

with positive real residue $\overline{k}/2$ and bounded coefficient array through both openings.

Use the circuit coordinates $U, V, Z_*, Q_*, B_*, R_*, \Psi$ of SH1–SH2, and $b = v - uz/j = \sqrt{y}B_*$. Before spending any scalar n-mass, complete the exact correlated block, as in RFN13–RFN16:

$$h_0 N_0^2 + n(N_0 + jb)^2 = (n + h_0)(N_0 + c_b b)^2 + M_e b^2, \quad c_b = \frac{nj}{n + h_0}, \quad M_e = \frac{nh_0 j^2}{n + h_0}. \tag{RL13}$$

Keep $nv^2(N_0 + pb)^2$ and all three additional positive hard squares RFN15 whole, including after the change $\widehat{N} = \sqrt{n + h_0}(N_0 + c_b b)$. At $n = 0$, put $c_b = M_e = 0$ and omit the current row. For $n > 0$, $c_b^2/M_e = n/[h_0(n + h_0)]$ is bounded.

The exact shifted scalar. Let F be RFN12. The shifted scalar is $F - 2HqL_M + H^2q^2H_M$. Use the whole $d_cV^2Q_*^2$ term in its q^2 diagonal. Using the exact actual-source identity $S_0 = 0$ in RFN22, the resulting coefficient on $(\sqrt{S}q)^2$ is exactly

$$D_q = S\{A\bar{h}^2 + \ell^2(C + D + \tfrac{1}{2})D_u\} - \frac{f_c}{Tx} - TH^2C\,u^2v^2 - \frac{k_cv^2}{yTx}. \tag{RL14}$$

It has bounded array and positive real residue $\overline{k}/2$. Indeed $f_c/(Tx) = O(jy)$ is infinitesimal, $TH^2C = \ell CH$ is bounded, and $\kappa/(yTx) = \kappa/(jy^2)$ is bounded. Its equivalent Q_*^2 mass has order jy. The only extra scalar cross terms are

$$-2Hq\{\mu uvb + v(\alpha - yCHu^2)R\}, \qquad R = z - uv/(Sy). \tag{RL15}$$

The budgets below allocate their entire expressions.

The remaining signed scalar terms are exactly $e_0V^2B_*^2 + k_cZ_*^2(V^2 - 1)^2$, where $e_0 = p^2y^2/2 - g_c$. The negative part of e_0 is at most g_c. SH2 purchases it at relative circuit cost

$$\frac{g_c}{f_c}(1 + V^2) = O(S(y + v^2)) \in \mathrm{I}. \tag{RL16}$$

Use $Z_*(V^2 - 1) = -VQ_* - R_*$ for the other signed term. If p is infinitesimal, (RL8) pays its opposite part directly from $r_cR_*^2$, and its VQ_* cost is $O(v^2)$. Otherwise p, j, M_e are positive units. Young's exact square identity with parameter $1/6$, followed by SH1 at $a = 3/2$, gives the legal SOS domination

$$\kappa Z_*^2(V^2 - 1)^2 \preceq_{\mathrm{sos}} \frac{7}{4}\kappa\Psi + \frac{21}{2}\kappa V^2Q_*^2 + \frac{7}{2}\kappa V^2B_*^2. \tag{RL17}$$

The circuit cost is at most $7/8 + o(1)$, by (RL9). The two other relative costs are $O(v^2)$. Thus fixed positive real fractions of $D_q(\sqrt{S}q)^2$, $r_cR_*^2$, $f_c\Psi$, and M_eb^2 remain; the last is absent only at $n = 0$. The opposite mass r_c was not spent in the second case.

210.1.3 All helper rows and both complete normals

As usual, a free row permits any fixed real square pivot at infinitesimal cost from the preceding retained scalar forms. SH2 proves that both $E_u = u(uvz - jy)$ and $E_v = v(uvz - jy)$ are free. The row $R_u = v(u^2 - x)$ is free as well. For p infinitesimal, use the actual RFN19 rational Gram: its three costs are $O(S^2y)$, $O(Syu^2)$, and $O(Su^2/(1 - p))$. When p, j, M_e are units, apply SH1 to the R_*^2 term in RFN19. The resulting costs are

$$O(S^2y + Su^2/j), \quad O(Syu^2 + Su^2v^2), \quad O(jSu^2v^2/M_e), \tag{RL18}$$

all infinitesimal.

The rows pv^2b and $\sqrt{S}\,\mu uvb$ are free. If η_2 is infinitesimal, (RL6) and (RL13) imply

$$\frac{p^2}{M_e} = \frac{p^2}{h_0j^2} + \frac{p^2}{nj^2} = O(1). \tag{RL19}$$

They then cost $O(v^4)$ and $O(Su^2v^2)$. If η_2 is a positive unit, the actual source gives $p^2y/f_c = O(1)$ and $\mu = O(p)$. The second identity in SH2 buys the same rows with costs

$$O(v^2(y+v^2)), \qquad O(Su^2(y+v^2)), \tag{RL20}$$

including $n = 0$.

The full opposite term $\sqrt{S}\,v(\alpha - yCHu^2)R$ is free by SH4–SH8. Also $j\alpha b$ is free:

$$j\alpha b = -(\alpha/B)R_u - \alpha uR, \qquad \alpha/B = O(y(j+1)) \in \mathrm{I}_c. \tag{RL21}$$

For p infinitesimal, the direct r_c cost for αuR is $O(Sy^2(j+1)^2u^2)$. In the unit p, j, M_e case use SH1; the three costs are $O(Sy^2u^2)$, $O(Sy^2u^2v^2)$, and $O(Sjy^2u^2v^2/M_e)$. All are infinitesimal. These statements account for the whole source α, not only its leading coefficient.

The first normal. The exact RFN21 source cancellations give, after (RL11),

$$L'_N = [j(n-\alpha) + p(n+\tfrac{1}{2})v^2]b + K_sR_u + \kappa_N E_v, \tag{RL22}$$

$$K_s = -\mu + (\alpha+\tfrac{1}{2})(\ell+p-1), \qquad \kappa_N = -(n+\tfrac{1}{2})(1-p/j).$$

Both coefficients are bounded. For κ_N, use the exact numerator of n and $n(j-p)/j = O(p/[j(j+1)])$; this cancels the r opening. After the entire correlated completion (RL13), the remaining normal is $-j\alpha b + \frac{1}{2}pv^2b + K_sR_u + \kappa_N E_v$. Every row is free by (RL18)–(RL21). Division by $\sqrt{n+h_0}$ is bounded. The shift generates couplings to $\sqrt{M_e}b$ with bounded coefficients $c_b/\sqrt{M_e}$; they are absent at $n = 0$. Large fixed real helper pivots absorb these couplings by exact Young identities.

The second normal and the entire hard block. The complete RFN22 normal after (RL11) is

$$L_L = (C_q - HH_M)q + \mu uvb + v(\alpha - yCHu^2)R, \qquad C_q - HH_M = -D - H(C+D+\tfrac{1}{2})u^2. \tag{RL23}$$

The coefficient of $\sqrt{S}q$ after the normalization of L is infinitesimal; D is infinitesimal and $H(C + D + 1/2)$ is bounded. The other two terms are free by (RL20) and SH6–SH8. Exactly the same rows pay (RL15), whose first factor is $\sqrt{S}q$ with a bounded source multiplier.

The remaining hard off-diagonal is $(\alpha + 1/2)uv\sqrt{S\ell\bar{\bar{h}}}/\sqrt{n+h_0}$, which is infinitesimal. Its c_bb correction has the same bound after division by $\sqrt{M_e}$. The complete finite Gram therefore has a positive definite real residue, with principal normal pivots

$$1, \quad \overline{k}/2, \quad \overline{k}/2 \tag{RL24}$$

on $\widehat{N}, \widehat{L}, \sqrt{S}q$, together with the retained strict scalar and purchased helper pivots. All extra positive hard squares and both original retained current packets are restored whole. The strict bounded-Gram lemma proves tensor SOS for the entire parent.

210.1.4 State and outer-chart exhaustion

The fresh full-state sign lemma SJ1–SJ10 permits positive u, v, and on the remaining negative-core chart gives $wz > 0$. Divide by the nonzero pure-testing w to obtain the normalization used above with arbitrary $z > 0$. No estimate in this proof uses its size. Literal axes and homogeneous endpoints have the direct original-core SOS exit. Real and imaginary state parts separate. The exact outer theorem R2O1–R2O24 covers every other testing pair on all original R2 sources. This proves the stated all-tests conclusion on (RL1).

The complementary source a_s unbounded and p a positive unit is not inferred from this proof. It requires the separate TU source comparison and additional-current conductor.

Chapter 211

Two exact circuit budgets and the complete R2 opposite normal

Component `p09b2107cc05e`
Source: `NEW/weak_c2/r2_shifted_current_helpers.tex`
SHA256: `09b2107cc05e6bc59cd3b6cace84175aba218efc77c894335db8ff8bdf34fa77`
Inventory use: SH1-SH8

211.1 Two exact circuit budgets and the complete R2 opposite normal

Use the compact actual-source dictionary of RSR1–RSR8. In the R2 source put $S = 1/T$, $x = jSy$, $B = Sy$, and $H = S\ell$, where $0 < \ell < 1$. Both testing angles u, v are positive infinitesimals. Normalize the rotated testing coordinate $w = 1$; the other rotated coordinate $z > 0$ is arbitrary, including unbounded values. No coefficient estimate below involves z.

Set

$$U = \frac{u}{\sqrt{x}}, \quad V = \frac{v}{\sqrt{y}}, \quad Z_* = \frac{z}{\sqrt{j/S}}, \quad Q_* = U - VZ_*, \quad B_* = V - UZ_*, \quad R_* = Z_* - UV,$$

$$\Psi = U^2 + V^2 + Z_*^2 + U^2V^2Z_*^2 - 4UVZ_*.$$

All monomials and radicals separate between the two fields.

A parameterized opposite-square identity. For every positive real $a > 1$, let $K = a/(a-1)$, $d = U - V$, $t = UV$, $E = Z_*(1-t)$, $T_* = t(Z_* - 1)$. Exact multiplication gives

$$\begin{aligned} a\Psi + KV^2(Q_*^2 + B_*^2) - R_*^2 = {} & ad^2 + (a-1)(E - T_*/(a-1))^2 \\ & + (2at + KV^4)(1 - Z_*)^2 + 2KZ_*(t - V^2)^2. \end{aligned} \tag{SH1}$$

The right side is a finite legal separated SOS. For example $a = 3/2$ gives $K = 3$. This is a different budget from the older one with $U^2B_*^2$; neither budget dominates the other uniformly at independent source and testing rates.

The two symmetric current budgets are

$$\begin{aligned} (1+U^2)\Psi &= R_*^2 + B_*^2 + U^2Q_*^2 + U^2(1 - UVZ_*)^2, \\ (1+V^2)\Psi &= Q_*^2 + R_*^2 + V^2B_*^2 + V^2(1 - UVZ_*)^2. \end{aligned} \tag{SH2}$$

In particular the two actual rows

$$E_u = u(uvz - jy), \qquad E_v = v(uvz - jy) \tag{SH3}$$

are free from a fixed positive fraction of $f_c\Psi$ whenever $f_c \asymp j^2y^2$. Their respective relative circuit costs are $O(x+u^2)$ and $O(y+v^2)$, both infinitesimal. Here *free* means that an arbitrarily large fixed real multiple of the row square can be inserted at infinitesimal cost in the strict circuit Gram. It is not a pointwise bound without an SOS witness: multiply (SH2) by j^2y^2x or j^2y^3.

The full opposite row after the harmonic shift. Let

$$q = vz - u/S, \quad R = z - \frac{uv}{Sy}, \quad R_v = u(v^2 - y),$$
$$d_s = \frac{BCH(1-p)(1+x)}{E}, \qquad E = 1 + B(HT + p - 1).$$

The exact actual source identity is

$$\frac{\alpha}{y} - CHx = d_s. \tag{SH4}$$

Indeed it is equivalent to $(K + Hjy) - jE = (1-p)(1+x)$, after canceling the displayed positive source factors in the RSR dictionary. The two exact testing identities

$$u(u^2 - x) = SE_u - Su^2q, \qquad v(u^2 - x) = SE_v - Suvq \tag{SH5}$$

now give the complete identity

$$\begin{aligned}\sqrt{S}\, v(\alpha - yCHu^2)R = {} & [\alpha - yCHu^2 - CHu^2(v^2 - y)]\,(\sqrt{S}q) \\ & + \sqrt{S}\, CH(v^2 - y)E_u - \frac{d_s}{\sqrt{S}}R_v.\end{aligned} \tag{SH6}$$

There is no restriction on z, and no normal has been set to zero. The first coefficient in (SH6) is infinitesimal, because $CH = O(1)$, y, u, v are infinitesimal, and $\alpha = O(Sy^2(j+1))$ is infinitesimal by $jy \in \mathrm{I}$. The second coefficient is infinitesimal and (SH3) is free.

For the last term write

$$R_v = -\sqrt{x}\, y(Q_* + VR_*), \qquad \frac{d_s}{\sqrt{S}} = \sqrt{S}\, yCH(1-p)(1+x)/E. \tag{SH7}$$

Suppose the scalar has retained strict pivots of respective orders jy and $jy^2(1-p)$ on Q_*^2, R_*^2, as in the actual R2 source. Twice Young's identity and (SH7) buy this row at relative costs

$$O(S^2y^4(1-p)^2), \qquad O(S^2y^2(1-p)v^2), \tag{SH8}$$

respectively. Both are infinitesimal even when $1-p$ has arbitrarily small positive order. No inverse of that opening remains. Multiplication by the additional bounded source factor $\sqrt{\ell(1-\ell)}$, used to normalize the complete second normal, preserves the estimate.

Thus the entire opposite term in that normal is controlled uniformly through all R2 source ratios, once the stated scalar pivots have been retained. This lemma resolves that normal allocation only. It does not assert the signed scalar allocation or all-source R2 positivity.

Verification. The companion checker proves (SH1)–(SH7), including (SH4) from the actual rational source module, as exact identities. The ordered-field estimates and the conditional scalar-pivot hypotheses in (SH8) are explicit analytical statements.

Chapter 212

R3 and R4 at infinitesimal e, with all source ratios and tests

Component `p1de8b5ffab69`
Source: `NEW/weak_c2/r34_e_inf_all_sources_all_tests.tex`
SHA256: `1de8b5ffab6931a4c77de0ccdd50f8bd5de94b1fcbbce66030129714ae3bbd84`
Inventory use: EN1-EN25

212.1 R3 and R4 at infinitesimal e, with all source ratios and tests

Statement and exact domain. Use the original common-II stationary source, its six native weights n, m, C, A, D, α, all its strict source gaps, $d = 1$, $\lambda = \beta = 0$, and $n \geq 0$. Suppose B_s, a_sB_s, t_sB_s are positive infinitesimals, t_s, h are positive bounded units, and $e = 1 - r$ is positive infinitesimal. In addition impose either

$$\begin{aligned} &\text{R3}: \quad 1 - h \text{ is infinitesimal,} \\ &\text{R4}: \quad 1 - h \text{ is a positive unit and } 1 + a_sB_sr - (1-h)/t_s \text{ is infinitesimal.} \end{aligned} \tag{EN1}$$

There is no condition on the size of a_s, $a_s^2B_s$, a_se, or their ratios. The conclusion is finite separated tensor SOS for the entire original parent at every independent projective testing pair and every state, including signs, axes, and literal $n = 0$. The already recovered positive-unit-e proofs then complete the whole original R3 and R4 rows. No other weak-C2 row is inferred.

The source and testing fields are independent real closed extensions $K_c, K_t \supset \mathbb{R}$. A bounded coefficient means an element of $B_c \otimes B_t$; its residue is the homomorphism to $\mathbb{R}$. The strict bounded-Gram lemma is used only for matrices with positive definite real residue. Row entries may be arbitrarily large. Every inverse and radical below is pure source, pure testing, or a product of such factors.

212.1.1 One exact compact source for both ends

Use the exact compact variables of the reconstructed actual-source dictionary RSR2–RSR6:

$$\begin{aligned} &c_s = 1 + a_sB_sr, \quad R_s = 1 + ha_sB_s, \quad j = a_sc_s/R_s, \quad B = B_sR_s/c_s, \quad T = t_sc_s/R_s, \\ &p = a_se/R_s, \quad H = (1-h)/(t_sc_s), \qquad x = jB = a_sB_s, \quad y = TB = t_sB_s, \quad q_z = jT. \end{aligned} \tag{EN2}$$

Thus $0 < p < \min(1, j)$, $0 < H < 1$, and

$$B, x, y \in \mathrm{I}_c, \quad T \in \mathrm{U}_c, \quad 1 - HT \in \mathrm{U}_c, \quad p/j \in \mathrm{I}_c. \tag{EN3}$$

The last assertion is exact source entry: $e = p(1+x)/(j+px) \sim p/j$. In R3, H is infinitesimal. In R4, $1 - H$ is infinitesimal and $0 < \overline{T} < 1$. In particular $H(1-H) \in \mathrm{I}_c$ in both rows. The native inverse is

$$r = \frac{j-p}{j+px}, \quad h = \frac{1-HT}{1+HTx}, \quad a_s = \frac{j+px}{1+HTx}, \quad B_s = \frac{B(1+HTx)}{1+pB}, \quad t_s = \frac{T(1+pB)}{1+HTx}. \tag{EN4}$$

The exact six weights are RSR6 at this substitution. Those rational identities, including every contact first jet, were checked against the native source without size restrictions. Their former R1 size hypothesis is not imported.

Put $\sigma = 1 + (1-H)T$, $K = j + 1 - p$, $E = 1 + B(HT + p - 1)$, and use the explicit polynomials $\mathcal{P}, \mathcal{L}, \mathcal{N}, \mathcal{F}, \mathcal{K}$ in RSR5, RSR15, RSR18. Then

$$\frac{\mathcal{P}}{j+1} \sim 1, \qquad \frac{\mathcal{L}}{j+1} \sim \sigma, \qquad \frac{\mathcal{F}}{j+1} \sim \sigma. \tag{EN5}$$

Here $T, \sigma, (1-HT)^{-1}$ are bounded positive units. To prove these uniform statements, divide the printed finite polynomials by $j+1$. The corrections in $\mathcal{P}$ contain $B + x$ or TB; those in $\mathcal{L}$ after $(j+1)\sigma$ contain $p/(j+1)$ or $B + x$. In $\mathcal{F}$ the only additional quotient is $p(1-p)/(j(j+1))$, infinitesimal by (EN3). Every other term contains B or $Bj/(j+1)$. Thus no restriction on j or jT is needed.

The exact physicality numerator has the grouping

$$\mathcal{N} = p\{1+B(3j+1-p)+BT(j+1)\} - Bj(j+1)\{1+(1-H)T\} + \mathcal{R}, \quad |\mathcal{R}| \le C_0 BT\, Bj(j+1). \tag{EN6}$$

The bound follows termwise from RSR5 using $p < \min(1, j)$ and bounded T. Since $n = \mathcal{N}/[2(j-p)\mathcal{P}] \ge 0$,

$$\begin{aligned} &\eta_* = \frac{Bj(j+1)}{p} = O(1), \qquad 0 \le n = O\Big(\frac{p}{j(j+1)}\Big) \in \mathrm{I}_c, \\ &\eta_* \in \mathrm{I}_c \Longrightarrow n \sim \frac{p}{2(j-p)(j+1)} > 0. \end{aligned} \tag{EN7}$$

This includes $\mathcal{N} = 0$ literally.

Write $\mu = m(1-p)$, $h_0 = \mu + 1/2$. The full source formulas give

$$\begin{aligned} &\mu = O(p), \quad h_0 \asymp 1, \quad \alpha/[B^2(j+1)] \sim \sigma/2, \quad \alpha/B \in \mathrm{I}_c, \\ &A \sim \frac{T}{2(1-HT)}, \qquad CH \sim \frac{\sigma}{2T}, \qquad D(1-H) \sim \frac{1}{2T}. \end{aligned} \tag{EN8}$$

Consequently A and the finite one of C, D have positive real lower bounds. The other coefficient is unbounded. The exact normalized source limits are

	C	A	D
$R3$	$+\infty$	$T/2$	$1/(2T)$
$R4$	$1/(2T)$	$T/[2(1-T)]$	$+\infty$.

(EN9)

Only real residues of the bounded entries are meant in this table. The coefficient m may be infinitesimal, a unit, or unbounded.

212.1.2 The scalar allocation in the complete two-normal frame

Use the entire parent RFN4 and the exact full-state change RFN5–RFN9, with $w = (W + uY)/(1 + u^2) = 1$ and arbitrary $z = (Z - uX)/(1 + u^2) > 0$. The identities are rational identities of (EN4), so they retain both complete normals and every state term in the present source. The tests in this subsection satisfy $u, v > 0$ infinitesimal. For clarity, the inverse is

$$X = N_0 + pv - uz, \quad Z = z + uN_0 + puv, \quad W = 1 + u(M_0 + Hvz), \quad Y = u - M_0 - Hvz. \tag{EN10}$$

Retain the actual positive packet $mpD_u^2 N_0^2$, and divide the remaining entire parent by the pure testing factor $D_u = 1 + u^2$. Its exact expansion is RFN7–RFN9. In particular the hard entries are

$$\begin{aligned} &H_N = h_0 D_u + \tfrac{1}{2}v^2 + \alpha u^2 + nD_v, \\ &H_M = D_u(CD_v + D + \tfrac{1}{2}) + AD_v + (\alpha + \tfrac{1}{2})v^2, \qquad H_{NM} = (\alpha + \tfrac{1}{2})uv. \end{aligned} \tag{EN11}$$

Keep the separated rows $U = u/\sqrt{x}$, $V = v/\sqrt{y}$, $Z_* = z/\sqrt{q_z}$, $Q_* = U - VZ_*$, $B_* = V - UZ_*$, $R_* = Z_* - UV$, and

$$\Psi = U^2 + V^2 + Z_*^2 + U^2V^2Z_*^2 - 4UVZ_* = (U-V)^2 + Z_*^2(1-UV)^2 + 2UV(1-Z_*)^2. \quad \text{(EN12)}$$

The last radical is separated, for example $\sqrt{UV} = (xy)^{-1/4}\sqrt{uv}$. The exact scalar identity is RFN12 with the same actual circuit masses. From (EN5), the closed formulas RSR16–RSR20 now give

$$\begin{aligned} &f_c \sim Tx^2\sigma/2, \quad q_c \sim \frac{Tx}{2(1-HT)}, \quad r_c \sim TjB^2(1-p)\sigma/2, \\ &g_c/f_c = Hy\,\mathcal{L}/\mathcal{F} \in \mathrm{I}_c, \quad k_c < 0, \quad |k_c|/(q_c y) = O(p+B). \end{aligned} \quad \text{(EN13)}$$

In particular all three displayed positive masses are positive. The sign of k_c follows from the same exact positive-surplus identity RSR19: its error bound uses $B(j+1) \in \mathrm{I}_c$, not T infinitesimal.

Before spending any current mass, perform the exact source-only completion

$$\begin{aligned} h_0N_0^2 + n(N_0+jb)^2 &= (n+h_0)(N_0+c_bb)^2 + M_eb^2, \\ b = v - uz/j, \quad c_b &= \frac{nj}{n+h_0}, \quad M_e = \frac{nh_0j^2}{n+h_0}. \end{aligned} \quad \text{(EN14)}$$

Keep $nv^2(N_0+pb)^2$ and the positive extra hard terms RFN15 as whole squares. This reserves exactly the correlated n packets of the complete parent. If $n = 0$, set $c_b = M_e = 0$ and omit that row. For $n > 0$, $c_b^2/M_e = n/[h_0(n+h_0)]$ is bounded.

The signed scalar allocation differs from R1 in one source split. The e_0 cost is unchanged: its negative part is at most g_c, and RFN18b buys it at cost $H(\mathcal{L}/\mathcal{F})(y+v^2) \in \mathrm{I}$. Also $d_c > 0$. Use $Z_*(V^2-1) = -VQ_* - R_*$ for k_c. The VQ_* cost is $O((p+B)v^2)$, infinitesimal. For its opposite-row part the exhaustive cases are:

$$\begin{array}{c|l} p \in \mathrm{I}_c & |k_c|/r_c = T\mathcal{K}/[(1-p)(1+Hy)\mathcal{L}] = O(p+B) \in \mathrm{I}_c \\ p \in \mathrm{U}_c,\ \eta_* \in \mathrm{U}_c & j \text{ unbounded},\ j^2B \in \mathrm{U}_c,\ |k_c|/f_c \in \mathrm{I}_c,\ |k_c|/(f_cx) = O(1) \\ p \in \mathrm{U}_c,\ \eta_* \in \mathrm{I}_c & j \text{ unbounded},\ M_e \in \mathrm{U}_c,\ |k_c|/f_c \in \mathrm{I}_c, \\ & |k_c|/(q_cy) = O(1),\ |k_c|/(M_eyx) = O(1). \end{array} \quad \text{(EN15)}$$

In the second case j is unbounded because p/j is infinitesimal; $j^2B \asymp p\eta_*$. In the third, $nj^2 \sim p/2$ and n is infinitesimal, so $M_e \sim p/2$. The closed formulas show $|k_c|/f_c = O((p+B)/j)$ in the last two cases and $|k_c|/(M_eyx) = O((p+B)(1+1/n)/j^2)$ in the third. These bounds are uniform through $p \to 1$.

Case two uses RFN18a: $R_*^2 \preceq_{\mathrm{sos}} (1+U^2)\Psi$. Case three uses RFN18c: $R_*^2 \preceq_{\mathrm{sos}} 2(\Psi + V^2Q_*^2 + U^2B_*^2)$. The non-infinitesimal source ratios in (EN15) acquire the testing factors u^2 or v^2. Thus all costs are infinitesimal. Literal $n = 0$ lies in the first or second case; it is not discarded. After this finite allocation, fixed positive fractions of

$$q_cQ_*^2, \quad r_cR_*^2, \quad f_c\Psi, \quad M_eb^2 \quad \text{(EN16)}$$

remain, with the last row absent exactly at $n = 0$.

212.1.3 Both complete normal arrays, with the new endpoint weights

The exact derivatives RFN21–RFN22, and their exact source cancellations $S_N = S_0 = S_1 = S_2 = 0$, hold unchanged. Let $q = vz - Tu$, $R = z - uv/B$, $R_u = v(u^2 - x)$. In particular

$$\begin{aligned} L_N^{\mathrm{rem}} &= -j\alpha b + \tfrac{1}{2}pv^2b + K_RR_u + (K_c/j)uvq, \\ L_M &= C_qq + \mu uvb + v(\alpha - BTCHu^2)R, \\ K_c &= p/2 + j\{H(\alpha+1/2) - 1/2\} + n(p-j), \\ K_R &= -\mu - (\alpha+1/2)(1-p) + (T/j)K_c, \\ K_d &= CH - (D+1/2)(1-H) = -\mu/j - BTCH, \\ C_q &= K_d + AH + 1/2 + v^2\{CH(1+u^2) + AH + (\alpha+1/2)H\}. \end{aligned} \quad \text{(EN17)}$$

Every displayed coefficient other than a retained row is bounded: p/j and n are infinitesimal, $\mu/j = O(p/(j+1))$ is bounded, and $A, CH, D(1-H), T$ are bounded by (EN8). In particular K_R, K_c/j, and C_q are bounded.

Here is an explicit audit of all costs, to avoid importing the former $T \in \mathrm{I}_c$ hypothesis of RFN. The exact radial Gram RFN19–RFN20 purchases R_u^2 at infinitesimal cost from (EN16). Its three cases are still $1-p$ a unit; $p \sim 1, \eta_*$ a unit; and $p \sim 1, \eta_*$ infinitesimal. Their costs are respectively

$$\begin{array}{r|l} 1-p \in \mathrm{U}_c & O(B),\ O(Bu^2),\ O(u^2/(1-p)) \\ p \sim 1, \eta_* \in \mathrm{U}_c & O(B + u^2/j + u^4/(j^2 B)),\ O(Bu^2) \\ p \sim 1, \eta_* \in \mathrm{I}_c & O(u^2/j),\ O(u^4/j),\ [8u^4 + 2(u^2-x)^2]/M_e. \end{array} \tag{EN18}$$

They require only T bounded above and below, because T cancels between $y = TB$ and the masses (EN13). The small-current budgets NSR3–NSR5 likewise apply unchanged: both $pv^2 b$ and $\sqrt{T}\,\mu uvb$ are free. For η_* infinitesimal their coefficients are $O(v^4)$ and $O(Tu^2v^2)$, using $p^2/M_e = O(1)$. For η_* a unit the coefficients are $O(v^2(y+v^2))$ and $O(Tu^2(y+v^2))$, using $p^2 y/f_c = O(1)$. Each is infinitesimal.

The row $q/\sqrt{T}$ is strict by (EN13). Its N coupling has coefficient $(K_c/j)uv\sqrt{T}$ infinitesimal. The identity $j\alpha b = -(\alpha/B)R_u - \alpha u R$ handles the remaining N term. For $1-p$ a unit the latter square costs $O(B^2(j+1)^2u^2/(1-p))$ against $r_c R_*^2$. For $p \sim 1$, RFN18a bounds the entire $j\alpha b$ square by $O(B^3(j+1)^2 + B^2(j+1)^2u^2/j)$ of the circuit. These are infinitesimal. A sufficiently large fixed real radial pivot pays the bounded K_R coupling, including the bounded shift $c_b/\sqrt{M_e}$ from (EN14).

Normalize the complete second normal by $M_0 = \sqrt{TH(1-H)}\,\widehat{M}$. The full hard diagonal (EN11) then has residue

$$\overline{TH(1-H)H_M} = \begin{cases} (1+\overline{T})/2, & R3, \\ 1/2, & R4. \end{cases} \tag{EN19}$$

Every term of the complete diagonal is bounded: the only divergent factors occur in CH and $D(1-H)$. The normalized q coupling is $T\sqrt{H(1-H)}C_q$, now infinitesimal because $H(1-H)$ is infinitesimal. This is the replacement for the T-infinitesimal argument in R1. The normalized current coupling is a bounded multiple of the already purchased $\sqrt{T}\,\mu uvb$.

For the entire remaining opposite row put $J = B(j+1) + u^2$ and $a_R = Tq_z v^2(\alpha - BTCHu^2)^2$. Since $\alpha - BTCHu^2 = O(BJ)$, its costs are

$$\begin{array}{ll} a_R/f_c = O((T/j)v^2J^2), & a_R/(q_c y) = O(v^2J^2), \\ a_R/r_c = O(Tv^2J^2/(1-p)), & a_R/(M_e y x) = O(T(1+1/n)v^2J^2/j^2). \end{array} \tag{EN20}$$

The direct r_c cost applies for $1-p$ a unit. For $p \sim 1, \eta_*$ a unit, RFN18a applies using $j^2 B$ a unit. For $p \sim 1, \eta_*$ infinitesimal, RFN18c applies using $(1+1/n)/j^2 = O(1)$. Thus the entire square, including every higher testing degree, is free. No state ratio occurs in a cost.

Finally the complete mixed hard term is $2(\alpha+1/2)uv\sqrt{TH(1-H)}(\widehat{N}/\sqrt{n+h_0} - c_b b)\widehat{M}$. Its bounded coefficients have zero residue. The strict bounded-Gram argument therefore applies exactly as in the final paragraph of RFN: choose a large fixed radial pivot, buy the finitely many free rows, and retain positive fractions of (EN16). The two hard pivots are 1 and the positive number (EN19). Restore all whole packets in (EN14), RFN15, and $mpD_u^2N_0^2$, and the pure testing factor D_u. This proves the full local parent for all infinitesimal $u, v > 0$ and arbitrary $z > 0$, with both unrestricted state normals. The exact SJ state/sign reduction supplies every state and the literal axes in this local chart.

212.1.4 Every other testing angle, with weak m retained

This outer join uses only (EN9), positivity of the six native weights, and exact full-parent identities. Set $\widetilde{m} = m/(1+m)$, a bounded positive source coefficient below m, and retain the difference as the whole original m square. Keep the entire n and α events throughout. For the single unbounded coefficient C or D, retain its excess over a sufficiently large positive real cap. Other selected coefficients have the actual bounded positive residues (EN9).

Finite Bob with nonzero residue and bounded Alice. Let v have nonzero finite residue, u be bounded, and use the five full-state rows

$$P = uW, \quad Y, \quad X, \quad Z, \quad S = \sqrt{\widetilde{m}}\, W. \tag{EN21}$$

The selected m event is the entire square $[vS - \sqrt{\widetilde{m}}(X - uvY + uZ)]^2$. All other terms use P, Y, X, Z with bounded coefficients. In R3 the strong row is $P - Y$; on its kernel the selected remaining form on (Y, X, Z) is

$$\frac{T}{2} D_u D_v Y^2 + \frac{1+T}{2T} v^2 (uX - Z)^2 + K_0.$$

In R4 the strong row is $P + uvX - Y - vZ$; on its kernel it is

$$\frac{D_v v^2}{2T}(Z - uX)^2 + \frac{T}{2(1-T)} D_u D_v Y^2 + K_0, \quad K_0 = \frac{D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_u D_v Z^2}{2}.$$

The leading principal minors in the order (Y, X, Z) are

	$R3$	$R4$
1	$\dfrac{D_u(TD_v + v^2)}{2}$	$\dfrac{D_u(T + v^2)}{2(1-T)}$
2	$\dfrac{D_u(TD_v + v^2)(Tu^2v^2 + Tv^2 + T + u^2v^2)}{4T}$	$\dfrac{D_u D_v(T + v^2)(T + u^2v^2)}{4T(1-T)}$
3	$\dfrac{v^2(1+T)D_u^2 D_v(Tu^2 + v^2)}{8T}$	$\dfrac{v^2 D_u^2 D_v^2(Tu^2 + v^2)}{8T(1-T)}$.

(EN22)

All are strictly positive at the real residues, even $u = 0$. The S direction has the positive coefficient v^2 in the selected m event. A sufficiently large real cap on the strong row makes the entire five-row residue positive definite. This is the real finite-cap Schur lemma: if a bounded real quadratic form is positive on the kernel of a linear map, adding a sufficiently large real multiple of the squared map makes it positive definite. The unbounded actual coefficient permits every such cap. This argument includes m infinitesimal.

Infinitesimal Bob and Alice a finite positive unit. Use $R = uW - Y$, $B_t = X + uZ$, $V_Z = vZ$, and Y; the testing inverse $1/u$ is bounded. The full selected parent is a bounded quadratic on these four rows because $v(W + uY) = (v/u)(R + D_u Y)$ and $J = R + uvB_t - D_u V_Z$. Its residue splits off $(\widetilde{m} + 1/2)B_t^2$ and the form

$$CR^2 + AD_u Y^2 + (D + 1/2)(R - D_u V_Z)^2 - D_u Y V_Z - \tfrac{1}{2} D_u V_Z^2. \tag{EN23}$$

In R3 restrict to the strong-row kernel $R = 0$. After removing a positive factor D_u, the matrix on (Y, V_Z) is $\left(\begin{smallmatrix} T/2 & -1/2 \\ -1/2 & (1+T)D_u/(2T) - 1/2 \end{smallmatrix}\right)$; its determinant is $(1+T)u^2/4 > 0$. In R4 the strong kernel is $R = D_u V_Z$. On (R, Y) its first diagonal is $1/(2T) - 1/(2D_u) > 0$, and its determinant is $u^2/[4(1-T)] > 0$. A large real cap again makes (EN23) strict.

Infinite Bob and bounded Alice. Put $w = 1/v$, divide the homogeneous parent by v^2, and use the five rows (EN21). Select any fixed positive real lower caps c_0, a_0, d_0 below C, A, D, and the actual $\widetilde{m}$ event. The full quadratic is bounded; its residue is

$$[S + \sqrt{\widetilde{m}}\, uY]^2 + c_0(P - Y)^2 + (a_0 + \tfrac{1}{2}) D_u Y^2 + d_0(uX - Z)^2 + \tfrac{1}{2} D_u X^2. \tag{EN24}$$

This is positive definite on the five rows, for every bounded residue of u. It includes $u = 0$ and m infinitesimal.

Infinite Alice. Let $s = 1/u$ and keep the entire m, n, α events. For finite v of nonzero residue, after division by u^2 the selected residue on (W, Y, X, Z) is

$$c_0 D_v W^2 + a_0 D_v Y^2 + (d_0 + \tfrac{1}{2})(W + vX)^2 + \tfrac{1}{2}(vY - Z)^2,$$

which is positive definite. For v infinitesimal use instead $(N, W, Y, V) = (sX + Z, W, Y, vX)$. The complete signed core after division by u^2 is

$$\frac{1}{2}\{W+(1+s^2)V-sY-svN\}^2+\frac{1}{2}(1+s^2)v^2Y^2-(1+s^2)Y(vN-sV)-\frac{1}{2}(1+s^2)(vN-sV)^2+\frac{1}{2}D_vN^2.$$

Adding the selected C, A, D packets gives strict residue $c_0W^2 + a_0Y^2 + (d_0 + 1/2)(W + V)^2 + N^2/2$. Every full coefficient is bounded; no s/v is used. If also Bob is infinite, divide by u^2v^2 and use $(W, X, Y, S_Z, T_Z) = (W, X, Y, sZ, wZ)$. The complete selected residue is

$$c_0 W^2 + a_0 Y^2 + d_0(X - S_Z)^2 + \tfrac{1}{2}X^2 + \tfrac{1}{2}(Y - T_Z)^2, \tag{EN25}$$

again positive definite. The full quadratic follows directly from $J/(uv) = wW + X - swY - S_Z$ and the exact signed core; its coefficients are bounded polynomials in s, w.

These charts exhaust all nonnegative projective angles: both infinitesimal are the complete local theorem, and every other pair is one of (EN21)–(EN25) or (EN23). All strict-matrix coefficients use only bounded pure source or pure testing factors. Restore every selected excess, the untouched m, n, α packets, and the homogeneous testing squares. The exact SJ sign gauges, literal projective endpoints and real/imaginary state split finish the asserted all-tests theorem.

Audit scope and dependencies. The source dictionary and exact scalar/normal identities are the freshly reconstructed RSR, NSR, RFN sources; only their explicitly rechecked algebraic identities and finite SOS budgets are used. Equations (EN3)–(EN20) prove all changed source and coefficient bounds afresh. The entire outer join (EN21)–(EN25) is new. The focused checker corroborates source coefficient limits, full-parent expressions, complete hard entries, and every printed positive kernel matrix. Analytical uniform estimates and the exhaustive join are the proof above, not an inference from its receipt.

Chapter 213

The complete uniform infinitesimal C1 source, at all tests

Component `p845b3cb8b114`
Source: `NEW/weak_c2/uniform_c1_all_tests_join_rederived.tex`
SHA256: `845b3cb8b114ecf6bfee1273bc12f4cc753b5b4dc230831c485c8691a9b4f33e`
Inventory use: fresh complete uniform source/all-testing join UJ1–5

213.1 The complete uniform infinitesimal C1 source, at all tests

Theorem and unchanged source scope. Use the original physical canonical C1 stationary source with $\lambda = n = 0$, $d > 0$, every original nonnegative event coefficient, and every original strict source gap. Suppose

$$k, f, g, e = kf/g > 0 \text{ are infinitesimal}, \qquad r = 1 - af^2, \quad R = 1 + k^2 r, \quad T_0 = 1 + k^2 - tk^2 g^2,$$
$$a, t > 0, \qquad J = T_0 - a > 0, \quad M = R - t > 0. \tag{UJ1}$$

Then the entire original parent is a finite tensor sum of squares for every independent projective testing pair and every real or complex testing state. There is no restriction on $b = g/k$, g/f, or any relative infinitesimal rate. In particular the original $\beta = 0$ face is retained.

This is a mathematical join of the actual bounded-b endpoint product-repair proof, the fresh whole Z-chart proof EZ1–EZ19, the fresh inverse-b local proof IW1–IW24, and the two full-parent outer-current identities below. No unavailable historical EZ or IW coefficient witness is used as a premise.

213.1.1 Exact original-source bounds for the two outer identities

Normalize $d = 1$ by a positive pure-source scale. The retained `endpoint_source.tex` prints seven original coquotients. Their respective residues on (UJ1) are

$$(c_2, c_\beta, c_3, c_5, c_4, c_\alpha, c_d) \sim (2, 2, 2, 2, 2, 2, 4).$$

Their arguments are bounded, so they are positive source units. The actual weights satisfy

$$C = \frac{c_3}{af^2 c_d}, \qquad D = \frac{B_c}{2Jf^2 c_d}, \qquad A = \frac{f^2 c_4}{c_d}, \qquad B_c = 2c_5 - Jf^2 c_d \sim 4.$$

Consequently

$$C^{-1}, D^{-1} \in \mathfrak{m}_c, \qquad g_B = \frac{CD}{C + D} > 0, \qquad \gamma = A g_B = \frac{c_4 c_3 B_c}{c_d^2 (aB_c + 2Jc_3)} \sim \frac{1}{4}. \tag{UJ2}$$

This is uniform in both openings a, J: the last denominator has residue $16 \cdot 4(\bar{a} + \bar{J}) = 64$, since $a + J = T_0 \sim 1$. No positive lower bound on either opening is assumed. These bounds use the actual original C, D, A, not selected substitute weights. Any other positive event is restored whole.

213.1.2 The two exact outer-current identities

Here are the retained identities with the $d = 1$ normalization made explicit. Write $D_u = 1 + u^2$, $D_v = 1 + v^2$, and let the original core be

$$\mathcal{H} = 2(uW + uvX - Y - vZ)^2 + 2D_u(vY - Z)^2 + 2D_v(X + uZ)^2 - 2D_uD_vZ^2.$$

Set

$$R_t = uW - Y, \quad C_t = v(uX - Z), \quad B_t = X + uZ, \quad h_B = \frac{D}{C + D}, \quad N_t = R_t + h_BC_t,$$

$$c_u = D_u^{-1/2}, \quad t_u = uD_u^{-1/2}.$$

All angle normalizations are pure testing.

If $1/v$ is bounded, put $\eta_v = \sqrt{D_v}/v$ and

$$K_0 = 2\sqrt{g_B}\, C_t, \qquad B_0 = \sqrt{2}\sqrt{D_v/D_u}\, B_t.$$

Then the entire selected parent has the exact identity

$$\begin{aligned} 4CD_vR_t^2 + 4D(R_t + C_t)^2 + \mathcal{H} &= 4(C + D)N_t^2 + 4Cv^2R_t^2 + 2(R_t + C_t)^2 \\ &\quad + 2D_u(vY - Z)^2 + (K_0, B_0)Q(K_0, B_0)^T, \\ Q &= \begin{pmatrix} 1 - \dfrac{c_u^2\eta_v^2}{2g_B} & \sqrt{\dfrac{1}{2g_B}}\, t_u\eta_v \\ \sqrt{\dfrac{1}{2g_B}}\, t_u\eta_v & 1 \end{pmatrix}. \end{aligned} \tag{UJ3}$$

Its bounded Gram has residue I_2, by (UJ2), since c_u, t_u, η_v are bounded. This proves the selected parent tensor-SOS for every other angle and every state. Restoring the other nonnegative original events proves the whole-parent result.

It remains outside the small-angle corner that v is infinitesimal and $1/u$ is bounded. Put $\epsilon = 1/g_B$, $q = \sqrt{\epsilon}$, and

$$K_1 = 2\sqrt{g_B}\, C_t + \sqrt{\epsilon}\, Y, \quad B_1 = B_t - uvY, \quad V_0 = vY, \quad Y_0 = \sqrt{\epsilon}\sqrt{D_u}\, Y,$$
$$L_0 = \frac{qc_u}{2}(K_1 - c_uY_0) - t_uvB_1 + c_uvV_0.$$

Direct multiplication gives $L_0 = \sqrt{D_u}\, v(vY - Z)$ and the exact selected-parent identity

$$\begin{aligned} &4CD_vR_t^2 + 4D(R_t + C_t)^2 + 4AD_uD_vY^2 + \mathcal{H} \\ &\quad = 4(C + D)N_t^2 + 4Cv^2R_t^2 + 2(R_t + C_t)^2 + \mathcal{G}, \\ \mathcal{G} &= K_1^2 + 2D_v(B_1^2 + V_0^2) + (4\gamma D_v - c_u^2)Y_0^2 \\ &\quad + 2qv(K_1 - c_uY_0)V_0 - 2L_0^2. \end{aligned} \tag{UJ4}$$

The Gram residue on (K_1, B_1, V_0, Y_0) is

$$\operatorname{diag}(1, 2, 2, 4\bar{\gamma} - \bar{c}_u^2) = \operatorname{diag}(1, 2, 2, \bar{t}_u^2) \succ 0.$$

Here $\bar{t}_u^2 > 0$ follows from bounded $1/u$. Every correction contains q or v and has zero residue. Again restore all other positive original events.

Both identities are whole-parent identities, not state restrictions. They use bounded normalized homogeneous angle coordinates, and $\eta_v^2 = 1 + (1/v)^2$; clearing the original pure-testing homogeneous factors gives the same certificates at literal Alice or Bob infinity. The exact sign gauges of SJ1–SJ10 preserve all event packets and reduce finite angle signs to the nonnegative ones used here. Literal zero angles have the direct core SOS exits. Thus (UJ3)–(UJ4) cover every testing chart except $u, v > 0$ both infinitesimal.

213.1.3 Every remaining source and state chart

At u, v infinitesimal normalize a nonzero testing state by a coordinate of largest absolute value. This is a pure-testing operation; all four coordinates are bounded and at least one is a unit. If X or Y is a unit, the original pure-testing core has positive real residue $2(X^2 + Y^2)$. It is a positive testing square, and every original event remains a positive separated square. We can therefore assume X, Y infinitesimal.

If Z is a unit, normalize $Z = 1$. The remaining W is arbitrary bounded and may have either sign. EZ1–EZ19 applies in each of its two actual-source coordinate rows:

source	exact coordinates	actual proof
$b = g/k$ bounded	$f = eb, \quad g = kb$	EZ1−EZ19, bounded b
$b = g/k$ unbounded	$(k, f, g), \quad k/g \in \mathfrak{m}_c$	EZ1−EZ19, inverse b.

(UJ5)

The extra inverse-b source inequality used by EZ is derived from the actual $\beta \geq 0$: $tb^2 \leq 2c_\beta/c_d$. Thus it adds no source hypothesis to (UJ1). Its full proof retains both state normals and the testing-angle normal, including unbounded scaled normal coordinates through their exact sN_y, sN_v coefficients.

If Z is infinitesimal, W must be a unit. Normalize $W = 1$; then u, v, X, Y, Z are all infinitesimal. For bounded b, use the actual retained `product_repair.tex` proof. Its source variables are exactly (e, k, b, a, t) with $f = eb$, $g = kb$; its gaps $R_c = 1 + k^2(1 - ae^2b^2)$ and $T_0 = 1 + k^2 - tk^4b^2$ coincide with (UJ1). It allows every bounded positive b, including infinitesimal b, both strict-gap openings, and all original coefficient faces. Its whole seven-packet polynomial, complete first normals, higher-normal support and nine-root product identity are printed with `endpoint_source.tex` and `endpoint_hermite.tex`. Its final original-parent restoration is an exact sum of complete positive packets. There is no remaining condition on k^3/e or k/e in that proof.

For unbounded b, use IW1–IW24. Its exact source hypotheses $k, f, g, k/g \in \mathfrak{m}_c$, physical $\beta \geq 0$, and the original gaps follow directly from (UJ1) and this ratio class. It proves the complete original parent with both unrestricted state normals for every positive rotated state ratio, so it includes the present W-unit chart. The retained β packet, including its zero face, and both source openings are part of that fresh proof. The state and sign reduction is the actual SJ1–SJ10 identity, not a historical chart label.

These cases exhaust every source ratio and every state. The zero state is immediate. A complex state splits into its real and imaginary evaluations because the parent matrix is real. Undo all pure-testing normalizations and the positive source scale d. This proves the theorem on (UJ1) at all independent projective tests.

Exact dependency scope. The retained bounded-b endpoint proof and its two printed source inputs are available bodies in the recovered checkpoint. The outer identities above are printed and checked as finite algebra. The fresh local premises are EZ1–EZ19 and IW1–IW24, each with its own current analytical review and whole-parent checker. This theorem closes only the uniform infinitesimal C1 source. It neither imports a missing historical witness nor asserts that every original C1 source enters (UJ1). That source-routing question belongs to the separate global assembly.

Chapter 214

The actual hypotheses used by the retained weak-C2 leaves

Component `p49f367f1b5e8`
Source: `NEW/weak_c2/weak_c2_bare_source_scope_audit.tex`
SHA256: `49f367f1b5e891824f752e7d13b23dc2db0000e8948a5758a4bc7a4926120028`
Inventory use: BSR1-BSR15

214.1 The actual hypotheses used by the retained weak-C2 leaves

This section proves the scope extension needed for the source join. Several retained theorem statements were written on a complement of earlier admissions. Such a complement is not a mathematical assumption in the coefficient calculations below. The extension is justified by checking the entire proofs, including the state normals and the remote joins; it is not inferred from their historical admission labels.

Common source hypotheses. Throughout this section use the original physical stationary C2 source, with $d = 1$, $\lambda_{\mathrm{event}} = \beta = 0$, its actual contact, and all original strict gaps and nonnegative event coefficients. In particular $n \geq 0$, including equality. Put

$$\begin{gathered} z = aB, \quad y = tB, \quad H = hB, \quad L = lB, \quad l = 1 - h, \quad e = 1 - r, \\ g = 1 - ae + haB, \quad j = 1 + aBr - l/t. \end{gathered} \tag{BSR1}$$

Here B, z, y are positive source infinitesimals, $0 < h, r < 1$, $g, j > 0$, and $B - zyr > 0$. The full quarter-parent is the polynomial HJ2 in the following source-assembly section. All the retained identities cited here are identities for that polynomial, not only for its value on a contact graph. A *unit* below is positive, bounded, and has positive real residue. The source and testing fields are arbitrarily chosen, separately labelled real closed extensions of $\mathbb{R}$; no algebraic-disjointness assumption is made. They may be two labelled copies of the same extension. The source and testing choices are independent in the universal quantifiers. The positive-infinitesimal total-scale condition may be retained; on every row with an unbounded original coefficient it follows already from that coefficient.

214.1.1 Regular A-opening exits

On t, g, j unit and h infinitesimal, the four actual source exits have exactly the following hypotheses:

	additional source conditions	complete bodies
WA	r, e, a unit	`five_root_local`, `five_root_remote`
WE	r, e unit, a infinitesimal	`extension`
SL	r infinitesimal, a bounded	`local_small_r`, `remote_small_r`
WC	e infinitesimal, a unit	`centered_circuit_closure`, `remote_reduction`.

(BSR2)

The WA, WE and SL statements explicitly require no residual membership premise. Their source bounds use the original formulas, and their complete five-root identities keep both state normals. WE uses the exact monic a-division of $(b_2 - \alpha/B^2)/a$; its canceled denominator is a unit. It requires no bound on h/a. SL uses nr, Ah and the normalized roots $\sqrt{n}\,\xi, \sqrt{A}\,\eta, aR_2$; it requires no bound on r/h, a/r, or a testing/source ratio. Their printed homogeneous remote identities restore every projective test.

For WC the precise extra scale bound is proved, rather than assumed. The exact original identity is

$$\begin{aligned} 2r\Delta n &= eQ - BK, \qquad K = (1+a)(h+t)(1+aB), \\ Q &= 1 + B(1 + t + ah + 2ar + at) \\ &\quad + B^2\{a(h+t) + a^2[h(r+1) + t(1 + r - r^2)]\}. \end{aligned} \tag{BSR3}$$

On the WC domain Q, K, Δ are positive units. Thus B/e and n/e are bounded, including $n = 0$. These are exactly BS1–BS3, not an unprinted assumption about the relative rates of B, e, h. WC3 and WC13–WC18 prove the full scalar and normal coquotient bounds; its reciprocal testing-ratio argument uses the same physical n/m bound. The finite three-circuit lift is used only after these complete coefficient checks. GR1–GR7 give the remote parent, including the additional $\sqrt{m}W$ root. None of these four exits calls a later weak-C2 row or the present source join.

214.1.2 Unit-e parts of R3 and R4

The actual bodies `r3_e_unit_all_tests` and `r4_e_unit_all_tests` apply on their explicit row conditions together with e unit, without an earlier-exit premise. Indeed $g > 0$ first gives bounded a. In their source calculations all denominators are units after replacing the potentially divergent coefficients by

$$nr, \quad mg, \quad Cl \text{ in R3}, \qquad nr, \quad mg, \quad Dj \text{ in R4}. \tag{BSR4}$$

The full ξ-row is normalized by $\sqrt{n + 1/2}$. R3 keeps $\sqrt{C}\,\eta$; R4 uses the exact complete centered-D identity RD3–RD6 and solves its l-unit coefficient. The only apparent a-pole is the same exact WE monic quotient used in BSR2. For g infinitesimal the centered m excess is retained and the remainder depends on bounded mg, rather than on $1/g$. The strict two-by-two blocks R3E5–R3E6 and R4E7–R4E8 include all these openings. The remote proofs use only the printed positive lower bounds and the complete selected-parent identities R4E14–R4E16. Their source estimate $nr \sim e/[2(1+a)] > 0$ proves that $n = 0$ is impossible on these particular domains. It is not removed from a neighboring source branch.

214.1.3 R5

On bare R5, t, h, l, j are units and r is infinitesimal. Positivity of g gives bounded a with real residue $\rho \in [0, 1]$. Writing $\chi = h_0$, $\ell = 1 - \chi$, and $\tau = t_0 > \ell$, the actual source calculation R5.1 gives

$$\begin{aligned} (nr)_0 &= \frac{1}{2(1+\rho)}, \qquad A_0 = \frac{\tau}{2\chi(1+\rho)}, \\ C_0 &= \frac{\chi + \tau - \rho\ell}{2\ell(1+\rho)}, \qquad D_0 = \frac{1}{2(1+\rho)(\tau - \ell)}, \\ (\alpha/B^2)_0 &= \frac{\chi + \tau - \rho\ell}{2(1+\rho)}. \end{aligned} \tag{BSR5}$$

Every displayed pivot is positive. For $\rho < 1$, m/a^2 is bounded; for $\rho = 1$, $mg \sim 1/4$ and m is unbounded. These alternatives are exhaustive, and nr proves $n = 0$ impossible here.

The complete local proof uses $\sqrt{n}\,\xi$, the exact WE quotient when $\rho = 0$, and the centered excess R5.2–R5.5 when $\rho = 1$. For clarity, that last identity is

$$mM^2 - (m-K)(M - gvW)^2 = KM^2 + 2(m-K)g\,vWM - (m-K)g^2v^2W^2, \tag{BSR6}$$

where $M = vW - X + uvY - uZ$ and $K > 0$ is real. It is an entire-event identity; it does not change the source coefficients in any other packet. The full (η, R_1) determinant is $A_0(C_0 + D_0 + 1/2) > 0$. The finite remote determinants R5.8 and R5.10, the $\rho = 0$ extra root aW, and R5.13–R5.17 cover all remaining states and projective tests. Their entries use precisely BSR5, nr, and mg. No earlier-admission condition or relative opening rate occurs in this chain. Therefore it proves bare R5.

214.1.4 R6

On bare R6, t, h, l, j, r are units and g is infinitesimal. The exact equality $a(e - H) = 1 - g$ excludes infinitesimal a. For bounded a, e is a unit and $a_0 = 1/e_0$. R6.1 gives positive-unit $n, A, C, D, \alpha/B^2, mg$. The entire proof consists of BSR6 followed by the strict local block R6.2, the full kernel determinant R6.4, and the homogeneous matrices R6.5–R6.7. Only these actual residues are used; no prior-complement condition is needed.

For unbounded a, put $s = 1/a$ and $b = a^2B$. The exact physical identity BSR3, in the original variables, proves b bounded before any contact normalization. The literal zero of n is retained in that inequality. The positive bounded b is either a unit or infinitesimal:

$$\begin{array}{r|l} b \text{ unit} & \texttt{r6_b_unit_all_tests} \quad (\text{RB1}-\text{RB24}) \\ b \text{ infinitesimal} & \texttt{r6_b_inf_all_tests} \quad (\text{RI1}-\text{RI20}). \end{array} \tag{BSR7}$$

Both proofs retain the actual centered m packet and, where required, the whole actual n packet before forming the bounded remainder. RB5–RB9 give its full polynomial inverse and actual contact. RB13–RB18 and RI8–RI15 verify the raw scalar support and both complete first-normal modules before applying the restricted CC1–CC6 conductor. The infinitesimal-b proof verifies the additional ε, j filtrations on the full coefficients; it does not import an unrestricted contact-ideal inclusion. Its strict leading block is RI19. The adjacent unit-state charts retain the original n packet when that is the needed pivot.

The remote inputs in the two dependency manifests are selected full-parent identities: BE4, the BE infinity/zero identity, R5.5, RT10, and R6.5–R6.7. Their lower bounds are verified from the actual R6 source. The C1 source theorem and a bounded-a R6 source theorem are not transported to this domain. Hence the three actual R6 bodies prove its entire bare source. $n = 0$ remains in the unit-b component; where a positive residue excludes it, that exclusion is proved by the original formula.

214.1.5 R7 and its one genuine source exclusion

Bare R7 has t unit and h, g infinitesimal, with r, j not both infinitesimal. This last condition is used and must be kept. For bounded a, $a_0 = 1/e_0$, and R7.1 gives

$$C_0 = \frac{t_0 - e_0}{2(1 + e_0)}, \qquad (\alpha/B^2)_0 = \frac{t_0 - e_0}{2e_0(1 + e_0)}, \qquad (Ah)_0 = \frac{t_0}{2(1 + e_0)}. \tag{BSR8}$$

Here $t_0 \geq 1$, $0 < e_0 \leq 1$. The retained exclusion is exactly what prevents $t_0 = e_0 = 1$; it makes the first two pivots strict. The full proof `r7_bounded_a_all_tests` uses the centered m identity and the exact centered D event

$$L_D = uW + uvX - Y - vZ + jvZ. \tag{BSR9}$$

Its five-root matrix identity R7.3 includes the zero syzygy and both normals. After extracting DL_D^2, the remaining coefficients involve bounded nr, Ah, mg, Dj. The appropriate hard rows are normalized jointly. The remote cap lemma, full kernel matrices, and R7.6–R7.12 use the same bounds, with no comparison between the four divergent coefficients.

For unbounded a, the actual source gives $r \sim 1$; thus the retained r, j exclusion is automatic. The body `r7_unbounded_a_body` derives bounded $b = a^2B$, $n = O(s^2)$, and bounded positive $mg, Ah, D(1 - c)$ directly from the source formulas. RU7 is the entire bounded remainder after all four actual positive packets have been removed. RU13–RU16 check its full raw support and normal modules; RU20 verifies the coefficient filtrations before reusing the restricted RI14–RI15 algorithm.

The complete strict block RU24 has determinant $(t_0+2)/(4t_0) > 0$, including $t_0 = 1$. RU26–RU30 and the selected BE identities supply the remote and projective join. No earlier-exit condition is used.

The source join below proves exactly

$$r, g, h, j \text{ infinitesimal, } t \text{ unit} \quad \Longrightarrow \quad C \text{ infinitesimal.} \tag{BSR10}$$

Consequently the excluded R7 intersection cannot occur on the directed strong-C input. One must not extend BSR8 across that intersection by deleting the condition.

214.1.6 R8

Bare R8 has t, g units and h, j infinitesimal; the exact gap gives $t_0 = 1$. There are two exhaustive cases for r. For r a unit, `joint_D_R8_closure` gives the complete local proof JD1–JD18. The centered-D identity JD4 removes the whole divergent rank-one block, including both normal terms. It then splits the actual source into bounded a with e unit, bounded a with e infinitesimal, and unbounded a. The latter uses the positive source quantity $\rho = n/e^2$: its noninfinitesimal part has the strict weighted pivot, its positive infinitesimal part has the exact paired-D thin lift, and $n = 0$ has the physical map described below. JD14–JD17 verify the new D-dependent coquotients; no inverse of j remains in their bounded arrays. This is an algebraic comparison with a formal source, not application of a physical C1 theorem at negative beta.

For r infinitesimal, the independent body `r8_small_r_local` derives

$$(nr)_0 = (Ah)_0 = (Dj)_0 = \frac{1}{2(1+a_0)}, \qquad 0 \le a_0 < 1. \tag{BSR11}$$

It jointly normalizes $\sqrt{n}\,\xi$, $\sqrt{A}\,\eta$, and $\sqrt{D}\,L_D$, using the exact matrix syzygy RD3–RD6. RD7 checks every full remaining coefficient, including the monic a-quotient at $a_0 = 0$. Its five strict residue pivots are printed in RD9–RD11. No relative rate among r, h, j is required. JD19–JD24 and the exact homogeneous selected-parent identities cover the remote complement for both cases of r. Thus their union proves bare R8; it uses no historical residual-exclusion premise.

214.1.7 R9 and R10

Bare R9 has t, g, j units and h, e, a infinitesimal. Its exact source denominator has unit residue $2(t_0 - 1) > 0$. The R9 source appendix proves the cancellations

$$\lambda = \frac{m - a^2 q}{t a^2 B} \sim \frac{1}{2}, \qquad c_F \sim \frac{1}{2t_0}, \qquad C \sim Ah \sim \frac{t_0}{2}, \qquad D \sim \frac{1}{2(t_0-1)}. \tag{BSR12}$$

The body `reciprocal_contact_closure`, RC1–RC13, contains the entire scalar, both complete first normals, and the hard block. Its coupled circuit removes the apparent unbounded testing-ratio coefficient; the final homogeneous roots have no such inverse. The R13 frontier's R9 extension of GR1–GR7 uses the actual bounds $B/e, n/e$ and $m/(a^2 e)$, the full additional $\sqrt{m}W$ row, and the printed homogeneous endpoints. All these estimates are on the explicit bare source, including $n = 0$.

Bare R10 has t, g, j units, h, e infinitesimal, and a unbounded. The R10 source-and-weighted proof puts $s = 1/a, w = ae, b = a^2 B$ and proves

$$b/w = O(1), \qquad n = O(s^2 w), \qquad m/w \sim \frac{1}{2(1-w_0)}, \qquad w_0 < 1. \tag{BSR13}$$

The bound on b/w comes from the original inequality BSR3 before any normalized circuit is introduced. The same source proof gives the full remote parent, retaining bounded m as a whole packet where needed. The actual all-tests join has three local source cases: $\rho = n/e^2$ noninfinitesimal, $n = 0$, and $0 < \rho$ infinitesimal. The first is the complete weighted identity R10.9 and strict source Schur cases R10.11–R10.12. The last is the whole thin-parent identity TL4–TL10 together with

`r10_unit_W_join`. The latter proves the exact critical deficit divided by m equals s times a bounded source coefficient; it includes every bounded normalized Z, including zero. The maximal-coordinate state partition supplies exactly that chart and its complementary Z-unit chart. The restricted CA/CS/SE conductors are used only after the full raw supports, both normal arrays, and the required scaled divisibilities have been checked. No bound on e/B, h/e, or an independent testing/source ratio is inserted.

The literal zero boundary in R8 and R10. The only physical C1 source import just used is at literal $n = 0$. The exact OP4 map is

$$x_1 = z, \quad \eta_1 = t, \quad \zeta_1 = 1/a, \quad r_1 = h, \quad \theta_1 = 1 - r. \tag{BSR14}$$

It gives the identical four contact state coordinates. OP2 and OP5 prove that the formal beta is exactly zero on this boundary. The original C2 parent is therefore the actual C1 parent, up to its allowed positive coefficient normalization. In the unbounded-a cases where this import is used,

$$k_1 = \sqrt{z},\ \zeta_1,\ r_1 \text{ are infinitesimal}, \qquad \eta_1 = t \text{ is a unit}, \qquad e^2_{1,\mathrm{scale}} = t\,a^2 B \text{ is positive bounded}. \tag{BSR15}$$

If the last scale is infinitesimal, apply the actual theorem `small_e_all_tests`. Otherwise apply `centered_a_bounded_delta`. The former covers every positive native r_1; the latter covers the present infinitesimal native r_1. Both statements retain every original positive opening and literal $\beta_1 = 0$, and neither has an earlier-complement premise. Their full centered-parent and both-normal proofs, outer/state lemmas, and homogeneous endpoints are the stated imports. No whole-C1 routing theorem or later weak-C2 result is a dependency. For $n > 0$, OP5 gives negative formal beta; BSR14 is then never used as a physical-source theorem.

Conclusion of the scope proof. Every use of a retained source theorem has now been matched to its explicit algebraic hypotheses. Every use outside its old source label is an exact selected-parent identity or a restricted conductor whose complete coefficient hypotheses have been proved in the importing body. In particular all full hard blocks and both state-normal rows remain in the respective identities. The only retained source exclusion needed by the ten-row join is the simultaneous r, j condition in R7, and BSR10 dispatches it explicitly. Thus the listed bodies prove the bare-source versions needed for HJ, without a circular appeal to the union being assembled. Every file and the exact imported locations are pinned in the accompanying used-input manifest.

Chapter 215

The complete directed weak-C2 source and testing join

Component `pec6c617d3a3d`
Source: `NEW/weak_c2/weak_c2_complete_source_and_testing_join_rederived.tex`
SHA256: `ec6c617d3a3da5fb604976cb80c903ef1b029248819a4b4f91fd59e5206193c4`
Inventory use: HJ1-HJ16 and row/state restoration

215.1 The complete directed weak-C2 source and testing join

Theorem. Let K_c, K_t be arbitrarily chosen, separately labelled real closed extensions of $\mathbb{R}$; no algebraic-disjointness assumption is made. In particular they may be two labelled copies of the same extension. Use an original physical stationary C2 source with $d > 0$, $\lambda_{\text{event}} = \beta = 0$, all event coefficients nonnegative, and every original strict source gap. Normalize $d = 1$. Suppose the positive actual source chart has two infinitesimal contact half-angles and

$$C \notin \operatorname{Inf}_c, \qquad \alpha \in \operatorname{Inf}_c, \qquad \delta_{\text{tot}} = \frac{1}{1+n+m+C+A+D+\alpha} \in \operatorname{Inf}_{c,>0}. \tag{HJ1}$$

Then the entire original parent is a finite sum of squares in $K_c \otimes_{\mathbb{R}} K_t$ at every independent projective testing pair and every real or complex state. There is no earlier-admission-complement hypothesis. The theorem retains literal $n = 0$ and every allowed independent source opening. By the exact reflection HJ3 it also proves the original directed weak-C, strong-α HH source. The entry into the physical positive source chart is a separate global source-entry theorem, not a conclusion inferred from this join.

The full parent and the reflection. For finite testing half-angles write

$$D_u = 1 + u^2, \quad D_v = 1 + v^2, \quad M = vW - X + uvY - uZ, \quad J = uW + uvX - Y - vZ.$$

The quarter-parent is exactly

$$\begin{aligned}\mathcal{P} = {} & nD_uD_vX^2 + mM^2 + CD_v(uW - Y)^2 + AD_uD_vY^2 + DJ^2 + \alpha D_u(vY - Z)^2 \\ & + \tfrac{1}{2}\{J^2 + D_u(vY - Z)^2 + D_v(X + uZ)^2 - D_uD_vZ^2\}.\end{aligned} \tag{HJ2}$$

Every proof imported below certifies this entire polynomial. The omission-preserving reflection is

$$\begin{aligned}&(u, v; W, X, Y, Z; n, m, C, A, D, \alpha) \\ &\quad \longmapsto (1/v, 1/u; Z, X, Y, W; n, D, \alpha, A, m, C).\end{aligned} \tag{HJ3}$$

Under HJ3, $\mathcal{P} = u^2v^2\mathcal{P}'$ exactly. This is verified packet by packet and for the full signed core. Its degree-$(2,2)$ homogeneous form exchanges the corresponding angle coordinates, so it includes the literal projective endpoints. It preserves n, A, and interchanges m, D and C, α. A weak-C HH input therefore becomes strong-C, weak-α II. No transformation $t \mapsto 1/t$ of the subsequent source shape is asserted.

215.1.1 Actual source equations and the weak face

Normalize the positive actual contact by $Z_c = 1$, and put

$$\begin{aligned} &z = u_c^2, \quad y = v_c^2, \quad H = v_c Y_c, \quad L = v_c(u_c W_c - Y_c), \quad B = H + L, \quad r = -X_c/u_c, \\ &c_a = 1 + zr, \quad S_c = c_a + L, \quad T_c = B + zr, \quad j = c_a - L/y, \quad G = B + z(H + r - 1). \end{aligned} \tag{HJ4}$$

The unchanged source conditions are

$$z, y > 0, \quad 0 < H, r < 1, \quad L > 0, \quad j, G, B - zyr > 0, \quad n \geq 0.$$

In particular

$$(u_c, v_c; W_c, X_c, Y_c, Z_c) = \left(\sqrt{z}, \sqrt{y}; \frac{B}{\sqrt{zy}}, -r\sqrt{z}, \frac{H}{\sqrt{y}}, 1\right).$$

The exact source denominator and positive numerator are

$$\begin{aligned} \delta &= z + (1+z)B + (1+z)B^2 + 2z(1+z)Br + z^3r^2 + 2z^2r \\ &\quad + y\{(1+z)B + z + z^2r(2-r)\}, \\ M_s &= (1+z)T_c(H+y) - z(1+y)(1-r)L \\ &= (1+z)T_cH + yG + yzj + zrL(1+y). \end{aligned} \tag{HJ5}$$

The six original coefficients, before any size specialization, are

$$\begin{aligned} n &= \frac{T_c c_a S_c - r\delta}{2r\delta}, & m &= \frac{z^2(1+y)(1-r)c_a S_c}{2G\delta}, \\ C &= \frac{c_a M_s}{2L\delta}, & A &= \frac{y c_a T_c S_c}{2H\delta}, \\ D &= \frac{(1+z)T_c(B - zyr)S_c}{2yj\delta}, & \alpha &= \frac{T_c M_s}{2(1-H)\delta}. \end{aligned} \tag{HJ6}$$

All inequalities and all contact identities below refer to these actual coefficients. In particular the numerator of n is never replaced by a limiting source whose sign has not been proved.

Since z, y are infinitesimal, the exact j gap gives $L < y(1 + zr)$, hence L infinitesimal and B bounded. If B is infinitesimal, then H is infinitesimal and $\delta \geq B + z \geq T_c$. The first expression for M_s gives

$$0 < \alpha \leq \frac{T_c(1+z)(H+y)}{2(1-H)} \in \mathrm{Inf}_c. \tag{HJ7}$$

If B is a unit, $H = B - L$, δ, and M_s are units by the positive expression in HJ5. Then α is noninfinitesimal and $CL = c_a M_s/(2\delta)$ is a unit. Thus α is weak exactly when B is infinitesimal. The input HJ1 therefore has

$$B, z, y \in \mathrm{Inf}_{c,>0}. \tag{HJ8}$$

This conclusion is an actual source estimate. It imposes no power-law relation or unprinted rate comparison between B, z, y.

215.1.2 Uniform source alternatives

Put

$$a = z/B, \quad t = y/B, \quad h = H/B, \quad l = L/B = 1 - h, \quad e = 1 - r, \quad g = G/B. \tag{HJ9}$$

The quantities h, l, r, e, g, j are positive and bounded; a, t may be infinitesimal, units, or unbounded. Their exact relations give

$$\begin{gathered} g = 1 - ae + hz > 0, \qquad j = 1 + zr - l/t > 0, \\ ae < 1 + hz, \qquad e \text{ unit} \Longrightarrow a \text{ bounded}, \\ a \text{ unbounded} \Longrightarrow e \text{ infinitesimal}, \qquad a \text{ infinitesimal} \Longrightarrow g \sim 1, \\ g \text{ infinitesimal} \Longrightarrow ae \sim 1. \end{gathered} \tag{HJ10}$$

Here is the source estimate behind the finite partition. From HJ5, $\delta/(B+z) \sim 1$. Moreover $T_c \geq B$, $T_c > z(1-H)$, and $T_c \leq B+z$, so T_c/δ is a positive unit. Consequently

$$r(2n+1) = \frac{T_c c_a S_c}{\delta} \asymp 1, \qquad \frac{AH}{y} = \frac{c_a T_c S_c}{2\delta} \asymp 1. \tag{HJ11}$$

The m formula can be rewritten exactly as

$$m = \frac{(1+y)c_a S_c}{2} \frac{z}{\delta} \frac{1 + zH/B - g}{g}. \tag{HJ12}$$

For unit g this is bounded. For infinitesimal g, $ae \sim 1$, z/δ is a unit, and $m \asymp 1/g$. Thus no independent $a = z/B$ rate creates an omitted intermediate m case.

The positive decomposition of M_s, and its first expression in HJ5, similarly show

original coefficient is unbounded	equivalent source condition
n	r infinitesimal
m	g infinitesimal
A	H/y infinitesimal
C	$L/(H+y)$ infinitesimal
D	y/B infinitesimal or j infinitesimal.

(HJ13)

For completeness, the C upper bound is a bounded multiple of $(H+y)/L$. If $L/(H+y)$ is infinitesimal, the subtracted term in the first formula for M_s is infinitesimal relative to $T_c(H+y)$; this gives the reverse estimate. For D, if B/y is infinitesimal, then $j \sim 1$ and the exact formula gives $D = O(B/y)$. Otherwise zyr/B is infinitesimal, and $D \asymp B/(yj)$. The original gap $B - zyr > 0$ is thereby retained in the region where it cannot be replaced by B.

It follows that all six original coefficients are bounded if and only if

$$t, h, l, j, r, g \text{ are units.} \tag{HJ14}$$

Indeed t infinitesimal makes C, D unbounded; t unbounded makes C, A unbounded. For unit t, HJ13 gives exactly the other five unit conditions. HJ1 rules out HJ14, since the sum in the denominator of δ_{tot} would then be bounded. No positive-d source theorem is needed to remove this empty intersection.

There is also the exact source equivalence

$$C \text{ infinitesimal} \quad \Longleftrightarrow \quad r, g, H/y, j \text{ infinitesimal.} \tag{HJ15}$$

To verify the forward direction, the four positive terms of $M_s/(L\delta)$ first force H/L infinitesimal. Bounded C and HJ13 give $L \asymp y$, hence H/y infinitesimal and $B \asymp y$. The same four summands give $G/\delta, zr/\delta, zj/\delta$ infinitesimal. The factor z/δ cannot be infinitesimal: that would imply $B/\delta \sim 1$ and $G/\delta \sim 1$. Hence r, j are infinitesimal, and $G > 0$ together with z/δ unit gives B/z unit. It follows that g is infinitesimal. Conversely r, g infinitesimal imply $B/z \sim 1$. With $H/y, j$ infinitesimal the gap gives $L/y \sim 1$, hence $B/y \sim 1$. Every positive summand of $M_s/(L\delta)$ is then infinitesimal, proving the converse. On the directed input HJ1 this entire intersection is empty. In particular it is not an allowed unproved continuation of the R7 residue pivots.

215.1.3 The ten rows and every remaining exit

The symbol I denotes positive infinitesimal, U a positive unit, and O positive unbounded. Every row below includes HJ4–HJ10 and the actual coefficient formulas HJ6.

$$\begin{array}{r|l}
\text{R1} & t : \text{I} \\
\text{R2} & t : \text{O} \\
\text{R3} & t, h : \text{U},\ l : \text{I} \\
\text{R4} & t, h, l : \text{U},\ j : \text{I} \\
\text{R5} & t, h, l, j : \text{U},\ r : \text{I} \\
\text{R6} & t, h, l, j, r : \text{U},\ g : \text{I} \\
\text{R7} & t : \text{U},\ h, g : \text{I}, \quad r, j \text{ not both I} \\
\text{R8} & t, g : \text{U},\ h, j : \text{I} \\
\text{R9} & t, g, j : \text{U},\ h, e, a : \text{I} \\
\text{R10} & t, g, j : \text{U},\ h, e : \text{I},\ a : \text{O}.
\end{array} \tag{HJ16}$$

The rows are disjoint. Their complements have the following explicit destinations, not a blanket earlier-admission clause.

Split t first. Its infinitesimal and unbounded cases are R1 and R2. For t unit split h. If h is a unit, test in order whether l, j, r, g is infinitesimal. These are exactly R3–R6. If none is, HJ14 contradicts the total-scale condition. If h is infinitesimal, $l \sim 1$. For infinitesimal g, simultaneous infinitesimal r, j contradicts HJ15 and HJ1; otherwise this is R7. For unit g, infinitesimal j gives R8.

Only t, g, j units and h infinitesimal remain. If e is a unit, HJ10 makes a bounded. For unit r, unit a gives WA and infinitesimal a gives WE. For infinitesimal r, either bounded type of a gives SL. If e is infinitesimal, $r \sim 1$. Then unit a gives WC, infinitesimal a gives R9, and unbounded a gives R10. The complete physical source conditions of WA/WE/SL/WC are exactly BSR2–BSR3. No h/a, e/B, or source/testing comparison has been inserted. This proves exhaustion of the actual source, without asserting that every formal type tuple is physically realized.

215.1.4 Actual complete proof for each row

The mathematical imports in this paragraph are the specified proof bodies, not admission receipts. The bare-scope argument BSR1–BSR15 proves precisely why their earlier residual-complement wording is not a needed hypothesis.

R1 and R2. R1N1–R1N24 in `r1_all_sources_all_tests.tex` prove all original R1 sources. The exact compact-source and full-normal derivation permits arbitrary positive normalized state ratio; its full six-root cap identities and projective entry supply all outer tests. R2A1–R2A4 in `r2_all_original_sources_all_tests.tex` prove all original R2 sources. That theorem explicitly splits bounded a, unbounded a with bounded $n/(tB)$, and unbounded $a, n/(tB)$, using respectively RL/SH, TU, and UL. Its R2O/SJ join supplies every test and state. The formal C1 comparison in TU/UL is implemented by exact coefficients and the complete actual n packet; it never assumes a negative-beta source is physical.

R3 and R4. Every positive $e < 1$ is either a unit or infinitesimal. The unit cases are the actual R3E and R4E all-tests bodies, with BSR4's bare-scope verification. Every infinitesimal e case is EN1–EN25 in `r34_e_inf_all_sources_all_tests.tex`. The latter keeps both complete state normals, the endpoint H or $1 - H$ opening, and the entire source partition before its remote join. These alternatives exhaust each original row.

R5–R7. Bare R5 is BSR5–BSR6. R6 splits into bounded a and unbounded a; in the latter case physicality bounds $b = a^2 B$, whose unit and infinitesimal cases are RB and RI. Thus BSR7 supplies every source of R6. R7 splits into bounded and unbounded a; BSR8–BSR10 and RU supply both. Its sole genuine simultaneous-r, j exclusion has already been dispatched by HJ15. These statements preserve the actual $n = 0$ boundary wherever it is possible.

R8–R10. R8 is the union of the unit-r local JD proof, the infinitesimal-r local RD proof, and JD's remote join valid for both cases. R9 is the complete reciprocal-contact RC1–RC13 body with its actual source appendix and the explicit R9 extension of GR's full remote proof. R10 is the actual three-component all-tests join: its weighted full-parent proof, its positive-thin contact proof, and its unit-W complement. The source subcases in BSR11–BSR15 are exhaustive. The literal $n = 0$ import is exactly OP4 followed by the native C1 small-e or centered-A body. It contains no whole-C1 or whole-weak-C2 routing premise.

215.1.5 State, sign, and projective restoration

Each row proof certifies the full HJ2, either directly on four states or after an exact inverse with both state normals present. All normalizations use pure-source, pure-testing, or positive separated factors. At a nonzero finite chart they are reversed on the SOS roots. Literal zero and infinite testing angles use the actual homogeneous parent, not a limiting division by a zero coordinate. The exact state sign gauges preserve every event packet and the core; the negative-core state normalizations and their complementary nonnegative-core branches are part of the named all-tests bodies. No arbitrary state ratio is bounded by a source scale in this assembly.

Every bounded complete Gram used in those bodies has strict positive real residue on its retained root list. The strict bounded-Gram lemma, including its WA10 polynomial version, therefore gives a finite tensor SOS; it does not take a square root or inverse of a mixed tensor. Restoring all original positive excess packets preserves that conclusion. The original measurement matrices are real, so a complex-state value is the sum of its real and imaginary state values. The source partition HJ16 and its four explicit regular exits now prove the theorem.

Dependency boundary. This is a proof of the complete directed weak-C2 leaf and its exact reflected HH orientation. The positive-total-scale interior is excluded algebraically by HJ1 and HJ14, and the two-weak intersection is excluded algebraically by HJ1 and HJ15. Hence neither global A5 nor a global C2 union is a dependency. Global source entry, other omission patterns, and other source-scale regimes remain the responsibility of their own stated theorems. The accompanying manifest lists actual proof bodies and prerequisite parent/conductor identities with their precise uses; status documents are not mathematical inputs.