

Have You Made Full Use of the OCR Feature?

Make a scan, enhance it and save it. Are these all the features you know about CamScanner? If so, you have missed too many cool experiences. CamScanner offers you lots of features rather than scanning. What we are sharing today is the OCR (Optical Character Recognition) feature.



What can you do with OCR feature?

1. Searching

What can you do if you want to search for a document but just can't remember the names of some docs? Use this feature to recognize all the texts on your scans. Next time you just need to enter some key words in the search box and all the documents within the words will be found.

2. Text extraction

Just purchase the one-time paid version and you can enjoy the text extraction for lifetime! Ever want to edit some texts on a paper document or a PDF file? Import it into CamScanner and all texts can be extracted as .txt file after OCR!

Why wait? Follow the steps to start using OCR!

1. Sign in to CamScanner to sync all your docs -> All texts will be auto recognized after syncing.
2. If you don't want to sign in, you can open one single page of any doc -> Tap the Recognize button -> All recognized texts will be shown in a dialog box -> Tap Share to export the texts.

Saathi

Date ___ / ___ / ___

8d

Page No.

Discrete Structures

well defined collection

- Cantor's def
- Intuitive principle : If $p(x)$ is a property

Then

$$S = \{x \mid p(x)\} \rightarrow \text{Logical } p(x) \text{ is true}$$

↓
proposition or logical statement
either true or false

$p(x) = x$ is a natural no
between 1 and 10

$$\begin{aligned} S &= \{x \mid p(x)\} \\ &= \{2, 3, 4, 5, 6, 7, 8, 9\} \end{aligned}$$

$$\text{Let } f = \{x_1, x_2, \dots, x_n\}$$

we define a function

$$X_A(x) = \begin{cases} 0, & x \notin A \\ 1, & x \in A \end{cases}$$

Here, $X_A(x)$ is called a characteristic function.

$$A_1 \subseteq A$$

$$A_1 = \{x_1, x_2\}$$

$$\underline{f_{A_1}(x)}$$

$$X_{A_1}(x) = \{1, 1, 0, 0, \dots\}$$

$$X_n(x) = \{1, 1, 1, 1, 1, \dots\}$$

We can write sets or its subsets in binary form.

$$X = \{x_1, x_2, \dots, x_n\}$$

$$|X| = n$$

$$P(X) = \text{Power of } X.$$

= The set of all subsets of X

$$\text{eg. } X = \{1, 2, 3\}$$

$$P(X) = \{\emptyset, \{1, 2, 3\}, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}\}$$

Total no. of elements in $P(X) = 2^n$.

$$X_n(x) = \{0, 0, 0, \dots, 0\}$$

↓
2 choices

$$\text{Total} = 2 \times 2 \times 2 \times \dots \text{ntimes} = 2^n$$

$$X = \{82 \text{ students}\}$$

$$A \subseteq X$$

↓
intelligent students

$$A = \{0.9, 0.8, 0.95, \dots\}$$

↳ logic

ambiguity

$$M_A = \begin{cases} 0 & \text{if } x \notin A \\ 1 & \text{if } x \in A \end{cases} \quad X_n(x)$$

→ Membership function
(Artificial Intelligence)

Notation

Date / /

Relations:Ordered pair:Let $A \neq \emptyset$, $B \neq \emptyset$ Let $a \in A$ & $b \in B$.Then (a, b) is ordered pair.Cartesian product:Let $A \neq \emptyset$, $B \neq \emptyset$ Then $A \times B = \{(a, b) \mid a \in A, b \in B\}$ is called the cartesian product of A and B .Let no. of elements in $A = m$ " " in $B = n$

Then, no. of elements in

$$A \times B = m \times n$$

No. of subsets of $A \times B = 2^m \times 2^n$ Function:Let $X \neq \emptyset$ & $Y \neq \emptyset$ thena function f from X to Y is a rule according to which each element of X is associated with a unique element of Y .Defn: A function f from X to Y is denoted as

$$f: X \rightarrow Y$$

$$X = \{1, 2, 3\}, Y = \{3, 5, 7\}$$

$$f: X \rightarrow Y \text{ such that } f(x) = 2x + 1$$

Domain = X

Codomain = Y

Range set / Image set = {3, 5, 7}

- Binary Relation :- A binary relation / operation on a set A is a function such that $f: X \times X \rightarrow Y$

$$f(x, y) = t, \quad x, y, t \in X$$

$$* : X \times X \rightarrow X$$

$$x + y = t \in X$$

i.e. $x + y \in X$


(1) operation addition is binary operation on set of natural no.

(2) " multiplication "

(3) subtraction " is not "

- Unary Operation :-

$$c : P(X) \rightarrow P(X)$$



- Relation : A relation between two sets A and B (or A to B) is a subset of $A \times B$ which consists of ordered pair (a, b) such that a is related to b according to some rule.

$$A = \{1, 3, 5, 7, 9\} \quad |A| = 5$$

$$B = \{2, 4, 6, 8, 11\} \quad |B| = 5$$

$$A \times B = 25$$

Date _____

 $R \subseteq A \times B$ Def: (a, b) iff $a \leq b$

$$R = \{ (1, 2), (1, 4), (1, 6), (1, 8), (1, 11), \\ \cancel{(2, 1)}, \cancel{(2, 3)}, \cancel{(2, 5)}, \cancel{(2, 7)}, \cancel{(2, 9)}, \cancel{(2, 11)}, \\ (3, 6), (3, 8), (3, 11), (5, 8), (5, 11), (7, 8), (7, 11) \}$$

Reflexive Relation:

Let $A \neq \emptyset$ then a relation R is said to be reflexive if $(a, a) \in R \forall a \in A$ or $aRa \forall a \in A$.

Cartesian product of A with itself $(A \times A)$ Symmetric Relation: $a R b \Rightarrow b R a$

$$(a, b) \leftrightarrow (b, a)$$

Transitive: $a R b$ and $b R c$ then $a R c$.

if $(a, b) \in R$ and $(b, c) \in R$
then, $(a, c) \in R$

Let $A = \{a_1, a_2, a_3, \dots, a_n\}$ and $A \times A = \{ (a_i, a_1), (a_i, a_2), \dots, (a_i, a_n), \\ (a_2, a_1), (a_2, a_2), \dots, (a_2, a_n), \dots \}$

$$(a_n, a_1), (a_n, a_2), \dots, (a_n, a_n) \}$$

$$M = A \times A = \begin{bmatrix} (a_1, a_1) & (a_1, a_2) & \dots & (a_1, a_n) \\ (a_2, a_1) & (a_2, a_2) & \dots & (a_2, a_n) \\ \vdots & \vdots & \ddots & \vdots \\ (a_n, a_1) & (a_n, a_2) & \dots & (a_n, a_n) \end{bmatrix}$$

No. of entries = $n \times n$ Total no. of diagonal elements = n

Let R be a reflexive relation. Then the set R will have include diagonal elements of matrix M .

Reflexive element in $M = n^2 - n$

$$R = \{ (a_1, a_1), \dots, (a_n, a_n), \underbrace{\quad \quad \quad}_{\text{n times}} \underbrace{\quad \quad \quad}_{2 \times 2 \times 2} \dots \}$$

$$2^{n^2-n} = 2^{n(n-1)}$$

→ In the reflexive relation apart from diagonal elements of matrix some more elements out of $n^2 - n$ elements will appear. Now, from remaining $n^2 - n$ elements every element have two choices either it can be included or may not be included. Thus, for remaining $n^2 - n$ elements we have $\underbrace{2 \cdot 2 \cdot 2 \cdot \dots \cdot 2}_{n^2-n \text{ times}} \text{ choices} = 2^{n^2-n}$ choices.

Consequently, the no of reflexive relations on a set A of n elements = $2^{n^2-n} = 2^{n(n-1)}$

Q1 No. of symmetric relation on a set of n elements.

<u>Ans</u>	$M = A \times A =$	$(a_1, a_1), (a_1, a_2), \dots, (a_1, a_n)$ $(a_2, a_1), (a_2, a_2), \dots, (a_2, a_n)$ \dots $(a_n, a_1), (a_n, a_2), \dots, (a_n, a_n)$
------------	--------------------	--

Date _____

Ex 1) Let $A = \{a_1, a_2, a_3, \dots, a_n\}$ we can write $A \times A$ in the matrix form

$$M = \begin{bmatrix} (a_1, a_1) & (a_1, a_2) & \cdots & (a_1, a_n) \\ (a_2, a_1) & (a_2, a_2) & \cdots & (a_2, a_n) \\ \vdots & \vdots & \ddots & \vdots \\ (a_n, a_1) & (a_n, a_2) & \cdots & (a_n, a_n) \end{bmatrix}$$

↓
principal
diagn

nn

Any relation R is a subset of $A \times A$ i.e. $R \subseteq A \times A$
 i.e. elements of R , are some entries from matrix M .

We know that R is symmetric if $(a, b) \in R \Rightarrow (b, a) \in R$.
 We have $\frac{n^2-n}{2}$ elements above and below the
 principle diagonal.

$$R = \left\{ 2, 2, 2, \dots, \frac{n^2-n}{2} \right\} = \boxed{\frac{n^2-n}{2}}$$

Total choices of elements below the diagonal = $2^{\frac{n^2-n}{2}}$
 for diagonal elements:

$$2 \times 2 \times 2 \times \dots \text{times} = 2^n$$

$$\boxed{\text{Total choices} = 2^{\frac{n^2-n}{2}} \cdot 2^n}$$

The elements below the principle diagonal & above the principle diagonal will be present in the symmetric relation in the similar manner. Therefore, we check the presence of elements below the principal diagonal in the symmetric relation R .

- No. of ways in which we can check the belongingness of an element in $R = 2$ (either it can be in R or it cannot be in R).
- Total no. of ways in which elements below the principal diagonal can be present in $R = 2 \cdot 2 \cdot 2 \cdot 2 \cdots 2 = 2^{\frac{n(n-1)}{2}}$
- further, n diagonal elements of matrix M can be present in $R = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdots n$ times $= 2^n$

Total no. of elements in a ~~symmetric~~ symmetric relation are $2^n + 2^{\frac{n(n-1)}{2}}$

a) No. of elements in antisymmetric relation on a set of n elements = $2^n \cdot 3^{\frac{n(n-1)}{2}}$

→ Partition of a set :

Let $X \neq \emptyset$

and suppose $S_1, S_2, S_3, \dots, S_n$ be n -subsets of X . Then the set $P = \{S_1, S_2, S_3, \dots, S_n\}$ is called partition of X .

If (1) $S_1 \cup S_2 \cup S_3 \cup \dots \cup S_n = X$

(2) $S_i \cap S_j = \emptyset$ when $i \neq j$.

Rule: partition cannot contain \emptyset .

Ex) (1) $X = \{1, 2, 3\}$

$S_1 = \{1\}$ & $S_2 = \{2\}$

$S_3 = \{3\}$ & $S_4 = \{1, 2, 3\}$

$P = \{S_1, S_2, S_3, S_4\}$

Date _____

$$S_1 = \{1\} \quad S_2 = \{2\} \quad S_3 = \{3\}$$

$$P = \{1, 2, 3\}$$

\rightarrow Union of $S_1, S_2, S_3 = P$

\rightarrow Intersection of any two = \emptyset .

Equivalence relation: If it satisfies three properties, then it is said to be equivalence relation.

- (1) Reflexive
- (2) Symmetric
- (3) Transitive

Equivalence class: Let A be any set and R be a relation defined on A .

Let $a \in A$, then equivalence class of a in A is defined as

$$[a] = \{x \in A \mid a R x\}$$

Example :-

$$\text{Let } A = \{1, 2, 3, 4\}$$

$$\text{Let } R = \{(1, 1), (2, 2), (3, 3), (4, 4), (1, 2), (1, 3), (2, 1), (2, 3), (3, 1), (3, 2)\}$$

Clearly, R is equivalence relation.

Equivalence class

$$[1] = \{1, 2, 3\} \quad R = \{(1, 2)\}$$

$$[2] = \{1, 3\}$$

$$[3] = \{1, 2\}$$

$$[4] = \{4\}$$



There are two equivalent classes,

Partition This with
subset.

Page No. _____

Date 1/1

No. of equivalence relations = No. of partition of that set.

(1)

→ Fundamental Theorem of Equivalence Relation: There is one-to-one correspondence between the number of equivalence relations defined on a set of n -elements and no. of partition of the set.

(2)

→ Partial Order Relations:

Let $A \neq \emptyset$, then R is said to be a partially ordered relation.

- (1) R is reflexive
- (2) R is anti-symmetric
- (3) R is transitive

(3)

Ex) Let $X \neq \emptyset$ and $P(X)$ be power set of X . we define a relation ' \leq ' on $P(X)$ such that $A \leq B$ iff $A \subseteq B$.

→

(1)

Reflexive

we have, $A \subseteq A$ & $A \in P(X)$.
 $A \leq A$

Example

∴ ' \leq ' is Reflexive

(2)

Anti-symmetric

($A R B$ and $B R A$)

$\Rightarrow A = B$)

if $A \leq B$ and $B \leq A$

$\Rightarrow A \subseteq B$ and $B \subseteq A$

$\Rightarrow A = B$

∴ Anti-symmetr. is satisfied

Transitive

Let $A \subseteq B$ and $B \subseteq C$

$\Rightarrow A \subseteq B$ and $B \subseteq C$

$\Rightarrow A \subseteq C \Rightarrow A \subseteq C$

\therefore 'Transitive' is satisfied

1. Give two examples of each of the following.

① Reflexive Relation

② Symmetric

③ Antisymmetric

④ Transitive

⑤ Equivalence

⑥ Partial Order

	1	2	3	4
1	11	12	13	14
2	21	22	23	24
3	31	32	33	34
4	41	42	43	44

2. Find all partitions of the set $\{1, 2, 3, 4\}$.
~~Ans - 7~~ (Incomplete)

Partially ordered set (POSET)

Let $X \neq \emptyset$ and ' \geq ' is a partial order on X .

Then (X, \geq) is called as partially ordered set.

Example: Let $X \neq \emptyset$, then $(P(X), \subseteq)$ is a partially ordered set

when $A \subseteq B$ iff $A \subseteq B \wedge A, B \in P(X)$

Divisibility of two numbers:

Let $a, b \in \mathbb{N}$, then we say that $\frac{a}{b}$ (pronounced as 'a' divides 'b')

if \exists an integer l such that

$$b = a \cdot l$$

Denominator
↑
 a/b
Numerator

Date / /

eg : $\frac{5}{30}$ ✓ $\frac{5}{29} \times$

Ex 2) Let N be the set of natural numbers we define a relation ' \leq ' on N such that $a \leq b$ iff $a|b$.

① Reflexive:

$a|a \quad \forall a \in N$

$\Rightarrow a \leq b \quad \forall a \in N$

$\therefore \leq$ is reflexive

② Antisymmetric:

Let $a \leq b$ and $b \leq a$

$\therefore a|b$ and $b|a$ (By definition)

$\Rightarrow a = b$

$\therefore \leq$ is antisymmetric

③ Transitive:

Let $a \leq b$ and $b \leq c$

$\Rightarrow a|b$ and $b|c$

$\Rightarrow a|c$

$\Rightarrow a \leq c$

$\therefore \leq$ is transitive

$\therefore (N, \leq)$ is POSET

Date _____

m^n

Let D_{14} = The set of divisor of 14.

$$\text{ie } D_{14} = \{1, 2, 7, 14\}$$

Let $a \leq b$ iff $a|b$ & $a, b \in D_{14}$.

It is POSET


Functions

$$f: A \longrightarrow B$$


$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6, 8, 9, 10\}$$

$$f(x) = 2x$$



Let $|A|=m$ $|B|=n$
How many functions can be defined from A to B




$$(\text{no. of choices})^{\text{no. of total elements}} = \text{no. of choice} = n \times n \times \dots \times n = n^m.$$

→ One - One function:

A function $f: A \rightarrow B$ is said to be one-one if distinct elements of A have distinct images
ie Let $x, y \in A$ such that $x \neq y$
then $f(x) \neq f(y)$

or

$$\begin{aligned}f(x) &= f(y) \\ \Rightarrow x &= y\end{aligned}$$




Q) How many one-one functions can be defined from A to B ?

- First condition for

one-one function $n \geq m$

Ans $\begin{cases} \text{No. of choices } P_m \text{ if } n \geq m \\ 0 \text{ otherwise} \end{cases}$



$$\text{Total choices} = [(n - (m - 1))][(n - (m - 2))] \cdots (n - 1)n$$

$$\Rightarrow [1 \cdot 2 \cdot 3 \cdot 4 \cdots (n-m)][(n-(m-1))][n-(m-2)] \cdots [n-(1)]$$

$$= nP_{n-m} = \frac{n!}{(n-m)!}$$

Date / /



Saathi

No. of one-one and onto functions.

$${}^n P_n = \frac{n!}{(n-n)!} = \frac{n!}{0!} = n! \quad (n=m)$$

No. of onto functions that can be defined.

Ans

Answer

1

2

3



4

5

6

7

8



R₃

R₄

R₅

No. of choices in which

one room can be

$$\text{kept vacant} = {}^4 C_1 \times 3^8$$

$$\text{" two rooms } = {}^4 C_2 \times 2^8$$

$$\text{" three rooms } = {}^4 C_3 \times 1^8$$

$$\text{Total choices } = 4^8$$

No. of ways in which 8 guests
can be accommodated in

$$\begin{aligned} & \text{9 room provided} \\ & \text{no room is left vacant} = 4^8 - ({}^4 C_1 \times 3^8 + {}^4 C_2 \times 2^8) \\ & \quad + {}^4 C_3 \times 1^8 \end{aligned}$$

Date / /

Onto: Each element of co-domain should have a pre-image.

Ans: $n^m - \sum_{x=1}^{n-1} n(n-x)^m$

Exercise ① Give two examples of the following :-

- ① Partially Ordered Set
- ② One-One f^n
- ③ Onto f^n
- ④ One-one \hookrightarrow Onto f^n

→ Lattice :-

lower bound and upper bound of a set.

Lower Bound: Let $S \neq \emptyset$ Then lower bound l of S is a real number such that $l \leq x \forall x \in S$.

Upper Bound: Let $S \neq \emptyset$ Then upper bound u of S is a real number such that $u \geq x \forall x \in S$.

finite set

$$S = \{1, 2, 3, 4\}$$

Q) What is lower bound of S ?

A) $-1 \leq x \forall x \in S$

$0 \leq x \forall x \in S$

~~lower bound~~ $-1 \leq x \forall x \in S$

~~of~~ $2 \leq x \forall x \in S$

i) A given set can have infinite number of lower bounds

ii) 1 is greatest among all lower bounds of set S . This is called greatest lower bound (GLB).

Page No. _____

Date / /

Greatest Lower Bound or Infimum of a Set:

Let $S \neq \emptyset$, Then a real no. l is said to be a greatest lower bound of S if :-

$l < u \forall u \in S$
If l' is another ^{lower} bound among ^{bound} of S then $l' \leq l$.

$$\text{Q1} \quad \text{lub } S = \{1, 2, 4\}$$

$$\text{Then, glb}(S) = 1$$

and set S has infinite no. of lower bounds.

$$\text{Q2} \quad S = (0, 1)$$

$$= \{u \mid 0 < u < 1\}$$

$$\begin{array}{c} \hline - & & + \\ & 0 & \\ \hline \end{array}$$

$$\text{glb}(S) = 0$$

To : The glb of a set may or may not belong to the set itself.

Least Upper Bound or Supremum of a Set:

Let $S \neq \emptyset$, then a real no. u is said to be lub S if

$$u \leq v \quad \forall v \in S$$

If v' is another upper bound of S then $v \leq v'$

$$\text{Q1} \quad S = \{1, 2, 3, 4\}$$

$$\text{lub}(S) = 4$$

$$S = (0, 1)$$

$$\text{lub} = 1$$

Date _____ / _____ / _____

Note : The least upper bound or greatest lower bound of a set if exists is unique.

→ lattice

Let $\langle L, \leq \rangle$ be a partially ordered set then
 \downarrow
 usual sum relation

- ① L is said to be a lattice if
 $\text{lub}\{a, b\} = \text{Inf}\{a, b\} = a \wedge b$ exists in L
 & $a, b \in L$
- ② $\text{lub}\{a, b\} = \text{Sup}\{a, b\} = a \vee b$ exists in L
 & $a, b \in L$

Ex) ① Let $P(S)$ be the power set of S

Then $(P(S), \subseteq)$ is a POSET

$$P(S) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \dots\}$$

\swarrow All subsets of S

A B

$$\text{Let } A \vee B = A \cup B = \text{Sup}\{A, B\} \in P(S)$$

$$A \wedge B = A \cap B = \text{Inf}\{A, B\} \in P(S)$$

$A \subseteq A \cup B \Rightarrow A \leq A \cup B$
$B \subseteq A \cup B \Rightarrow B \leq A \cup B$

$\therefore P(S)$ is a lattice

Ex Let L be a lattice then

① $a \vee b = b$ iff $a \leq b$

* ② $a \wedge b = b$ iff $a \geq b$

Date _____ / _____ / _____

$$a \leq b$$

$$b \leq b$$

$$\sup \{a, b\} \leq b$$

$$[a \vee b \leq b]$$

$$a \leq a \vee b$$

$$b \leq a \vee b$$

$$\sup \{a, b\} \geq a \vee b$$

$$a \leq b$$

$$b \leq b$$

$$\sup \{a, b\} \leq b$$

Saath!

Proof

Let $a \leq b$

we have $a \vee b = \sup \{a, b\} \geq \inf \{a, b\}$

$$\therefore a \leq a \vee b$$

and $b \leq a \vee b$ —① (By def of sup)

By Given cond.

$$a < b$$

$$b \leq b$$

$$\Rightarrow \sup \{a, b\} < b$$

$$\Rightarrow a \vee b \leq b$$
 —②

From ① & ②

$$[a \vee b = b]$$

$$a \geq b$$

$$a \leq b$$

$$\Rightarrow a = b$$

Ex)

$$(N, \leq)$$

↓
usual sense

Let $a, b \in N$

$$a \wedge b = \max \{a, b\}$$

$$a \wedge b = \min \{a, b\}$$

' \leq ' in usual sense

Is (N, \leq) a lattice?

$$(N, \leq)$$

① Reflexive

$$a \leq a \quad \forall a \in N$$

② Transitive

If $a \leq b$ and $b \leq c$

then $a \leq c$

③ Antisymmetric

If $a \leq b$ and $b \leq a$

$$\text{Then } a = b$$

$\Rightarrow \therefore (N, \leq)$ is POSET

Let $a, b \in \mathbb{N}$

$$a \wedge b = \max\{a, b\} \in \mathbb{N}$$

$$\max\{2, 3\} = 3 \in \mathbb{N}$$

$$a \wedge b = \min\{a, b\} \in \mathbb{N}$$

$\therefore (\mathbb{N}, \leq)$ is a lattice.

(Q) (\mathbb{N}, \mid) is a lattice
divisibility

Ans) $a \leq b \text{ iff } a/b \in \mathbb{Z}$

① Reflexive

let we have

$$a/a \in \mathbb{Z} \text{ and } a \in \mathbb{N}$$

$$\therefore a \leq a \text{ and } a \in \mathbb{N}$$

$\therefore \leq$ is Reflexive

② Antisymmetric

We have

$$a \leq b \text{ and } b \leq a$$

$$\Rightarrow a/b \text{ and } b/a$$

which is possible iff $a = b$

$\therefore \leq$ is antisymmetric

③ Transitive

Let $a \leq b$ and $b \leq c$ and $a, b, c \in \mathbb{N}$

$$\therefore a/b \text{ and } b/c$$

$$\Rightarrow a/c$$

$$\Rightarrow a \leq c \quad \therefore \leq \text{ is transitive}$$

Date / /

 $\therefore (N, \leq)$ is poset

$$2 \leq 5 \Rightarrow 2/5 \times$$

$$\sup\{a, b\} = a \vee b \Rightarrow 5 \leq 5 \Rightarrow 5/5 \checkmark$$

$$\inf\{a, b\} = a \wedge b \Rightarrow \begin{matrix} 2 \leq 2 \Rightarrow 2/2 \checkmark \\ 5 \leq 2 \Rightarrow 5/2 \times \end{matrix}$$

$$\sup\{a, b\} = \text{lcm}\{a, b\} \in N$$

$$\inf\{a, b\} = \text{gcd}\{a, b\} \in N$$

$$\sup\{2, 5\} = 2 \vee 5 = \text{lcm}\{2, 5\} = 10 \in N$$

$$2 \leq 10 \Rightarrow 2/10 -$$

$$5 \leq 10 \Rightarrow 5/10 -$$

$$\inf\{2, 5\} = \text{gcd}\{2, 5\} = 1 \in N$$

$$1 \leq 2 \Rightarrow 1/2 -$$

$$1 \leq 5 \Rightarrow 1/5 -$$

 $\therefore (N, \leq)$ is a Lattice

→ Properties of Lattice :-

Let (L, \leq, \vee, \wedge) be a lattice then following properties are satisfied:

(1) Idempotent Property:

$$a \vee a = a \quad | \quad a \wedge a = a \quad \# \quad a \in L$$

(2) Absorption Law:

$$a \vee (a \wedge b) = a \quad | \quad a \wedge (a \vee b) = a$$

↓
b is absorbed

(1) Associative :

$$a \vee (a \vee c) = (a \vee b) \vee c \quad | \quad a \wedge (b \wedge c) = (a \wedge b) \wedge c$$

(2) Commutative :

$$a \vee b = b \vee a \text{ and } a \wedge b = b \wedge a \quad \forall a, b \in L$$

(3) Distributive law follows from Absorption Law.

(4) From the absorption law, we have

$$a \vee (a \wedge b) = a \quad \text{--- (1) } \forall a, b \in L$$

If $a, b \in L$ then $a \vee b \in L$ ∴ In (1) we can replace b by $a \vee b$

$$a \vee (a \wedge (a \vee b)) = a$$

$$a \vee a = a \quad \forall a \in L \quad (\because a \wedge (a \vee b) = a \text{ using absorption law})$$

Similarly $a \wedge a = a \quad \forall a \in L$

∴ Idempotent Law follows from absorption law

(5) Let $(L_1, \leq, \wedge, \vee)$ and $(L_2, \leq, \wedge, \vee)$ be two lattices.Then define $L_1 \times L_2$ is also a lattice① $L_1 \times L_2$ is a POSET② $\text{Sup } \{(a_1, b_1), (a_2, b_2)\} = \{a_1 \vee a_2, b_1 \vee b_2\}$ ③ $\text{Inf } \{(a_1, b_1), (a_2, b_2)\} = \{a_1 \wedge a_2, b_1 \wedge b_2\}$

Date / /

→ Another definition of Lattice :

A non empty set L with two binary operations ' \vee ' and ' \wedge ' is said to be a lattice if it satisfies the following properties:

① Commutative Property :

$$\begin{aligned} a \vee b &= b \vee a \\ a \wedge b &= b \wedge a \quad \forall a, b \in L \end{aligned}$$

② Associative Property :

$$\begin{aligned} (a \vee b) \vee c &= a \vee (b \vee c) \\ \text{and } (a \wedge b) \wedge c &= a \wedge (b \wedge c) \quad \forall a, b, c \in L \end{aligned}$$


③ Absorption Law :-

$$a \vee (a \wedge b) = a, \quad a \wedge (a \vee b) = a$$

$$\quad \forall a, b \in L$$

→ Sub lattice : Let L be a lattice, Then a subset S of L is said to be a sub lattice of L .

if $a \vee b \in S$ & $a \wedge b \in S$
and $a \vee b \in S$ & $a \wedge b \in S$



Example)Let $X = \{a, b, c\}$,
 $P(X) = \text{power set} = \{\emptyset, \{a\}, \{b\}, \{c\}, \{ab\}, \{bc\}, \{ac\}, \{abc\}\}$
 $(P(X), \cup, \cap)$ is a lattice.Take $\vee \rightarrow \cup$ $\wedge \rightarrow \cap$ ① Let $A, B \in P(X)$ then

$$A \cup B \in P(X)$$

$$A \cap B \in P(X)$$

 $\therefore \cup, \cap$ are Binary Operations on $P(X)$ ② Commutative :

$$A \cup B = B \cup A, A \cap B = B \cap A$$

$$\forall A, B \in P(X)$$

③ Associativity :

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

④ Absorption :

$$A \cup (A \cap B) = A$$

$$A \cap (A \cup B) = A$$

 \therefore Absorption is satisfied $\therefore (P(X), \cup, \cap)$ is a lattice.

Now, Take $S = \{\emptyset, x, f_a y, f_b y, f_{ab} y\}$

\cup	$\emptyset \times f_a y \quad f_b y \quad f_{ab} y$
\emptyset	$\emptyset \times f_a y \quad f_b y \quad f_{ab} y$
x	$x \quad x \quad x \quad x \quad x$
$f_a y$	$f_a y \times f_a y \quad f_b y \quad f_{ab} y$
$f_b y$	$f_b y \quad x \quad f_{ab} y \quad f_b y \quad f_{ab} y$
$f_{ab} y$	$f_{ab} y \quad x \quad f_{ab} y \quad f_{ab} y \quad f_{ab} y$

$\therefore A \cup B \in S$ for $A, B \in S$

and $A \cap B \in S$ & $A, B \in S$

$\therefore S$ is a sublattice of $P(X)$

Now, Take $S_1 = \{\emptyset, x, f_a y, f_b y, f_{ab} y\}$

when we take $\neg f_{ab} y \notin S_1$
Hence it is ^{union} _{not} a sub lattice

Qx → Lattice Homomorphism :-
 \downarrow
 function

Let L_1 and L_2 be two lattices then
 a function $f: L_1 \rightarrow L_2$ is said to be
 a homomorphism if two properties are
 satisfied.

$$\text{① } f(a \vee b) = f(a) \vee f(b)$$


$$\text{② } f(a \wedge b) = f(a) \wedge f(b)$$

$\forall a, b \in L_1$

Also, the lattices L_1 & L_2 are said to be Homomorphic

Date _____

Saathi

 L_1 & L_2 are homomorphic f is a lattice homomorphism→ Lattice Isomorphism: A Homomorphism $f: L_1 \rightarrow L_2$ issaid to be an isomorphism if f is one-one and onto.

Structure 1



Structure - 2



Q1) Give two examples of lattice Homomorphism / Isomorphism

Q2) Give two examples of lattices which are not
@ HomomorphicQ3) Give ^{one} example of two lattices which are homomorphic
but not isomorphic→ Bounded Lattice: A lattice L is said to be boundedif $\exists l \in L$ such that $l \leq x \forall x \in L$ and
 $\exists u \in L$ such that $u \geq x \forall x \in L$.

(P(S), ⊂, ∩)

we have $\emptyset \in P(S)$

st

 $\emptyset \subset A \wedge A \in P(S)$ ($\because \emptyset \subseteq A$)

Date ___ / ___ / ___

we have $S \in P(S)$ \vdash $A \subseteq S \rightarrow A \in P(S)$

→ Boolean Algebra:

Let $B \neq \emptyset$ and ' $+$ ', ' $*$ ' be two binary operations on B . Let ' 1 ' is a unary operation on B . Also, B has at least two elements ' 0 ' and ' 1 '. Then, B is said to be a Boolean Algebra if it satisfies following axioms.

B_1 : Commutative law:

$$a+b = b+a, \quad a*b = b*a \quad \forall a, b \in B$$

B_2 : Identity law:

$$a+0 = a, \quad a*1 = a \quad \forall a \in B$$

B_3 : Distributive law:

$$a*(b+c) = a*b + a*c, \quad a+(b*c) = (a+b) * (a+c) \quad \forall a, b, c \in B$$

B_4 : Complement law:

$$a+a' = 1, \quad a*a' = 0 \quad \forall a \in B$$

0 stands for identity element wrt $+$ → does not mean usual plus

1 stands for identity element wrt $*$ → does not mean usual multiplication

$0 \rightarrow " " \text{ zero.}$

$1 \rightarrow " " \text{ one.}$

Date / /

$$S \neq \emptyset \rightarrow P(S)$$

$$\emptyset, S \rightarrow \text{universal set}$$

$$A \cup B \in P(S) \quad \forall A, B \in P(S)$$


$$A \cap B \in P(S) \quad \forall A, B \in P(S)$$

\cup, \cap are binary operation.

$$A \in P(S)$$

$$\rightarrow A' \in P(S)$$

C10C



$$\text{B}_1: A + B = A \cup B = B \cup A = B + A$$

$$\Rightarrow A + B = B + A \quad \forall A, B \in P(S)$$

$$\text{B}_2: A \in P(S)$$

$$A + \emptyset = A \cup \emptyset = A$$

$$A + S = A \cap (S) = A$$

Similarly B_3, B_4 \checkmark
 $\therefore P(S)$ is Boolean Algebra

Notation: $B(1, +, ', \cap, \cup)$

$$\text{eg } P(S)(U, 0, 1, \emptyset, S)$$

$$\text{Ex. } D_{14} = \{1, 2, 3, 14\}$$

$\uparrow \rightarrow \text{LCM}$

$\downarrow \rightarrow \text{GCD}$

$$o' = \frac{14}{2}$$

Date _____

Saathi

lcm | 1 2 7 14

1	(1)	2	7	14	?	all the elements
2	(2)	2	14	14	?	96 98
7	(7)	14	7	14		
14	(14)	14	14	14		

gcd | 1 2 7 14

1	1	1	1	(1)	?	all the elements of D_{14}
2	1	2	1	(2)		
7	1	1	7	(7)		
14	1	2	7	(14)		

$a = \frac{14}{1}$ | 1 2 7 14

$$B_1 : \text{lcm}(a, b) = \text{lcm}(b, a) \quad \forall a, b \in D_{14}$$

$$B_2 : \text{gcd}(a, b) = \text{gcd}(b, a) \quad \forall a, b \in D_{14}$$

$$B_3 : a \in D_{14}$$

$$a + (1) = a(1, 1) = a$$

$$a + (14) = \text{gcd}(a, 14) = a$$

$$0 \rightarrow 1$$

$$1 \rightarrow 14$$

$$D = \{1, 3, 9\}$$

Saathi

Date _____

B.

Complement

$$a + (14) = \text{lcm}(a, 14) = 14$$

$$a + (1) = \text{gcd}(a, 1) = 1$$

(Q)

D_{35} is a Boolean Algebra.

(Q)

Let D_n be the set of all divisors of n .
Find the least value of n st D_n is not a Boolean Algebra. ($n=3$)

(Q)

$B = \{0, 1\}$ is the smallest Boolean Algebra

→ Properties of Boolean Algebra:

Let B be a Boolean algebra, then elements of B satisfies the following properties:

① Idempotent Law:

$$a + a = a \quad \text{and} \quad a \cdot a = a \quad \forall a \in B$$

② Boundary Law:

$$a + 1 = 1 \quad \text{and} \quad a \cdot 0 = 0 \quad \forall a \in B$$

③ Absorption Law:

$$a + (a \cdot b) = a \quad \text{and} \quad a \cdot (a + b) = a \\ \forall a, b \in B$$

④ Associative Law:

$$a + (b + c) = (a + b) + c$$

Date _____ / _____ / _____

$$\text{and } a * (b + c) = (a * b) + c$$

$\forall a, b, c \in B$

Ex 1) Let $B (+, *, ', 0, 1)$ be a Boolean Algebra. Then
 $0' = 1$ and $1' = 0$

Then, let $B (+, *, ', 0, 1)$ be a Boolean algebra,
then

$$(i) (a + b)' = a' + b' \quad \forall a, b \in B$$

$$(ii) (a * b)' = a' * b' \quad \forall a, b \in B$$

To prove

Proof. (i) $(a + b)' = a' + b'$

We know

$$a + a' = 1$$

$$a + a' = 0$$

Identity Complement

↓ ↓

Commutative

Distributive

$$(a + b) * (a' + b') = 0$$

Proof:

$$\Rightarrow a + (a' + b) + b * (a' + b') \quad \because (\text{Distributive law})$$

$$\Rightarrow (a + a') + b + b * (a' + b') \quad \because (\text{Associative law})$$

$$\Rightarrow (a + a') + b' + b * (b' + a') \quad \because (\text{Commutative law})$$

$$\Rightarrow (a + a') + b' + (b * b') + a' \quad \because (\text{Associative law})$$

$$\Rightarrow \overbrace{0 + b'} + \overbrace{0 + a'} \quad \because (\text{Complement law})$$

$$\Rightarrow 0 + 0 \quad \because (\text{Boolean law})$$

$$\Rightarrow 0$$

 $\therefore Q.E.D.$

$$(a + b) * (a' + b') = 0 \quad \text{--- Q.E.D.}$$

Date _____

$$(a+b)' + (a'+b') = 1$$

$$\begin{aligned} &\Rightarrow (a+b+a') + (a+b+b') \quad (\because \text{Distributive law}) \\ &\Rightarrow (1+b) + (1+a) \quad (\because \text{Complement law}) \\ &\Rightarrow 1 + 1 \quad (\because \text{Boolean law}) \end{aligned}$$

$$(a+b)' + (a'+b') = 1 \quad \text{--- (1)}$$

*"Proved"*Using $a \wedge b \Leftrightarrow$ we have

$$(a+b)' = a' \cdot b'$$

By principle of duality
 $(a+b)' = a' \cdot b'$ \therefore De Morgan's law hold in Boolean Algebra

$$1' = 0 \quad \text{and} \quad 0' = 1$$

We have

$$a + a' = 1$$

$$\text{i.e. } 1 = a + a'$$

Taking complement on both sides

$$1' = (a + a')'$$

$$1' = a' \cdot (a')'$$

$$1' = a' \cdot a$$

$$\Rightarrow 1' = 0$$

"Proved"

We have

$$a + a' = 0$$

$$\text{i.e. } 0 = a + a'$$

Taking complement on both sides

$$0' = (a + a')'$$

$$0' = a' \cdot (a')'$$

$$0' = a' \cdot a$$

$$0' = 1 \quad \text{"Proved"}$$

Date _____


→ Boolean Homomorphism:

Let B_1 and B_2 be two Boolean Algebras. Then a mapping $f: B_1 \rightarrow B_2$ is said to be Boolean Homomorphism if

- (i) $f(a \wedge b) = f(a) \wedge f(b)$ & $a, b \in B_1$,
- (ii) $f(a \vee b) = f(a) \vee f(b)$ & $a, b \in B_1$,
- (iii) $f(\bar{a}) = [\bar{f(a)}]'$ & $a \in B_1$.

If the Boolean homomorphism f is 1-1 and onto, then it is called a Boolean isomorphism.

Ex 1) $B_1 = \{1, 2, 3, 4\}$ (Divisors of 12)
 and $B_2 = \{a, b, c, d\}$ (Powerset of $\{a, b\}$)
 Define $f: B_1 \rightarrow B_2$.
 one-to-one & onto.



Now

$$\begin{aligned}
 f(1') &= f(4) = \{a, b\} & f(1 \wedge 2) &= f(2) \\
 f(1) &= \{a\} & \text{and } f(2) &= \{a\} \\
 (f(1))' &= \{\emptyset\} = \{a, b\}' & f(1) \wedge f(2) &= \emptyset + \{a\} \\
 & & &= \emptyset \cup \{a\} \\
 \therefore f &\text{ is Boolean isomorphism} & \therefore f(1 \wedge 2) &= f(1) \wedge f(2)
 \end{aligned}$$

Date _____

(Q)

Give an example of Boolean algebra which are not isomorphic and explain the reason.

(Q)

Let B be a Boolean algebra. Then show that following conditions are equivalent

$$\textcircled{1} \quad a + b = b \quad (\text{ii}) \Rightarrow (\text{iii})$$

$$\textcircled{2} \quad a + b = a \quad (\text{iii}) \Rightarrow (\text{ii}),$$

$$\textcircled{3} \quad a' + b = 1 \quad (\text{iii}), \Rightarrow (\text{iv}),$$

$$\textcircled{4} \quad a' + b' = 0 \quad (\text{iv}) \Rightarrow (\text{ii})$$

Assignment - 1

(Q) \rightarrow Boolean algebra cannot have 3 distinct elements

(Q) A Boolean " " "

(Q) Let B be a Boolean algebra. Then prove the following:

$$\textcircled{1} \quad (a+b)' = a' + b'$$

$$\textcircled{2} \quad (a+b) + (a+c) + (b+c) = a+c = a+b + b+c$$

$$\textcircled{3} \quad (a+b)(b+c)(c+a) = ab + bc + ca$$

$$\textcircled{4} \quad (a+b')(b+c')(c+a') = (a+b')(b'+c')(c+a')$$

(Q) Let L_1 and L_2 be two Lattices, then show

$L_1 \cap L_2$ is also a sub-lattice of L .

$L \cup L_2$ is also a sub-lattice? Justify.

(Q) Let B_1 and B_2 be two sub-algebras of a Boolean algebra B . Then show that $B_1 \cap B_2$ is also a sub-algebra. What about $B_1 \cup B_2$?

(Q) Explain the application of Boolean operations in simplification of switching circuits.

- ④ Let $(G, +)$ be a group non-empty set and $a+b = ab/2 \forall a, b \in G$
then show that G is a group w.r.t operator \oplus
- ⑤ Give an example of two non-isomorphic Boolean algebra
- ⑥ What is Hasse diagram? Give two examples
- ⑦ Give an example of a relation which is not reflexive

Sub-algebra: Let R be a Boolean algebra and $S \subseteq R$.

Theo S is said to be a sub-algebra of R if

$$a+b \in S \quad \forall a, b \in S$$

$$a \cdot b \in S \quad \forall a, b \in S$$

$$a \in S \Rightarrow a' \in S \quad \forall a \in S$$

$$\subseteq R \subseteq S$$

$$S_1, S_2$$

$$S_1 \cap S_2$$

Let a, b be two elements of $S_1 \cap S_2$,

Since $a \in S_1$ & $b \in S_1$ and $a, b \in S_2$

Since S_1 & S_2 are sub-algebras of R

$$a+b \in S_1 \Rightarrow a+b \in S_2$$

$$\Rightarrow a+b \in S_1 \cap S_2$$

$$(iii) a \cdot b \in S_1 \cap S_2$$

$$\Rightarrow a \in S_1 \text{ & } b \in S_2$$

$$\Rightarrow a' \in S_1 \text{ & } b' \in S_2$$

$$\Rightarrow a' \cdot b' \in S_1 \cap S_2$$