


A Journey Through Moduli Spaces

Part 1: Foundations for Understanding Moduli Spaces

Prepared by: **Anuj Sharma**

Reading Project

Date: November 25, 2024



*“Mathematics is the music of reason.” – James Joseph
Sylvester*

Abstract

The first part of this reading project explores the foundations necessary to understand moduli spaces. We begin with the building blocks of algebraic geometry, focusing on concepts such as algebraic curves, elliptic curves, hyperelliptic curves, and the genus—a measure of a curve’s complexity. Further, we delve into rational maps, unirationality, and Jacobians, which are essential tools for studying the structure of moduli spaces. This section aims to provide a comprehensive foundation, enabling readers to engage with the advanced topics in later parts.

Contents

1	Introduction	3
2	Building Blocks of Algebraic Geometry	3
2.1	Algebraic Curves	3
2.2	Elliptic Curves	3
2.3	Hyperelliptic Curves	3
3	Key Concepts for Moduli Spaces	4
3.1	Genus	4
3.2	Rational Maps and Unirationality	4
3.3	Jacobians	4
4	Conclusion	4
5	Foundations of Moduli Spaces	4
5.1	Algebraic Curves	4
5.2	Rational Maps and Morphisms	5
5.3	Unirationality	5
5.4	Elliptic Curves	5
5.5	Hyperelliptic Curves	6
5.6	Jacobians of Curves	6
5.7	Moduli Spaces as Classifiers	6
6	Foundations of Moduli Spaces	6
6.1	Algebraic Curves	7
6.2	Rational Maps and Morphisms	7
6.3	Unirationality	7
6.4	Elliptic Curves	8
6.5	Hyperelliptic Curves	8
6.6	Jacobians of Curves	8
6.7	Moduli Spaces as Classifiers	8
7	Exploring Moduli Spaces of Small Genus	9
7.1	The Moduli Space \mathcal{M}_1	9

7.2	The Moduli Space \mathcal{M}_2	9
7.3	The Moduli Space \mathcal{M}_3	10
7.4	Higher Genus Moduli Spaces	10
7.5	Exploring Large Genus: \mathcal{M}_{23}	10
7.6	Compactification and Stable Curves	11
7.7	Applications of Small Genus Moduli Spaces	11
8	Exploring the Moduli Space \mathcal{M}_{23}	11
8.1	Properties of \mathcal{M}_{23}	11
8.2	Significance of \mathcal{M}_{23}	12
8.3	An Intuitive Analogy	12
8.4	Insights from Gavril Farkas' Work	12
8.5	Conclusion	13
8.6	Further Reading	13
8.7	Further Reading	13

1 Introduction

The concept of *moduli spaces* is central to modern algebraic geometry, encapsulating the idea of classifying mathematical objects based on their intrinsic properties. This reading project begins by introducing the fundamental building blocks required to understand these spaces. The discussion is rooted in the theory of algebraic curves, with an emphasis on elliptic and hyperelliptic curves, their genus, and associated invariants.

Moduli spaces arise naturally when one seeks to understand how these curves vary within families. For instance, the moduli space of genus 1 curves, denoted \mathcal{M}_1 , provides a framework to study elliptic curves up to isomorphism. To appreciate the depth and utility of such spaces, one must first grasp foundational concepts like rational maps, unirationality, and Jacobians.

2 Building Blocks of Algebraic Geometry

2.1 Algebraic Curves

An *algebraic curve* is a one-dimensional variety defined as the set of zeros of a polynomial equation $f(x, y) = 0$ over an algebraically closed field k . These curves can be classified by their **genus** g , which serves as a topological measure of their complexity. For example:

- A genus 0 curve is birationally equivalent to \mathbb{P}^1 , the projective line.
- A genus 1 curve is an elliptic curve, provided it has a rational point.
- Higher-genus curves are more intricate and can exhibit hyperelliptic properties.

2.2 Elliptic Curves

Elliptic curves are smooth projective curves of genus 1, typically expressed in Weierstrass form:

$$y^2 = x^3 + ax + b, \quad \text{where } \Delta = 4a^3 + 27b^2 \neq 0.$$

The condition $\Delta \neq 0$ ensures non-singularity. Elliptic curves play a pivotal role in various fields, from cryptography to number theory.

2.3 Hyperelliptic Curves

A *hyperelliptic curve* is a smooth projective curve of genus $g \geq 2$ that admits a double cover of \mathbb{P}^1 . It can be expressed as:

$$y^2 = f(x),$$

where $f(x)$ is a polynomial of degree $2g + 1$ or $2g + 2$. These curves generalize elliptic curves and are of particular interest in the study of higher-genus moduli spaces.

3 Key Concepts for Moduli Spaces

3.1 Genus

The genus g of an algebraic curve is a topological invariant representing the number of "holes" in its surface. It is calculated using the Riemann-Hurwitz formula or derived from the dimension of the space of holomorphic differentials:

$$g = \dim H^0(C, \Omega_C^1).$$

3.2 Rational Maps and Unirationality

A *rational map* between two varieties X and Y is a morphism defined by rational functions. A variety is said to be *unirational* if there exists a dominant rational map from a projective space \mathbb{P}^n to the variety. For instance, \mathcal{M}_1 is unirational.

3.3 Jacobians

The Jacobian $J(C)$ of a curve C is an abelian variety that parametrizes degree-zero divisors modulo linear equivalence. It plays a crucial role in the study of moduli spaces, serving as a bridge between the geometry of curves and their function spaces.

4 Conclusion

This foundational exploration sets the stage for understanding the moduli spaces \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{M}_3 in the subsequent parts. By delving into the intricate relationship between algebraic curves, their genus, and associated invariants, we prepare to navigate the more advanced landscapes of algebraic geometry.

5 Foundations of Moduli Spaces

Moduli spaces are central objects in geometry and topology that serve as classifiers for families of geometric or algebraic structures. This section delves into the foundational concepts necessary to understand moduli spaces, with examples to illustrate the underlying principles.

5.1 Algebraic Curves

An **algebraic curve** is the zero set of a polynomial $f(x, y) = 0$ in two variables, typically studied over an algebraically closed field such as \mathbb{C} . These curves vary in complexity, which is often measured by their *genus*.

Example: The equation

$$y^2 = x^3 - x$$

defines an algebraic curve. When visualized, this curve has a toroidal topology, signifying its genus is $g = 1$.

The genus g is a key invariant, defined topologically as the number of "holes" in the surface associated with the curve. For projective smooth curves, it is calculated as:

$$g = \frac{(d-1)(d-2)}{2},$$

where d is the degree of the curve.

Classification by Genus:

- **Genus 0:** Curves that are topologically equivalent to the Riemann sphere \mathbb{P}^1 , such as $y = mx + c$.
- **Genus 1:** Elliptic curves, such as $y^2 = x^3 + ax + b$, which are smooth curves with one hole.
- **Genus $g \geq 2$:** Higher genus curves, such as hyperelliptic curves, which exhibit increasing complexity.

5.2 Rational Maps and Morphisms

A **rational map** between two varieties X and Y is a map defined by rational functions. Rational maps allow us to compare curves and classify them up to *birational equivalence*.

Example: Consider the map

$$\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1 \quad \text{defined by} \quad \phi(t) = \frac{t^2 + 1}{t - 1}.$$

This map demonstrates a birational equivalence between two varieties.

5.3 Unirationality

A space is **unirational** if there exists a dominant rational map from \mathbb{P}^n to the space. For instance, the moduli space \mathcal{M}_1 of elliptic curves is unirational because all elliptic curves can be parameterized by their j -invariant.

5.4 Elliptic Curves

Elliptic curves are genus-1 smooth projective curves with a specified base point O . They are central in the study of moduli spaces due to their well-understood properties and group structure.

Weierstrass Form: The general equation for an elliptic curve is:

$$y^2 = x^3 + ax + b, \quad \text{where } \Delta = 4a^3 + 27b^2 \neq 0.$$

The discriminant Δ ensures smoothness.

Example: j -Invariant: Elliptic curves are classified by their j -invariant:

$$j = \frac{1728 \cdot 4a^3}{\Delta}.$$

Curves with the same j -invariant are isomorphic.

5.5 Hyperelliptic Curves

A **hyperelliptic curve** is a smooth projective curve of genus $g \geq 2$, defined as a double cover of \mathbb{P}^1 . These curves generalize elliptic curves to higher genera.

Example: The curve

$$y^2 = x^5 - x + 1$$

is hyperelliptic with genus $g = 2$, calculated as:

$$g = \left\lfloor \frac{\deg(f) - 1}{2} \right\rfloor.$$

5.6 Jacobians of Curves

The **Jacobian** $J(C)$ of a curve C is an abelian variety representing degree-zero divisors modulo linear equivalence. Jacobians are critical in understanding moduli spaces, as they enable the parametrization of families of curves.

Example: For the elliptic curve $y^2 = x^3 - x$, the Jacobian is the curve itself, viewed as a group under point addition.

5.7 Moduli Spaces as Classifiers

The moduli space \mathcal{M}_g classifies all genus g curves up to isomorphism. Each point in \mathcal{M}_g corresponds to an equivalence class of curves.

Visualization of \mathcal{M}_1 : - A point in \mathcal{M}_1 represents an isomorphism class of elliptic curves.
- The j -invariant serves as a coordinate for \mathcal{M}_1 .

Example: Consider the family:

$$y^2 = x^3 + tx + t, \quad t \in \mathbb{C}.$$

For different values of t , the curves vary, but \mathcal{M}_1 organizes them by isomorphism equivalence.

6 Foundations of Moduli Spaces

Moduli spaces are central objects in geometry and topology that serve as classifiers for families of geometric or algebraic structures. This section delves into the foundational concepts necessary to understand moduli spaces, with examples to illustrate the underlying principles.

6.1 Algebraic Curves

An **algebraic curve** is the zero set of a polynomial $f(x, y) = 0$ in two variables, typically studied over an algebraically closed field such as \mathbb{C} . These curves vary in complexity, which is often measured by their *genus*.

Example: The equation

$$y^2 = x^3 - x$$

defines an algebraic curve. When visualized, this curve has a toroidal topology, signifying its genus is $g = 1$.

The genus g is a key invariant, defined topologically as the number of "holes" in the surface associated with the curve. For projective smooth curves, it is calculated as:

$$g = \frac{(d-1)(d-2)}{2},$$

where d is the degree of the curve.

Classification by Genus:

- **Genus 0:** Curves that are topologically equivalent to the Riemann sphere \mathbb{P}^1 , such as $y = mx + c$.
- **Genus 1:** Elliptic curves, such as $y^2 = x^3 + ax + b$, which are smooth curves with one hole.
- **Genus $g \geq 2$:** Higher genus curves, such as hyperelliptic curves, which exhibit increasing complexity.

6.2 Rational Maps and Morphisms

A **rational map** between two varieties X and Y is a map defined by rational functions. Rational maps allow us to compare curves and classify them up to *birational equivalence*.

Example: Consider the map

$$\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1 \quad \text{defined by} \quad \phi(t) = \frac{t^2 + 1}{t - 1}.$$

This map demonstrates a birational equivalence between two varieties.

6.3 Unirationality

A space is **unirational** if there exists a dominant rational map from \mathbb{P}^n to the space. For instance, the moduli space \mathcal{M}_1 of elliptic curves is unirational because all elliptic curves can be parameterized by their j -invariant.

6.4 Elliptic Curves

Elliptic curves are genus-1 smooth projective curves with a specified base point O . They are central in the study of moduli spaces due to their well-understood properties and group structure.

Weierstrass Form: The general equation for an elliptic curve is:

$$y^2 = x^3 + ax + b, \quad \text{where } \Delta = 4a^3 + 27b^2 \neq 0.$$

The discriminant Δ ensures smoothness.

Example: j -Invariant: Elliptic curves are classified by their j -invariant:

$$j = \frac{1728 \cdot 4a^3}{\Delta}.$$

Curves with the same j -invariant are isomorphic.

6.5 Hyperelliptic Curves

A **hyperelliptic curve** is a smooth projective curve of genus $g \geq 2$, defined as a double cover of \mathbb{P}^1 . These curves generalize elliptic curves to higher genera.

Example: The curve

$$y^2 = x^5 - x + 1$$

is hyperelliptic with genus $g = 2$, calculated as:

$$g = \left\lfloor \frac{\deg(f) - 1}{2} \right\rfloor.$$

6.6 Jacobians of Curves

The **Jacobian** $J(C)$ of a curve C is an abelian variety representing degree-zero divisors modulo linear equivalence. Jacobians are critical in understanding moduli spaces, as they enable the parametrization of families of curves.

Example: For the elliptic curve $y^2 = x^3 - x$, the Jacobian is the curve itself, viewed as a group under point addition.

6.7 Moduli Spaces as Classifiers

The moduli space \mathcal{M}_g classifies all genus g curves up to isomorphism. Each point in \mathcal{M}_g corresponds to an equivalence class of curves.

Visualization of \mathcal{M}_1 : - A point in \mathcal{M}_1 represents an isomorphism class of elliptic curves.
- The j -invariant serves as a coordinate for \mathcal{M}_1 .

Example: Consider the family:

$$y^2 = x^3 + tx + t, \quad t \in \mathbb{C}.$$

For different values of t , the curves vary, but \mathcal{M}_1 organizes them by isomorphism equivalence.

7 Exploring Moduli Spaces of Small Genus

Moduli spaces of curves provide a geometric framework for classifying algebraic curves of fixed genus g . In this section, we explore the properties and significance of moduli spaces \mathcal{M}_g for small values of g , focusing on their structure, dimension, and compactifications.

7.1 The Moduli Space \mathcal{M}_1

The moduli space \mathcal{M}_1 represents isomorphism classes of elliptic curves. Its rich structure stems from the connection between elliptic curves and modular forms.

Dimension: \mathcal{M}_1 is a 1-dimensional space, as each elliptic curve is determined up to isomorphism by a single parameter, the j -invariant.

Parameterization: The j -invariant is defined as:

$$j(E) = \frac{1728g_2^3}{\Delta}, \quad \text{where } g_2 = 4a^3 \text{ and } \Delta = 4a^3 + 27b^2.$$

Here, $E : y^2 = x^3 + ax + b$ is a Weierstrass equation of an elliptic curve.

Properties: - \mathcal{M}_1 is isomorphic to the affine line \mathbb{A}^1 . - It has a natural compactification, $\overline{\mathcal{M}}_1$, obtained by adding a point at infinity corresponding to the nodal cubic curve.

Applications: Elliptic curves are critical in cryptography, number theory, and complex geometry. The structure of \mathcal{M}_1 forms the foundation for modular forms and modular curves.

7.2 The Moduli Space \mathcal{M}_2

\mathcal{M}_2 classifies smooth projective curves of genus 2. These curves are hyperelliptic and can be described as double covers of \mathbb{P}^1 , branched at six points.

Dimension: \mathcal{M}_2 is 3-dimensional, as the configuration of six branch points modulo projective transformations determines the curve.

Coordinate Description: Hyperelliptic genus-2 curves are described by the equation:

$$y^2 = f(x), \quad \deg(f) = 6.$$

The moduli space \mathcal{M}_2 is thus associated with the space of degree-6 polynomials modulo the action of $\text{PGL}(2, \mathbb{C})$.

Properties: - \mathcal{M}_2 is a smooth quasi-projective variety. - The Torelli theorem relates \mathcal{M}_2 to the Siegel modular variety \mathcal{A}_2 , via the Jacobian of genus-2 curves.

Compactification: The Deligne-Mumford compactification $\overline{\mathcal{M}}_2$ includes stable curves, allowing for a complete moduli space.

7.3 The Moduli Space \mathcal{M}_3

\mathcal{M}_3 corresponds to smooth projective curves of genus 3. Unlike genus 2, not all genus-3 curves are hyperelliptic; they can also be plane quartics.

Dimension: The dimension of \mathcal{M}_3 is 6, reflecting the higher complexity of genus-3 curves.

Hyperelliptic vs. Non-Hyperelliptic: - Hyperelliptic genus-3 curves are a proper subset of \mathcal{M}_3 , forming a divisor. - Non-hyperelliptic genus-3 curves are represented as plane quartics:

$$f(x, y, z) = 0, \quad \deg(f) = 4.$$

Compactification and Properties: - \mathcal{M}_3 is connected, smooth, and quasi-projective. - The Jacobians of genus-3 curves embed \mathcal{M}_3 into \mathcal{A}_3 , the Siegel modular variety of dimension 6.

7.4 Higher Genus Moduli Spaces

\mathcal{M}_4 : - Dimension: 9. - Includes hyperelliptic curves as a codimension-1 locus. - Non-hyperelliptic curves are plane quintics.

\mathcal{M}_g for $g \geq 5$: - For $g \geq 5$, \mathcal{M}_g becomes increasingly complex, with dimension $3g - 3$. - The Kodaira dimension of \mathcal{M}_g provides insights into its birational geometry.

7.5 Exploring Large Genus: \mathcal{M}_{23}

The moduli space \mathcal{M}_{23} represents the threshold where the moduli spaces of curves are of general type. This marks a transition in the geometric properties of these spaces:

- Dimension: 66 (calculated as $3g - 3$).
- General type: \mathcal{M}_{23} and all higher genus moduli spaces have maximal Kodaira dimension.

Significance: - \mathcal{M}_{23} plays a key role in studying the birational classification of moduli spaces. - Its structure influences research in algebraic geometry, string theory, and mathematical physics.

7.6 Compactification and Stable Curves

The Deligne-Mumford compactification $\overline{\mathcal{M}}_g$ adds stable nodal curves to \mathcal{M}_g , ensuring a complete moduli space. Stable curves satisfy:

The arithmetic genus g remains constant, and every rational component intersects others at least 3 times.

Examples of Stable Curves: - For $g = 1$: The nodal cubic curve. - For $g = 2$: A genus-1 curve with a rational tail.

7.7 Applications of Small Genus Moduli Spaces

Moduli spaces of small genus have applications in enumerative geometry, string theory, and mathematical physics:

- \mathcal{M}_1 : Connections with modular forms and elliptic curves.
- \mathcal{M}_2 : Siegel modular varieties and Jacobians.
- \mathcal{M}_3 : Plane quartics and Prym varieties.
- \mathcal{M}_{23} : Study of moduli spaces as varieties of general type.

8 Exploring the Moduli Space \mathcal{M}_{23}

The moduli space \mathcal{M}_{23} represents the parameter space for smooth algebraic curves of genus 23. This particular space plays a pivotal role in the study of higher-dimensional algebraic geometry and moduli theory, marking a transition in the behavior and complexity of such spaces. In this section, we delve into the properties, significance, and intuitive understanding of \mathcal{M}_{23} .

8.1 Properties of \mathcal{M}_{23}

The space \mathcal{M}_{23} exhibits the following notable properties:

1. Kodaira Dimension:

- The Kodaira dimension of \mathcal{M}_{23} is known to be at least 2, suggesting that the space is of *general type*. This classification implies that \mathcal{M}_{23} has a richly intricate structure akin to higher-dimensional varieties in algebraic geometry.
- For moduli spaces of genus $g > 23$, \mathcal{M}_g is definitively of general type. However, for $g \leq 22$, these spaces are conjectured to be *uniruled*, meaning they can be covered by rational curves.

2. Boundary Divisors:

- \mathcal{M}_{23} contains boundary divisors representing degenerate or singular curves. These divisors play a critical role in compactifying the space, ensuring that it remains a complete and well-defined parameter space.

- These boundary components correspond to nodal curves (curves with specific singularities), which are fundamental in understanding the geometry of \mathcal{M}_{23} .

3. Multicanonical Divisors:

- The study of \mathcal{M}_{23} relies heavily on analyzing multicanonical divisors derived from Brill-Noether theory. These divisors provide constraints on embeddings and syzygies of curves, revealing deep geometric insights.

4. Transition Behavior:

- \mathcal{M}_{23} is a transitional space between lower-genus moduli spaces (which are typically unirational) and higher-genus moduli spaces (which are generally of general type).
- This dual nature makes \mathcal{M}_{23} a cornerstone for understanding the global geometry of moduli spaces.

8.2 Significance of \mathcal{M}_{23}

The moduli space \mathcal{M}_{23} is particularly important for the following reasons:

- **Bridge in Moduli Theory:** As the first moduli space of general type in the genus sequence, \mathcal{M}_{23} helps bridge our understanding of spaces that are uniruled and those of higher complexity.
- **Applications to Algebraic Geometry:** Insights from \mathcal{M}_{23} inform the classification of algebraic varieties, the behavior of canonical divisors, and the structure of high-dimensional varieties.
- **Interdisciplinary Relevance:** The tools developed to study \mathcal{M}_{23} , such as divisor theory and syzygies, have applications in string theory and mathematical physics.

8.3 An Intuitive Analogy

To visualize \mathcal{M}_{23} , imagine it as a **3D sculpture gallery**:

- Each sculpture in the gallery represents an algebraic curve, uniquely defined by its genus and other intrinsic properties.
- The **floorplan** of the gallery corresponds to the "map" of \mathcal{M}_{23} , organizing these sculptures by their parameters.
- The gallery also contains "**hidden corners**"—boundary divisors that describe singular or nodal curves. These regions are critical for compactifying \mathcal{M}_{23} and understanding its global geometry.

8.4 Insights from Gavril Farkas' Work

Research by Gavril Farkas has further illuminated the structure of \mathcal{M}_{23} :

- His analysis of multicanonical divisors demonstrates the role of syzygies in embedding genus 23 curves in projective space.

- The work reveals that \mathcal{M}_{23} balances between uniruled and general type behaviors, providing a model for understanding this critical transition in moduli spaces.

8.5 Conclusion

The moduli space \mathcal{M}_{23} stands as a pivotal object in algebraic geometry, bridging the study of simpler, unirational spaces with the complex, general type spaces of higher genus. Its properties, from the Kodaira dimension to boundary divisors, showcase the rich interplay between geometry, topology, and arithmetic. Understanding \mathcal{M}_{23} not only advances mathematical theory but also opens pathways for interdisciplinary applications, making it a cornerstone in the study of moduli spaces.

8.6 Further Reading

For a deeper understanding, refer to:

- R. Hartshorne, *Algebraic Geometry*.
- J. Silverman, *The Arithmetic of Elliptic Curves*.
- Joe Harris, *Moduli of Curves*.

8.7 Further Reading

For a deeper understanding, refer to:

- R. Hartshorne, *Algebraic Geometry*.
- J. Silverman, *The Arithmetic of Elliptic Curves*.
- Joe Harris, *Moduli of Curves*.

References

- R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, 1977.
- J. Silverman, *The Arithmetic of Elliptic Curves*, Springer, 2009.
- <https://mathworld.wolfram.com/ModuliSpace.html>