

Physics Short Notes

Detailed Revision: Unit I (Quantum) & Unit V (Nanotech) with Derivations


UNIT I : Quantum Computing

1. Bits vs Qubits & Quantum Entanglement

Bit: Classical unit of info. Exists strictly as a deterministic state of **0 or 1**. Implemented physically via voltage levels in transistors.

Qubit: Quantum unit of info. Can exist as 0, 1, or a **coherent superposition** of both simultaneously ($|\Psi\rangle = \alpha|0\rangle + \beta|1\rangle$). This allows quantum computers to process an exponential number of states (2^n) simultaneously.

Quantum Entanglement: A physical phenomenon where a pair of particles interact such that their quantum states are universally linked. Measuring the state of one instantly determines the state of the other ("spooky action at a distance").



Bloch Sphere representing a Qubit.

2. Wave-Particle Duality & Matter Waves

Concept: Radiation and matter exhibit both wave-like (interference, diffraction) and particle-like (photoelectric effect) properties depending on the experiment.

Davisson-Germer Experiment: Provided the first experimental proof of matter waves by firing a beam of electrons at a nickel crystal and observing a diffraction pattern similar to X-rays.

IMP Derivation: De-Broglie Wavelength of an Accelerated Electron

1. By De-Broglie's hypothesis, the wavelength of a moving particle is: $\lambda = \frac{h}{p}$
2. If an electron (charge e , mass m) is accelerated from rest through a potential difference V , the work done becomes its kinetic energy (K):

$$K = eV$$

3. Relating Kinetic Energy to momentum ($p = mv$):

$$K = \frac{1}{2}mv^2 = \frac{m^2v^2}{2m} = \frac{p^2}{2m} \implies p = \sqrt{2mK}$$

4. Substitute $K = eV$ into the momentum equation:

$$p = \sqrt{2meV}$$

5. Substitute p back into the De-Broglie equation:

$$\lambda = \frac{h}{\sqrt{2meV}}$$

6. Substituting standard constants ($h = 6.626 \times 10^{-34}$, $m = 9.1 \times 10^{-31}$, $e = 1.6 \times 10^{-19}$):

$$\lambda = \frac{12.27}{\sqrt{V}} \text{ \AA}$$

3. Wave Packets & Heisenberg Uncertainty Principle

Wave Packets: A single matter wave extends to infinity, meaning position is completely uncertain. To localize a particle, we superimpose many waves of slightly different wavelengths to form a "wave packet" that is localized in space.

Heisenberg Uncertainty Principle: It is fundamentally impossible to simultaneously determine both the exact position (x) and exact momentum (p) of a microscopic particle.

$$\Delta x \cdot \Delta p \geq \frac{\hbar}{2} \quad \text{or} \quad \Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

4. Schrödinger Wave Equation & Wave Function (Ψ)

Wave Function (Ψ): A complex mathematical function that contains all dynamical information about the system. Physically, $|\Psi|^2$ (Probability Density) gives the probability of finding the particle at a specific coordinate.

Normalization Condition: The particle must exist somewhere in the universe, so the total probability over all space must be 1 (100%).

$$\int_{-\infty}^{\infty} \Psi \Psi^* dV = 1$$

Time-Independent Equation (1D): Used for stationary states where energy does not change with time.

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2}(E - V)\psi = 0$$

Eigenvalues & Eigenfunctions: Solving the Schrödinger equation yields physically acceptable wave functions called *Eigenfunctions*, which correspond to specific, discrete energy levels called *Eigenvalues*.

5. Application: 1D Infinite Potential Well (Particle in a Box)

Concept: A particle of mass m trapped in a 1D box of length L with impenetrable walls ($V = \infty$ outside, $V = 0$ inside). Because $V = \infty$ outside, $\psi(x) = 0$ at the boundaries $x = 0$ and $x = L$.

IMP Derivation: Quantization of Energy in 1D Well

1. Inside the box ($V = 0$), the Schrödinger equation is:

$$\frac{d^2\psi}{dx^2} + \frac{2mE}{\hbar^2}\psi = 0$$

2. Let $k^2 = \frac{2mE}{\hbar^2}$. The general solution is a harmonic wave:

$$\psi(x) = A \sin(kx) + B \cos(kx)$$

3. Apply Boundary Condition 1 ($x = 0, \psi = 0$):

$$0 = A \sin(0) + B \cos(0) \implies B = 0$$

So, the equation reduces to $\psi(x) = A \sin(kx)$.

4. Apply Boundary Condition 2 ($x = L, \psi = 0$):

$$0 = A \sin(kL)$$

Since $A \neq 0$ (particle exists), $\sin(kL) = 0$, which implies $kL = n\pi$. Therefore, $k = \frac{n\pi}{L}$ (where $n = 1, 2, 3 \dots$).

5. Equate the two expressions for k^2 :


$$k^2 = \left(\frac{n\pi}{L}\right)^2 = \frac{n^2\pi^2}{L^2}$$

$$\frac{2mE}{\hbar^2} = \frac{n^2\pi^2}{L^2} \implies E = \frac{n^2\pi^2\hbar^2}{2mL^2}$$

6. Since $\hbar = \frac{h}{2\pi}$, substitute to get the final discrete energy levels:

$$E_n = \frac{n^2 h^2}{8mL^2}$$

(Energy is proportional to n^2 , proving that energy is quantized).



Quantized wave functions for a particle in a 1D box.

UNIT V : Nanotechnology

1. Concept of Nanotechnology & Bulk vs. Nanomaterials

Definition: Engineering and manipulating materials at the atomic or molecular scale (typically 1 to 100 nm). At this scale, classical physics gives way to quantum mechanics.

Massive Surface Area to Volume Ratio: As a particle shrinks, the percentage of atoms on its surface increases exponentially. *Result:* Nanomaterials are highly chemically reactive (excellent catalysts) and have lower melting points.

Quantum Confinement: When a material is reduced to nanoscale, the continuous energy bands of bulk materials break into discrete, quantized energy levels. *Result:* Optical and electrical properties change drastically (e.g., Gold nanoparticles can appear red or purple depending on size).

2. Approaches to Synthesis

Top-Down Approach (Destructive)

Breaking down bulk materials into nanomaterials using mechanical, chemical, or laser forces (e.g., Lithography, Ball milling).

Pros: Excellent for large-scale production.

Cons: Generates surface defects and impurities.

Bottom-Up Approach (Constructive)

Assembling materials atom-by-atom or molecule-by-molecule (e.g., Sol-gel, Chemical Vapor Deposition).

Pros: High purity, molecular precision, fewer defects.

Cons: Slower process, harder to scale up.

3. Key Synthesis Processes (Detailed)

A. Ball Mill Process (Top-Down)

- **Mechanism:** Uses a rotating cylindrical drum filled with heavy, hard grinding media (steel or ceramic balls) and the bulk powder to be ground.
- **Action:** As the drum rotates, the balls drop from the top of the drum, causing high-energy impacts. Through repeated severe plastic deformation, cold welding, and fracturing, the bulk material is reduced to nanoparticles.
- **Use:** Synthesis of metallic and ceramic nanocomposites.

B. Sol-Gel Process (Bottom-Up)

- **Step 1 (Hydrolysis):** Metal alkoxide precursors are mixed with water and alcohol, reacting to form a colloidal suspension of solid particles in liquid, called a **Sol**.
- **Step 2 (Condensation):** The particles link together, eliminating water/alcohol, forming a continuous, integrated microscopic 3D network called a **Gel**.
- **Step 3 (Aging & Drying):** Solvent is removed (via evaporation or supercritical drying) to form a porous solid (xerogel or aerogel).
- **Step 4 (Calcination):** The material is heated to high temperatures to remove organic residues, leaving pure nanoscale metal oxide powders or films.

4. Special Types of Materials

Zeolites: Microporous, crystalline aluminosilicate minerals.

Structure: They possess a 3D framework of pores and channels of precise, uniform nanoscale dimensions.

Applications: Act as molecular sieves, highly useful in water purification (ion exchange), chemical filters, and catalysis (oil refining).

Graphene: A 2D allotrope of carbon consisting of a single atomic layer arranged in a hexagonal (honeycomb) lattice.

Properties: Extremely high electrical & thermal conductivity (zero effective mass electrons), 200 times stronger than steel by weight, highly transparent, and flexible.



Graphene Hexagonal Lattice

5. Applications of Nanotechnology

Medicine: Targeted drug delivery (using gold nanoparticles to attack cancer cells specifically), highly sensitive nano-biosensors, and tissue engineering.

Electronics: Quantum dot displays (vibrant colors based on particle size), smaller/faster transistors, and ultra-high-density memory storage.

Energy: More efficient solar cells, high-capacity lithium-ion batteries, and better catalysts for hydrogen fuel cells.

Materials: Self-cleaning surfaces (Lotus effect), stain/water-resistant fabrics, and lightweight but ultra-strong composites for aerospace.