

Opioid Education & Naloxone Training

NAMES

Registered Opioid Overdose Prevention Program #787

Date

Objectives

- Learn about opioid use disorder, withdrawal, and overdose response
- Understand how to access Naloxone kits
- Complete the opioid overdose response steps
- Complete post-administration naloxone reports


Columbia Connections

- History of Leadership & Public Health
- Help reduce stigma
- Referral to appropriate resources
- Supportive community
- **Save lives**




South Bronx and East Harlem Continued to Experience High Rates of Overdose Deaths in 2024


Rate of unintentional drug overdose death per 100,000 residents, New York City, 2023



Rate of unintentional drug overdose death per 100,000 residents, New York City, 2024



Rate of unintentional drug poisoning (overdose) death per 100,000 residents, New York City



Top 5 neighborhoods with highest rates of overdose death




Source: NYC Office of Chief Medical Examiner and NYC Health Department, 2023-2024

Data for 2023 and 2024 are provisional and subject to change.

Harm Reduction

- A set of practical strategies and ideas aimed at reducing negative consequences associated with drug use
- A movement for social justice built on a belief in, and respect for, the rights of people who use drugs



Talking About Substance Use

Stigmatizing Language to Avoid

- Addict
- Substance abuser, dope fiend, junkie, tecato/a
- Substance abuse
- Clean
- Doctor shopper or drug seeker
- Replacement or substitution therapy

Person – Centered Language to Use Instead


- Person who uses or injects drugs
- Person living with substance use disorder
- Substance use or misuse
- Decreasing or stopping drug use
- Patient, participant, client
- Medication (such as suboxone) for opioid use disorder; acronym MOUD

What are opioids?


About Opioids

- Opioids are a class of drugs commonly used to reduce pain
- Types of opioids include:
- **Prescription Opioids:** Oxycodone (OxyContin), hydrocodone (Vicodin,) morphine, methadone, and percocet
 - **Synthetic Opioids:** Fentanyl, Carfentanil, Nitazines
 - **Illicit Opioids:** Heroin
- Short-acting (immediate-release) opioid and long-acting (extended-release) opioid

Opioid Use Disorder and Withdrawal



Opioid use disorder



Opioid withdrawal symptoms

Medication for Opioid Use Disorder

- Medication used to treat opioid use disorder
 - Manage withdrawal symptoms and limit the effects of opioids
 - Some medication, such as suboxone, contains naloxone in it, but it is not the same as the naloxone spray
- Current approved medications:
 - Buprenorphine, naltrexone, and methadone
 - Students can learn more information about MOUD through campus health services
 - Staff and Faculty can learn more through the Employee Assistance Program and health insurance benefits
 - NYC Residents can call 3-1-1 for resources near them.

Risk Factors and Harm Reduction

Risk Factors	Harm Reduction
Mixing drugs • Fentanyl • Extremely strong opioid • Used to: enhance heroin, imitate commercial pills • Other drugs can be laced with fentanyl • Synergistic effect	• Use fentanyl test strips or drug testing kit to test drugs • Start low, go slow • Avoid mixing substances • Get prescription medication from licensed pharmacies • Only take the medication and dose that is prescribed
Using alone • Increase risk for amount consumed • No awareness of overdose onset	• Start low, go slow • Have someone with you who is sober or not using • Never Use Alone Hotline (877- 696- 1996)

Naloxone (Narcan)

Naloxone (Narcan®)

What

- Medication that temporarily reverses an opioid overdose
- Works for 30-90 minutes
- As of Sept. 2023, Naloxone is available OTC in NYC

When

- Anytime you believe someone may be experiencing an opioid overdose


Why

- May save a life

Harmless if not an opioid overdose







Signs of an Overdose

Signs of an Opioid Overdose	Signs of an Opioid Overdose – Fentanyl
• Unresponsiveness or unconsciousness • Slowed or stopped breathing • Blue, purple, gray, or white lips or fingernails* • Snoring or gurgling sounds • Cold and clammy skin • Pinpoint eyes • Shout in the person's ear and use a sternal rub to check for responsiveness	• All signs of an opioid overdose • Standing seizures • Wooden chest and locked jaw • *Those with darker skin are more likely to have gray or ashen lips or fingernails • *Those with lighter skin are more likely to have blue or purplish lips or fingernails

Review: Overdose Response Steps

Overdose Response Steps


1. Call **911** for medical help. Tell them someone is not breathing.
2. Give naloxone. Do not test the device, as it can only be used once.



Peel



Place



Press

3. CPR (if you are trained)

Always administer naloxone first

Resuscitation should only be performed if the person is not breathing.

#1: If you are trained in CPR

You may do **both** rescue breathing and chest compressions

#2: Perform chest compressions alone


Some stimulation is likely better than none.

#3: Rescue Breathing (mouth-to-mouth):

1. Face mask is included in naloxone kit
 - One-way mask: be sure you can read printing clearly
2. Tilt head back, pinch nose
3. Start with 2 breaths, one after the other
4. If the chest does not rise
 1. Reposition head and check mouth for food, gum etc. and try again.
 2. If the chest is not rising, you may be blowing up their stomach
5. Give one breath every 5 seconds
6. Give normal sized breaths so you don't blow up the stomach or get dizzy

What is the recovery position?

4. When not giving naloxone or rescue breaths, put the person on their side to prevent choking.



5. Follow-up When Person Responds:

- Tell them that there is a medical emergency
- Advise against immediately using any drugs

If 9-1-1 was not called:



Option 1:
If the person is at least partly
able to communicate:

ACTIONS:

- *Ask if ok to call 9-1-1*
- *Offer to take them to the ER/Urgent Care*

OR

Option 2:
If not seeking medical care:

ACTIONS:

- *Stay with the person for at least 3 hours until after the naloxone has fully worn off*
- *Watch for return of overdose*

The Good Samaritan Law

Take-Away:

If someone is experiencing a life-threatening emergency (not just an overdose), they will not be prosecuted for drugs or alcohol found by the police (with a few exceptions)

Additional information:

Offers **protection** from:

- **Drugs up to an A2 felony offense** (possession of 8oz of narcotics)
- **Alcohol** (for underage drinkers)
- **Marijuana** (any amount)
- **Paraphernalia offenses**
- **Sharing of drugs** (in NYS: sharing constitutes a “sales” offense).

Does **NOT** offer **protection** from:

- **Other crimes on the scene** (weapons, stolen goods, drug sales)
- **Parole or probation violation** (but calling 9-1-1 would be considered favorably)
- **Previous conviction for class A-1, A-2, B felony possession** (B=over ½ lb. of marijuana)

Columbia's Version: Responsible Community Action Policy

Take-Away:

If a student makes a call for help (e.g. CU-EMS), they will not face campus discipline for the substance-related issue.

Additionally, Barnard, at its discretion, offers employees amnesty from minor policy violations related to the incident (e.g. when reporting harassment or discrimination they have experienced).

Does **NOT** offer protection from:

- **Other policy violations** (e.g. violence, vandalism, etc.)

Resources and Referrals

Morningside/Manhattanville (including JTS, TC, and UTS)	CUIMC	Barnard
#787 Columbia Health OOPP 212-854-8975 health@columbia.edu Alice! Health Promotion Go Ask Alice! And BASICS John Jay Hall, 3rd Floor 212-854-5453 or alice@columbia.edu Medical Services John Jay Hall, 4th Floor 212-854-7426 or Health@columbia.edu Counseling and Psychological Services Lerner Hall, 8th Floor 212-854-2878 or Health@columbia.edu Sexual Violence Response Lerner Hall, 7th Floor, Suite 700 Helpline: 212-854-HELP (212-854-4357) or svrresponse@columbia.edu	Student Health on Haven 100 Haven Avenue, 2nd Floor, Suite 230 212-305-3400 Studenthealth.cuimc.columbia.edu Addiction Information + Management Strategies 100 Haven Avenue, 2nd Floor, Suite 230 212-305-3400 aims@cumc.columbia.edu Sexual Violence Response 50 Haven Ave, Suite 206 Helpline: 212-854-HELP (212-854-4357) svrresponse@columbia.edu	Primary Care Health Service Brooks Hall, Lower Level 212-854-2091 barnard.edu/primarycare Furman Counseling Center 100 Hewitt Hall, First Floor 212-854-2092 or counseling@barnard.edu Denise LeFrak Alcohol & Substance Awareness Program 100 Hewitt Hall, First Floor 212-854-2092 or asap@barnard.edu The Wellness Spot LeFrak Center for Well-Being, 106B Barnard Hall 212-854-3063 or TheWellnessSpot@barnard.edu Being Barnard LeFrak Center for Well-Being, 106A Barnard Hall 212-853-8195 or BeingBarnard@barnard.edu

Off-Campus Resources

Local

Alliance LES Harm Reduction Center

(212) 645-0875

<https://alliance.nyc/leshrc>

NYC 988 Suicide and Crisis Lifeline

Call or Text 988

<https://nyc988.cityofnewyork.us/en/>

NYC Health – Department of Health and Mental Hygiene

<https://www.nyc.gov/site/doh/health/health-topics/alcohol-and-drug-use.page>

Office of Addiction Services and Support*

877-846-7369

<https://oasas.ny.gov/>

OnPoint NYC

(212) 828-8464

<https://onpointnyc.org/>

* Medication for opioid use disorder (MOUD) available

National

DanceSafe

888-636-2411

<https://dancesafe.org/>

MATTERS Network*

765-628-8377

<https://mattersnetwork.org/>

National Harm Reduction Coalition

<https://harmreduction.org/>

Never Use Alone Hotline

877-696-1996

<https://neverusealone.com/>

Next Distro

<https://nextdistro.org/>

North America Syringe Exchange Network (NASEN)

(253) 272-4857

<https://nasen.org/>

Reporting and Refills

- **Complete a Naloxone Recipient Form (NRF)**

- Only required if you are taking a kit
- Names and information are not shared with the city or the state
- You can either pick up a kit in-person at your nearest site or by mail (we cover the cost of shipping)

Report all administrations

- Report contact information on blue card; can also email health@columbia.edu
- Minimal information is 1 dose or 2 doses
- We do not report names or details to city or state
- **Return for another kit, if:**
 - You use your kit (anonymous reporting form)
 - Near expiration date

Certificate of Completion

Date _____

This certifies that _____ has been trained in opioid overdose prevention including the use of naloxone for the purpose of preventing death from an opioid overdose. This practice is legal under New York State Public Health Law Section 3309 and under 10 N.Y.C.R.R. Section 80.138.

Prescribed by (name): Melanie J Bernitz, MD
Under a non-patient specific order

Check formulation(s), if any

- ☒ NARCAN® Nasal Spray (naloxone HCl) 4 mg units, 2 units. Administer IN prn as directed
- ☐ Naloxone 2mg/2ml, 2 prefilled syringes with atomizers. Administer IN prn as directed
- ☐ Naloxone 0.4mg/ml, 2 vials and 2 3cc syringes. Administer IM prn as directed
- ☐ Other, specify: _____

NYSDOH Registered Opioid Overdose Program
Get a kit at your pharmacy or program.

Columbia Health
1116 Amsterdam Ave, 125 Wallach Hall, New York, NY 10027
212-854-8975 health@columbia.edu

New York State Department of Health Opioid Overdose Initiative
1-800-692-8528 | www.health.ny.gov/overdose

Syringe Access, Drug & Syringe Disposal Resources
www.health.ny.gov/syringes | www.thepointny.org

OASAS HOPEline
1-877-846-7369 | www.oasas.ny.gov

9808 8/17

Information on reporting and requesting refills is on your Certificate of Completion (blue) card

Questions & Thank You

- Remember to submit the Naloxone Recipient Form (NRF) to get a kit
- Remember to report if you administer (anonymous) and get a refill
- Tell others about the opportunity to be trained
- Contact us for more information
 - health@columbia.edu
 - 212-854-8975
 - health.columbia.edu

