

Mechanical Processing in Internally Coupled Ears

Anupam Prasad Vedurmudi

TMP Thesis Defence
July 5, 2013



Auditory Systems



Independent Ears

Eustachian tubes typically very narrow.

Effectively independent eardrum vibrations.



Coupled Ears


Eardrums connected through wide eustachian tubes and a large mouth cavity.

Eardrums vibrations influence each other.

Advantages of Low Frequency Hearing


Mouth Cavity

Previous Model




► $V_{\text{cyl}} = \pi a_{\text{tymp}}^2 L$

Our Model



- V_{cyl} based on anatomical data.
- $a_{\text{cyl}} = \sqrt{V_{\text{cyl}}/\pi L}$

Acoustic Head Model



- ▶ I - Ipsilateral C - Contralateral
- ▶ Sound source far enough away from the animal ("Infinity").
- ▶ Phase difference between sound at both ears - $\Delta = 1.5kL \sin \theta$.
- ▶ No appreciable amplitude difference, $|p_0| = |p_L|$.

Cavity Pressure

3D Wave Equation


$$\frac{1}{c^2} \partial_t^2 p(x, r, \phi, t) = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial p(x, r, \phi, t)}{\partial r} \right) + \frac{1}{r^2} \frac{\partial p(x, r, \phi, t)}{\partial \phi^2} + \frac{\partial p(x, r, \phi, t)}{\partial x^2} \quad (1)$$

No-penetration at all solid boundaries

$$-j\rho\omega\mathbf{v} = \nabla p(x, r, \phi; t) = 0 \quad (2)$$

Eardrum


Sketch of a Tokay eardrum as seen from the outside^a.



COL - approximate position opposite the extracolumella insertion.

^aG. A. Manley, "The middle ear of the tokay gecko," *Journal of Comparative Physiology*, vol. 81, no. 3, pp. 239–250, 1972

The ICE eardrum.



Extracolumella (dark) - rigid, stationary.

Tympanum - assumed linear elastic.

Rigidly clamped at the boundaries ($r = a_{\text{tym}}$ and $\phi = \beta, 2\pi - \beta$)

Membrane Vibrations

Membrane EOM

$$-\partial_t^2 u(r, \phi; t) - 2\alpha \partial_t u(r, \phi; t) + c_M^2 \nabla^2 u(r, \phi; t) = \frac{1}{\rho_m d} \Psi(r, \phi; t) \quad (3)$$

Evaluation

Conclusion

Thank You

