

Announcements

1. Lab Exp 2 due on Jan 2

2. Test 2

Recap:

White Noise (Discrete)

Flat p.s.d \Leftrightarrow WSS, uncorrelated, zero mean.

$$X = \text{White Noise} \xrightarrow{\quad} \boxed{h[n]} \longrightarrow Y \quad S_X(f) = \sigma^2$$

$$S_Y(f) = \sigma^2 \cdot |H(f)|^2$$

$$R_Y(k) = \sigma^2 \sum_{n=k}^{\infty} h[n] h[n-k]$$

— (*) h is causal

Example: Suppose an uncorrelated WSS process with mean m , variance σ^2 is input to an LTI system with impulse response.

$$h[k] = r^k; \quad k \geq 0 \\ |r| < 1$$

Find mean and autocorrelation of output?

$$Y[n] = \sum_{k=0}^{\infty} h[k] X[n-k]$$

$$E[Y[n]] = \sum_{k=0}^{\infty} h[k] \underbrace{E[X[n-k]]}_m$$

$$= m \sum_{k=0}^{\infty} h[k]$$

$$= m \cdot \sum_{k=0}^{\infty} r^k$$

$$m_y = \frac{m}{1-r} //$$

$$C_y[k] = \sigma^2 \sum_{n=k}^{\infty} h[n] h[n-k]$$

Equation (*)

$$= \sigma^2 \sum_{n=k}^{\infty} r^n r^{n-k}$$

$$= \sigma^2 \cdot r^{-k} \sum_{n=k}^{\infty} r^{2n}$$

$$= \sigma^2 r^{-k} \cdot r^{2k} \underbrace{\sum_{n=k}^{\infty} r^{2(n-k)}}_{\sum_{l=0}^{\infty} r^{2l} \quad [l=n-k]}$$

$$= \sigma^2 r^k \cdot \sum_{l=0}^{\infty} r^{2l}$$

$$= \sigma^2 \cdot r^k \cdot \frac{1}{1-r^2} //$$

$$C_y[k] = R_y[k] - m_y^2 //$$

White Noise (Continuous)

RP with the following auto-correlation function

$$K_x(\tau) = \sigma^2 \delta(\tau)$$

$$\boxed{S_x(f) = \sigma^2 \quad \forall f}$$

\Leftrightarrow Uncorrelated, WSS with zero mean.

$$R_x(0) = E[X^2(t)]$$

Good model
for

"thermal noise"

of

electronic components

$$= \int_{-\infty}^{\infty} S_x(f) df$$

$$E[X^2(t)] = \int_{-\infty}^{\infty} \sigma^2 df$$

\rightarrow Infinite power

Example: Suppose the transfer function of an LTI system is given by

$$H(\omega) = \text{sgn}(\omega) \left(\frac{\omega}{2\pi} \right)^2 \exp(-j(\omega \tau / \pi))$$

$H(\omega)$

ω -domain
(not in
f-domain)
 $\omega = 2\pi f$

$$W(\omega) = \begin{cases} 1, & |\omega| \leq 40\pi \\ 0, & \text{else.} \end{cases}$$

(input)

Let X be a WSS process with autocorrelation function

$$R_x(\tau) = \frac{\sigma^2}{2} \delta(\tau) + 2$$

Compute total average output power?

$$S_x(\omega) = F[R_x]$$

$$= \frac{\sigma^2}{2} + 4\pi \delta(\omega)$$

$$S_y(\omega) = |H(\omega)|^2 S_x(\omega)$$

$$= \left[\frac{\sigma^2}{2} + 4\pi \delta(\omega) \right] |H(\omega)|^2$$

$$= \frac{5}{2} |H(\omega)|^2$$

$$= \frac{5}{2} \left(\frac{\omega}{2\pi}\right)^4 W(\omega)$$

$$R_Y(0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} S_Y(\omega) d\omega$$

$$= \frac{1}{2\pi} \int_{-40\pi}^{40\pi} \frac{5}{2} \left(\omega/2\pi\right)^4 d\omega$$

$$= 3.2 \times 10^6 //$$

Special Case: White Gaussian Noise

(White Gaussian Process)

White Noise (Continuous time)

$$X(t) \sim \mathcal{N}(0, \sigma^2)$$

White Noise (Discrete time)


$$X[n] \sim \mathcal{N}(0, \sigma^2)$$

Theorem: The derivative of Brownian motion process is white Gaussian noise.

More than one random process

"Vector random process"

$$x(t)$$



$$Z(t) = (X(t), Y(t))$$

$$SSS: F_{Z(t_1) Z(t_2) \dots Z(t_k)} = F_{Z(t_1+\tau) Z(t_2+\tau) \dots Z(t_k+\tau)}$$

Jointly Wide Sense Stationary (2 random
processes)

Cross-correlation function

$$R_{xy}(t_1, t_2) = E[X(t_1) Y(t_2)]$$

$$R_{yx}(t_1, t_2) = E[Y(t_1) X(t_2)]$$