

Total No. of Questions: 6

Total No. of Printed Pages: 2



Enrollment No.....

Faculty of Pharmacy

End Sem Examination Dec 2024

PY3RC02 Remedial Mathematics

Programme: B. Pharm.

Branch/Specialisation: Pharmacy

Duration: 3 Hrs.**Maximum Marks: 35**

Note: All questions are compulsory. Internal choices, if any, are indicated. Assume suitable data if necessary. Notations and symbols have their usual meaning.

	Marks	BL	PO	CO	PSO
Q.1 i. What is the value of $\lim_{x \rightarrow 0} \frac{\sin x}{x}$?	1	1	1	1	
ii. If $A = \begin{bmatrix} 3 & 5 \\ -4 & 2 \end{bmatrix}$ then write the minor of M_{22} .	1	1	1	1	
iii. Differentiation of x^n with respect to x is ____.	1	1	1	1	
iv. Find the slope of line joining the points (2, -5) and (4,1).	1	2	1	2	
v. Degree of differential equation $\frac{dy}{dx} + 2y = x$ is-	1	1	1	1	
Q.2 i. Resolve into partial fraction- $\frac{2x+3}{(x+1)(x-3)}$	6	3	1	3	
OR ii. (a) Prove that $\log 72 = 2 \log 3 + 3 \log 2$. (b) If $f(x) = x^3 + 14x^2 - 2x + 4$ then find the value of $f(3)$.	6	3	1	3	
Q.3 i. If $A = \begin{pmatrix} 3 & 2 & 1 \\ 0 & -5 & 6 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 5 & -2 \\ 3 & 1 & 8 \end{pmatrix}$ then show that $(A + B)^T = A^T + B^T$, where T → Transpose. OR ii. Evaluate the following-	6	3	1	3	
(a) $\begin{vmatrix} -1 & 2 & 3 \\ 2 & -3 & 1 \\ 3 & -1 & 1 \end{vmatrix}$ (b) $\begin{vmatrix} 5 & 4 \\ -2 & 3 \end{vmatrix}$					

Q.4 i. Find $\frac{dy}{dx}$ if-	6	3	1	4
(a) $y = e^{4x} \cdot \sin x$ (b) $y = x^3 + \tan x$				
OR ii. If $y = ae^{mx} + be^{-mx}$ then show that $\frac{d^2y}{dx^2} = m^2y$.	6	3	1	3
Q.5 i. Write distance formula and find the distance between the points (10,15) and (7, 10).	6	3	1	3
OR ii. Find the following-	6	3	1	3
(a) $\int x \sin x dx$				
(b) $\int_{-1}^2 x^3 dx$				
Q.6 i. Solve the equation- $\frac{dy}{dx} - y = e^{2x}$	6	3	1	3
OR ii. Find the Laplace transform of- $F(t) = 1 + \sin 2t + e^t$	6	3	1	3

(1)

Medi-Caps University
 Faculty of Pharmacy
 End Sem Examination Dec (2024)

Solutions of PY3RC02 Remedial Mathematics

Q1. 1) $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

+1

2) If $A = \begin{bmatrix} 3 & 5 \\ -4 & 2 \end{bmatrix}$ then minor $M_{22} = 2$

+1

3) $\frac{d}{dx} x^n = n x^{n-1}$

+1

4) Slope of line joining $(2, -5)$ & $(4, 1)$ is 3

+1

5) Degree of differential equation $\frac{dy}{dx} + 2y = x^2$ is 1

+1

Q2. 1) Resolve into partial fraction $\frac{2x+3}{(x+1)(x-3)}$

$$\frac{2x+3}{(x+1)(x-3)} = \frac{A}{(x+1)} + \frac{B}{(x-3)} \rightarrow ①$$

+1

$$2x+3 = A(x-3) + B(x+1)$$

+2

$$2x+3 = Ax + Bx - 3A + B$$

$$2x = (A+B)x \quad -3A + B = 3$$

$$A+B=2$$

$$\begin{array}{r} A+B=2 \\ -3A+B=3 \\ \hline + - - \end{array}$$

$$4A = -1$$

$$A = -1/4, B = 9/4$$

+2

(2)

put in equation ①

$$\begin{aligned}\frac{2x+3}{(x+1)(x-3)} &= \frac{-1/4}{(x+1)} + \frac{9/4}{(x-3)} & +1 \\ &= \frac{9}{4(x-3)} - \frac{1}{4(x+1)} & \text{Ans} \\ &\quad \underline{\text{or}}\end{aligned}$$

ii) Prove that $\log_{12} = 2\log_3 + 3\log_2$

a) $\Rightarrow \log(2 \times 2 \times 2 \times 3 \times 3)$ +1

$\Rightarrow \log(2^3 \times 3^2)$ +1.

$\Rightarrow \log 2^3 + \log 3^2$ formula $\log(m \cdot n) = \log m + \log n$

formula $\log a^b = b \log a$ +1.

$\Rightarrow 3\log_2 + 2\log_3 = 2\log_3 + 3\log_2$ Ans. +1.

b) if

$f(x) = 2x^3 + 14x^2 - 2x + 4$ then $f(3) = ?$

$$\begin{aligned}f(3) &= (3)^3 + 14(3)^2 - 2(3) + 4 \\ &= 27 + 122 - 6 + 4\end{aligned}$$

$$= 147 \quad \underline{\text{Ans}}$$

Q3

$$\text{i) } A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & -5 & 6 \end{bmatrix} \quad B = \begin{bmatrix} 4 & 5 & -2 \\ 3 & 1 & 8 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 3 & 0 \\ 2 & -5 \\ 1 & 6 \end{bmatrix} \quad B^T = \begin{bmatrix} 4 & 3 \\ 5 & 1 \\ -2 & 8 \end{bmatrix}$$

$$A^T + B^T = \begin{bmatrix} 7 & 3 \\ 7 & -4 \\ -1 & 14 \end{bmatrix}$$

$$A+B = \begin{bmatrix} 3 & 2 & 1 \\ 0 & -5 & 6 \end{bmatrix} + \begin{bmatrix} 4 & 5 & -2 \\ 3 & 1 & 8 \end{bmatrix} = \begin{bmatrix} 7 & 7 & -1 \\ 3 & -4 & 14 \end{bmatrix}$$

$$(A+B)^T = \begin{bmatrix} 7 & 3 \\ 7 & -4 \\ -1 & 14 \end{bmatrix}$$

Hence

$$A^T + B^T = (A+B)^T$$

OR

$$\text{ii) a) } \begin{vmatrix} -1 & 2 & 3 \\ 2 & -3 & 1 \\ 3 & -1 & 1 \end{vmatrix} = -1(-3+1) - 2(2-3) + 3(-2+9)$$

$$= -1(-2) - 2(-1) + 3(7)$$

$$= 2 + 2 + 21$$

$$= 25 \quad \text{Ans}$$

$$\text{b) } \begin{vmatrix} 5 & 4 \\ -2 & 3 \end{vmatrix} = 15 - (-8) = 15 + 8 = 23 \quad \text{Ans}$$

Q4 i) find $\frac{dy}{dx}$.

a) $y = e^{4x} \cdot \sin x$

formula : $u \cdot v = uv' + vu'$

$$\begin{aligned}\frac{dy}{dx} &= e^{4x} (\cos x) + \sin x (4e^{4x}) \\ &= \cos x e^{4x} + 4 \sin x e^{4x} \\ &= e^{4x} (\cos x + 4 \sin x) \quad \text{Ans}\end{aligned}$$

b) $y = x^3 + \tan x$

$$\frac{dy}{dx} = \frac{d(x^3)}{dx} + \frac{d(\tan x)}{dx}$$

$$= 3x^2 + \sec^2 x. \quad \text{Ans}$$

ii) $y = ae^{mx} + be^{-mx}$. Then

$$\frac{dy}{dx} = ame^{mx} - bme^{-mx}$$

$$\frac{d^2y}{dx^2} = am^2e^{mx} + bm^2e^{-mx}$$

$$\frac{d^2y}{dx^2} = m^2(ae^{mx} + be^{-mx})$$

$$\frac{d^2y}{dx^2} = m^2y. \quad \text{Ans}$$

Q5 i) distance formula = $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$. +2

distance b/w points (10, 15) and (7, 10)

$$= \sqrt{(10-7)^2 + (15-10)^2}$$

$$= \sqrt{9 + 25}$$

$$= \sqrt{34} \quad \text{Ans}$$

or

ii)
a) $\int x \sin x dx$.

formula of integration by parts

$$\int u \cdot v' = u \int v - \int [u' \int v].$$

$$u = x \quad v = \sin x$$

$$\int x \sin x dx = x(-\cos x) - \int [\frac{dx}{dx} \int \sin x] dx + 1$$

$$= -x \cos x - \int (-\cos x) dx.$$

$$= -x \cos x + \int \cos x dx.$$

$$= -x \cos x + \sin x + C \text{ Ans}$$

b) $\int_{-1}^2 x^3 dx = \left[\frac{x^{3+1}}{3+1} \right]_1^2$

$$= \left[\frac{x^4}{4} \right]_1^2$$

(6)

$$= \frac{1}{4} (x^4 - 1)^2$$

$$= \frac{1}{4} (2^4 - (-1)^4)$$

$$= \frac{1}{4} (16 - 1) = \frac{15}{4} \quad \underline{\text{Ans}}$$

Q6 i)

$$\frac{dy}{dx} - y = e^{2x}$$

$$\frac{dy}{dx} + Py = Q$$

$$P = -1$$

$$Q = e^{2x}$$

$$IF = e^{\int P dx} = e^{-\int dx} = e^{-x}$$

$$y \cdot IF = \int Q \cdot IF dx$$

$$y \cdot e^{-x} = \int e^{2x} \cdot e^{-x} dx.$$

$$y \cdot e^{-x} = \int e^{x} dx.$$

$$y \cdot e^{-x} = e^x + C \quad \underline{\text{Ans}}$$

$$\text{ii)} \quad f(t) = 1 + \sin 2t + e^t$$

$$L\{f(t)\} = L\{1\} + L\{\sin 2t\} + L\{e^t\}. \quad +3$$

$$= \frac{1}{s} + \frac{2}{s^2 + 4} + \frac{1}{s-1}$$

+3