

# 1 PCA

For the following problems, we have  $N$  *zero-mean* data points  $\mathbf{x}_i \in \mathbb{R}^{D \times 1}$  and  $\mathbf{S} = \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i \mathbf{x}_i^T \in \mathbb{R}^{D \times D}$  is the sample covariance matrix of the dataset.

## 1.1 Derivation of Second Principal Component

(a) (5 points) Let cost function

$$J = \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i - p_{i1} \mathbf{e}_1 - p_{i2} \mathbf{e}_2)^T (\mathbf{x}_i - p_{i1} \mathbf{e}_1 - p_{i2} \mathbf{e}_2)$$

with  $\mathbf{e}_1$  and  $\mathbf{e}_2$  are the orthonormal vector basis for the dimensionality reduction, i.e.  $\|\mathbf{e}_1\|_2 = 1$ ,  $\|\mathbf{e}_2\|_2 = 1$ , and  $\mathbf{e}_1^T \mathbf{e}_2 = 0$ , and some coefficients  $p_{i1}$  and  $p_{i2}$ .

Show that  $\frac{\partial J}{\partial p_{i2}} = 0$  yields  $p_{i2} = \mathbf{e}_2^T \mathbf{x}_i$ , i.e. the projection length of data point  $\mathbf{x}_i$  along vector  $\mathbf{e}_2$ .

**Answer:**

$$\begin{aligned} J &= \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i - p_{i1} \mathbf{e}_1 - p_{i2} \mathbf{e}_2)^T (\mathbf{x}_i - p_{i1} \mathbf{e}_1 - p_{i2} \mathbf{e}_2) \\ &= \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i^T - p_{i1} \mathbf{e}_1^T - p_{i2} \mathbf{e}_2^T) (\mathbf{x}_i - p_{i1} \mathbf{e}_1 - p_{i2} \mathbf{e}_2) \\ &= \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i^T \mathbf{x}_i + p_{i1}^2 \mathbf{e}_1^T \mathbf{e}_1 + p_{i2}^2 \mathbf{e}_2^T \mathbf{e}_2 - 2p_{i1} \mathbf{e}_1^T \mathbf{x}_i + 2p_{i1} p_{i2} \mathbf{e}_1^T \mathbf{e}_2 - 2p_{i2} \mathbf{e}_2^T \mathbf{x}_i) \\ &= \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i^T \mathbf{x}_i + p_{i1}^2 (1) + p_{i2}^2 (1) - 2p_{i1} \mathbf{e}_1^T \mathbf{x}_i + 2p_{i1} p_{i2} (0) - 2p_{i2} \mathbf{e}_2^T \mathbf{x}_i) \\ &= \frac{1}{N} \sum_{i=1}^N (\mathbf{x}_i^T \mathbf{x}_i + p_{i1}^2 + p_{i2}^2 - 2p_{i1} \mathbf{e}_1^T \mathbf{x}_i - 2p_{i2} \mathbf{e}_2^T \mathbf{x}_i) \end{aligned}$$

$$\frac{\partial J}{\partial p_{i2}} = 2p_{i2} - 2\mathbf{e}_2^T \mathbf{x}_i = 0$$

$$p_{i2} = \mathbf{e}_2^T \mathbf{x}_i$$

(b) (5 points) Show that the value of  $\mathbf{e}_2$  that minimizes cost function

$$\tilde{J} = -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + \lambda_2 (\mathbf{e}_2^T \mathbf{e}_2 - 1) + \lambda_{12} (\mathbf{e}_2^T \mathbf{e}_1 - 0)$$

is given by the eigenvector associated with the second largest eigenvalue of  $\mathbf{S}$ .

$\lambda_2$  is the Lagrange Multiplier for equality constraint  $\mathbf{e}_2^T \mathbf{e}_2 = 1$  and  $\lambda_{12}$  is the Lagrange Multiplier for equality constraint  $\mathbf{e}_2^T \mathbf{e}_1 = 0$ .

*Hint:* Recall that  $\mathbf{S} \mathbf{e}_1 = \lambda_1 \mathbf{e}_1$  ( $\mathbf{e}_1$  is the normalized eigenvector associated with the largest eigenvalue  $\lambda_1$  of  $\mathbf{S}$ ) and  $\frac{\partial \mathbf{y}^T \mathbf{A} \mathbf{y}}{\partial \mathbf{y}} = (\mathbf{A} + \mathbf{A}^T) \mathbf{y}$ . Also notice that  $\mathbf{S}$  is a symmetric matrix.

**Answer:**

Taking partial derivative of  $\tilde{J}$  with respect to Lagrange Multiplier  $\lambda_2$  yields:

$$\begin{aligned} \frac{\partial \tilde{J}}{\partial \lambda_2} &= \mathbf{e}_2^T \mathbf{e}_2 - 1 = 0 \\ \mathbf{e}_2^T \mathbf{e}_2 &= 1 \end{aligned}$$

Taking partial derivative of  $\tilde{J}$  with respect to Lagrange Multiplier  $\lambda_{12}$  yields:

$$\frac{\partial \tilde{J}}{\partial \lambda_{12}} = \mathbf{e}_2^T \mathbf{e}_1 = 0$$

Taking partial derivative of  $\tilde{J}$  with respect to Lagrange Multiplier  $\mathbf{e}_2$  yields:

$$\begin{aligned} \frac{\partial \tilde{J}}{\partial \mathbf{e}_2} &= -(\mathbf{S} + \mathbf{S}^T) \mathbf{e}_2 + 2\lambda_2 \mathbf{e}_2 + \lambda_{12} \mathbf{e}_1 = 0 \\ &\quad - 2\mathbf{S} \mathbf{e}_2 + 2\lambda_2 \mathbf{e}_2 + \lambda_{12} \mathbf{e}_1 = 0 \end{aligned} \tag{1}$$

Pre-multiply (or left-multiply) the equation 1 with  $\mathbf{e}_2^T$  yields:

$$\begin{aligned} -2\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + 2\lambda_2 \mathbf{e}_2^T \mathbf{e}_2 + \lambda_{12} \mathbf{e}_2^T \mathbf{e}_1 &= 0 \\ -2\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + 2\lambda_2 (1) + \lambda_{12} (0) &= 0 \\ \lambda_2 &= \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 \end{aligned}$$

Pre-multiply (or left-multiply) the equation 1 with  $\mathbf{e}_1^T$  yields:

$$\begin{aligned} -2\mathbf{e}_1^T \mathbf{S} \mathbf{e}_2 + 2\lambda_2 \mathbf{e}_1^T \mathbf{e}_2 + \lambda_{12} \mathbf{e}_1^T \mathbf{e}_1 &= 0 \\ -2\mathbf{e}_1^T \mathbf{S} \mathbf{e}_2 + 2\lambda_2 (0) + \lambda_{12} (1) &= 0 \\ \lambda_{12} &= 2\mathbf{e}_1^T \mathbf{S} \mathbf{e}_2 \\ \lambda_{12} &= 2\mathbf{e}_1^T \mathbf{S}^T \mathbf{e}_2 \\ \lambda_{12} &= 2(\mathbf{S} \mathbf{e}_1)^T \mathbf{e}_2 \\ \lambda_{12} &= 2(\lambda_1 \mathbf{e}_1)^T \mathbf{e}_2 \\ \lambda_{12} &= 2\lambda_1 \mathbf{e}_1^T \mathbf{e}_2 \\ \lambda_{12} &= 2\lambda_1 (0) \\ \lambda_{12} &= 0 \end{aligned}$$

Substituting  $\lambda_{12} = 0$  into equation 1:

$$\begin{aligned} -2\mathbf{S} \mathbf{e}_2 + 2\lambda_2 \mathbf{e}_2 + (0) \mathbf{e}_1 &= 0 \\ \mathbf{S} \mathbf{e}_2 &= \lambda_2 \mathbf{e}_2 \end{aligned}$$

Thus,  $\mathbf{e}_2$  is an eigenvector associated with eigenvalue  $\lambda_2$  of  $\mathbf{S}$ .  
 Substituting  $\lambda_2 = \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2$  and  $\lambda_{12} = 0$  into the definition of cost function  $\tilde{J}$  yields:

$$\begin{aligned}
 \tilde{J} &= -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + \lambda_2 (\mathbf{e}_2^T \mathbf{e}_2 - 1) + \lambda_{12} (\mathbf{e}_2^T \mathbf{e}_1 - 0) \\
 &= -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 (\mathbf{e}_2^T \mathbf{e}_2 - 1) + (0) (\mathbf{e}_2^T \mathbf{e}_1 - 0) \\
 &= -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 \mathbf{e}_2^T \mathbf{e}_2 - \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 \\
 &= -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 + \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 (1) - \mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 \\
 &= -\mathbf{e}_2^T \mathbf{S} \mathbf{e}_2 \\
 &= -\lambda_2
 \end{aligned}$$

Thus, to minimize  $\tilde{J}$ , we should pick the maximum possible value for  $\lambda_2$ . Since  $\lambda_1$  is the largest eigenvalue of  $\mathbf{S}$ ,  $\lambda_2$  should be the second largest eigenvalue of  $\mathbf{S}$ , and  $\mathbf{e}_2$  is the eigenvector associated with eigenvalue  $\lambda_2$ .

## 1.2 Derivation of PCA Residual Error

(a) (5 points) Prove that for a data point  $\mathbf{x}_i$ :

$$\|\mathbf{x}_i - \sum_{j=1}^K p_{ij} \mathbf{e}_j\|_2^2 = \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j$$

*Hint:* The most common method to prove a mathematical equation of this flavor is by using mathematical induction. To perform a proof by mathematical induction in this case, first show that the equation above holds for the base case  $K = 1$ , and then using the assumption that the equation holds for  $K = k - 1$ , show that the equation also holds for  $K = k$ , for any  $1 \leq k \leq D$ .

Use the fact that  $\mathbf{e}_j^T \mathbf{e}_j = 1$  (length of eigenvector  $\mathbf{e}_j$  is 1) and  $\mathbf{e}_j^T \mathbf{e}_m = 0$  for  $j \neq m$  (eigenvectors are perpendicular each other for square symmetric matrix  $\mathbf{S}$ ). Also, use definition  $p_{ij} = \mathbf{e}_j^T \mathbf{x}_i$ .

**Answer:**

Proof by mathematical induction:

- Base case  $K = 1$ :

$$\begin{aligned}
 \|\mathbf{x}_i - p_{i1} \mathbf{e}_1\|_2^2 &= (\mathbf{x}_i - p_{i1} \mathbf{e}_1)^T (\mathbf{x}_i - p_{i1} \mathbf{e}_1) \\
 &= (\mathbf{x}_i^T - p_{i1} \mathbf{e}_1^T) (\mathbf{x}_i - p_{i1} \mathbf{e}_1) \\
 &= \mathbf{x}_i^T \mathbf{x}_i - p_{i1} \mathbf{x}_i^T \mathbf{e}_1 - p_{i1} \mathbf{e}_1^T \mathbf{x}_i + p_{i1}^2 \mathbf{e}_1^T \mathbf{e}_1 \\
 &= \mathbf{x}_i^T \mathbf{x}_i - 2p_{i1} \mathbf{x}_i^T \mathbf{e}_1 + p_{i1}^2 (1) \\
 &= \mathbf{x}_i^T \mathbf{x}_i - 2\mathbf{e}_1^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_1 + \mathbf{e}_1^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_1 \\
 &= \mathbf{x}_i^T \mathbf{x}_i - \mathbf{e}_1^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_1
 \end{aligned}$$

Thus the equation holds for base case  $K = 1$

- Now, assuming that the equation holds for  $K = k - 1$ , i.e.:

$$\|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 = \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^{k-1} \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j$$

we will show that the equation also holds for  $K = k$ , as follows:

$$\begin{aligned}
\|\mathbf{x}_i - \sum_{j=1}^k p_{ij} \mathbf{e}_j\|_2^2 &= \left( \mathbf{x}_i - \sum_{j=1}^k p_{ij} \mathbf{e}_j \right)^T \left( \mathbf{x}_i - \sum_{j=1}^k p_{ij} \mathbf{e}_j \right) \\
&= \left( \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) - p_{ik} \mathbf{e}_k \right)^T \left( \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) - p_{ik} \mathbf{e}_k \right) \\
&= \left( \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right)^T - p_{ik} \mathbf{e}_k^T \right) \left( \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) - p_{ik} \mathbf{e}_k \right) \\
&= \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right)^T \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) - \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right)^T p_{ik} \mathbf{e}_k \\
&\quad - p_{ik} \mathbf{e}_k^T \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) + p_{ik}^2 \mathbf{e}_k^T \mathbf{e}_k \\
&= \|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 - \left( \mathbf{x}_i^T - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j^T \right) p_{ik} \mathbf{e}_k - p_{ik} \mathbf{e}_k^T \left( \mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j \right) + p_{ik}^2 (1) \\
&= \|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 - p_{ik} \mathbf{x}_i^T \mathbf{e}_k + \sum_{j=1}^{k-1} p_{ij} p_{ik} \mathbf{e}_j^T \mathbf{e}_k - p_{ik} \mathbf{e}_k^T \mathbf{x}_i + \sum_{j=1}^{k-1} p_{ij} p_{ik} \mathbf{e}_k^T \mathbf{e}_j + p_{ik}^2 \\
&= \|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 - 2p_{ik} \mathbf{x}_i^T \mathbf{e}_k + \sum_{j=1}^{k-1} p_{ij} p_{ik} (0) + \sum_{j=1}^{k-1} p_{ij} p_{ik} (0) + p_{ik}^2 \\
&= \|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 - 2p_{ik} \mathbf{x}_i^T \mathbf{e}_k + p_{ik}^2 \\
&= \|\mathbf{x}_i - \sum_{j=1}^{k-1} p_{ij} \mathbf{e}_j\|_2^2 - 2\mathbf{e}_k^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_k + \mathbf{e}_k^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_k \\
&= \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^{k-1} \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j - \mathbf{e}_k^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_k \\
&= \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^k \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j
\end{aligned}$$

Thus the equation holds for any  $1 \leq K \leq D$

(b) **(5 points)** Now show that

$$J_K \triangleq \frac{1}{N} \sum_{i=1}^N \left( \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j \right) = \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \lambda_j$$

*Hint:* recall that  $\mathbf{e}_j^T \mathbf{S} \mathbf{e}_j = \lambda_j \mathbf{e}_j^T \mathbf{e}_j = \lambda_j$

**Answer:**

$$\begin{aligned}
 J_K &\triangleq \frac{1}{N} \sum_{i=1}^N \left( \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j \right) \\
 &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^K \mathbf{e}_j^T \mathbf{x}_i \mathbf{x}_i^T \mathbf{e}_j \\
 &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \mathbf{e}_j^T \left( \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i \mathbf{x}_i^T \right) \mathbf{e}_j \\
 &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \mathbf{e}_j^T \mathbf{S} \mathbf{e}_j \\
 &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \lambda_j \mathbf{e}_j^T \mathbf{e}_j \\
 &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \lambda_j
 \end{aligned}$$

- (c) (**5 points**) If  $K = D$  principal components are used, there is no truncation, so  $J_D = 0$ . Use this to show that the error from only using  $K < D$  principal components is given by

$$J_K = \sum_{j=K+1}^D \lambda_j$$

**Answer:**

When  $K = D$ :

$$\begin{aligned}
 J_D &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^D \lambda_j = 0 \\
 \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i &= \sum_{j=1}^D \lambda_j
 \end{aligned}$$

Thus for  $K < D$ :

$$\begin{aligned}
 J_K &= \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^T \mathbf{x}_i - \sum_{j=1}^K \lambda_j \\
 &= \left( \sum_{j=1}^D \lambda_j \right) - \left( \sum_{j=1}^K \lambda_j \right) \\
 &= \sum_{j=K+1}^D \lambda_j
 \end{aligned}$$

### 1.3 A Real Example

(a) The eigenvectors and values are as follows:

$$u_1 = \begin{bmatrix} 0.22 \\ 0.41 \\ 0.88 \end{bmatrix}$$

$$u_2 = \begin{bmatrix} 0.25 \\ 0.85 \\ -0.46 \end{bmatrix}$$

$$u_3 = \begin{bmatrix} 0.94 \\ -0.32 \\ -0.08 \end{bmatrix}$$

$$\lambda_1 = 1626.52 \quad \lambda_2 = 128.99 \quad \lambda_3 = 7.10$$

- (b)  $u_2$  and  $u_3$  can be omitted because the corresponding eigenvalues for these two directions are contributing a small amount to the total variation of the data. In fact  $u_1$  accounts for  $\frac{\lambda_1}{\lambda_1 + \lambda_2 + \lambda_3} = 92.8\%$  of the data variation and  $u_2$  accounts for 7.32% of variation in data. The remaining principal component, explaining only 0.40% of the data, is negligible compared to the first two.
- (c) We might think of  $u_1$  as giving a generalized notion of “size” that incorporates length, wingspan, and weight. Indeed, all three entries of  $u_1$  have the same sign, indicating that birds with larger “size” tend to have larger length, wingspan, and weight.

## 2 Hidden Markov Model (25 Points)

In this problem, you will implement Hidden Markov Model. First, please read forward, backward, and Viterbi algorithm in the lecture note.

A simple DNA sequence is  $\mathbf{O} = \overline{O_1 O_2 \cdots O_T}$ , with each component  $O_i$  takes from  $\{A, C, G, T\}$ . Assume it is generated from a Hidden Markov Model controlled by a hidden variable  $X$ , which takes two possible states  $S_1, S_2$ .

This HMM has the following parameters  $\Theta = \{\pi_i, a_{ij}, b_{ik}\}$  for  $i, j = 1, 2$  and  $k \in \{A, C, G, T\}$ :

- Initial state distribution  $\pi_i$  for  $i = 1, 2$ :

$$\pi_1 = P(X_1 = S_1) = 0.6; \pi_2 = P(X_1 = S_2) = 0.4.$$

- Transition probabilities  $a_{ij} = P(X_{t+1} = S_j | X_t = S_i)$  for any  $t \in \mathbb{N}^+$ ,  $i = 1, 2$ , and  $j = 1, 2$ :

$$a_{11} = 0.7, a_{12} = 0.3; a_{21} = 0.4, a_{22} = 0.6.$$

- Emission probabilities  $b_{ik} = P(O_t = k | X_t = S_i)$  for any  $t \in \mathbb{N}^+$ ,  $i = 1, 2$ , and  $k \in \{A, C, G, T\}$ :

$$b_{1A} = 0.4, b_{1C} = 0.2, b_{1G} = 0.3, b_{1T} = 0.1;$$

$$b_{2A} = 0.2, b_{2C} = 0.4, b_{2G} = 0.1, b_{2T} = 0.3;$$

Assume we have an observed sequence  $\mathbf{O} = \overline{O_1 O_2 \cdots O_6} = ACCGTA$ , please answer the following questions with step-by-step computations and explanation for full credits. Your code should return all following answers when we run it.

- (a) **(5 points)** *Probability of an observed sequence.* Calculate  $P(\mathbf{O}; \Theta)$ .
- (b) **(5 points)** *Filtering.* Calculate  $P(X_6 = S_i | \mathbf{O}; \Theta)$  for  $i = 1, 2$ .
- (c) **(5 points)** *Smoothing.* Calculate  $P(X_4 = S_i | \mathbf{O}; \Theta)$  for  $i = 1, 2$ .
- (d) **(5 points)** *Most likely explanation.* Compute  $\mathbf{X} = \overline{X_1 X_2 \cdots X_6} = \arg \max_{\mathbf{X}} P(\mathbf{X} | \mathbf{O}; \Theta)$ .
- (e) **(5 points)** *Prediction.* Compute  $P(O_7 | \mathbf{O}; \Theta)$ . Then, which observation is most likely after  $o_{1:6}$ ? ( $O_7 = \arg \max_O P(O | \mathbf{O}; \Theta)$ ).

Answer:

- (a) **(5 points)**  $P(\mathbf{O}; \Theta) = 0.0002738928(\log: -8.20277376901)$ .

$$\begin{aligned}\alpha_1(j) &= P(X_1 = S_j, o_1) = P(o_1 | X_1 = S_j) P(X_1 = S_j) \\ \alpha_t(j) &= P(X_t = S_j, o_{1:t}) = P(o_t | X_t = S_j) \sum_i a_{ij} \alpha_{t-1}(i) \\ P(o_{1:T}) &= \sum_j \alpha_T(j)\end{aligned}$$

- (b) **(5 points)**  $P(X_6 = S_1 | \mathbf{O}; \Theta) = 0.67355987452$ ,  $P(X_6 = S_2 | \mathbf{O}; \Theta) = 0.32644012548$ .

$$\begin{aligned}\beta_T(j) &= 1 \text{ for any } j \\ \beta_{t-1}(i) &= P(o_{t:T} | X_{t-1} = S_i) = \sum_j \beta_t(j) a_{ij} P(o_t | X_t = S_j) \\ \gamma_t(j) &= P(X_t = S_j | o_{1:T}) = \frac{\alpha_t(j) \beta_t(j)}{\sum_j' \alpha_t(j') \beta_t(j')}\end{aligned}$$

- (c) **(5 points)**  $P(X_4 = S_1 | \mathbf{O}; \Theta) = 0.705017437479$ ,  $P(X_4 = S_2 | \mathbf{O}; \Theta) = 0.294982562521$ .

$$\gamma_t(j) = P(X_t = S_j | o_{1:T}) = \frac{\alpha_t(j) \beta_t(j)}{\sum_j' \alpha_t(j') \beta_t(j')}$$

- (d) **(5 points)**  $\mathbf{X} = \overline{X_1 X_2 \cdots X_6} = \arg \max_{\mathbf{X}} P(\mathbf{X} | \mathbf{O}; \Theta) = S_1 S_1 S_2 S_1 S_2 S_1$ .  $\delta_t(j)$  is the probability of the most likely path ending with  $j$  at time  $t$ .

$$\begin{aligned}\delta_t(j) &= \max_{x_1, x_2, \dots, x_{t-1}} P(X_1 = x_1, X_2 = x_2, \dots, X_{t-1} = x_{t-1}, X_t = S_j, o_{1:t} | \Theta) \\ &= \max_i \delta_{t-1}(i) a_{ij} P(o_t | X_t = S_j)\end{aligned}$$

Thus,  $\arg \max_j \delta_t(j)$  tells which state is most likely at time  $t$  given  $o_{1:t}$ .

- (e) **(5 points)**  $O_7 = \arg \max_O P(O | \mathbf{O}; \Theta) = A$ .  
 $P(O_7 = A) = 0.3204$ ,  $P(O_7 = C) = 0.2796$ ,  $P(O_7 = G) = 0.2204$ ,  $P(O_7 = T) = 0.1796$ .  $P(O_7 = A)$  is the highest probability

alpha:

```
[[ 2.40000000e-01,  8.00000000e-02],
 [ 4.00000000e-02,  4.80000000e-02],
```

```

[ 9.44000000e-03, 1.63200000e-02],
[ 3.94080000e-03, 1.26240000e-03],
[ 3.26352000e-04, 5.81904000e-04],
[ 1.84483200e-04, 8.94096000e-05]]
beta:
[[ 7.99764000e-04, 1.02436800e-03],
 [ 2.87580000e-03, 3.30960000e-03],
 [ 1.22100000e-02, 9.72000000e-03],
 [ 4.90000000e-02, 6.40000000e-02],
 [ 3.40000000e-01, 2.80000000e-01],
 [ 1.00000000e+00, 1.00000000e+00]]
gamma:
[[ 0.70079739, 0.29920261],
 [ 0.41998913, 0.58001087],
 [ 0.42083034, 0.57916966],
 [ 0.70501744, 0.29498256],
 [ 0.40512084, 0.59487916],
 [ 0.67355987, 0.32644013]]
delta:
[[ 2.40000000e-01, 8.00000000e-02],
 [ 3.36000000e-02, 2.88000000e-02],
 [ 4.70400000e-03, 6.91200000e-03],
 [ 9.87840000e-04, 4.14720000e-04],
 [ 6.91488000e-05, 8.89056000e-05],
 [ 1.93616640e-05, 1.06686720e-05]]
Prob (d): [0 0 1 0 1 0] 1.93616640e-05
Prob (e):
Prob of next state: [0.6021, 0.3979]
Prob of next observatoin: [0.3204, 0.2796, 0.2204, 0.1796]

```