**Mrs. Rahima Begam**
Age / Sex : 61 YRS / F
Referred by : INDRA KIDNEY CARE & SURGICAL CENTER
Reg. no. : 6960

6960
Registered on : 19/10/2026 07:30 PM
Collected on : 19/10/2026 06:00 PM
Received on : 19/10/2026 06:02 PM
Reported on : 19/10/2026 07:30 PM

Scan to download

## SEROLOGY & IMMUNOLOGY

### VIRAL MARKER

| TEST | VALUE | UNIT | REFERENCE |
|---|---|---|---|
| HIV (CARD TEST) | | | |
| HIV - 1 | NEGATIVE | | |
| HIV - 2 | NEGATIVE | | |
| **HEPATITIS C VIRUS, HCV** | **WEAKLY POSITIVE** | | |
| HBSAG | NEGATIVE | | |
| *Notes* | *RECOMMENDA HCV ELISA/ RNA FOR CONFIRMATION* | | |

**HbsAg** is the first serologic marker appearing in the serum 6-16 weeks following hepatitis B viral infection. In typical HBV infection, HBsAg will be detected 2-4 weeks before the liver enzyme levels (ALT) become abnormal and 3-5 weeks before the patient develops jaundice. In acute cases HbsAg usually disappears 1-2 months after the onset of symptoms. Persistence of HbsAg for more than 6 months indicates development of either a chronic carrier state or chronic liver disease. The presence of HbsAg is frequently associated with infectivity. HbsAg when accompanied by Hepatitis Be antigen and/or hepatitis B viral DNA almost always indicates infectivity.

**Anti HCV card test interpretation**
Reactive test result indicates presence of Hepatitis C virus infection. Active infection to be confirmed by HCV RNA PCR test. It cannot differentiate between the stages of Hepatitis C viral infection nor used to monitor the efficacy of treatment. Non-Reactive test result indicates Hepatitis C virus infection is unlikely.

**HIV Interpretation**
A negative result implies that no Anti HIV-1 & HIV-2 antibodies have been detected in the sample by this method. This means that either the patient has not been exposed to HIV-1 or HIV-2 infection or the sample has been tested during the “WINDOW PHASE” (before the development of detectable levels of antibodies). A positive result suggests the possibilities of HIV-I and / or HIV-II infection. However these results must be verified by a confirmatory test (IFA / WESTERN BLOT I-II) before pronouncing the patient positive for HIV-1 and / or HIV-2 infection. All reactive samples should be confirmed by using HIV Western Blot/PCR.

~~~ End of report ~~~

Powered by Labsmart
~~~