Reg. No. RMEE2552994
PURE PATH LAB
Add: Infront of Cantabil, Beside Distt. Hospital-Gonda
7703021000, 7897319900
www.purepathlab.in Gmail: help@purepathlab.in
PURE PATH LAB
Palce of Accuracy & Your Satisfaction
Dr. Abha Mishra
M.B.B.S (M.D. PATH)
Reg. No-MP-8779

**Mr. Buddhu**
Age / Sex : 49 YRS / Male
Reg. no. : 6950

Registered on : -
Collected on : -
Received on : -
Reported on : -

## SEROLOGY & IMMUNOLOGY
### VIRAL MARKER

| TEST | VALUE | UNIT | REFERENCE |
|---|---|---|---|
| HIV - 1 | NEGATIVE | - | - |
| HIV - 2 | NEGATIVE | - | - |
| HEPATITIS C VIRUS, HCV | NEGATIVE | - | - |
| HBSAG | NEGATIVE | - | - |

**HbsAg**

HbsAg is the first serologic marker appearing in the serum 6-16 weeks following hepatitis B viral infection. In typical HBV infection, HBsAg will be detected 2-4 weeks before the liver enzyme levels (ALT) become abnormal and 3-5 weeks before the patient develops jaundice. In acute cases HbsAg usually disappears 1-2 months after the onset of symptoms. Persistence of HbsAg for more than 6 months indicates development of either a chronic carrier state or chronic liver disease. The presence of HbsAg is frequently associated with infectivity. HbsAg when accompanied by Hepatitis Be antigen and/or hepatitis B viral DNA almost always indicates infectivity.

**Anti HCV card test interpretation**

Reactive test result indicates presence of Hepatitis C virus infection. Active infection to be confirmed by HCV RNA PCR test. It cannot differentiate between the stages of Hepatitis C viral infection nor used to monitor the efficacy of treatment. Non-Reactive test result indicates Hepatitis C virus infection is unlikely.

**HIV Interpretation**

A negative result implies that no Anti HIV-1 & HIV-2 antibodies have been detected in the sample by this method. This means that either the patient has not been exposed to HIV-1 or HIV-2 infection or the sample has been tested during the “WINDOW PHASE” (before the development of detectable levels of antibodies). A positive result suggests the possibilities of HIV-I and / or HIV-II infection. However these results must be verified by a confirmatory test (IFA / WESTERN BLOT I-II) before pronouncing the patient positive for HIV-1 and / or HIV-2 infection. All reactive samples should be confirmed by using HIV Western Blot/PCR.

## BIOCHEMISTRY
### BIOCHEMISTRY

| TEST | VALUE | UNIT | REFERENCE |
|---|---|---|---|
| RANDOM BLOOD SUGAR | 105.6 | mg/dl | 70 - 140 |

## BIOCHEMISTRY
### KIDNEY FUNCTION TEST (KFT)

| TEST | | VALUE | UNIT | REFERENCE |
|---|---|---|---|---|
| **SERUM UREA** | **H** | **116.97** | **mg/dl** | **19 - 45** |
| **SERUM CREATININE** | **H** | **10.46** | **mg/dl** | **0.72 - 1.43** |
| **SERUM PHOSPHORUS** | **H** | **7.39** | **mg/dl** | **2.5 - 4.5** |
| **SERUM URIC ACID** | **H** | **7.24** | **mg/dl** | **3.5 - 7.2** |
| SERUM SODIUM | | 136.5 | mmol/L | 136 - 146 |
| **SERUM POTASSIUM** | **H** | **5.25** | **mmol/L** | **3.5 - 5.1** |
| **ICALCIUM** | **L** | **1.08** | **mmol/l** | **1.13 - 1.33** |

Creatinine is a nitrogenous waste product formed in muscle from creatine phosphate. Endogenous production of creatinine is proportional to muscle mass and body weight.

| Causes of Increased Serum Creatinine Level | Causes of Decreased Serum Creatinine Level |
|---|---|
| 1. Pre-renal, renal, and post-renal azotemia | 1. Pregnancy |
| 2. Large amount of dietary meat | 2. Increasing age (reduction in muscle mass) |
| 3. Active acromegaly and gigantism | |

~~~ End of report ~~~
~~~