

# Lax-Idempotent 2-Monads, Degrees of Relatedness, and Multilevel Type Theory

**Andreas Nuyts**

KU Leuven, Belgium

TYPES '23  
Valencia, Spain  
June 12, 2023

# Degrees of Relatedness (RelDTT)

Nuyts and Devriese (2018) @ LICS

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright t$  indexed by degree  $i$ :
  - Just one for small types ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - More for larger types ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  $i$ -edges.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

- Parametricity is about **relations**,
- Equip types with multiple, **proof-relevant relations**  $s \rightsquigarrow_i t$  indexed by degree  $i$ :
  - Just one for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - More for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  $i$ -edges.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by degree  $i$ :
  - Just one for small types ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - More for larger types ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  $i$ -edges.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by **degree**  $i$ :
  - Just one for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - More for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  $i$ -edges.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by **degree**  $i$ :
  - Just **one** for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - **More** for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  $i$ -edges.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by **degree  $i$** :
  - Just **one** for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - **More** for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  **$i$ -edges**.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by **degree  $i$** :
  - Just **one** for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - **More** for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  **$i$ -edges**.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - parametricity
  - ad hoc polymorphism
  - . irrelevance
  - .. shape-irrelevance
  - aspects of algebra, unions, intersections,  $\text{Prop}$ , ...

# Degrees of Relatedness: Overview

- Parametricity is about **relations**,
- Equip types with **multiple, proof-relevant relations**  $s \curvearrowright_i t$  indexed by **degree  $i$** :
  - Just **one** for **small types** ( $\text{Bool}, \mathbb{N} \rightarrow \mathbb{N}, \dots$ ),
  - **More** for **larger types** ( $\mathbf{U}_0 \rightarrow \mathbf{U}_0, \text{Grp}, \dots$ ).
  - Proofs called  **$i$ -edges**.
- Describe **function behaviour** by saying how functions **influence degree** of relatedness,
- This explains
  - **parametricity**
  - **ad hoc polymorphism**
  - **. irrelevance**
  - **.. shape-irrelevance**
  - aspects of **algebra, unions, intersections, Prop, ...**

# Degrees of Relatedness: Four Laws

- **Reflexivity:**  $(a : A) \sim_i^A (a : A)$   
(Semantically, prop. eq. = def. eq.)
- **Degradation:**  $((a : A) \sim_i^R (b : B)) \rightarrow ((a : A) \sim_{i+1}^R (b : B))$
- **Dependency:**  $(a : A) \sim_i^R (b : B)$  presumes  $R : A \sim_{i+1}^U B$
- **Identity extension:**  $(a : A) \sim_0^A (b : A)$  means  $a = b : A$ .  
~ heterogeneous  $\sim_0$  serves as heterogeneous equality.

# Degrees of Relatedness: Four Laws

- **Reflexivity:**  $(a = b : A) \rightarrow (a : A) \sim_i^A (b : A)$   
(Semantically, prop. eq. = def. eq.)
- **Degradation:**  $((a : A) \sim_i^R (b : B)) \rightarrow ((a : A) \sim_{i+1}^R (b : B))$
- **Dependency:**  $(a : A) \sim_i^R (b : B)$  presumes  $R : A \sim_{i+1}^U B$
- **Identity extension:**  $(a : A) \sim_0^A (b : A)$  means  $a = b : A$ .  
~ heterogeneous  $\sim_0$  serves as heterogeneous equality.

# Degrees of Relatedness: Four Laws

- **Reflexivity:**  $(a = b : A) \rightarrow (a : A) \sim_i^A (b : A)$   
(Semantically, prop. eq. = def. eq.)
- **Degradation:**  $((a : A) \sim_i^R (b : B)) \rightarrow ((a : A) \sim_{i+1}^R (b : B))$
- **Dependency:**  $(a : A) \sim_i^R (b : B)$  presumes  $R : A \sim_{i+1}^U B$
- **Identity extension:**  $(a : A) \sim_0^A (b : A)$  means  $a = b : A$ .  
~ heterogeneous  $\sim_0$  serves as heterogeneous equality.

# Degrees of Relatedness: Four Laws

- **Reflexivity:**  $(a = b : A) \rightarrow (a : A) \sim_i^A (b : A)$   
(Semantically, prop. eq. = def. eq.)
- **Degradation:**  $((a : A) \sim_i^R (b : B)) \rightarrow ((a : A) \sim_{i+1}^R (b : B))$
- **Dependency:**  $(a : A) \sim_i^R (b : B)$  **presumes**  $R : A \sim_{i+1}^U B$
- **Identity extension:**  $(a : A) \sim_0^A (b : A)$  means  $a = b : A$ .  
~ heterogeneous  $\sim_0$  serves as heterogeneous equality.

# Degrees of Relatedness: Four Laws

- **Reflexivity:**  $(a = b : A) \rightarrow (a : A) \sim_i^A (b : A)$   
(Semantically, prop. eq. = def. eq.)
- **Degradation:**  $((a : A) \sim_i^R (b : B)) \rightarrow ((a : A) \sim_{i+1}^R (b : B))$
- **Dependency:**  $(a : A) \sim_i^R (b : B)$  **presumes**  $R : A \sim_{i+1}^U B$
- **Identity extension:**  $(a : A) \sim_0^A (b : A)$  means  $a = b : A$ .  
~ heterogeneous  $\sim_0$  serves as heterogeneous equality.

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation R.

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation P.

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation Q along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

V could specify that Q must respect e **and** \*  
(but it could ask Q to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation R.

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation P.

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation Q along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

V could specify that Q must respect e **and** \*  
(but it could ask Q to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation R.

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation P.

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation Q along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

V could specify that Q must respect e **and** \*  
(but it could ask Q to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation  $R$ .

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation  $P$ .

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation  $Q$  along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

$V$  could specify that  $Q$  must respect  $e$  and  $*$  (but it could ask  $Q$  to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation  $R$ .

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation  $P$ .

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation  $Q$  along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

$V$  could specify that  $Q$  must respect  $e$  and  $*$  (but it could ask  $Q$  to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation  $R$ .

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation  $P$ .

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation  $Q$  along ...

$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

$V$  could specify that  $Q$  must respect  $e$  **and**  $*$   
(but it could ask  $Q$  to be anything).

# Understanding degrees

$$(a : A) \sim_0^A (b : A)$$

Equality.

$$(a : A) \sim_0^R (b : B)$$

Heterogeneous equality along ...

$$R : (A : U_0) \sim_1^{U_0} (B : U_0)$$

Any relation  $R$ .

$$P : (G : \text{Grp}) \sim_1^{\text{Grp}} (H : \text{Grp})$$

Any logical/algebraic relation  $P$ .

$$Q : (G : \text{Grp}) \sim_1^V (M : \text{Monoid})$$

Any logical/algebraic relation  $Q$  along ...


$$V : (\text{Grp} : U_1) \sim_2^{U_1} (\text{Monoid} : U_1)$$

$V$  could specify that  $Q$  must respect  $e$  **and**  $*$   
(but it could ask  $Q$  to be anything).

# Understanding modalities (2)

**par** : types  $\rightarrow$  values


$$if : (\text{par} \dashv X : U_0) \rightarrow B X$$



**con** : types  $\rightarrow$  types

$$B : U_0 \rightarrow U_0$$


$$B X = \text{Bool} \rightarrow X \rightarrow X \rightarrow X$$



# Understanding modalities (2)

**par** : types → values


$$if : (\text{par} \dashv X : U_0) \rightarrow B X$$



**con** : types → types

$$B : U_0 \rightarrow U_0$$


$$B X = \text{Bool} \rightarrow X \rightarrow X \rightarrow X$$



# Understanding modalities (2)

**par** : types  $\rightarrow$  values


$$if : (\text{par} \dashv X : U_0) \rightarrow B X$$



**con** : types  $\rightarrow$  types

$$B : U_0 \rightarrow U_0$$


$$B X = \text{Bool} \rightarrow X \rightarrow X \rightarrow X$$



# Understanding modalities (2)

**par** : types  $\rightarrow$  values


$$if : (\text{par} \dashv X : U_0) \rightarrow B X$$



**con** : types  $\rightarrow$  types

$$B : U_0 \rightarrow U_0$$


$$B X = \text{Bool} \rightarrow X \rightarrow X \rightarrow X$$



## Understanding modalities (2)

**par** : types  $\rightarrow$  values


$$if : (\text{par} \dashv X : U_0) \rightarrow B X$$



**con** : types  $\rightarrow$  types

$$B : U_0 \rightarrow U_0$$


$$B X = \text{Bool} \rightarrow X \rightarrow X \rightarrow X$$



# Understanding modalities (1)

**irr** : values → values

$$[] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types


$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# Understanding modalities (1)

**irr** : values → values

$$[] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types


$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# Understanding modalities (1)


**irr** : values → values

$$[ ] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types


$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# Understanding modalities (1)


**irr** : values → values

$$[ ] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types


$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# Understanding modalities (1)


**irr** : values → values

$$[ ] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types


$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# Understanding modalities (1)


**irr** : values → values

$$[ ] : (\mathbf{irr} \perp n : \mathbb{N}) \rightarrow \text{List}_n A$$



**shi** : values → types

$$\lambda n. \text{List}_n A : (\mathbf{shi} \perp n : \mathbb{N}) \rightarrow U_0$$



# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$
- Modalities  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$  where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \rightsquigarrow_{i,\mu}^A y) \rightarrow f(x) \rightsquigarrow_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \rightsquigarrow_i^{(\mu|A)} \text{mod}_\mu y = x \rightsquigarrow_{i,\mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p+1$  different dimension flavours.

# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$

- Modalities  $\mu : p \rightarrow q$  are

functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$

where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \rightsquigarrow_{i \cdot \mu}^A y) \rightarrow f(x) \rightsquigarrow_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \rightsquigarrow_i^{(\mu|A)} \text{mod}_\mu y = x \rightsquigarrow_{i \cdot \mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p+1$  different dimension flavours.

# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$
- Modalities  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$  where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \frown_{i \cdot \mu}^A y) \rightarrow f(x) \frown_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \frown_i^{\langle \mu | A \rangle} \text{mod}_\mu y = x \frown_{i \cdot \mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p+1$  different dimension flavours.

# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$
- Modalities  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$  where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \frown_{i \cdot \mu}^A y) \rightarrow f(x) \frown_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \frown_i^{\langle \mu | A \rangle} \text{mod}_\mu y = x \frown_{i \cdot \mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p+1$  different dimension flavours.

# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$
- Modalities  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$  where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \frown_{i \cdot \mu}^A y) \rightarrow f(x) \frown_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \frown_i^{\langle \mu | A \rangle} \text{mod}_\mu y = x \frown_{i \cdot \mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p+1$  different dimension flavours.

# The mode theory DoR

ReIDTT can be built on an instance of MTT (Multimode Type Theory) with mode theory DoR:

- Modes are **depths**  $p \in \mathbb{Z}_{\geq -1}$
- Modalities  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq \top\} : i \mapsto i \cdot \mu$  where  $f : (\mu \dashv x : A) \rightarrow B(x)$  sends

$$(r : x \frown_{i \cdot \mu}^A y) \rightarrow f(x) \frown_i^{B(r)} f(y).$$

Modal types:

$$\text{mod}_\mu x \frown_i^{\langle \mu | A \rangle} \text{mod}_\mu y = x \frown_{i \cdot \mu}^A y$$

- 2-cells are degree-wise inequalities.

Depth  $p$  is modelled in cubical sets with  $p + 1$  different dimension flavours.

Depth

0

(Cubical  
sets)

-1


(Sets)

Depth

0  
(Cubical sets)




-1  
(Sets)




discrete  
 $\Delta$

+



global  
sections  
 $\Gamma$




+




codiscrete  
 $\nabla$

Depth

0  
(Cubical  
sets)



-1  
(Sets)




discrete  
 $\Delta$

global  
sections  
 $\Gamma$

codiscrete  
 $\nabla$


Depth

$0$   
(Cubical  
sets)




$-1$   
(Sets)

discrete  
 $\Delta$



global  
sections  
 $\Gamma$



codiscrete  
 $\nabla$


Depth

0  
(Cubical sets)




$$\{ \circ \} \quad \downarrow -\Delta \quad \{ (=) \leq T \}$$


-1  
(Sets)



discrete  
 $\Delta$



$$\{ (=) \leq 0 \leq T \} \quad \downarrow \quad \{ \}$$



$$\{ \circ \}$$

$$\{ (=) \leq T \}$$

codiscrete  
 $\nabla$

Depth


0  
(Cubical sets)




$$\{\circ\} \xrightarrow{[-\Delta]} \{(\equiv) \leq T\}$$



-1  
(Sets)




discrete  
 $\Delta$



$$\{\Gamma| \xrightarrow{T} \} \xleftarrow{[-\Gamma]} \{\}\}$$



global  
sections  
 $\Gamma$



$$\{\circ\}$$

$$\{(\equiv) \leq T\}$$

codiscrete  
 $\nabla$

Depth


0  
(Cubical sets)



$$\{\circ\} \xrightarrow{[-\Delta]} \{(\equiv) \leq T\}$$




-1  
(Sets)




discrete  
 $\Delta$



$$\{\Gamma| \xrightarrow{[-\Gamma]} \{(\equiv) \leq 0 \leq T\}\}$$



global  
sections  
 $\Gamma$



$$\{\circ\}$$

$$\nabla \xrightarrow{[-\nabla]} \{(\equiv) \leq T\}$$

$$\{(\equiv) \leq T\}$$

codiscrete  
 $\nabla$


Depth

0  
(Cubical sets)



$$\{\circ\} \xrightarrow{-\Delta} \{(\equiv) \leq T\}$$

-1  
(Sets)




discrete  
 $\Delta$



$$\{(\equiv) \leq 0 \leq T\} \xrightarrow{-\Gamma} \{\}\xrightarrow{\quad}$$

global  
sections  
 $\Gamma$




$$\{\circ\} \xrightarrow{-\nabla} \{(\equiv) \leq T\}$$

codiscrete  
 $\nabla$

Depth

$0$   
(Cubical sets)




$$\langle \Delta I \rightarrow \uparrow \rangle \quad \downarrow \quad \{ \circ \}$$

$$\{ (=) \leqslant T \}$$

$-1$   
(Sets)




discrete  
 $\Delta$



$$\langle \Gamma I \rightarrow \downarrow \rangle \quad \uparrow \quad \{ (=) \leqslant 0 \leqslant T \}$$

$$\{ \} \quad \downarrow \quad \langle -\Gamma \rangle$$

global  
sections  
 $\Gamma$



$$\langle \nabla I \rightarrow \uparrow \rangle \quad \downarrow \quad \{ \circ \}$$

$$\{ (=) \leqslant T \} \quad \uparrow \quad \langle -\nabla \rangle$$

codiscrete  
 $\nabla$

## Two-Level Type Theory (2LTT)

Voevodsky (2013)

Altenkirch, Capriotti & Kraus (2016)

Annenkov, Capriotti, Kraus & Sattler (2017)

More specifically, its general presheaf model

## Two-Level Type Theory (2LTT)

Voevodsky (2013)

Altenkirch, Capriotti & Kraus (2016)

Annenkov, Capriotti, Kraus & Sattler (2017)

More specifically, **its general presheaf model**

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\mathbf{CwF}(\mathcal{C}, \mathbf{Ty}^i, \mathbf{Tm}^i)$

$\mathbf{Ty}^i \in \mathbf{Psh}(\mathcal{C})$

$\mathbf{Tm}^i \in \mathbf{Psh}(\int_{\mathcal{C}} \mathbf{Ty}^i)$

$\Gamma \mathbf{ctx}^i$

$\Gamma \vdash^i T \mathbf{type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\mathbf{CwF}(\mathbf{Psh}(\mathcal{C}), \mathbf{Ty}^o, \mathbf{Tm}^o)$

$\vdash^o \mathbf{Ty}^i \mathbf{type}$

$X : \mathbf{Ty}^i \vdash^o \mathbf{Tm}^i(X) \mathbf{type}$

$\Rightarrow \mathbf{y}\Gamma \mathbf{ctx}^o$

$\Leftrightarrow \mathbf{y}\Gamma \vdash^o T : \mathbf{Ty}^i$

$\Leftrightarrow \mathbf{y}\Gamma \vdash^o t : \mathbf{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\mathbf{CwF}(\mathcal{C}, \mathbf{Ty}^i, \mathbf{Tm}^i)$

$\mathbf{Ty}^i \in \mathbf{Psh}(\mathcal{C})$

$\mathbf{Tm}^i \in \mathbf{Psh}(\int_{\mathcal{C}} \mathbf{Ty}^i)$

$\Gamma \mathbf{ctx}^i$

$\Gamma \vdash^i T \mathbf{type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\mathbf{CwF}(\mathbf{Psh}(\mathcal{C}), \mathbf{Ty}^o, \mathbf{Tm}^o)$

$\vdash^o \mathbf{Ty}^i \mathbf{type}$

$X : \mathbf{Ty}^i \vdash^o \mathbf{Tm}^i(X) \mathbf{type}$

$\Rightarrow \mathbf{y} \Gamma \mathbf{ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \mathbf{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \mathbf{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow y \Gamma \text{ ctx}^o$

$\Leftrightarrow y \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow y \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$

$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality

## Inner Level

$\text{CwF}(\mathcal{C}, \text{Ty}^i, \text{Tm}^i)$

$\text{Ty}^i \in \text{Psh}(\mathcal{C})$

$\text{Tm}^i \in \text{Psh}(\int_{\mathcal{C}} \text{Ty}^i)$

$\Gamma \text{ ctx}^i$

$\Gamma \vdash^i T \text{ type}$

$\Gamma \vdash^i t : T$

E.g inner system = HoTT

## Outer Level

$\text{CwF}(\text{Psh}(\mathcal{C}), \text{Ty}^o, \text{Tm}^o)$

$\vdash^o \text{Ty}^i \text{ type}$

$X : \text{Ty}^i \vdash^o \text{Tm}^i(X) \text{ type}$


$\Rightarrow \mathbf{y} \Gamma \text{ ctx}^o$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o T : \text{Ty}^i$

$\Leftrightarrow \mathbf{y} \Gamma \vdash^o t : \text{Tm}^i(T)$

Talk about strict equality


## Lifting Functors



## Multilevel Type Theory

- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .

## Lifting Functors




## Multilevel Type Theory

- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .


## Lifting Functors

## Multilevel Type Theory




- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .

## Lifting Functors




## Multilevel Type Theory




- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .

## Lifting Functors




## Multilevel Type Theory




- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .

## Lifting Functors



- $F^*$  = precomposition by  $F$ .
- $F_!$  extends  $F$  along  $y$ , i.e.  $F_!y \cong yF$ .

## Multilevel Type Theory



## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders  
are

simplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $y : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $y^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders  
are

simplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{x}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders  
are

simplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders

are

simplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders

are

simplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders  
are

semisimplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders sets

are

symmetric  
semisimplicial sets

## Multilevel TT: Case Study

- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $y : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $y^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?

Presheaves over  
finite  
non-empty  
linear orders sets

are

symmetric  
semisimplicial sets  
with  $\infty$ -dimensional simplices

## Multilevel TT: Case Study


- Level 0:  $\star$  (point category is a CwF)
- Level 1:  $\text{Psh}(\star) \cong \text{Set}$ 
  - $\mathbf{y} : \star \rightarrow \text{Set} : \bullet \mapsto \top$
- Level 2:  $\text{Psh}(\text{Psh}(\star)) \cong \text{Psh}(\text{Set})$ 
  - $\mathbf{y}^* : \text{Psh}(\text{Set}) \rightarrow \text{Set}$  extracts points.

## What is $\text{Psh}(\text{Set})$ ?


Presheaves over  
finite  
non-empty  
linear orders sets


are


symmetric  
semisimplicial sets  
with  $\infty$ -dimensional simplices




# Cohesion in Multilevel TT









Degrees of Relatedness  $\stackrel{?}{\sim}$  Multilevel TT

Goal: Formalize this correspondence, using **interface** of

Lax-idempotent 2-monads

Degrees of Relatedness  $\stackrel{?}{\sim}$  Multilevel TT

Goal: Formalize this correspondence, using interface of

Lax-idempotent 2-monads

Degrees of Relatedness  $\stackrel{?}{\sim}$  Multilevel TT

Goal: Formalize this correspondence, using **interface** of

Lax-idempotent 2-monads

# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad


- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like  $\text{List } X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

It's a  $\mathbb{2}$ -monad!

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_*$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
$$\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$$


# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like  $\text{List } X$  is freely generated by  $[\cdot] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .
- It's a  $(\mathbb{2}\text{-})$ -monad!
  - Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$



# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
  - Presheaves are colimits of codiscrete presheaves.
  - $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**. (Free cocompletion.)
    - Like  $\text{List } X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .
- It's a  $\mathbb{2}$ -monad!
- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_*$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like  $\text{List } X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

It's a (2-)monad!

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like  $\text{List } X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**. (Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**. (Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$




# Why $\text{Psh} : \text{Cat} \rightarrow \text{Cat}$ is a Monad

- In  $\text{Psh}(\text{Set})$ , Yoneda-embeddings (representables) are **codiscrete**.
- Presheaves are colimits of codiscrete presheaves.
- $\text{Psh}(\mathcal{C})$  is **freely** generated by  $\mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$  and **colimits**.  
(Free cocompletion.)
  - Like List  $X$  is freely generated by  $[-] : X \rightarrow \text{List } X$  and  $[]$  and  $++$ .

**It's a (2-)monad!**

- Unit  $\eta = \mathbf{y} : \mathcal{C} \rightarrow \text{Psh}(\mathcal{C})$ ,
  - (Pseudo)functorial action:  $F \mapsto F_!$ ,
  - $\mu = \mathbf{y}^* : \text{Psh}(\text{Psh}(\mathcal{C})) \rightarrow \text{Psh}(\mathcal{C})$
- It's a **lax-idempotent** 2-monad:  
 $\text{Psh} \eta \dashv \mu \dashv \eta \text{ Psh}$



## Definition

Define 2-category DoR:

- Objects are  $p \in \mathbb{Z}_{\geq -1}$
- Morphisms  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq T\} : i \mapsto i \cdot \mu$
- 2-cells are degree-wise inequalities.
- Freely add  $\perp =: -2$ .

## Definition

Let LIM be the 2-category freely generated by:

- $\mathbf{C} \in \text{Obj}(\text{LIM})$ ,
- Lax-idempotent 2-monad  $\mathbf{M} : \text{LIM} \rightarrow \text{LIM}$

Clearly, we get  $[-]_{\mathcal{C}} : \text{LIM} \rightarrow \text{Cat}$  for any category  $\mathcal{C}$ :

- $\mathbf{C} \mapsto \mathcal{C}$ ,
- $\mathbf{M} \mapsto \text{Psh}$ .

Main theorem (formal proof WIP)

$$\begin{array}{ccccc} \text{DoR} & \cong & \text{LIM} & \rightarrow & \text{Cat} \\ p & \mapsto & \mathbf{M}^{p+2}(\mathbf{C}) & \mapsto & \text{Psh}^{p+2}(\mathcal{C}) \end{array}$$

## Definition

Define 2-category DoR:

- Objects are  $p \in \mathbb{Z}_{\geq -1}$
- Morphisms  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq T\} : i \mapsto i \cdot \mu$
- 2-cells are degree-wise inequalities.
- Freely add  $\perp =: -2$ .

## Definition

Let LIM be the 2-category freely generated by:

- $\mathbf{C} \in \text{Obj(LIM)}$ ,
- Lax-idempotent 2-monad  $\mathbf{M} : \text{LIM} \rightarrow \text{LIM}$

Clearly, we get  $[-]_{\mathcal{C}} : \text{LIM} \rightarrow \text{Cat}$  for any category  $\mathcal{C}$ :

- $\mathbf{C} \mapsto \mathcal{C}$ ,
- $\mathbf{M} \mapsto \text{Psh}$ .

Main theorem (formal proof WIP)

$$\begin{array}{ccccc} \text{DoR} & \cong & \text{LIM} & \rightarrow & \text{Cat} \\ p & \mapsto & \mathbf{M}^{p+2}(\mathbf{C}) & \mapsto & \text{Psh}^{p+2}(\mathcal{C}) \end{array}$$

## Definition

Define 2-category DoR:

- Objects are  $p \in \mathbb{Z}_{\geq -1}$
- Morphisms  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq T\} : i \mapsto i \cdot \mu$
- 2-cells are degree-wise inequalities.
- Freely add  $\perp =: -2$ .

## Definition

Let LIM be the 2-category freely generated by:

- $\mathbf{C} \in \text{Obj}(\text{LIM})$ ,
- Lax-idempotent 2-monad  $\mathbf{M} : \text{LIM} \rightarrow \text{LIM}$

Clearly, we get  $[-]_{\mathcal{C}} : \text{LIM} \rightarrow \text{Cat}$  for any category  $\mathcal{C}$ :

- $\mathbf{C} \mapsto \mathcal{C}$ ,
- $\mathbf{M} \mapsto \text{Psh}$ .

Main theorem (formal proof WIP)

$$\begin{array}{ccccc} \text{DoR} & \cong & \text{LIM} & \rightarrow & \text{Cat} \\ p & \mapsto & \mathbf{M}^{p+2}(\mathbf{C}) & \mapsto & \text{Psh}^{p+2}(\mathcal{C}) \end{array}$$

## Definition

Define 2-category DoR:

- Objects are  $p \in \mathbb{Z}_{\geq -1}$
- Morphisms  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq T\} : i \mapsto i \cdot \mu$
- 2-cells are degree-wise inequalities.
- Freely add  $\perp =: -2$ .

## Definition

Let LIM be the 2-category freely generated by:

- $\mathbf{C} \in \text{Obj}(\text{LIM})$ ,
- Lax-idempotent 2-monad  $\mathbf{M} : \text{LIM} \rightarrow \text{LIM}$

Clearly, we get  $[-]_{\mathcal{C}} : \text{LIM} \rightarrow \text{Cat}$  for any category  $\mathcal{C}$ :

- $\mathbf{C} \mapsto \mathcal{C}$ ,
- $\mathbf{M} \mapsto \text{Psh}$ .

Main theorem (formal proof WIP)

$$\begin{array}{ccccc} \text{DoR} & \cong & \text{LIM} & \rightarrow & \text{Cat} \\ p & \mapsto & \mathbf{M}^{p+2}(\mathbf{C}) & \mapsto & \text{Psh}^{p+2}(\mathcal{C}) \end{array}$$

## Definition

Define 2-category DoR:

- Objects are  $p \in \mathbb{Z}_{\geq -1}$
- Morphisms  $\mu : p \rightarrow q$  are functions  $\{0 \leq \dots \leq q\} \rightarrow \{(=) \leq 0 \leq \dots \leq p \leq T\} : i \mapsto i \cdot \mu$
- 2-cells are degree-wise inequalities.
- Freely add  $\perp =: -2$ .

## Definition

Let LIM be the 2-category freely generated by:


- $\mathbf{C} \in \text{Obj}(\text{LIM})$ ,
- Lax-idempotent 2-monad  $\mathbf{M} : \text{LIM} \rightarrow \text{LIM}$

Clearly, we get  $[-]_{\mathcal{C}} : \text{LIM} \rightarrow \text{Cat}$  for any category  $\mathcal{C}$ :


- $\mathbf{C} \mapsto \mathcal{C}$ ,
- $\mathbf{M} \mapsto \text{Psh}$ .

## Main theorem (formal proof WIP)

$$\begin{array}{ccccc} \text{DoR} & \cong & \text{LIM} & \rightarrow & \text{Cat} \\ p & \mapsto & \mathbf{M}^{p+2}(\mathbf{C}) & \mapsto & \text{Psh}^{p+2}(\mathcal{C}) \end{array}$$



# Sketch of Proof



Implications of main theorem:

- **Degrees of Relatedness\*** can serve as an **internal language** for **multilevel type theory**,
- A model for **parametricity, irrelevance, ... found in the wild.**

To do:

- Formalize proof.
- Study **discreteness** and **internal parametricity** in this setting.
- Directify :-)

\*Or a reasonable adaptation.

Implications of main theorem:

- **Degrees of Relatedness\*** can serve as an **internal language** for **multilevel type theory**,
- A model for **parametricity, irrelevance, ...** found **in the wild**.

To do:

- Formalize proof.
- Study discreteness and internal parametricity in this setting.
- Directify :-)

\*Or a reasonable adaptation.

Implications of main theorem:

- **Degrees of Relatedness\*** can serve as an **internal language** for **multilevel type theory**,
- A model for **parametricity, irrelevance, ...** found **in the wild**.

To do:

- Formalize proof.
- Study discreteness and internal parametricity in this setting.
- Directify :-)

\*Or a reasonable adaptation.

Implications of main theorem:

- **Degrees of Relatedness\*** can serve as an **internal language** for **multilevel type theory**,
- A model for **parametricity, irrelevance, ...** found **in the wild**.

To do:

- Formalize proof.
- Study **discreteness** and **internal parametricity** in this setting.
- Directify :-)

\*Or a reasonable adaptation.

Implications of main theorem:

- **Degrees of Relatedness\*** can serve as an **internal language** for **multilevel type theory**,
- A model for **parametricity, irrelevance, ...** found **in the wild**.

To do:

- Formalize proof.
- Study **discreteness** and **internal parametricity** in this setting.
- Directify :-)

\*Or a reasonable adaptation.

**Thanks!**

**Questions?**