

**Alberto Andrés Valdés González.**

**Degree:** Mathematical Engineer.

**Work position:** Data Scientist.


**Mail:** [anvaldes@uc.cl](mailto:anvaldes@uc.cl)/[alberto.valdes.gonzalez.96@gmail.com](mailto:alberto.valdes.gonzalez.96@gmail.com)

**Location:** Santiago, Chile.

---

## Fourier Series

A Fourier series is an expansion of a periodic function into a sum of trigonometric functions. The Fourier series is an example of a trigonometric series, but not all trigonometric series are Fourier series.



$$f(t) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[ a_n \cdot \cos \left( \frac{2n\pi t}{T} \right) + b_n \cdot \sin \left( \frac{2n\pi t}{T} \right) \right]$$

$$a_0 = \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt$$

$$a_n = \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \cos \left( \frac{2n\pi t}{T} \right) dt$$

$$b_n = \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \sin \left( \frac{2n\pi t}{T} \right) dt$$

**Integration by parts:**

$$\int_a^b h_1'(t) \cdot h_2(t) dt = h_1(t) \cdot h_2(t) \Big|_{t=a}^{t=b} - \int_a^b h_1(t) \cdot h_2'(t) dt$$

**Example:** Find the fourier series for  $f(t) = t$  for  $t \in [-\pi, \pi]$ .

Solution:

We have  $T = 2\pi$ .

$$a_0 = \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t dt = \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t dt = \frac{1}{\pi} \cdot \frac{t^2}{2} \Big|_{t=-\pi}^{t=\pi} = 0$$

$\Rightarrow$

$$\boxed{a_0 = 0} \quad (1)$$

$$\begin{aligned} a_n &= \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \cos \left( \frac{2n\pi t}{T} \right) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t \cdot \cos \left( \frac{2n\pi t}{2\pi} \right) dt \\ &= \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t \cdot \cos(nt) dt \end{aligned}$$

You can see that  $g(t) = t \cdot \cos(nt)$  is odd function:

$$g(-t) = (-t) \cdot \cos(-nt) = -t \cdot \cos(nt) = -g(t)$$

$$\boxed{g(t) = -g(-t)}$$

For this reason:

$$\boxed{a_n = 0} \quad (2)$$

$$\begin{aligned} b_n &= \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \sin\left(\frac{2n\pi t}{T}\right) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t \cdot \sin\left(\frac{2n\pi t}{2\pi}\right) dt \\ &= \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t \cdot \sin(nt) dt \end{aligned}$$

$$\boxed{b_n = \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t \cdot \sin(nt) dt}$$

Using integral by parts:

$$h_1(t) = t \Rightarrow h_1'(t) = 1$$

$$h_2'(t) = \sin(nt) \Rightarrow h_2(t) = -\frac{\cos(nt)}{n}$$

$\Rightarrow$


$$\begin{aligned} b_n &= \frac{1}{\pi} \cdot \left[ -\frac{t \cdot \cos(nt)}{n} - \left( -\frac{\sin(nt)}{n^2} \right) \right] \Big|_{t=-\pi}^{t=\pi} = \frac{1}{\pi} \cdot \left[ -\frac{t \cdot \cos(nt)}{n} + \frac{\sin(nt)}{n^2} \right] \Big|_{t=-\pi}^{t=\pi} \\ &= \frac{1}{\pi} \cdot \left[ -\frac{\pi \cdot \cos(n \cdot \pi)}{n} + \frac{\sin(n \cdot \pi)}{n^2} + \frac{(-\pi) \cdot \cos(-n \cdot \pi)}{n} - \frac{\sin(-n \cdot \pi)}{n^2} \right] \\ &= \frac{1}{\pi} \cdot \left[ -\frac{\pi \cdot \cos(n \cdot \pi)}{n} + \frac{(-\pi) \cdot \cos(-n \cdot \pi)}{n} \right] = \frac{1}{\pi} \cdot \left[ -\frac{\pi \cdot \cos(n \cdot \pi)}{n} - \frac{\pi \cdot \cos(n \cdot \pi)}{n} \right] \\ &= -\frac{2 \cdot \pi \cdot \cos(n \cdot \pi)}{n \cdot \pi} = -\frac{2 \cdot \cos(n \cdot \pi)}{n} = -\frac{2 \cdot (-1)^n}{n} = \frac{2 \cdot (-1)^{(n+1)}}{n} \end{aligned}$$

$\Rightarrow$

$$\boxed{b_n = \frac{2 \cdot (-1)^{(n+1)}}{n}} \quad (3)$$

$$f(t) = \sum_{n=1}^{\infty} \frac{2 \cdot (-1)^{(n+1)}}{n} \cdot \sin(n \cdot t)$$

$$f(t) = 2\sin(t) - \sin(2t) + \frac{2}{3} \cdot \sin(3t) - \frac{1}{2} \cdot \sin(4t) + \frac{2}{5} \cdot \sin(5t) + \dots$$



**Example:** Find the fourier series for  $f(t) = t^2$  for  $t \in [-\pi, \pi]$ .

Solution:

We have  $T = 2\pi$ .

$$\begin{aligned} a_0 &= \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t^2 dt = \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t^2 dt = \frac{1}{\pi} \cdot \left. \frac{t^3}{3} \right|_{t=-\pi}^{t=\pi} \\ &= \frac{1}{\pi} \cdot \frac{\pi^3}{3} - \frac{1}{\pi} \cdot \frac{(-\pi)^3}{3} = \frac{1}{\pi} \cdot \frac{\pi^3}{3} + \frac{1}{\pi} \cdot \frac{\pi^3}{3} = \frac{2\pi^2}{3} \end{aligned}$$

$\Rightarrow$

$$\boxed{a_0 = \frac{2\pi^2}{3} \quad , \quad \frac{a_0}{2} = \frac{\pi^2}{3}} \quad (1)$$

$$\begin{aligned}
a_n &= \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \cos\left(\frac{2n\pi t}{T}\right) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t^2 \cdot \cos\left(\frac{2n\pi t}{2\pi}\right) dt \\
&= \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t^2 \cdot \cos(nt) dt
\end{aligned}$$

$\Rightarrow$

$$a_n = \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t^2 \cdot \cos(nt) dt$$

Now we going to compute the integral using integration by parts.

$$h_1(t) = t^2 \Rightarrow h_1'(t) = 2t$$

$$h_2'(t) = \cos(nt) \Rightarrow h_2(t) = \frac{\sin(nt)}{n}$$

$\Rightarrow$

$$\int t^2 \cos(nt) dt = \frac{t^2 \cdot \sin(nt)}{n} - \int \frac{2t \cdot \sin(nt)}{n} dt = \frac{t^2 \cdot \sin(nt)}{n} - \frac{2}{n} \int t \cdot \sin(nt) dt$$

$\Rightarrow$

$$\boxed{\int t^2 \cos(nt) dt = \frac{t^2 \cdot \sin(nt)}{n} - \frac{2}{n} \cdot \int t \cdot \sin(nt) dt} \quad (i)$$

Now we going to use again integral by parts:

$$h_1(t) = t \Rightarrow h_1'(t) = 1$$

$$h_2'(t) = \sin(nt) \Rightarrow h_2(t) = -\frac{\cos(nt)}{n}$$

$$\begin{aligned}
\int t \cdot \sin(nt) dt &= -\frac{t \cdot \cos(nt)}{n} - \int -\frac{\cos(nt)}{n} dt \\
&= -\frac{t \cdot \cos(nt)}{n} + \frac{1}{n} \cdot \int \cos(nt) dt = -\frac{t \cdot \cos(nt)}{n} + \frac{1}{n} \cdot \frac{\sin(nt)}{n} \\
&= -\frac{t \cdot \cos(nt)}{n} + \frac{\sin(nt)}{n^2}
\end{aligned}$$

$\Rightarrow$

$$\begin{aligned}\int t^2 \cos(nt) dt &= \frac{t^2 \cdot \sin(nt)}{n} - \frac{2}{n} \cdot \left( -\frac{t \cdot \cos(nt)}{n} + \frac{\sin(nt)}{n^2} \right) \\ &= \frac{t^2 \cdot \sin(nt)}{n} + \frac{2 \cdot t \cdot \cos(nt)}{n^2} - \frac{2 \cdot \sin(nt)}{n^3}\end{aligned}$$

$\Rightarrow$

$$\boxed{\int t^2 \cos(nt) dt = \frac{t^2 \cdot \sin(nt)}{n} + \frac{2 \cdot t \cdot \cos(nt)}{n^2} - \frac{2 \cdot \sin(nt)}{n^3}}$$

$$\begin{aligned}a_n &= \frac{1}{\pi} \cdot \left[ \frac{\pi^2 \sin(n\pi)}{n} + \frac{2\pi \cdot \cos(n\pi)}{n^2} - \frac{2 \cdot \sin(n\pi)}{n^3} - \frac{(-\pi)^2 \cdot \sin(-n\pi)}{n} - \frac{2(-\pi) \cdot \cos(-n\pi)}{n^2} + \frac{2\sin(-n\pi)}{n^3} \right] \\ &= \frac{1}{\pi} \cdot \left[ \frac{2\pi \cdot \cos(n\pi)}{n^2} + \frac{2\pi \cdot \cos(-n\pi)}{n^2} \right] = \frac{1}{\pi} \cdot \left[ \frac{2\pi \cdot \cos(n\pi)}{n^2} + \frac{2\pi \cdot \cos(n\pi)}{n^2} \right] \\ &= \frac{1}{\pi} \cdot \frac{4\pi \cdot \cos(n\pi)}{n^2} = \frac{4\cos(n\pi)}{n^2} = \frac{4 \cdot (-1)^n}{n^2}\end{aligned}$$

$\Rightarrow$

$$\boxed{a_n = \frac{4 \cdot (-1)^n}{n^2}} \quad (2)$$

$$\begin{aligned}b_n &= \frac{2}{T} \cdot \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cdot \sin\left(\frac{2n\pi t}{T}\right) dt = \frac{2}{2\pi} \cdot \int_{-\pi}^{\pi} t^2 \cdot \sin\left(\frac{2n\pi t}{2\pi}\right) dt \\ &= \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} t^2 \cdot \sin(nt) dt\end{aligned}$$

We can define  $g(t) = t^2 \cdot \sin(nt)$ . This function is odd:

$$g(-t) = (-t)^2 \cdot \sin(-nt) = -t^2 \cdot \sin(nt) = -g(t)$$

For this is clear that:

$$\boxed{b_n = 0} \quad (3)$$

Thus:

$$f(t) = \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4 \cdot (-1)^n}{n^2} \cdot \cos(nt)$$

$$f(t) = \frac{\pi^2}{3} - 4\cos(t) + \cos(2t) - \frac{4}{9} \cdot \cos(3t) + \frac{1}{4} \cdot \cos(4t) - \frac{4}{25} \cdot \cos(5t) + \dots$$

