

Languages Not Regular

Lin Chen

Email: Lin.Chen@ttu.edu

Grader: zulfi.khan@ttu.edu



TEXAS TECH
UNIVERSITY.

Non-regular languages

- Can you create a DFA/NFA that accepts the following:
 $L = \{0^n 1^n : n > 0\}$

Non-regular languages

- Can you create a DFA/NFA that accepts the following:

$$L = \{0^n 1^n : n > 0\}$$

- a DFA/NFA for $L' = \{(01)^n : n > 0\}$ is easy, but L ...
Oblivious!
- we need to keep track of the total number of 0, then check whether there is the same number of 1s

Non-regular languages


- Can you create a DFA/NFA that accepts the following:
 $L = \{0^n 1^n : n > 0\}$
 - a DFA/NFA for $L' = \{(01)^n : n > 0\}$ is easy, but L ...
 - we need to keep track of the total number of 0, then check whether there is the same number of 1s
 - The information (number) has to be stored using states
 - Finite states and infinite different numbers...

Non-regular languages

- Can you create a DFA/NFA that accepts the following:
 $L = \{0^n 1^n : n > 0\}$
 - a DFA/NFA for $L' = \{(01)^n : n > 0\}$ is easy, but L ...
 - we need to keep track of the total number of 0, then check whether there is the same number of 1s
 - The information (number) has to be stored using states
 - Finite states and infinite different numbers...
- Probably the answer is “no” , but how can we show it?


Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$




Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- The machine only has finite states, so if n is sufficiently large, there must be some loop (pigeonhole principle).




Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$



Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$
- The DFA also accepts xz , $xyyz$, $xyyyz$, ...



Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$
- The DFA also accept xz , $xyyz$, $xyyyz$, ...
- What can be the possible y ?

Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$
- The DFA also accept xz , $xyyz$, $xyyyz$, ...
- What can be the possible y ?
 - $y = 000 \dots 0$
 - $y = 111 \dots 1$
 - $y = 000 \dots 0111 \dots 1$

Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
 - For sufficiently large n , $0^n 1^n = xyz$
 - The DFA also accept xz , $xyyz$, $xyyyz$, ...
 - What can be the possible y ?
 - $y = 000 \dots 0$
 - $y = 111 \dots 1$
 - $y = 000 \dots 0111 \dots 1$
- Impossible, because then xz will have different numbers of 0 and 1

Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$
- The DFA also accept xz , $xyyz$, $xyyyz$, ...
- What can be the possible y ?
 - $y = 000 \dots 0$
 - $y = 111 \dots 1$
 - $y = 000 \dots 0111 \dots 1$

Impossible, because then $xyyz$ will have 10 as substring

Non-regular languages

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- For sufficiently large n , $0^n 1^n = xyz$
- The DFA also accept xz , $xyyz$, $xyyyz$, ...
- What can be the possible y ?
 - $y = 000 \dots 0$
 - $y = 111 \dots 1$
 - $y = 000 \dots 0111 \dots 1$
- Impossible to have such a DFA, $L = \{0^n 1^n : n > 0\}$ is non-regular

Pumping lemma

- If a language L is regular, then there exists some $n \geq 1$ such that
 - for any string $w \in L$, $|w| \geq n$, we have $w = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^iz \in L$ for any $i \geq 0$

Pumping lemma

- If a language L is regular, then there exists some $n \geq 1$ such that
 - for any string $w \in L$, $|w| \geq n$, we have $w = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^iz \in L$ for any $i \geq 0$

Repeating started once the length exceeds the total number of states

Pumping lemma

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- Exists some n , such that if $2m \geq n$, then $0^m 1^m = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^i z \in L$ for any $i \geq 0$

Pumping lemma

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- Exists some n , such that if $2m \geq n$, then $0^m 1^m = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^i z \in L$ for any $i \geq 0$
- Choose $m = n$
- $y = 00 \cdots 0$

Pumping lemma

- Suppose you have a magic DFA to accept $L = \{0^n 1^n : n > 0\}$
- Exists some n , such that if $2m \geq n$, then $0^m 1^m = xyz$ such that
 - $y \neq \epsilon$
 - $|xy| \leq n$
 - $xy^i z \in L$ for any $i \geq 0$
- Choose $m = n$
- $y = 00 \dots 0$
- The DFA should also accept xz , contradiction.

Pumping lemma

- Example: prove that $L = \{ww : w \in (a + b)^*\}$ is not regular

Pumping lemma

- Suppose $L = \{ww : w \in (a + b)^*\}$ is regular
- Exists some n , such that if $v \in L$, $|v| \geq n$, then $v = xyz$ such that
 - $y \neq e$
 - $|xy| < n$
 - $xy^iz \in L$ for any $i \geq 0$

Pumping lemma

- Suppose $L = \{ww : w \in (a + b)^*\}$ is regular
- Exists some n , such that if $v \in L$, $|v| \geq n$, then $v = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^iz \in L$ for any $i \geq 0$
- Choose $v = a^nba^n$
 - $x = a^i, y = a^k$ for some i, k
 - $xy^2z = a^{n+k}ba^n \notin L$

The way to prove- apply Pumping lemma

- You have an adversary who thinks L is regular. You need to prove that your adversary is wrong.
 - you: Language L is not regular!
 - adv: Yes it is! I have a DFA to prove it!
 - you: Oh really? How many states are in your DFA?
 - adv: n
 - you: OK, here's a string $w \in L$ with $|w| > n$. Your machine must accept w – but since $|w| > n$, there must be a loop in your computation. Where's the loop?
 - adv: Right here! (breaks w into xyz , where y is the part of the string that goes through the loop)
 - you: Ah hah! If we go through the loop 2 times instead of 1, we get a string not in L that your machine will accept!
 - adv: ...

The way to prove - apply Pumping lemma

- Your adversary picks an n
 - because you do not know this n
- You pick a $w \in L$ (such that $|w| > n$)
- Your adversary breaks w into xyz (subject to $|xy| \leq n, |y| > 0$)
 - because you do not know how exactly it is split
- You pick an i such that $xy^iz \notin L$

The way to prove- apply Pumping lemma

- Example: prove that $L = \{w: w = a^p, p \text{ is prime}\}$ is not regular

The way to prove- apply Pumping lemma

- Suppose $L = \{w: w = a^p, p \text{ is prime}\}$ is regular
- Exists some n , such that if $w \in L$, $|w| \geq n$, then $w = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^iz \in L$ for any $i \geq 0$

The way to prove- apply Pumping lemma

- Suppose $L = \{w: w = a^p, p \text{ is prime}\}$ is regular
- Exists some n , such that if $w \in L$, $|w| \geq n$, then $w = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^iz \in L$ for any $i \geq 0$
- Pick $|w| = n + 2$
 - $x = a^\alpha, y = a^\beta, z = a^\gamma$ such that
 - $\beta \geq 1, \alpha + \beta < n, \alpha + i\beta + \gamma$ is always prime for any $i \geq 0$
 - choose $i = \alpha + \gamma$, then $\alpha + i\beta + \gamma = (\alpha + \gamma)(\beta + 1)$

The way to prove- apply Pumping lemma

- What happens if we try to apply Pumping lemma to a regular language

The way to prove- apply Pumping lemma

- What happens if we try to apply Pumping lemma to a regular language
- Consider $L = \{w: w = a^p, p \text{ is even}\}$
 - Exists some n , such that if $w \in L$, $|w| \geq n$, then $w = xyz$ such that
 - $y \neq e$
 - $|xy| \leq n$
 - $xy^i z \in L$ for any $i \geq 0$
 - Choose some $w = a^{2k}$ for $2k \geq n$
 - $x = a^\alpha, y = a^\beta, z = a^\gamma$ such that
 - $\beta \geq 1, \alpha + \beta + \gamma = 2k, \alpha + i\beta + \gamma$ is always even for any $i \geq 0$

The way to prove- apply Pumping lemma

- What happens if we try to apply Pumping lemma to a regular language
- Consider $L = \{w: w = a^p, p \text{ is even}\}$
 - Exists some n , such that if $w \in L$, $|w| \geq n$, then $w = xyz$ such that
 - $y \neq e$
 - $|xy| < n$
 - $xy^iz \in L$ for any $i \geq 0$
 - Choose some $w = a^{2k}$ for $2k \geq n$
 - $x = a^\alpha, y = a^\beta, z = a^\gamma$ such that
 - $\beta \geq 1, \alpha + \beta + \gamma = 2k, \alpha + i\beta + \gamma$ is always even for any $i \geq 0$

This can be correct if β is even – we have no control on the values of α, β, γ , so no contradiction

The way to prove - apply Pumping lemma

- Not able to apply pumping lemma to show a language is non-regular **does not** necessarily mean this language is regular
 - To show it is regular, create a regular expression or DFA/NFA

The way to prove- apply Pumping lemma

- Not able to apply pumping lemma to show a language is non-regular **does not** necessarily mean this language is regular
 - To show it is regular, create a regular expression or DFA/NFA
- Not able to create a DFA/NFA for a language **does not** necessarily mean this language is non-regular
 - To show it is non-regular, apply pumping lemma suitably

Closure Properties

- Besides creating DFA/NFA or using pumping lemma, is there any other way to assert a language is regular or non-regular

Closure Properties

- Besides creating DFA/NFA or using pumping lemma, is there any other way to assert a language is regular or non-regular
- We may utilize the closure properties
 - Is L_{REG} closed under union?
 $L1 = L[r1], L2 = L[r2]$, then $L1 \cup L2 = L[(r1 + r2)]$

Closure Properties

- Besides creating DFA/NFA or using pumping lemma, is there any other way to assert a language is regular or non-regular
- We may utilize the closure properties
 - Is L_{REG} closed under union?
 $L1 = L[r1], L2 = L[r2]$, then $L1 \cup L2 = L[(r1 + r2)]$
 - Is L_{REG} closed under complementation?
Given a DFA $M = (K, \Sigma, \delta, s, F)$ for L_{REG} , we create $M' = (K, \Sigma, \delta, s, K - F)$

Closure Properties

- Besides creating DFA/NFA or using pumping lemma, is there any other way to assert a language is regular or non-regular
- We may utilize the closure properties

- Is L_{REG} closed under union?

$$L1 = L[r1], L2 = L[r2], \text{ then } L1 \cup L2 = L[(r1 + r2)]$$

- Is L_{REG} closed under complementation?

Given a DFA $M = (K, \Sigma, \delta, s, F)$ for L_{REG} , we create $M' = (K, \Sigma, \delta, s, K - F)$

- Is L_{REG} closed under intersection?

$$A \cap B = \overline{\bar{A} \cup \bar{B}}$$

Closure Properties

- Prove that $L = \{w \in \{a, b\}^* : w \text{ has an equal number of } a \text{ and } b\}$

Closure Properties

- Prove that $L = \{w \in \{a, b\}^* : w \text{ has an equal number of } a \text{ and } b\}$
 - a^*b^* is regular
 - $L \cap a^*b^* = \{a^n b^n : n \geq 0\}$

Closure Properties

- Prove that $L = \{w \in \{a, b\}^* : w \text{ has an equal number of } a \text{ and } b\}$
 - a^*b^* is regular
 - $L \cap a^*b^* = \{a^n b^n : n \geq 0\}$
 - we have shown $\{a^n b^n : n \geq 0\}$ is not regular
 - if L is regular, then by closure property $L \cap a^*b^*$ is regular
 - so L is not regular