

Sets, Relations and Languages

Lin Chen

Email: Lin.Chen@ttu.edu



TEXAS TECH
UNIVERSITY.

Ordered Pair

- (x, y) is an ordered pair
- x, y are called components (of the ordered pair)

Ordered Pair

- (x, y) is an ordered pair
- x, y are called components (of the ordered pair)
- Order matters
 - $(x, y) \neq (y, x)$ if $x \neq y$
 - $\{x, y\} = \{y, x\}$
- Components can be identical
 - (x, x) is a valid ordered pair
 - $(x, x) \neq x$

Cartesian Product

- $A \times B = \{(x, y) : x \in A \text{ and } y \in B\}$
- $\{a, b\} \times \{c, d\} = \{(a, c), (a, d), (b, c), (b, d)\}$
- $2^{\{a\} \times \{b\}} = ?$

Cartesian Product

- $A \times B = \{(x, y) : x \in A \text{ and } y \in B\}$
- $\{a, b\} \times \{c, d\} = \{(a, c), (a, d), (b, c), (b, d)\}$
- $2^{\{a\} \times \{b\}} = 2^{\{(a, b)\}} = \{\emptyset, (a, b)\}$
- $2^{\{a\}} \times 2^{\{b\}} = ?$

Cartesian Product

- $A \times B = \{(x, y) : x \in A \text{ and } y \in B\}$
- $\{a, b\} \times \{c, d\} = \{(a, c), (a, d), (b, c), (b, d)\}$
- $2^{\{a\} \times \{b\}} = 2^{\{(a, b)\}} = \{\emptyset, \{(a, b)\}\}$
- $2^{\{a\}} \times 2^{\{b\}} = \{\{a\}, \emptyset\} \times \{\{b\}, \emptyset\} = \{(\{a\}, \{b\}), (\{a\}, \emptyset), (\emptyset, \{b\}), (\emptyset, \emptyset)\}$

Ordered Tuple

- (a_1, a_2, \dots, a_n) is an ordered (n -)tuple
- Each a_i is called the (i -th) component (of the ordered tuple)
- Ordered 2-tuple is ordered pair

Ordered Tuple

- (a_1, a_2, \dots, a_n) is an ordered $(n-)$ tuple
- Each a_i is called the $(i\text{-th})$ component (of the ordered tuple)
- Ordered 2-tuple is ordered pair
- $A_1 \times A_2 \times A_3 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for all } i\}$
 - $\{a\} \times \{b, c\} \times \{d\} = \{(a, b, d), (a, c, d)\}$

Ordered Tuple

- (a_1, a_2, \dots, a_n) is an ordered $(n-)$ tuple
- Each a_i is called the $(i\text{-th})$ component (of the ordered tuple)
- Ordered 2-tuple is ordered pair
- $A_1 \times A_2 \times A_3 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for all } i\}$
 - $\{a\} \times \{b, c\} \times \{d\} = \{(a, b, d), (a, c, d)\}$
 - The definition implies that $((a, b), c) = (a, b, c)$. It is true since we use $()$ to indicate that objects are ordered. Both $((a, b), c)$ and (a, b, c) denote the 3 objects are placed in the order of a, b, c .

Relations

- A relation R is a set of ordered pairs
 - Attendance sheet produces a relation: $\{(Alice, X), (Bob, -), \dots\}$
 - $<$ over natural numbers is also a relation
 - $\{ (0,1), (0,2), (0,3), \dots$
 - $(1,2), (1,3), (1,4), \dots$
 - $(2,3), (2,4), (2,5), \dots$
 - $\dots \}$

Functions

- A function is a special kind of relation (all functions are relations, but not all relations are functions)
- A relation $R \subseteq A \times B$ is a function if:
 - For each $a \in A$, there is exactly one ordered pair in R with the first component a

Functions

- A function is a special kind of relation (all functions are relations, but not all relations are functions)
- A relation $R \subseteq A \times B$ is a function if:
 - For each $a \in A$, there is exactly one ordered pair in R with the first component a
- Two relations (C : set of cities, S : set of states)
 - $R_1 = \{(x, y): x \in C, y \in S, x \text{ is a city in state } y\}$
 - $R_2 = \{(x, y): x \in S, y \in C, y \text{ is a city in state } x\}$

Who is a function?

Functions

- A function is a special kind of relation (all functions are relations, but not all relations are functions)
- A relation $R \subseteq A \times B$ is a function if:
 - For each $a \in A$, there is exactly one ordered pair in R with the first component a
- Two relations (C : set of cities, S : set of states)
 - $R_1 = \{(x, y): x \in C, y \in S, x \text{ is a city in state } y\}$
 - $R_2 = \{(x, y): x \in S, y \in C, y \text{ is a city in state } x\}$

Who is a function?

Functions

- A function f that is a subset of $A \times B$ is written as:
 - $f : A \mapsto B$
- $(a, b) \in f$ is written $f(a) = b$
- A is the domain of the function
- if $A' \subseteq A$, $f(A') = \{b : f(a) = b \text{ for some } a \in A'\}$ is the image of A'
- The range of a function is the image of its domain

Functions

- A function $f : A \mapsto B$ is:
 - one-to-one, if no two elements in A match to the same element in B
 - onto, if each element in B is mapped to by at least one element in A
 - bijection, if it is both one-to-one and onto

Inverse


- The inverse of a binary relation $R \subseteq A \times B$ is $\{(b, a): (a, b) \in R\}$, and denoted as R^{-1} .

Inverse

- The inverse of a binary relation $R \subseteq A \times B$ is $\{(b, a): (a, b) \in R\}$, and denoted as R^{-1} .
 - $R_1 = \{(x, y): x \in C, y \in S, x \text{ is a city in state } y\}$
 - $R_2 = \{(y, x): x \in C, y \in S, x \text{ is a city in state } y\}$
- The inverse of a relation, which is not a function, can be a function
- The inverse of a function may fail to be a function
- If a function is bijection, then its inverse is also a function

Relation Graph


- Representing a relation $R \subseteq A \times A$ using a **directed graph**
 - A directed graph (or digraph) is a graph that is made up of a set of vertices (nodes) connected by edges, where the edges have a direction associated with them



Relation Graph

- Representing a relation $R \subseteq A \times A$ using a **directed graph**
 - A directed graph (or digraph) is a graph that is made up of a set of vertices (nodes) connected by edges, where the edges have a direction associated with them
 - Use nodes to represent elements
 - Use directed edge to represent an ordered pair


$$R = \{(a, b), (b, c), (c, b), (c, a)\}$$



Relation Graph



- Representing a relation $R \subseteq A \times A$ using a **directed graph**
 - A directed graph (or digraph) is a graph that is made up of a set of vertices (nodes) connected by edges, where the edges have a direction associated with them
 - Use nodes to represent elements
 - Use directed edge to represent an ordered pair

$$R = \{(a, b), (b, c), (c, b), (c, a), (c, c)\}$$





Relation Types

- A relation $R \subseteq A \times A$ is **reflexive** if
 - $(a, a) \in R$ for every $a \in A$
 - Equivalently, every node has a self-loop





Relation Types

- A relation $R \subseteq A \times A$ is **symmetric** if
 - $(a, b) \in R$ if $(b, a) \in R$
 - Equivalently, every edge between two nodes is “two-ways”





Relation Types

- A relation $R \subseteq A \times A$ is **antisymmetric** if
 - If $(a, b) \in R$ and $a \neq b$, then $(b, a) \notin R$
 - Equivalently, every edge between two nodes is “one-way”




Relation Types

- A relation $R \subseteq A \times A$ is **transitive** if
 - If $(a, b) \in R$ and $(b, c) \in R$, then $(a, c) \in R$
 - Equivalently, you can always “shortcut”




Relation Types

- A relation $R \subseteq A \times A$ is an **equivalence relation** if
 - If it is reflexive, symmetric and transitive




Relation Types

- A relation $R \subseteq A \times A$ is an **equivalence relation** if
 - If it is reflexive, symmetric and transitive
- Example: A (All English words), $(w_1, w_2) \in R$ if they start with same letter



Relation Types


- A relation $R \subseteq A \times A$ is an **equivalence relation** if
 - If it is reflexive, symmetric and transitive
- Each connected component form an equivalence class



Relation Types

- A relation $R \subseteq A \times A$ is an **equivalence relation** if
 - If it is reflexive, symmetric and transitive
- Each connected component form an equivalence class

We can pick any element of an equivalent class to represent this class and denote it as $[a]$



Relation Types

- A relation $R \subseteq A \times A$ is a **partial order** if
 - If it is reflexive, antisymmetric and transitive
- Example:
 1. allow a person to be considered as an ancestor of himself/herself, then
$$R = \{(a, b): a, b \text{ are persons and } a \text{ is an ancestor of } b\}$$
 2. \leq defined on natural numbers

Relation Types

- A relation $R \subseteq A \times A$ is a **total order** if
 - If it is a partial order, and
 - For all $a, b \in A$, either $(a, b) \in R$ or $(b, a) \in R$
- Example:
 1. Ancestor relationship is not a total order
$$R = \{(a, b): a, b \text{ are persons and } a \text{ is an ancestor of } b\}$$
 2. \leq defined on natural numbers is a total order

Closure

- A set $A \subseteq B$ is closed under a relation $R \subseteq ((B \times B) \times B)$ if:
 - $a_1, a_2 \in A$ and $((a_1, a_2), c) \in R \Rightarrow c \in A$
 - That is, if a_1, a_2 are both in A , and $((a_1, a_2), c)$ is in the relation, then c is also in A
- N is closed under addition
- N is not closed under subtraction or division

Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal

Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal
- What if I don't do the calculation?
 - Pick one apple from one basket, then pick one from another
 - If both of them become empty at the same step \rightarrow same cardinality



Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal
- What if I don't do the calculation?
 - Pick one apple from one basket, then pick one from another
 - Create a bijection $f : A \mapsto B$

Why it has to be a bijection?

Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal
- What if I don't do the calculation?
 - Pick one apple from one basket, then pick one from another
 - Create a bijection $f : A \mapsto B$

Why it has to be a bijection?

- Bijection is one-one: if I pick one apple from A , I must also pick one from B , thus no two apples in A are mapped to the same apple in B

Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal
 - What if I don't do the calculation?
 - Pick one apple from one basket, then pick one from another
 - Create a bijection $f : A \mapsto B$
- Why it has to be a bijection?
- Bijection is onto: Eventually, the basket B should be empty.

Cardinality

- How can we tell if two sets A and B have the same cardinality?
 - Calculate $|A|$ and $|B|$, see if the two numbers equal
- What if I don't do the calculation?
 - Pick one apple from one basket, then pick one from another
 - Create a bijection $f : A \mapsto B$
- If there is a bijection, we say A and B are **equinumerous**

Cardinality

- A set is **finite** if it is equinumerous with $\{1, 2, \dots, n\}$ for some n
- Otherwise, the set is **infinite**

Countable sets

- A set is **finite** if it is equinumerous with $\{1, 2, \dots, n\}$ for some n
- Otherwise, the set is **infinite**
- Not all infinite sets are equinumerous
 - A set is **countably infinite** if it is equinumerous with \mathbb{N}
 - A set is **countable** if it is finite or countably infinite
 - Otherwise, it is **uncountable**

Countable sets

- A set is **countable** if it is equinumerous with \mathbb{N}
 - Even numbers of \mathbb{N}
$$f(x) = 2x$$

Countable sets


- A set is **countable** if it is equinumerous with \mathbb{N}

- Even numbers of \mathbb{N}

$$f(x) = 2x$$


- All integers

$$f(x) = \left\lfloor \frac{x}{2} \right\rfloor * (-1)^x$$



Countable sets


- A set is **countable** if it is equinumerous with \mathbb{N}
 - Union of two disjoint countable sets



$$f(x) = \begin{cases} a_{\frac{x}{2}} & \text{if } x \text{ is even} \\ b_{\frac{x-1}{2}} & \text{if } x \text{ is odd} \end{cases}$$

Countable sets

- A set is **countable** if it is equinumerous with N
 - $N \times N$



$$f(x, y) = \frac{(x + y)(x + y + 1)}{2} + x$$

Uncountable sets

- A set is **countable** if it is equinumerous with N
 - There exist uncountable sets, e.g., the real number within $(0,1)$
 - How can we show it is uncountable?

Uncountable sets

- A set is **countable** if it is equinumerous with N
 - There exist uncountable sets, e.g., the real number within $(0,1)$
 - How can we show it is uncountable?
 - Proof by contradiction

Suppose it is countable, then there exists some bijection, but given the bijection, we can find some real number which is not mapped to any element in N

Uncountable sets

- A set is **countable** if it is equinumerous with N
 - There exist uncountable sets, e.g., the real number within $(0,1)$

Suppose there is a bijection:

0	0.1 3 5 7 2 8 0 0 0 6 7 ...
1	0.2 3 7 5 0 6 2 1 1 4 5 ...
2	0.9 4 4 8 2 3 1 8 9 0 2 ...
3	0.6 2 2 6 3 3 7 7 9 3 9 ...
...	...

Uncountable sets

- A set is **countable** if it is equinumerous with N
 - There exist uncountable sets, e.g., the real number within $(0,1)$

Suppose there is a bijection:

0	0. 1 3 5 7 2 8 0 0 0 6 7 ...
1	0.2 3 7 5 0 6 2 1 1 4 5 ...
2	0.9 4 4 8 2 3 1 8 9 0 2 ...
3	0.6 2 2 6 3 3 7 7 9 3 9 ...
...	...

Consider 0. 2457... Does this number appear in the list?

Proof Techniques

- Three basic proof techniques used in this class
 - Induction
 - Diagonalization
 - Pigeonhole Principle

Induction

- Claim: Can create exact postage for any amount $\geq \$0.08$ using only 3 cent and 5 cent stamps

Induction

- Claim: Can create exact postage for any amount $\geq \$0.08$ using only 3 cent and 5 cent stamps
 - Base case:
Can create postage for 0.08 using one 5-cent and one 3-cent stamp

Induction

- Claim: Can create exact postage for any amount $\geq \$0.08$ using only 3 cent and 5 cent stamps

- Inductive case

To show: if we can create exact postage for $\$x$ using only 3-cent and 5-cent stamps, we can create exact postage for $\$x + \0.01 using 3-cent and 5-cent stamps

- Two cases:

1. Exact postage for $\$x$ uses at least one 5-cent stamp
2. Exact postage for $\$x$ uses no 5-cent stamps

Induction

- Claim: Can create exact postage for any amount $\geq \$0.08$ using only 3 cent and 5 cent stamps

-Inductive case

To show: if we can create exact postage for $\$x$ using only 3-cent and 5-cent stamps, we can create exact postage for $\$x + \0.01 using 3-cent and 5-cent stamps

- Two cases:

1. Exact postage for $\$x$ uses at least one 5-cent stamp
2. Exact postage for $\$x$ uses no 5-cent stamps

Replace one 5-cent with two 3-cent stamps

Induction

- Claim: Can create exact postage for any amount $\geq \$0.08$ using only 3 cent and 5 cent stamps

-Inductive case

To show: if we can create exact postage for $\$x$ using only 3-cent and 5-cent stamps, we can create exact postage for $\$x + \0.01 using 3-cent and 5-cent stamps

- Two cases:

1. Exact postage for $\$x$ uses at least one 5-cent stamp
2. Exact postage for $\$x$ uses no 5-cent stamps

Replace three 3-cent with two 5-cent stamps

Induction

What is wrong with the following purported proof that all horses are the same color?

Proof by induction on the number of horses:

Basis Step. There is only one horse. Then clearly all horses have the same color.

Induction Hypothesis. In any group of up to n horses, all horses have the same color.

Induction Step. Consider a group of $n + 1$ horses. Discard one horse; by the induction hypothesis, all the remaining horses have the same color. Now put that horse back and discard another; again all the remaining horses have the same color. So all the horses have the same color as the ones that were not discarded either time, and so they all have the same color.

Pigeonhole Principle

- A, B are finite sets, with $|A| > |B|$, then there is no one-to-one function from A to B
- If you have n pigeonholes, and $> n$ pigeons, and every pigeon is in a pigeonhole, there must be at least one hole with > 1 pigeon.

Pigeonhole Principle

- Claim: Let $R \subseteq A \times A$ be a relation. If there exists a path in the directed graph representation of R , then there exists a path of length at most $|A|$.

Pigeonhole Principle

- Claim: Let $R \subseteq A \times A$ be a relation. If there exists a path in the directed graph representation of R , then there exists a path of length at most $|A|$.
 - Suppose on the contrary the shortest path has length $\geq |A| + 1$
 - In this path (a_1, a_2, \dots, a_n) there must be some node appears at least twice (pigeonhole principle)

Pigeonhole Principle

- Claim: Let $R \subseteq A \times A$ be a relation. If there exists a path in the directed graph representation of R , then there exists a path of length at most $|A|$.
 - Suppose on the contrary the shortest path has length $\geq |A| + 1$
 - In this path (a_1, a_2, \dots, a_n) there must be some node appears at least twice (pigeonhole principle)
 - $(a_1, a_2, \dots, a_i, a_{i+1}, \dots, a_j = a_i, a_{j+1}, \dots, a_n)$
 - $(a_1, a_2, \dots, a_i, a_{j+1}, \dots, a_n)$ is shorter

Strings

- An alphabet Σ is a finite set of symbols
 - $\Sigma_1 = \{0,1\}$
 - $\Sigma_2 = \{a, b, c, \dots, z\}$
- A string is a finite sequence of symbols from an alphabet
 - apple, banana are both strings over $\Sigma_2 = \{a, b, c, \dots, z\}$
 - 100110 is a string over $\Sigma_1 = \{0,1\}$

Strings

- An alphabet Σ is a finite set of symbols
 - $\Sigma_1 = \{0,1\}$
 - $\Sigma_2 = \{a, b, c, \dots, z\}$
- A string is a finite sequence of symbols from an alphabet
 - apple, banana are both strings over $\Sigma_2 = \{a, b, c, \dots, z\}$
 - 100110 is a string over $\Sigma_1 = \{0,1\}$
- Length of a string = number of symbols in the string
 - $|apple|=5$, $|100110|=6$

Strings

- Empty string: e
 - $|e| = 0$

Strings

- For string w , we use $w(j)$ to denote the symbol on j -th position
 - $w = \text{apple}$, $w(2) = w(3) = \text{p}$
 - $\Sigma_2 = \{a, b, c, \dots, z\}$

Strings

- For string w , we use $w(j)$ to denote the symbol on j -th position
 - $w = \text{apple}$, $w(2) = w(3) = \text{p}$
 - $\Sigma_2 = \{a, b, c, \dots, z\}$

Strings

- String operation: ◦ Contatenation
 - apple ◦ banana = applebanana
 - 110 ◦ 101 = 110101
- We may also drop ◦ if it is clear from context
 - $w_1 \circ w_2 = w_1 w_2$ for two strings w_1, w_2

Strings

- String operation: ◦ Contatenation
 - apple ◦ banana = applebanana
 - 110 ◦ 101 = 110101
- We may also drop ◦ if it is clear from context
 - $w_1 \circ w_2 = w_1 w_2$ for two strings w_1, w_2
- $w \circ e = we = ew = w$

Strings

- Substring:
 - v is a substring of w if and only if $w = xvy$ for some strings x, y
- A string is a substring of itself (by taking $x = y = e$)
- e is a substring of any string

Strings

- If $w = xv$, v is a **suffix** of w
- If $w = xv$, x is a **prefix** of w

Strings

- A string can be concatenated with itself
 - $w \circ w = ww = w^2$
 - $w^{i+1} = w^i w$
 - $w^0 = e$
- Example: $\text{apple}^2 = \text{appleapple}$

Strings

- Reversal: writing backwards
 - $w = \text{apple}$, $w^R = \text{elppa}$
 - $e^R = e$

Formal language

- The set of all strings over alphabet Σ is denoted as Σ^*
 - $e \in \Sigma^*$
- Any subset of Σ^* is called a language
 - English is a subset of $\{a, b, \dots, z\}^*$

Language operations

- Concatenation
 - $L_1, L_2 \subseteq \Sigma^*$, $L_1 \circ L_2 = L_1 L_2 = \{w \in \Sigma^* : w = xy, x \in L_1, y \in L_2\}$
 - Example: $\{a, ab\}\{b, bb\} = \{abb, ab, abbb\}$

Language operations

- Concatenation
 - $L_1, L_2 \subseteq \Sigma^*$, $L_1 \circ L_2 = L_1 L_2 = \{w \in \Sigma^* : w = xy, x \in L_1, y \in L_2\}$
 - Example: $\{a, ab\}\{b, bb\} = \{abb, ab, abbb\}$
- A language can be concatenated with itself
 - $L^2 = LL = \{w^2 : w \in L\}$
 - $L^0 = \{e\}$

Language operations

- Complement
 - For $L \in \Sigma^*$, $\bar{L} = \Sigma^* - L$

Language operations

- Complement
 - For $L \in \Sigma^*$, $\bar{L} = \Sigma^* - L$
- Kleene star
 - L^* : the set of all strings obtained by concatenating zero or more strings of L
 - $L^* = \{w \in \Sigma^* : w = w_1 w_2 \cdots w_k \text{ for some } k \geq 0, w_i \in L, 1 \leq i \leq k\}$
 - $\Sigma^* = \{w \in \Sigma : w = w_1 w_2 \cdots w_k \text{ for some } k \geq 0, w_i \in \Sigma, 1 \leq i \leq k\}$ = all the strings over alphabet Σ