

Intuitionist Implicational logic (IIL)

Definition:-

Implicational formulae are built from propositional variables using the implicational connective " \rightarrow ": if σ and τ are formulas, then so is $(\sigma \rightarrow \tau)$. An infinite sequence of propositional variables (denoted a, b, c, \dots) is assumed to be given. There are no propositional constants and no other connectives than " \rightarrow ". " \rightarrow " is right-associative.

Defn: IIL has the foll. rules:-

$$(\rightarrow E) : \frac{\sigma \rightarrow \tau \quad \sigma}{\tau} \quad (\rightarrow I) : \frac{\begin{array}{c} [\sigma] \\ \vdots \\ \tau \end{array}}{\sigma \rightarrow \tau}$$

Each application of $(\rightarrow I)$ is said to discharge (cancel) some, all, or none of the occurrences of σ above τ , and must be accompanied by a discharge label that lists all the occurrences of σ that it discharges. If none are discharged then the appⁿ of $\rightarrow I$ is vacuous.

Example:

1. A proof of $(a \rightarrow a \rightarrow c) \rightarrow a \rightarrow c$

$$\begin{array}{c} \frac{\frac{\frac{[a \rightarrow a \rightarrow c] \quad (0011) \quad [a] \quad (0012)}{a \rightarrow c \quad (001)} \quad (\rightarrow E)}{c \quad (00)} \quad (\rightarrow I) \quad \{ \text{discharges 'a' at } 0012, 002 \}} \\ \frac{a \rightarrow c \quad (0)}{(a \rightarrow a \rightarrow c) \rightarrow a \rightarrow c} \quad (\rightarrow I) \quad \{ \text{discharges 'a' at } 0011 \} \end{array}$$

2. A proof of $(a \rightarrow a \rightarrow c) \rightarrow a \rightarrow a \rightarrow c$

$$\begin{array}{c} \frac{\frac{\frac{[a \rightarrow a \rightarrow c] \quad (00011) \quad [a] \quad (00012)}{a \rightarrow c \quad (0001)} \quad (\rightarrow E)}{c \quad (000)} \quad (\rightarrow I) \quad \{ \text{discharges 'a' at } 00012 \}} \\ \frac{a \rightarrow c \quad (00)}{a \rightarrow a \rightarrow c} \quad (\rightarrow I) \quad \{ \text{discharges 'a' at } 0002 \}} \\ \frac{(a \rightarrow a \rightarrow c) \rightarrow a \rightarrow a \rightarrow c}{} \quad (\rightarrow I) \quad \{ \text{discharges 'a' at } 00011 \} \end{array}$$

The Curry-Howard isomorphism :-

Defn : (C-H mapping from lambda to logic)

If Δ is a TA_{λ} -deduction of $\Gamma \vdash M : \tau$,
the corresponding logic deduction Δ_L is defined thus.

(i) $M \equiv x$ and $\Delta : x : \tau \vdash x : \tau$

Δ_L : just τ

(ii) $M \equiv P Q$ and $\Gamma = \Gamma_1 \cup \Gamma_2$ and last step in Δ
has the form

$$\frac{\begin{array}{c} (\Delta_1) \\ \Gamma_1 \vdash P : \sigma \rightarrow \tau \end{array} \quad \begin{array}{c} (\Delta_2) \\ \Gamma_2 \vdash Q : \sigma \end{array} \quad (\rightarrow E)}{\Gamma_1 \cup \Gamma_2 \vdash (PQ) : \tau}$$

let Δ_{1L} correspond to Δ_1
 Δ_{2L} " " Δ_2

Δ_L " " to Δ

Δ_L is obtained by applying $(\rightarrow E)$ of IIL to Δ_{1L}, Δ_{2L} .

(iii) $M \equiv \lambda x. P$, $\tau \equiv \rho \rightarrow \sigma$, $\Gamma = \Gamma' - x$ and the
last step in Δ is

$$\frac{\begin{array}{c} \Delta' \\ \Gamma' \vdash P : \sigma \end{array} \quad (\rightarrow I)}{\Gamma' - x \vdash (\lambda x. P) : \rho \rightarrow \sigma}$$

Δ_L is obtained from Δ'_L by discharging all occurrences
of ρ in Δ'_L whose positions are the same as the
positions of the free occurrences of x in P .

Example'

$$\begin{array}{lcl}
 \Delta & x : a \rightarrow a \rightarrow c \vdash x : a \rightarrow a \rightarrow c & z : a \vdash z : a & 1 \\
 \rightarrow E & \frac{x : a \rightarrow a \rightarrow c, z : a \vdash (xz) : a \rightarrow c}{x : a \rightarrow a \rightarrow c, z : a \vdash (xz)y : c} & y : a \vdash y : a & 2 \\
 (\rightarrow E) & \frac{x : a \rightarrow a \rightarrow c, z : a, y : a \vdash (xz)y : c}{x : a \rightarrow a \rightarrow c, y : a \vdash \lambda z. (xz)y : a \rightarrow c} & & 3 \\
 (\rightarrow I) & \frac{x : a \rightarrow a \rightarrow c, y : a \vdash \lambda z. (xz)y : a \rightarrow c}{x : a \rightarrow a \rightarrow c \vdash \lambda y. \lambda z. (xz)y : a \rightarrow a \rightarrow c} & \text{durch } z^4 & 4 \\
 (\rightarrow I) & \frac{x : a \rightarrow a \rightarrow c \vdash \lambda y. \lambda z. (xz)y : a \rightarrow a \rightarrow c}{x : a \rightarrow a \rightarrow c \vdash \lambda x. \lambda y. \lambda z. (xz)y : (a \rightarrow a \rightarrow c) \rightarrow a \rightarrow a \rightarrow c} & \text{durch } y^5 & 5 \\
 (\rightarrow I) & \vdash \lambda x. \lambda y. \lambda z. (xz)y : (a \rightarrow a \rightarrow c) \rightarrow a \rightarrow a \rightarrow c & \text{durch } x^6 & 6
 \end{array}$$

$$\begin{array}{lcl}
 \Delta_L & \frac{a \rightarrow a \rightarrow c \ [\#] \quad a \ [\#]}{a \rightarrow c \quad a \ [\#*]} & \frac{1}{2} \\
 & \frac{a \rightarrow c \quad a \ [\#*]}{c} & \text{durch } a \text{ at } [\#] \quad 3 \\
 & \frac{c}{a \rightarrow c} & \text{durch } a \text{ at } [\#*] \quad 4 \\
 & \frac{a \rightarrow c}{a \rightarrow a \rightarrow c} & \frac{5}{6} \\
 & \frac{a \rightarrow a \rightarrow c}{(a \rightarrow a \rightarrow c) \rightarrow a \rightarrow a \rightarrow c} & \text{durch } a \text{ at } [\#] \quad 6
 \end{array}$$