

# Solution to Homework 2

## Bayesian Methods

Anna Pavlova

### Problem 1

To find a minimum of the error-function

$$E = \frac{1}{2} \sum_n r_n (t_n - \boldsymbol{\theta}^T \mathbf{x}_n)^2$$

(where  $r_n > 0, \forall n$ ), set the gradient to zero

$$\nabla_{\boldsymbol{\theta}} E = \sum_n r_n (\boldsymbol{\theta}^T \mathbf{x}_n - t_n) \mathbf{x}_n = 0$$

now since  $\mathbf{a}^T \mathbf{b} = \mathbf{b}^T \mathbf{a}$ , define a matrix  $(\mathbf{X})_{ni} = (\mathbf{x}_n^T)_i$  and  $\mathbf{R} = \text{diag}(r_1, \dots, r_N)$  have

$$\mathbf{X}^T \mathbf{R} \mathbf{X} \boldsymbol{\theta} - \mathbf{R} \mathbf{X}^T \mathbf{t} = 0$$

so the argmin is

$$\boldsymbol{\theta}^* = (\mathbf{X}^T \mathbf{R} \mathbf{X})^{-1} \mathbf{R} \mathbf{X}^T \mathbf{t}.$$

This is truly a minimum, since all  $r_n > 0$  and the functional is thus positive-definite (this is actually some weighted least-squares method). This weighting can be interpreted in two ways (i): different data points  $t_n$  may be affected by different level of noise (thus having different dispersions) and (ii): if some data points  $t_n$  appear several times, this can be simply tracked by considering one point weighted appropriately.

### Problem 2

Need to proof inequality

$$\sigma_{N+1}^2(x) \leq \sigma_N^2(x)$$

We have variance  $\sigma_N^2(x) = \frac{1}{\beta} + \phi(x)^T S_N \phi(x)$  and  $S_N^{-1} = \alpha^{-1} I + \beta \Phi^T \Phi$  for calculation  $\sigma_{N+1}^2$  found  $S_{N+1}^{-1}$  by Woodbury matrix identity

$$\begin{aligned} \sigma_{N+1}^2(x) : \\ \sigma_{N+1}^2(x) &= \frac{1}{\beta} + \phi(x)^T S_{N+1} \phi(x) \\ S_{N+1}^{-1} &= \alpha^{-1} I + \beta \Phi_{N+1}^T \Phi_{N+1} \end{aligned}$$

$\Phi_{N+1}^T \Phi_{N+1} = \Phi_N^T \Phi_N + \phi_{N+1} \phi_{N+1}^T$  accordingly

$$S_{N+1}^{-1} = \alpha^{-1} I + \beta(\Phi_N^T \Phi_N + \phi_{N+1} \phi_{N+1}^T) = S_N^{-1} + \beta \phi_{N+1} \phi_{N+1}^T$$

found  $S_{N+1}$  using Woodbury matrix identity

$$(S_N^{-1} + \beta \phi_{N+1} \phi_{N+1}^T)^{-1} = S_N - \beta \frac{(S_N \phi_{N+1})(\phi_{N+1}^T S_N)}{1 + \beta \phi_{N+1}^T S_N \phi_{N+1}}$$

Let us prove that denominator  $\geq 0$ , for that we have to note that  $\phi^T S_N^{-1} \phi = \frac{\|\phi\|^2}{\alpha} + \beta \|\Phi_N \phi\|^2 \geq 0$ .  $S_N$  is hermitian matrix, so  $\phi^T (S_N \phi_{N+1})(\phi_{N+1}^T S_N) \phi = (\phi_{N+1}^T S_N \phi)^2 \geq 0$

$$\sigma_{N+1}(x) = \sigma_N(x) - \beta \frac{(\phi_{N+1}^T S_N \phi)^2}{1 + \beta \phi_{N+1}^T S_N \phi_{N+1}} \leq \sigma_N(x)$$