Class 11: alphafold pt 1

Ayanna Kerr (PID:A17143404)

Section 1: Proportion of G|G in a population

```
mxl <- read.csv("373531-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")</pre>
  head(mxl)
  Sample..Male.Female.Unknown. Genotype..forward.strand. Population.s. Father
1
                   NA19648 (F)
                                                      A|A ALL, AMR, MXL
                                                      G|G ALL, AMR, MXL
2
                   NA19649 (M)
3
                   NA19651 (F)
                                                      A|A ALL, AMR, MXL
4
                   NA19652 (M)
                                                      G|G ALL, AMR, MXL
5
                                                      G|G ALL, AMR, MXL
                   NA19654 (F)
                   NA19655 (M)
                                                      A|G ALL, AMR, MXL
 Mother
1
2
3
4
  table(mxl$Genotype..forward.strand.)
A|A A|G G|A G|G
22 21 12
  table(mxl$Genotype..forward.strand.)/nrow(mxl) *100
```

```
A|A A|G G|A G|G
34.3750 32.8125 18.7500 14.0625
```

Now let's look at a different population. I picked the GBR.

```
gbr <- read.csv("373522-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")
head(gbr)</pre>
```

```
Sample..Male.Female.Unknown. Genotype..forward.strand. Population.s. Father
1
                   HG00096 (M)
                                                       A|A ALL, EUR, GBR
2
                   HG00097 (F)
                                                       G|A ALL, EUR, GBR
3
                   HG00099 (F)
                                                       G|G ALL, EUR, GBR
4
                                                       A|A ALL, EUR, GBR
                   HG00100 (F)
5
                   HG00101 (M)
                                                       A|A ALL, EUR, GBR
6
                   HG00102 (F)
                                                       A|A ALL, EUR, GBR
 Mother
2
3
4
5
```

Find the proportion of G|G

```
round(table(gbr$Genotype..forward.strand.)/nrow(gbr) *100, 2)
```

```
A|A A|G G|A G|G
25.27 18.68 26.37 29.67
```

This variant that is associated with childhood asthma is more frequent in the GBR population than the MXL population.