




DEEP CERVICAL FASCIA



Introduction

Deep cervical fascia of the neck is condensed to form the following layers:

- **Investing layer**
- **Pretracheal layer**
- **Prevertebral layer**
- **Carotid sheath**




1. Investing layer:

- It lies deep to platysma and wrapped around the neck like a collar.

Attachments:

- **Superiorly:** external occipital protuberance, superior nuchal line, mastoid process and base of mandible
- **Inferiorly:** spine of scapula, acromion process, clavicle and manubrium
- **Posteriorly:** ligamentum nuchae and spine of seventh cervical vertebra
- **Anteriorly:** symphysis menti and hyoid bone



Modifications of the investing layer:


ENCLOSES 2 SPACES-

1.SUPRACLAVICULAR SPACE

- splitting of investing layer medial 3rd of clavicle & gets attached to anterior & posterior border of upper surface of clavicle

Content of the space

Supraclavicular nerves,
external jugular vein





2.SUPRASTERNAL SPACE

- Splitting of investing layer at the upper border of sternum which gets attached to anterior & posterior border of supra-sternal notch
- known as space of burns


Content of the space

- Inter-clavicular ligament, sternal head of sternocleidomastoid muscle, jugular venous arch, loose areolar tissue



Stylomandibular ligament:

It is formed by the thickening of deep cervical fascia




2. Pretracheal layer:

- It encloses and suspends thyroid gland and forms its false capsule.

Attachments:

- **Superiorly:** hyoid bone, oblique line on thyroid cartilage and cricoid cartilage
- **Inferiorly:** passes in front of trachea and blends with arch of aorta
- **Laterally:** fuses with carotid sheath
- **Posteriorly:** continuous with buccopharyngeal fascia of pharynx



Vertical extent





Fig. 4.13. Vertical disposition of pretracheal fascia.

Modification of pretracheal layer:

Suspensory ligament for thyroid/ligament of Berry:

- Posterior layer of thyroid capsule is thick and it forms suspensory ligament which is mainly attached to cricoid cartilage.
- **Significance:** it supports the thyroid gland and prevent it from sinking into the mediastinum.





3. Prevertebral layer:


- It lies in front of prevertebral muscles and forms floor of posterior triangle of the neck.

Attachments:

- **Superiorly:** base of the skull
- **Inferiorly:** extends into superior mediastinum and attached to the anterior longitudinal ligament and body of fourth thoracic vertebra
- **Anteriorly:** separated from buccopharyngeal fascia and pharynx by retropharyngeal space
- **Laterally:** lost deep to trapezius




Prevertebral muscles:



© www.kenhub.com

KEN
HUB



© www.kenhub.com


KEN
HUB

Carotid sheath:


- It is a condensation of fibroareolar tissue around the main vessels of the neck.

Contents:

- I. Common carotid arteries
- II. Internal carotid arteries
- III. Internal jugular vein(s)
- IV. Vagus nerve(s)




© www.kenhub.com
KEN HUB





Figs 3.8a and b: Right carotid sheath with its contents: (a) Surface view, and (b) sectional view

Buccopharyngeal fascia:

- Covers superior constrictor muscle of pharynx and extends upto superficial aspect of buccinator muscle.





**THANK
YOU**

