


Trained profile 5,172 (12.7%)


Ambiguous family 2,905 (7.1%)

Discovery layer only 32,613 (80.1%)

a Support strength of the 40,690 candidates



c Reference anchors



Ledger rows (bars, log-spaced) · positives · families reaching ≥3.
All 30 ledger positives trace to one DOI (Knoll 2009); the amendment mapped them to 24 primary papers — only hfam_52/hfam_70 reach ≥3.

Architecture-evidence coverage (% of each superfamily's rows)

b

	GxSxG lipase box	Catalytic domain type	PF06850 (Cys marker)	lid loop (MCL)	AHSMG (PhaZ7)	Secretion SignalP6
Intracellular Cys-type nPHASCL (n = 29,974)	0	0	100	0	0	0
Extracellular dPHASCL type 1 (n = 5,018)	100	100	0	0	0	100
Extracellular dPHASCL type 2 (n = 4,348)	100	100	0	0	0	100
Intracellular nPHAMCL (n = 987)	100	0	0	100	0	0
Extracellular dPHAMCL (n = 280)	100	0	0	0	0	100
Extracellular PhaZ7-like (n = 82)	45	0	0	0	100	100
Periplasmic periplasmic (n = 1)	100	100	0	0	0	100

PF06850 is the intracellular Cys-type C-terminal marker, not an extracellular SBD. SignalP coverage is 100%: the 29,469 rows still labelled 'pending' were re-tested and predicted intracellular (99.82%); they are counted as predicted non-secreted here. Candidate-only evidence.