

1. A league has 16 teams. In the regular season each team plays each other 6 times. The season is played in 90 “rounds”. In each round every team plays a game. Each team’s rank is determined by number of wins. If two teams end up with the same record, their rank is determined via a new tie-breaker game. (What’s relevant is that teams that end up with the same record at the end of the regular season have unresolved relative ranks).

During the season every team does their best to increase their rank, until possibly reaching a time when they realize that their rank can’t change.

- What is the earliest possible round at which all teams in the league no longer care about improving their rank? Equivalently, how large could r be, so that with r regular season rounds remaining none of the teams care?
- You should include an example of all team win records and justify why all teams no longer care.
- You should explain why your value of r couldn’t be made larger. Ideally you should mention the pigeonhole principle.

Largest valid value of r :

Example of win records:

Justification:

Larger values of r wouldn’t work because ...

2. Simplify the following expression, making one modification per line.

On each line you should also briefly explain what rule was used to get that line from the one above.

$$\neg(\neg(p \wedge q) \vee \neg(\neg q \wedge r))$$

1. $\neg((\neg p \vee \neg q) \vee \neg(\neg q \wedge r)) \rightarrow$ Demorgan's on $\neg(p \wedge q)$

2. $\neg((\neg p \vee \neg q) \vee (q \vee \neg r)) \rightarrow$ Demorgan's on $\neg(\neg q \wedge r)$

3. $\neg(\neg p \vee \neg q \vee q \vee \neg r) \rightarrow$ Associative Law

4. $p \wedge q \wedge \neg q \wedge r \rightarrow$ Demorgan's on outer parenthesis

5. $p \wedge F \wedge r \rightarrow q \wedge \neg q$ annihilates to F

6. $F \rightarrow p \wedge F \wedge r$ is always F

3. Definition: $aXb = \neg(a \vee b)$

Show that we can simulate the following three expressions, with other expressions in which the only operator used is X . The expressions shouldn't include $\neg$, $\vee$ or negation either. You should provide step-by-step derivation with comments. Each change should be on a new line. Each line should be equivalent to the preceding line.

a) $\neg p$

$\neg p \leftrightarrow \neg p \vee \neg p$ as $p \vee p$ is always p

$\neg p \vee \neg p \leftrightarrow \neg(p \vee p)$ by deMorgan's laws

$\neg(p \vee p) \leftrightarrow pXp$ by definition

b) $p \vee q$

$p \vee q \iff \neg(\neg(p \vee q))$ a double logical not cancels out via deMorgan's laws

$\iff \neg(pXq)$ the X operator is defined with a logical not

$\iff (pXq)X(pXq)$ as $pXp \iff \neg p$ as proved above

c) $p \wedge q$

All of this is wrong, but the answer is $(pXp)X(qXq)$, I need another take to prove it though

$p \wedge q \iff \neg(\neg p \vee \neg q)$ as the preposition $p \wedge q \leftrightarrow \neg p \vee \neg q$ is false in all cases

$\neg(\neg p \vee \neg q)$