

# On the boundary of the ample cone of a hyperkähler manifold

Andrey Soldatenkov

UNICAMP

Geometry, algebra, and combinatorics  
IMECC, April 24, 2025

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\begin{aligned}\omega_I(u, v) &= g(Iu, v), \\ \omega_J(u, v) &= g(Ju, v), \\ \omega_K(u, v) &= g(Ku, v).\end{aligned}$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$

# Hyperkähler manifolds

## Definition

A *hyperkähler structure* on a  $C^\infty$ -manifold  $X$  is a tuple  $(g, I, J, K)$ , where:

- $g$  is a Riemannian metric;
- $I, J$  and  $K$  are complex structures s.t.  $IJ = -JI = K$ ;
- $g$  is Kähler w.r.t.  $I, J$  and  $K$ .

We have two-forms  $\omega_I, \omega_J$  and  $\omega_K$ :

$$\omega_I(u, v) = g(Iu, v),$$

$$\omega_J(u, v) = g(Ju, v),$$

$$\omega_K(u, v) = g(Ku, v).$$

These forms are closed:

$$d\omega_I = d\omega_J = d\omega_K = 0.$$



# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

Equivalently:  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed (2,0)-form on  $X_I$ ,  
i.e. a *holomorphic symplectic form*.

Today we assume: a hyperkähler manifold  $X$  is compact and of *maximal holonomy*, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed (2,0)-form on  $X_I$ ,  
i.e. a *holomorphic symplectic form*.

Today we assume: a hyperkähler manifold  $X$  is compact and of *maximal holonomy*, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed (2,0)-form on  $X_I$ ,  
i.e. a *holomorphic symplectic form*.

Today we assume: a hyperkähler manifold  $X$  is compact and of *maximal holonomy*, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed (2,0)-form on  $X_I$ ,  
i.e. a *holomorphic symplectic form*.

Today we assume: a hyperkähler manifold  $X$  is compact  
and of *maximal holonomy*, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\mathrm{Hol}(\nabla^g) \subset \mathrm{Sp}(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$\mathrm{Sp}(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed  $(2,0)$ -form on  $X_I$ ,  
i.e. a **holomorphic symplectic form**.

Today we assume: a hyperkähler manifold  $X$  is compact and of **maximal holonomy**, i.e.  $\mathrm{Hol}(\nabla^g) = \mathrm{Sp}(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed  $(2,0)$ -form on  $X_I$ ,  
i.e. a **holomorphic symplectic form**.

Today we assume: a hyperkähler manifold  $X$  is compact and of **maximal holonomy**, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

## Definition

A Riemannian metric  $g$  as above is called *hyperkähler*.

**Equivalently:**  $g$  is hyperkähler if  $\text{Hol}(\nabla^g) \subset Sp(n)$ ,

$\nabla^g$  is the Levi-Civita connection for  $g$ .

$Sp(n)$  = the group of quaternionic-linear transformations of  $\mathbb{H}^n$  that preserve the quaternionic-Hermitian scalar product.

Consider the 2-form  $\sigma_I = \omega_J + \sqrt{-1}\omega_K$ .

$\sigma_I$  is a non-degenerate closed  $(2,0)$ -form on  $X_I$ ,  
i.e. a **holomorphic symplectic form**.

Today we assume: a hyperkähler manifold  $X$  is compact and of **maximal holonomy**, i.e.  $\text{Hol}(\nabla^g) = Sp(n)$ .

This implies:  $\pi_1(X) = 1$  and  $H^0(X_I, \Omega_{X_I}^2) = \mathbb{C}\sigma_I$ .

# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .



# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .

# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .

# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .

# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .

# Hyperkähler manifolds

Since  $\sigma_I$  is symplectic, we have:

- $\dim_{\mathbb{C}}(X_I) = 2n$ ,
- $\sigma_I^n$  is a nowhere vanishing section of  $K_{X_I} = \Omega_{X_I}^{2n}$ .

Theorem (Beauville, Bogomolov, Fujiki)

*There exists  $c_X \in \mathbb{Q}$  such that for all  $a \in H^2(X, \mathbb{Q})$*

$$\int_X a^{2n} = c_X q(a)^n,$$

*where  $q$  is a quadratic form on  $H^2(X, \mathbb{Q})$ ,  
the Beauville–Bogomolov–Fujiki form, or the BBF form.*

We may assume:  $q$  is primitive and integral on  $H^2(X, \mathbb{Z})$ .

The signature of  $q$  is  $(3, b_2(X) - 3)$ .

# Examples of hyperkähler manifolds

- Complex  $K3$ -surfaces:  $S$  a compact simply connected complex surface with  $K_S \simeq \mathcal{O}_S$ . For example:

$$S = \{(x_0 : x_1 : x_2 : x_3) \in \mathbb{C}P^3 \mid x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0\}.$$

- $K3^{[n]}$ -type. Let  $S$  be a complex K3 surface.  
 $S^{[n]}$  = the Hilbert scheme of length  $n$  subschemes of  $S$ .
- Kum <sup>$n$</sup> -type. Let  $T = \mathbb{C}^2/\mathbb{Z}^4$ . The Albanese morphism:

$$a: T^{[n+1]} \rightarrow T, \quad (x_0, \dots, x_n) \mapsto \sum x_i$$

$K^n T = a^{-1}(0)$  = the generalized Kummer variety.

- $OG6$  and  $OG10$ -types. O'Grady's exceptional hyperkähler manifolds of dimensions 6 and 10.

# Examples of hyperkähler manifolds

- Complex *K3*-surfaces:  $S$  a compact simply connected complex surface with  $K_S \simeq \mathcal{O}_S$ . For example:

$$S = \{(x_0 : x_1 : x_2 : x_3) \in \mathbb{C}P^3 \mid x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0\}.$$

- $K3^{[n]}$ -type. Let  $S$  be a complex K3 surface.  
 $S^{[n]}$  = the Hilbert scheme of length  $n$  subschemes of  $S$ .
- Kum <sup>$n$</sup> -type. Let  $T = \mathbb{C}^2/\mathbb{Z}^4$ . The Albanese morphism:

$$a: T^{[n+1]} \rightarrow T, \quad (x_0, \dots, x_n) \mapsto \sum x_i$$

$K^n T = a^{-1}(0)$  = the generalized Kummer variety.

- *OG6* and *OG10*-types. O'Grady's exceptional hyperkähler manifolds of dimensions 6 and 10.

# Examples of hyperkähler manifolds

- Complex  $K3$ -surfaces:  $S$  a compact simply connected complex surface with  $K_S \simeq \mathcal{O}_S$ . For example:

$$S = \{(x_0 : x_1 : x_2 : x_3) \in \mathbb{C}P^3 \mid x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0\}.$$

- $K3^{[n]}$ -type. Let  $S$  be a complex K3 surface.  
 $S^{[n]}$  = the Hilbert scheme of length  $n$  subschemes of  $S$ .
- Kum <sup>$n$</sup> -type. Let  $T = \mathbb{C}^2/\mathbb{Z}^4$ . The Albanese morphism:

$$a: T^{[n+1]} \rightarrow T, \quad (x_0, \dots, x_n) \mapsto \sum x_i$$

$K^n T = a^{-1}(0)$  = the generalized Kummer variety.

- $OG6$  and  $OG10$ -types. O'Grady's exceptional hyperkähler manifolds of dimensions 6 and 10.



# Examples of hyperkähler manifolds

- Complex  $K3$ -surfaces:  $S$  a compact simply connected complex surface with  $K_S \simeq \mathcal{O}_S$ . For example:

$$S = \{(x_0 : x_1 : x_2 : x_3) \in \mathbb{C}P^3 \mid x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0\}.$$

- $K3^{[n]}$ -type. Let  $S$  be a complex K3 surface.  
 $S^{[n]}$  = the Hilbert scheme of length  $n$  subschemes of  $S$ .
- Kum <sup>$n$</sup> -type. Let  $T = \mathbb{C}^2/\mathbb{Z}^4$ . The Albanese morphism:

$$a: T^{[n+1]} \rightarrow T, \quad (x_0, \dots, x_n) \mapsto \sum x_i$$

$K^n T = a^{-1}(0)$  = the generalized Kummer variety.

- $OG6$  and  $OG10$ -types. O'Grady's exceptional hyperkähler manifolds of dimensions 6 and 10.

# Examples of hyperkähler manifolds

- Complex  $K3$ -surfaces:  $S$  a compact simply connected complex surface with  $K_S \simeq \mathcal{O}_S$ . For example:

$$S = \{(x_0 : x_1 : x_2 : x_3) \in \mathbb{C}P^3 \mid x_0^4 + x_1^4 + x_2^4 + x_3^4 = 0\}.$$

- $K3^{[n]}$ -type. Let  $S$  be a complex K3 surface.  
 $S^{[n]}$  = the Hilbert scheme of length  $n$  subschemes of  $S$ .
- Kum <sup>$n$</sup> -type. Let  $T = \mathbb{C}^2/\mathbb{Z}^4$ . The Albanese morphism:

$$a: T^{[n+1]} \rightarrow T, \quad (x_0, \dots, x_n) \mapsto \sum x_i$$

$K^n T = a^{-1}(0)$  = the generalized Kummer variety.

- $OG6$  and  $OG10$ -types. O'Grady's exceptional hyperkähler manifolds of dimensions 6 and 10.

# The ample cone

From now on:  $X$  is a projective hyperkähler manifold.

The Néron–Severi group:

$$\mathrm{NS}(X) = \mathrm{im} \left( \mathrm{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \right) \subset H^{1,1}(X, \mathbb{Z})$$

The restriction of  $q$  to  $\mathrm{NS}_{\mathbb{R}}(X) = \mathrm{NS}(X) \otimes \mathbb{R} \subset H^2(X, \mathbb{R})$  has signature

$$(1, \rho - 1)$$

where  $\rho$  is the Picard number of  $X$ .

The positive cone:

$$\mathcal{C}_X = \{x \in \mathrm{NS}_{\mathbb{R}}(X) \mid q(x) > 0\}^\circ$$

The ample cone:  $\mathcal{A}_X \subset \mathcal{C}_X$  is the convex cone spanned by the classes  $c_1(L)$  of ample line bundles  $L$ .

# The ample cone

From now on:  $X$  is a projective hyperkähler manifold.

The Néron–Severi group:

$$\mathrm{NS}(X) = \mathrm{im} \left( \mathrm{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \right) \subset H^{1,1}(X, \mathbb{Z})$$

The restriction of  $q$  to  $\mathrm{NS}_{\mathbb{R}}(X) = \mathrm{NS}(X) \otimes \mathbb{R} \subset H^2(X, \mathbb{R})$  has signature

$$(1, \rho - 1)$$

where  $\rho$  is the Picard number of  $X$ .

The positive cone:

$$\mathcal{C}_X = \{x \in \mathrm{NS}_{\mathbb{R}}(X) \mid q(x) > 0\}^\circ$$

The ample cone:  $\mathcal{A}_X \subset \mathcal{C}_X$  is the convex cone spanned by the classes  $c_1(L)$  of ample line bundles  $L$ .

# The ample cone

From now on:  $X$  is a projective hyperkähler manifold.

The Néron–Severi group:

$$\mathrm{NS}(X) = \mathrm{im} \left( \mathrm{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \right) \subset H^{1,1}(X, \mathbb{Z})$$

The restriction of  $q$  to  $\mathrm{NS}_{\mathbb{R}}(X) = \mathrm{NS}(X) \otimes \mathbb{R} \subset H^2(X, \mathbb{R})$  has signature

$$(1, \rho - 1)$$

where  $\rho$  is the Picard number of  $X$ .

The positive cone:

$$\mathcal{C}_X = \{x \in \mathrm{NS}_{\mathbb{R}}(X) \mid q(x) > 0\}^\circ$$

The ample cone:  $\mathcal{A}_X \subset \mathcal{C}_X$  is the convex cone spanned by the classes  $c_1(L)$  of ample line bundles  $L$ .

# The ample cone

From now on:  $X$  is a projective hyperkähler manifold.

The Néron–Severi group:

$$\mathrm{NS}(X) = \mathrm{im} \left( \mathrm{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \right) \subset H^{1,1}(X, \mathbb{Z})$$

The restriction of  $q$  to  $\mathrm{NS}_{\mathbb{R}}(X) = \mathrm{NS}(X) \otimes \mathbb{R} \subset H^2(X, \mathbb{R})$  has signature

$$(1, \rho - 1)$$

where  $\rho$  is the Picard number of  $X$ .

The positive cone:

$$\mathcal{C}_X = \{x \in \mathrm{NS}_{\mathbb{R}}(X) \mid q(x) > 0\}^\circ$$

The ample cone:  $\mathcal{A}_X \subset \mathcal{C}_X$  is the convex cone spanned by the classes  $c_1(L)$  of ample line bundles  $L$ .

# The ample cone

From now on:  $X$  is a projective hyperkähler manifold.

The Néron–Severi group:

$$\mathrm{NS}(X) = \mathrm{im} \left( \mathrm{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \right) \subset H^{1,1}(X, \mathbb{Z})$$

The restriction of  $q$  to  $\mathrm{NS}_{\mathbb{R}}(X) = \mathrm{NS}(X) \otimes \mathbb{R} \subset H^2(X, \mathbb{R})$  has signature

$$(1, \rho - 1)$$

where  $\rho$  is the Picard number of  $X$ .

The positive cone:

$$\mathcal{C}_X = \{x \in \mathrm{NS}_{\mathbb{R}}(X) \mid q(x) > 0\}^\circ$$

The ample cone:  $\mathcal{A}_X \subset \mathcal{C}_X$  is the convex cone spanned by the classes  $c_1(L)$  of ample line bundles  $L$ .

# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called monodromy birationally minimal or MBM classes with the following properties.

- MBM is  $\mathcal{D}iff(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.



# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called monodromy birationally minimal or MBM classes with the following properties.

- MBM is  $\mathcal{D}\text{iff}(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.

# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called **monodromy birationally minimal** or **MBM** classes with the following properties.

- **MBM** is  $\mathcal{D}iff(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.

# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called **monodromy birationally minimal** or **MBM** classes with the following properties.

- **MBM** is  $\mathcal{D}iff(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.

# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called monodromy birationally minimal or MBM classes with the following properties.

- MBM is  $\mathcal{D}iff(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.

# The ample cone

We have:  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is a hyperbolic space. What is  $\mathbb{P}(\mathcal{A}_X)$ ?

Amerik, Verbitsky: there is a collection of integral classes

$$\text{MBM} \subset H^2(X, \mathbb{Z})$$

called **monodromy birationally minimal** or **MBM** classes with the following properties.

- **MBM** is  $\mathcal{D}iff(X)$ -invariant;
- There exists a constant  $M > 0$  such that for all  $x \in \text{MBM}$  we have  $-M \leq q(x) < 0$ ;
- Let  $\text{MBM}^{1,1} = \text{MBM} \cap \text{NS}(X)$ . Then the hyperplanes  $x^\perp$ , where  $x \in \text{MBM}^{1,1}$ , cut  $\mathcal{C}_X$  into open chambers.
- $\mathcal{A}_X$  is one of the chambers.

# The ample cone of a K3 surface

Let  $S$  be a projective K3 surface.

Then  $q$  is the intersection form and

$$H^2(S, \mathbb{Z}) \simeq (-E_8)^{\oplus 2} \oplus U^{\oplus 3},$$

where  $U$  is the rank two hyperbolic lattice  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

We have

$$\text{MBM} = \{x \in H^2(S, \mathbb{Z}) \mid q(x) = -2\}.$$

Theorem (Nikulin)

*Let  $\Lambda$  be an even lattice of signature  $(1, \rho - 1)$  where  $\rho \leq 10$ .  
Then there exists a projective K3 surface  $S$  with  $\text{NS}(S) \simeq \Lambda$ .*

# The ample cone of a K3 surface

Let  $S$  be a projective K3 surface.

Then  $q$  is the intersection form and

$$H^2(S, \mathbb{Z}) \simeq (-E_8)^{\oplus 2} \oplus U^{\oplus 3},$$

where  $U$  is the rank two hyperbolic lattice  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

We have

$$\text{MBM} = \{x \in H^2(S, \mathbb{Z}) \mid q(x) = -2\}.$$

## Theorem (Nikulin)

*Let  $\Lambda$  be an even lattice of signature  $(1, \rho - 1)$  where  $\rho \leq 10$ .  
Then there exists a projective K3 surface  $S$  with  $\text{NS}(S) \simeq \Lambda$ .*

# The ample cone of a K3 surface

Let  $S$  be a projective K3 surface.

Then  $q$  is the intersection form and

$$H^2(S, \mathbb{Z}) \simeq (-E_8)^{\oplus 2} \oplus U^{\oplus 3},$$

where  $U$  is the rank two hyperbolic lattice  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

We have

$$\text{MBM} = \{x \in H^2(S, \mathbb{Z}) \mid q(x) = -2\}.$$

## Theorem (Nikulin)

*Let  $\Lambda$  be an even lattice of signature  $(1, \rho - 1)$  where  $\rho \leq 10$ .  
Then there exists a projective K3 surface  $S$  with  $\text{NS}(S) \simeq \Lambda$ .*



# The ample cone of a K3 surface

Let  $S$  be a projective K3 surface.

Then  $q$  is the intersection form and

$$H^2(S, \mathbb{Z}) \simeq (-E_8)^{\oplus 2} \oplus U^{\oplus 3},$$

where  $U$  is the rank two hyperbolic lattice  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

We have


$$\text{MBM} = \{x \in H^2(S, \mathbb{Z}) \mid q(x) = -2\}.$$

## Theorem (Nikulin)

Let  $\Lambda$  be an even lattice of signature  $(1, \rho - 1)$  where  $\rho \leq 10$ .  
Then there exists a projective K3 surface  $S$  with  $\text{NS}(S) \simeq \Lambda$ .

# The ample cone of a K3 surface

For example, when  $\rho = 3$ , we have  $\mathbb{P}(\mathcal{C}_S) = \mathbb{H}^2$   
and  $\mathbb{P}(\mathcal{A}_S)$  may look like this:



# The ideal boundary of the ample cone

Identify  $\mathrm{NS}_{\mathbb{R}}(X)$  with  $\mathbb{R}^{1,\rho-1}$ , so that

$$q(x) = x_0^2 - x_1^2 - \dots - x_{\rho-1}^2.$$

Then  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is identified with the unit ball  $\mathbb{B}^{\rho-1}$  via the stereographic projection from the point  $(-1, 0, \dots, 0)$  and

$$\partial\mathbb{P}(\mathcal{C}_X) = \partial\mathbb{H}^{\rho-1} \simeq \mathbb{S}^{\rho-2}$$

is the sphere at infinity.

## Definition

*The ideal boundary of the ample cone:*

$$\mathcal{B}_X = \overline{\mathbb{P}(\mathcal{A}_X)} \cap \partial\mathbb{P}(\mathcal{C}_X)$$

# The ideal boundary of the ample cone

Identify  $\mathrm{NS}_{\mathbb{R}}(X)$  with  $\mathbb{R}^{1,\rho-1}$ , so that

$$q(x) = x_0^2 - x_1^2 - \dots - x_{\rho-1}^2.$$

Then  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is identified with the unit ball  $\mathbb{B}^{\rho-1}$  via the stereographic projection from the point  $(-1, 0, \dots, 0)$  and

$$\partial\mathbb{P}(\mathcal{C}_X) = \partial\mathbb{H}^{\rho-1} \simeq \mathbb{S}^{\rho-2}$$

is the sphere at infinity.

## Definition

*The ideal boundary of the ample cone:*

$$\mathcal{B}_X = \overline{\mathbb{P}(\mathcal{A}_X)} \cap \partial\mathbb{P}(\mathcal{C}_X)$$

# The ideal boundary of the ample cone

Identify  $\mathrm{NS}_{\mathbb{R}}(X)$  with  $\mathbb{R}^{1,\rho-1}$ , so that

$$q(x) = x_0^2 - x_1^2 - \dots - x_{\rho-1}^2.$$

Then  $\mathbb{P}(\mathcal{C}_X) = \mathbb{H}^{\rho-1}$  is identified with the unit ball  $\mathbb{B}^{\rho-1}$  via the stereographic projection from the point  $(-1, 0, \dots, 0)$  and

$$\partial\mathbb{P}(\mathcal{C}_X) = \partial\mathbb{H}^{\rho-1} \simeq \mathbb{S}^{\rho-2}$$

is the sphere at infinity.

## Definition

*The ideal boundary of the ample cone:*

$$\mathcal{B}_X = \overline{\mathbb{P}(\mathcal{A}_X)} \cap \partial\mathbb{P}(\mathcal{C}_X)$$

# The ideal boundary of the ample cone

The following result was classically known for K3 surfaces due to S. Kovács, and was more recently generalized to an arbitrary hyperkähler manifold by F. Denisi.

Theorem (Kovács, Denisi)

*Assume that  $X$  is a projective hyperkähler manifold with Picard number  $\rho > 2$ . Then we have the following dichotomy.*

- *If  $\mathcal{B}_X$  contains an open subset of  $\mathbb{S}^{\rho-2}$ , then  $\mathcal{B}_X = \mathbb{S}^{\rho-2}$  and  $\mathcal{A}_X = \mathcal{C}_X$ . This happens if and only if  $\mathrm{NS}(X)$  does not contain MBM classes.*
- *Otherwise  $\mathcal{B}_X$  is nowhere dense in  $\mathbb{S}^{\rho-2}$ .*

# The ideal boundary of the ample cone

The following result was classically known for K3 surfaces due to S. Kovács, and was more recently generalized to an arbitrary hyperkähler manifold by F. Denisi.

## Theorem (Kovács, Denisi)

*Assume that  $X$  is a projective hyperkähler manifold with Picard number  $\rho > 2$ . Then we have the following dichotomy.*

- *If  $\mathcal{B}_X$  contains an open subset of  $\mathbb{S}^{\rho-2}$ , then  $\mathcal{B}_X = \mathbb{S}^{\rho-2}$  and  $\mathcal{A}_X = \mathcal{C}_X$ . This happens if and only if  $\mathrm{NS}(X)$  does not contain MBM classes.*
- *Otherwise  $\mathcal{B}_X$  is nowhere dense in  $\mathbb{S}^{\rho-2}$ .*

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$



# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

In general,  $\mathcal{B}_X$  is a fractal in  $\mathbb{S}^{\rho-2}$ .

Let  $v \in MBM \cap NS(X) = MBM^{1,1}$ . Then:

- $\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a hyperbolic subspace  $\mathbb{H}_v^{\rho-2} \subset \mathbb{H}^{\rho-1}$ .
- $\partial\mathbb{P}(v^\perp \cap \mathcal{C}_X)$  is a sphere  $\mathbb{S}_v^{\rho-3} \subset \mathbb{S}^{\rho-2}$ .

Denote by  $H_v \subset \mathbb{R}^{\rho-1}$  the hyperplane such that

$$\mathbb{S}_v^{\rho-3} = \mathbb{S}^{\rho-2} \cap H_v.$$

Let  $H_v^+$  and  $H_v^-$  be the two open half-spaces in  $\mathbb{R}^{\rho-1}$  separated by  $H_v$ , and such that  $\mathcal{A}_X \subset H_v^+$ .

Then

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

# The ideal boundary of the ample cone

We have

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

Consider the stereographic projection of  $\mathbb{S}^{\rho-2}$  onto  $\mathbb{R}^{\rho-2}$ . Then  $D_v = \mathbb{S}^{\rho-2} \cap H_v^-$  is an open disc in  $\mathbb{R}^{\rho-2}$ .

Under the stereographic projection:

$$\mathcal{B}_X = \mathbb{R}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} D_v.$$

**Remark:** If  $\rho > 2$  and  $MBM^{1,1} \neq \emptyset$ , then  $MBM^{1,1}$  is infinite, because it is stable under the action of a finite index subgroup in  $O(NS(X), q)$ .

# The ideal boundary of the ample cone

We have

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

Consider the stereographic projection of  $\mathbb{S}^{\rho-2}$  onto  $\mathbb{R}^{\rho-2}$ . Then  $D_v = \mathbb{S}^{\rho-2} \cap H_v^-$  is an open disc in  $\mathbb{R}^{\rho-2}$ .

Under the stereographic projection:

$$\mathcal{B}_X = \mathbb{R}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} D_v.$$

**Remark:** If  $\rho > 2$  and  $MBM^{1,1} \neq \emptyset$ , then  $MBM^{1,1}$  is infinite, because it is stable under the action of a finite index subgroup in  $O(NS(X), q)$ .



# The ideal boundary of the ample cone

We have

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

Consider the stereographic projection of  $\mathbb{S}^{\rho-2}$  onto  $\mathbb{R}^{\rho-2}$ . Then  $D_v = \mathbb{S}^{\rho-2} \cap H_v^-$  is an open disc in  $\mathbb{R}^{\rho-2}$ .

Under the stereographic projection:

$$\mathcal{B}_X = \mathbb{R}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} D_v.$$

**Remark:** If  $\rho > 2$  and  $MBM^{1,1} \neq \emptyset$ , then  $MBM^{1,1}$  is infinite, because it is stable under the action of a finite index subgroup in  $O(NS(X), q)$ .

# The ideal boundary of the ample cone

We have

$$\mathcal{B}_X = \mathbb{S}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} H_v^-.$$

Consider the stereographic projection of  $\mathbb{S}^{\rho-2}$  onto  $\mathbb{R}^{\rho-2}$ . Then  $D_v = \mathbb{S}^{\rho-2} \cap H_v^-$  is an open disc in  $\mathbb{R}^{\rho-2}$ .


Under the stereographic projection:

$$\mathcal{B}_X = \mathbb{R}^{\rho-2} \setminus \bigcup_{v \in MBM^{1,1}} D_v.$$

**Remark:** If  $\rho > 2$  and  $MBM^{1,1} \neq \emptyset$ , then  $MBM^{1,1}$  is infinite, because it is stable under the action of a finite index subgroup in  $O(NS(X), q)$ .


# The ideal boundary of the ample cone

For example, if  $\rho = 4$ , then  $\mathcal{B}_X$  may look like this:



# The ideal boundary of the ample cone

The discs  $D_v$  in the above example look like this:



# The ideal boundary of the ample cone

In the above example we used the lattice  $\Lambda$  with the intersection matrix

$$A = \begin{pmatrix} -2 & 4 & 0 & 0 \\ 4 & -2 & 2 & 0 \\ 0 & 2 & -4 & 2 \\ 0 & 0 & 2 & -4 \end{pmatrix}$$

If we start from the intersection matrix

$$A = \begin{pmatrix} -2 & 2 & 2 & 2 \\ 2 & -2 & 2 & 2 \\ 2 & 2 & -2 & 2 \\ 2 & 2 & 2 & -2 \end{pmatrix},$$

we get the Apollonian gasket.

# The ideal boundary of the ample cone

In the above example we used the lattice  $\Lambda$  with the intersection matrix


$$A = \begin{pmatrix} -2 & 4 & 0 & 0 \\ 4 & -2 & 2 & 0 \\ 0 & 2 & -4 & 2 \\ 0 & 0 & 2 & -4 \end{pmatrix}$$

If we start from the intersection matrix

$$A = \begin{pmatrix} -2 & 2 & 2 & 2 \\ 2 & -2 & 2 & 2 \\ 2 & 2 & -2 & 2 \\ 2 & 2 & 2 & -2 \end{pmatrix},$$

we get the Apollonian gasket.


# The Apollonian gasket as the ideal boundary



## Definition

*The Apollonian carpet of  $X$  is the union of all positive-dimensional real-analytic subvarieties contained in  $\mathcal{B}_X$ .*

# The Apollonian gasket as the ideal boundary



## Definition

The *Apollonian carpet* of  $X$  is the union of all positive-dimensional real-analytic subvarieties contained in  $\mathcal{B}_X$ .



# The Apollonian carpet

Theorem (Amerik, S., Verbitsky)

Let  $X$  be a projective hyperkähler manifold with  $\rho \geq 4$ , and  $Z$  a germ of a positive-dimensional irreducible real-analytic subset of  $\mathcal{B}_X$ . Then:

- There exists a sublattice  $\Lambda \subset \mathrm{NS}(X)$  of signature  $(1, d)$  for some  $d \leq \rho$ , such that

$$Z \subset \mathbb{S}_\Lambda^d \subset \mathcal{B}_X,$$

where  $\mathbb{S}_\Lambda^d = \partial \mathbb{P}((\Lambda \otimes \mathbb{R}) \cap \mathcal{C}_X)$  is the ideal boundary of the corresponding hyperbolic subspace;

- The Apollonian carpet of  $X$  is the union of the spheres  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above;
- If the Apollonian carpet of  $X$  is non-empty, then its closure is equal to  $\mathcal{B}_X$ ;
- For any  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above,  $\mathrm{Aut}(X)$  contains a subgroup commensurable with  $\mathrm{O}(\Lambda)$ .

# The Apollonian carpet

Theorem (Amerik, S., Verbitsky)

Let  $X$  be a projective hyperkähler manifold with  $\rho \geq 4$ , and  $Z$  a germ of a positive-dimensional irreducible real-analytic subset of  $\mathcal{B}_X$ . Then:

- There exists a sublattice  $\Lambda \subset \mathrm{NS}(X)$  of signature  $(1, d)$  for some  $d \leq \rho$ , such that

$$Z \subset \mathbb{S}_{\Lambda}^d \subset \mathcal{B}_X,$$

where  $\mathbb{S}_{\Lambda}^d = \partial \mathbb{P}((\Lambda \otimes \mathbb{R}) \cap \mathcal{C}_X)$  is the ideal boundary of the corresponding hyperbolic subspace;

- The Apollonian carpet of  $X$  is the union of the spheres  $\mathbb{S}_{\Lambda}^d \subset \mathcal{B}_X$  as above;
- If the Apollonian carpet of  $X$  is non-empty, then its closure is equal to  $\mathcal{B}_X$ ;
- For any  $\mathbb{S}_{\Lambda}^d \subset \mathcal{B}_X$  as above,  $\mathrm{Aut}(X)$  contains a subgroup commensurable with  $\mathrm{O}(\Lambda)$ .

# The Apollonian carpet

Theorem (Amerik, S., Verbitsky)

Let  $X$  be a projective hyperkähler manifold with  $\rho \geq 4$ , and  $Z$  a germ of a positive-dimensional irreducible real-analytic subset of  $\mathcal{B}_X$ . Then:

- There exists a sublattice  $\Lambda \subset \mathrm{NS}(X)$  of signature  $(1, d)$  for some  $d \leq \rho$ , such that

$$Z \subset \mathbb{S}_\Lambda^d \subset \mathcal{B}_X,$$

where  $\mathbb{S}_\Lambda^d = \partial\mathbb{P}((\Lambda \otimes \mathbb{R}) \cap \mathcal{C}_X)$  is the ideal boundary of the corresponding hyperbolic subspace;

- The Apollonian carpet of  $X$  is the union of the spheres  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above;
- If the Apollonian carpet of  $X$  is non-empty, then its closure is equal to  $\mathcal{B}_X$ ;
- For any  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above,  $\mathrm{Aut}(X)$  contains a subgroup commensurable with  $\mathrm{O}(\Lambda)$ .

# The Apollonian carpet

Theorem (Amerik, S., Verbitsky)

Let  $X$  be a projective hyperkähler manifold with  $\rho \geq 4$ , and  $Z$  a germ of a positive-dimensional irreducible real-analytic subset of  $\mathcal{B}_X$ . Then:

- There exists a sublattice  $\Lambda \subset \text{NS}(X)$  of signature  $(1, d)$  for some  $d \leq \rho$ , such that

$$Z \subset \mathbb{S}_\Lambda^d \subset \mathcal{B}_X,$$

where  $\mathbb{S}_\Lambda^d = \partial\mathbb{P}((\Lambda \otimes \mathbb{R}) \cap \mathcal{C}_X)$  is the ideal boundary of the corresponding hyperbolic subspace;

- The Apollonian carpet of  $X$  is the union of the spheres  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above;
- If the Apollonian carpet of  $X$  is non-empty, then its closure is equal to  $\mathcal{B}_X$ ;
- For any  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above,  $\text{Aut}(X)$  contains a subgroup commensurable with  $\text{O}(\Lambda)$ .

# The Apollonian carpet

Theorem (Amerik, S., Verbitsky)

Let  $X$  be a projective hyperkähler manifold with  $\rho \geq 4$ , and  $Z$  a germ of a positive-dimensional irreducible real-analytic subset of  $\mathcal{B}_X$ . Then:

- There exists a sublattice  $\Lambda \subset \mathrm{NS}(X)$  of signature  $(1, d)$  for some  $d \leq \rho$ , such that

$$Z \subset \mathbb{S}_\Lambda^d \subset \mathcal{B}_X,$$


where  $\mathbb{S}_\Lambda^d = \partial\mathbb{P}((\Lambda \otimes \mathbb{R}) \cap \mathcal{C}_X)$  is the ideal boundary of the corresponding hyperbolic subspace;

- The Apollonian carpet of  $X$  is the union of the spheres  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above;
- If the Apollonian carpet of  $X$  is non-empty, then its closure is equal to  $\mathcal{B}_X$ ;
- For any  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$  as above,  $\mathrm{Aut}(X)$  contains a subgroup commensurable with  $\mathrm{O}(\Lambda)$ .

## The boundary spheres


There are two types of boundary spheres  $\mathbb{S}_\Lambda^d \subset \mathcal{B}_X$ .

The first type corresponds to the case when there exists  $v \in \text{MBM}^{1,1} \cap \Lambda^\perp$ .




## The boundary spheres

The second type: there is no  $v \in \text{MBM}^{1,1} \cap \Lambda^\perp$ .




# Examples






# Examples



# Examples



Thank you!