MATH 110 — Week 10

Notes Chapter 18; Homework 10

§1 Orthogonal and orthonormal lists

The question. When do squared lengths add without mixed terms? For $(a, 0)$ and $(0, b)$ in the standard product on $\mathbb{R}^2$, the squared length of their sum is $a^2 + b^2$.

Tools and conventions. Throughout, $F = \mathbb{R}$ or $\mathbb{C}$, and inner products are linear in the first argument.

- Week 9: $\|v\|^2 = \langle v, v\rangle$, $\langle u, av\rangle = \overline{a}\langle u, v\rangle$, and $|\langle u, v\rangle| \le \|u\|\|v\|$ by expanding a nonnegative squared length.
- Weeks 2–5: independent lists extend to bases; dimensions add in a direct sum; $\dim V' = \dim V$ by dual bases; finite-dimensional injective maps between equal-dimensional spaces are isomorphisms by rank-nullity.
- Weeks 6–8: polynomial evaluation commutes with T; m_T divides every annihilating polynomial by division with remainder; its roots are eigenvalues. Week 7 characterizes upper-triangular form by invariant initial spans.
- Week 9: trace is the sum of the diagonal in any basis, because cyclic interchange of matrix factors proves invariance under similarity.

Definition (Notes Definition 46; LADR 6.10, 6.22, 6.27)**.** Vectors u, v are orthogonal, written $u \perp v$, if $\langle u, v\rangle = 0$. A list is orthogonal if every two distinct members are orthogonal; its members need not be nonzero unless stated. It is orthonormal if $\langle e_i, e_j\rangle = \delta_{ij}$, where δ_{ij} is 1 for $i = j$ and 0 otherwise. An orthonormal basis is a vector-space basis which is orthonormal; vector-space basis means every vector is a finite linear combination.

Proposition (LADR 6.11–6.13: elementary orthogonality)

Only zero is orthogonal to every vector. If $u \perp v$, then $\|u + v\|^2 = \|u\|^2 + \|v\|^2$ (Pythagorean theorem). For nonzero u, every v has a unique orthogonal decomposition $v = au + z$ with $a = \langle v, u\rangle/\|u\|^2$ and $z \perp u$.

Where the proof comes from

Want. Remove mixed terms and separate a prescribed vector direction.
Know. Expanding a squared length gives a pair of conjugate mixed terms.
Key step. Orthogonality makes both terms zero; choose a by solving $\langle v - au, u\rangle = 0$.

Proof. 1. If w is orthogonal to every vector, take the vector w itself: $\|w\|^2 = 0$, so $w = 0$. Zero is orthogonal to everything by Week 9.

2. If $\langle u, v\rangle = 0$, conjugate symmetry also gives $\langle v, u\rangle = 0$. Expanding $\langle u+v, u+v\rangle$ proves the squared-length identity.

3. For the stated a, $\langle v - au, u\rangle = \langle v, u\rangle - a\|u\|^2 = 0$. Thus $z = v - au$ gives the claimed decomposition. In any such decomposition, taking the inner product with u forces $a = \langle v, u\rangle/\|u\|^2$, and then $z = v - au$ is forced as well.

□

Proposition (Notes Proposition 84; LADR 6.24)

For an orthonormal list $e_1, \ldots, e_m$ and scalars a_j, $\|\sum_j a_j e_j\|^2 = \sum_j |a_j|^2$.

Where the proof comes from

Want. Compute a linear combination's length from its coefficients.
Know. Every cross inner product in an orthonormal list is zero.
Key step. Expand the finite double sum and retain only diagonal terms.

Proof. 1. Sesquilinearity gives $\langle \sum_i a_i e_i, \sum_j a_j e_j \rangle = \sum_{i,j} a_i \overline{a_j} \langle e_i, e_j \rangle$.
2. Substitute $\langle e_i, e_j \rangle = \delta_{ij}$ to obtain $\sum_j a_j \overline{a_j} = \sum_j |a_j|^2$.

□

Proposition (Notes Proposition 85; LADR 6.25, 6.28)
An orthonormal list is independent. On an n-dimensional space, an orthonormal list of length n is therefore an orthonormal basis.

Where the proof comes from

Want. Recognize a basis without solving a general dependence system.
Know. A sum of nonnegative squared moduli is zero only when each modulus is zero.
Key step. Take the squared norm of a proposed dependence.

Proof. 1. If $\sum_j a_j e_j = 0$, Proposition 84 gives $0 = \sum_j |a_j|^2$, hence every $a_j = 0$.
2. An independent length-n list in dimension n is a basis by Week 2.

□

Example (LADR 6.23: coordinate and function examples) 1. The standard coordinate vectors are orthonormal because their dot products are δ_{ij}. The vectors $(1,1,1)/\sqrt{3}$, $(-1,1,0)/\sqrt{2}$ and $(1,1,-2)/\sqrt{6}$ also form an orthonormal basis of F^3: their squared norms are 1 and their pairwise dot products are 0.
2. On $[-\pi, \pi]$ with the integral product, the real list $1/\sqrt{2\pi}$, $\cos(kx)/\sqrt{\pi}$, $\sin(kx)/\sqrt{\pi}$ for $k = 1, 2, \ldots$ is orthonormal.
3. Indeed, the product-to-sum trigonometric identities and integration give $\int \cos(kx)\cos(\ell x) = \pi\delta_{k\ell}$, $\int \sin(kx)\sin(\ell x) = \pi\delta_{k\ell}$ and every sine–cosine integral 0 for positive integers k, ℓ; each integral here is over $[-\pi, \pi]$. Also $\int 1 = 2\pi$ and every nonconstant sine/cosine has integral 0. These calculations use elementary calculus and trigonometric identities.

Proposition (Notes Proposition 86; LADR 6.30)
For an orthonormal basis $e_1, \ldots, e_n$, $v = \sum_j \langle v, e_j \rangle e_j$ and $\|v\|^2 = \sum_j |\langle v, e_j \rangle|^2$.

Where the proof comes from

Want. Recover coordinates by inner products instead of elimination.
Know. A basis gives unique coefficients and orthonormality removes all but one coefficient in a mixed product.
Key step. Take the inner product with e_k in the second position.

Proof. 1. Write $v = \sum_j a_j e_j$. Then $\langle v, e_k \rangle = \sum_j a_j \langle e_j, e_k \rangle = a_k$. Substitute these coefficients.
2. Apply Proposition 84 to the resulting expansion to obtain the norm formula.

□

Read coordinates in an orthonormal basis (LADR 6.29, 6.31)

1. Use the four columns $e_1 = (1,1,1,1)/2$, $e_2 = (1,1,-1,-1)/2$, $e_3 = (1,-1,-1,1)/2$, $e_4 = (-1,1,-1,1)/2$. Direct dot products give the identity matrix of pairwise inner products, hence an orthonormal basis.
2. For $v = (1,2,4,7)$, its four dot products are $7, -4, 1, 2$, so $v = 7e_1 - 4e_2 + e_3 + 2e_4$.

3. The norm check is $\|v\|^2 = 1 + 4 + 16 + 49 = 70 = 49 + 16 + 1 + 4$, as Proposition 86 predicts.

Proposition (LADR 6.26: Bessel's inequality)
For an orthonormal list $e_1, \ldots, e_m$, $\sum_j |\langle v, e_j\rangle|^2 \le \|v\|^2$, with equality exactly when $v \in \operatorname{span}(e_1, \ldots, e_m)$.

Where the proof comes from

Want. Bound the coefficients even when the list is not a basis.
Know. Subtracting the indicated components leaves a vector orthogonal to the list.
Key step. Apply Pythagoras to that remainder and the finite combination.

Proof. 1. Put $u = \sum_j \langle v, e_j\rangle e_j$ and $z = v - u$. Taking inner products with e_k gives $\langle z, e_k\rangle = 0$; by conjugate linearity, $z \perp u$.
2. Pythagoras and Proposition 84 give $\|v\|^2 = \|u\|^2 + \|z\|^2 = \sum_j |\langle v, e_j\rangle|^2 + \|z\|^2$.
3. Equality holds exactly when $z = 0$. This implies v lies in the span, and the coefficient formula proves the converse for such v.

□

Proposition (LADR 8.55: trace in an orthonormal basis)
For finite-dimensional V, $\operatorname{tr} T = \sum_j \langle Te_j, e_j\rangle$ in any orthonormal basis.

Where the proof comes from

Want. Compute trace using inner products.
Know. Trace sums the diagonal matrix coefficients; Proposition 86 extracts coefficients.
Key step. Extract coefficient j from the image of basis vector j.

Proof. 1. Proposition 86 applied to Te_j shows its jth coordinate is $\langle Te_j, e_j\rangle$.
2. These are precisely the matrix's diagonal entries; summing gives its trace, which is basis independent by Week 9.

□

Cited analytic tools for the notes' Fourier aside. In $L^2[0,1]$, the exponential functions $e^{2\pi inx}$, $n \in \mathbb{Z}$, are a complete orthonormal system: their finite combinations approximate each L^2 function in norm. The analytic Parseval theorem states that $\|f\|^2 = \sum_{n\in\mathbb{Z}} |c_n|^2$ for $c_n = \int_0^1 f(x)e^{-2\pi inx}\,dx$. This infinite-series assertion requires analytic completeness and convergence theory; it is cited, rather than inferred from the finite identity above.

Apply the cited Parseval theorem (Notes Chapter 18)

1. For $f(x) = x$, integration gives $c_0 = 1/2$. Integration by parts gives $c_n = -1/(2\pi i n)$ for $n \neq 0$, since $e^{-2\pi i n} = 1$.
2. The corresponding Fourier series is $x = 1/2 - (2\pi i)^{-1} \sum_{n \neq 0} e^{2\pi i n x}/n$ for $0 < x < 1$ (the interior convergence statement is also a cited Fourier theorem).
3. Parseval gives $1/4 + (4\pi^2)^{-1} \sum_{n \neq 0} 1/n^2 = \int_0^1 x^2\, dx = 1/3$. The terms at n and $-n$ agree, so $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$.

Shortcuts

- In an orthonormal basis, coordinates are $\langle v, e_j \rangle$ and squared length is their squared-modulus sum.
- Bessel's inequality detects whether the list accounts for all of a given vector: equality means it lies in the list's span.

Traps

- An orthogonal list containing zero need not be independent; the list $(e_1, 0)$ is a counterexample. Unit norm excludes zero for orthonormal lists.
- In a complex space, the Pythagorean equality does not force orthogonality: $u = 1, v = i$ give $|1 + i|^2 = 2 = |1|^2 + |i|^2$, but $\langle 1, i \rangle = -i \neq 0$.

§2 Gram–Schmidt and orthonormal triangular bases

The question. How can one turn a basis into an orthonormal basis without changing its initial spans? Starting with $1, x$ on $[0, 1]$, subtracting the component $1/2$ from x gives $x - 1/2$, orthogonal to 1.

Proposition (Notes Proposition 87; LADR 6.32, 6.35)

For an independent list $v_1, \ldots, v_n$, define $r_j = v_j - \sum_{i<j} \langle v_j, e_i \rangle e_i$ and $e_j = r_j / \|r_j\|$, with the empty sum at $j = 1$. These vectors are well defined and orthonormal, and their first j spans equal $\operatorname{span}(v_1, \ldots, v_j)$. In particular a finite-dimensional inner-product space has an orthonormal basis.

Where the proof comes from

Want. Orthogonalize one vector while preserving all earlier directions.
Know. Orthonormal coefficients are obtained by inner products.
First try. Normalizing alone leaves mixed products: $(1, 0)$ and $(1, 1)/\sqrt{2}$ have product $1/\sqrt{2}$.
Key step. Subtract those coefficients and normalize; independence ensures the remainder is nonzero.

Proof. 1. Induct on j. For $j = 1$, $v_1 \neq 0$ and division by its norm produces a unit vector with the same span.
2. Assume the preceding list is orthonormal with the required spans. For $k < j$, $\langle r_j, e_k \rangle = \langle v_j, e_k \rangle - \langle v_j, e_k \rangle = 0$.
3. If $r_j = 0$, then v_j is in the preceding span, contradicting independence. Thus e_j is defined, unit length, and orthogonal to the preceding list.
4. The formula puts e_j in the first j original span and gives $v_j = \|r_j\| e_j + \sum_{i<j} \langle v_j, e_i \rangle e_i$. Both span inclusions follow, completing the induction.

□

Normalize shifted Legendre polynomials (Notes Example 70)

1. On $\mathcal{P}_2(\mathbb{C})$ with $\langle p, q \rangle = \int_0^1 p\bar{q}$, start with $1, x, x^2$. Since $\|1\| = 1$, $e_1 = 1$.
2. $r_2 = x - \langle x, 1 \rangle = x - 1/2$, with $\|r_2\|^2 = 1/12$. Thus $e_2 = \sqrt{3}(2x - 1)$.
3. The coefficients for x^2 are $\langle x^2, e_1 \rangle = 1/3$ and $\langle x^2, e_2 \rangle = \sqrt{3}/6$. Hence $r_3 = x^2 - 1/3 - (2x - 1)/2 = x^2 - x + 1/6$.
4. Its squared norm is $1/180$, so $e_3 = \sqrt{5}(6x^2 - 6x + 1)$. The three pairwise products and norms give the identity matrix of pairwise inner products.

Use unnormalized orthogonal vectors (LADR 6.34; HW10 P1)

1. One may first form $u_j = v_j - \sum_{i<j} \langle v_j, u_i \rangle u_i / \|u_i\|^2$, then normalize each u_j. This is the same subtraction as Gram–Schmidt because $e_i = u_i / \|u_i\|$.
2. On $[-1, 1]$ for $1, x, x^2$, symmetry gives $u_1 = 1$, $u_2 = x$, $u_3 = x^2 - 1/3$, with squared norms $2, 2/3, 8/45$.
3. Normalizing gives $1/\sqrt{2}$, $\sqrt{3/2}\, x$, $\sqrt{5/8}(3x^2 - 1)$. Their first spans agree with the monomial initial spans.

Proposition (LADR 6.36: extension)

Every orthonormal list in a finite-dimensional inner-product space extends to an orthonormal basis.

Where the proof comes from

Want. Retain a given orthonormal list while filling the remaining dimensions.

Know. It is independent and thus extends to an ordinary basis.

Key step. Apply Gram–Schmidt in that order; the initial orthonormal vectors are unchanged.

Proof. 1. Extend the independent list to a basis by Week 2.

2. Apply Proposition 87. On each given initial vector, all preceding coefficients are zero and the norm is 1, so it remains unchanged; the full output is the desired basis.

□

Proposition (LADR 5.44: splitting and triangular form)

An operator on finite-dimensional V has an upper-triangular matrix in some basis if and only if its minimal polynomial splits into linear factors over F. The factors may repeat; the zero-space polynomial is the empty product 1.

Where the proof comes from

Want. Identify exactly when real triangularization is possible.

Know. The minimal polynomial divides annihilators, and a split nonconstant polynomial has a root in the field.

Key step. In one direction use the invariant initial spans; in the other induct on a proper shifted range.

Proof. 1. Suppose the diagonal entries in a triangular basis are $\lambda_1, \ldots, \lambda_n$, and let U_j be its initial spans. The map $T - \lambda_j I$ sends U_j into U_{j-1}, with $U_0 = 0$.

2. Applying these factors from $j = n$ down to 1 sends all of V to zero. They commute, so $\prod_j (T - \lambda_j I) = 0$. Its polynomial is split; divisibility shows m_T also splits.

3. Conversely induct on dimension, with the empty basis at dimension 0. For nonzero V a root λ of m_T is an eigenvalue (Week 7); hence $U = \operatorname{range}(T - \lambda I)$ is proper and invariant.

4. The restriction's minimal polynomial divides m_T, so splits. Inductively triangulate $T|_U$ and extend that basis to V. For any added v_j, $Tv_j = (T - \lambda I)v_j + \lambda v_j \in U + Fv_j$, giving triangular form by invariant initial spans.

□

Proposition (LADR 6.37–6.38: orthonormal triangular form and Schur's theorem)

If m_T splits over F, an orthonormal basis can be chosen with upper-triangular matrix for T. In particular this holds for every complex finite-dimensional operator.

Where the proof comes from

Want. Retain triangular form while making the basis orthonormal.

Know. Triangularity is invariance of initial spans, and Gram–Schmidt preserves those spans.

Key step. Apply Gram–Schmidt to an existing triangular basis.

Proof. 1. The preceding result supplies a triangular basis when m_T splits. Its initial spans are invariant by Week 7.
2. Proposition 87 gives an orthonormal basis with the same initial spans, so its matrix is still upper triangular.
3. Over $\mathbb{C}$, m_T splits by the fundamental theorem of algebra from Week 6, so the criterion always applies.

□

Orthogonal polynomial context from the notes. Repeating Gram–Schmidt on the monomials with the $[0,1]$ integral product produces the shifted Legendre family; $[-1,1]$ gives the usual Legendre family. A Gaussian weight on the real line gives the Hermite family. The family identifications and their differential-equation properties are cited classical facts here; the first three Legendre calculations above are explicitly proved.

Example (Notes Chapter 18: no orthogonal Hamel basis in ℓ^2)

Let ℓ^2 consist of sequences a with $\sum_j |a_j|^2 < \infty$, and use $\langle a, b\rangle = \sum_j a_j \overline{b_j}$. A Hamel basis means a vector-space basis with only finite sums allowed. This space has no orthogonal Hamel basis.

1. The product series converges absolutely by applying finite Cauchy–Schwarz to tails and taking limits. The standard coordinate sequences are independent, so ℓ^2 has infinite dimension. We cite the analytic completeness of ℓ^2: every Cauchy sequence in its norm converges there.
2. Suppose an orthogonal Hamel basis existed. Normalize its nonzero vectors to unit length and choose a countably infinite sublist $e_1, e_2, \ldots$. The partial sums $s_N = \sum_{j=1}^{N} 2^{-j} e_j$ are Cauchy, because Proposition 84 gives $\|s_M - s_N\|^2 = \sum_{j=N+1}^{M} 4^{-j} \to 0$. Completeness supplies a limit $v \in \ell^2$.
3. For fixed j, Cauchy–Schwarz gives $|\langle v - s_N, e_j\rangle| \le \|v - s_N\| \to 0$, so $\langle v, e_j\rangle = 2^{-j} \neq 0$.
4. Any finite expansion in the entire orthonormal family has zero inner product with all but finitely many members of this sublist. Thus it cannot equal v, contradicting the Hamel-basis property. The argument applies to a sublist of any cardinal-sized basis.

Shortcuts

- For polynomial Gram–Schmidt, use parity on symmetric intervals before integrating; odd products have zero integral.
- Delay square roots by forming unnormalized orthogonal remainders, then normalize once at the end.
- Gram–Schmidt preserves invariant initial spans, so it preserves upper-triangular form.

Traps

- Dividing by a zero remainder is invalid. For the dependent input $v_1 = (1,0), v_2 = (2,0)$, the second remainder is zero; remove dependent inputs before applying the independent-list theorem.
- Schur form is triangular, and need not be diagonal: $J_2(1)$ has no eigenbasis, even though its standard basis is orthonormal and its matrix triangular.

§3 Functionals and orthogonal complements

The question. How can an inner product represent a linear functional? On $\mathbb{C}$, $\varphi(z) = iz$ equals $\langle z, -i\rangle$, so conjugating the representative's coefficient is essential.

Proposition (Notes Proposition 88; LADR 6.42: Riesz representation)
On finite-dimensional V, every $\varphi \in V'$ has a unique $r \in V$ with $\varphi(u) = \langle u, r\rangle$. The map $\Gamma : V \to V'$, $\Gamma(r)(u) = \langle u, r\rangle$, is a real-linear isomorphism (regarding complex spaces as real). Over $\mathbb{C}$ it is conjugate-linear, $\Gamma(ar) = \overline{a}\Gamma(r)$.

Where the proof comes from

Want. Identify each functional with one vector without choosing coordinates permanently.
Know. A vector orthogonal to itself is zero; the dual has the same dimension.
Key step. Prove injectivity by evaluating at the proposed representative, then use equal real dimensions.

Proof. 1. First-position linearity makes $\Gamma(r)$ a linear functional. Second-position conjugate linearity gives $\Gamma(ar+bs) = \overline{a}\Gamma(r) + \overline{b}\Gamma(s)$, hence real linearity.
2. If $\Gamma(r) = 0$, evaluate it at r: $0 = \langle r, r\rangle$ implies $r = 0$. Thus Γ is injective.
3. Over $\mathbb{R}$, V and V' have the same dimension. Over $\mathbb{C}$, both underlying real dimensions are $2\dim_{\mathbb{C}} V$. Rank-nullity makes the real-linear injection surjective and proves uniqueness.

□

Proposition (Riesz computing formula; LADR 6.43)
In an orthonormal basis, the representative is $r = \sum_j \overline{\varphi(e_j)} e_j$.

Where the proof comes from

Want. Compute the Riesz vector explicitly.
Know. Testing a functional on basis vectors determines it, and the second position conjugates coefficients.
Key step. Solve $\langle e_j, r\rangle = \varphi(e_j)$ one coefficient at a time.

Proof. 1. If $r = \sum_j a_j e_j$, then $\langle e_k, r\rangle = \overline{a_k}$. Thus $a_k = \overline{\varphi(e_k)}$.
2. For $u = \sum_k c_k e_k$, first-position linearity gives $\langle u, r\rangle = \sum_k c_k \varphi(e_k) = \varphi(u)$, verifying the formula.

□

Compute coordinate and polynomial representatives (LADR 6.40–6.41, 6.44)

1. For $\varphi(z_1, z_2, z_3) = 2z_1 - 5z_2 + z_3$, the standard orthonormal basis gives $r = (2, -5, 1)$. If instead $\varphi(z) = iz_1$, its representative is $(-i, 0, 0)$, because the formula conjugates i.
2. For $\varphi(p) = \int_{-1}^{1} p(x)\cos(\pi x)\,dx$ on $\mathcal{P}_2(\mathbb{R})$, the monomial values are $0, 0, -4/\pi^2$, by integration by parts and parity.
3. Seek $r = a + bx + cx^2$. The equations $\langle 1, r\rangle = 0$, $\langle x, r\rangle = 0$, $\langle x^2, r\rangle = -4/\pi^2$ are $2a + 2c/3 = 0$, $2b/3 = 0$, $2a/3 + 2c/5 = -4/\pi^2$. Solving gives $r = 15(1 - 3x^2)/(2\pi^2)$.
4. On $\mathcal{P}_5(\mathbb{R})$, the same functional from LADR 6.41 has an even representative of degree 4. Solving the even-monomial equations gives
$$r(x) = \frac{105(27 - 2\pi^2)}{8\pi^4} + \frac{315(4\pi^2 - 45)}{4\pi^4}x^2 + \frac{1575(21 - 2\pi^2)}{8\pi^4}x^4.$$
Its products with the odd monomials vanish by parity; the three even products equal the corresponding integrals against $\cos(\pi x)$, verifying all six basis tests.

Definition (Notes Definition 47; LADR 6.46)**.** For any subset $S \subseteq V$, define $S^{\perp} = \{v : \langle v, s\rangle = 0 \text{ for every } s \in S\}$. By conjugate symmetry the reversed inner products also vanish. For a subspace U, this is its orthogonal complement.

Proposition (Notes Definition 47; LADR 6.48: complement properties)

$S^\perp$ is a subspace and equals $(\operatorname{span} S)^\perp$. Also $\{0\}^\perp = V$, $V^\perp = \{0\}$; a subspace U has $U \cap U^\perp = \{0\}$; and $S \subseteq T$ implies $T^\perp \subseteq S^\perp$.

Where the proof comes from

Want. Reduce orthogonal-complement calculations to basis tests.

Know. Inner products are linear in one position and conjugate-linear in the other.

Key step. Zero equations are preserved under linear combinations.

Proof. 1. If v, w are orthogonal to every s, then $\langle av + bw, s\rangle = a0 + b0 = 0$; zero also satisfies the equations. Thus $S^\perp$ is a subspace.

2. If v is orthogonal to each $s \in S$, conjugate linearity makes it orthogonal to every finite combination of the s. This proves the span equality.

3. Zero is orthogonal to all vectors; a vector orthogonal to all of V, or one in $U \cap U^\perp$, is orthogonal to itself and therefore zero.

4. More vectors in T impose at least all the equations imposed by $S \subseteq T$, proving the reversed inclusion.

□

Proposition (Notes Proposition 89; LADR 6.49, 6.51)

For a finite-dimensional subspace U of any inner-product space V, $V = U \oplus U^\perp$. If V is finite-dimensional, $\dim U^\perp = \dim V - \dim U$.

Where the proof comes from

Want. Decompose every vector into a subspace component and an orthogonal remainder.

Know. U has an orthonormal basis and $U \cap U^\perp = 0$.

Key step. Subtract the coefficients in that orthonormal basis.

Proof. 1. Choose an orthonormal basis $e_1, \dots, e_m$ of U, explicitly using Proposition 87. For $v \in V$ put $u = \sum_j \langle v, e_j\rangle e_j$.

2. For every k, $\langle v - u, e_k\rangle = 0$, so $v - u \in U^\perp$ by the span property. Thus $v = u + (v - u)$ is in the sum.

3. The zero intersection proves directness and uniqueness. When V is finite-dimensional, the direct-sum dimension formula gives the stated dimension.

□

Proposition (Notes Proposition 90; LADR 6.52, 6.54)

For finite-dimensional U, $(U^\perp)^\perp = U$. In particular $U^\perp = \{0\}$ exactly when $U = V$. In finite-dimensional V, for subspaces U, W, $(U + W)^\perp = U^\perp \cap W^\perp$ and $(U \cap W)^\perp = U^\perp + W^\perp$.

Where the proof comes from

Want. Reverse orthogonal-complement operations and simplify sums/intersections.

Know. Every vector has the preceding direct decomposition.

Key step. Its unwanted orthogonal component must be orthogonal to itself.

Proof. 1. Conjugate symmetry gives $U \subseteq (U^\perp)^\perp$. If v belongs to the latter, decompose $v = u+w$ with $u \in U, w \in U^\perp$. Then $w = v - u$ also belongs to $(U^\perp)^\perp$, so $\langle w, w\rangle = 0$ and $w = 0$.

2. This proves equality. If $U^\perp = 0$, take complements to get $U = \{0\}^\perp = V$; the converse follows from $V^\perp = 0$.
3. A vector is orthogonal to every $u + w$ exactly when it is orthogonal separately to U and W (set one summand zero, then use conjugate linearity). This proves the sum identity.
4. Apply the sum identity to $U^\perp, W^\perp$ and then take complements. In finite-dimensional V every subspace involved is finite-dimensional, so the double-complement equalities give the intersection identity.

□

Compute a complement by equations (LADR 6.47)

1. In $\mathbb{R}^3$, the complement of the point set $\{(2,3,5)\}$ is the plane $2x + 3y + 5z = 0$, by its defining dot product. The plane's complement is the line $\mathbb{R}(2,3,5)$: the vector belongs to it, and the dimension formula gives dimension 1.
2. In F^5, the coordinate plane $U = \{(a,b,0,0,0)\}$ has $U^\perp = \operatorname{span}(e_3, e_4, e_5)$, since orthogonality to e_1, e_2 sets the first two coordinates to zero.
3. If an orthonormal basis is $e_1, \ldots, e_m, f_1, \ldots, f_k$, the same coefficient test gives $\operatorname{span}(e_1, \ldots, e_m)^\perp = \operatorname{span}(f_1, \ldots, f_k)$.

Use the kernel complement to find a representative (LADR 6.58)

1. For nonzero φ, choose $w \neq 0$ in $(\ker \varphi)^\perp$, possible because the finite-dimensional kernel is proper and has a complementary nonzero space. Then $\varphi(w) \neq 0$, since the kernel meets its complement only at zero.
2. For any u, the vector $u - \varphi(u)w/\varphi(w)$ is in the kernel and orthogonal to w. Thus $\langle u, w\rangle = \varphi(u)\|w\|^2/\varphi(w)$.
3. Therefore $r = \overline{\varphi(w)}w/\|w\|^2$ satisfies $\langle u, r\rangle = \varphi(u)$; for $\varphi = 0$ use $r = 0$. For example $\varphi(x,y) = 2x - y$ has kernel $\mathbb{R}(1,2)$, choose $w = (2,-1)$, and obtain $r = 5w/5 = (2,-1)$.

Annihilators and complements. In finite dimension, Γ from Proposition 88 identifies $U^\perp$ with the annihilator U^0 from Week 5: $\Gamma(v)(u) = \langle u, v\rangle = 0$ for all $u \in U$ exactly when $v \in U^\perp$. This is a real-linear identification over $\mathbb{C}$, with the same underlying conjugate-linear convention as Γ.

Shortcuts

- Test orthogonality against a basis of the subspace; linearity extends the zero equations to every vector.
- Use a full orthonormal basis extending one for U; its remaining vectors give a basis of $U^\perp$.
- In a complex Riesz calculation conjugate the functional's values on an orthonormal basis.

Traps

- Complex Γ is not complex-linear: $\Gamma(ir) = -i\Gamma(r)$. For $\varphi(z) = iz$, the representative is $-i$, not i.
- Complement reverses sums: for the two coordinate lines in $\mathbb{R}^2$, $(U \cap W)^\perp = \mathbb{R}^2$, whereas $U^\perp \cap W^\perp = \{0\}$.

§4 Orthogonal projections and best approximation

The question. Which vector in a subspace is closest to a given vector? For $v = (2,3)$ and $U = \mathbb{R}(1,0)$, the closest point is $(2,0)$, with orthogonal error $(0,3)$.

Definition (Notes Definition 48; LADR 6.55)**.** For finite-dimensional U, write the unique decomposition $v = u + w$ with $u \in U, w \in U^\perp$. The orthogonal projection onto U is $P_U v = u$. A projector is a linear operator P with $P^2 = P$.

Proposition (Notes Proposition 91)

Every projector satisfies $V = \ker P \oplus \operatorname{range} P$. It is the projection onto its range along its kernel.

Where the proof comes from

Want. Relate idempotence to a direct-sum decomposition.
Know. $P^2 = P$ makes P the identity on its range.
Key step. Split v as $(v - Pv) + Pv$.

Proof. 1. $P(v - Pv) = Pv - P^2v = 0$, so $v - Pv \in \ker P$ and $Pv \in \operatorname{range} P$. Thus the two subspaces sum to V.
2. If $w = Pv \in \ker P$, then $w = P^2v = P(Pv) = Pw = 0$, so the intersection is zero. On its range P is identity; on its kernel it is zero, identifying the projection.

□

Proposition (Notes Proposition 92; LADR 6.57: orthogonal projection properties)
The map P_U is linear, has range U and kernel $U^\perp$, fixes U, and is zero on $U^\perp$. Also $P_U^2 = P_U$, $v - P_Uv \in U^\perp$, and $\|P_Uv\| \le \|v\|$. For an orthonormal basis $e_1, \dots, e_m$ of U, $P_Uv = \sum_j \langle v, e_j\rangle e_j$.

Where the proof comes from

Plan. Use the direct decomposition for the map properties, coefficients for the formula, and Pythagoras for the norm bound.
Want. Compute the projection and verify all its defining features.
Know. $V = U \oplus U^\perp$ uniquely, with an orthonormal basis of the finite-dimensional summand.
Key step. Addition and scaling respect the two subspaces, while the orthogonal remainder has nonnegative squared norm.

Proof. 1. For $v_i = u_i + w_i$ and scalars a, b, $av_1 + bv_2 = (au_1 + bu_2) + (aw_1 + bw_2)$ is the unique decomposition. Thus $P_U(av_1 + bv_2) = aP_Uv_1 + bP_Uv_2$.
2. The decompositions $u = u + 0$ and $w = 0 + w$ show that the map fixes U and is zero on $U^\perp$, giving range, kernel and idempotence. The definition gives $v - P_Uv \in U^\perp$.
3. Since $\langle v, e_j\rangle = \langle P_Uv, e_j\rangle$, Proposition 86 on U gives the stated coefficient formula.
4. Pythagoras gives $\|v\|^2 = \|P_Uv\|^2 + \|v - P_Uv\|^2 \ge \|P_Uv\|^2$, proving the norm bound.

□

Example (Notes Chapter 18; LADR 6.56) 1. For $U = Fu$ with $u \neq 0$, normalize $e = u/\|u\|$ in Proposition 92 to get $P_Uv = \langle v, u\rangle u/\|u\|^2$.
2. On $\mathbb{R}^2$, the notes' $P(x, y) = (0, y)$ is the orthogonal projection onto $\mathbb{R}e_2$, with kernel $\mathbb{R}e_1$. In contrast $N(x, y) = (y, 0)$ has $N^2 = 0$ and is not a projector since $N \neq 0$.

Project a polynomial (Notes Example 71)

1. On $[0, 1]$ project x^3 onto $U = \operatorname{span}(1, x)$ using the orthonormal basis $1, \sqrt{3}(2x - 1)$ from Example 70.
2. The two coefficients are $1/4$ and $3\sqrt{3}/20$, so $P_Ux^3 = 1/4 + (3\sqrt{3}/20)\sqrt{3}(2x - 1) = 9x/10 - 1/5$.
3. Direct integration gives $\langle x^3 - (9x/10 - 1/5), 1\rangle = 0$ and $\langle x^3 - (9x/10 - 1/5), x\rangle = 0$, confirming the remainder is orthogonal to U. It is therefore also orthogonal to the projected polynomial.

Project using any basis, without orthonormalizing

1. For a basis $u_1, \dots, u_m$ of U, solve $\sum_j c_j\langle u_j, u_i\rangle = \langle v, u_i\rangle$ for each i. The equations say $v - \sum_j c_ju_j \in U^\perp$, so their solution gives $P_Uv = \sum_j c_ju_j$.
2. The coefficient matrix is invertible: a solution of the homogeneous system would give a vector in $U \cap U^\perp = 0$, whose basis coefficients are all zero. Finite-dimensional invertibility then gives a unique solution.
3. For the preceding example with $u_1 = 1, u_2 = x$, the system is $c_1 + c_2/2 = 1/4$, $c_1/2 + c_2/3 = 1/5$. It gives $(c_1, c_2) = (-1/5, 9/10)$, matching the orthonormal calculation.


Proposition (LADR 6.61: best approximation)
For finite-dimensional U, $\|v - P_U v\| \le \|v - u\|$ for all $u \in U$, with equality exactly when $u = P_U v$.

Where the proof comes from

Want. Prove a global minimum and its uniqueness.
Know. The residual $v - P_U v$ is orthogonal to U.
Key step. Decompose the error at any other candidate into this residual plus a vector in U.

Proof. 1. Write $v - u = (v - P_U v) + (P_U v - u)$; the two terms are orthogonal, because the first is in $U^\perp$ and the second in U.
2. Pythagoras gives $\|v - u\|^2 = \|v - P_U v\|^2 + \|P_U v - u\|^2$. The last term is nonnegative and is zero exactly at $u = P_U v$, proving both claims.

□




The error from v to its projection is orthogonal to the subspace. Any different candidate in U adds a nonnegative squared error.

Best degree-5 sine approximation (LADR 6.63)

1. In $C[-\pi, \pi]$ with $\langle f, g\rangle = \int_{-\pi}^{\pi} fg$, project $\sin x$ onto $\mathcal{P}_5(\mathbb{R})$. Odd and even polynomials are orthogonal by parity, and $\sin x$ is orthogonal to every even polynomial, so the projection has the form $c_1 x + c_3 x^3 + c_5 x^5$.
2. For $r, s \in \{1, 3, 5\}$ the coefficient matrix entries are $G_{rs} = 2\pi^{r+s+1}/(r + s + 1)$. Integration by parts gives the right side $b = (2\pi, 2\pi(\pi^2 - 6), 2\pi(\pi^4 - 20\pi^2 + 120))^{\mathsf{T}}$. Solve $Gc = b$ as in the preceding method.
3. The exact solution is

$$c_1 = \frac{105(\pi^4 - 153\pi^2 + 1485)}{8\pi^6},$$
$$c_3 = -\frac{315(\pi^4 - 125\pi^2 + 1155)}{4\pi^8},$$
$$c_5 = \frac{693(\pi^4 - 105\pi^2 + 945)}{8\pi^{10}}.$$

4. Thus the best polynomial is approximately $0.987862136x - 0.155271411x^3 + 0.005643118x^5$. It minimizes the squared integral error by LADR 6.61; its value is about 0.000116169.
5. The degree-5 Taylor polynomial $x - x^3/6 + x^5/120$ has squared integral error about 0.118702 on this interval. At $x = 3$ the absolute errors are respectively about 0.001416 and 0.383880, illustrating the difference between approximation over an interval and matching derivatives at a point.



The best polynomial approximates sine over the entire interval in squared integral error. The Taylor polynomial is designed by derivatives at zero; its endpoint errors are larger.

Shortcuts

- For a line, divide by $\|u\|^2$ when the spanning vector is not normalized.
- A candidate projection is verified by membership in U and an orthogonal residual; uniqueness follows from the direct sum.
- A general basis gives a square system of inner-product equations, avoiding square roots from orthonormalization.

Traps

- A projector need not be orthogonal: $P(x, y) = (x + y, 0)$ has $P^2 = P$ but range $\mathbb{R}e_1$ and kernel $\mathbb{R}(-1, 1)$ are not orthogonal. It sends $(1, 1)$ to $(2, 0)$, increasing norm from $\sqrt{2}$ to 2.
- Omitting the denominator for a nonunit line vector gives a wrong projection: for $u = (2, 0)$ and $v = (1, 1)$, $\langle v, u\rangle u = (4, 0)$, whereas dividing by $\|u\|^2 = 4$ gives $(1, 0)$.

§5 Pseudoinverse and least-squares solutions

The question. How can one select a best approximate solution when a linear system is inconsistent? For $T(a, b, c, d) = (a + b + c, 2c + d, 0)$, a target (x, y, z) with $z \neq 0$ cannot be attained, but $(x, y, 0)$ can.

Proposition (LADR 6.67)

For finite-dimensional V and $T \in L(V, W)$ between inner-product spaces, the restriction $T|_{(\ker T)^\perp} : (\ker T)^\perp \to \operatorname{range} T$ is an isomorphism.

Where the proof comes from

Want. Obtain an invertible map from an arbitrary linear map.
Know. The kernel has an orthogonal complement and their direct sum is V.
Key step. Remove the kernel from the domain without losing any range values.

Proof. 1. A vector in the restriction's kernel belongs to both $\ker T$ and its orthogonal complement, so is zero. Thus the restriction is injective.

2. If $w = Tv$, decompose $v = k + u$ with $k \in \ker T$ and $u \in (\ker T)^\perp$. Then $Tu = Tv = w$, proving surjectivity onto $\operatorname{range} T$.

□

Definition (LADR 6.68)**.** For such T, its pseudoinverse is the linear map $T^\dagger : W \to V$ given by $T^\dagger = (T|_{(\ker T)^\perp})^{-1} P_{\operatorname{range} T}$. The projection exists because $\operatorname{range} T$ is finite-dimensional.

Proposition (LADR 6.69)

If T is invertible, $T^\dagger = T^{-1}$. In general $TT^\dagger = P_{\operatorname{range} T}$ and $T^\dagger T = P_{(\ker T)^\perp}$. Consequently surjective T has $TT^\dagger = I$, and injective T has $T^\dagger T = I$.

Where the proof comes from

Want. Compare the pseudoinverse with a genuine inverse.
Know. Its definition first projects to the range, then uses the invertible restriction.
Key step. Evaluate separately on each orthogonal direct-sum component.

Proof. 1. Applying T after its restricted inverse gives $TT^\dagger w = P_{\operatorname{range} T} w$.
2. For $v = k + u$ with $k \in \ker T$ and $u \in (\ker T)^\perp$, $Tv = Tu \in \operatorname{range} T$. The restricted inverse gives $T^\dagger T v = u = P_{(\ker T)^\perp} v$.
3. If T is invertible, its kernel is zero and its range is W, so the definition reduces to T^{-1}. In the surjective or injective cases the corresponding projection is the identity, proving the final claims.

□

Proposition (LADR 6.70)

For $b \in W$, the vector $x_0 = T^\dagger b$ minimizes $\|Tx - b\|$. Every minimizer is $x_0 + k$, $k \in \ker T$. Among these minimizers, x_0 is the unique one with least norm.

Where the proof comes from

Want. Solve both least-error and least-norm selection problems.
Know. $Tx_0 = P_{\operatorname{range} T} b$ and $x_0 \perp \ker T$.
Key step. Use Pythagoras once for the residual in W, then again for kernel changes in V.

Proof. 1. Decompose $Tx - b = (Tx - P_{\operatorname{range} T} b) + (P_{\operatorname{range} T} b - b)$. The first term lies in $\operatorname{range} T$, the second in its orthogonal complement.
2. Pythagoras gives $\|Tx - b\|^2 = \|Tx - P_{\operatorname{range} T} b\|^2 + \|P_{\operatorname{range} T} b - b\|^2$. The minimum occurs exactly when $Tx = Tx_0$, equivalently $x - x_0 \in \ker T$.
3. For $x = x_0 + k$ with $k \in \ker T$, orthogonality gives $\|x\|^2 = \|x_0\|^2 + \|k\|^2$, minimized uniquely at $k = 0$.

□

Compute the pseudoinverse by equations (LADR 6.71)

1. For $T : F^4 \to F^3$, $T(a,b,c,d) = (a+b+c, 2c+d, 0)$, the range is the first coordinate plane. Its projection sends (x,y,z) to $(x,y,0)$. Solving $T(a,b,c,d) = 0$ gives the kernel basis $(-1,1,0,0), (-1,0,1,-2)$.
2. The least-norm preimage of $(x,y,0)$ satisfies the image equations $a+b+c = x$, $2c+d = y$, and the two orthogonality equations $-a+b = 0$, $-a+c-2d = 0$.
3. Solving gives
$$T^\dagger(x,y,z) = \frac{1}{11}(5x-2y,\ 5x-2y,\ x+4y,\ -2x+3y).$$
4. For example at $(1,2,3)$ this is $(1,1,9,4)/11$, whose image is $(1,2,0)$ and residual norm is 3. Every other minimizer adds a vector in the displayed kernel, increasing squared norm by that vector's squared norm.

Shortcuts

- To compute $T^\dagger b$, project b to the range, solve the image equation, and impose orthogonality to a kernel basis.
- All least-squares solutions differ by a kernel vector, because their images equal the unique best range approximation.

Traps

- A pseudoinverse is defined even when an inverse is not: the displayed T is neither injective nor surjective.
- $TT^\dagger$ need not be identity: in this example it sends $(0,0,1)$ to zero, because that vector lies outside the range.

§6 Problem set

The question. Construct orthonormal bases, solve complement equations and verify projections by orthogonal errors. For a line $F(1,0)$ in F^2, the complement is $F(0,1)$ and the projection sends (a,b) to $(a,0)$.

HW10 P1. On $\mathcal{P}_2(\mathbb{R})$ with $\langle p,q\rangle = \int_{-1}^{1} pq$, apply Gram–Schmidt to $1, x, x^2$ to obtain an orthonormal basis.
Hint: Use Proposition 87 or its unnormalized version, with parity.

HW10 P2. In standard $\mathbb{R}^4$, let $W = \operatorname{span}((1,-1,1,0),(0,2,1,0))$. Find a basis of $W^\perp$.
Hint: Use the basis-test equations from Definition 47.

HW10 P3. On $\mathcal{P}_2(\mathbb{R})$ with the same product, let $W = \operatorname{span}(x+1)$. Find a basis of $W^\perp$.
Hint: Set $\langle a+bx+cx^2, x+1\rangle = 0$, then use the complement dimension formula.

HW10 P4. Prove $(U \cap W)^\perp = U^\perp + W^\perp$ for subspaces of finite-dimensional V.
Hint: Apply the sum-complement identity and Proposition 90 twice.

HW10 P5. For an orthonormal basis $e_1, \dots, e_m$ of U, prove (a) $P_U v = \sum_j \langle v, e_j\rangle e_j$; (b) $\|v - P_U v\| \le \|v - u\|$ for every $u \in U$, and determine equality.
Hint: Use the direct decomposition and Pythagoras as in Proposition 92 and LADR 6.61.

HW10 P6 (optional). On $\mathcal{P}_1(\mathbb{R})$ with $\langle f,g\rangle = \int_{-1}^{1} fg$, find the unique p such that $\int_0^2 f(x)\,dx = \langle f,p\rangle$ for all $f \in \mathcal{P}_1(\mathbb{R})$.
Hint: Use Riesz representation, testing $1, x$.

HW10 P7 (optional). For a linear projector P on finite-dimensional V, prove it is an orthogonal projection exactly when $\|Pv\| \le \|v\|$ for all v.
Hint: In Proposition 91's decomposition, test $v = u + tw$ with $u \in \operatorname{range} P$, $w \in \ker P$, choosing t to make any nonzero mixed inner product contradict the norm bound.

Additional practice (LADR 6A Exercise 11). Find $u, v \in \mathbb{R}^2$ with $u \in \mathbb{R}(1,3)$, $v \perp (1,3)$ and $(1,2) = u + v$.
Hint: Apply the line-projection formula, then subtract.

Additional practice (LADR 6A Exercise 6). Prove $\langle u, v \rangle = 0$ if and only if $\|u\| \leq \|u + av\|$ for every $a \in F$.
Hint: Expand the squared norm; if $v \neq 0$, choose $a = -\langle u, v \rangle / \|v\|^2$.