

Class 8: ML Mini Project

Aparajita Pranjal

Exploratory Data Analysis

The .csv file was downloaded and read in as a new variable.

```
# Save your input data file into your Project directory
fna.data <- "WisconsinCancer.csv"

# Complete the following code to input the data and store as wisc.df
wisc.df <- read.csv(fna.data, row.names=1)
# We can use -1 here to remove the first column
wisc.data <- wisc.df[,-1]
# Create diagnosis vector for later
diagnosis <- factor(wisc.df[,1])
```

- **Q1.** How many observations are in this dataset? **569**

```
dim(wisc.data)
```

```
[1] 569  30
```

- **Q2.** How many of the observations have a malignant diagnosis? **212**

```
table(diagnosis, exclude = "B")
```

```
diagnosis
```

```
  M
```

```
212
```

- **Q3.** How many variables/features in the data are suffixed with `_mean`? **10 variables**

```
col_names <- colnames(wisc.data)
length(grep("_mean", col_names))
```

```
[1] 10
```

PCA Analysis

```
# Check column means and standard deviations
colMeans(wisc.data)
```

radius_mean	texture_mean	perimeter_mean
1.412729e+01	1.928965e+01	9.196903e+01
area_mean	smoothness_mean	compactness_mean
6.548891e+02	9.636028e-02	1.043410e-01
concavity_mean	concave.points_mean	symmetry_mean
8.879932e-02	4.891915e-02	1.811619e-01
fractal_dimension_mean	radius_se	texture_se
6.279761e-02	4.051721e-01	1.216853e+00
perimeter_se	area_se	smoothness_se
2.866059e+00	4.033708e+01	7.040979e-03
compactness_se	concavity_se	concave.points_se
2.547814e-02	3.189372e-02	1.179614e-02
symmetry_se	fractal_dimension_se	radius_worst
2.054230e-02	3.794904e-03	1.626919e+01
texture_worst	perimeter_worst	area_worst
2.567722e+01	1.072612e+02	8.805831e+02
smoothness_worst	compactness_worst	concavity_worst
1.323686e-01	2.542650e-01	2.721885e-01
concave.points_worst	symmetry_worst	fractal_dimension_worst
1.146062e-01	2.900756e-01	8.394582e-02

```
apply(wisc.data,2,sd)
```

radius_mean	texture_mean	perimeter_mean
3.524049e+00	4.301036e+00	2.429898e+01
area_mean	smoothness_mean	compactness_mean
3.519141e+02	1.406413e-02	5.281276e-02
concavity_mean	concave.points_mean	symmetry_mean

7.971981e-02	3.880284e-02	2.741428e-02
fractal_dimension_mean	radius_se	texture_se
7.060363e-03	2.773127e-01	5.516484e-01
perimeter_se	area_se	smoothness_se
2.021855e+00	4.549101e+01	3.002518e-03
compactness_se	concavity_se	concave.points_se
1.790818e-02	3.018606e-02	6.170285e-03
symmetry_se	fractal_dimension_se	radius_worst
8.266372e-03	2.646071e-03	4.833242e+00
texture_worst	perimeter_worst	area_worst
6.146258e+00	3.360254e+01	5.693570e+02
smoothness_worst	compactness_worst	concavity_worst
2.283243e-02	1.573365e-01	2.086243e-01
concave.points_worst	symmetry_worst	fractal_dimension_worst
6.573234e-02	6.186747e-02	1.806127e-02

```
# Perform PCA on wisc.data by completing the following code
wisc.pr <- prcomp(wisc.data, scale = TRUE)

# Look at summary of results
summary(wisc.pr)
```

Importance of components:


	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					

Cumulative Proportion 1.00000 1.00000

- **Q4.** From your results, what proportion of the original variance is captured by the first principal components (PC1)? **0.4427**
- **Q5.** How many principal components (PCs) are required to describe at least 70% of the original variance in the data? **3 components**
- **Q6.** How many principal components (PCs) are required to describe at least 90% of the original variance in the data? **7 components**

Interpreting PCA Results


```
biplot(wisc.pr)
```



- **Q7.** What stands out to you about this plot? Is it easy or difficult to understand? Why?


The function plots all the points on the graph and labels them which makes it difficult to read and understand the different groups/trends.

```
plot(wisc.pr$x , col = diagnosis,  
      xlab = "PC1", ylab = "PC2")
```



- **Q8.** Generate a similar plot for principal components 1 and 3. What do you notice about these plots? **They have a larger range on the y-axis for PCA3 as the data points are a bit more spread out.**

```
# Repeat for components 1 and 3
plot(wisc.pr$x[,1],wisc.pr$x[,3], col = diagnosis,
     xlab = "PC1", ylab = "PC3")
```




Graphing with ggplot

```
# Create a data.frame for ggplot
df <- as.data.frame(wisc.pr$x)
df$diagnosis <- diagnosis

# Load the ggplot2 package
library(ggplot2)

# Make a scatter plot colored by diagnosis
ggplot(df) +
  aes(PC1, PC2, col= diagnosis) +
  geom_point()
```




Variance Explained

```
# Calculate variance of each component
pr.var <- wisc.pr$sdev^2
head(pr.var)
```

```
[1] 13.281608  5.691355  2.817949  1.980640  1.648731  1.207357
```


```
# Variance explained by each principal component: pve
pve <- pr.var / sum(pr.var)

# Plot variance explained for each principal component
plot(pve, xlab = "Principal Component",
     ylab = "Proportion of Variance Explained",
     ylim = c(0, 1), type = "o")
```



Scree Plot

```
# Alternative scree plot of the same data, note data driven y-axis
barplot(pve, ylab = "Precent of Variance Explained",
        names.arg=paste0("PC",1:length(pve)), las=2, axes = FALSE)
axis(2, at=pve, labels=round(pve,2)*100 )
```

- **Q9.** For the first principal component, what is the component of the loading vector (i.e. `wisc.pr$rotation[,1]`) for the feature `concave.points_mean`? This tells us how much this original feature contributes to the first PC. **-0.26085376**

```
wisc.pr$rotation[,1]
```

radius_mean	texture_mean	perimeter_mean
-0.21890244	-0.10372458	-0.22753729
area_mean	smoothness_mean	compactness_mean
-0.22099499	-0.14258969	-0.23928535
concavity_mean	concave.points_mean	symmetry_mean
-0.25840048	-0.26085376	-0.13816696
fractal_dimension_mean	radius_se	texture_se
-0.06436335	-0.20597878	-0.01742803
perimeter_se	area_se	smoothness_se
-0.21132592	-0.20286964	-0.01453145
compactness_se	concavity_se	concave.points_se
-0.17039345	-0.15358979	-0.18341740
symmetry_se	fractal_dimension_se	radius_worst
-0.04249842	-0.10256832	-0.22799663
texture_worst	perimeter_worst	area_worst
-0.10446933	-0.23663968	-0.22487053

smoothness_worst	compactness_worst	concavity_worst
-0.12795256	-0.21009588	-0.22876753
concave.points_worst	symmetry_worst	fractal_dimension_worst
-0.25088597	-0.12290456	-0.13178394

Hierarchical Clustering

```
# Scale the wisc.data data using the "scale()" function
data.scaled <- scale(wisc.data)
```


```
data.dist <- dist(data.scaled)
```

```
wisc.hclust <- hclust(data.dist, method = "complete")
```

- **Q10.** Using the `plot()` and `abline()` functions, what is the height at which the clustering model has 4 clusters? **Height is 19**

```
plot(wisc.hclust)
abline(h = 19, col="red", lty=2, )
```

Cluster Dendrogram



```
data.dist
hclust (*, "complete")
```

```
wisc.hclust.clusters <- cutree(wisc.hclust, h = 19)
table(wisc.hclust.clusters, diagnosis)
```


	diagnosis		
wisc.hclust.clusters	B	M	
1	12	165	
2	2	5	
3	343	40	
4	0	2	

- **Q12.** Which method gives your favorite results for the same `data.dist` dataset? Explain your reasoning.

Ward.D2 gives the best results for the same dataset as it makes visualizing the differences between clusters much easier as it squares the differences whereas complete takes in all the data points and creates a large dendrogram with tighter clusters as we go further down the tree whereas ward.D@ distinguishes them higher up.

Clustering PCA Results

```
wisc.pr.hclust <- hclust(data.dist, method = "ward.D2")
plot(wisc.pr.hclust)
```




```
grps <- cutree(wisc.pr.hclust, k=2)
table(grps)
```

```
grps
  1  2
184 385
```


```
table(grps, diagnosis)
```

```
      diagnosis
grps    B    M
  1    20 164
  2   337  48
```

```
plot(wisc.pr$x[,1:2], col=grps)
```



```
plot(wisc.pr$x[,1:2], col=diagnosis)
```




```
g <- as.factor(grps)
levels(g)
```

```
[1] "1" "2"
```

```
g <- relevel(g,2)
levels(g)
```

```
[1] "2" "1"
```

```
# Plot using our re-ordered factor
plot(wisc.pr$x[,1:2], col=g)
```



New Plot

```
#install.packages("rgl")
#library(rgl)
#plot3d(wisc.pr$x[,1:3], xlab="PC 1", ylab="PC 2", zlab="PC 3", cex=1.5, size=1, type="s",
```

Clustering Continued

```
## Use the distance along the first 7 PCs for clustering i.e. wisc.pr$x[, 1:7]
wisc.pr.hclust <- hclust(dist(wisc.pr$x[,1:7]), method="ward.D2")

wisc.pr.hclust.clusters <- cutree(wisc.pr.hclust, k=2)

table(wisc.pr.hclust.clusters , diagnosis)
```

```
          diagnosis
wisc.pr.hclust.clusters  B  M
1      28 188
2     329  24
```

```
table(wisc.hclust.clusters, diagnosis)
```

```

              diagnosis
wisc.hclust.clusters  B  M
              1 12 165
              2  2   5
              3 343  40
              4  0   2

```

- **Q13.** How well does the newly created model with four clusters separate out the two diagnoses? **The new model further sub-categorizes the two diagnoses providing a more detailed breakdown of benign versus malignant.**
- **Q14.** How well do the hierarchical clustering models you created in previous sections (i.e. before PCA) do in terms of separating the diagnoses? Again, use the `table()` function to compare the output of each model (`wisc.km$cluster` and `wisc.hclust.clusters`) with the vector containing the actual diagnoses. **Before PCA the clustering does not provide information on outliers or potential false readings which after PCA there is a difference if a few individuals that was separated which is more useful in obtaining specific details from the dataset and not just looking it as it overall.**

Prediction

```

#url <- "new_samples.csv"
url <- "https://tinyurl.com/new-samples-CSV"
new <- read.csv(url)
npc <- predict(wisc.pr, newdata=new)
npc

```


```

      PC1      PC2      PC3      PC4      PC5      PC6      PC7
[1,]  2.576616 -3.135913  1.3990492 -0.7631950  2.781648 -0.8150185 -0.3959098
[2,] -4.754928 -3.009033 -0.1660946 -0.6052952 -1.140698 -1.2189945  0.8193031
      PC8      PC9      PC10     PC11     PC12     PC13     PC14
[1,] -0.2307350 0.1029569 -0.9272861 0.3411457  0.375921 0.1610764 1.187882
[2,] -0.3307423 0.5281896 -0.4855301 0.7173233 -1.185917 0.5893856 0.303029
      PC15     PC16     PC17     PC18     PC19     PC20
[1,] 0.3216974 -0.1743616 -0.07875393 -0.11207028 -0.08802955 -0.2495216
[2,] 0.1299153  0.1448061 -0.40509706  0.06565549  0.25591230 -0.4289500
      PC21     PC22     PC23     PC24     PC25     PC26
[1,] 0.1228233 0.09358453 0.08347651 0.1223396 0.02124121 0.078884581
[2,] -0.1224776 0.01732146 0.06316631 -0.2338618 -0.20755948 -0.009833238
      PC27      PC28      PC29      PC30

```

```
[1,] 0.220199544 -0.02946023 -0.015620933 0.005269029
[2,] -0.001134152 0.09638361 0.002795349 -0.019015820
```

```
plot(wisc.pr$x[,1:2], col=g)
points(npc[,1], npc[,2], col="blue", pch=16, cex=3)
text(npc[,1], npc[,2], c(1,2), col="white")
```



- **Q16.** Which of these new patients should we prioritize for follow up based on your results? **Group 2 as they belong to the malignant category and should therefore be brought in for further testing before the benign patients.**