Transparent Screening for LLM Inference and Training Impacts

Arnault Pachot[1]* and Thierry Petit[1]

Emotia, Paris, France
{apachot,tpetit}@emotia.com

Abstract. This paper presents the transparent screening framework used in ImpactLLM to estimate inference and training impacts for current large language models. It documents the algorithms that convert natural-language application descriptions into bounded environmental estimates, and it also reports the comparative tables and visual results published in the observatory. The objective is not to claim direct measurement for opaque proprietary services, but to provide an auditable, source-linked proxy framework that improves comparability and public discussion under conditions of limited observability.

Keywords: Large language models · environmental impact · screening method · transparent estimation

1 Introduction

This paper documents the transparent screening framework used in the *ImpactLLM Observatory*. The observatory now covers 41 models, and this manuscript explains how the interface extracts scenarios from natural-language descriptions, propagates them through bounded multi-factor screening proxies, and publishes the resulting comparative tables and figures. The goal is to keep those proxies auditable while keeping the application workflow fast and interpretable.

The remainder of this paper covers the technical design of the inference estimator, the training proxy, the comparative outputs currently exposed in the observatory, and the limits of those assumptions.

The methodological premise is straightforward. For the dominant hosted LLM services, direct provider-side environmental telemetry is generally unavailable, even though those systems dominate practical use and public debate. In that context, avoiding approximation altogether does not produce a more rigorous discussion; it leaves room for opaque claims, unsupported comparisons, and contradictory numerical narratives. The role of the present paper is therefore not to claim direct measurement where none exists, but to define a bounded, source-linked, inspectable proxy framework that is methodologically preferable to unverifiable assertions.

* Corresponding author: Arnault Pachot, apachot@emotia.com

2 Inference Screening Method

The main practical question addressed by the tool is straightforward: given a software feature using an LLM, can we produce an inference estimate that is useful for screening and comparison when direct provider telemetry is absent?

The current answer is deliberately limited. The estimator is *inference-only*. It excludes model training, embodied impacts, application-side software consumption, and ancillary infrastructure from the displayed result. This is not because those dimensions are unimportant, but because the available inference anchors are structured enough to support a transparent screening method, whereas mixing them with broader lifecycle terms would blur the meaning of the result.

The design objective is also practical speed. Instead of asking users to fill a large technical form, the web interface starts from a natural-language description such as "We use GPT-4o-mini for customer support, around 4,000 uses per month in France." A parser then maps this text to a compact scenario with model, request type, approximate tokens, usage volume, and country assumptions. The output is therefore fast to obtain, but every inferred parameter remains visible to the user and can be inspected or challenged.

The current market-model release follows five rules.

- It starts from an observed literature anchor rather than from provider claims. The retained prompt-energy anchor is the Gemini Apps median prompt energy reported by Elsworth et al.: 0.24 Wh/prompt [7].
- It makes token volume explicit. Prompt compute is approximated from input and output tokens, with a larger weight for output generation than for input processing.
- It does not scale on raw parameters alone. It constructs *effective active parameters* by adjusting the target profile with context-window class, serving mode, modality support, and architecture notes.
- It reports a bounded low-central-high interval rather than a single falsely precise point.
- It derives carbon from retained energy and the electricity mix of the retained country, which is a contextual assumption rather than a model measurement.

Formally, let $E_a = 0.24$ Wh be the anchor value and $P_a = 180$ the active-parameter proxy retained for the anchor model. For a target model t, let P_t denote active parameters, T_{in} input tokens, and T_{out} output tokens. We define weighted prompt-compute volume:

$$V_t = T_{in} + \omega T_{out}, \tag{1}$$

with $\omega = 1.8$ in the current release. The project reference scenario is:

$$V_{ref} = 1000 + 1.8 \times 550 = 1990. \tag{2}$$

Effective active parameters are then defined by:

$$P_{t,s}^{eff} = P_t \times F_{t,s}^{ctx} \times F_{t,s}^{srv} \times F_{t,s}^{mod} \times F_{t,s}^{arch}, \tag{3}$$

where $s \in \{\text{low}, \text{central}, \text{high}\}$ indexes the screening scenario. The final per-request energy estimate is:

$$\hat{E}_{t,s} = E_a \times \left(\frac{P_{t,s}^{eff}}{P_a}\right)^{\alpha_s} \times \left(\frac{V_t}{V_{ref}}\right)^{\beta_s}. \tag{4}$$

The current implementation uses $\alpha_{low} = 0.85$, $\alpha_{central} = 0.95$, $\alpha_{high} = 1.05$, and $\beta_{low} = 0.85$, $\beta_{central} = 0.92$, $\beta_{high} = 1.0$. Carbon is then derived from the retained country mix:

$$\hat{C}_{t,s,c} = \frac{\hat{E}_{t,s}}{1000} \times CI_c, \tag{5}$$

where CI_c is the carbon intensity of country c in gCO_2e/kWh.

This method is not presented as a physical law of inference scaling. It is a traceable screening proxy anchored in observed prompt energy and explicit contextual assumptions. Its purpose is to provide a fast and inspectable estimate for comparative reasoning, not an audited declaration of real provider-side energy use.

3 Training Screening Method

The observatory also exposes training orders of magnitude for current market models. Here the uncertainty is even higher than for inference because published values are sparse, heterogeneous, and often reported only as aggregate emissions. A pure parameter-only scaling is therefore too brittle. The current release instead uses a second bounded proxy that combines retained parameter count with additional training priors.

The training proxy starts from literature anchors that directly report training CO_2e, and from training-energy reconstructions derived from those emissions when the source-country electricity mix is documented. For a target model t, the central training estimate is:

$$\hat{E}_{t,s}^{train} = E_a^{train} \times \left(\frac{P_t}{P_a}\right)^{\alpha_s} \times \left(\frac{Tok_t}{Tok_a}\right)^{\beta_s} \times F_{t,s}^{reg} \times F_{t,s}^{arch-tr} \times F_{t,s}^{hw}, \tag{6}$$

where P_t is the retained parameter count, Tok_t is a training-token prior, F^{reg} is a training-regime prior, $F^{arch-tr}$ captures architecture and multimodality assumptions, and F^{hw} is a hardware-class proxy. The current release uses a simple prior of 20 training tokens per retained parameter when no better public estimate is available, and defaults market models to a foundation-pretraining regime unless a narrower public indication exists.

This second proxy is not used in the application estimator shown to users, which remains inference-only. It is used in the observatory and the comparative tables to avoid suggesting that training impacts can be projected from parameter count alone. The result should still be read as a screening order of magnitude rather than as an audited declaration of model-development impact.

4 Results and Comparative Outputs

The underlying dataset and derived tables were refreshed for the March 2026 release, so the published values mirror the recalculated inference and training columns for the expanded market-model catalog. For inference, the observatory now emphasizes the central prompt-level proxy per standardized request before any application-specific annualization.

Table 1 summarizes a few illustrative outputs. The first block shows how the observatory makes model orders of magnitude legible at the request level. The second block shows how the same proxy can be annualized at the software-feature level from compact, natural-language scenarios.

Case	Energy	Carbon	Interpretation
Ministral 3B, one standardized request	0.0056 Wh	0.0002 gCO_2e	Small hybrid model under a French provider-country proxy
GPT-5 mini, one standardized request	0.1706 Wh	0.0657 gCO_2e	Medium proprietary hosted model under a US provider-country proxy
Claude Opus 4.1, one standardized request	2.9922 Wh	1.1520 gCO_2e	Very large proprietary estimate illustrating the steep growth of screening orders of magnitude
Support chatbot, Ministral 8B, 20,000 conversations/month	2.38 kWh/year	96 gCO_2e/year	Low-carbon annual result despite high usage because the retained electricity factor is favorable
Retrieval assistant, GPT-5 mini, 4,000 uses/month	12.31 kWh/year	4.74 $kgCO_2e$/year	Higher annualized result driven by hosted-service assumptions and US electricity contextualization

Table 1. Illustrative outputs from the current observatory and application estimator. These values are screening estimates, not audited declarations.

These examples illustrate three practical properties of the method. First, annualization matters: apparently small unit values become relevant when multiplied by real usage. Second, the chosen model profile matters materially. Third, the retained electricity mix can dominate the carbon interpretation, which is

why country contextualization is explicitly shown in the interface. Just as importantly, these results can be obtained quickly from natural-language inputs while keeping the extracted scenario and assumptions visible in the interface.

For completeness, Table 2 reports the current calculated values for all 40 market models tracked in the March 2026 release. The purpose of this table is documentary: it makes the observatory outputs directly inspectable in the paper, while the web interface remains the primary medium for interactive exploration.

Table 2: Calculated environmental indicators for the 40 market models currently tracked by ImpactLLM. Inference values correspond to the central prompt-level proxy retained by the observatory, expressed per standardized request. Training energy values are multi-factor screening estimates, and training CO_2e is contextualized from retained training energy using the model-country electricity mix.

Model	Inference Wh/request	Inference gCO_2e/request	Training GWh	Training $ktCO_2e$
Claude 2	0.1738	0.0669	0.31	0.12
Claude 3.5 Haiku	0.0412	0.0159	0.02	0.01
Claude 3.5 Sonnet	0.2957	0.1139	0.96	0.37
Claude 3.7 Sonnet	0.2957	0.1139	0.96	0.37
Claude Opus 4.1	2.9922	1.1520	125.63	48.37
Claude Sonnet 4	0.6486	0.2497	5.03	1.93
Codestral 22B	0.0347	0.0014	0.01	0.00
DeepSeek R1	0.0836	0.0451	12.30	6.64
DeepSeek V3	0.0798	0.0431	12.30	6.64
Gemini 2.0 Flash	0.0594	0.0229	0.03	0.01
Gemini 2.5 Flash	0.1508	0.0581	0.20	0.08
Gemini 2.5 Flash-Lite	0.0594	0.0229	0.03	0.01
Gemini 2.5 Pro	0.3601	0.1387	1.26	0.48
GLaM 130B	0.1845	0.0710	0.64	0.25
Gopher 280B	0.4136	0.0744	2.29	0.41
GPT-3.5 Turbo	0.2646	0.1019	0.98	0.38
GPT-4	0.7524	0.2897	14.93	5.75
GPT-4o mini	0.0155	0.0060	0.00	0.00
GPT-5 mini	0.1706	0.0657	0.28	0.11
GPT-5 nano	0.1706	0.0657	0.28	0.11
GPT-5.2	2.7897	1.0740	101.76	39.18
GPT-5.2-pro	2.7897	1.0740	101.76	39.18
GPT-OSS 120B	0.0116	0.0045	0.39	0.15
GPT-OSS 20B	0.0074	0.0029	0.01	0.00
Grok 1	0.1590	0.0612	2.79	1.07
Grok 2	0.2389	0.0920	2.06	0.79

Model	Inference Wh/request	Inference gCO_2e/request	Training GWh	Training $ktCO_2e$
Grok 4	1.0994	0.4233	11.31	4.35
Jurassic-1 Jumbo	0.2689	0.1035	1.42	0.55
LaMDA 1	0.2097	0.0807	0.56	0.22
Llama 3.1 405B	0.5530	0.2129	5.23	2.01
Llama 3.1 70B	0.1043	0.0402	0.16	0.06
Llama 3.1 8B	0.0133	0.0051	0.00	0.00
Megatron-Turing NLG 530B	0.7583	0.2919	14.48	5.57
Ministral 3B	0.0056	0.0002	0.00	0.00
Ministral 8B	0.0142	0.0006	0.00	0.00
Mistral Large	0.2074	0.0083	0.48	0.02
Mistral Small 3.1 24B	0.0413	0.0017	0.02	0.00
OPT 175B	0.2337	0.0900	1.12	0.43
Qwen2.5 32B	0.0496	0.0268	0.03	0.02
Qwen2.5 72B	0.1072	0.0579	0.17	0.09
Qwen2.5 7B	0.0117	0.0063	0.00	0.00

5 Literature and Source Base

The results above are not produced from a single benchmark, but from a layered source base combining direct environmental anchors, estimator literature, infrastructure and electricity context, and broader analytical framing. This structure corresponds to the source logic exposed in the site's *Sources* page, where each value is linked to a document, a system boundary, and a use role in the estimator.

5.1 Direct Environmental Anchors for LLMs

The primary literature anchors for training and inference come from work that reports environmental indicators directly for language-model systems or adjacent production systems. The core training anchors include Strubell et al. on transformer training costs [26], Luccioni et al. on BLOOM [14], Morrison et al. on holistic language-model impacts [18], and recent lifecycle or training-oriented extensions such as Fernandez et al. [8] and d'Orgeval et al. [5]. For inference, the most operational prompt-level anchor in the current release is Elsworth et al. [7], complemented by broader framing from Ren et al. [23] and water-related contextualization from Li et al. [13].

5.2 Measurement, Tracking, and Proxy Literature

The methodological design also builds on the literature and tooling ecosystem for measuring or estimating AI impacts when direct telemetry is available or partially missing. This includes CarbonTracker [2], CodeCarbon [17],

the ML.ENERGY benchmark [3], and proxy-oriented tooling such as EcoLogits [9]. These references do not provide one transferable number for all current hosted LLMs, but they define the design space within which a screening proxy must remain explicit about assumptions, units, and uncertainty.

5.3 Infrastructure, Electricity, and Cloud Context

A second family of sources informs the infrastructural and contextual layers needed to convert energy proxies into carbon estimates or to interpret datacenter scale effects. This includes cloud and infrastructure accounting references such as Cloud Carbon Footprint [4], AWS CCFT [1], the Microsoft Emissions Impact Dashboard [16], Google Cloud carbon reporting [10], and the OVHcloud environmental tracker [19]. It also includes system-level or electricity-demand framing from EPRI [6], Lawrence Berkeley National Laboratory [12,25], the International Energy Agency [11], and Joule-scale discussion by de Vries-Gao [27].

5.4 Model Documentation and Conceptual Framing

Some source families play a different role: they do not provide direct environmental measurements but document model properties or frame the interpretation of the results. Examples include model and provider documentation such as the Llama 3.1 model card [15], broader environmental framing from Rillig et al. [24], and earlier conceptual work on sustainable AI and environmental decision support [21,20]. The observatory and the site itself are part of this transparency layer because the project republishes those linked records in reusable interfaces [22].

6 Discussion and Perspectives

Beyond the catalog table and the static release timelines, the observatory can also be read as a dynamic screening signal about the acceleration of frontier-model impacts. Figure 1 summarizes that intuition with a simple log-linear reading of the GPT, Claude, and Grok flagship series for retained training impacts. The purpose is not to claim a physical law or a universal growth constant. It is to provide a compact interpretation of the retained central estimates under the current observatory assumptions.

Two lessons follow from this perspective view. First, training remains the dimension where screening values grow fastest. This is consistent with the current screening structure: training combines retained parameter count, token priors, architecture effects, and hardware assumptions, so frontier-model scaling quickly reaches very high orders of magnitude. Inference, by contrast, is represented in this paper at the per-request level, where comparisons remain useful but less visually dominant than the corresponding training orders of magnitude.

Second, these visuals are useful precisely because they remain interpretive rather than declarative. They help communicate that the observatory does not



Fig. 1. Illustrative reading of the observatory's central training-screening estimates for flagship GPT, Claude, and Grok models. The visual condenses the current retained values into an approximate doubling-time interpretation.

only contain isolated point estimates; it also suggests a market trajectory in which the environmental stakes of frontier models may be rising rapidly. But the value of that reading depends entirely on the transparency of the underlying assumptions. If those assumptions change, the apparent doubling times change as well. The figure should therefore be read as a discussion tool and policy prompt, not as a direct provider-side measurement or immutable forecasting law.

From a policy and governance perspective, this is arguably one of the most useful roles an open observatory can play. Even without direct industrial telemetry, it can surface the possibility that model-scale growth is outpacing the public visibility of environmental reporting. In that sense, the discussion value of the observatory is not limited to ranking models; it also lies in making structural trends legible enough to motivate better disclosure, more granular reporting, and more disciplined debate.

7 Limitations

The main limitations are methodological. First, the estimator depends on scarce literature anchors, so the uncertainty intervals rely on wide parameter- and token-exponent bounds to keep the screening range honest. Second, proprietary service configurations remain opaque, forcing the method to rely on contextual proxies for regime, architecture, and hardware. Third, training-level emissions are reconstructed from sparse reports and assumed token priors; more published training anchors would reduce that uncertainty. Finally, the natural-language parser must interpret ambiguous descriptions, which is why every extracted assumption is surfaced to the user along with the final estimate.

8 Implementation Notes

The prototype is implemented as a Python/Flask web application that builds on the observatory dataset stored in `ImpactLLM/data/market_models.csv`. The front-end sends the user description to OpenAI for parsing, then queries the estimator to compute the inference and training proxies. The same dataset drives the observatory tables, which remain independent of the estimator logic.

9 Conclusion

The contribution of this manuscript is both methodological and documentary: it explains how ImpactLLM transforms natural-language descriptions into bounded inference and training estimates, and it makes the resulting comparative tables and figures for the tracked models directly inspectable in one place.

That contribution should not be read as an attempt to eliminate uncertainty. For the most widely used proprietary LLM services, uncertainty is structurally unavoidable because the decisive telemetry remains inaccessible. The relevant methodological task is therefore to organize approximation rather than to pretend to escape it. This paper argues that a transparent, source-linked, reproducible screening proxy is a better basis for comparison and discussion than the current situation in which strong environmental claims often circulate without explicit assumptions, traceable derivation, or shared comparison rules.

Acknowledgments. The authors thank the ImpactLLM project contributors and reviewers of earlier internal drafts for helping clarify the presentation of the screening framework.

Disclosure of Interests. The authors have no competing interests to declare that are relevant to the content of this article.

References

1. Amazon Web Services: What is the customer carbon footprint tool? (2026), https://docs.aws.amazon.com/awsaccountbilling/latest/aboutv2/what-is-ccft.html, aWS documentation, accessed March 2026
2. Anthony, L., Kanding, B., Selvan, R., Christensen, E., Andersson, O., et al.: Carbontracker: Tracking and predicting the carbon footprint of training deep learning models. arXiv preprint arXiv:2007.03051 (2020), https://arxiv.org/abs/2007.03051
3. Chung, J.W., Ma, J.J., Wu, R., Liu, J., Kweon, O.J., Xia, Y., Wu, Z., Chowdhury, M.: The ml.energy benchmark: Toward automated inference energy measurement and optimization. arXiv preprint arXiv:2505.06371 (2025). https://doi.org/10.48550/arXiv.2505.06371
4. Cloud Carbon Footprint: Cloud carbon footprint methodology (2026), https://www.cloudcarbonfootprint.org/docs/methodology/, open-source software documentation, accessed March 2026

5. d'Orgeval, T., Azaïs, C., Michaux, J.L., Perrin, G., Trévisan, V., et al.: Life cycle assessment of data centres for generative artificial intelligence. Applied Energy **392**, 126617 (2026). https://doi.org/10.1016/j.apenergy.2025.126617
6. Electric Power Research Institute: Power demand for data centers and artificial intelligence (2024), https://www.publicpower.org/periodical/article/epri-report-examines-power-demand-data-centers-artificial-intelligence
7. Elsworth, C., Huang, K., Patterson, D., Schneider, I., et al.: Measuring the environmental impact of delivering ai at google scale. arXiv preprint arXiv:2508.15734 (2025), https://arxiv.org/abs/2508.15734
8. Fernandez, C., Pérez-Lombard, L., Ruiz, G., Gutiérrez, E., et al.: Life cycle assessment of large language models: A methodological proposal. Tackling Climate Change with Machine Learning (2025), https://www.climatechange.ai/papers/iclr2025/24
9. GenAI Impact: Ecologits documentation and methodology (2026), https://ecologits.ai/latest/, open-source software documentation, accessed March 2026
10. Google Cloud: View carbon emissions reports (2026), https://docs.cloud.google.com/carbon-footprint/docs/view-carbon-data, product documentation, accessed March 2026
11. International Energy Agency: Energy and ai (2025), https://www.iea.org/reports/energy-and-ai/energy-demand-from-ai
12. Lawrence Berkeley National Laboratory: Report evaluates increase in electricity demand from data centers (2025), https://newscenter.lbl.gov/2025/01/15/berkeley-lab-report-evaluates-increase-in-electricity-demand-from-data-centers/
13. Li, P., Yang, J., Islam, M.A., Ren, S.: Making ai less "thirsty": Uncovering and addressing the secret water footprint of ai models. Communications of the ACM **68**(4), 46–54 (2025), https://cacm.acm.org/research/making-ai-less-thirsty/
14. Luccioni, A.S., Viguier, S., Ligozat, A.L.: Estimating the carbon footprint of BLOOM, a 176b parameter language model. Journal of Machine Learning Research **24**(253), 1–15 (2023), https://www.jmlr.org/papers/v24/23-0069.html
15. Meta: Llama 3.1 model card (2024), https://huggingface.co/meta-llama/Llama-3.1-405B
16. Microsoft: Emissions impact dashboard (2026), https://www.microsoft.com/en/sustainability/emissions-impact-dashboard, product documentation, accessed March 2026
17. MLCO2 Project: Codecarbon documentation and methodology (2026), https://mlco2.github.io/codecarbon/, open-source software documentation, accessed March 2026
18. Morrison, J., Na, C., Fernandez, J., Dettmers, T., et al.: Holistically evaluating the environmental impact of creating language models. arXiv preprint arXiv:2503.05804 (2025), https://arxiv.org/abs/2503.05804
19. OVHcloud: Ovhcloud launches environmental impact tracker (2024), https://corporate.ovhcloud.com/es/newsroom/news/environmental-impact-tracker/, product announcement
20. Pachot, A., Patissier, C.: Toward sustainable artificial intelligence: An overview of environmental protection uses and issues. Green and Low-Carbon Economy **3**(2), 105–112 (2023), https://ojs.bonviewpress.com/index.php/GLCE/article/view/608
21. Pachot, A., Patissier, C., Studio, O.: Intelligence artificielle et environnement : alliance ou nuisance ? L'IA face aux défis écologiques d'aujourd'hui et de demain. Dunod (2022), https://www.dunod.com/entreprise-et-economie/intelligence-artificielle-et-environnement-alliance-ou-nuisance-ia-face-aux

22. Pachot, A., Petit, T.: Impactllm (2026), https://github.com/apachot/ImpactLLM, gitHub repository and online demo: https://dev.emotia.com/impact-llm
23. Ren, S., Tomlinson, B., Black, R.W., Torrance, A.W.: Reconciling the contrasting narratives on the environmental impact of large language models. Scientific Reports **14**, 28180 (2024). https://doi.org/10.1038/s41598-024-76682-6
24. Rillig, M.C., Ågerstrand, M., Bi, M., Gould, K.A., Sauerland, U.: Risks and benefits of large language models for the environment. Environmental Science & Technology **57**(9), 3464–3466 (2023). https://doi.org/10.1021/acs.est.3c01106
25. Shehabi, A., Newkirk, A., Smith, S.J., Hubbard, A., Lei, N., et al.: 2024 united states data center energy usage report. Tech. rep., Lawrence Berkeley National Laboratory (2024), https://escholarship.org/uc/item/32d6m0d1
26. Strubell, E., Ganesh, A., McCallum, A.: Energy and policy considerations for deep learning in nlp. In: Proceedings of the 57th Annual Meeting of the Association for Computational Linguistics. pp. 3645–3650 (2019). https://doi.org/10.18653/v1/P19-1355
27. de Vries-Gao, A.: Artificial intelligence: Supply chain constraints and energy implications. Joule **9**(6), 1153–1156 (2025). https://doi.org/10.1016/j.joule.2025.101961