

Triple-Higgs production as a probe of the Higgs potential

John M. Campbell¹, Giuseppe De Laurentis², R. Keith Ellis³, Michelle Gensmann^{4,5} and Ulrich Haisch^{4*}

¹ Fermilab, PO Box 500, Batavia IL 60510-5011, USA

² Higgs Centre for Theoretical Physics, University of Edinburgh, Edinburgh, EH9 3FD, UK

³ Institute for Particle Physics Phenomenology, Durham University, Durham, DH1 3LE, UK

⁴ Max Planck Institute for Physics, Boltzmannstr. 8, 85748 Garching, Germany

⁵ Institute for Theoretical Physics, Karlsruhe Institute of Technology, Wolfgang-Gaede-Str. 1, 76131 Karlsruhe, Germany.

★ haisch@mpp.mpg.de

Preprint: FERMILAB-PUB-26-0464-T, LHCHWG-2026-xxx, MPP-2026-xxx

Abstract

We study triple-Higgs production in the κ -framework as a direct probe of the Higgs potential. We derive compact parametrisations of the inclusive gluon-fusion production cross section that capture the dependence on the Higgs trilinear and quartic self-couplings together with anomalous two-top-two-Higgs and two-top-three-Higgs interactions. These results provide a practical framework for interpreting current and future LHC measurements. We use the parametrisations to assess the sensitivity of LHC Run 2 and HL-LHC data to Higgs self-couplings and show that differential kinematic distributions can provide additional discrimination between scenarios with nearly identical inclusive production rates.

Copyright attribution to authors.

This work is a submission to SciPost Physics.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

Contents

1	Introduction	2
2	Triple-Higgs production in the κ-framework	3
2.1	Including the two-top–three-Higgs interaction	8
3	Conclusions	12
	References	13

1 Introduction

The discovery of the Higgs boson in 2012 [1, 2] completed the particle content of the Standard Model (SM) and established the mechanism of electroweak (EW) symmetry breaking. Since then, a major objective of the LHC physics programme has been to determine the properties of the Higgs boson with increasing precision. While its couplings to EW gauge bosons and third-generation fermions are already constrained at the few- to ten-percent level, its self-interactions remain largely unexplored. Measuring the Higgs self-couplings is therefore one of the most important outstanding goals of collider physics, since they directly probe the shape of the Higgs potential and the dynamics of EW symmetry breaking.

In the SM, the Higgs potential is determined by a single quartic coupling $\lambda = m_h^2/(2v^2)$. After EW symmetry breaking, it predicts unique values for the Higgs trilinear and quartic self-couplings,

$$\lambda_3^{\text{SM}} = \lambda, \quad \lambda_4^{\text{SM}} = \frac{\lambda}{4}, \quad (1)$$

once the Higgs boson mass $m_h \simeq 125 \text{ GeV}$ and vacuum expectation value $v \simeq 246 \text{ GeV}$ are fixed. Any deviation from these predictions would provide evidence for physics beyond the SM, such as extended scalar sectors, composite Higgs models, or mechanisms for EW baryogenesis. Reviews of such beyond-the-SM (BSM) scenarios can be found, for example, in Refs. [3, 4].

At the LHC, double-Higgs production provides the primary direct probe of the Higgs trilinear self-coupling, while loop-induced effects in single-Higgs processes yield complementary constraints [5–13]. These measurements are typically interpreted in the κ -framework [14–16], which parametrises deviations from the SM through

$$\kappa_3 = \frac{\lambda_3}{\lambda_3^{\text{SM}}}, \quad (2)$$

assuming all other couplings take their SM values. Using the full $\sqrt{s} = 13 \text{ TeV}$ LHC Run 2 data set, the ATLAS and CMS collaborations obtain the combined constraint

$$-0.71 < \kappa_3 < 6.1 \quad (\text{ATLAS} + \text{CMS}) \quad (3)$$

at the 95% confidence level (CL) [17]. The HL-LHC, with an integrated luminosity of 6 ab^{-1} , is expected to improve this sensitivity to

$$0.5 < \kappa_3 < 1.7 \quad (\text{HL-LHC}). \quad (4)$$

Constraining the quartic Higgs self-coupling is considerably more challenging. It is conventionally parametrised as

$$\kappa_4 = \frac{\lambda_4}{\lambda_4^{\text{SM}}}, \quad (5)$$

and triple-Higgs production provides its only direct probe at hadron colliders. However, the tiny SM cross section of approximately 100 ab at $\sqrt{s} = 14$ TeV, calculated at next-to-next-to-leading order (NNLO) in QCD [18], renders such measurements experimentally very challenging. Current ATLAS [19] and CMS [20] searches yield

$$-230 < \kappa_4 < 240 \quad (\text{ATLAS}), \quad -190 < \kappa_4 < 190 \quad (\text{CMS}), \quad (6)$$

at the 95% CL, assuming all other couplings are SM-like, in particular $\kappa_3 = 1$. A recent HL-LHC study [21] projects these bounds to improve to

$$-81 < \kappa_4 < 89 \quad (\text{HL-LHC}) \quad (7)$$

with an integrated luminosity of 6 ab^{-1} . In addition to triple-Higgs production, double-Higgs production is also sensitive to κ_4 through two-loop EW corrections [21–24], although these constraints are generally weaker.

In this note, we present a comprehensive study of triple-Higgs production within the κ -framework. We derive compact parametrisations of the inclusive gluon-fusion production cross section that capture its dependence on the Higgs trilinear and quartic self-couplings, the top-quark Yukawa coupling modifier κ_t , and the **two-top-two-Higgs and two-top-three-Higgs interactions** κ_{2t} and κ_{3t} . These expressions provide practical tools for interpreting current and future LHC measurements. Using them, we investigate the sensitivity of current and projected LHC data to the Higgs self-couplings and study the impact of correlated modifications of the **top-Higgs interactions**. We further demonstrate that differential kinematic distributions retain significant sensitivity even in scenarios with nearly degenerate inclusive cross sections, highlighting the importance of shape information in future triple-Higgs searches.

2 Triple-Higgs production in the κ -framework

Triple-Higgs production at hadron colliders has been the subject of extensive theoretical investigation over the past two decades [18, 21, 22, 25–41]. Within the SM, the gluon-fusion process $gg \rightarrow 3h$ is dominated by the one-loop topologies shown in the upper and middle rows of Figure 1. In the figure, green (blue) boxes denote insertions of the Higgs trilinear (quartic) self-coupling, while orange circles represent the top-quark Yukawa coupling. The two diagrams in the lower row illustrate representative BSM contributions involving a contact interaction between two top quarks and two Higgs bosons. In the SM, bottom-quark loops **modify** the $gg \rightarrow 3h$ production cross section **by only about 0.2% at LHC energies [39]; they are nevertheless retained in the numerical calculations below.**

The κ -formalism Lagrangian, which parametrises the interaction vertices highlighted in Figure 1, takes the form

$$\mathcal{L}_\kappa \supset -\kappa_3 v \lambda h^3 - \kappa_4 \frac{\lambda}{4} h^4 - \kappa_t \frac{y_t}{\sqrt{2}} \bar{t} t h - \kappa_{2t} \frac{y_t}{\sqrt{2}v} \bar{t} t h^2 - \kappa_{3t} \frac{y_t}{2\sqrt{2}v^2} \bar{t} t h^3, \quad (8)$$

where κ_3 and κ_4 were introduced in Eqs. (2) and (5), respectively, and $y_t = \sqrt{2}m_t/v$, with m_t denoting the top-quark mass. The parameter κ_t thus encodes modifications of the top-quark Yukawa coupling, while κ_{2t} and κ_{3t} **parametrise contact interactions** between two top




Figure 1: Representative one-loop Feynman diagrams contributing to triple-Higgs production via gluon fusion, illustrating the triangle, box, and pentagon topologies. Green (blue) boxes denote the Higgs trilinear (quartic) self-coupling, while orange (magenta) circles indicate top-quark couplings involving one (two) Higgs bosons.

quarks and two or three Higgs bosons, respectively. The final term can equivalently be written as $-\kappa_{3t}m_t\bar{t}th^3/(2v^3)$ [35]. The SM is recovered from Eq. (8) for $\kappa_3 = \kappa_4 = \kappa_t = 1$ and $\kappa_{2t} = \kappa_{3t} = 0$.

To determine the dependence of the inclusive $gg \rightarrow 3h$ production cross section on κ_3 , κ_4 , κ_t , and κ_{2t} , we employ the leading-order (LO) matrix elements calculated in Ref. [39] and implemented in the MCFM [42–45] parton-level event generator. We use $m_h = 125$ GeV and $m_t = 173$ GeV as input parameters together with the PDF4LHC21 parton distribution functions (PDFs) [46]. The renormalisation and factorisation scales, μ_R and μ_F , are chosen dynamically as $\mu_R = \mu_F = m_{3h}/2$, where m_{3h} denotes the invariant mass of the three-Higgs system. For collision energies of $\sqrt{s} = 13$ TeV, $\sqrt{s} = 13.6$ TeV, and $\sqrt{s} = 14$ TeV, we parametrise the LO cross section as

$$\sigma_{\text{LO}}^{\sqrt{s}} = (\sigma_{\text{LO}}^{\sqrt{s}})_{\text{SM}} \sum_{n,m} c_{nm}^{\sqrt{s}}(\kappa_t, \kappa_{2t}) (\Delta\kappa_3)^n (\Delta\kappa_4)^m, \quad (9)$$

where $\Delta\kappa_i = \kappa_i - 1$ for $i = 3, 4$. The corresponding LO SM cross sections are

$$\sigma_{\text{LO}}^{13\text{TeV}} = 42.1 \text{ ab}, \quad \sigma_{\text{LO}}^{13.6\text{TeV}} = 47.9 \text{ ab}, \quad \sigma_{\text{LO}}^{14\text{TeV}} = 51.0 \text{ ab}. \quad (10)$$




Figure 2: Constraints in the κ_3 - κ_4 plane at LHC Run 2 (left) and the HL-LHC (right). The red contours indicate the preferred 95% CL regions obtained from triple-Higgs production in gluon fusion for three representative values of κ_t , as specified by the contour labels. We assume $\kappa_{2t} = 0$. The SM prediction is denoted by black points. The yellow regions, enclosed by black dotted contours, represent the perturbative unitarity bounds derived from tree-level $hh \rightarrow hh$ scattering.

For $\sqrt{s} = 13$ TeV, the coefficients $c_{nm}^{\sqrt{s}}(\kappa_t, \kappa_{2t})$ appearing in Eq. (9) are

$$c_{00}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = -6.05 \cdot 10^{-4} + 0.0384\kappa_t^2 - 0.350\kappa_t^3 + 1.47\kappa_t^4 - 3.04\kappa_t^5 + 2.89\kappa_t^6 \\ + \kappa_{2t} (1.61\kappa_t^2 - 7.82\kappa_t^3 + 15.6\kappa_t^4 - 14.8\kappa_t^5) \\ + \kappa_{2t}^2 (6.96 - 22.3\kappa_t + 37.0\kappa_t^2), \quad (11)$$

$$c_{01}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0829\kappa_t^2 - 0.264\kappa_t^3 + 0.0895\kappa_t^4 + \kappa_{2t} (0.673\kappa_t - 1.12\kappa_t^2), \quad (12)$$

$$c_{02}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0169\kappa_t^2, \quad (13)$$

$$c_{10}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.211\kappa_t^2 - 1.45\kappa_t^3 + 3.65\kappa_t^4 - 3.28\kappa_t^5 \\ + \kappa_{2t} (3.49\kappa_t - 13.4\kappa_t^2 + 15.6\kappa_t^3) + \kappa_{2t}^2 (13.9 - 22.3\kappa_t) \quad (14)$$

$$c_{11}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0978\kappa_t^2 - 0.264\kappa_t^3 + 0.673\kappa_{2t}\kappa_t, \quad (15)$$

$$c_{20}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.288\kappa_t^2 - 1.25\kappa_t^3 + 1.84\kappa_t^4 + \kappa_{2t} (2.81\kappa_t - 6.70\kappa_t^2) + 6.96\kappa_{2t}^2, \quad (16)$$

$$c_{21}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0489\kappa_t^2 \quad (17)$$

$$c_{30}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.159\kappa_t^2 - 0.417\kappa_t^3 + 0.938\kappa_{2t}\kappa_t, \quad (18)$$

$$c_{40}^{13\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0396\kappa_t^2. \quad (19)$$

At $\sqrt{s} = 13.6$ TeV, the corresponding coefficients read

$$c_{00}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = -9.34 \cdot 10^{-3} + 0.0572\kappa_t^2 - 0.446\kappa_t^3 + 1.68\kappa_t^4 - 3.17\kappa_t^5 + 2.89\kappa_t^6 \\ + \kappa_{2t} (1.59\kappa_t^2 - 7.69\kappa_t^3 + 15.3\kappa_t^4 - 14.6\kappa_t^5) \\ + \kappa_{2t}^2 (6.89 - 22.1\kappa_t + 36.8\kappa_t^2), \quad (20)$$




Figure 3: As Figure 2, but in the κ_3 - κ_{2t} plane at LHC Run 2 (left) and the HL-LHC (right). The parameters not shown explicitly are fixed to their SM values, i.e. $\kappa_4 = 1$ and $\kappa_t = 1$.

$$c_{01}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0814\kappa_t^2 - 0.260\kappa_t^3 + 0.0866\kappa_t^4 + \kappa_{2t}(0.665\kappa_t - 1.10\kappa_t^2), \quad (21)$$

$$c_{02}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0167\kappa_t^2, \quad (22)$$

$$c_{10}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.204\kappa_t^2 - 1.40\kappa_t^3 + 3.56\kappa_t^4 - 3.21\kappa_t^5 + \kappa_{2t}(3.43\kappa_t - 13.2\kappa_t^2 + 15.3\kappa_t^3) + \kappa_{2t}^2(13.8 - 22.1\kappa_t) \quad (23)$$

$$c_{11}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0961\kappa_t^2 - 0.260\kappa_t^3 + 0.665\kappa_{2t}\kappa_t, \quad (24)$$

$$c_{20}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.283\kappa_t^2 - 1.23\kappa_t^3 + 1.81\kappa_t^4 + \kappa_{2t}(2.77\kappa_t - 6.59\kappa_t^2) + 6.89\kappa_{2t}^2, \quad (25)$$

$$c_{21}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0480\kappa_t^2 \quad (26)$$

$$c_{30}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.155\kappa_t^2 - 0.408\kappa_t^3 + 0.922\kappa_{2t}\kappa_t, \quad (27)$$

$$c_{40}^{13.6\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0388\kappa_t^2. \quad (28)$$

Finally, for $\sqrt{s} = 14\text{TeV}$, we obtain

$$c_{00}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = -2.18 \cdot 10^{-3} + 0.0323\kappa_t^2 - 0.311\kappa_t^3 + 1.35\kappa_t^4 - 2.92\kappa_t^5 + 2.84\kappa_t^6 + \kappa_{2t}(1.61\kappa_t^2 - 7.79\kappa_t^3 + 15.5\kappa_t^4 - 14.7\kappa_t^5) + \kappa_{2t}^2(6.99 - 22.4\kappa_t + 37.4\kappa_t^2), \quad (29)$$

$$c_{01}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0822\kappa_t^2 - 0.262\kappa_t^3 + 0.0865\kappa_t^4 + \kappa_{2t}(0.672\kappa_t - 1.12\kappa_t^2), \quad (30)$$

$$c_{02}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0169\kappa_t^2, \quad (31)$$

$$c_{10}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.194\kappa_t^2 - 1.39\kappa_t^3 + 3.58\kappa_t^4 - 3.24\kappa_t^5 + \kappa_{2t}(3.46\kappa_t - 13.3\kappa_t^2 + 15.5\kappa_t^3) + \kappa_{2t}^2(14.0 - 22.4\kappa_t) \quad (32)$$

$$c_{11}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0968\kappa_t^2 - 0.262\kappa_t^3 + 0.672\kappa_{2t}\kappa_t, \quad (33)$$




Figure 4: As Figure 2, but in the κ_4 - κ_{2t} plane at LHC Run 2 (left) and the HL-LHC (right). The remaining parameters are fixed to their SM values, i.e. $\kappa_3 = 1$ and $\kappa_t = 1$.

$$c_{20}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.284\kappa_t^2 - 1.23\kappa_t^3 + 1.82\kappa_t^4 + \kappa_{2t}(2.79\kappa_t - 6.65\kappa_t^2) + 6.99\kappa_{2t}^2, \quad (34)$$

$$c_{21}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0484\kappa_t^2 \quad (35)$$

$$c_{30}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.156\kappa_t^2 - 0.411\kappa_t^3 + 0.931\kappa_{2t}\kappa_t, \quad (36)$$

$$c_{40}^{14\text{TeV}}(\kappa_t, \kappa_{2t}) = 0.0391\kappa_t^2. \quad (37)$$

Notice that the coefficients $c_{nm}^{\sqrt{s}}(\kappa_t, \kappa_{2t})$ depend only mildly on the collision energy over the range considered. Likewise, the choice of PDFs is expected to have a negligible numerical impact on the parameterisations given in Eqs. (11) to (37). Higher-order QCD corrections to the coefficients $c_{nm}^{\sqrt{s}}(\kappa_t, \kappa_{2t})$ are currently unknown, although they could, in principle, be estimated in the infinite top-quark mass limit. In the SM, these corrections increase the inclusive cross section by a K -factor of approximately 1.7 at NNLO in QCD [18]. Parameterisations of the LHC triple-Higgs gluon-fusion production cross section similar to Eqs. (9) to (37) have also been presented in Refs. [35, 36, 38, 39].

The two panels in Figure 2 illustrate the expected sensitivity to the Higgs self-couplings κ_3 and κ_4 using current LHC Run 2 data and HL-LHC projections, assuming an integrated luminosity of 3 ab^{-1} . The red contours indicate the 95% CL constraints derived from triple-Higgs production in gluon fusion using the parameterisations in Eqs. (11) to (19) and Eqs. (29) to (37), with the normalisation to the corresponding LO SM cross section given in Eq. (10). For LHC Run 2, the triple-Higgs constraints are obtained by imposing $\mu_{3h}^{\text{LHC Run 2}} < 588$ [20], while for the HL-LHC we assume a projected upper limit of $\mu_{3h}^{\text{HL-LHC}} < 125$ [47]. Throughout this analysis, we set $\kappa_{2t} = 0$, and the black dots indicate the SM prediction. The left panel demonstrates that modifications of the top-quark Yukawa coupling of $\Delta\kappa_t = \pm 20\%$, where $\Delta\kappa_t = \kappa_t - 1$, which remain allowed at the 95% CL after LHC Run 2 [48, 49], have a significant impact on the resulting constraints. This behaviour was already pointed out in Ref. [41] and originates from the strong power-law dependence of the coefficients $c_{nm}^{\sqrt{s}}(\kappa_t, \kappa_{2t})$ appearing in Eqs. (11) to (37) on κ_t . By the end of the HL-LHC, the top-quark Yukawa coupling is expected to be measured with a precision of $\Delta\kappa_t = \pm 5\%$ at the 95% CL [50]. Consequently, its impact on the $gg \rightarrow 3h$ constraints in the κ_3 - κ_4 plane will be substantially reduced, as illustrated in the right panel. Finally, we note that the projected HL-LHC constraints from $gg \rightarrow 3h$

are comparable to those imposed by perturbative unitarity [34, 51, 52], shown as the yellow region enclosed by the black-dotted contour in both panels.

The panels in Figures 3 and 4 show the expected sensitivity in the κ_3 – κ_{2t} and κ_4 – κ_{2t} planes, respectively. The red contours represent the 95% CL constraints from triple-Higgs production in gluon fusion, obtained following the procedure described above. Parameters not shown explicitly are fixed to their SM values, and the black dots indicate the SM prediction. As in Figure 2, the left panels demonstrate that variations of the top-quark Yukawa coupling of $\Delta\kappa_t = \pm 20\%$ can have a significant impact on the resulting constraints, whereas the smaller variations $\Delta\kappa_t = \pm 5\%$ considered in the right panels lead to only minor changes. For κ_{2t} , triple-Higgs production at LHC Run 2 yields $-5.1 < \kappa_{2t} < 5.3$, while the HL-LHC is expected to improve this range to $-2.3 < \kappa_{2t} < 2.5$ at 95% CL. These limits are obtained by fixing $\kappa_3 = \kappa_4 = \kappa_t = 1$. We note that, under the same assumptions, the 95% CL constraint on the double-Higgs production signal strength obtained at LHC Run 2, $\mu_{2h}^{\text{LHC Run 2}} < 2.5$ [17], already implies $-0.19 < \kappa_{2t} < 0.72$. The quoted bound is obtained using the expressions given in Refs. [53, 54]. Consequently, triple-Higgs production is not expected to provide a stronger constraint on κ_{2t} than double-Higgs production in a single-parameter analysis. It can, however, provide complementary information in multi-parameter fits, in particular when these involve κ_4 , which can only be weakly constrained by double-Higgs production [21–24].

2.1 Including the two-top–three-Higgs interaction

We next extend the inclusive-rate parametrisation to the contact interaction between two top quarks and three Higgs bosons in Eq. (8). We generate loop-induced LO parton-level samples with MadGraph5_aMC@NLO [55], using the same m_h , m_t , PDFs, and scale choice as above. In this scan $\kappa_t = 1$ and $\kappa_{2t} = 0$ are fixed.

Following the form of Eq. (9), we write the extended cross section as

$$\sigma_{\text{LO}}^{\sqrt{s}} = (\sigma_{\text{LO}}^{\sqrt{s}})_{\text{SM}} \sum_{n,m} c_{nm}^{\sqrt{s}}(\kappa_{3t}) (\Delta\kappa_3)^n (\Delta\kappa_4)^m. \quad (38)$$

This is an exact regrouping of the LO coupling basis at the fixed values $\kappa_t = 1$ and $\kappa_{2t} = 0$. Since the amplitude is linear in κ_{3t} , c_{00} is at most quadratic in κ_{3t} , while c_{01} , c_{10} , and c_{20} are at most linear; the remaining nonzero coefficient functions are constant. The fitted SM cross sections are

$$(\sigma_{\text{LO}}^{13\text{TeV}})_{\text{SM}} = 42.12 \text{ ab}, \quad (\sigma_{\text{LO}}^{13.6\text{TeV}})_{\text{SM}} = 47.40 \text{ ab}, \quad (\sigma_{\text{LO}}^{14\text{TeV}})_{\text{SM}} = 51.08 \text{ ab}. \quad (39)$$

They agree at the percent level with the independent values in Eq. (10).

The normalisation fixes $c_{00}^{\sqrt{s}}(0) = 1$. The fitted coefficient functions are listed below.

For $\sqrt{s} = 13 \text{ TeV}$, the non-vanishing coefficient functions are

$$c_{00}^{13\text{TeV}}(\kappa_{3t}) = 1 - 2.83183 \kappa_{3t} + 15.81829 \kappa_{3t}^2, \quad (40)$$

$$c_{01}^{13\text{TeV}}(\kappa_{3t}) = -0.0914560 + 0.793591 \kappa_{3t}, \quad (41)$$

$$c_{02}^{13\text{TeV}}(\kappa_{3t}) = 0.0168979, \quad (42)$$

$$c_{10}^{13\text{TeV}}(\kappa_{3t}) = -0.865038 - 3.92841 \kappa_{3t}, \quad (43)$$

$$c_{11}^{13\text{TeV}}(\kappa_{3t}) = -0.167158, \quad (44)$$

$$c_{20}^{13\text{TeV}}(\kappa_{3t}) = 0.872294 + 0.900894 \kappa_{3t}, \quad (45)$$

$$c_{21}^{13\text{TeV}}(\kappa_{3t}) = 0.0487935, \quad (46)$$

$$c_{30}^{13\text{TeV}}(\kappa_{3t}) = -0.259375, \quad (47)$$

$$c_{40}^{13\text{TeV}}(\kappa_{3t}) = 0.0395354. \quad (48)$$

For $\sqrt{s} = 13.6 \text{ TeV}$, they are

$$c_{00}^{13.6 \text{ TeV}}(\kappa_{3t}) = 1 - 2.87872 \kappa_{3t} + 16.21306 \kappa_{3t}^2, \quad (49)$$

$$c_{01}^{13.6 \text{ TeV}}(\kappa_{3t}) = -0.0909154 + 0.800038 \kappa_{3t}, \quad (50)$$

$$c_{02}^{13.6 \text{ TeV}}(\kappa_{3t}) = 0.0168638, \quad (51)$$

$$c_{10}^{13.6 \text{ TeV}}(\kappa_{3t}) = -0.866564 - 3.95165 \kappa_{3t}, \quad (52)$$

$$c_{11}^{13.6 \text{ TeV}}(\kappa_{3t}) = -0.166501, \quad (53)$$

$$c_{20}^{13.6 \text{ TeV}}(\kappa_{3t}) = 0.871502 + 0.897150 \kappa_{3t}, \quad (54)$$

$$c_{21}^{13.6 \text{ TeV}}(\kappa_{3t}) = 0.0485042, \quad (55)$$

$$c_{30}^{13.6 \text{ TeV}}(\kappa_{3t}) = -0.257216, \quad (56)$$

$$c_{40}^{13.6 \text{ TeV}}(\kappa_{3t}) = 0.0390715. \quad (57)$$

For $\sqrt{s} = 14 \text{ TeV}$, they are

$$c_{00}^{14 \text{ TeV}}(\kappa_{3t}) = 1 - 2.95187 \kappa_{3t} + 16.47338 \kappa_{3t}^2, \quad (58)$$

$$c_{01}^{14 \text{ TeV}}(\kappa_{3t}) = -0.0921387 + 0.803365 \kappa_{3t}, \quad (59)$$

$$c_{02}^{14 \text{ TeV}}(\kappa_{3t}) = 0.0168553, \quad (60)$$

$$c_{10}^{14 \text{ TeV}}(\kappa_{3t}) = -0.853992 - 3.93891 \kappa_{3t}, \quad (61)$$

$$c_{11}^{14 \text{ TeV}}(\kappa_{3t}) = -0.166145, \quad (62)$$

$$c_{20}^{14 \text{ TeV}}(\kappa_{3t}) = 0.863440 + 0.899872 \kappa_{3t}, \quad (63)$$

$$c_{21}^{14 \text{ TeV}}(\kappa_{3t}) = 0.0482948, \quad (64)$$

$$c_{30}^{14 \text{ TeV}}(\kappa_{3t}) = -0.255934, \quad (65)$$

$$c_{40}^{14 \text{ TeV}}(\kappa_{3t}) = 0.0390078. \quad (66)$$




Figure 5: Constraints in the κ_3 – κ_{3t} plane at LHC Run 2 (left) and the HL-LHC (right), obtained from Eq. (38) using $\mu_{3h} < 588$ and $\mu_{3h} < 125$, respectively. The red regions are allowed by the inclusive triple-Higgs rate, the black points denote the SM, and the yellow vertical bands show the projection of the tree-level $hh \rightarrow hh$ perturbative-unitarity constraint. We fix $\kappa_4 = \kappa_t = 1$ and $\kappa_{2t} = 0$. The yellow band constrains κ_3 only; no independent unitarity condition on κ_{3t} is imposed.




Figure 6: As in Figure 5, but in the κ_4 – κ_{3t} plane with $\kappa_3 = \kappa_t = 1$ and $\kappa_{2t} = 0$. The yellow band is the projection of the $hh \rightarrow hh$ perturbative-unitarity condition onto κ_4 .

Figures 5 and 6 are the κ_{3t} analogues of Figures 3 and 4. Fixing all other couplings to their SM values yields

$$-6.00 < \kappa_{3t} < 6.18 \quad (\text{LHC Run 2}), \quad -2.66 < \kappa_{3t} < 2.83 \quad (\text{HL-LHC}). \quad (67)$$

The quadratic form also exposes an inclusive-rate degeneracy. At $\kappa_3 = \kappa_4 = 1$, the two solutions of $c_{00}^{\sqrt{s}}(\kappa_{3t}) = 1$ are $\kappa_{3t} = 0$ and $\kappa_{3t} \simeq 0.179$ at all three energies to the accuracy of the fit. Differential information is therefore essential for distinguishing these two solutions.

Modifications of the Higgs trilinear and quartic self-couplings affect not only the inclusive triple-Higgs production cross section but also the kinematic properties of the $gg \rightarrow 3h$ final state. To illustrate this point, Figure 7 shows normalised kinematic distributions at the HL-LHC for the SM and two BSM benchmark scenarios. Both benchmark points yield the same inclusive signal strength, $\mu_{3h} = 65$, with inclusive cross sections differing by only about 5%,




Figure 7: Normalised kinematic distributions for $gg \rightarrow 3h$ production at the HL-LHC in the SM (solid black line) and in two benchmark scenarios defined by κ_3 and κ_4 (red dashed and blue dotted lines). The left and right panels show the invariant mass of the three-Higgs system, m_{3h} , and the scalar sum of the transverse momenta of the Higgs bosons, $\sum p_{T,h}$, respectively. All results are obtained for $\kappa_t = 1$ and $\kappa_{2t} = 0$.




Figure 8: As in Figure 7, but for two other κ -benchmark scenarios indicated in the panels. The remaining parameters are fixed to their SM values.

while remaining consistent with the perturbative unitarity bounds obtained from tree-level $hh \rightarrow hh$ scattering. The left and right panels show the invariant mass of the three-Higgs system, m_{3h} , and the scalar sum of the Higgs transverse momenta, $\sum p_{T,h}$, respectively. All results are obtained for $\kappa_t = 1$ and $\kappa_{2t} = 0$. This choice is motivated by the observation of Ref. [41] that variations of κ_t approximately induce an overall rescaling of the normalised distributions, without significantly altering their shapes. The comparison demonstrates that values of κ_3 and κ_4 yielding nearly degenerate inclusive cross sections can nevertheless lead to sizeable distortions of the differential spectra, reaching up to about 50% in both the m_{3h} and $\sum p_{T,h}$ distributions, in the peak regions as well as in the high-energy tails. Consequently, exploiting differential information in future HL-LHC analyses of $gg \rightarrow 3h$ production has the potential to improve the constraint quoted below Eq. (6), which currently relies on the inclusive rate only [21].




Figure 9: Normalised m_{3h} and $\sum p_{T,h}$ distributions at the HL-LHC for the SM and the two $\mu_{3h} \simeq 65$ benchmarks $(\kappa_3, \kappa_4, \kappa_{3t}) = (2.1, 23, -2.3)$ and $(2.2, 0, 2.2)$. The distributions use the same 40 GeV binning and displayed ranges as Figures 7 and 8. We fix $\kappa_t = 1$ and $\kappa_{2t} = 0$.

Figure 8 presents additional normalised kinematic distributions at the HL-LHC for the SM and two further BSM benchmark scenarios within the κ -framework. The benchmark points are selected such that both lead to an enhanced signal strength of $\mu_{3h} = 75$, while their inclusive cross sections differ by less than about 5%. The corresponding values of κ_3 and κ_4 are consistent with perturbative unitarity constraints, and all remaining parameters are fixed to their SM values. As in Figure 7, sizeable distortions of the differential spectra arise despite the near-degeneracy of the inclusive rates. This further demonstrates that exploiting differential information in future HL-LHC searches for $gg \rightarrow 3h$ production can help lift parameter degeneracies within the κ -framework.

The two mixed-coupling benchmarks in Figure 9 provide more dramatic examples at an enhanced but nearly degenerate inclusive rate. Both have $\mu_{3h} \simeq 65$, yet produce large distortions of the normalised distributions relative to the SM, including a pronounced migration towards larger invariant masses and transverse momenta. The samples use the same PDFs and scale choice as the inclusive-rate parametrisation.

3 Conclusions

Unlike the Higgs couplings to EW gauge bosons and third-generation fermions, which are already being measured with increasing precision, the Higgs self-interactions remain largely unconstrained by LHC Run 2 data. This leaves considerable room for BSM scenarios that modify the shape of the Higgs potential. Indeed, both theoretical considerations and explicit ultraviolet-complete constructions demonstrate that deviations of the Higgs self-couplings exceeding 200% can be consistent with existing single-Higgs measurements without requiring fine-tuned parameter choices [4]. Measurements of multi-Higgs production at the HL-LHC and future colliders therefore provide a unique opportunity to probe largely unexplored regions of parameter space and retain significant discovery potential, even in scenarios where single-Higgs observables remain close to their SM predictions.

In this note, we have derived compact “pocket formulas” for triple-Higgs production within the κ -framework, enabling constraints on the Higgs trilinear and quartic self-couplings to be obtained from current and future measurements of the $gg \rightarrow 3h$ signal strength. The origi-

nal parameterisation retains the full dependence on κ_3 , κ_4 , κ_t , and κ_{2t} at LO in perturbation theory. We have supplemented it with Eq. (38), which includes the complete LO dependence on κ_{3t} for $\kappa_t = 1$ and $\kappa_{2t} = 0$ at 13, 13.6, and 14 TeV. Since the derivation relies on flexible matrix-element implementations, further extensions to incorporate additional coupling modifications — such as those arising at next-to-leading order in the Higgs Effective Field Theory [54] — are comparatively straightforward.

Using these parameterisations, we have investigated the sensitivity to κ_3 and κ_4 using current and projected LHC data. We find that the presently allowed uncertainty in κ_t can have a substantial impact on the constraints in the κ_3 – κ_4 plane. This dependence is significantly reduced at the HL-LHC, where the expected percent-level precision of the top-quark Yukawa coupling measurement will strongly restrict the corresponding parameter space. For κ_{2t} , which parametrises the interaction between two top quarks and two Higgs bosons, we find that triple-Higgs production is unlikely to provide stronger bounds than double-Higgs production in a single-parameter analysis. Nevertheless, it can yield complementary information in global multi-parameter fits, especially once κ_4 is included, as this coupling remains only weakly constrained by double-Higgs measurements [21–24]. In a single-parameter interpretation, the present and projected triple-Higgs rate assumptions give $-6.00 < \kappa_{3t} < 6.18$ and $-2.66 < \kappa_{3t} < 2.83$, respectively. The nonzero solution near $\kappa_{3t} = 0.179$ is rate-degenerate with the SM. We further demonstrate that differential information lifts inclusive-rate degeneracies and identify mixed $(\kappa_3, \kappa_4, \kappa_{3t})$ benchmarks with $\mu_{3h} \simeq 65$ but substantially different kinematic distributions from the SM.

The projected HL-LHC sensitivity to κ_4 in a single-parameter analysis remains modest, with constraints reaching only $|\kappa_4| \lesssim 80$, comparable to those derived from perturbative unitarity arguments [34, 51, 52]. An improvement by approximately one order of magnitude in the determination of the Higgs quartic self-coupling could be achieved at a hadron-collider option of the Future Circular Collider [22, 28–30, 32, 33, 35–38, 40]. Reaching a precision at the level of 50%, however, would likely require the capabilities of a future muon collider [56].

Acknowledgements

UH acknowledges Konstantin Schmid for helpful discussions. This manuscript has been authored by Fermi Forward Discovery Group, LLC under Contract No. 89243024CSC000002 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. RKE acknowledges receipt of a Leverhulme Emeritus Fellowship from the Leverhulme Trust.

References

- [1] G. Aad *et al.*, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Phys. Lett. B **716**, 1 (2012), doi:10.1016/j.physletb.2012.08.020, 1207.7214.
- [2] S. Chatrchyan *et al.*, *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, Phys. Lett. B **716**, 30 (2012), doi:10.1016/j.physletb.2012.08.021, 1207.7235.
- [3] J. Alison *et al.*, *Higgs boson potential at colliders: Status and perspectives*, Rev. Phys. **5**, 100045 (2020), doi:10.1016/j.revip.2020.100045, 1910.00012.
- [4] G. Durieux, M. McCullough and E. Salvioni, *Charting the Higgs self-coupling bound-*

- aries, JHEP **12**, 148 (2022), doi:10.1007/JHEP12(2022)148, [Erratum: JHEP **02**, 165 (2023)], 2209.00666.
- [5] M. Gorbahn and U. Haisch, *Indirect probes of the trilinear Higgs coupling: $gg \rightarrow h$ and $h \rightarrow \gamma\gamma$* , JHEP **10**, 094 (2016), doi:10.1007/JHEP10(2016)094, 1607.03773.
- [6] G. Degrandi, P. P. Giardino, F. Maltoni and D. Pagani, *Probing the Higgs self coupling via single Higgs production at the LHC*, JHEP **12**, 080 (2016), doi:10.1007/JHEP12(2016)080, 1607.04251.
- [7] W. Bizon, M. Gorbahn, U. Haisch and G. Zanderighi, *Constraints on the trilinear Higgs coupling from vector boson fusion and associated Higgs production at the LHC*, JHEP **07**, 083 (2017), doi:10.1007/JHEP07(2017)083, 1610.05771.
- [8] F. Maltoni, D. Pagani, A. Shivaji and X. Zhao, *Trilinear Higgs coupling determination via single-Higgs differential measurements at the LHC*, Eur. Phys. J. C **77**(12), 887 (2017), doi:10.1140/epjc/s10052-017-5410-8, 1709.08649.
- [9] M. Gorbahn and U. Haisch, *Two-loop amplitudes for Higgs plus jet production involving a modified trilinear Higgs coupling*, JHEP **04**, 062 (2019), doi:10.1007/JHEP04(2019)062, 1902.05480.
- [10] U. Haisch and G. Koole, *Off-shell Higgs production at the LHC as a probe of the trilinear Higgs coupling*, JHEP **02**, 030 (2022), doi:10.1007/JHEP02(2022)030, 2111.12589.
- [11] J. Gao, X.-M. Shen, G. Wang, L. L. Yang and B. Zhou, *Probing the Higgs boson trilinear self-coupling through Higgs boson+jet production*, Phys. Rev. D **107**(11), 115017 (2023), doi:10.1103/PhysRevD.107.115017, 2302.04160.
- [12] U. Haisch and M. Niggetiedt, *Exact two-loop amplitudes for Higgs plus jet production with a cubic Higgs self-coupling*, JHEP **10**, 236 (2024), doi:10.1007/JHEP10(2024)236, 2408.13186.
- [13] A. Ghosh, M. Griesse, U. Haisch and T. H. Park, *Neural simulation-based inference of the Higgs trilinear self-coupling via off-shell Higgs production*, Eur. Phys. J. C **86**(4), 415 (2026), doi:10.1140/epjc/s10052-026-15605-3, 2507.02032.
- [14] A. David, A. Denner, M. Duehrssen, M. Grazzini, C. Grojean, G. Passarino, M. Schumacher, M. Spira, G. Weiglein and M. Zanetti, *LHC HXSWG interim recommendations to explore the coupling structure of a Higgs-like particle* (2012), 1209.0040.
- [15] J. R. Andersen et al., *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties* (2013), doi:10.5170/CERN-2013-004, 1307.1347.
- [16] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, CERN Yellow Rep. Monogr. **2**, 1 (2017), doi:10.23731/CYRM-2017-002, 1610.07922.
- [17] G. Aad et al., *Combination of ATLAS and CMS searches for Higgs boson pair production at $\sqrt{s} = 13$ TeV* (2026), 2602.23991.
- [18] D. de Florian, I. Fabre and J. Mazzitelli, *Triple Higgs production at hadron colliders at NNLO in QCD*, JHEP **03**, 155 (2020), doi:10.1007/JHEP03(2020)155, 1912.02760.
- [19] G. Aad et al., *Search for triple Higgs boson production in the $6b$ final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Phys. Rev. D **111**(3), 032006 (2025), doi:10.1103/PhysRevD.111.032006, 2411.02040.

- [20] *Search for nonresonant triple Higgs boson production in the six b -quark final state in proton-proton collisions at 13 TeV*, CERN, Geneva (2025), <https://cds.cern.ch/record/2945361>.
- [21] J.-L. Ding *et al.*, *Constraining the Higgs potential using multi-Higgs production* (2026), doi:10.21468/SciPostPhysCommRep.24, 2601.13248.
- [22] W. Bizoń, U. Haisch and L. Rottoli, *Constraints on the quartic Higgs self-coupling from double-Higgs production at future hadron colliders*, JHEP **10**, 267 (2019), doi:10.1007/JHEP02(2024)170, 1810.04665.
- [23] S. Borowka, C. Duhr, F. Maltoni, D. Pagani, A. Shivaji and X. Zhao, *Probing the scalar potential via double Higgs boson production at hadron colliders*, JHEP **04**, 016 (2019), doi:10.1007/JHEP04(2019)016, 1811.12366.
- [24] H. T. Li, Z.-G. Si, J. Wang, X. Zhang and D. Zhao, *Improved constraints on Higgs boson self-couplings with quartic and cubic power dependencies of the cross section*, Chin. Phys. C **49**(2), 023107 (2025), doi:10.1088/1674-1137/ad9d1d, 2407.14716.
- [25] T. Plehn and M. Rauch, *The quartic Higgs coupling at hadron colliders*, Phys. Rev. D **72**, 053008 (2005), doi:10.1103/PhysRevD.72.053008, hep-ph/0507321.
- [26] T. Binoth, S. Karg, N. Kauer and R. Ruckl, *Multi-Higgs boson production in the Standard Model and beyond*, Phys. Rev. D **74**, 113008 (2006), doi:10.1103/PhysRevD.74.113008, hep-ph/0608057.
- [27] F. Maltoni, E. Vryonidou and M. Zaro, *Top-quark mass effects in double and triple Higgs production in gluon-gluon fusion at NLO*, JHEP **11**, 079 (2014), doi:10.1007/JHEP11(2014)079, 1408.6542.
- [28] A. Papaefstathiou and K. Sakurai, *Triple Higgs boson production at a 100 TeV proton-proton collider*, JHEP **02**, 006 (2016), doi:10.1007/JHEP02(2016)006, 1508.06524.
- [29] C.-Y. Chen, Q.-S. Yan, X. Zhao, Y.-M. Zhong and Z. Zhao, *Probing triple-Higgs productions via $4b2\gamma$ decay channel at a 100 TeV hadron collider*, Phys. Rev. D **93**(1), 013007 (2016), doi:10.1103/PhysRevD.93.013007, 1510.04013.
- [30] B. Fuks, J. H. Kim and S. J. Lee, *Probing Higgs self-interactions in proton-proton collisions at a center-of-mass energy of 100 TeV*, Phys. Rev. D **93**(3), 035026 (2016), doi:10.1103/PhysRevD.93.035026, 1510.07697.
- [31] D. de Florian and J. Mazzitelli, *Two-loop corrections to the triple Higgs boson production cross section*, JHEP **02**, 107 (2017), doi:10.1007/JHEP02(2017)107, 1610.05012.
- [32] W. Kilian, S. Sun, Q.-S. Yan, X. Zhao and Z. Zhao, *New Physics in multi-Higgs boson final states*, JHEP **06**, 145 (2017), doi:10.1007/JHEP06(2017)145, 1702.03554.
- [33] B. Fuks, J. H. Kim and S. J. Lee, *Scrutinizing the Higgs quartic coupling at a future 100 TeV proton-proton collider with taus and b -jets*, Phys. Lett. B **771**, 354 (2017), doi:10.1016/j.physletb.2017.05.075, 1704.04298.
- [34] P. Stylianou and G. Weiglein, *Constraints on the trilinear and quartic Higgs couplings from triple Higgs production at the LHC and beyond*, Eur. Phys. J. C **84**(4), 366 (2024), doi:10.1140/epjc/s10052-024-12722-9, 2312.04646.
- [35] A. Papaefstathiou and G. Tetlalmatzi-Xolocotzi, *Multi-Higgs boson production with anomalous interactions at current and future proton colliders*, JHEP **06**, 124 (2024), doi:10.1007/JHEP06(2024)124, 2312.13562.

- [36] H. Abouabid *et al.*, *HHH whitepaper*, Eur. Phys. J. C **84**, 1183 (2024), doi:10.1140/epjc/s10052-024-13376-3, 2407.03015.
- [37] Z. Dong, X. Sun, B. Guo, L. Zhang, Z. Li, J. Wang, Z. Li, Y. Ban and Y. Mao, *Probing triple Higgs production via $4\tau 2b$ decay channel at a 100 TeV hadron collider*, JHEP **08**, 040 (2025), doi:10.1007/JHEP08(2025)040, 2504.04037.
- [38] U. Haisch, A. Sankar and G. Zanderighi, *A new probe of the quartic Higgs self-coupling* (2025), 2505.20463.
- [39] J. M. Campbell, G. De Laurentis and R. K. Ellis, *An analytic result for the $0 \rightarrow ggHHH$ amplitude*, JHEP **01**, 157 (2026), doi:10.1007/JHEP01(2026)157, 2507.19313.
- [40] B. Fuks, A. Papaefstathiou and G. Tetlalmatzi-Xolocotzi, *Extracting Higgs self-coupling constraints through triple Higgs boson production at future hadron colliders*, Eur. Phys. J. C **85**(11), 1309 (2025), doi:10.1140/epjc/s10052-025-15051-7, 2509.16364.
- [41] L. Panizzi, *Role of the top Yukawa coupling in triple Higgs production at the LHC*, Phys. Rev. D **113**(5), L051706 (2026), doi:10.1103/5ssz-3nh7, 2512.03003.
- [42] J. M. Campbell and R. K. Ellis, *An Update on vector boson pair production at hadron colliders*, Phys. Rev. D **60**, 113006 (1999), doi:10.1103/PhysRevD.60.113006, hep-ph/9905386.
- [43] J. M. Campbell, R. K. Ellis and C. Williams, *Vector Boson Pair Production at the LHC*, JHEP **07**, 018 (2011), doi:10.1007/JHEP07(2011)018, 1105.0020.
- [44] J. M. Campbell, R. K. Ellis and W. T. Giele, *A Multi-Threaded Version of MCFM*, Eur. Phys. J. C **75**(6), 246 (2015), doi:10.1140/epjc/s10052-015-3461-2, 1503.06182.
- [45] J. Campbell and T. Neumann, *Precision Phenomenology with MCFM*, JHEP **12**, 034 (2019), doi:10.1007/JHEP12(2019)034, 1909.09117.
- [46] R. D. Ball *et al.*, *The PDF4LHC21 combination of global PDF fits for the LHC Run III*, J. Phys. G **49**(8), 080501 (2022), doi:10.1088/1361-6471/ac7216, 2203.05506.
- [47] *HL-LHC prospects for the measurement of triple-Higgs production in the $6b$ final state at the ATLAS experiment*, CERN, Geneva, (2025), <https://cds.cern.ch/record/2924772>.
- [48] G. Aad *et al.*, *A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery*, Nature **607**(7917), 52 (2022), doi:10.1038/s41586-022-04893-w, [Erratum: Nature **612**, E24 (2022)], 2207.00092.
- [49] A. Tumasyan *et al.*, *A portrait of the Higgs boson by the CMS experiment ten years after the discovery*, Nature **607**(7917), 60 (2022), doi:10.1038/s41586-022-04892-x, [Erratum: Nature **623**, (2023)], 2207.00043.
- [50] *Highlights of the HL-LHC physics projections by ATLAS and CMS*, CERN, Geneva, (2025), <https://cds.cern.ch/record/2929200>.
- [51] T. Liu, K.-F. Lyu, J. Ren and H. X. Zhu, *Probing the quartic Higgs boson self-interaction*, Phys. Rev. D **98**(9), 093004 (2018), doi:10.1103/PhysRevD.98.093004, 1803.04359.
- [52] L. Di Luzio, R. Gröber and M. Spannowsky, *Maxi-sizing the trilinear Higgs self-coupling: how large could it be?*, Eur. Phys. J. C **77**(11), 788 (2017), doi:10.1140/epjc/s10052-017-5361-0, 1704.02311.

- [53] G. Buchalla, M. Capozzi, A. Celis, G. Heinrich and L. Scyboz, *Higgs boson pair production in non-linear Effective Field Theory with full m_t -dependence at NLO QCD*, JHEP **09**, 057 (2018), doi:10.1007/JHEP09(2018)057, [Erratum: JHEP **06**, 094 (2025)], 1806.05162.
- [54] I. Brivio, R. Gröber and K. Schmid, *Higgs pair production in gluon fusion to higher orders in Higgs Effective Field Theory*, JHEP **05**, 031 (2026), doi:10.1007/JHEP05(2026)031, 2511.23411.
- [55] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, JHEP **07**, 079 (2014), doi:10.1007/JHEP07(2014)079, 1405.0301.
- [56] M. Chiesa, F. Maltoni, L. Mantani, B. Mele, F. Piccinini and X. Zhao, *Measuring the quartic Higgs self-coupling at a multi-TeV muon collider*, JHEP **09**, 098 (2020), doi:10.1007/JHEP09(2020)098, 2003.13628.