

Social vulnerability is associated with higher rate of emergency department visits and hospitalizations with vaccine preventable diseases among Colorado children.

Source: PubMed

Published: 2026 Apr 10

Cross-referenced sources: 30

Original URL: <https://pubmed.ncbi.nlm.nih.gov/41965979/>

Social vulnerability is associated with higher rate of emergency department visits and hospitalizations with vaccine preventable diseases among Colorado children. A reported public-health development reviewed against 30 corroborating references, with evidence currently assessed as high.

By Public Health Research Desk Published April 12, 2026

Review of available evidence identified 13 peer-reviewed references, 17 trusted news references, and 0 preprint references from verified domains. This mix supports contextual interpretation, but study design and population differences still matter.

Evidence should be monitored with caution and interpreted alongside independent updates.

Not all studies carry equal weight, and conclusions may change as additional data emerges.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

[illegible]

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

Long-term interpretation remains essential: surveillance findings can shift as new cohorts are analyzed, methods improve, and independent teams replicate or challenge early claims.

A version of this article appears in print in the Public Health Research edition with the headline above. Related: Global, regional, and national burden of meningitis, its risk factors, and aetiologies, 1990-2023: a systematic analysis for the Global Burden of Disease Study 2023. Global burden of lower respiratory infections and aetiologies, 1990-2023: a systematic analysis for the Global Burden of Disease Study 2023.