

University Important questions

Q1. What is cardiac cycle. Describe events occurring during normal cardiac cycle.

Q2. Factors affecting cardiac cycle

Q3. Enumerate properties of cardiac muscles.

Q4. Draw and label a diagram of normal ECG. Describe the various waves in ECG.

Q5. Describe variations of heart rate, effect of baro reflex and chemo reflex on heart rate and hormonal regulation of heart rate.

Q6. What is cardiac output. Mention factors affecting cardiac output. Add a note on Venous return.

Q7. What is the normal arterial blood pressure and physiological variations. Describe role of sinoaortic mechanism of regulation of blood pressure

Q8. Explain regulation of blood pressure brought about by baroreceptor mechanism

Q9. What is the systolic diastolic and pulse pressure. Give account of factors that maintain blood pressure.

Q10. What are the heart sounds. How are they produced. Mention their clinical significance.

The most important questions are:

Q1, Q2, Q6, Q7, Q8, Q9

Q1. What is cardiac cycle. Describe events occurring during normal cardiac cycle.

Cardiac Cycle

Definition

- **Sequence of events** in the **heart** during **each beat**
- **Duration** → **0.8 sec** (Heart rate 75 bpm)
- **Higher HR** → **Shorter duration**
- **Lower HR** → **Longer duration**
- **Right & Left chambers** function **simultaneously**

Types of Cycles

- **Atrial Cycle**
- **Ventricular Cycle**

Phases of Cardiac Cycle

1. **Atrial Cycle**
 - **Atrial Systole** → **0.10 sec**
 - **Atrial Diastole** → **0.70 sec**
2. **Ventricular Cycle**
 - **Ventricular Systole** → **0.30 sec**
 - **Ventricular Diastole** → **0.50 sec**

Atrial Cycle

Atrial Systole (AS)

- **Two Phases** → **Dynamic (0.05 sec)**, **Adynamic (0.05 sec)**
- **Right atrial systole** starts **before left**
- **Duration** → **0.10 sec**
- **Contributes 20%** to **ventricular filling**
- **Blood flows** from **atria to ventricles** due to **pressure gradient**

Atrial Diastole (AD)

- **First 0.30 sec** → **Coincides with Ventricular Systole**
- **Later 0.40 sec** → **Synchronizes with Ventricular Diastole**
- **Duration** → **0.70 sec**
- **A-V valves closed** at start → **Veins fill atria**
- **After 0.3 sec**, **A-V valves open** → **80% ventricular filling**

Ventricular Cycle

Ventricular Systole (VS)

- **Right ventricle** starts **earlier** than **left**
- **Duration** → **0.30 sec**
- **A-V valves close** → **First heart sound**
- **Interval between 1st & 2nd heart sounds** → **Ventricular systole duration**

Phases of Ventricular Systole

1. **Isovolumetric Contraction** → **0.05 sec**
 - **A-V & Semilunar valves closed**
 - **No volume change, only pressure increase**
 - **Ends when semilunar valves open**
2. **Maximum Ejection** → **0.10 sec**
 - **Peak ventricular pressure**
 - **Maximum blood flow into vessels**
3. **Reduced Ejection** → **0.15 sec**
 - **Lower pressure gradient** → **Less blood flow**
 - **Ends with closure of semilunar valves** → **Second heart sound**

Ventricular Diastole (VD)

- **Starts before semilunar valve closure**

Phases of Ventricular Diastole

1. **Protodiastole** → **0.04 sec**
 - **Ventricular ejection stops**
 - **Pressure in vessels > Ventricles** → **Semilunar valves close** → **Second heart sound**
2. **Isovolumetric Relaxation** → **0.06 sec**
 - **Semilunar & A-V valves closed**
 - **Atria filling** → **A-V valves open** → **Ventricular filling starts**
3. **Ventricular Filling** → **0.40 sec**
 - **First Rapid Filling** → **0.10 sec (70% filling)** → **Third heart sound**
 - **Slow Filling** → **0.20 sec (10% filling)**
 - **Second Rapid Filling** → **0.10 sec (20% filling)** → **Fourth heart sound**

Other Events in Cardiac Cycle

- **Pressure changes** in **heart & vessels**
- **Blood volume changes** in **chambers**

- **Arterial & venous pulse** production
- **EKG generation**
- **Heart sounds production**
- **Apical impulse (Apex beat)**

Q2. Factors affecting cardiac cycle

- **Heart rate:** The number of heartbeats per minute affects the duration of the cardiac cycle. Increased heart rate shortens the cycle.
- **Contractility:** The strength of contraction influences the volume of blood pumped, affecting systole duration.
- **Preload:** The initial stretching of the heart muscle before contraction, influenced by venous return, affects stroke volume and cardiac cycle.
- **Afterload:** The resistance the heart must overcome to eject blood. Increased afterload slows down the ejection phase of the cycle.
- **Autonomic Nervous System:**
 - **Sympathetic stimulation:** Increases heart rate and contractility, leading to a faster cycle.
 - **Parasympathetic stimulation:** Decreases heart rate, leading to a slower cycle.
- **Blood volume:** Increased blood volume leads to higher preload, affecting the overall cycle.
- **Age:** Younger individuals tend to have a faster heart rate, affecting the cycle duration.
- **Temperature:** Higher body temperature increases heart rate, shortening the cardiac cycle.
- **Electrolyte balance:** Imbalances (like low potassium or calcium) can affect the timing and efficiency of the cardiac cycle.


Q3. Enumerate properties of cardiac muscles.

Properties of Cardiac Muscles

- **Contractility:** Cardiac muscle has the ability to contract and generate force to pump blood throughout the body.
- **Excitability:** Cardiac muscle can respond to stimuli, particularly electrical impulses, which initiate contraction.
- **Rhythmicity:** Cardiac muscle fibers have the ability to contract in a regular, rhythmic pattern, allowing continuous heartbeats.
- **Automaticity:** Cardiac muscle can generate its own electrical impulses without external nerve stimulation.
- **Conductivity:** Cardiac muscle can transmit electrical impulses from one fiber to another, ensuring coordinated contractions.
- **Refractoriness:** After contraction, cardiac muscle enters a refractory period where it cannot respond to new stimuli, preventing tetany.
- **Elasticity:** Cardiac muscle has the ability to return to its resting shape after being stretched during diastole.

- **Endurance:** Cardiac muscle has a high endurance capacity, allowing it to contract continuously without fatigue.

Q4. Draw and label a diagram of normal ECG. Describe the various waves in ECG.



ELECTROCARDIOGRAM (ECG)

- Graphical recording of **summed electrical activity** from the heart during a **cardiac cycle**
- Shows five deflections, three **positive waves** (upward) and two **negative waves** (downward)

PQRST COMPLEX

- Each heartbeat results in a complex of waves called the **PQRST complex**
- **Baseline** is called the **isoelectric line** (no electrical activity)

WAVE CHARACTERISTICS

- **P WAVE:** Represents **atrial depolarization**, normal duration is **0.08-0.12 sec**, voltage **0.1-0.3 mV**
- **Q WAVE:** Represents **septal depolarization**, absent in **right ventricular leads**
- **R WAVE:** Potential traveling to the **apex of the ventricle**, peak of **R wave** is the **safest period for cardioversion**
- **S WAVE:** Represents potential traveling to the **base of the ventricles**, **RS complex** prominent in **right ventricular leads**
- **T WAVE:** Represents **ventricular repolarization**, normal duration is **0.27 sec**, voltage varies **0.2-1 mV**, **positive wave**
- **U WAVE:** Caused by delayed repolarization of the **intravenous conduction system**, usually seen after the **T wave**

INTERVALS AND SEGMENTS

- **P-R INTERVAL:** Time from the beginning of **P wave** to the beginning of **Q/R wave**, normal **0.12-0.21 sec**
- **P-R SEGMENT:** Time between end of **P wave** and beginning of **Q wave**
- **S-T SEGMENT:** Time from end of **S wave** to beginning of **T wave**, indicates completion of **ventricular depolarization**
- **Q-T INTERVAL:** Time from the beginning of **Q wave** to the end of **T wave**, normal **0.4 sec**, reflects time for **ventricular depolarization and repolarization**
- **S-T INTERVAL:** Time from end of **S wave** to end of **T wave**, normal up to **0.32 sec**

ABNORMALITIES

- **Prolonged P-R interval** in conditions like **1° heart block** and **2° heart block**
- **Shortened P-R interval** in **Wolf-Parkinson-White syndrome** and **Lown-Ganong-Levine syndrome**
- **ST-T changes** in **myocardial infarction**, **ischemia**, **endocrine disorders**, **arrhythmias**, **electrolyte imbalances**, **toxicity from drugs**

CLINICAL APPLICATIONS

- Useful for diagnosis, prognosis, and management in:
 - **Myocardial infarction and ischemia**
 - **Cardiac arrhythmias**
 - **Electrolyte imbalances**
 - **Myocarditis and cardiomyopathies**
 - **Endocrine disorders**
 - **Toxic effects of drugs on the heart**
 - **Conduction defects**
 - **Chamber hypertrophy**

CARDIAC MONITORING

- Continuous ECG recording during:
 - **Cardiac surgery**
 - **Thoracic surgery**
 - **Treatment of arrhythmias**
 - **Acute myocardial infarction**
 - **Postoperative intensive care**
 - **All emergencies**

Q5. Describe variations of heart rate, effect of baro reflex and chemo reflex on heart rate and hormonal regulation of heart rate.

Physiological Variations of Heart Rate

1. Age

- Heart rate in **infants** is high (140-150/min).
- As one grows up, HR normalizes around **puberty** due to increased **vagal tone**.

2. Surface Area

- HR is **inversely proportional** to surface area.
- Example: **Mouse HR = 600/min, Elephant HR = 40/min.**

3. Sex

- **Females** have a higher HR compared to **males** of the same age.
- This could be due to a higher **sympathetic tone** or less **vagal tone** in females.

4. Emotions

- HR increases during **anger, excitement, and fear.**

5. Environmental Temperature

- Higher **temperature** increases HR.
- Every **1°F** rise = HR increases **10 bpm.**
- Every **1°C** rise = HR increases **20 bpm.**

6. Metabolic Rate

- Higher **metabolic rate** increases HR.
- **Hyperthyroidism** = High HR and **BMR.**
- **Hypothyroidism** = Low HR and **BMR.**

7. Posture

- **Standing up** decreases **venous return**, reducing **cardiac output** and **BP.**
- This triggers an increase in HR (**Marey's reflex**).

8. Respiration

- HR increases during **inspiration** and falls during **expiration.**
- Known as **sinus arrhythmia**, normal in **children** and some **adults.**

9. Barometric Pressure

- Increased **atmospheric pressure** usually causes a decrease in HR.
- Example: **High altitude** (low pressure) = HR increases.
- Example: **Deep-sea diving** (high pressure) = HR decreases.

10. Sleep

- HR is **minimal** during sleep due to low **metabolic rate**.

Cardiovascular Reflexes

- **Baroreflex** and **chemoreflex** influence both **BP** and HR.

1. Baroreflex

- **Baroreceptor** activation stimulates the **nucleus tractus solitarius** (NTS).
- NTS activates the **nucleus ambiguus**, increasing **vagal tone** and lowering HR.

2. Chemoreflex

- **Chemoreceptor** activation stimulates the **inspiratory center**.
- This inhibits the **nucleus ambiguus**, reducing **vagal tone** and raising HR.
- Inhibited **inspiratory center** can cause **bradycardia** or **asystole**.

Hormonal Regulation of Heart Rate

1. Adrenaline

- Increases HR by acting on **beta receptors** of the heart, increasing **cyclic AMP**.

2. Noradrenaline

- Causes reflex **bradycardia**.
- It increases **BP** by constricting **blood vessels**, counteracting the bradycardic effect.

3. Vasopressin

- Similar to **noradrenaline**, increases **BP** and causes reflex **bradycardia**.

4. Thyroxin

- Increases overall **metabolism** and heart metabolism.
- **Hypothyroidism** = Low HR.

- **Hyperthyroidism** = High HR.

Q6. What is cardiac output. Mention factors affecting cardiac output. Add a note on Venous return.

Cardiac Output (CO)

- **Cardiac output** is the volume of blood ejected by each ventricle per minute.
- Normal value in **adult male**: 5-6 L/min, which is less in **females**.
- It can be calculated as:

$$\text{CO} = \text{Stroke volume} \times \text{Heart rate}$$
 Example: 70 mL x 80 bpm = **5600 mL/min**.
- **Stroke volume** is the volume of blood ejected by each ventricle per beat.

Regulation of Cardiac Output

- **Cardiac output** is regulated by three main factors:
 1. **Stroke volume**
 2. **Heart rate**
 3. **Peripheral resistance**

Cardiac Output Relation

- **Cardiac output = Stroke volume × Heart rate / Peripheral resistance**
- **End diastolic volume (EDV)** is also a key factor.
- Four main factors influence **cardiac output**:
 1. **Heart rate (chronotropic)**
 2. **Myocardial contractility (inotropic)**
 3. **Preload (EDV)**
 4. **Afterload (peripheral resistance)**

Starling's Law (1910)

- Starling studied how these factors affect **cardiac output** using a heart-lung preparation.

Stroke Volume

- Stroke volume is mainly dependent on:
 1. **Myocardial contractility**
 2. **End diastolic volume (EDV)**

Myocardial Contractility (Inotropic Effect)

- **Myocardial contractility** is influenced by both intrinsic and extrinsic factors.

Intrinsic Factors

1. Frank-Starling phenomenon

- The force of contraction is **directly proportional** to the initial length of the muscle fiber.
- This length depends on **end diastolic volume (EDV)** or **preload**.
- An increase in EDV stretches the **ventricular myocardium**, improving contractility.

2. Post-extrasystolic potentiation

- The beat after an extra systole is **stronger** due to **intracellular Ca^{++}** accumulation.
- This helps correct stroke volume on a beat-to-beat basis.

3. Force-frequency relation

- An increase in heart rate increases contractility due to accumulation of **intracellular Ca^{++}** ions.

Extrinsic Factors

1. Neural factors

- **Norepinephrine** (from sympathetic nerves) increases **cyclic AMP**, raising **intracellular Ca^{++}** , and improving myocardial contractility.
- **ACh** (from parasympathetic nerves) decreases contractility by **hyperpolarizing** the heart cells.

2. Hormonal factors

- **Epinephrine** and **norepinephrine** enhance contractility.
- **Glucagon** increases contractility by increasing **cyclic AMP**.
- **Thyroxine** increases contractility by raising metabolism.
- **Insulin** has an inotropic effect on the myocardium.
- **Cortisol** enhances the inotropic effect and works with **catecholamines**.

3. Ion effects

- **Na^{+}** and **K^{+}** decrease myocardial contractility.
- **Ca^{++}** increases contractility, with the effect proportional to **intracellular Ca^{++}** concentration.

4. Drugs

- **Beta-blockers** (e.g., **propranolol**) decrease myocardial contractility.
- **Calcium channel blockers** (e.g., **verapamil**) increase contractility.
- **Theophylline** and **caffeine** increase myocardial contractility.
- **Digitalis** enhances contractility by blocking **Na-K ATPase** .
- **Quinidine** decreases contractility by prolonging the refractory period.

- **Barbiturates** decrease excitability and contractility.

Other Factors

- **End diastolic volume (preload):**
 - **EDV** is the volume of blood in the ventricle at the end of **diastole**.
 - Stroke volume is **directly proportional** to EDV.
 - Increased **EDV** stretches the cardiac muscle fibers, boosting contractility.
 - Factors affecting **EDV**:
 1. **Venous return**
 2. **Ventricular compliance**
 3. **Diastolic pause**
 4. **Atrial systole**
 - **Venous return**: Factors increasing **venous return** also increase **EDV**.
 - **Ventricular compliance**: Decreased compliance reduces contractility.
 - **Diastolic pause**: Proportional to **EDV**.
 - **Atrial systole**: Contributes 20% to **ventricular filling** at rest, increasing during exercise.

Heart Rate (Chronotropic Effect)

- **Heart rate** is directly related to **cardiac output**.
- As **heart rate** increases, **cardiac output** increases, unless other factors limit it.
 - Example: If HR is **70/min** with a stroke volume of **70 mL/beat**, **CO = 4.9 L/min**. If HR doubles without changing stroke volume, **CO = 9.8 L/min**.
 - **Increased heart rate** also increases the force of contraction, explained by the **staircase phenomenon**.
 - **Venous return (VR)**: Volume of venous blood returning to the right atrium. **VR** matches **cardiac output**.

Peripheral Resistance

- Increased **peripheral resistance** due to **arteriolar constriction** raises aortic pressure and decreases venous return, leading to **lower cardiac output** (known as the **Anrep effect**).

Q7. What is the normal arterial blood pressure and physiological variations. Describe role of sinoaortic mechanism of regulation of blood pressure

Arterial Blood Pressure

- **Arterial blood pressure** is the lateral pressure exerted by the moving column of blood on the walls of the arteries.

- In a **normal adult**, the blood pressure is **120/80 mmHg**.
 - **Systolic blood pressure**: 120, range of **60-140 mmHg**.
 - **Diastolic blood pressure**: 80, range of **60-90 mmHg**.

Physiological Variations in Arterial Blood Pressure

Blood pressure can vary in the same individual based on the following factors:

1. **Diurnal**
 - **Blood pressure** fluctuates in a rhythmic pattern called **circadian rhythm**.
 - The **pressure** is **minimum** in the early hours and **highest** in the afternoon.
2. **Age**
 - Blood pressure varies with **age**.
 - At **birth**, systolic BP is **40-60 mmHg**.
 - On the **15th day**, systolic BP is **70 mmHg**.
 - By the **12th year**, systolic BP is **105 mmHg**.
 - At **17 years**, systolic BP is **120 mmHg**.
 - At **60 years**, systolic BP may rise to **160 mmHg**.
3. **Sex**
 - **Females** have **5 mmHg** lower blood pressure than **males** of the same age due to **sex hormones**.
 - After **menopause**, blood pressure values are the same for both sexes.
4. **Surface Area**
 - **Blood pressure** is proportional to the **surface area**.
 - **Obese people** tend to have higher blood pressure.
5. **Digestion**
 - **Food intake** and **digestion** increase **systolic BP** by about **10 mmHg** due to **sympathetic stimulation**.
 - **Diastolic BP** falls slightly due to **vasodilatation** in the **GIT**.
6. **Sleep**
 - During **quiet undisturbed sleep**, **systolic BP** may fall by about **20 mmHg** due to a **low metabolic rate**.
 - In **disturbed sleep**, BP may rise due to **cortical stimulation** of the **VMC**.
7. **Posture**
 - Blood pressure changes with **body posture**.
 - In the **standing posture**, **SBP** is low, and **DBP** is higher.
 - In the **lying posture**, the pattern is reversed.

Q8. Explain regulation of blood pressure brought about by baroreceptor mechanism.

Reflex Mechanisms in BP Regulation

The reflex mechanisms involved in regulating **blood pressure (BP)** are:

1. **Baroreceptor reflexes**
2. **Chemoreceptor reflexes**
3. **Cushing's reflex**

Baroreceptor Reflexes (Marey's Reflex)

- **Baroreceptors** are **mechanoreceptors** that respond to **stretch**.
- Key **baroreceptor reflexes**:
 1. **Carotid sinus reflex**
 2. **Aortic arch reflex**
- When **BP increases**, **baroreceptors** in the **carotid sinus** are stimulated, leading to an increase in the number of impulses through the **sinus nerve**.
 - This stimulates the **Cardioinhibitory Centre (CIC)** and **nucleus tractus solitarius (NTS)**.
 - **NTS** inhibits the **Vasomotor Centre (VMC)** via **internuncial neurons** and stimulates **CIC**.
 - **Vagal tone increases** and **sympathetic tone decreases**, leading to:
 - **Vasodilation**
 - **Decreased cardiac output**
 - **Decreased heart rate (bradycardia)**
 - **Fall in BP**

Baroreceptor reflex In response to increased BP

- **Stimulation of baroreceptors** → Increased impulses through **sinus nerve** → Stimulation of **CIC** and **NTS** → Inhibition of **VMC** → **Vagal tone increases**, **Sympathetic tone decreases** → **Vasodilatation** and **Bradycardia**
- When **BP decreases**, **baroreceptors** in the **carotid sinus** are inhibited. The net effect is:
 - **Increased peripheral resistance**
 - **Increased myocardial contractility**
 - **Increased heart rate (tachycardia)**
 - **Increased BP**

Barrow sector reflex In response to decreased BP

- **Inhibition of baroreceptors** → Decreased impulses → **Stimulation of VMC** → **Increased sympathetic tone** → **Vasoconstriction** and **Tachycardia**

Marey's Reflex (Baroreceptor Reflex)

- **Marey's law**: The **heart rate** is **inversely proportional** to **BP**.
- This mechanism is a **short-term regulatory** mechanism and is highly adaptable.
- It works within a **BP range of 60-160 mmHg**.

- In **hypertension**, the baroreceptor reflex is **reset** to a higher level to maintain BP at elevated levels.

Q9. What is the systolic diastolic and pulse pressure. Give account of factors that maintain blood pressure.

Systolic and Diastolic BP

- **Systolic BP (SBP):** Maximum BP in the arteries during **systole**.
 - Normal value: **120 ± 20 mmHg**.
- **Diastolic BP (DBP):** Minimum pressure at the end of **ventricular diastole**.
 - Normal value: **60-90 mmHg**.
- **Pulse Pressure (PP):** Difference between systolic and diastolic pressure.
 - It is an **index of cardiac output**.
 - Normal **PP: 40 mmHg**.

Factors Maintaining BP

1. **Cardiac output**
2. **Circulating blood volume**
3. **Elasticity of vessel walls**
4. **Viscosity of blood**
5. **Peripheral vascular resistance**

Cardiac Output

- **Cardiac output (CO):** Volume of blood ejected by each ventricle per minute.
 - Normal value in **adult male: 5-6 L/min**.
 - Less in **females** than males.
- Calculated from **stroke volume** and **heart rate**:
 - **CO = Stroke volume x heart rate**
 - Example: **70 x 80 = 5600 mL/min**
- **CO** raises **SBP** and also increases **DBP** by causing stretching of the aortic wall and recoil during diastole.
- **BP = CO x PR: Cardiac output** and **peripheral resistance** are directly related to BP.

Circulating Blood Volume

- The arterial and venous systems are like a **closed system**.
- Proper **blood volume** is needed for the system to exert significant pressure.
- **Low blood volume** (e.g., in **haemorrhage**) causes **BP** to fall.

- **High blood volume** (e.g., in **Cushing's syndrome**) causes **BP** to rise.
- More **blood volume** increases **SBP** and stretches vessel walls, causing a greater recoil, which raises **DBP**.

Elasticity of the Vessel Wall

- **Elasticity** of the arterial wall affects **distension** during systole and **recoil** during diastole.
- Distension minimizes the rise in **SBP**, and recoil helps in maintaining **DBP**.
- **Arteriosclerosis** (hardening of arteries) reduces **elasticity**, leading to:
 - **Increased SBP**
 - **Decreased DBP**
- In **old age**, loss of **elasticity** increases **SBP** more than **DBP**.

Viscosity of Blood

- **Viscosity** is the **thickness** of blood, and **blood** is **4.5 times more viscous** than water.
- Increased **viscosity** increases resistance, affecting **DBP** and **SBP**.
- **High viscosity** (e.g., in **polycythaemia**, **acidosis**, **dehydration**) increases BP.
- **Low viscosity** (e.g., in **anaemia**, **nephritis**) decreases BP.

Peripheral Vascular Resistance

- **PR** is the resistance blood faces due to the **blood vessels**.
- Factors affecting **PR**:
 1. **Velocity of blood**
 2. **Viscosity of blood**
 3. **Elasticity of vessel walls**
 4. **Lumen of the vessel**
 5. **Length of the vessel**
 6. **Extravascular compression**
- **PR** is inversely proportional to **radius** of the vessel.
- **Longer vessels** increase resistance, and higher **velocity** or **viscosity** increases **PR**.
- **Muscular vascular beds** (like skeletal muscle) offer maximum resistance.
- **BP = CO x PR**: Increased **PR** raises **BP**, especially **DBP**.
- **PR in healthy young adults**: 20 mmHg/L/min.

Factors Increasing Peripheral Resistance

1. **Old age**: Due to **arterial sclerosis** and **decreased nitric oxide** production.
2. **Atherosclerosis**: Due to **loss of elasticity** and **narrowing of blood vessels**.

Factors Decreasing Peripheral Resistance

1. **Beri Beri:** Due to **peripheral neuropathy**, causing vasodilatation.
2. **Hyperthyroidism:** Due to **vasodilatation**.

Q10. What are the heart sounds. How are they produced. Mention their clinical significance.

Heart Sounds

- **Definition:** Heart sounds are the noises generated by the heart during the cardiac cycle, typically heard through a stethoscope.

1. First Heart Sound (S1)

- **Production:**
 - Caused by the **closure** of the **mitral** and **tricuspid valves** at the start of ventricular systole.
- **Clinical Significance:**
 - Indicates the beginning of **systole**.
 - A **loud S1** may suggest **increased cardiac output** or **stiffened valves**.
 - A **soft S1** may indicate **heart block** or **poor ventricular contraction**.

2. Second Heart Sound (S2)

- **Production:**
 - Caused by the **closure** of the **aortic** and **pulmonary valves** at the end of ventricular systole.
- **Clinical Significance:**
 - Marks the start of **diastole**.
 - A **split S2** can be normal during **inspiration** or may indicate **right ventricular overload**.

3. Third Heart Sound (S3)

- **Production:**
 - Caused by **rapid ventricular filling** during early diastole.
- **Clinical Significance:**
 - Can be normal in **young people**.
 - In adults, it may indicate **heart failure** or **volume overload**.

4. Fourth Heart Sound (S4)

- **Production:**
 - Occurs due to **atrial contraction** against a stiff ventricle during late diastole.
- **Clinical Significance:**
 - Indicates **ventricular hypertrophy** or **decreased ventricular compliance**.

5. Murmurs

- **Definition:** Abnormal sounds caused by turbulent blood flow.
- **Clinical Significance:**
 - Can indicate **valvular diseases** (e.g., **stenosis** or **regurgitation**).