

Lecture 5A: Cointegration — Using OLS

Vitor Possebom

EESP-FGV

Econometrics 2

- Recommended Reading: Hamilton's Chapters 19.1 and 19.2
- Problem Set 4 - Deadline: June 13th at 9:00 am

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

Motivation

Motivation

Cointegration Models are useful to understand the relationship between many economic variables and test economic theories.

- Marginal Propensity to Consume: Consumption and Income
- Purchasing Power Parity: Price indexes in different countries and exchange rate
- Law of One Price
- Commodities' Spot and Future Prices

Definitions and Examples

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

Cointegrated Time Series Vector

Cointegrated Time Series Vector

An $(n \times 1)$ vector time series $\{Y_t\}$ is said to be **cointegrated** if

Cointegrated Time Series Vector

An $(n \times 1)$ vector time series $\{Y_t\}$ is said to be **cointegrated** if

1. each of the series taken individually is $I(1)$,

Cointegrated Time Series Vector

An $(n \times 1)$ vector time series $\{Y_t\}$ is said to be **cointegrated** if

1. each of the series taken individually is $I(1)$, and
2. there exist a linear combination of the series $a'Y_t$ that is $I(0)$ for some nonzero $(n \times 1)$ vector a .

Cointegrated Time Series Vector

An $(n \times 1)$ vector time series $\{Y_t\}$ is said to be **cointegrated** if

1. each of the series taken individually is $I(1)$, and
2. there exist a linear combination of the series $a'Y_t$ that is $I(0)$ for some nonzero $(n \times 1)$ vector a .

- a is called a **cointegrating vector**.

Cointegrated Time Series Vector

An $(n \times 1)$ vector time series $\{Y_t\}$ is said to be **cointegrated** if

1. each of the series taken individually is $I(1)$, and
2. there exist a linear combination of the series $a'Y_t$ that is $I(0)$ for some nonzero $(n \times 1)$ vector a .

- a is called a **cointegrating vector**.
- If $n > 2$, then there may be more than one linearly independent cointegrating vector.

Basis for the space of cointegrating vectors

Basis for the space of cointegrating vectors

Suppose that there exist h linearly independent cointegrating vectors (a_1, \dots, a_h) such that,

Basis for the space of cointegrating vectors

Suppose that there exist h linearly independent cointegrating vectors (a_1, \dots, a_h) such that, if c is a $(n \times 1)$ vector and is linearly independent from (a_1, \dots, a_h) , then

Basis for the space of cointegrating vectors

Suppose that there exist h linearly independent cointegrating vectors (a_1, \dots, a_h) such that, if c is a $(n \times 1)$ vector and is linearly independent from (a_1, \dots, a_h) , then $c'Y_t$ is a nonstationary scalar.

Basis for the space of cointegrating vectors

Suppose that there exist h linearly independent cointegrating vectors (a_1, \dots, a_h) such that, if c is a $(n \times 1)$ vector and is linearly independent from (a_1, \dots, a_h) , then $c'Y_t$ is a nonstationary scalar.

In this case, we say that there are **exactly h cointegrating relations** among the elements of $\{Y_t\}$ and

Basis for the space of cointegrating vectors

Suppose that there exist h linearly independent cointegrating vectors (a_1, \dots, a_h) such that, if c is a $(n \times 1)$ vector and is linearly independent from (a_1, \dots, a_h) , then $c'Y_t$ is a nonstationary scalar.

In this case, we say that there are exactly h cointegrating relations among the elements of $\{Y_t\}$ and that (a_1, \dots, a_h) form a basis for the space of cointegrating vectors.

We combine the cointegrating vectors in one $(n \times h)$ matrix:

$$A := \begin{bmatrix} a_1 & a_2 & \cdots & a_h \end{bmatrix}.$$

Definitions

We combine the cointegrating vectors in one $(n \times h)$ matrix:

$$A := \begin{bmatrix} a_1 & a_2 & \cdots & a_h \end{bmatrix}.$$

- $A'Y_t$ is a stationary $(h \times 1)$ vector.

Definitions

We combine the cointegrating vectors in one $(n \times h)$ matrix:

$$A := \begin{bmatrix} a_1 & a_2 & \cdots & a_h \end{bmatrix}.$$

- $A'Y_t$ is a stationary $(h \times 1)$ vector.
- For any $k \in \{1, \dots, h\}$ and any $b \in \mathbb{R} \setminus \{0\}$, the vector $\pi := b \cdot a_k$ is also a cointegrating vector.

Definitions

We combine the cointegrating vectors in one $(n \times h)$ matrix:

$$A := \begin{bmatrix} a_1 & a_2 & \cdots & a_h \end{bmatrix}.$$

- $A'Y_t$ is a stationary $(h \times 1)$ vector.
- For any $k \in \{1, \dots, h\}$ and any $b \in \mathbb{R} \setminus \{0\}$, the vector $\pi := b \cdot a_k$ is also a cointegrating vector.
- We normalize the first element in a_1 to be 1.

Numerical Example

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} =$

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} = \gamma \cdot (Y_{2,t-1} + U_{2,t}) + U_{1,t} =$

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} = \gamma \cdot (Y_{2,t-1} + U_{2,t}) + U_{1,t} = \gamma \cdot Y_{2,t-1} + U_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} = \gamma \cdot (Y_{2,t-1} + U_{2,t}) + U_{1,t} = \gamma \cdot Y_{2,t-1} + U_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$
 $Y_{1,t} = Y_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} = \gamma \cdot (Y_{2,t-1} + U_{2,t}) + U_{1,t} = \gamma \cdot Y_{2,t-1} + U_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$
 $Y_{1,t} = Y_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$
Hence, $\{Y_{1,t}\}$ is an $I(1)$ process too.

Numerical Example

Claim: The following system is a cointegrated vector process:

$$Y_{1,t} = \gamma \cdot Y_{2,t} + U_{1,t}$$

$$Y_{2,t} = Y_{2,t-1} + U_{2,t}$$

with $U_{1,t}$ and $U_{2,t}$ uncorrelated white noise processes.

Proof: Three steps.

1. $\{Y_{2,t}\}$ is an $I(1)$ process by definition.
2. $Y_{1,t} = \gamma \cdot (Y_{2,t-1} + U_{2,t}) + U_{1,t} = \gamma \cdot Y_{2,t-1} + U_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$
 $Y_{1,t} = Y_{1,t-1} - U_{1,t-1} + \gamma \cdot U_{2,t} + U_{1,t}$
Hence, $\{Y_{1,t}\}$ is an $I(1)$ process too.
3. $a' := (1, -\gamma)$ is a cointegrating vector because $Y_{1,t} - \gamma Y_{2,t} = U_{1,t}$ is an $I(0)$ process.

Cointegration means that,

Cointegration means that,

- although many developments can cause permanent changes in the individual elements of $\{Y_t\}$,

Cointegration means that,

- although many developments can cause permanent changes in the individual elements of $\{Y_t\}$,
- there is some long-run equilibrium relation tying the individual components together.

Cointegration means that,

- although many developments can cause permanent changes in the individual elements of $\{Y_t\}$,
- there is some long-run equilibrium relation tying the individual components together.
- This long-run equilibrium is represented by the linear combination $a'Y_t$.

Empirical Example 1

Empirical Example 1

Marginal Propensity to Consume:

Empirical Example 1

Marginal Propensity to Consume:

- Both consumption and income exhibit a unit root.

Empirical Example 1

Marginal Propensity to Consume:

- Both consumption and income exhibit a unit root.
- Over the long-run, consumption tends to be a roughly constant proportion of income.

Empirical Example 1

Marginal Propensity to Consume:

- Both consumption and income exhibit a unit root.
- Over the long-run, consumption tends to be a roughly constant proportion of income.
- The difference between the log of consumption and the log of income appears to be a stationary process.

Empirical Example 2

Empirical Example 2

Purchasing Power Parity:

Empirical Example 2

Purchasing Power Parity:

- Apart from transportation costs, goods should sell for the same effective price in two countries.

Empirical Example 2

Purchasing Power Parity:

- Apart from transportation costs, goods should sell for the same effective price in two countries.
- p_t = log of the price level in the U.S., p_t^* = log of the price level in Italy, and s_t = log of exchange rate are $I(1)$ processes.

Empirical Example 2

Purchasing Power Parity:

- Apart from transportation costs, goods should sell for the same effective price in two countries.
- p_t = log of the price level in the U.S., p_t^* = log of the price level in Italy, and s_t = log of exchange rate are $I(1)$ processes.
- The purchasing power parity implies that $p_t = s_t + p_t^*$.

Empirical Example 2

Purchasing Power Parity:

- Apart from transportation costs, goods should sell for the same effective price in two countries.
- p_t = log of the price level in the U.S., p_t^* = log of the price level in Italy, and s_t = log of exchange rate are $I(1)$ processes.
- The purchasing power parity implies that $p_t = s_t + p_t^*$.
- In practice, measurement error, transportation costs and quality differences prevent purchasing power parity to hold exactly.

Empirical Example 2

Purchasing Power Parity:

- Apart from transportation costs, goods should sell for the same effective price in two countries.
- p_t = log of the price level in the U.S., p_t^* = log of the price level in Italy, and s_t = log of exchange rate are $I(1)$ processes.
- The purchasing power parity implies that $p_t = s_t + p_t^*$.
- In practice, measurement error, transportation costs and quality differences prevent purchasing power parity to hold exactly.
- A weaker version of this hypothesis imposes that $Z_t := p_t - s_t - p_t^*$ is stationary.

Empirical Example 3

Zero Lower Bound (ZLM): Duffy et al. [2022]

Empirical Example 3

Zero Lower Bound (ZLM): Duffy et al. [2022]

- IS Curve + Phillips Curve + Taylor Rule

Zero Lower Bound (ZLM): Duffy et al. [2022]

- IS Curve + Phillips Curve + Taylor Rule
- Interest rate has to be positive.

Empirical Example 3

Zero Lower Bound (ZLM): Duffy et al. [2022]

- IS Curve + Phillips Curve + Taylor Rule
- Interest rate has to be positive.
- Unconventional monetary policies when the economy reaches the ZLM

Zero Lower Bound (ZLM): Duffy et al. [2022]

- IS Curve + Phillips Curve + Taylor Rule
- Interest rate has to be positive.
- Unconventional monetary policies when the economy reaches the ZLM
- Non-linear Cointegration between interest rate, unconventional policy and inflation.

Why does cointegration matter?

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate,

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate, we can analyze $\{\Delta Y_t\}$ directly through a *VAR* model.

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate, we can analyze $\{\Delta Y_t\}$ directly through a *VAR* model.
- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that cointegrate,

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate, we can analyze $\{\Delta Y_t\}$ directly through a *VAR* model.
- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that cointegrate, we need a different model

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate, we can analyze $\{\Delta Y_t\}$ directly through a *VAR* model.
- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that cointegrate, we need a **different model** because there does not exist a *VAR* representation for $\{\Delta Y_t\}$ in this case.

Why does cointegration matter?

- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that do not cointegrate, we can analyze $\{\Delta Y_t\}$ directly through a *VAR* model.
- If $\{Y_{1,t}\}, \dots, \{Y_{1,n}\}$ are $I(1)$ processes that cointegrate, we need a **different model** because there does not exist a *VAR* representation for $\{\Delta Y_t\}$ in this case.
 - Intuitively, the level of $Y_{j,t}$ contains information that is useful for forecasting $Y_{j',t}$ beyond that contained in a finite number of lagged changes in $Y_{j,t}$ alone.

Our Different Model: Vector Error Correction Model

Our Different Model: Vector Error Correction Model

Suppose that the vector process $\{Y_t\}$ can be represented as a nonstationary p -order autoregression:

$$Y_t = \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \dots + \Phi_p \cdot Y_{t-p} + \epsilon_t.$$

Our Different Model: Vector Error Correction Model

Suppose that the vector process $\{Y_t\}$ can be represented as a nonstationary p -order autoregression:

$$Y_t = \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \dots + \Phi_p \cdot Y_{t-p} + \epsilon_t.$$

We can rewrite this model as

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t,$$

where

Our Different Model: Vector Error Correction Model

Suppose that the vector process $\{Y_t\}$ can be represented as a nonstationary p -order autoregression:

$$Y_t = \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \dots + \Phi_p \cdot Y_{t-p} + \epsilon_t.$$

We can rewrite this model as

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t,$$

where

- $\rho := \Phi_1 + \Phi_2 + \dots + \Phi_p$

Our Different Model: Vector Error Correction Model

Suppose that the vector process $\{Y_t\}$ can be represented as a nonstationary p -order autoregression:

$$Y_t = \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \dots + \Phi_p \cdot Y_{t-p} + \epsilon_t.$$

We can rewrite this model as

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t,$$

where

- $\rho := \Phi_1 + \Phi_2 + \dots + \Phi_p$
- $\zeta_s := -[\Phi_{s+1} + \Phi_{s+2} + \dots + \Phi_p]$ for any $s \in \{1, 2, \dots, p-1\}$

Mathematical Details

Our Different Model: Vector Error Correction Model

We have that:

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t.$$

Our Different Model: Vector Error Correction Model

We have that:

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t.$$

Subtracting Y_{t-1} from both sides produces

Our Different Model: Vector Error Correction Model

We have that:

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t.$$

Subtracting Y_{t-1} from both sides produces

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t,$$

where

Our Different Model: Vector Error Correction Model

We have that:

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t.$$

Subtracting Y_{t-1} from both sides produces

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t,$$

where

$$\zeta_0 := \rho - I_n.$$

Our Different Model: Vector Error Correction Model

We have that:

$$Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \rho \cdot Y_{t-1} + \epsilon_t.$$

Subtracting Y_{t-1} from both sides produces

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t,$$

where

$$\zeta_0 := \rho - I_n.$$

Observation: It looks like an Augmented Dickey-Fuller test for vectors!

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t.$$

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t.$$

Unproven Lemma (See Hamilton's page 579 for a proof.)

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t.$$

Unproven Lemma (See Hamilton's page 579 for a proof.)

There exists a $(n \times h)$ matrix B such that $\zeta_0 = -BA'$, where A is a basis for the cointegrating vector space.

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha + \zeta_0 \cdot Y_{t-1} + \epsilon_t.$$

Unproven Lemma (See Hamilton's page 579 for a proof.)

There exists a $(n \times h)$ matrix B such that $\zeta_0 = -BA'$, where A is a basis for the cointegrating vector space.

We can rewrite our model as

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha - BA' \cdot Y_{t-1} + \epsilon_t.$$

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha - BA' \cdot Y_{t-1} + \epsilon_t.$$

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha - BA' \cdot Y_{t-1} + \epsilon_t.$$

Define the stationary $(h \times 1)$ vector $Z_t := A' Y_t$.

Our Different Model: Vector Error Correction Model

We have that:

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha - BA' \cdot Y_{t-1} + \epsilon_t.$$

Define the stationary $(h \times 1)$ vector $Z_t := A'Y_t$. Then, we have that

$$\Delta Y_t = \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \dots + \zeta_{p-1} \cdot \Delta Y_{t-p+1} + \alpha - BZ_{t-1} + \epsilon_t.$$

The last equation is known as **the vector error-correction representation** of the cointegrated system.

Testing the Null Hypothesis of No Cointegration

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

Testing the Null Hypothesis of No Cointegration

Testing for Cointegration when the Cointegrating Vector is Known

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

1. Test whether each of the elements of $\{Y_t\}$ is individually $I(1)$.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

1. Test whether each of the elements of $\{Y_t\}$ is individually $I(1)$.
2. If the null hypothesis of a unit root in each series is individually not rejected, construct the scalar $Z_t := a'Y_t$.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

1. Test whether each of the elements of $\{Y_t\}$ is individually $I(1)$.
2. If the null hypothesis of a unit root in each series is individually not rejected, construct the scalar $Z_t := a'Y_t$.
3. Test whether $\{Z_t\}$ is an $I(1)$ process.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

1. Test whether each of the elements of $\{Y_t\}$ is individually $I(1)$.
2. If the null hypothesis of a unit root in each series is individually not rejected, construct the scalar $Z_t := a'Y_t$.
3. Test whether $\{Z_t\}$ is an $I(1)$ process.
4. If the null hypothesis that $\{Z_t\}$ is $I(1)$ is rejected, then a is a cointegrating vector.

Testing for Cointegration when the Cointegrating Vector is Known

Sometimes, a theory not only implies that some variables should be cointegrated, but also implies a known cointegrating vector a .

- The purchasing power parity implies that $a = (1, -1, -1)'$.

In those cases, we should use this information to directly test for cointegration.

1. Test whether each of the elements of $\{Y_t\}$ is individually $I(1)$.
2. If the null hypothesis of a unit root in each series is individually not rejected, construct the scalar $Z_t := a'Y_t$.
3. Test whether $\{Z_t\}$ is an $I(1)$ process.
4. If the null hypothesis that $\{Z_t\}$ is $I(1)$ is rejected, then a is a cointegrating vector.
5. If the null hypothesis that $\{Z_t\}$ is $I(1)$ is not rejected, then a is not a cointegrating vector.

Testing for Cointegration when the Cointegrating Vector is Known

We will implement this test in Question 1 in Problem Set 4.

In this question, we will test the purchasing power parity between Brazil and the United States.

Testing the Null Hypothesis of No Cointegration

What to do when the cointegrating vector is unknown?

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

Estimating the Cointegration Vector by OLS

Sometimes, our theory suggest a vector $\{Y_t\}$ of stochastic processes is cointegrated, but does not imposes restrictions on the cointegrating vector a .

Estimating the Cointegration Vector by OLS

Sometimes, our theory suggest a vector $\{Y_t\}$ of stochastic processes is cointegrated, but does not imposes restrictions on the cointegrating vector a .

In those cases, we can try to estimate the cointegrating vector a by OLS and, then, use this estimate in our cointegration test.

Estimating the Cointegration Vector by OLS

Sometimes, our theory suggest a vector $\{Y_t\}$ of stochastic processes is cointegrated, but does not imposes restrictions on the cointegrating vector a .

In those cases, we can try to estimate the cointegrating vector a by OLS and, then, use this estimate in our cointegration test.

This approach will give you consistent estimates.

Estimating the Cointegration Vector by OLS

Sometimes, our theory suggest a vector $\{Y_t\}$ of stochastic processes is cointegrated, but does not imposes restrictions on the cointegrating vector a .

In those cases, we can try to estimate the cointegrating vector a by OLS and, then, use this estimate in our cointegration test.

This approach will give you consistent estimates. But we will learn alternative methods to estimate the cointegrating vector a .

Estimating the Cointegration Vector by OLS

The best approach will depend on your study's goal.

Estimating the Cointegration Vector by OLS

The best approach will depend on your study's goal.

- The OLS approach may be best if your goal is estimating IRFs. You will impose exclusion (or ordering) assumptions anyway.

Estimating the Cointegration Vector by OLS

The best approach will depend on your study's goal.

- The OLS approach may be best if your goal is estimating IRFs. You will impose exclusion (or ordering) assumptions anyway.
- The alternative approach may be better if your focus is on cointegration by itself (e.g., testing a theory).

Estimating the Cointegration Vector by OLS

Step-by-step procedure:

Estimating the Cointegration Vector by OLS

Step-by-step procedure:

1. Assuming that the cointegrating vector has a nonzero coefficient for the first element of Y_t ($a_1 \neq 0$), normalize $a_1 = 1$ and represent the subsequent entries as the negatives of a set of unknown parameters:

$$(a_1, a_2, \dots, a_n)' = (1, -\gamma_2, \dots, -\gamma_n)'.$$

Estimating the Cointegration Vector by OLS

Step-by-step procedure:

1. Assuming that the cointegrating vector has a nonzero coefficient for the first element of Y_t ($a_1 \neq 0$), normalize $a_1 = 1$ and represent the subsequent entries as the negatives of a set of unknown parameters:

$$(a_1, a_2, \dots, a_n)' = (1, -\gamma_2, \dots, -\gamma_n)'.$$

2. Consistent estimates of $\gamma_2, \dots, \gamma_n$ are obtained by estimating the following model by OLS:

$$Y_{1,t} = \alpha + \gamma_2 \cdot Y_{2,t} + \dots + \gamma_n \cdot Y_{n,t} + U_t. \quad (1)$$

What is happening when there is more than one cointegrating vector?

What is happening when there is more than one cointegrating vector?

When there is only one cointegrating vector, the previous procedure consistently estimates the only cointegrating vector.

What is happening when there is more than one cointegrating vector?

When there is only one cointegrating vector, the previous procedure consistently estimates the only cointegrating vector.

But what is happening when there is more than one cointegrating vector?

What is happening when there is more than one cointegrating vector?

When there is only one cointegrating vector, the previous procedure consistently estimates the only cointegrating vector.

But what is happening when there is more than one cointegrating vector?

The previous procedure consistently estimate the cointegrating vector whose residuals are uncorrelated with any $I(1)$ linear combination of $Y_{2,t}, \dots, Y_{n,t}$.

1. Motivation
2. Definitions and Examples
3. Testing the Null Hypothesis of No Cointegration
 - 3.1 Testing for Cointegration when the Cointegrating Vector is Known
 - 3.2 What to do when the cointegrating vector is unknown?
 - Estimating the Cointegration Vector by OLS
 - Testing for Cointegration when the Cointegrating Vector is Unknown

Testing for Cointegration when the Cointegrating Vector is Unknown

Now, we want to test for cointegration using the OLS-estimated cointegrating vector.

Testing for Cointegration when the Cointegrating Vector is Unknown

Now, we want to test for cointegration using the OLS-estimated cointegrating vector.

Our null hypothesis is that there is no cointegrating relation among our variables Y_t .

Testing for Cointegration when the Cointegrating Vector is Unknown

Now, we want to test for cointegration using the OLS-estimated cointegrating vector.

Our null hypothesis is that there is no cointegrating relation among our variables Y_t .

We will estimate the OLS regression (1) and use its residuals to test for our null hypothesis of no cointegration.

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

The Phillips-Ouliaris-Hansen explore this property of the OLS residuals as a test for cointegration.

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

The Phillips-Ouliaris-Hansen explore this property of the OLS residuals as a test for cointegration.

1. We estimate Equation (1) and save the estimated residuals \hat{U}_t .

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

The Phillips-Ouliaris-Hansen explore this property of the OLS residuals as a test for cointegration.

1. We estimate Equation (1) and save the estimated residuals \hat{U}_t .
2. We test whether \hat{U}_t is a $I(1)$ process using the Augmented Dickey-Fuller test.

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

The Phillips-Ouliaris-Hansen explore this property of the OLS residuals as a test for cointegration.

1. We estimate Equation (1) and save the estimated residuals \hat{U}_t .
2. We test whether \hat{U}_t is a $I(1)$ process using the Augmented Dickey-Fuller test.
3. The critical values of this test are different from the standard critical values of the Augmented Dickey-Fuller test.

Testing for Cointegration when the Cointegrating Vector is Unknown

Under the null of no cointegration, Equation (1) is a regression of an $I(1)$ variable on a set of $I(1)$ variables for which no coefficients produce an $I(0)$ error term.

The Phillips-Ouliaris-Hansen explore this property of the OLS residuals as a test for cointegration.

1. We estimate Equation (1) and save the estimated residuals \hat{U}_t .
2. We test whether \hat{U}_t is a $I(1)$ process using the Augmented Dickey-Fuller test.
3. The critical values of this test are different from the standard critical values of the Augmented Dickey-Fuller test.
4. The null of our test is that \hat{U}_t is a $I(1)$ process, i.e., there is no cointegrating relation among our variables.

Testing for Cointegration when the Cointegrating Vector is Unknown

The **Phillips-Ouliaris-Hansen** test uses the Augmented Dickey-Fuller test with no constant term,

$$\hat{U}_t = \rho \cdot \hat{U}_{t-1} + \sum_{i=2}^p \zeta_{i-1} \cdot \Delta \hat{U}_{t-i+1} + V_t,$$

and tests the null hypothesis that

$$H_0 : \rho = 1.$$

Testing for Cointegration when the Cointegrating Vector is Unknown

The **Phillips-Ouliaris-Hansen** test uses the Augmented Dickey-Fuller test with no constant term,

$$\hat{U}_t = \rho \cdot \hat{U}_{t-1} + \sum_{i=2}^p \zeta_{i-1} \cdot \Delta \hat{U}_{t-i+1} + V_t,$$

and tests the null hypothesis that

$$H_0 : \rho = 1.$$

Critical values for this test depend on whether the original stochastic processes $\{Y_t\}$ had a time trend or not.

Phillips-Ouliaris-Hansen Critical Values: No Time Trend

Phillips-Ouliaris-Hansen Critical Values: No Time Trend

TABLE B.9

Critical Values for the Phillips Z_t Statistic or the Dickey-Fuller t Statistic When Applied to Residuals from Spurious Cointegrating Regression

Number of right-hand variables in regression, excluding trend or constant ($n - 1$)	Sample size (T)	Probability that $(\hat{\rho} - 1)/\hat{\sigma}_{\hat{\rho}}$ is less than entry						
		0.010	0.025	0.050	0.075	0.100	0.125	0.150
Case 2								
1	500	-3.96	-3.64	-3.37	-3.20	-3.07	-2.96	-2.86
2	500	-4.31	-4.02	-3.77	-3.58	-3.45	-3.35	-3.26
3	500	-4.73	-4.37	-4.11	-3.96	-3.83	-3.73	-3.65
4	500	-5.07	-4.71	-4.45	-4.29	-4.16	-4.05	-3.96
5	500	-5.28	-4.98	-4.71	-4.56	-4.43	-4.33	-4.24

Phillips-Ouliaris-Hansen Critical Values: With Time Trend

Phillips-Ouliaris-Hansen Critical Values: With Time Trend

TABLE B.9

Critical Values for the Phillips Z_t Statistic or the Dickey-Fuller t Statistic When Applied to Residuals from Spurious Cointegrating Regression

Number of right-hand variables in regression, excluding trend or constant ($n - 1$)	Sample size (T)	Probability that $(\hat{\rho} - 1)/\hat{\sigma}_{\hat{\rho}}$ is less than entry						
		0.010	0.025	0.050	0.075	0.100	0.125	0.150
Case 3								
1	500	-3.98	-3.68	-3.42	—	-3.13	—	—
2	500	-4.36	-4.07	-3.80	-3.65	-3.52	-3.42	-3.33
3	500	-4.65	-4.39	-4.16	-3.98	-3.84	-3.74	-3.66
4	500	-5.04	-4.77	-4.49	-4.32	-4.20	-4.08	-4.00
5	500	-5.36	-5.02	-4.74	-4.58	-4.46	-4.36	-4.28

Testing for Cointegration when the Cointegrating Vector is Unknown

We implement the **Phillips-Ouliaris-Hansen** in Question 2 in Problem Set 4.

In this question, we will test whether the following variables cointegrate: Brazilian IPCA, American CPI and the R\$/US\$ exchange rate.

Phillips-Ouliaris-Hansen: Drawbacks

Phillips-Ouliaris-Hansen: Drawbacks

The Phillips-Ouliaris-Hansen test can give different answers depending on which variable is labeled Y_1 .

In the next lecture, we will learn a test for cointegration that is invariant to the ordering of our variables.

This procedure is based on a full-information maximum likelihood test and is known as Johansen's test.

Thank you!

Contact Information:

Vitor Possebom

E-mail: vitor.possebom@fgv.br

Website: sites.google.com/site/vitorapossebom/

References

J. A. Duffy, S. Mavroeidis, and S. Wycherley. Cointegration with Occasionally Binding Constraints. Available at <https://arxiv.org/abs/2211.09604>., Nov. 2022.

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$.

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$Y_t = \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \underbrace{(\Phi_1 + \Phi_2 + \Phi_3)}_{=: \rho} \cdot Y_{t-1} + \underbrace{(-\Phi_2) \cdot (Y_{t-1} - Y_{t-2})}_{=: \Delta Y_{t-1}} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned}Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\&= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\&\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\&= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\&= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-2} \\&\quad - \Phi_3 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t\end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \Phi_1 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-1} - \Phi_2 \cdot Y_{t-1} + \Phi_2 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} - \Phi_3 \cdot Y_{t-1} + \Phi_3 \cdot Y_{t-2} \\ &\quad - \Phi_3 \cdot Y_{t-2} + \Phi_3 \cdot Y_{t-3} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \underbrace{(Y_{t-1} - Y_{t-2})}_{=\Delta Y_{t-1}} \\ &\quad + (-\Phi_3) \cdot \underbrace{(Y_{t-2} - Y_{t-3})}_{=\Delta Y_{t-2}} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$Y_t = \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-2} + \epsilon_t$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-2} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + \underbrace{(-(\Phi_2 + \Phi_3))}_{=:\zeta_1} \cdot \Delta Y_{t-1} + \underbrace{(-\Phi_3)}_{=:\zeta_2} \cdot \Delta Y_{t-2} + \epsilon_t \end{aligned}$$

Mathematical Details

We will show that the former representation is valid for a $VAR(3)$. We know that

$$\begin{aligned} Y_t &= \alpha + \rho \cdot Y_{t-1} + (-\Phi_2) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-1} + (-\Phi_3) \cdot \Delta Y_{t-2} + \epsilon_t \\ &= \alpha + \rho \cdot Y_{t-1} + \zeta_1 \cdot \Delta Y_{t-1} + \zeta_2 \cdot \Delta Y_{t-2} + \epsilon_t \end{aligned}$$

[Back](#)