Term	OpenMM class
Energy of one summand (Hwang et al. 1994, Eq. 1)	
bond	`CustomBondForce`
${}^2K_b(b-b_0)^2 + {}^3K_b(b-b_0)^3 + {}^4K_b(b-b_0)^4$	
angle	`CustomAngleForce`
${}^2K_\theta(\theta-\theta_0)^2 + {}^3K_\theta(\theta-\theta_0)^3 + {}^4K_\theta(\theta-\theta_0)^4$	
torsion	`CustomTorsionForce`
${}^1K_\phi(1-\cos\phi) + {}^2K_\phi(1-\cos 2\phi) + {}^3K_\phi(1-\cos 3\phi)$	
bond–bond	`CustomCompoundBondForce (3 atoms)`
$K_{bb'}(b-b_0)(b'-b'_0)$	
bond–angle	`CustomCompoundBondForce (3 atoms)`
$K_{b\theta}(b-b_0)(\theta-\theta_0)$	
angle–angle	`CustomCompoundBondForce (4 atoms)`
$K_{\theta\theta'}(\theta-\theta_0)(\theta'-\theta'_0)$	
bond–torsion	`CustomCompoundBondForce (4 atoms)`
$(b-b_0)[\ {}^1K_{\phi b}\cos\phi + {}^2K_{\phi b}\cos 2\phi + {}^3K_{\phi b}\cos 3\phi\ ]$	
angle–torsion	`CustomCompoundBondForce (4 atoms)`
$(\theta-\theta_0)[\ {}^1K_{\phi\theta}\cos\phi + {}^2K_{\phi\theta}\cos 2\phi + {}^3K_{\phi\theta}\cos 3\phi\ ]$	
angle–angle–torsion	`CustomCompoundBondForce (4 atoms)`
$K_{\phi\theta\theta'}(\theta-\theta_0)(\theta'-\theta'_0)\cos\phi$	