Term	Energy of one summand (Hwang et al. 1994, Eq. 1)	OpenMM
bond	${}^2K_b(b-b_0)^2 + {}^3K_b(b-b_0)^3 + {}^4K_b(b-b_0)^4$	`CustomBondForce`
angle	${}^2K_\theta(\theta-\theta_0)^2 + {}^3K_\theta(\theta-\theta_0)^3 + {}^4K_\theta(\theta-\theta_0)^4$	`CustomAngleForce`
torsion	${}^1K_\phi(1-\cos\phi) + {}^2K_\phi(1-\cos 2\phi) + {}^3K_\phi(1-\cos 3\phi)$	`CustomTorsionForce`
bond–bond	$K_{bb'}(b-b_0)(b'-b'_0)$	`CustomCompoundBondForce (3 atoms)`
bond–angle	$K_{b\theta}(b-b_0)(\theta-\theta_0)$	`CustomCompoundBondForce (3 atoms)`
angle–angle	$K_{\theta\theta'}(\theta-\theta_0)(\theta'-\theta'_0)$	`CustomCompoundBondForce (4 atoms)`
bond–torsion	$(b-b_0)\left[{}^1K_{\phi b}\cos\phi + {}^2K_{\phi b}\cos 2\phi + {}^3K_{\phi b}\cos 3\phi\right]$	`CustomCompoundBondForce (4 atoms)`
angle–torsion	$(\theta-\theta_0)\left[{}^1K_{\phi\theta}\cos\phi + {}^2K_{\phi\theta}\cos 2\phi + {}^3K_{\phi\theta}\cos 3\phi\right]$	`CustomCompoundBondForce (4 atoms)`
angle–angle–torsion	$K_{\phi\theta\theta'}(\theta-\theta_0)(\theta'-\theta'_0)\cos\phi$	`CustomCompoundBondForce (4 atoms)`