Problem 001: blacking out a grid so that every rectangle can still be found

Notes on everything done so far

Arjun Pemmasani

October 2026

These are working notes on Problem 001 of Peter Kagey's Open Problem Collection [1]. They go through every result so far, slowly, with pictures and small examples. The proofs are complete unless a result is marked as computer-assisted, and in that case the notes say exactly what the computer checked. The code is in the repository `github.com/apemm/Kagey-Problems`, folder `problem001`.

grid	maximum blackout	status
$2 \times m$, $m \geq 3$	$m+1$, with $m \cdot 2^{m-1}$ optima	proved (Section 3)
$3 \times m$, $m \geq 3$	$m+2$	proved (Section 7)
4×4, 4×5	7, 8	proved, finite computer check (Section 8.2)
$4 \times m$, $m \geq 6$	$m+4$	proved (Section 8)
5×5	10, with exactly 8 optima	value proved, count by computer (Section 9)
$n \times m$, $n, m \geq 3$	at most $m + \lfloor n^2/4 \rfloor$	proved (Section 5)
$n, m \geq 5$, up to 10×10	at least $n+m$	by computer (Section 10)

Contents

1 The game **2**

1.1 The rules . . . 2

1.2 Conventions . . . 3

1.3 What the player sees . . . 4

1.4 A first example: the 2×3 strip . . . 4

2 From the game to a hitting set **5**

2.1 When do 2 rectangles look the same? . . . 5

2.2 Pruning . . . 6

3 Two rows: the Strip Theorem **7**

4 The row–column graph **8**

4.1 The graph of a blackout . . . 8

4.2 Three corners are enough . . . 8

4.3 Short cycles . . . 9

5 An upper bound for every grid **10**
5.1 Building the graph H . . . 10
5.2 Mantel's theorem . . . 11

6 Tilted rectangles on 3 and 4 rows **12**

7 Three rows **13**

8 Four rows **14**
8.1 The construction for $m \geq 6$ . . . 14
8.2 Widths 4 and 5 . . . 16

9 The 5×5 grid **17**
9.1 The upper bound . . . 17
9.2 What the optimum looks like . . . 19

10 Bigger grids **20**

11 Computations **20**
11.1 The programs . . . 20
11.2 The values . . . 21
11.3 Predictions that failed . . . 21

12 Open questions **22**


1 The game

1.1 The rules

The problem is posed here with some changes chosen for clarity. Suppose you, the player, are shown an $n \times m$ grid of points. Each point is either white or blacked out, and a blacked-out point can't tell you anything. Before the game starts you choose which points to black out. We call the set of blacked-out points the *blackout*. Then the game master draws a rectangle with its 4 corners on grid points. You can't see the rectangle. You are only told which of its corners are white. The rectangle doesn't have to line up with the grid: tilted rectangles count too. You must be able to say where all 4 corners are with complete confidence, whatever rectangle the game master draws. Kagey's question is how many points you can black out and still do this.

The order of play matters. The blackout is fixed before the rectangle is drawn, so a blackout is only good if *no* 2 rectangles could ever look the same to you.

Example 1.1. Figure 1 plays one round. In (b) we black out all of row 1 and all of column 1, which is 6 points. In (c) the game master draws the tilted square with corners $(2, 2)$, $(3, 1)$, $(4, 2)$ and $(3, 3)$. The corner $(3, 1)$ is in row 1, so it is black, and in (d) we are only shown $(2, 2)$, $(4, 2)$ and $(3, 3)$. That is enough. Three corners of a rectangle always determine the fourth (Lemma 4.2



(a) the point $(3, 2)$ is ringed (b) our blackout (c) the rectangle (d) what we are shown

Figure 1: One round on the 3×4 grid. Black dots are blacked out and open dots are white. In (d) the grey specks are points we are told nothing about.


below), and here the fourth corner has to be $(3, 1)$. We will see in Section 7 that this blackout works against every rectangle on the 3×4 grid, and that 6 is the most anyone can black out there.

1.2 Conventions

We use these conventions all the way through.

1. An $n \times m$ grid has n rows and m columns, so nm points. A $2 \times m$ grid is called a *strip*.
2. The point (i, j) is in column i and row j, and $(1, 1)$ is the top-left point. So i grows to the right and j grows downwards, as in Figure 1(a). The set of all points is V.
3. A *rectangle* is any rectangle whose 4 corners are grid points, tilted ones included. It must have positive area. We write $C(R)$ for the set of corners of R. Two different rectangles have different corner sets, so we can think of a rectangle as its set of 4 corners.
4. A *blackout* is a set $S \subseteq V$, and the *kept set* is $K = V \setminus S$, the points left white.

How do we tell whether 4 points form a rectangle? Pair them up into 2 "diagonals". The 4 points form a parallelogram exactly when the 2 diagonals have the same midpoint, and the parallelogram is a rectangle exactly when the 2 diagonals also have the same length. So 4 points are a rectangle if and only if some pairing gives 2 segments with the same midpoint and the same length. The verification script finds all rectangles this way: it sorts all pairs of points by midpoint and length and takes 2 pairs from the same class.



(a) axis-aligned (b) a tilted square (c) tilted, sides $1 : 2$ (d) not a rectangle

Figure 2: Which 4-point sets are rectangles, on the 4×4 grid. In (c) and (d) the dashed lines are the diagonals.

Example 1.2. In Figure 2(c) the diagonals join $(1, 2)$ to $(4, 3)$ and $(2, 1)$ to $(3, 4)$. Both have midpoint $(\frac{5}{2}, \frac{5}{2})$, and both have squared length $3^2 + 1^2 = 1^2 + 3^2 = 10$, so the 4 points are a rectangle, with sides of length $\sqrt{2}$ and $2\sqrt{2}$. In (d) the diagonals join $(1, 1)$ to $(4, 3)$ and $(3, 1)$ to $(2, 3)$. They

have the same midpoint $(\frac{5}{2}, 2)$, so the shape is a parallelogram, but the squared lengths are 13 and 5, so it is not a rectangle.

The number of rectangles grows quickly. The 3×3 grid has 10 (9 axis-aligned and 1 tilted square), the 4×4 grid has 44 (36 and 8), and the 5×5 grid has 130 (100 and 30). The counts on square grids, $1, 10, 44, 130, 313, \ldots$, are OEIS A085582 [2], and the counts on $n \times m$ grids are in OEIS A289832 [3]. Our counts agree with both everywhere we computed them.

1.3 What the player sees

For a blackout S and a rectangle R, what we are shown is the set of white corners,

$$P_S(R) = C(R) \setminus S.$$

We call $P_S(R)$ the *presentation* of R.

Definition 1.3. A blackout S is *valid* if different rectangles always have different presentations, that is, if $P_S(R_1) \neq P_S(R_2)$ whenever $R_1 \neq R_2$. The *maximum blackout* of a grid is the largest size of a valid blackout.

The empty set is allowed as a presentation. It just means every corner was black. We don't need a separate rule about it.

Remark 1.4. Suppose 2 distinct rectangles $R_1 \neq R_2$ are both fully hidden, that is, $C(R_1) \subseteq S$ and $C(R_2) \subseteq S$. Then $P_S(R_1) = \emptyset = P_S(R_2)$, so S is not valid. Hence at most 1 rectangle can be fully hidden under a valid blackout. This follows from the definition; it isn't an extra rule.

1.4 A first example: the 2×3 strip

On 2 rows every rectangle is axis-aligned, because a tilted rectangle has its 4 corners in at least 3 different rows. An axis-aligned rectangle on a strip uses both rows and 2 columns $a < b$, and we call it $R_{a,b}$. The 2×3 strip has the 3 rectangles of Figure 3.




Figure 3: The 3 rectangles of the 2×3 strip.



(a) valid (b) not valid

Figure 4: Two blackouts of 4 points on the 2×3 strip. In (b) the rectangles $R_{1,3}$ (solid) and $R_{2,3}$ (dashed) both show exactly the ringed points.

Example 1.5. Figure 4 shows 2 blackouts with 4 points each. Here are the presentations:

rectangle	blackout (a)	blackout (b)
$R_{1,2}$	$\{(1,1),(2,1)\}$	$\emptyset$
$R_{1,3}$	$\{(1,1)\}$	$\{(3,1),(3,2)\}$
$R_{2,3}$	$\{(2,1)\}$	$\{(3,1),(3,2)\}$

In (a) the 3 presentations are different, so (a) is valid. In (b) the rectangles $R_{1,3}$ and $R_{2,3}$ look the same. If we are shown $(3,1)$ and $(3,2)$ we can't tell whether the left side is in column 1 or column 2, so (b) is not valid. Both blackouts have 4 points, so the size alone doesn't decide anything; where the points go matters.

Can we black out 5 points? Then only 1 point is kept, and every presentation is either empty or that 1 point. That is 2 possible presentations for 3 rectangles, so 2 of them must agree. So the maximum blackout of the 2×3 strip is 4.

2 From the game to a hitting set

2.1 When do 2 rectangles look the same?

Lemma 2.1 (difference sets)**.** *For any sets A, B, S: $A \setminus S = B \setminus S$ if and only if $A \triangle B \subseteq S$.*

Here $A \triangle B = (A \setminus B) \cup (B \setminus A)$ is the set of points in exactly 1 of A and B. In words: 2 rectangles look the same exactly when every corner that belongs to only 1 of them is black. The corners they share don't matter, since they are shown or hidden for both.

Proof. ($\Rightarrow$) Assume $A \setminus S = B \setminus S$, and let $x \in A \triangle B$ be arbitrary. The statement is symmetric in A and B, so we may assume $x \in A \setminus B$. Suppose $x \notin S$. Then $x \in A \setminus S = B \setminus S$, so $x \in B$, which contradicts $x \in A \setminus B$. Therefore $x \in S$. As x was arbitrary, $A \triangle B \subseteq S$.

($\Leftarrow$) Assume $A \triangle B \subseteq S$. By symmetry it is enough to prove $A \setminus S \subseteq B \setminus S$. Let $x \in A \setminus S$ be arbitrary, so $x \in A$ and $x \notin S$. Suppose $x \notin B$. Then $x \in A \setminus B \subseteq A \triangle B \subseteq S$, which contradicts $x \notin S$. Hence $x \in B$, and with $x \notin S$ we get $x \in B \setminus S$. □



(a) $R_{1,2}$ and $R_{1,3}$ (b) $R_{1,2}$ and $R_{3,4}$

Figure 5: Difference sets on the 2×4 strip. The shaded points are $C(R_1) \triangle C(R_2)$ for the solid and the dashed rectangle.

Example 2.2. In Figure 5(a), $R_{1,2}$ and $R_{1,3}$ share column 1. Their corners in column 1 cancel, and the difference set is the 4 points of columns 2 and 3. By Lemma 2.1, the 2 rectangles look the same exactly when all 4 shaded points are black. To keep them apart we must keep at least 1 of those 4 points. In (b) the rectangles $R_{1,2}$ and $R_{3,4}$ share nothing, and the difference set is all 8 points.

Applying the lemma to every pair of rectangles gives the following.

Corollary 2.3. *A blackout S is valid if and only if the kept set $K = V \setminus S$ contains at least 1 point of $C(R_1) \triangle C(R_2)$ for every pair of distinct rectangles R_1, R_2.*

Proof. S is valid if and only if $C(R_1) \setminus S \neq C(R_2) \setminus S$ for every pair of distinct rectangles. By Lemma 2.1 this holds if and only if $C(R_1) \triangle C(R_2) \not\subseteq S$ for every pair, which says exactly that some point of the difference set lies in K. □

So the problem is really about the *difference hypergraph*: the collection of all difference sets $C(R_1) \triangle C(R_2)$. A set that meets every difference set is a *hitting set*, and

$$\text{maximum blackout} = nm - (\text{smallest size of a hitting set}).$$

Small blackouts are easy; the work is in keeping very few points and still meeting every difference set. The computer program in Section 11 solves exactly this hitting set problem.

2.2 Pruning

Before computing anything we shrink the list of difference sets. Think of it as a table with one line for each pair of rectangles. To avoid confusing these lines with rows of the grid, we call them *constraints*.

1. *Repeats.* Two identical constraints say the same thing, so we keep 1 copy.
2. *Containment.* If $D_1 \subsetneq D_2$, any K that meets D_1 also meets D_2, so D_2 can be deleted without changing which kept sets are valid.
3. *No empty constraints.* $C(R_1) \triangle C(R_2) = \emptyset$ would mean $C(R_1) = C(R_2)$, and different rectangles have different corners, so this never happens.

After pruning we are left with the minimal constraints, and a set K meets every constraint if and only if it meets every minimal one.

Example 2.4. The 2×4 strip has 6 rectangles, so 15 pairs. Every rectangle is made of whole columns, so each difference set is a set of whole columns too. Here is the full table, listing the columns in each difference set.

pair	columns	points	after pruning
$R_{1,2}$, $R_{1,3}$	$\{2,3\}$	4	kept
$R_{1,2}$, $R_{1,4}$	$\{2,4\}$	4	kept
$R_{1,2}$, $R_{2,3}$	$\{1,3\}$	4	kept
$R_{1,2}$, $R_{2,4}$	$\{1,4\}$	4	kept
$R_{1,2}$, $R_{3,4}$	$\{1,2,3,4\}$	8	deleted, contains columns $\{1,2\}$
$R_{1,3}$, $R_{1,4}$	$\{3,4\}$	4	kept
$R_{1,3}$, $R_{2,3}$	$\{1,2\}$	4	kept
$R_{1,3}$, $R_{2,4}$	$\{1,2,3,4\}$	8	deleted, contains columns $\{1,2\}$
$R_{1,3}$, $R_{3,4}$	$\{1,4\}$	4	repeat
$R_{1,4}$, $R_{2,3}$	$\{1,2,3,4\}$	8	deleted, contains columns $\{1,2\}$
$R_{1,4}$, $R_{2,4}$	$\{1,2\}$	4	repeat
$R_{1,4}$, $R_{3,4}$	$\{1,3\}$	4	repeat
$R_{2,3}$, $R_{2,4}$	$\{3,4\}$	4	repeat
$R_{2,3}$, $R_{3,4}$	$\{2,4\}$	4	repeat
$R_{2,4}$, $R_{3,4}$	$\{2,3\}$	4	repeat

Of the 15 lines, 6 survive, one for each pair of columns. So a kept set is valid on the 2×4 strip if and only if, for every 2 columns, it keeps a point in at least 1 of them.

3 Two rows: the Strip Theorem

Example 2.4 already contains the whole idea for strips.

Theorem 3.1 (Strip Theorem)**.** *Let $m \geq 3$. The maximum blackout of the $2 \times m$ strip is $m+1$. There are exactly $m \cdot 2^{m-1}$ maximum blackouts: black out both points of 1 column, and exactly 1 of the 2 points of every other column. For $m = 2$ there is only 1 rectangle, so every blackout is valid and the maximum is 4.*

Proof. Step 1: the constraints. Take 2 distinct rectangles $R_{a,b}$ and $R_{c,d}$. They can't share both columns. If they share exactly 1 column, the shared column cancels and the difference set is the 4 points of the other 2 columns p, q. We call this the *pair constraint* on $\{p, q\}$. If they share no column, the difference set is all 8 points of 4 columns, and it contains a pair constraint, so pruning deletes it. Every pair $\{p, q\}$ does occur: since $m \geq 3$ there is a third column z, and $R_{z,p}$, $R_{z,q}$ (with the columns put in order) have difference set exactly the pair constraint on $\{p, q\}$. So after pruning the constraints are exactly the $\binom{m}{2}$ pair constraints.

Step 2: at most 1 empty column. Let K be a valid kept set and call a column empty if K has no point in it. If 2 columns $p \neq q$ were empty, K would miss the pair constraint on $\{p, q\}$. So at most 1 column is empty.

Step 3: the bound. Every non-empty column has at least 1 kept point, so $|K| \geq m - 1$ and $|S| = 2m - |K| \leq m + 1$. If equality holds then $|K| = m - 1$ exactly, so exactly 1 column is empty (if none were, $|K| \geq m$) and every other column has exactly 1 kept point.

Step 4: these all work. Take any K with 1 empty column and exactly 1 point in every other column. Any 2 columns include at least 1 non-empty one, whose kept point meets the pair constraint. So K is valid by Corollary 2.3, and $S = V \setminus K$ has $m + 1$ points. There are m choices for the empty column and 2 choices in each of the other $m - 1$ columns, so $m \cdot 2^{m-1}$ maximum blackouts. □




Figure 6: A maximum blackout of the 2×6 strip: column 4 (shaded) is fully black and every other column has 1 black point. It has $m + 1 = 7$ points.

Example 3.2. For $m = 3$ the theorem says the maximum is 4 with 12 optima, matching Example 1.5, and Figure 7 shows all 12 (found by the computer, by checking all 15 blackouts of size 4). For $m = 4$ it gives 5 and $4 \cdot 8 = 32$, and for $m = 5$ it gives 6 and 80. The prediction of 80 for the 2×5 strip was written down before the program first ran on that grid, and the program found 80.




Figure 7: All $3 \cdot 2^2 = 12$ maximum blackouts of the 2×3 strip, grouped by the column that is fully black. Blackout (a) of Figure 4 is one of the 4 on the right.

4 The row–column graph


From 3 rows on, there are both axis-aligned and tilted rectangles. The axis-aligned ones are handled by a graph, and this section sets it up.

4.1 The graph of a blackout

Think of a black point (i, j) as a link between row j and column i.

Definition 4.1. The *row–column graph* G_S of a blackout S has a vertex r_j for each row and a vertex c_i for each column, and an edge $r_j c_i$ for each black point $(i, j) \in S$.

So G_S is bipartite (every edge joins a row to a column), it has $n + m$ vertices, and it has $|S|$ edges. We draw rows as squares and columns as circles.



(a) the blackout (b) its graph G_S

Figure 8: The blackout of Example 1.1 and its graph. Row 1 is black in every column, and column 1 is black in every row.

An axis-aligned rectangle uses 2 rows j, j' and 2 columns i, i', and its 4 corners are the 4 possible edges between them. If all 4 corners are black, these 4 edges form a cycle $r_j\, c_i\, r_{j'}\, c_{i'}$ of length 4 in G_S. The main lemma of this section says that cycles of length 4 and 6 are exactly what makes axis-aligned rectangles look alike.

4.2 Three corners are enough

Lemma 4.2 (three corners suffice)**.** *Three corners of a rectangle determine the rectangle. So if 2 rectangles have the same presentation and it has at least 3 points, the 2 rectangles are equal.*

Proof. Let p, q, r be 3 corners of a rectangle. Exactly 1 of them, say q, is next to both of the others, and the angle at q in the triangle pqr is a right angle. A triangle has at most 1 right angle, so the 3

points alone tell us which one is q. The fourth corner is then $p + r - q$. For the second claim, a presentation with at least 3 points contains 3 corners of each rectangle, the same 3 points, so the 2 rectangles have the same corners. □





Figure 9: The right angle is at q, so the missing corner is $p + r - q$.

So 2 rectangles can only be confused if both show at most 2 corners. When we check a blackout, we only need to look at rectangles with at least 2 black corners.

4.3 Short cycles

Lemma 4.3 (axis-aligned rectangles and short cycles)**.** *Suppose $m \geq 3$. No 2 axis-aligned rectangles have the same presentation if and only if G_S has no cycle of length 4 and no cycle of length 6.*

Before the proof, here are the 2 bad cases.




(a) a black 4-cycle (b) G_S

Figure 10: A black 4-cycle: rows 1, 2 and columns 1, 2 are all black. The rectangles $R_{1,3}$ and $R_{2,3}$ both show the 2 points of column 3.

Example 4.4 (4-cycles)**.** The bad blackout (b) of Example 1.5 is a black 4-cycle (Figure 10). The rectangles $R_{1,3}$ and $R_{2,3}$ agree in column 3, and their other columns are fully black, so they look the same. The same thing happens in any grid with a black 4-cycle on rows j, j' and columns i, i': pick any third column w, and the 2 rectangles on rows j, j' with columns $\{i, w\}$ and $\{i', w\}$ both show only what is white in column w. This is where $m \geq 3$ is used.

Example 4.5 (6-cycles)**.** In Figure 11 the black points $(1,1), (1,2), (2,2), (2,3), (3,3), (3,1)$ form the cycle $r_1 c_1 r_2 c_2 r_3 c_3$ in G_S. The rectangle on rows $\{1,2\}$ and columns $\{1,2\}$ has corners $(1,1), (2,1), (1,2), (2,2)$, and only $(2,1)$ is white. The rectangle on rows $\{1,3\}$ and columns $\{2,3\}$ has corners $(2,1), (3,1), (2,3), (3,3)$, and again only $(2,1)$ is white. So both show just $(2,1)$.

Proof of Lemma 4.3. The 2 examples show the "only if" direction in general. For a 4-cycle, see the example above. For a 6-cycle $r_a\, c_x\, r_b\, c_y\, r_c\, c_z$, the black points are $(x,a), (x,b), (y,b), (y,c), (z,c), (z,a)$. The rectangle on rows $\{a,b\}$ and columns $\{x,y\}$ and the rectangle on rows $\{a,c\}$ and columns $\{y,z\}$



(a) a black 6-cycle (b) G_S

Figure 11: A black 6-cycle on the 3×3 grid. The solid and the dashed rectangle both show only the ringed point $(2, 1)$.

are different, and every corner of each is black except possibly the shared corner (y, a). So they have the same presentation.

For the "if" direction, assume G_S has no 4-cycle and no 6-cycle. Let $R_1 \neq R_2$ be axis-aligned rectangles with the same presentation P. We get a contradiction in each case.

- $|P| \geq 3$: Lemma 4.2 gives $R_1 = R_2$.
- $|P| = 0$: all 4 corners of R_1 are black, and they form a 4-cycle.
- $|P| = 2$, the 2 points in different rows and different columns: then they are opposite corners of both rectangles, and 2 opposite corners determine an axis-aligned rectangle, so $R_1 = R_2$.
- $|P| = 2$, the 2 points in 1 row a, at columns x and y: then R_1 uses rows $\{a, b\}$ and R_2 uses rows $\{a, c\}$ with $b \neq c$, and the points $(x, b), (y, b), (x, c), (y, c)$ are all black. That is the 4-cycle $r_b\, c_x\, r_c\, c_y$. The case of 1 column is the same with rows and columns swapped.
- $|P| = 1$, say $P = \{(x, a)\}$: write R_1 as rows $\{a, b\}$, columns $\{x, y\}$ and R_2 as rows $\{a, c\}$, columns $\{x, z\}$. The black points are $(y, a), (x, b), (y, b)$ and $(z, a), (x, c), (z, c)$. If $b = c$ then $y \neq z$, and rows a, b with columns y, z are all black, a 4-cycle. If $y = z$, then in the same way rows b, c with columns x, y give a 4-cycle. Otherwise the 6 black points form the 6-cycle $r_a\, c_y\, r_b\, c_x\, r_c\, c_z$.

Every case is impossible, so R_1 and R_2 can't have the same presentation. □

Note that the lemma is only about axis-aligned rectangles against each other. A tilted rectangle could still be confused with something, and we deal with that separately for each grid.

5 An upper bound for every grid


Theorem 5.1 (row bound)**.** *Let $m \geq 3$. Every valid blackout of an $n \times m$ grid has at most $m + \lfloor n^2/4 \rfloor$ points. If also $n \geq 3$, then by swapping rows and columns it has at most $n + \lfloor m^2/4 \rfloor$ points.*

The plan is to turn G_S into a graph H on the rows alone, with no triangles, and then use Mantel's theorem, which says a triangle-free graph on n vertices has at most $n^2/4$ edges.

5.1 Building the graph H

For each column i let B_i be the set of rows where column i is black. If B_i is not empty, join the rows in B_i by any tree (for example a path from top to bottom). A tree on $|B_i|$ vertices has $|B_i| - 1$

edges, so a column with 1 black point adds no edges. Put the trees of all the columns together on the same n row vertices, and call the result H.



(a) a blackout (b) its graph H (c) another blackout (d) its H has a triangle

Figure 12: (a) A blackout of the 5×4 grid, with the black points of each column joined. (b) The graph H: each column with 2 or more black points contributes a path on its black rows, in the same colour. (c), (d) The same with column 4 moved to rows 2 and 5, which creates a triangle in H.

Example 5.2. In Figure 12(a), column 1 is black in rows $1, 3, 5$, column 2 in rows $2, 3$, column 3 in row 4 and column 4 in rows $4, 5$. In (b), column 1 gives the path $r_1r_3r_5$, column 2 gives the edge r_2r_3, column 3 gives nothing and column 4 gives r_4r_5. The result is a tree with 4 edges. Count the black points column by column: each non-empty column has 1 point for its first row and 1 more for each edge of its tree, so $|S| = 4 + 4 = 8$ (4 non-empty columns, 4 edges of H).

In (c), column 4 is black in rows 2 and 5 instead, and H gets the edge r_2r_5. Now H has the triangle $r_2r_3r_5$, with its 3 edges coming from 3 different columns. Go back to the grid and this triangle is the black 6-cycle $r_3\, c_1\, r_5\, c_4\, r_2\, c_2$, through the points $(1,3), (1,5), (4,5), (4,2), (2,2), (2,3)$. By Lemma 4.3 that blackout is not valid.

The example shows the general facts we need.

Lemma 5.3. *If S is valid and $m \geq 3$, then H has no repeated edges and no triangles, and $|S| = t + e(H)$, where t is the number of non-empty columns and $e(H)$ the number of edges of H.*

Proof. By Lemma 4.3, G_S has no 4-cycle and no 6-cycle.


No repeated edges. One tree never uses the same edge twice. If 2 different columns i, i' both gave the edge $r_jr_{j'}$, both columns would be black in rows j and j', which is the 4-cycle $r_j\, c_i\, r_{j'}\, c_{i'}$.

No triangles. Suppose $r_ar_br_c$ is a triangle in H. The 3 edges can't all come from 1 column, because a tree has no cycles. If 2 come from column i and the third from column $i' \neq i$, then all of a, b, c are in B_i, and the third edge puts 2 of them in $B_{i'}$; so columns i, i' share 2 black rows, a 4-cycle. If the 3 edges come from 3 different columns i (r_ar_b), i' (r_br_c) and i'' (r_ar_c), then $r_a\, c_i\, r_b\, c_{i'}\, r_c\, c_{i''}$ is a 6-cycle in G_S.

The count. $e(H) = \sum_{B_i \neq \emptyset}(|B_i| - 1) = |S| - t$. □

5.2 Mantel's theorem

Lemma 5.4 (Mantel)**.** *A graph on n vertices with no triangle and no repeated edges has at most $\lfloor n^2/4 \rfloor$ edges.*



$n = 3$: 2 edges | $n = 4$: 4 edges | $n = 5$: 6 edges

Figure 13: Triangle-free graphs with the most edges for $n = 3, 4, 5$: split the vertices into 2 nearly equal groups and join every pair across.

Proof. Let the graph have e edges, and write $d(v)$ for the degree of v. If $e = 0$ there is nothing to prove. For an edge uv, the neighbours of u and the neighbours of v don't overlap (a common neighbour would make a triangle), so $d(u) + d(v) \le n$. Add this up over all e edges. Each vertex v appears once for each of its $d(v)$ edges, so the left side adds up to $\sum_v d(v)^2$, and

$$\sum_v d(v)^2 \le ne.$$

On the other hand $\sum_v d(v) = 2e$, so by the Cauchy–Schwarz inequality $\sum_v d(v)^2 \ge (2e)^2/n$. Together, $4e^2/n \le ne$, so $e \le n^2/4$, and since e is a whole number, $e \le \lfloor n^2/4 \rfloor$. □

Proof of Theorem 5.1. By Lemma 5.3, $|S| = t + e(H)$ with $t \le m$, and H is triangle-free on n vertices, so $e(H) \le \lfloor n^2/4 \rfloor$ by Mantel. Hence $|S| \le m + \lfloor n^2/4 \rfloor$. Everything so far is symmetric in rows and columns except where Lemma 4.3 needed a third column, so if $n \ge 3$ the same argument with rows and columns swapped gives $|S| \le n + \lfloor m^2/4 \rfloor$. □

For 2, 3 and 4 rows this gives $m + 1$, $m + 2$ and $m + 4$. On 2 rows that is the Strip Theorem's value. On 3 rows and on 4 rows with $m \ge 6$ it is also exact, and the next sections show it.

6 Tilted rectangles on 3 and 4 rows

To use the row bound on narrow grids we still need to handle tilted rectangles. On 3 or 4 rows there are only a few kinds.

For $1 \le i \le m - 2$ and $1 \le j \le n - 2$, the *diamond* is the tilted square

$$D(i, j) = \{(i, j + 1),\ (i + 1, j),\ (i + 2, j + 1),\ (i + 1, j + 2)\}.$$

For $1 \le i \le m - 3$, define 4 more kinds on 4 rows:

$$A_i = \{(i, 3), (i + 1, 4), (i + 2, 1), (i + 3, 2)\}, \qquad B_i = \{(i, 3), (i + 2, 4), (i + 1, 1), (i + 3, 2)\},$$
$$C_i = \{(i, 2), (i + 1, 4), (i + 2, 1), (i + 3, 3)\}, \qquad E_i = \{(i, 2), (i + 2, 4), (i + 1, 1), (i + 3, 3)\}.$$

Lemma 6.1 (tilted rectangles on narrow grids)**.** *On 3 rows, the tilted rectangles are exactly the diamonds $D(i, 1)$. On 4 rows, they are exactly the diamonds $D(i, 1)$ and $D(i, 2)$ and the rectangles A_i, B_i, C_i, E_i.*

Proof. First, a tilted rectangle has exactly 1 leftmost corner. Suppose 2 corners both have the smallest column number x_0. They aren't next to each other, since no side of a tilted rectangle is




Figure 14: The diamond $D = D(1,1)$ and the 4 other kinds A_1, B_1, C_1, E_1, in a 4×4 window. The ringed corner is the leftmost one.

vertical, so they are opposite corners and the centre of the rectangle is in column x_0. The other 2 corners have the same centre, so their column numbers add up to $2x_0$. Both are at least x_0, so both equal x_0, and then all 4 corners are on 1 vertical line, which is impossible.

From the leftmost corner both sides go to the right. They are at right angles, so one goes up and the other goes down. Write them as $(a, -b)$ (up and right) and (c, d) (down and right), where a, b, c, d are positive integers. Being at right angles means $ac - bd = 0$, so $ac = bd$. The corners use the rows from (leftmost row $-\,b$) to (leftmost row $+\,d$), so the rectangle covers $b + d + 1$ rows.

On 3 rows, $b + d + 1 \leq 3$, so $b = d = 1$, then $ac = 1$ gives $a = c = 1$, and the rectangle is a diamond. On 4 rows, either $b = d = 1$ again (a diamond), or $b + d = 3$, so $\{b, d\} = \{1, 2\}$, $ac = bd = 2$ and $\{a, c\} = \{1, 2\}$. These are 4 choices, and they are the 4 kinds:

	A	B	C	E
side going up, $(a, -b)$	$(2, -2)$	$(1, -2)$	$(2, -1)$	$(1, -1)$
side going down, (c, d)	$(1, 1)$	$(2, 1)$	$(1, 2)$	$(2, 2)$

The leftmost corner is then forced into row 3 for A, B (it needs 2 rows above and 1 below) and row 2 for C, E, which gives the formulas above. □

So the 4×4 grid has 8 tilted rectangles: $D(1,1), D(2,1), D(1,2), D(2,2), A_1, B_1, C_1, E_1$. Note that B and C are tilted squares with side $\sqrt{5}$, while A and E have sides $\sqrt{2}$ and $2\sqrt{2}$.

7 Three rows


Theorem 7.1 (Three-Row Theorem)**.** *For every $m \geq 3$, the maximum blackout of the $3 \times m$ grid is $m + 2$.*

The upper bound is the row bound, $m + \lfloor 9/4 \rfloor = m + 2$. For the lower bound we black out all of row 1 and all of column 1:

$$S = \{(i, 1) : 1 \leq i \leq m\} \cup \{(1, 2), (1, 3)\}.$$

This has $m + 2$ points. It is the blackout of Example 1.1, now on a longer grid.

Proof that S is valid. Axis-aligned rectangles. In G_S the vertex r_1 is joined to every column and c_1 is also joined to r_2 and r_3 (Figure 15(b)). That is $m + 3$ vertices, $m + 2$ edges and connected, so G_S is a tree. A tree has no cycles at all, so by Lemma 4.3 no 2 axis-aligned rectangles look the same.



(a) row 1 and column 1, with $D(1,1)$ and $D(3,1)$

(b) G_S is a tree

Figure 15: The three-row construction for $m = 6$. The diamond $D(1,1)$ (solid) shows only $(3,2)$ and $(2,3)$; the diamond $D(3,1)$ (dashed) shows 3 corners.

Tilted rectangles. By Lemma 6.1 they are the diamonds $D(i,1) = \{(i,2),(i+1,1),(i+2,2),(i+1,3)\}$, $1 \le i \le m-2$. Check each corner. The corner $(i+1,1)$ is in row 1, so it is black. The corner $(i,2)$ is black only if $i = 1$. The corners $(i+2,2)$ and $(i+1,3)$ are in rows 2, 3 and columns at least 2, so they are white.

So for $i \ge 2$ the diamond has 1 black corner and shows 3, and by Lemma 4.2 nothing else shows the same 3 points. For $i = 1$ the diamond shows $\{(3,2),(2,3)\}$. Could another rectangle show exactly these 2 points? It would have both points as corners. An axis-aligned rectangle with corners $(3,2)$ and $(2,3)$ has them as opposite corners, so it is the square with corners $(2,2),(3,2),(2,3),(3,3)$, and $(2,2)$, $(3,3)$ are white, so it shows 4 points. Every other diamond shows 3 points. So no other rectangle shows $\{(3,2),(2,3)\}$.

Finally an axis-aligned rectangle and a diamond $D(i,1)$ with $i \ge 2$ can only agree if both show the same 3 points, which Lemma 4.2 rules out. So S is valid. □

Remark 7.2. The same "row 1 plus column 1" blackout has $n + m - 1$ points on any grid, and it is valid on every grid up to 10×10 (checked by `verify_narrow.py`). It is the maximum on 3 rows and on 4×4 and 4×5 (Section 8.2), but from 4×6 on we can do 1 better.

8 Four rows

8.1 The construction for $m \ge 6$


Theorem 8.1 (Four-Row Theorem)**.** *For every $m \ge 6$, the maximum blackout of the $4 \times m$ grid is $m + 4$.*

The upper bound is the row bound, $m + \lfloor 16/4 \rfloor = m + 4$. For the lower bound take

$$S = \{(i,1) : 2 \le i \le m-1\} \cup \{(m-1,2),(m,2),(1,3),(2,3),(1,4),(m,4)\},$$

which has $(m-2) + 6 = m + 4$ points.

Proof that S is valid. Axis-aligned rectangles. Only 4 columns have 2 black points: column 1 (rows 3, 4), column 2 (rows 1, 3), column $m-1$ (rows 1, 2) and column m (rows 2, 4). Every other column is black only in row 1. So G_S is the 8-cycle $r_1\, c_2\, r_3\, c_1\, r_4\, c_m\, r_2\, c_{m-1}$ with the columns $c_3, \ldots, c_{m-2}$ hanging off r_1 (Figure 16). It has no 4-cycle and no 6-cycle, so Lemma 4.3 handles the axis-aligned rectangles.



(a) the blackout, $m = 7$ (b) G_S

Figure 16: The four-row construction for $m = 7$. The bold path in (a) turns only at black points and alternates between rows and columns; it is the 8-cycle in (b). The columns $3, 4, 5$ hang off r_1 (dashed).

Tilted rectangles with at most 1 black corner. These show 3 or 4 corners and are safe by Lemma 4.2. This covers $D(i, 2)$, C_i and E_i. For example, take $D(i, 2) = \{(i, 3), (i + 1, 2), (i + 2, 3), (i + 1, 4)\}$. In rows 2 to 4 the black points are $(m - 1, 2), (m, 2), (1, 3), (2, 3), (1, 4), (m, 4)$. So $(i, 3)$ is black only if $i \le 2$, $(i + 1, 2)$ only if $i = m - 2$, and $(i + 2, 3)$ and $(i + 1, 4)$ are never black (since $3 \le i + 2$ and $2 \le i + 1 \le m - 1$). Since $m \ge 6$, $i \le 2$ and $i = m - 2$ can't both hold, so at most 1 corner is black. For C_i and E_i the same kind of check shows that the only black corner is the one in row 1.

The other tilted rectangles. Checking each corner of each $D(i, 1)$, A_i and B_i in the same way gives the following list of the tilted rectangles that show at most 2 corners. Here F stands for A (with $s = 1$) or B (with $s = 2$).

rectangle	which i	corners shown
$D(1, 1)$		$(1, 2), (3, 2)$
$D(i, 1)$	$i = m - 3,\ m - 2$	$(i, 2), (i + 1, 3)$
F_i	$i \in \{1, 2\},\ i \notin \{m - 4, m - 3\}$	$(i + s, 4), (i + 3, 2)$
F_i	$i \in \{m - 4, m - 3\},\ i \notin \{1, 2\}$	$(i, 3), (i + s, 4)$
F_i	$i \in \{1, 2\} \cap \{m - 4, m - 3\}$	$(i + s, 4)$

The last line only happens when $m = 6$ and $i = 2$. As a sample check, take A_1, with corners $(1, 3)$, $(2, 4)$, $(3, 1)$ and $(4, 2)$. The corners $(1, 3)$ and $(3, 1)$ are black, $(2, 4)$ is white (row 4 is black only in columns 1 and m), and $(4, 2)$ is white (row 2 is black only in columns $m - 1, m$, and $4 < m - 1$). So A_1 shows $(2, 4), (4, 2)$, which is the third line with $s = 1$. Figure 17 shows the cases for $m = 7$. (The verification script also checks the whole construction against every rectangle for each width from 6 to 40 and for 64, 100 and 128.)

We have to show that each presentation in the table belongs to only 1 rectangle.


Against each other. The 2 shown points are in rows $\{2, 2\}$ for $D(1, 1)$, rows $\{2, 3\}$ for the other diamonds, rows $\{2, 4\}$ in line 3 and rows $\{3, 4\}$ in line 4, so different lines never agree. Within a line, the columns of the 2 points give i, and their horizontal distance ($3 - s$ in line 3 and s in line 4) gives whether it is A or B. The one-point presentations in the last line are $(3, 4)$ and $(4, 4)$, which are different.

Against axis-aligned rectangles.

- $D(1, 1)$ shows $(1, 2), (3, 2)$, both in row 2. An axis-aligned rectangle showing exactly these would use columns 1 and 3 and one more row in which both $(1, \cdot)$ and $(3, \cdot)$ are black. Column 1 is black in rows 3, 4 and column 3 only in row 1, so there is no such row.

- $D(i,1)$ with $i \geq m-3$ shows $(i,2),(i+1,3)$, which are in different rows and columns. The only axis-aligned rectangle with these corners also has the corner $(i,3)$, which is white because $i \geq m-3 \geq 3$. So that rectangle shows at least 3 points.
- Line 3: the other axis-aligned corner $(i+s,2)$ is white, because $i+s \leq 4 < m-1$.
- Line 4: the other axis-aligned corner $(i,4)$ is white, because $3 \leq i \leq m-3$.
- The last line ($m=6$): an axis-aligned rectangle showing only $(z,4)$ with $z \in \{3,4\}$ would need another column c and another row r with $(c,4)$, (z,r) and (c,r) all black. Then $c \in \{1,6\}$ (row 4) and $r=1$ (columns 3, 4 are black only in row 1), but $(1,1)$ and $(6,1)$ are white.

Finally, no rectangle is fully black: an axis-aligned one would be a 4-cycle, and every tilted one shows at least 1 point by the table and the paragraph before it. So S is valid. □



(a) $D(1,1)$ (b) A_1 and B_1 (c) A_4 and B_4

Figure 17: Tilted rectangles with 2 black corners in the construction for $m=7$, with the shown corners ringed. Each pair of shown points belongs to no other rectangle.

8.2 Widths 4 and 5

On 4×4 and 4×5 the row bound says $m+4$, but the answer is $m+3$. To prove it we first narrow down what a blackout of $m+4$ points would have to look like.


Lemma 8.2 (equality in the row bound)**.** *Let $m \geq 4$. If a valid blackout of the $4 \times m$ grid has $m+4$ points, then every column is non-empty, exactly 4 columns have 2 black points and the others have 1, and the graph H of Section 5 is a 4-cycle on the 4 rows.*

Proof. By Lemma 5.3, $m+4 = |S| = t + e(H)$ with $t \leq m$ and $e(H) \leq 4$, so $t = m$ (every column is non-empty) and $e(H) = 4$. Recall $e(H) = \sum_i (|B_i| - 1)$.

A column black in all 4 rows gives 3 edges, and then every other column has at most 1 black point (2 would share 2 rows with it, a 4-cycle), so $e(H) = 3$. A column black in 3 rows gives 2 edges. Any other column with 2 or more black points can share at most 1 row with it, so it is a pair made of the fourth row and 1 of the 3. There is at most 1 such column: 2 with the same pair give a 4-cycle, and 2 with different pairs give a 6-cycle. So $e(H) \leq 2+1 = 3$. Hence every column has 1 or 2 black points, and exactly 4 columns have 2.

These 4 columns give 4 edges of H, with no repeats and no triangle. A graph on 4 vertices with 4 edges misses 2 of the 6 possible edges. If the 2 missing edges shared a vertex v, the other 3 vertices would form a triangle. So the missing edges don't touch, and what is left is a 4-cycle. □

Corollary 8.3 (computer-assisted)**.** *The maximum blackout is 7 on the 4×4 grid and 8 on the 4×5 grid.*



(a) the 3 possible graphs H (b) a candidate with 8 points (c) valid with 7 points

Figure 18: (a) The 3 possible 4-cycles H on the rows. (b) One of the 72 candidates on the 4×4 grid. The tilted rectangles $D(2,1)$ (solid) and A_1 (dashed) are both completely black, so both have empty presentations. (c) Row 1 plus column 1: valid, with 7 points.

Proof. The row bound gives at most $m+4$, and Lemma 8.2 says what a blackout of size $m+4$ looks like. So we can list all of them: choose 1 of the 3 four-cycles on the rows (Figure 18(a)), choose the 4 columns with 2 black points, match the 4 edges of the cycle to those 4 columns, and choose 1 black row for each remaining column. That is

$$3 \cdot \binom{m}{4} \cdot 4! \cdot 4^{m-4}$$

candidates, which is 72 for $m = 4$ and 1,440 for $m = 5$. The script `verify_narrow.py` checks every candidate against all rectangles, and every one of them has 2 rectangles with the same presentation. Figure 18(b) shows one. So the maximum is at most $m+3$. Row 1 plus column 1 has $m+3$ points and the script checks that it is valid on both grids, so the maximum is $m+3$. □

In Figure 18(b) the axis-aligned rectangles are fine (the graph has no short cycle), and the trouble comes from 2 tilted rectangles that are both completely hidden. This is Remark 1.4 at work. Of the 72 candidates on 4×4, the first collision the script finds involves only tilted rectangles in 9 cases and 1 axis-aligned rectangle in the other 63; on 4×5 the split is 118 and 1,322.

9 The 5×5 grid

The 5×5 grid is the first square where $n+m-1$ is not the answer. It is also the smallest grid with both sides at least 5. The row bound only gives $5 + \lfloor 25/4 \rfloor = 11$ here. The answer is 10, and the proof is a counting argument on the graph G_S.

9.1 The upper bound

On a 5×5 grid the graph G_S has only 10 vertices. Lemma 4.3 rules out cycles of length 4 and 6, and G_S is bipartite, so every cycle has length at least 8. On 10 vertices there is room for 1 such cycle but not for 2 that overlap, and 2 that don't overlap need at least 16 vertices. So G_S can't have more edges than vertices.

Lemma 9.1 (few edges without short cycles)**.** *Let G be a bipartite graph with $v \le 10$ vertices and no cycle of length 4 or 6. Then G has at most v edges. If it has exactly v edges, then G is connected and has exactly 1 cycle, of length 8 or 10.*




Figure 19: The count in Step 1. Around a vertex x of degree at least 3, the vertices at distance at most 3 form a tree, so there are at least $1+3+3+3$ of them, and each vertex at distance 3 leads on to a vertex at distance 4. Black and white show the 2 sides of the bipartite graph.

Proof. Since G is bipartite, every cycle has even length, so every cycle has length at least 8.

Step 1: no connected piece of G has more edges than vertices. Suppose a piece G' does. Remove a vertex of degree 1 from G', together with its edge, and repeat until there are none. Each step removes 1 vertex and 1 edge, so what is left, H, still has more edges than vertices. In particular H isn't empty and every vertex of H has degree at least 2. If every degree were exactly 2, H would have as many edges as vertices, so some vertex x has degree at least 3.

Let D_k be the set of vertices of H at distance k from x. Two different paths from x to the same vertex, each of length at most 3, would together contain a cycle of length at most 6, which is impossible. So for $k = 1, 2, 3$ each vertex of D_k has exactly 1 neighbour in D_{k-1}, its *parent*. It has no neighbour in D_k itself, because D_k is on one side of the bipartite graph. Its degree is at least 2, so it has a neighbour in D_{k+1}. Two different vertices of D_k (with $k \le 2$) can't share such a neighbour, since that neighbour would have 2 parents. So $|D_1| \ge 3$, $|D_2| \ge |D_1| \ge 3$, $|D_3| \ge |D_2| \ge 3$ and $|D_4| \ge 1$ (Figure 19). That is at least $1+3+3+3+1 = 11$ vertices, but $v \le 10$. This contradiction proves Step 1.


Step 2: equality. By Step 1 each piece has at most as many edges as vertices, so G has at most v edges. If it has exactly v, each piece has exactly as many edges as vertices, and a connected graph like that has exactly 1 cycle. Two pieces would mean 2 separate cycles of length at least 8, which needs 16 vertices. So G is connected, and its 1 cycle has even length between 8 and 10. □

Theorem 9.2 (the 5×5 grid)**.** *The maximum blackout of the 5×5 grid is 10. For every maximum blackout S, the graph G_S is connected with exactly 1 cycle, of length 8 or 10. By computer, there are exactly 8 maximum blackouts, they are the 8 images of Figure 20(a) under the symmetries of the square, and each graph is an 8-cycle with 2 pendant edges.*

Proof. Upper bound. If S is valid, G_S has no 4-cycle or 6-cycle (Lemma 4.3, with $m = 5 \ge 3$). It is bipartite with 10 vertices and $|S|$ edges, so Lemma 9.1 gives $|S| \le 10$, and if $|S| = 10$ it gives the structure in the statement.

Lower bound. The blackout in Figure 20(a) has 10 points. Its graph is an 8-cycle with 2 pendant edges, so Lemma 4.3 handles the 100 axis-aligned rectangles. The script checks all 130 presentations directly and finds them all different.

The count. A maximum blackout has a graph with 10 edges and no 4- or 6-cycle. The script lists all 44,640 such sets of 10 points (adding points one at a time and only keeping a point if no short cycle appears), and checks each one against all 130 rectangles. Exactly 8 are valid, and they are the 8 images in Figure 21. The count of 8 had been found earlier, in a completely separate way, by the hitting set program and by checking every blackout straight from the definition (Section 11). □



(a) the blackout (b) G_S

Figure 20: The 5×5 optimum. (a) The 10 black points and the 15 kept points, which form the 2 shaded blocks of 5 and 10. The bold path turns only at black points and alternates between rows and columns; its 8 corners are the 8-cycle of (b). The 2 grey points are the 2 pendant edges of (b). (b) The graph G_S: an 8-cycle with 2 pendant edges.




Figure 21: The 8 maximum blackouts of the 5×5 grid. They are all different, so the optimum is unique up to symmetry but has no symmetry of its own.

9.2 What the optimum looks like

The optimum in Figure 20(a) is a staircase of 9 points from the top-right corner to the bottom-left corner, plus the corner $(5, 5)$. The staircase alone has $9 = n + m - 1$ points and its graph is a tree. The extra corner closes a cycle of length 8, the shortest the lemma allows. The kept points form 2 blocks of 5 and 10 on either side of the staircase.

What about the other shape Lemma 9.1 allows, a single 10-cycle? That means 2 black points in every row and every column, and there are 1,440 such sets. None of them is valid, and the trouble always involves a tilted rectangle. Figure 22 shows one.

So the open question of a hand proof of uniqueness comes down to blackouts of 10 points whose graph is connected with 1 cycle of length 8 or 10, and to the tilted rectangles on them.




Figure 22: A 10-cycle blackout: 2 black points in every row and every column, and no short cycle. The tilted rectangle (solid) and the axis-aligned one (dashed) both show only the ringed points $(1, 3)$ and $(3, 5)$.

10 Bigger grids

What happens beyond 4 rows? The pattern on 5×5, a staircase from one corner to the opposite corner plus 1 more point, keeps working.




Figure 23: Valid blackouts of $n + m$ points on 5×6, 6×6 and 6×7: a staircase from one corner to the opposite corner, plus 1 more point.

While building the game (`game/`), a search found a valid blackout of $n+m$ points of this kind on every grid from 5×5 up to 10×10, and the game stores 1 for each size. The script `verify_narrow.py` rechecks all 21 of them with its own rectangle list, which shares no code with the game, and they are all valid. So on every grid with both sides between 5 and 10,

$$n + m \;\le\; \text{maximum blackout} \;\le\; \min\bigl(m + \lfloor n^2/4 \rfloor,\; n + \lfloor m^2/4 \rfloor\bigr).$$

For 5×5 both ends meet only after Lemma 9.1. For 5×6, 5×7 and 6×6 the hitting set program found the value $n + m$ exactly. Everything we have seen fits the following guess.

Conjecture 10.1 (Excess-1)**.** *If* $\min(n, m) \ge 4$*, the maximum blackout of the* $n \times m$ *grid is* $n + m$*, except for* 4×4 *and* 4×5 *(and their transposes), where it is* $n + m - 1$*.*

The four-row case is now Theorem 8.1 with Corollary 8.3, and 5×5 is Theorem 9.2. The rest is open. Neither the row bound nor Lemma 9.1 (which needs $n + m \le 10$) is strong enough there.

11 Computations

11.1 The programs

Three separate programs were used, and they agree wherever more than 1 of them can run.

- *The hitting set program* (`solver/engine.py`) lists the rectangles, builds the difference sets, prunes them as in Section 2.2, and finds the smallest hitting set with an integer linear program: one 0–1 variable x_v for each point, minimize $\sum_v x_v$, subject to $\sum_{v \in D} x_v \geq 1$ for every constraint D. The value is nm minus the optimum. To count the optima it then lists every kept set of the smallest size and checks it.
- *The brute force check* (`solver/brute.py`) uses no lemma at all. It finds rectangles by testing every 4-point set with the diagonal test, and it checks each blackout straight from the definition by computing all presentations and looking for a repeat. It is slow on purpose and only feasible up to about 30 points.
- *The narrow-grid verifier* (`verify_narrow.py`) checks the lemmas and constructions of Sections 4 to 10: Lemma 4.3 on all 74,304 blackouts of 5 small grids, the list of tilted rectangles, the constructions at widths up to 128, the finite four-row checks, the 5×5 search and the stored large blackouts. It uses only the Python standard library and runs in about 20 seconds.

11.2 The values

grid	rectangles	max blackout	kept	optima	checked by
2×2	1	4	0	1	hand, program, brute force
2×3	3	4	2	12	hand, program, brute force
2×4	6	5	3	32	hand, program, brute force
2×5	10	6	4	80	program, brute force (predicted)
2×6	15	7	5	192	program, brute force
2×7	21	8	6	448	program, brute force
3×3	10	5	4	53	program, brute force
3×4	20	6	6	224	program, brute force
3×5	33	7	8	892	program, brute force
3×6	49	8	10	3420	program, brute force
3×7	68	9	12	12704	program, brute force
3×8	90	10	14	45864	program, brute force
3×9	115	11	16	161992	program, brute force
3×10	143	12	18	–	program (value only)
4×4	44	7	9	316	program, brute force
4×5	74	8	12	1810	program, brute force
4×6	110	10	14	26	program, brute force
4×7	152	11	17	292	program, brute force
4×8	200	12	20	–	program (value only)
5×5	130	10	15	8	program, brute force
5×6	198	11	19	–	program (value only)
5×7	276	12	23	–	program (value only)
6×6	313	12	24	–	program (value only)

Every value for 2, 3 and 4 rows and the value for 5×5 now also follow from proofs in these notes. The values for 5×6, 5×7 and 6×6 rest on the program, with the lower bound $n + m$ also confirmed by the stored blackouts of Section 10. The counts of optima beyond 2 rows are only computed.

11.3 Predictions that failed

Before most computations a prediction was written in the ledger (`data/ledger.md`), so that the computation could prove it wrong. Several did go wrong.




Figure 24: The maximum blackout of each computed $n \times m$ grid. Shaded cells beat $n + m - 1$ by exactly 1, white cells equal it. A small "p" marks a value that is proved. (2×2 is the trivial grid with only 1 rectangle.)

guess	where it failed	what happened
kept $= (n-1)(m-1)$, so blackout $n+m-1$	5×5	10 instead of 9
blackout $= n+m-1+[\,n+m \geq 10\,]$	3×7	9 instead of 10
blackout $= n+m-1+[\,nm \geq 24\,]$	3×8	10 instead of 11
kept on 4 rows $= \lceil 5m/2 \rceil - 1$	4×8	12 instead of 13

The first guess fit all 8 values computed before 5×5. The 5×5 grid broke it, and it also has only 8 optima against 1,810 on 4×5, so something changes there. The next 2 guesses tried to say when the extra point appears, by the sum or by the area of the grid, and 3 rows killed both: 3 rows never get the extra point. The four-row formula fit 4 values and died on the fifth. What survived is what is now proved: $m+1$, $m+2$ and $m+4$ on 2, 3 and 4 rows, and the Excess-1 conjecture beyond.

12 Open questions

1. Prove Conjecture 10.1 when both sides are at least 5. The lower bound $n+m$ is known up to 10×10 by computer; a general construction and a matching upper bound are both missing.
2. Count the optima beyond 2 rows. Why does the number collapse to 26 on 4×6, when its neighbours have hundreds or thousands?
3. Prove that the 5×5 optimum is unique up to symmetry without a computer. Theorem 9.2 reduces this to blackouts whose graph is connected with 1 cycle of length 8 or 10.
4. Does the excess over $n+m-1$ ever reach 2? The next grids to try are 5×8 and 6×7, where the ledger predicts 13.
5. The row–column graph turns the axis-aligned part into a question about bipartite graphs with no 4- or 6-cycle. How does this compare with the literature on test covers and on extremal graphs of large girth?

References

[1] P. Kagey, Problem 001, Open Problem Collection, https://peterkagey.com/problems/001/ (accessed August 2026).

[2] OEIS Foundation, Sequence A085582: number of rectangles (orthogonal or not) with corners on an $n \times n$ grid of points, https://oeis.org/A085582.

[3] OEIS Foundation, Sequence A289832: number of rectangles with corners on an $n \times k$ grid of points, https://oeis.org/A289832. Here $T(n, k)$ counts rectangles on an $(n+1) \times (k+1)$ grid of points, so our $r \times c$ grid is $T(c-1, r-1)$; for example our 3×7 count is $T(6, 2) = 68$.