Maximum Blackouts That Keep Every Lattice Rectangle Identifiable


Arjun Pemmasani
Department of Mathematics
Harvey Mudd College
Claremont, CA 91711
USA
apemmasani@hmc.edu

Abstract

Kagey's Problem 001 concerns a grid of points, some of them blacked out. A rectangle with its corners on grid points, tilted or not, shows only its white corners. We ask how many points can be blacked out while every rectangle stays determined by what it shows. We study the maximum blackout and the number of optima, tabulated in A397315 and A399580 of the On-Line Encyclopedia of Integer Sequences. A row-column graph and Mantel's theorem bound the maximum by the width plus a quarter of the squared height. We find the maximum on up to five rows at every width, and on six rows from width 15 on. The five-row result refutes our earlier guess that the maximum is the sum of the two sides. We count the three-row optima exactly and find the growth rate on four rows. At every fixed height, the maximum minus the width is eventually periodic, and the optimum counts have a rational generating function. Some values rest on computer checks.

1 Introduction

Start with an $h \times m$ grid and black out some of its points. The game master chooses a rectangle (tilted ones are allowed) and tells you which of its corners



(a) the point $(3, 2)$ (b) our blackout (c) the rectangle (d) what we are shown

Figure 1: One round on a largest valid blackout of the 3×4 grid (Section 5). Points are (column, row). Gray crossed circles are black. Panel (c) shows the red rectangle, and (d) its yellow corners. The three yellow corners fix the fourth by Lemma 4.

are white. Can you recover all four corners? Kagey [17, 18] asks how many points can be blacked out while keeping the answer unique for every rectangle. Such a set is a *valid blackout* (Definition 1). One round is shown in Figure 1.

Write $f(h, m)$ for the most points we can black out, and $a_h(m)$ for the number of ways to attain it. The conjectures recorded in our OEIS entries A397315 and A399580 [30] were $f(3, m) = m + 2$ and $f(h, m) = h + m$ for $\min(h, m) \geq 4$, with exceptions $f(4, 4) = 7$ and $f(4, 5) = 8$. The first was proved in Lean by the Omega Institute, whose pull request was merged on 15 September 2026 [26]. Our independent proof combines the row bound (Theorem 8) and the boundary cross (Lemma 17). The second guess breaks at 5×8 (Theorem 40). Throughout, put $q_h = \lfloor h^2/4 \rfloor$. Computer-assisted claims are marked, with evidence in Table 3.

The main results follow, with proof locations in parentheses.

Theorem A (narrow grids)**.**

(a) We have $f(h, m) \geq h + m - 1$*, and* $f(h, m) \leq m + q_h$ *if* $m \geq 3$*.* (Lemma 17 and Theorem 8, Section 4.)

(b) We have $f(2, m) = m + 1$ *and* $f(3, m) = m + 2$ *for* $m \geq 3$*,* $f(4, m) = m+4$ *for* $m \geq 6$*, and* $f(5, 5) = 10$*.* (Theorem 5, Section 5, Theorem 31, and Theorem 44(a).)

(c) (Computer-assisted, with checked certificates for $f(5, 6)$ *and* $f(5, 7)$*, and the last claim checked in Lean.) We have* $f(4, 4) = 7$*,* $f(4, 5) = 8$*,* $f(5, 6) = 11$*,* $f(5, 7) = 12$*, and* $f(5, m) = m + 6$ *for* $m \geq 8$*.* (Corollary 32 and Theorem 40(a), (b).)

(d) *(Computer-assisted for $m \geq 9$, with checked certificates for $m \leq 14$, and checked in Lean for $m \geq 15$.) We have $f(6,m) \leq m+8$ for $8 \leq m \leq 14$, and $f(6,m) = m+9$ for $m \geq 15$.* (Theorem 42.)

(e) *(Computer-assisted, with checked certificates.) We have $f(6,6) = 12$, $f(6,7) = 13$, $f(7,7) = 14$, and $f(6,8) = 15$.* (Theorem 43.)

For (c) and (d), the collar theorem reduces each infinite family to one validity check (Theorem 39). Lemma 47 justifies the SAT bounds.

Theorem B (three-row optima)**.**

(a) *For $m \geq 4$ we have $a_3(m) = 4(m-1)(3m+2)\,3^{m-5} + E_m$, where $E_m > 0$ counts the optima with a column black in rows 1 and 3 (A398448), and $E_m = O(m^2(1+\sqrt{3})^m)$. So $a_3(m) \sim \frac{4}{81} m^2 3^m$, and a fraction tending to $4/27$ of the $m^2 3^{m-1}$ axis-valid candidates is valid. The series $\sum_{m \geq 3} a_3(m) z^m$ is rational, and its denominator divides $(1-3z)^3(1-2z-2z^2)^3(1+2z^2)^3$.* (Theorem 24, Proposition 19(c), the last claims of Theorem 26, and Theorem 25(a), Section 5.)

(b) *(Computer-assisted, exact computer algebra.) This is the reduced denominator, and $E_m \sim \frac{19+11\sqrt{3}}{432} m^2 (1+\sqrt{3})^m$.* (Theorem 25(b), proved in Section 5.2, and Theorem 26.)

Theorem C (four rows)**.** *If $m \geq 4$, no valid blackout of the $4 \times m$ grid hides all corners of a rectangle* (Theorem 33)*. (Computer-assisted.) The series $\sum_{m \geq 6} a_4(m) z^m$ is rational with reduced denominator of degree 420, and $a_4(m) \sim c\, m^4 \rho_4^m$ with $c \approx 8.1611 \times 10^{-5}$ and $\rho_4 = 3.35199\ldots$, an algebraic number of degree 20* (Proposition 36, Theorem 37, and Corollary 38, Section 6)*.*

Theorem D (every fixed height)**.** *Fix $h \geq 1$ and write a blackout as the word $B_1 \cdots B_m$ of the black rows of its columns. The valid words form a regular language. If $h \geq 2$, then $f(h,m) - m$ is eventually periodic in m, and $\sum_{m \geq 0} a_h(m) z^m$ is rational.* (Theorems 45 and 46, Section 8.)

Corollary 15 proves the conjecture in A338885 that its row n has largest entry A033638(n).

Prior work. Choosing white points to distinguish rectangles is a test-cover problem [4, 12, 19, 24] (Corollary 3). If we only forbid fully black axis-aligned rectangles, we get the Zarankiewicz problem [28]. To distinguish all axis-aligned rectangles, the row-column graph must avoid both four- and six-cycles (Lemma 7). Such graphs, including unbalanced parts, are studied by de Caen and Székely [7, 8]. Neuwirth's edge bounds [25, Theorem 1(ii) and Proposition 17] already give our row bound in graph form. Our Mantel proof [20] also classifies and counts the equality cases. The tilted rectangles require additional arguments. For fixed height, our column method uses transfer matrices [1, 9, 21], [10, Proposition V.6]. Theorem 1(3) of Guilhot, Little, and Parkinson [13] also yields Theorem D once regularity is proved.

Outline. Sections 2 to 4 set up the problem and its graph bounds. Sections 5 to 8 handle narrow grids and fixed height. Section 9 gives computational evidence and encoding proofs, Section 10 the open questions, and Appendix A the square-grid bound.

2 Conventions

We keep track of the black points column by column. The following notation will be used throughout.

(1) An $h \times m$ grid has h rows, m columns, and point set V. A $2 \times m$ grid is a *strip*. The point (i, j) lies in column i and row j, with $(1, 1)$ at the top left (Figure 1(a)).

(2) A *rectangle* has four grid points as corners and positive area. We write $C(R)$ for its corner set. It is *axis-aligned* if its sides follow the grid lines, and *tilted* otherwise.

(3) A *blackout* is the set $S \subseteq V$ of black points. Its *kept set* is $K = V \setminus S$, and the *presentation* $P_S(R) = C(R) \setminus S$ is what the rectangle shows, namely its white corners.

(4) Column i has *black rows* $B_i = \{j : (i, j) \in S\}$, and is *empty*, a *singleton*, a *double*, or a *triple* when $|B_i|$ is 0, 1, 2, or 3, and *full* when $|B_i| = h$. The *pair set* F is the set of row pairs contained in some B_i, and $q_h = \lfloor h^2/4 \rfloor$ is A002620(h).

Definition 1 (valid blackout)**.** A blackout S is *valid* if distinct rectangles have distinct presentations. We write $f(h, m)$ for the largest size of a valid blackout of the $h \times m$ grid, and $a_h(m)$ for the number of valid blackouts of that size. We count them with their coordinates (reflections and rotations are not identified).

Transposing the grid gives $f(h, m) = f(m, h)$ and $a_h(m) = a_m(h)$. There are no rectangles at width zero or height one, so we set $f(h, 0) = 0$, $a_h(0) = 1$, $f(1, m) = m$ and $a_1(m) = 1$. A rectangle is allowed to show no corners, but at most one can do so in a valid blackout.

Lemma 2 (difference sets)**.** *For sets A, B, and S, we have $A \setminus S = B \setminus S$ if and only if $A \triangle B \subseteq S$.*

Proof. The two visible sets agree exactly when every point belonging to just one of A and B is black, that is, belongs to S. □

Corollary 3 (hitting sets)**.** *A blackout S is valid if and only if its kept set K meets $C(R_1) \triangle C(R_2)$ for every pair of distinct rectangles R_1, R_2.*

This is Lemma 2 applied to the corner sets. Keeping an extra point cannot create an ambiguity, so every subset of a valid blackout is valid. Hence ruling out blackouts of size k also rules out every larger size.

Lemma 4 (three corners suffice)**.** *Three corners of a rectangle determine it, so distinct rectangles with the same presentation show at most two corners.*

Proof. Three corners form a right triangle, whose only right angle is at the corner q adjacent to the other two, p and r. So the fourth corner is $p + r - q$. □

3 Strips

On two rows every rectangle is axis-aligned. To see this, take a lowest corner of a tilted rectangle. Its two neighbors lie strictly above it, and the opposite corner lies above both, so the rectangle meets at least three rows. The strip problem therefore comes down to pairs of columns.

Theorem 5 (strip theorem)**.** *Let $m \geq 3$.*

(a) *A blackout of the* $2 \times m$ *strip is valid if and only if at most one of its columns is full.*

(b) *We have* $f(2, m) = m + 1$*, and the maximum blackouts are those with one full column and one black point in every other column.*

(c) *There are* $a_2(m) = m \cdot 2^{m-1}$ *maximum blackouts, so row* 2 *of* A399580 *is* A001787 *from* $m = 3$ *on.*

For $m = 2$ *there is one rectangle, so* $f(2, 2) = 4$ *and* $a_2(2) = 1$*.*

Proof. (a) Two distinct rectangles share at most one column. Their difference set therefore contains both points of at least two columns. Conversely, every pair of columns p, q occurs as a whole difference set. Choose a third column z (this is where $m \geq 3$ is used) and take the rectangles on $\{z, p\}$ and $\{z, q\}$. By Corollary 3, validity is thus equivalent to keeping a point in every pair of columns, or having at most one full column.

(b, c) We must keep at least one point in all but one column, so $|S| \leq 2m - (m - 1) = m + 1$. Equality requires one full column and exactly one black point in each other column. There are m choices for the full column, then two choices in each of the remaining $m - 1$ columns. □

Proposition 6 (orbits and kings)**.** *For* $m \geq 3$ *the number* o_m *of maximum blackouts of the* $2 \times m$ *strip up to the row reflection and the column reversal is* $m\,2^{m-3} + [m \text{ odd}]\,2^{(m-3)/2}$*. It equals* A322284$(m - 1)$*, which counts* $m - 1$ *nonattacking kings on a* $2 \times (2m - 2)$ *board up to symmetry.*

Proof. Apply Burnside's lemma to the four symmetries. The identity fixes all $m2^{m-1}$ optima, and swapping the rows fixes none. A column reversal or half turn can fix an optimum only when m is odd and the full column is central. Each then has $2^{(m-1)/2}$ fixed optima, since the singleton columns come in pairs. Dividing their sum by four gives the formula.

Put $n = m - 1$. Nonattacking kings on the $2 \times 2n$ board are columns $c_1 < \cdots < c_n$ with $c_{j+1} \geq c_j + 2$, and any rows r_j. The gaps $c_1 - 1$, $c_{j+1} - c_j - 2$, and $2n - c_n$ are nonnegative and sum to 1, so exactly one, say the eth, is 1, and $c_j = 2j - 1 + [j \geq e]$. Sending this to the optimum with full column e and other black points in rows $r_1, \ldots, r_n$ from left to right is a bijection. It commutes with the reflection, which swaps the rows on both sides, and with the reversal, which sends e to $n + 2 - e$ and c to $2n + 1 - c$ and reverses the order of the row labels, since $2n + 1 - c_{n+1-j} = 2j - 1 + [j \geq n + 2 - e]$. So the orbits correspond. □

4 The row-column graph and general bounds

A blackout S is *axis-valid* when it distinguishes axis-aligned rectangles. Call a rectangle *weak* if at most two corners show. In the *row-column graph* G_S, join row r_j to column c_i for each black point (i, j).

Lemma 7 (axis-aligned rectangles and short cycles)**.** *If G_S has no four-cycle and no six-cycle, then S is axis-valid. The converse holds for $m \geq 3$.*

Proof. For $m \geq 3$, a four-cycle $r_j\, c_i\, r_{j'}\, c_{i'}$ gives indistinguishable rectangles $\{i, w\} \times \{j, j'\}$ and $\{i', w\} \times \{j, j'\}$ using a third column w. A six-cycle $r_a\, c_x\, r_b\, c_y\, r_c\, c_z$ gives $\{x, y\} \times \{a, b\}$ and $\{y, z\} \times \{a, c\}$, both black except possibly at (y, a). Conversely, axis-aligned $R_1 \neq R_2$ with equal presentations have $C(R_1) \triangle C(R_2) \subseteq S$. They cannot share opposite corners, so by symmetry any common corners lie in one row a. With none, R_1 is fully black. With two, (x, a) and (y, a), the other rows $b \neq c$ give a black four-cycle $r_b\, c_x\, r_c\, c_y$. With just (x, a), write $R_1 = \{x, y\} \times \{a, b\}$ and $R_2 = \{x, z\} \times \{a, c\}$ to obtain $r_a\, c_y\, r_b\, c_x\, r_c\, c_z$. □

Theorem 8 (row bound)**.** *Let $m \geq 3$. Every axis-valid blackout of the $h \times m$ grid has at most $m + q_h$ points, and at most $h + q_m$ if also $h \geq 3$. Hence $f(h, m) \leq m + q_h$.*

Join each nonempty column's black rows B_i by a tree T_i. Their union H reduces the problem to Mantel's theorem.

Lemma 9 (row graph)**.** *Let $m \geq 3$ and let S be axis-valid. For any choice of the trees, H is simple and triangle-free, and $e(H) = |S| - t$, where t is the number of nonempty columns.*

Proof. A repeated edge of H would put the same row pair in two columns, giving a four-cycle in G_S. Lemma 7 rules this out, so $e(H) = \sum(|B_i| - 1) = |S| - t$. Now suppose H contains a triangle $r_a r_b r_c$. Its edges cannot all belong to one tree. If two columns i, i' supply them, two of a, b, c belong to $B_i \cap B_{i'}$, again giving a four-cycle. If three columns supply them, they give the six-cycle $r_a\, c_i\, r_b\, c_{i'}\, r_c\, c_{i''}$. Both possibilities contradict Lemma 7. □

Lemma 10 (Mantel [20])**.** *Let H be a simple triangle-free graph on h vertices, let a vertex of maximum degree Δ have neighbor set A, and let B be the other $h - \Delta$ vertices, spanning $e(B)$ edges. Then*

$$e(H) = \sum_{b \in B} d(b) - e(B) \leq (h - \Delta)\Delta - e(B) \leq q_h.$$

Proof. The set A is independent, so every edge meets B. Summing degrees over B counts its internal edges twice. Subtract those once, then use $d(b) \le \Delta$ and $(h-\Delta)\Delta \le q_h$. □

Proof of Theorem 8. Combining Lemmas 9 and 10 gives $|S| = t + e(H) \le m + q_h$. The need for a third column enters only through Lemma 7. Transposing therefore gives $|S| \le h + q_m$ when $h \ge 3$. □

Theorem 11 (equality structure)**.** *Let $h \ge 4$ and $m \ge 3$. A blackout of the $h \times m$ grid with $m + q_h$ points is axis-valid if and only if*

(a) every column is a singleton or a double,

(b) the q_h doubles have distinct row pairs, and

(c) these pairs form a balanced complete bipartite graph on the rows.

There are $p_h\,(m)_{q_h}\,h^{m-q_h}$ such blackouts, where $(m)_q$ is the falling factorial and the number p_h of balanced bipartitions of the rows is $\binom{h}{\lfloor h/2 \rfloor}$, halved for even h.

Proof. Equality in the row-bound proof forces $t = m$, $e(B) = 0$, and $(h - \Delta)\Delta = q_h$, with all of B joined to all of A. Thus H is the graph in (c). If $|B_i| \ge 3$, its tree T_i has a vertex v with neighbors u, w. Since $h \ge 4$, the part containing v has another vertex z. Replacing vw by uw preserves uz, wz and creates triangle uwz, contradicting Lemma 9. This proves (a), and the tree edges, now the row pairs of doubles, give (b) and (c). Conversely, (a) and (b) exclude four-cycles. A six-cycle $r_a\,c_x\,r_b\,c_y\,r_c\,c_z$ needs doubles $\{a, b\}$, $\{b, c\}$, and $\{a, c\}$, contradicting (c). Apply Lemma 7. To count, choose the bipartition, assign edges to distinct columns, and choose the remaining singleton rows. □

Corollary 12 (small widths)**.** *Let $h \ge 4$ and $3 \le m < q_h$. Then $f(h, m) \le m + q_h - 1$. In particular $f(5, 5) \le 10$.*

Proof. Attaining $m + q_h$ would require $q_h \le m$ double columns by Theorem 11. Since this is impossible here, Theorem 8 improves by one. Here $q_5 = 6$. □

Put $W_h = 1 + \lfloor (h-1)^2/4 \rfloor$, so $W_3 = 2$, $W_4 = 3$, and $W_5 = 5$.

Lemma 13 (leftmost corner)**.** *A tilted rectangle R has a unique corner $\ell = (x, r)$ in its smallest column, and a unique corner in its smallest row. The sides from ℓ are $(a, -b)$ and (c, d) with positive integers a, b, c, d and $ac = bd$, and the corners are*

$$(x, r), \qquad (x + a,\, r - b), \qquad (x + c,\, r + d), \qquad (x + a + c,\, r - b + d). \quad (1)$$

Its rows are $r - b$ to $r + d$, and its span *is $a + c$.*

Proof. There are no vertical sides. Two corners in the leftmost column x would be opposite, forcing the other two there as well by their mean column. Likewise for rows. The two sides from ℓ therefore point right, one up and one down. Perpendicularity gives $ac = bd$. □

Lemma 14 (span bound)**.** *Every tilted rectangle of the $h \times m$ grid has span $a + c \le W_h$.*

Proof. The rectangle fits in h rows, so $b + d \le h - 1$. Also $(a - 1)(c - 1) \ge 0$, whence $a + c \le bd + 1 \le W_h$, using the integer bound $bd \le (b + d)^2/4$. □

Thus W_h is A033638$(h - 1)$. For $h \ge 3$ and $m > W_h$, the sides $(1, -b)$ and (bd, d) attain it when $b + d = h - 1$ and $|b - d| \le 1$. Here a rectangle's *height* and *width* mean its row and column ranges (the distances between its extreme coordinates).

Corollary 15 (the conjecture in A338885)**.** *For $n \ge 2$, the largest width of a tilted lattice rectangle of height n is $\lfloor n^2/4 \rfloor + 1$. So row n of A338885 has largest entry A033638(n).*

Proof. In (1), set the height $b + d = n$. As in the span bound, the width is $a + c \le bd + 1 \le \lfloor n^2/4 \rfloor + 1$. Choose sides $(1, -b)$ and (bd, d) with $b = \lfloor n/2 \rfloor$ and $d = \lceil n/2 \rceil$ to attain it. □

The *diamond* $D(i, j)$ is the square with corners $(i, j + 1)$, $(i + 1, j)$, $(i + 2, j + 1)$, and $(i + 1, j + 2)$. The rectangles A_i, B_i, C_i, and E_i have one corner in each of the columns i to $i + 3$, in the rows 3412, 3142, 2413, and 2143 from left to right.

Lemma 16 (tilted rectangles on narrow grids)**.** *The tilted rectangles of the $3 \times m$ grid are the diamonds $D(i, 1)$. Those of the $4 \times m$ grid are $D(i, 1)$, $D(i, 2)$, A_i, B_i, C_i, and E_i.*

Proof. All the listed sets are rectangles. On three rows, necessarily $b = d = 1$, hence $a = c = 1$ and the leftmost corner ℓ is in row 2. On four rows, $b = d = 1$ again gives diamonds. The only other possibility is $\{b, d\} = \{1, 2\} = \{a, c\}$, with ℓ in row $1 + b$. The four choices of sides, $(2, -2), (1, 1)$, $(1, -2), (2, 1)$, $(2, -1), (1, 2)$, and $(1, -1), (2, 2)$, give A, B, C, and E. □

Lemma 17 (boundary cross)**.** *The first row and the first column of the $h \times m$ grid form a valid blackout, so $f(h, m) \geq h + m - 1$.*

Proof. Only rectangles showing at most two points need checking (Lemma 4). An axis-aligned rectangle $\{x_1, x_2\} \times \{y_1, y_2\}$ shows $X \times Y$, where $X = \{x_1, x_2\} \setminus \{1\}$ and $Y = \{y_1, y_2\} \setminus \{1\}$. These recover its columns and rows, since $|X| = 1$ precisely when $1 \in \{x_1, x_2\}$, and likewise for Y.

By Lemma 13, a tilted rectangle can have only two black corners, ℓ and $(x + a, r - b)$. When both are black, $\ell = (1, 1 + b)$ and the shown corners are $p = (1 + c, 1 + b + d)$ and $q = (1 + a + c, 1 + d)$. Their coordinates recover a, b, c, d. They also differ in both coordinates, so an axis-aligned rectangle could show this pair only if its other two corners were black, which they are not. □

Lemma 18 (axis-aligned rivals)**.** *Let no axis-aligned rectangle be fully black. A* rival *of a set P of at most two white points is an axis-aligned rectangle that shows exactly P. The empty set has no rival, and P has one exactly when it is a singleton $\{(x, r)\}$ with $\{r, s\} \in F$ for some $s \in B_x$, a vertical pair $\{(x, r), (x, s)\}$ with $\{r, s\} \in F$, a horizontal pair $\{(x, r), (z, r)\}$ with $B_x \cap B_z \neq \emptyset$, or an oblique pair $\{(x, r), (z, s)\}$ with $x \neq z$, $r \neq s$, and $(x, s), (z, r) \in S$.*

Proof. A rival of $P \neq \emptyset$ has rows $\{r, s\}$ and columns $\{x, z\}$, with every corner outside P black. Reading those corners gives the listed conditions, which force $z \neq x$ and $s \neq r$ since (x, r) is white. □

5 Three rows

Take $m \geq 3$ with rows $1, 2, 3$ downward. The tilted rectangles are just $D(i, 1)$ (Lemma 16). The row bound and cross give $f(3, m) = m + 2$, also obtained by the Omega Institute [26]. Denote the columns by $a = \{1\}$, $b = \{2\}$, $c = \{3\}$, $A = \{1, 2\}$, $B = \{1, 3\}$, $C = \{2, 3\}$, and $X = \{1, 2, 3\}$, with a and c *outer*. A blackout without empty columns is a *word* $B_1 \cdots B_m$. We also use each double letter for its pair in F (Section 2).

Proposition 19 (the two types)**.** *Let S be a blackout of the $3 \times m$ grid with $m + 2$ points.*

(a) If S is valid, its word is one X among singletons (type T) or two different doubles among singletons (type P).

(b) For every word of type T or P, G_S is a tree, so axis-aligned rectangles have distinct presentations.

(c) There are $m3^{m-1} + \binom{m}{2} \cdot 6 \cdot 3^{m-2} = m^2 3^{m-1}$ words of type T or P.

Proof. (a) Lemmas 9 and 10 give $|S| = t + e(H)$, with $t \leq m$ and $e(H) \leq 2$. Equality forces nonempty columns and $\sum_i (|B_i| - 1) = 2$, hence one triple or two doubles. Equal doubles would give a four-cycle in G_S. (b) The nonsingleton columns connect the three rows, and each singleton adds a leaf. (c) Place the triple or ordered pair of distinct doubles, then choose each remaining singleton. □

The nonsingleton letters sort *candidates* into X, AC, CA, AB, BA, BC, and CB, read left to right. Use AC_m for valid width-m words in AC, and likewise for the rest.

Lemma 20 (diamond test)**.** *A candidate S with pair set F is valid if and only if every diamond $D(i, 1)$ passes the test below, where $P = D(i, 1) \setminus S$.*

(a) If $|P| \geq 3$ or P is empty, the diamond passes.

(b) If $P = \{(i + 1, 1), (i + 1, 3)\}$, the diamond fails if and only if $B \in F$.

(c) If $P = \{(i, 2), (i + 2, 2)\}$, it fails if and only if $B_i \cap B_{i+2}$ meets $\{1, 3\}$.

(d) If P is an oblique pair $\{(x, r), (z, s)\}$, it fails if and only if $(x, s), (z, r) \in S$.

(e) If $P = \{(x, r)\}$, it fails if and only if $\{r, s\} \in F$ for some $s \in B_x$.

Proof. Proposition 19(b) settles axis-aligned pairs, and Lemma 4 settles presentations of at least three points. An empty diamond has no axis-aligned rival since G_S has no four-cycle. Lemma 18 gives the remaining tests, with $B \in F$ in (b) and $\{r, s\} \in F$ in (e). Two diamonds share at most a tip. For equal presentations, both middle columns would need black rows 1 and 3, impossible in a candidate. □

A *window* is three consecutive letters, in columns i to $i+2$. It *passes* or *fails* with $D(i,1)$. Its *tips* $(i,2)$ and $(i+2,2)$ are black exactly when their letters are b, A, C, or X.

Lemma 21 (singleton dichotomy)**.** *In a candidate, a window of three singletons fails if and only if $B \in F$ and it is bab, bbb, or bcb.*

Proof. With two black tips, the vertical pair or one middle-column corner remains. Tests (b) and (e) reject exactly when $B \in F$. With one black tip, three corners remain or test (d) passes because $(i+1,2)$ is white. With neither, at least three corners remain. □

Lemma 22 (six of nine in the class $\{A,C\}$)**.** *Let S be a candidate with $F=\{A,C\}$, and let s,t be singletons.*

(a) A window sAt fails if and only if st is bb, bc, or cb, also for $t=C$ read as $t=b$.

(b) A window sCt fails if and only if st is bb, ba, or ab.

(c) A window with one double, at a tip, passes.

Proof. (a) Only $(i+1,3)$ is white in the middle column. The right tip $(i+2,2)$ is black for $t \in \{b,C\}$. With both tips black, test (e) fails since $2 \in B_{i+1}$ and $C \in F$. With just the right tip black, the pair $(i,2),(i+1,3)$ fails test (d) exactly for $s=c$. The other tip is symmetric. With neither black, three corners show. (b) Reflect across the middle row. (c) Apply Lemma 21 to Ast with the left tip already black. No case fails since $B \notin F$. Reflection and reversal give the remaining positions. □

A word in AC_m has form $UAVCW$, with singleton strings U, V, and W. By the diamond test it is enough that every window passes. Set $G_{AC}(z)=\sum_{m\geq 2}|AC_m|z^m$. Define G_{AB} by counting $UAVBW$, and $G_X(z)=\sum_{m\geq 1}|X_m|z^m$ by counting UXW. Write $R=1-3z$. A nonempty singleton string with a specified letter next to a double contributes z/R.

Proposition 23 (the class $\{A,C\}$)**.** *The generating function of AC is*

$$G_{AC}(z)=\frac{z^2-4z^3+10z^4-8z^5-3z^6}{(1-3z)^3}, \tag{2}$$

and

$$|AC_m|=2(m-1)(3m+2)3^{m-5} \qquad (m\geq 4). \tag{3}$$

Proof. Lemma 22(a) and the half turn allow tAC only for $t = a$ and ACt only for $t = c$. Each AtC passes tests (b) and (e). Lemmas 21 and 22(c) leave only windows centered on a double.

Split according to the gap V. With no gap, a nonempty U must end in a and a nonempty W must start in c, contributing $z^2(1 + z/R)^2$. For $V = t$, the allowed last letters of U number $p_t = 3, 1, 2$, and the allowed first letters of W number $q_t = 2, 1, 3$, for $t = a, b, c$. Thus $\sum p_t = \sum q_t = 6$ and $\sum p_t q_t = 13$, giving $z^3 \sum_t (1+p_t z/R)(1+q_t z/R) = z^3(3+12z/R+13z^2/R^2)$. For $|V| \geq 2$, each end contributes $3 + 6z/R$ (a lone endpoint letter, or one of six passing pairs). The unrestricted middle contributes $1/R$, so this case gives $9z^4(1 - z)^2/R^3$. Adding proves (2). Extract coefficients using $[z^m]z^k R^{-3} = \binom{m-k+2}{2} 3^{m-k}$ to obtain (3). □

Theorem 24 (the number of three-row optima)**.** *Let E_m count the optima of the $3 \times m$ grid with a column black in rows 1 and 3 (A398448). For $m \geq 4$,*

$$a_3(m) = 4(m - 1)(3m + 2)3^{m-5} + E_m. \tag{4}$$

Proof. A column black in rows 1 and 3 must be B or X. Proposition 19(a) therefore identifies these optima with the classes other than AC and CA. Reversing columns pairs AC, AB, and BC with CA, BA, and CB, respectively. Reflection across the middle row also pairs AB with CB. Hence

$$a_3(m) = 2|AC_m| + 4|AB_m| + |X_m| = 2|AC_m| + E_m \qquad (m \geq 3), \tag{5}$$

and substituting (3) gives (4). □

5.1 The generating function

Put $L = 1 - 2z - 2z^2$, $N = 1 + 2z^2$, and $D = LN$.

Theorem 25 (three-row generating function, parts (b) and (c) computer-assisted)**.**

(a) The series $\sum_{m \geq 3} a_3(m) z^m$ equals $2G_{AC} + 4G_{AB} + G_X - z - 12z^2$, with G_{AB} and G_X from Propositions 29 and 28, of denominators dividing D^3 and D^2.

(b) Its reduced denominator is exactly $Q = R^3 L^3 N^3$.

(c) With

$$\begin{aligned} P(z) = z &- 3z^2 - 34z^3 + 228z^4 - 641z^5 + 1307z^6 - 2186z^7 \\ &+ 2056z^8 - 236z^9 - 1060z^{10} + 2584z^{11} - 4264z^{12} \\ &- 1440z^{13} + 704z^{14} - 1024z^{15} + 2080z^{16} + 2176z^{17} + 384z^{18}, \end{aligned}$$

we have

$$\sum_{m\geq 3} a_3(m) z^m = \frac{P(z)}{Q(z)} - z - 12z^2. \tag{6}$$

Proof. For (a), sum (5), removing widths one and two by subtracting $z+12z^2$. Part (b) is proved in Section 5.2. For (c), exact algebra in SymPy [22] gives P. The script `three_row_closed_form_check.py` independently checks (6) against a transfer matrix through width 100. □

Let $\rho = 1 + \sqrt{3}$ and $\sigma = 1 - \sqrt{3}$.

Theorem 26 (three-row closed form, computer-assisted exact algebra)**.** *For every $m \geq 3$,*

$$E_m = p_+(m)\rho^m + p_-(m)\sigma^m + T_m, \tag{7}$$

where $432\, p_\pm(m) = (19 \pm 11\sqrt{3})m^2 + (61 \pm 41\sqrt{3})m + 104 \pm 28\sqrt{3}$ and $T_m = (-2)^{\lfloor m/2 \rfloor}\, \kappa(m)$ with $\kappa(m) = (2m^2 + 41m - 104)/216$ for even m and $\kappa(m) = (7m^2 - 32m - 175)/432$ for odd m. Hence $E_m \sim \frac{19+11\sqrt{3}}{432} m^2 \rho^m$. The hand proof gives $E_m > 0$ and

$$a_3(m) = \frac{4}{81} m^2 3^m - \frac{4}{243}(m+2)3^m + O(m^2\rho^m). \tag{8}$$

Proof. Equation (5) makes E_m the coefficient of z^m in $4G_{AB} + G_X$, with denominator dividing L^3N^3. The possible poles are $1/\rho$, $1/\sigma$, and $\pm 1/(i\sqrt{2})$, each of order at most three. Apart from a polynomial part of degree one, their coefficients have the form (7). Exact algebra determines the twelve rational constants. Both sides obey the order-12 recurrence from L^3N^3 for $m = 14$ onward and agree at $m = 2, \ldots, 13$. The leading term follows from $|\sigma| < 1$ and $|T_m| = O(m^2 2^{m/2})$. For the hand bound, the pole $1/\rho$ of least modulus gives $E_m = O(m^2\rho^m)$, which yields (8) through (4). Finally, the boundary cross has word $Xa\cdots a$ (Lemma 17), so $E_m > 0$. □

5.2 The classes AB and X

It remains to count X and AB and rule out cancellation in Theorem 25(b). Keep $D = LN = 1 - 2z - 4z^3 - 4z^4$. A *restricted* singleton string has no two b letters at distance two. Reversal and reflection carry AB to BA, BC, and CB.

Lemma 27 (forbidden windows in the classes containing B)**.** *In classes AB and X, the table lists exactly the failing windows. Here s is any singleton.*

class	*windows*	*forbidden*
AB, X	*singletons only*	bsb
AB	*one double*	bsA, Asb, bAb, bAc, cAb, aBa, aBb, bBa, cBc
	two doubles	bAB, ABa
X	*one triple*	bsX, Xsb, aXa, aXb, bXa, bXc, cXb, cXc

Proof. Lemma 21 handles singleton windows. For sAt, use Lemma 22(a), also rejecting bAb by test (e) since $B \in F$. In sBt or sXt, the middle corners are black. For $s = t = b$ the diamond is empty. For outer letters, test (c) fails exactly for $s = t$. For $s = b \neq t$, only $(i+2, 2)$ shows, and test (e) fails exactly when $\{2, t\} \in F$. In uAB, the black corners $(i+2, 3), (i+1, 2)$ make test (d) fail for $u = b$, while otherwise three corners show. In ABw, $w = b$ gives an empty diamond, and test (e) rejects $w = a$. Tests (b) and (e) exclude Asb and Xsb. Other left-tip windows show three corners or need the white point $(i+1, 2)$ for an oblique rival. Reflect for the other cases. □

Encode outer letters by bit 0 with weight 2, and b by bit 1. Let h_{ij} count continuations after i, j, forbidding b after $1, j$. Read each run $w_1 \cdots w_n$, $n \geq 1$, away from its double. Let g_0, g_1 count a prescribed outer first letter and $w_1 = b$, and u, v the same with w_2 outer if present. A fictitious preceding letter, outer for g and b for u, v, gives $(g_0, g_1, u, v) = z(h_{00}, h_{01}, h_{10}, h_{11})$. The recurrence solves to

$$Dg_0 = z(1+z+3z^2+2z^3), \quad Dg_1 = z(1+z), \quad Du = z(1+2z^2+2z^3), \quad Dv = z. \tag{9}$$

Proposition 28 (the triple class)**.** *With u and v as in (9), $G_X(z) = z\big(1 + 4u + 2v + 2u^2 + v^2\big)$, and its denominator divides D^2.*

Proof. In UXW, both runs are restricted. Forbidding bsX and Xsb makes the second letter from X outer. A single nonempty side gives $2u + v$, on either side. Two nonempty sides allow neighbor pairs (a, c), (b, b), and (c, a), contributing $2u^2 + v^2$. □

In $UAVBW$, all singleton runs are restricted. Reading outward, U from A uses u or v, and W from B uses g_0 or g_1 (Lemma 27). Include empty runs in ℓ_t and r_t when t follows A or precedes B. The double-centered windows give

$$(\ell_a, \ell_b, \ell_c) = (1{+}2u{+}v, 1{+}u, 1{+}2u),\ (r_a, r_b, r_c) = (1{+}g_0, 1{+}g_0{+}g_1, 1{+}g_0{+}g_1). \tag{10}$$

We also need the final bit. For $s \in \{00, 10\}$, let $J_{s\beta}$ count continuations from s ending in β, assigning the empty continuation to $\beta = 0$. The same recurrence gives the matrix below, with rows indexed by s and columns by β

$$J = \frac{1}{D}\begin{pmatrix} 1 + 2z^2 + 2z^3 & z + z^2 \\ 1 & 2z^2 + 2z^3 \end{pmatrix}. \tag{11}$$

Proposition 29 (the class $\{A, B\}$, polynomial form computer-assisted)**.** *With (9) to (11),*

$$\begin{aligned} G_{AB}(z) = {} & z^2(1 + 2u)(1 + g_0 + g_1) + z^3(\ell_a r_a + \ell_b r_b + \ell_c r_c) \\ & + z^4\,(\ell_a + \ell_c,\ \ell_b)\,J\,(r_a + r_c,\ 2r_b)^T. \end{aligned} \tag{12}$$

Moreover $G_{AB} = P_B/D^3$ with

$$P_B(z) = z^2(1 + z)(1 + z^2 - 3z^3 + 4z^6 + 4z^7). \tag{13}$$

Proof. Let k be the length of V. For $k = 0$, the boundary windows uAB and ABw require $u \neq b$ and $w \neq a$, giving the first term. For $k = 1$, AtB always passes, while uAt and tBw contribute $\ell_t r_t$.

For $k \geq 2$, Av_1v_2 forces v_2 to be outer, and $v_{k-1}v_kB$ always passes. The remaining string $v_3 \cdots v_k$ continues from bits 00 or 10, according as v_1 is outer or b, with boundary weight $\ell_{v_1} r_{v_k}$. The vector entry $r_a + r_c$ already counts the two outer choices for v_k. Thus for $k \geq 3$, the outer-letter weight 2 in J, and the factor 2 in $2r_b$, account for the two choices of v_2. The constant term of J handles $k = 2$. This gives (12), and exact algebra gives (13). □

Proof of Theorem 25(b). The denominator divides $R^3D^3 = Q$ by part (a). Its factors share no roots, since $D(\frac{1}{3}) \neq 0$ and $L - N = -2z - 4z^2$ vanishes only where $N \neq 0$. At $z = \frac{1}{3}$, G_{AC} alone has a triple pole, its numerator being $\frac{4}{81}$. At roots of D, the remainders $P_B \bmod L = \frac{z}{2}$ and $P_B \bmod N = \frac{1}{4} - \frac{z}{2}$ are nonzero. Thus G_{AB} has triple poles there, which neither the double poles of G_X nor the regular G_{AC} can cancel. □

Remark 30 (optima up to symmetry). Counting up to row reflection and column reversal gives $o_3(m)$, with first values 15, 58, 227, 864, 3190, and 11502 for widths 3 to 8 (A401226). Burnside's lemma says that $4o_3(m)$ equals $a_3(m)$ plus the fixed counts of the three nonidentity symmetries. Row reflection allows only b, B, and X. An optimum must therefore have one X and otherwise b. For $m \geq 4$, one of Xbb, bbX, or bbb occurs and fails (Lemmas 20(b) and 21). The only fixed optimum is bXb.

For the half turn, put $m = 2k$ or $m = 2k + 1$. It reverses the word and exchanges A with C and a with c. A fixed two-double word has A and C in mirror columns $j \leq k$ and $m + 1 - j$, with b at the center for odd m. The first k letters determine the rest. Only the windows centered on A and C can fail (the proof of Proposition 23), and these are mirror images. Lemma 22(a) leaves 3^{k-1} words in AC when $j = 1$, 3^{k-2} when $j = k \geq 2$, and $2 \cdot 3^{k-2}$ when $1 < j < k$. The count in CA is equal. Summing gives $4k3^{k-2}$ for $k \geq 2$. At $m = 3$ there are just AbC and CbA. Their combined series is $W = 2z^3 + \frac{4}{9}(1 + z)\big(3z^2(1 - 3z^2)^{-2} - 3z^2\big)$.

Column reversal forces a palindrome. Every noncentral letter is repeated, so a fixed optimum must have odd width and form UXU^{rev}. The middle window tXt requires $t = b$, and the forbidden windows bsX and Xsb force the next letter outward to be outer (Lemma 27). Thus U is counted by v in (9). For a half-turn-fixed triple word $UX\iota(U^{\mathrm{rev}})$, where ι reflects rows, the allowed middle pairs are (a, c), (b, b), and (c, a), giving $2u+v$ for U. Combine the contribution z^3 of bXb, the paired-run contribution $2z\big(u(z^2) + v(z^2)\big)$, and the two-double contribution W to get

$$4\sum_{m\geq 3} o_3(m)z^m = \sum_{m\geq 3} a_3(m)z^m + z^3 + \frac{4z^3(1 + z^4 + z^6)}{M(1 + 2z^4)} + \frac{2z^3(1 - z)(1 + 5z + 3z^2 - 3z^3)}{(1 - 3z^2)^2},$$

Here $M = 1 - 2z^2 - 2z^4$, so $D(z^2) = M(1 + 2z^4)$. The independent check

`three_row_orbits.py` agrees with the fixed counts through width 24 and the orbit counts $o_3(m)$ for $m \le 10$.

6 Four rows

Theorem 31 (four rows)**.** *For every $m \ge 6$ we have $f(4, m) = m + 4$.*

To attain the upper bound of Theorem 8, use

$$S = \{(i, 1) : 2 \le i \le m - 1\} \cup \{(m - 1, 2), (m, 2), (1, 3), (2, 3), (1, 4), (m, 4)\}. \tag{14}$$

Here $|S| = m+4$ and G_S is the eight-cycle $r_1\, c_2\, r_3\, c_1\, r_4\, c_m\, r_2\, c_{m-1}$ with leaves $c_3, \ldots, c_{m-2}$ on r_1.

Proof of Theorem 31. The graph G_S has no four- or six-cycle. Thus Lemma 7 distinguishes the axis-aligned rectangles, and none is fully hidden.

For tilted rectangles, use Lemma 16. Their corners in row 1 are black and those in row 4 are white. For $D(i, 1)$ the additional black corners are $(i + 2, 2)$ when $i \ge m - 3$ and $(i + 1, 3)$ when $i = 1$. For $D(i, 2)$ they are $(i, 3)$ when $i \le 2$ and $(i + 1, 2)$ when $i = m - 2 \ge 4$. For A_i and B_i they are $(i, 3)$ when $i \le 2$ and $(i + 3, 2)$ when $i \ge m - 4$. Three shown corners suffice, leaving the weak rectangles. Below, $s = 1$ for A_i and $s = 2$ for B_i.

rectangle	which i	corners shown
$D(1, 1)$		$(1, 2), (3, 2)$
$D(i, 1)$	$i \in \{m - 3, m - 2\}$	$(i, 2), (i + 1, 3)$
A_i or B_i	$i \in \{1, 2\}$, $i \notin \{m - 4, m - 3\}$	$(i + s, 4), (i + 3, 2)$
A_i or B_i	$i \in \{m - 4, m - 3\}$, $i \notin \{1, 2\}$	$(i, 3), (i + s, 4)$
A_i or B_i	$i = 2$, $m = 6$	$(i + s, 4)$

The first four cases have different shown row sets. Within any one, the point in row 2 or 3 determines i, and the row-4 point determines s. For rivals, no row r has both $(1, r)$ and $(3, r)$ black, ruling out a rival of $\{(1, 2), (3, 2)\}$. In the next three cases, the axis-aligned rectangle through the pair has a white corner, respectively $(i, 3)$, $(i + s, 2)$, or $(i, 4)$. Here we use $i \ge 3$, $i + s \le 4 < m - 1$, and $3 \le i \le m - 3$, respectively. The final case has $m = 6$ and one shown point $(z, 4)$ with $z \in \{3, 4\}$. Column z is black only in row 1, and no column is black in rows 1 and 4, so this singleton also has no rival. □

For $m \geq 4$, Theorem 11 leaves $3\,(m)_4\,4^{m-4}$ *candidates* of size $m+4$. Each has four doubles forming a four-cycle F on the rows, with a singleton in every remaining column.

Corollary 32 (widths four and five, computer-assisted)**.** *We have* $f(4,4) = 7$ *and* $f(4,5) = 8$.

Proof. The boundary cross gives the lower bounds. For the upper bounds, the candidate search finds an ambiguous pair in every one of the 72 and 1,440 cases, respectively. □

Theorem 33 (no hidden rectangle)**.** *For* $m \geq 4$*, no valid blackout of the* $4 \times m$ *grid hides a rectangle.*

Proof. By Lemma 2, it is enough to express each rectangle's corner set as $C(R_1) \triangle C(R_2)$ for $R_1 \neq R_2$. For an axis-aligned rectangle, use its two rows and pair each column x, y with a third column w. For $1 \leq i \leq m-3$, put $\mathsf{Q}_i = \{(i,2),(i,3),(i+3,2),(i+3,3)\}$. The tilted cases follow from

$$\begin{aligned} \mathsf{C}_i &= \mathsf{A}_i \triangle \mathsf{Q}_i, \qquad & \mathsf{E}_i &= \mathsf{B}_i \triangle \mathsf{Q}_i, \\ D(i+1,1) &= D(i,2) \triangle \mathsf{A}_i, \qquad & D(i+1,2) &= D(i,1) \triangle \mathsf{E}_i, \end{aligned} \tag{15}$$

since any of the three sets in an identity is the difference of the other two. The first line handles A_i, B_i, C_i, and E_i. For a diamond $D(j,1)$ or $D(j,2)$, use the second line with $i = j-1$ or $i = j$. One choice is always available, since $j \geq 2$ or $j \leq m-3$ when $m \geq 4$. □

Theorem 34 (forbidden-block bound)**.** *For every* $m \geq 6$ *we have* $a_4(m) \leq 3\,(m)_4\,4^{m-4}(53/64)^k$*, where* $k = \max(0, \lfloor m/4 \rfloor - 4)$*. So* $a_4(m) = O\big(m^4\,212^{m/4}\big)$.

Proof. First choose F, the four double columns, and their pairs, in $3\,(m)_4$ ways. Call four consecutive singleton columns a *block*, recorded by their black rows. Starting at column x, the four *template words* 2413, 3412, 2143, and 3142 are the corner rows of C_x, A_x, E_x, and B_x. Fix one template rectangle T with word $t_1t_2t_3t_4$.

Swapping t_p and t_q gives a *swap word*. It cannot occur, since T and the axis-aligned rectangle through $(x+p-1, t_p)$ and $(x+q-1, t_q)$ would show the same pair. The templates are exactly the permutations whose end letters both lie in $\{2,3\}$. Swaps within the end or middle positions give templates

again. Swaps between these positions give the sixteen permutations with one end letter in $\{2, 3\}$. Together these are 20 forbidden swap words.

An *edit word* changes one letter $t_p = r$ to s with $\{r, s\} \in F$. This is also impossible. The rectangle T and the axis-aligned rectangle on rows r, s and columns $x + p - 1$ and d, where d is the double column for $\{r, s\}$, would show only $(x + p - 1, r)$. There are 8 edits per template. Two templates can share an edit only when two positions contain r, s in opposite orders. They then share two edits if $\{r, s\} \in F$. This happens for two template pairs with $\{r, s\} = \{2, 3\}$ and two with $\{1, 4\}$. Hence at least $32 - 8 = 24$ edit words are forbidden. They repeat a letter and so are separate from the swap words. A block therefore allows at most $256 - 20 - 24 = 212 = 4^4 \cdot 53/64$ words.

Partition the columns into the $\lfloor m/4 \rfloor$ groups $4j - 3, \ldots, 4j$. At most four contain a double, leaving at least k singleton blocks. All other singleton columns have at most 4 choices, which gives the bound. □

Lemma 35 (six columns of memory)**.** *Let $m \geq 6$, and let F be the edge set of a four-cycle on the rows. A word over the four singleton letters and the pairs of F, each pair used once, represents an optimum if and only if (i) no tilted rectangle is hidden, (ii) no tilted rectangle shares its presentation with an axis-aligned rectangle whose columns are among its own and the double columns, and (iii) no two distinct tilted rectangles in a block of at most seven consecutive columns share a presentation.*

Proof. Necessity follows from validity and Theorem 33. Conversely, the prescribed doubles and singletons give an axis-valid word by Theorem 11. No axis-aligned rectangle is hidden, since that would give a four-cycle in G_S, and (i) excludes hidden tilted rectangles.

If two rectangles nevertheless show the same P, then $P \neq \emptyset$ and at least one is tilted. If the other is axis-aligned, a column of it either meets P or has two black corners. In the first case it is a column of the tilted rectangle, and in the second it is a double column. Thus (ii) excludes the pair. If both are tilted, choose a common column c meeting P. Each rectangle spans at most four columns, so their union fits in $c - 3$ to $c + 3$, contradicting (iii). The word is valid and has $m + 4$ black points, hence is an optimum. □

For a word built from singleton letters and pairs of F, a tilted rectangle *fails locally* in any of three cases. It is hidden, its presentation has a rival under Lemma 18 with pair set F, or another tilted rectangle shows the same set.

Proposition 36 (the automaton)**.** *Let F be the edge set of a four-cycle on the rows. The automaton $\mathcal{A}_F$ reads the four singleton letters and the pairs in F. It starts at $(\epsilon, \emptyset)$, where ϵ is the empty word. A state (u, U) records the suffix of at most six letters read so far and the set $U \subseteq F$ of pairs already used.*

On reading y, it rejects if $y \in U$ or if a tilted rectangle in uy ending in the new column fails locally. Otherwise it replaces u by the suffix of uy of length $\min(6, |uy|)$ and U by $U \cup (\{y\} \cap F)$. A word is accepted if it is read to the end without rejection and the final register is $U = F$. For each $m \geq 6$, the accepted words of length m are exactly the optima of the $4 \times m$ grid with pair set F.

Proof. Take $m \geq 6$. At its last column, each tilted rectangle lies in uy, since its span is at most three. An accepted word uses every pair of F once, so it is a candidate. Presentations of at least three points are already unique (Lemma 4). The local tests rule out hidden tilted rectangles and axis-aligned rivals, giving (i) and (ii) of Lemma 35. For (iii), a pair of tilted rectangles within seven columns is present when the later last column is read. That step compares their presentations. The memory lemma therefore makes an accepted word an optimum.

In the other direction, an optimum with pair set F uses every pair once. Theorem 33 excludes hidden rectangles, and validity excludes every repeated presentation. Applied with F, Lemma 18 cannot find a rival. The word passes each step and finishes with register F, so it is accepted. □

The script `four_row_equivalence_check.py` builds $\mathcal{A}_F$ (60,975, 138,095, and 55,575 states) and checks equivalence with a smaller machine from an earlier search. Including the pair register, that machine has 2,519, 3,089, and 2,119 reachable states. Two transfer counters agree for $m \leq 1500$, and direct candidate tests agree for $m \leq 10$ (Table 1).

m	6	7	8	9	10	11	12
$\{1,2\} \mid \{3,4\}$	0	4	128	920	4,852	24,112	114,424
$\{1,3\} \mid \{2,4\}$	20	256	1,780	11,128	63,540	346,320	1,785,768
$\{1,4\} \mid \{2,3\}$	6	32	210	1,284	7,006	32,224	137,656
$a_4(m)$	26	292	2,118	13,332	75,398	402,656	2,037,848

Table 1: Four-row optima by pair set (computer-assisted).

Theorem 37 (the four-row generating function, computer-assisted)**.** *There are coprime integer polynomials $P_4 = 26z^6 - 826z^7 + \cdots$ of degree 444 and $Q_4 = \sum_{j=0}^{420} q_j z^j$ of degree 420 with $q_0 = 1$ and $\sum_{m\ge 6} a_4(m) z^m = P_4/Q_4$. The recurrence $\sum_j q_j\, a_4(n-j) = 0$ holds for every $n \ge 445$ and fails at $n = 444$. Over the integers*

$$Q_4 = (1-z)^2(1+z)^2\, g_5^3\, \tilde{g}_5^3\, g_6^3\, g_{11}^5\, g_{17}^3\, g_{19}^3\, g_{20}^5\, g_{21}^5,$$

where g_d and $\tilde{g}_d$ are irreducible of degree d with constant term 1.

Proof. The automaton proposition identifies the counted words with the optima. Running Berlekamp–Massey modulo eight primes on $a_4(m)$ for $m \le 1500$ produces P_4 and Q_4. To check all widths, let T_F be the transfer matrix on reachable states and write $v(n) = v(0)T_F^n$. The state vectors satisfy $\sum_j q_j v(n_F - j) = 0$ at $n_F = 436$, 435, and 445 for the three machines. Powers of T_F proves the recurrence for all $n \ge 445$. Since $\deg P_4 = 444$, it fails at $n = 444$. The displayed factorization is obtained in SymPy. □

Corollary 38 (growth, computer-assisted)**.** *Let $\rho_4 = 3.351996777310\ldots$ be the largest real root of $p(x) = x^{20} g_{20}(1/x) = x^{20} - 4x^{19} + 2x^{18} + \cdots - 26x^2 - 4x + 8$. There is a polynomial B of degree 4 with $a_4(m) = \rho_4^m B(m) + O((1000/341)^m)$, so $a_4(m) \sim c\, m^4 \rho_4^m$ with $c \approx 8.1611 \times 10^{-5}$.*

Proof. Isolate the roots of every factor of Q_4 exactly. The only root of modulus at most 0.341 is $r = 1/\rho_4 = 0.29832\ldots$, simple in g_{20} and therefore of order five in Q_4. Separate its principal part as $P_4/Q_4 = \sum_{j=1}^{5} c_j (1 - z/r)^{-j} + E(z)$. The coefficients of E are $O((1000/341)^m)$. Coprimality gives $c_5 = H(r) \ne 0$, where $H(z) = P_4(z)(1-z/r)^5/Q_4(z)$. Now $[z^m](1-z/r)^{-j} = \rho_4^m \binom{m+j-1}{j-1}$, so $B(m) = \sum_j c_j \binom{m+j-1}{j-1}$ and $c = c_5/24$. Interval arithmetic encloses c between $8.1610970775 \times 10^{-5}$ and $8.1610970776 \times 10^{-5}$. □

7 Five and six rows

7.1 Boundary padding

For a blackout S of the $h \times m$ grid and $L, R \ge 0$, form $S_{L,R}$ by adding columns $1-L, \ldots, 0$ and $m+1, \ldots, m+R$, black only in a chosen *padding row*, row 1 or row h.

Theorem 39 (finite collar)**.** *Let $h \geq 3$, fix the padding row and the padded sides, and let L^* and R^* be W_h of Lemma 14 on padded sides and 0 otherwise. Then $S_{L,R}$ is valid for all $L, R \geq 0$ vanishing on unpadded sides if and only if S_{L^*,R^*} is valid.*

Proof. Set $S^* = S_{L^*,R^*}$. Necessity follows by this choice. Suppose S^* is valid and put $U = S_{L,R}$.

A tilted rectangle has at most one corner in row 1 and at most one in row h (Lemma 13 and (1)). A weak tilted rectangle of U therefore has a black corner in an original column e. Its span is bounded by Lemma 14, so its columns lie in $[e - W_h, e + W_h] \cap [1 - L, m + R] \subseteq [1 - L^*, m + R^*]$. It lies in S^* and has the same presentation there.

Suppose distinct R_1, R_2 in U show the same P.

Both axis-aligned. Every added column has degree one in G_U, so any cycle already lies in G_{S^*}. Since S^* is valid and has at least three columns, Lemma 7 makes U axis-valid.

Both tilted. Lemma 4 gives $|P| \leq 2$. Both rectangles are weak, so both lie in S^* with presentation P, contradicting its validity.

One of each. Take R_1 tilted, hence already in S^*. A column of R_2 meeting P also belongs to R_1. A column not meeting P has two black corners, so it must be original. Thus R_2 also lies in S^* with the same presentation, again a contradiction. □

7.2 Five rows


For the constructions, list the black rows in each column from left to right. Start with the blackout S_8 given by $14, 45, 25, 5, 5, 34, 23, 12$ of the 5×8 grid, so that $B_1 = \{1, 4\}$ and $B_4 = \{5\}$.

Theorem 40 (five rows, (a) and (c) computer-assisted, (b) checked in Lean)**.**


(a) We have $f(5, m) = m + 5$ for $m = 5$, 6, and 7.

(b) We have $f(5, m) = m + 6$ for every $m \geq 8$, attained by S_8 followed by $m - 8$ columns black in row 1 only.

(c) The 5×8 grid has exactly four maximum blackouts, the images of S_8 under row reflection and column reversal.



(a)



(b)

Figure 2: (a) The 5×13 test board of S_8. (b) The 6×16 blackout S_{16}.

Proof. (b) The upper bound is Theorem 8. For the construction, $W_5 = 5$, so the collar theorem with right padding in row 1 reduces the whole family from S_8 to Figure 2(a). Three enumerators and Lean verify that the 954 rectangles of this test board have different presentations. (a) At $m = 5$, use Corollary 12 and the construction S_0 in Theorem 44. At $m = 6, 7$, the constructions are $15, 12, 23, 3, 34, 45$ and $1, 12, 23, 3, 34, 45, 15$, recorded in `jis/data/small_cells_sat_5x6.json` and `jis/data/small_cells_sat_5x7_b12.json`. They give the lower bounds. Checked SAT certificates rule out $m + 6$ points (Section 9). (c) Two SAT enumerations find the four images. A separate column search checks all $10\,(8)_6\,5^2 = 5{,}040{,}000$ equality candidates from Theorem 11 and finds the same four. □

Theorem A gives $f(h, m) \le h + m$ whenever a side is at most four, and on 5×5, 5×6, 5×7, and 6×6. Up to transposition, these exhaust the grids of area below 40. Thus 5×8 is the first failure of the guess $f(h, m) = h + m$.

Remark 41 (five-row counts, computer-assisted). For $m = 13, \ldots, 23$ the

counts $a_5(m)$ are

$$193{,}406,\quad 1{,}346{,}758,\quad 8{,}650{,}416,\quad 52{,}128{,}242,$$
$$299{,}349{,}210,\quad 1{,}653{,}637{,}492,\quad 8{,}828{,}301{,}226,\quad 45{,}743{,}165{,}894,$$
$$230{,}934{,}909{,}328,\quad 1{,}139{,}486{,}126{,}866,\quad 5{,}510{,}071{,}794{,}736.$$

Each count agrees in two independent transfer calculations using the equality structure of Theorem 11.

7.3 Six rows and small grids

Let S_{16} be the blackout of the 6×16 grid in Figure 2(b). Its doubles, in the columns $1, 2, 3, 6, 9, 10, 14, 15, 16$, carry the edges of $K_{3,3}$ between $\{1, 3, 5\}$ and $\{2, 4, 6\}$.

Theorem 42 (six rows, (a) computer-assisted, (b) checked in Lean)**.**

(a) For $8 \le m \le 14$ we have $f(6, m) \le m + 8$.

(b) For every $m \ge 15$ we have $f(6, m) = m + 9$.

Proof. (a) Corollary 12 handles $m = 8$. For $9 \le m \le 14$, an optimum with $m+9$ points would have double columns carrying $K_{3,3}$, by Theorem 11. There are ten possible row partitions. The structured encoding has checked LRAT certificates excluding every one (Lemma 47). (b) Again the row bound gives the upper bound. For $m = 15$, take $56, 45, 34, 23, 1, 1, 14, 1, 1, 12, 36, 1, 25, 6, 16$ from `jis/data/six_row_sat_w15_A.json`, which is valid with 24 points. For $m \ge 16$, apply the collar theorem to S_{16} in Figure 2(b). Since $W_6 = 7$, it suffices to add $(17, 1), \ldots, (23, 1)$. Three enumerators and Lean verify distinct presentations for all 4,627 rectangles of that test board. □

Theorem 43 (small grids, computer-assisted)**.** *We have $f(6, 6) = 12$, $f(6, 7) = 13$, $f(7, 7) = 14$, and $f(6, 8) = 15$. The blackout $15, 1, 123, 34, 4, 4, 456, 26$ of the 6×8 grid attains the last value.*

Proof. The other constructions are $14, 1, 12, 2, 23, 3456$ on the 6×6 grid, $16, 1, 1, 12, 2, 23, 3456$ on 6×7, and $17, 1, 12, 2, 23, 3456, 67$ on 7×7. Their records are `jis/data/small_cells_sat_6x6.json`, `jis/data/small_cells_sat_6x7.json`, and `jis/data/small_cells_sat_7x7.json`. All are valid. For each grid, a checked SAT certificate excludes one additional point. □

Theorem 44 (the 5×5 grid)**.**

(a) We have $f(5,5) = 10$.

(b) (Computer-assisted.) The $a_5(5) = 8$ maximum blackouts are the images of S_0 below under the square's symmetries.

Proof. (a) Corollary 12 gives the upper bound. To attain it, take S_0 to be $45, 34, 123, 1, 15$. Its row-column graph is an 8-cycle with two pendant edges, hence it is axis-valid. Lemma 13 lists 30 tilted rectangles, namely nine unit diamonds, sixteen others from Lemma 16, and five spanning the grid. The fifteen weak rectangles show the pairs below. After an oblique pair, the point following the bar is a white corner of the axis-aligned rectangle through it.

$(1,2),(2,1) \mid (1,1)$	$(1,3),(2,2) \mid (1,2)$	$(3,4),(4,3) \mid (4,4)$	$(2,5),(3,4) \mid (3,5)$
$(2,2),(4,2)$	$(1,2),(4,3) \mid (1,3)$	$(4,4),(5,2) \mid (4,2)$	$(3,5),(4,4) \mid (3,4)$
$(4,3),(5,2) \mid (4,2)$	$(3,4),(5,2) \mid (5,4)$	$(2,5),(5,2) \mid (2,2)$	$(4,5),(5,3) \mid (4,3)$
$(1,3),(4,2) \mid (1,2)$	$(2,5),(4,3) \mid (4,5)$		$(4,5),(5,4) \mid (4,4)$

These pairs are all different. The other tilted rectangles show at least three points, so Lemma 4 distinguishes them. The listed white corners rule out axis-aligned rivals for every oblique pair. The remaining pair $(2,2),(4,2)$ has no rival because $B_2 \cap B_4 = \emptyset$. This proves validity. (b) Three programs check all 44,640 sets of 10 points whose graph has no four- or six-cycle. Only the images of S_0 are valid. □

8 Every fixed height

Fix the height h and read the grid from left to right. Its blackout is the *column word* $w = B_1 \cdots B_m$, whose letter B_i records the black rows in column i. Write $|w| = m$ and $S(w)$ for its blackout, and let $\mathcal{L}_h$ be the valid words, including the empty word. We first show that validity needs only finite memory. A bounded counter will then pick out the words with the most black points.

Theorem 45 (regularity and rational series)**.** *For every $h \geq 1$ the language $\mathcal{L}_h$ is regular, and $G_h(z,u) = \sum_{w \in \mathcal{L}_h} z^{|w|} u^{|S(w)|}$ is rational in z and u.*

Proof. The *trace* of a rectangle records, in each column, the rows of its corners. It determines the corner set. By Lemma 2, two distinct traces Y, Y' give the same presentation exactly when $Y_i \triangle Y_i' \subseteq B_i$ in every column.

The possible traces form a regular language. For an axis-aligned rectangle, choose $r < s$, read a column $\{r, s\}$, then another such column after any number of empty columns. There can also be empty columns before and after. For a tilted rectangle, Lemmas 13 and 14 leave only finitely many templates (1), each of span at most W_h, again with arbitrary empty columns on either side.

An automaton can therefore read w while guessing two rectangle traces. It checks the containment above and remembers whether the traces have differed anywhere. Accepting when they differ recognizes the invalid words. Regular languages are closed under complementation, so $\mathcal{L}_h$ is regular too.

Choose a deterministic automaton for $\mathcal{L}_h$ and give a transition reading B the weight $u^{|B|}$. If $M(u)$ is its weighted transition matrix, α its start vector and β its accepting vector, then

$$G_h(z, u) = \alpha(I - zM(u))^{-1}\beta.$$

Each word is counted once because the automaton is deterministic. Cramer's rule makes the expression rational [10, Proposition V.6]. □

Theorem 46 (fixed height)**.** *Let $h \geq 2$. (a) There are integers $m_0 \geq 3$, $p \geq 1$, and $k_0, \ldots, k_{p-1}$ such that $f(h, m) = m + k_{m \bmod p}$ for every $m \geq m_0$. (b) The series $\sum_{m \geq 0} a_h(m) z^m$ is rational, with the conventions of Section 2, and the words of optimum blackouts form a regular language.*

Proof. Put $E(w) = |S(w)| - |w|$, the number of black points beyond one per column. Every consecutive block v of a valid word is valid on its own grid. Theorem 8 bounds $E(v)$ by q_h when $|v| \geq 3$. For a shorter block, $|S(v)| \leq h|v|$ gives the common bound

$$E(v) \leq C_h := \max(q_h, 2h - 2).$$

If a valid word $w = uv$ has excess k, applying this to the prefix and suffix gives

$$k - C_h \leq E(u) = k - E(v) \leq C_h.$$

Thus, for a fixed k, only finitely many prefix excesses need to be remembered. Add this counter to an automaton for $\mathcal{L}_h$, and accept when the final excess is k. It recognizes the regular language $\mathcal{L}_{h,k}$ of valid words with that excess.

The set of lengths of a regular language is eventually periodic. For $m \geq 3$, Lemma 17 and Theorem 8 put the optimal excess between $h-1$ and q_h. There

are only finitely many such k, so choose one common period p and threshold m_0 for their length sets. The largest available excess then depends only on $m \bmod p$. Calling it k_r proves (a).

For each residue r, keep the words of $\mathcal{L}_{h,k_r}$ whose lengths are at least m_0 and congruent to r modulo p. This is still a regular language. Their union, together with the finitely many shorter optimum words, is exactly the language of optimum blackouts. Its counting series is rational by the same transfer-matrix argument as above, proving (b). □

9 Computations and verification

Every computed maximum and exact count has two separately written checks. The bound $a_7(8) \geq 18$, obtained by one enumeration. The repository [27] stores the code and evidence. All paths here start from `kagey-problems/problem001`. For commands, SHA-256 hashes, and running times, see `jis/scripts/README.md`. Table 3 connects each computer-assisted claim to its checks.

We solve the hitting-set problem of Lemma 2 by ILP, using CBC [11] through PuLP [23], and check it by brute force. SAT enumeration uses PySAT [15]. The column searches grow the board from left to right. They use $f(6,6) \leq 12$ to prune 6×7 and 6×8, and $f(6,7) \leq 13$ to prune 6×8 and 7×7. Every saved construction and counted model is tested for validity.

As a check on rectangle generation, the totals match A085582 for squares and A289832 for rectangular grids.

Encoding A prevents the inclusion-minimal differences $C(R_1) \triangle C(R_2)$, $R_1 \neq R_2$, from being entirely black. Encoding B forbids $C(R_1) \triangle C(R_2)$ only for distinct rectangles with a common corner. Each rectangle R also has an indicator y_R, forced true if it is hidden, and at most one y_R may be true.

For the structured six-row instances, fix a partition $A \mid B$ into triples. A column type is its black-row set, either a singleton or one of the nine pairs meeting A and B. Use each pair once. Of encoding B's clauses, retain those involving a tilted rectangle whose forbidden set can be all black. Cardinality is enforced by Sinz's sequential counter [29], expanded by our generator for certified instances, or by a totalizer [3]. Both enforce exactly the specified size bound.

Lemma 47 (faithful encodings)**.** *Let S be a blackout of the $h \times m$ grid, and let $\mathcal{D} = \{C(R_1) \triangle C(R_2) \mid R_1 \neq R_2\}$, where R_1 and R_2 range over rectangles*

grid	f	$a_h(m)$	by	grid	f	$a_h(m)$	by	grid	f	$a_h(m)$	by
2×2	4	1	HIB	3×8	10	45,864	IB	5×8	14	4	TSDC
2×3	4	12	HIB	3×9	11	161,992	IB	5×9	15	44	TSC
2×4	5	32	HIB	3×10	12	562,912	ITD	5×10	16	350	TSC
2×5	6	80	IB	4×4	7	316	IB	5×11	17	3,004	TSC
2×6	7	192	TI	4×5	8	1,810	IB	5×12	18	24,706	TSCM
2×7	8	448	TI	4×6	10	26	IB	6×6	12	160	ISC
3×3	5	53	IB	4×7	11	292	IB	6×7	13	2,536	SC
3×4	6	224	IB	4×8	12	2,118	IMD	6×8	15	8	SC
3×5	7	892	IB	5×5	10	8	IB	6×15	24	–	TS
3×6	8	3,420	IB	5×6	11	76	ISC	7×7	14	7,300	SC
3×7	9	12,704	IB	5×7	12	1,184	ISC	7×8	16	≥ 18	S

Table 2: Computed values for $h \leq m$. H: hand, T: theorem, I: ILP (value only on 3×10, 4×8, 5×6, 5×7, and 6×6), B: brute force, D: equality candidates, C: column search, S: two SAT encodings, with complete enumeration for exact counts, M: transfer counter, –: not established. On 7×8 only a lower bound for the count is given.

of this grid.

(a) S is valid if and only if it contains no inclusion-minimal member of $\mathcal{D}$.

(b) S is valid if and only if it hides at most one rectangle and contains no $C(R_1) \triangle C(R_2)$ for distinct rectangles R_1, R_2 sharing a corner.

(c) Suppose $h = 6$, $m \geq 3$, and S is valid with $m + 9$ points. For some partition of the rows into two triples, selecting the black-row set B_i as the type of column i extends to a satisfying assignment of the structured instance.

Proof. (a) Any member of the finite family $\mathcal{D}$ contains an inclusion-minimal member. Thus a blackout contains a forbidden difference exactly when it contains a minimal one. Apply Corollary 3.

(b) Part (a) leaves only pairs with no shared corner to check. For these, the difference is $C(R_1) \cup C(R_2)$, contained in S exactly when both rectangles are hidden. Allowing at most one hidden rectangle excludes precisely these remaining pairs.

(c) Theorem 11 gives a partition into two row triples. Every B_i is a singleton or a pair crossing it, and all nine pairs occur once. Use these as the column types. Set y_R to 1 exactly for hidden rectangles. By (b), these choices satisfy encoding B. Dropping clauses preserves satisfaction in the structured instance. □

Reading the point variables of A or B therefore gives precisely the valid blackouts of the chosen size. For six rows, part (c) and the row bound imply $f(6, m) \leq m + 8$ once the ten structured instances of width m are all unsatisfiable.

CaDiCaL [5] produces LRAT certificates for $f(5, 6) < 12$, $f(5, 7) < 13$, $f(6, 6) < 13$, $f(6, 7) < 14$, $f(7, 7) < 15$, $f(6, 8) < 16$, and $f(7, 8) < 17$ using encoding A on all blackouts. It also certifies the 70 structured six-row instances, for widths 8 to 14 and ten partitions per width, in Theorem 42(a). The clauses agree between two independent generators. We split 6×8 by its first column into 64 cases, and 7×7 and 7×8 into 128 each. The second generator checks that these splits exhaust the first-column choices. Every one of the 394 proofs passes both drat-trim's `lrat-check` [6, 31] and the Lean LRAT checker.

The constructions in `jis/data/table_counts/sat_7x8_16.json` give $f(7, 8) \geq 16$, completing $f(7, 8) = 16$. The separate enumeration `jis/data/table_counts/sat2split_7x8_16.json` gives 18 valid constructions with 16 points. This proves only $a_7(8) \geq 18$.

statement	method; independent check	evidence
Cor. 32, Thm. 44(b)	candidates; brute force	`verify_narrow.py`
Thms. 25, 26, Rem. 30	SymPy; transfer matrix	`three_row_*.json`
Thm. 37, Cor. 38, Table 1	Prop. 36; candidates	`four_row_*.json`
Thms. 40(a), 43	encoding A, LRAT; encoding B	`unsat_certificates.json`, `small_cells_*`
Thms. 40(b), 42(b)	collar board; enumerators, Lean	`five_row_*family*`, `six_row_*w15*`
Thm. 40(c), Rem. 41	SAT, counter; candidate search, second counter	`five_row_*`
Thm. 42(a)	structured, LRAT; encoder, search	`unsat_certificates.json`, `six_row_sat_w*`
Table 2, open question 2	SAT; column search; collar test	`table_counts/*`, `six_row_collar_*`

Table 3: Computer-assisted statements, their method and check, and the evidence. Evidence files are in `jis/data`, and `verify_narrow.py` is in the problem folder.

The formal proofs are in `../lean/KageyProblems/Problem001`. They cover Lemmas 2, 4, and 14, Corollary 3, the strip theorem, both orientations of the row bound, and the collar theorem. They also show that at most one rectangle can be hidden.

Kernel-checked validity tests cover S_0, S_8, the construction in Theorem 43, the 6×15 construction in Theorem 42, and the 5×13 and 6×23

collar boards. The kernel also checks that $\{(1,2),(2,1),(3,2),(2,3)\}$ hides $D(1,1)$ on 4×3. Thus the restriction $m \geq 4$ in Theorem 33 is necessary. Together these prove $f(5,m) = m+6$ for $m \geq 8$, $f(6,m) = m+9$ for $m \geq 15$, $f(5,5) \geq 10$, and $f(6,8) \geq 15$ from only Lean's three standard axioms.

The remaining unformalized parts are Lemma 13, Theorem 11, $f(5,5) \leq 10$, Lemma 47, the other results in Sections 3 to 8, and Appendix A.

10 Open questions

(1) What is the rational series $\sum_{m \geq 0} a_5(m) z^m$ guaranteed by Theorem 46, and the growth rate of $a_5(m)$? See the counts in Remark 41.

(2) A valid 6×9 blackout of 16 points passes the collar test with seven columns black only in row 1 added on each side (`six_row_collar_6x9_check.py`). Theorems 39 and 42 therefore give $m+7 \leq f(6,m) \leq m+8$ for $9 \leq m \leq 14$. Which value holds at each width?

(3) For $h \geq 7$, does $f(h,m) = m + q_h$ hold eventually? At height seven, a valid $7 \times m$ construction with $m+12$ points would suffice if it passes the collar test (Theorem 39). This would settle $h = 7$.

(4) Lemma 7 and Neuwirth's bound for bipartite graphs of girth eight [25, Corollary 16] give $f(n,n) < n^{4/3} + \frac{2}{3}n - \frac{2}{9}n^{2/3} - \frac{20}{81}n^{1/3}$ for $n \geq 3$, and Theorem 50 gives $f(n,n) \geq 0.29\, n^{6/5}(1+\ln n)^{-2/5}$ for large n. Lavrov [16] used the same method for the triangular version. What is the true order of growth? What is the first n with $f(n,n) > 2n$?

11 Acknowledgments

I thank Peter Kagey for posing the problem, and for discussing its early stages with me, including the uniqueness of the 5×5 optimum and the first graph-theoretic approach to the narrow grids. The design of Figures 1 and 2 follows the figure in his posting of Problem 001 [18].

A note on the use of AI

I began this work in January of 2026 and wrote the first draft and its mathematics between August and October, together with most of the Python code. I used Claude Opus 5.5 (Anthropic) for the October 17–19 revision, including the cut of a draft from 45 to 30 pages and the language passes. It did not add any language, just proposed a cut version. Using this as reference, I rewrote the text myself and implemented the revisions as I saw fit. No sentence in this paper was written by an LLM. It also drew the figures in the style of Kagey's posting of the problem given my own hand-drawn sketches.

I used it to check my Python code, and it wrote further checks of the computer-assisted results. One of these, `six_row_collar_6x9_check.py`, gives the lower bound in the second open question of Section 10. It also wrote the Lean files, the LRAT certificate pipeline and the formal checks described in Theorem A and Section 9. The formalization did not change a statement of the paper or find an error or missing hypothesis. It changed how several results are established, as Section 9 describes.

I also used Claude Opus 5.5 to search the literature beyond the sources I had found. Several references came from that search. I read every cited source, including the parts supporting each citation. Later reviews with Claude Opus 5.5 and OpenAI Codex on October 23rd pointed out problems that I checked and corrected, such as typos and inconsistencies or redundancies. I checked these revisions and implemented them all myself, and take full responsibility for the contents of the paper.

A Square grids

For squares, $f(n, n)$ eventually exceeds every fixed multiple of n. Sample black points, then delete points to distinguish the rectangles.

Fix $n \geq 2$ and put $\Lambda = 1 + \ln n$. Write $\mathcal{R}$ for all rectangles on the $n \times n$ grid, including tilted ones, and d_v for the number through corner v.

Lemma 48. *If rectangles $R_1 \neq R_2$ have exactly two common corners p and q, and $\{p, q\}$ is a side of both or a diagonal of both, then $C(R_1) \triangle C(R_2)$ is the corner set of a rectangle.*

Proof. For a common side, the remaining corners $p+u, q+u$ and $p+v, q+v$ have $u \neq v$ nonzero and perpendicular to $q - p$. They form a rectangle with

sides $q-p$ and $v-u$. For a common diagonal, write them as $o \pm u, o \pm v$, with $o = (p+q)/2$, $|u| = |v|$, and $v \neq \pm u$. The equal segments $[o-u, o+u]$ and $[o-v, o+v]$ share a midpoint, so they are diagonals of a rectangle. □

Let $r_2(t)$ be the number of integer pairs (a, b) satisfying $a^2 + b^2 = t$, and $\tau(t)$ the divisor count of t. We use the estimates $r_2(t) \leq 4\tau(t)$ and, for each $\varepsilon > 0$, $\tau(t) \leq K_\varepsilon t^\varepsilon$ for $t \geq 1$, with a constant K_ε [14, Sections 16.9, 16.10, and 18.1].

Collect two kinds of four-point obstructions in $\mathcal{A}$, namely the full corner sets $C(R)$ for $R \in \mathcal{R}$, and the differences $C(R_1) \triangle C(R_2)$ when exactly two shared corners are a side of one rectangle and a diagonal of the other. Let $\mathcal{B}$ contain the differences $C(R_1) \triangle C(R_2)$ for pairs with exactly one shared corner.

Lemma 49.

(a) Each point v has $d_v \leq 8n^2\Lambda$, and $|\mathcal{R}| \leq 2n^4\Lambda$.

(b) For every $\varepsilon > 0$, $|\mathcal{A}| \leq A_\varepsilon n^{4+2\varepsilon}\Lambda$ with $A_\varepsilon = 2(1 + 16 \cdot 2^\varepsilon K_\varepsilon)$, and $|\mathcal{B}| \leq 32n^6\Lambda^2$.

(c) A blackout that contains no member of $\mathcal{A} \cup \mathcal{B}$ is valid.

Proof. (a) Write the sides at v as $k(a, b)$ and $l(-b, a)$, with $\gcd(a, b) = 1$, $k \geq 1$, and $l \neq 0$. For $d = \max(|a|, |b|)$, there are $8d$ primitive directions, at most $(n-1)/d$ choices for k, and $2(n-1)/d$ for l. This counts each rectangle twice, giving $2d_v \leq \sum_{d=1}^{n-1} 16(n-1)^2/d \leq 16n^2\Lambda$. Sum over corners to get $4|\mathcal{R}| = \sum_v d_v \leq 8n^4\Lambda$. (b) With diagonal $\{p, q\}$ fixed, the other diagonal determines the rectangle. It is an integer vector with squared length $t = |p-q|^2 < 2n^2$. Hence a side of any $R \in \mathcal{R}$ is the diagonal of at most $r_2(t) \leq 4K_\varepsilon(2n^2)^\varepsilon$ rectangles. Counting the four sides gives $|\mathcal{A}| \leq |\mathcal{R}|(1 + 16K_\varepsilon 2^\varepsilon n^{2\varepsilon}) \leq A_\varepsilon n^{4+2\varepsilon}\Lambda$. For $\mathcal{B}$, count the pairs of rectangles through a common corner, giving $|\mathcal{B}| \leq \sum_v \binom{d_v}{2} \leq \frac{n^2}{2}(8n^2\Lambda)^2$. (c) An invalid S contains $C(R_1) \triangle C(R_2)$ for some $R_1 \neq R_2$ by Lemma 2. They share at most two corners (Lemma 4). With two shared corners, the difference belongs to $\mathcal{A}$, either by Lemma 48 or by definition. With one, it belongs to $\mathcal{B}$. With none, $C(R_1) \subseteq S$ itself belongs to $\mathcal{A}$. □

Theorem 50 (deletion method)**.** *We have*

$$\liminf_{n\to\infty} \frac{f(n,n)\,(1+\ln n)^{2/5}}{n^{6/5}} \geq \frac{5}{6 \cdot 192^{1/5}} = 0.2911\ldots.$$

In particular $f(n,n) - 2n \to \infty$.

Proof. Use the deletion method of Alon and Spencer [2, Section 3.2]. Set $a = 192^{-1/5}$, $T = n^{6/5}\Lambda^{-2/5}$, and $\theta = a\,n^{-4/5}\Lambda^{-2/5} \in (0,1)$. Choose each point independently with probability θ. Let X count chosen points, and let Y and Z count fully chosen members of $\mathcal{A}$ and $\mathcal{B}$, respectively. Then

$$\mathbb{E}(X - Y - Z) = n^2\theta - |\mathcal{A}|\theta^4 - |\mathcal{B}|\theta^6.$$

Here $n^2\theta = aT$. From Lemma 49(b), with $\varepsilon = 1/10$, the two losses are bounded by $|\mathcal{B}|\theta^6 \le 32a^6T$ and $|\mathcal{A}|\theta^4 \le A_{1/10}a^4n^{-1/5}T$. Since $32a^5 = \frac{1}{6}$, some chosen set S' satisfies $X - Y - Z \ge (\frac{5}{6}a - A_{1/10}a^4n^{-1/5})T$. Remove one point from each forbidden member of $\mathcal{A}\cup\mathcal{B}$ contained in S'. At most $Y + Z$ points are lost, leaving at least $X - Y - Z$. Lemma 49(c) makes the remainder valid. Consequently $f(n,n)/T \ge \frac{5}{6}a - A_{1/10}a^4n^{-1/5} \to 5/(6\cdot 192^{1/5})$. Finally $T/n \to \infty$, so $f(n,n) - 2n \to \infty$. □

References

[1] S. Alanko, S. Crevals, A. Isopoussu, P. R. J. Östergård, and V. Pettersson, Computing the domination number of grid graphs, *Electron. J. Combin.* **18** (2011), Paper P141.

[2] N. Alon and J. H. Spencer, *The Probabilistic Method*, 4th ed., Wiley, 2016.

[3] O. Bailleux and Y. Boufkhad, Efficient CNF encoding of Boolean cardinality constraints, in F. Rossi, ed., *Principles and Practice of Constraint Programming – CP 2003*, Lecture Notes in Comput. Sci., Vol. 2833, Springer, 2003, pp. 108–122.

[4] T. Bellitto, Separating codes and traffic monitoring, *Theoret. Comput. Sci.* **717** (2018), 73–85.

[5] A. Biere, T. Faller, K. Fazekas, M. Fleury, N. Froleyks, and F. Pollitt, CaDiCaL 2.0, in A. Gurfinkel and V. Ganesh, eds., *Computer Aided Verification, CAV 2024*, Lecture Notes in Comput. Sci., Vol. 14681, Springer, 2024, pp. 133–152.

[6] L. Cruz-Filipe, M. J. H. Heule, W. A. Hunt, Jr., M. Kaufmann, and P. Schneider-Kamp, Efficient certified RAT verification, in L. de Moura, ed., *Automated Deduction – CADE 26*, Lecture Notes in Comput. Sci., Vol. 10395, Springer, 2017, pp. 220–236.

[7] D. de Caen and L. A. Székely, The maximum size of 4- and 6-cycle free bipartite graphs on m, n vertices, in G. Halász, L. Lovász, D. Miklós, and T. Szőnyi, eds., *Sets, Graphs and Numbers (Budapest, 1991)*, Colloq. Math. Soc. János Bolyai, Vol. 60, North-Holland, 1992, pp. 135–142.

[8] D. de Caen and L. A. Székely, On dense bipartite graphs of girth eight and upper bounds for certain configurations in planar point-line systems, *J. Combin. Theory Ser. A* **77** (1997), 268–278.

[9] R. Euler, The Fibonacci number of a grid graph and a new class of integer sequences, *J. Integer Sequences* **8** (2005), Article 05.2.6.

[10] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.

[11] J. Forrest et al., coin-or/Cbc, version 2.10.3, software, Zenodo, 2019, https://doi.org/10.5281/zenodo.3246628.

[12] F. Foucaud, Decision and approximation complexity for identifying codes and locating-dominating sets in restricted graph classes, *J. Discrete Algorithms* **31** (2015), 48–68.

[13] J. Guilhot, E. Little, and J. Parkinson, Bounded weight functions on regular languages and groups, *Groups Geom. Dyn.*, online 27 August 2026, https://doi.org/10.4171/GGD/983. Theorem numbers from arXiv:2510.17061v1.

[14] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, 2008.

[15] A. Ignatiev, A. Morgado, and J. Marques-Silva, PySAT: a Python toolkit for prototyping with SAT oracles, in O. Beyersdorff and C. M. Wintersteiger, eds., *Theory and Applications of Satisfiability Testing – SAT 2018*, Lecture Notes in Comput. Sci., Vol. 10929, Springer, 2018, pp. 428–437.

[16] P. Kagey, Removing points from a triangular array without losing information, Mathematics Stack Exchange, question 2465571, 2017, https://math.stackexchange.com/q/2465571.

[17] P. Kagey, Open problem collection, Problem 1, G4G14 gift exchange, 2020, https://www.gathering4gardner.org/g4g14gift/G4G14-PeterKagey-OpenishProblems.pdf.

[18] P. Kagey, Problem 001, Open Problem Collection, https://peterkagey.com/problems/001/ (accessed October 2026).

[19] G. Katona, On separating systems of a finite set, *J. Combin. Theory* **1** (1966), 174–194.

[20] W. Mantel, Problem 28, *Wiskundige Opgaven* **10** (1907), 60–61.

[21] S. Mertens, Domination polynomials of the grid, the cylinder, the torus, and the king graph, *J. Integer Sequences* **27** (2024), Article 24.7.6.

[22] A. Meurer, C. P. Smith, M. Paprocki, O. Čertík, S. B. Kirpichev, M. Rocklin, A. Kumar, S. Ivanov, J. K. Moore, S. Singh, T. Rathnayake, S. Vig, B. E. Granger, R. P. Muller, F. Bonazzi, H. Gupta, S. Vats, F. Johansson, F. Pedregosa, M. J. Curry, A. R. Terrel, Š. Roučka, A. Saboo, I. Fernando, S. Kulal, R. Cimrman, and A. Scopatz, SymPy: symbolic computing in Python, *PeerJ Comput. Sci.* **3** (2017), e103.

[23] S. Mitchell, M. O'Sullivan, and I. Dunning, PuLP: a linear programming toolkit for Python, technical report, 2011, https://optimization-online.org/2011/09/3178/.

[24] B. M. E. Moret and H. D. Shapiro, On minimizing a set of tests, *SIAM J. Sci. Statist. Comput.* **6** (1985), 983–1003.

[25] S. Neuwirth, The size of bipartite graphs with girth eight, preprint, 2001, https://arxiv.org/abs/math/0102210v1.

[26] The Omega Institute, Prove the exact Three-Row blackout maximum, pull request 8084, merged on 15 September 2026 into the branch `dev` as commit `ace272f`, https://github.com/the-omega-institute/trureturing/pull/8084.

[27] A. Pemmasani, Kagey-Problems repository, 2026, commit 9d6ac7fc7d76, https://github.com/apemm/Kagey-Problems/tree/9d6ac7fc7d76/kagey-problems/problem001.

[28] I. Reiman, Über ein Problem von K. Zarankiewicz, *Acta Math. Acad. Sci. Hungar.* **9** (1958), 269–273.

[29] C. Sinz, Towards an optimal CNF encoding of Boolean cardinality constraints, in P. van Beek, ed., *Principles and Practice of Constraint Programming – CP 2005*, Lecture Notes in Comput. Sci., Vol. 3709, Springer, 2005, pp. 827–831.

[30] N. J. A. Sloane and The OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, 2026. Available at https://oeis.org.

[31] N. Wetzler, M. J. H. Heule, and W. A. Hunt, Jr., DRAT-trim: efficient checking and trimming using expressive clausal proofs, in C. Sinz and U. Egly, eds., *Theory and Applications of Satisfiability Testing – SAT 2014*, Lecture Notes in Comput. Sci., Vol. 8561, Springer, 2014, pp. 422–429.

2020 *Mathematics Subject Classification*: Primary 05A15; Secondary 05A16, 05B40, 05C35, 52C05, 68Q45, 68R15, 68V05.
Keywords: blackout, lattice rectangle, test cover, girth, transfer matrix, Mantel's theorem, regular language, SAT certificate.
(Concerned with sequences A001787, A002620, A033638, A085582, A289832, A322284, A338885, A397315, A398448, A399580, and A401226.)