## The University of Western Australia DEPARTMENT OF MATHEMATICS & STATISTICS

## AMO/TT TRAINING SESSIONS

## Tournament of the Towns Problems Senior Paper: Years 11, 12 Northern Autumn 2012 (A Level)

**Note:** Each contestant is credited with the largest sum of points obtained for three problems.

1. You are given an infinite sequence  $a_1, a_2, a_3, \ldots \in \mathbb{R}$ , where for each  $k \in \mathbb{N}$  there exists  $t = t(k) \in \mathbb{N}$  such that  $a_k = a_{k+t} = a_{k+2t} = \cdots$ .

Is this sequence necessarily periodic?

(4 points)

2. Chip and Dale play a game. To start, Chip puts 1001 nuts into 3 piles.

Dale knows the way they have been divided and chooses an integer  $N \in \{1, ..., 1001\}$ . Then Chip moves, if necessary, one or more nuts to make a fourth pile such that one pile or more piles contains a total of exactly N nuts.

Dale then gets the nuts moved to the fourth pile by Chip.

What is the largest number of nuts that Dale can get for sure, no matter how Chip acts?
(5 points)

3. A car is driven clockwise around a circular track. At noon Peter and Paul took up different positions on the track. The car passed each of them 30 times. Peter observed that each successive lap by the car was 1 second faster than the previous lap, while Paul observed that each successive lap was 1 second slower than the previous lap.

Prove that Peter and Paul were observing for at least an hour and a half. (6 points)

4. In  $\triangle ABC$  with incentre I, points  $C_1$  and  $A_1$  are chosen on the sides AB and BC, respectively, such that they do not coincide with the vertices. Let K be the midpoint of  $A_1C_1$  and suppose  $A_1BC_1I$  is cyclic.

Prove  $\angle AKC$  is obtuse. (8 points)

5. Peter and Paul play a game. First, Peter chooses  $a \in \mathbb{N}$  such that the sum of its digits S(a) is 2012. Paul wants to determine a; at first, he knows only that S(a) = 2012. On each turn, Paul chooses an  $x \in \mathbb{N}$  and Peter responds with S(|x-a|).

What is the least number of turns, in which Paul can determine a for sure? (8 points)

- 6. (a) A point A is chosen inside a sphere. Three pairwise perpendicular lines drawn through A intersect the sphere at six points.
  - Prove the centre of mass of the 6 points on the circle does not depend on the choice of the 3 lines. (5 points)
  - (b) A regular icosahedron with centre A is drawn inside a sphere (A does not necessarily coincide with the centre of the sphere). The rays going from A through the vertices of the icosahedron intersect the sphere at 12 points. Then the icosahedron is rotated about its centre. New rays going from A through the vertices of the icosahedron intersect the sphere at 12 new points. Let O and N be the centres of mass of old and new points on the sphere, respectively.

Prove that O = N. (An icosahedron has 20 triangular faces with each vertex incident with 5 faces.) (5 points)

7. There are 1000000 soldiers in a line. The sergeant splits the line into 100 (not necessarily equal) segments and permutes the segments, preserving the order of the soldiers in each segment, to form a new line. The sergeant repeats this procedure several times, splitting the line into segments of the same lengths in the same order as the first time, and permuting the segments in exactly the same way each time. Each soldier who was in the first segment after the first division recorded the number of times the procedure was performed until the first time he returned to the first segment.

Prove that at most 100 of the numbers recorded are different.

(10 points)