


**2017 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS****CALCULUS BC****SECTION II, Part A****Time—30 minutes****Number of questions—2****A GRAPHING CALCULATOR IS REQUIRED FOR THESE QUESTIONS.**

$h$ (feet)	0	2	5	10
$A(h)$ (square feet)	50.3	14.4	6.5	2.9

1. A tank has a height of 10 feet. The area of the horizontal cross section of the tank at height  $h$  feet is given by the function  $A$ , where  $A(h)$  is measured in square feet. The function  $A$  is continuous and decreases as  $h$  increases. Selected values for  $A(h)$  are given in the table above.
- (a) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the volume of the tank. Indicate units of measure.
- (b) Does the approximation in part (a) overestimate or underestimate the volume of the tank? Explain your reasoning.
- (c) The area, in square feet, of the horizontal cross section at height  $h$  feet is modeled by the function  $f$  given by  $f(h) = \frac{50.3}{e^{0.2h} + h}$ . Based on this model, find the volume of the tank. Indicate units of measure.
- (d) Water is pumped into the tank. When the height of the water is 5 feet, the height is increasing at the rate of 0.26 foot per minute. Using the model from part (c), find the rate at which the volume of water is changing with respect to time when the height of the water is 5 feet. Indicate units of measure.
-

2017 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS



2. The figure above shows the polar curves  $r = f(\theta) = 1 + \sin \theta \cos(2\theta)$  and  $r = g(\theta) = 2 \cos \theta$  for  $0 \leq \theta \leq \frac{\pi}{2}$ . Let  $R$  be the region in the first quadrant bounded by the curve  $r = f(\theta)$  and the  $x$ -axis. Let  $S$  be the region in the first quadrant bounded by the curve  $r = f(\theta)$ , the curve  $r = g(\theta)$ , and the  $x$ -axis.
- (a) Find the area of  $R$ .
- (b) The ray  $\theta = k$ , where  $0 < k < \frac{\pi}{2}$ , divides  $S$  into two regions of equal area. Write, but do not solve, an equation involving one or more integrals whose solution gives the value of  $k$ .
- (c) For each  $\theta$ ,  $0 \leq \theta \leq \frac{\pi}{2}$ , let  $w(\theta)$  be the distance between the points with polar coordinates  $(f(\theta), \theta)$  and  $(g(\theta), \theta)$ . Write an expression for  $w(\theta)$ . Find  $w_A$ , the average value of  $w(\theta)$  over the interval  $0 \leq \theta \leq \frac{\pi}{2}$ .
- (d) Using the information from part (c), find the value of  $\theta$  for which  $w(\theta) = w_A$ . Is the function  $w(\theta)$  increasing or decreasing at that value of  $\theta$ ? Give a reason for your answer.

END OF PART A OF SECTION II

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC**  
**2017 SCORING GUIDELINES**

**Question 1**

	1 : units in parts (a), (c), and (d)
<p>(a) <math>\text{Volume} = \int_0^{10} A(h) \, dh</math></p> $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3 \text{ cubic feet}$	<p>2 : <math>\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}</math></p>
<p>(b) The approximation in part (a) is an overestimate because a left Riemann sum is used and <math>A</math> is decreasing.</p>	1 : overestimate with reason
<p>(c) <math>\int_0^{10} f(h) \, dh = 101.325338</math></p> <p>The volume is 101.325 cubic feet.</p>	<p>2 : <math>\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}</math></p>
<p>(d) Using the model, <math>V(h) = \int_0^h f(x) \, dx</math>.</p> $\left. \frac{dV}{dt} \right _{h=5} = \left[ \frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5}$ $= \left[ f(h) \cdot \frac{dh}{dt} \right]_{h=5}$ $= f(5) \cdot 0.26 = 1.694419$ <p>When <math>h = 5</math>, the volume of water is changing at a rate of 1.694 cubic feet per minute.</p>	<p>3 : <math>\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}</math></p>