

## 2004 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS

6. Let  $f$  be the function given by  $f(x) = \sin\left(5x + \frac{\pi}{4}\right)$ , and let  $P(x)$  be the third-degree Taylor polynomial for  $f$  about  $x = 0$ .
- (a) Find  $P(x)$ .
- (b) Find the coefficient of  $x^{22}$  in the Taylor series for  $f$  about  $x = 0$ .
- (c) Use the Lagrange error bound to show that  $\left|f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right| < \frac{1}{100}$ .
- (d) Let  $G$  be the function given by  $G(x) = \int_0^x f(t) dt$ . Write the third-degree Taylor polynomial for  $G$  about  $x = 0$ .
- 

**END OF EXAMINATION**

**AP<sup>®</sup> CALCULUS BC**  
**2004 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function given by  $f(x) = \sin\left(5x + \frac{\pi}{4}\right)$ , and let  $P(x)$  be the third-degree Taylor polynomial for  $f$  about  $x = 0$ .

- (a) Find  $P(x)$ .
- (b) Find the coefficient of  $x^{22}$  in the Taylor series for  $f$  about  $x = 0$ .
- (c) Use the Lagrange error bound to show that  $\left|f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right| < \frac{1}{100}$ .
- (d) Let  $G$  be the function given by  $G(x) = \int_0^x f(t) dt$ . Write the third-degree Taylor polynomial for  $G$  about  $x = 0$ .

$$\begin{aligned} \text{(a)} \quad f(0) &= \sin\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2} \\ f'(0) &= 5\cos\left(\frac{\pi}{4}\right) = \frac{5\sqrt{2}}{2} \\ f''(0) &= -25\sin\left(\frac{\pi}{4}\right) = -\frac{25\sqrt{2}}{2} \\ f'''(0) &= -125\cos\left(\frac{\pi}{4}\right) = -\frac{125\sqrt{2}}{2} \\ P(x) &= \frac{\sqrt{2}}{2} + \frac{5\sqrt{2}}{2}x - \frac{25\sqrt{2}}{2(2!)}x^2 - \frac{125\sqrt{2}}{2(3!)}x^3 \end{aligned}$$

$$\text{(b)} \quad \frac{-5^{22}\sqrt{2}}{2(22!)}$$

$$\begin{aligned} \text{(c)} \quad \left|f\left(\frac{1}{10}\right) - P\left(\frac{1}{10}\right)\right| &\leq \max_{0 \leq c \leq \frac{1}{10}} \left|f^{(4)}(c)\right| \left(\frac{1}{4!}\right) \left(\frac{1}{10}\right)^4 \\ &\leq \frac{625}{4!} \left(\frac{1}{10}\right)^4 = \frac{1}{384} < \frac{1}{100} \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad \text{The third-degree Taylor polynomial for } G \text{ about } \\ x = 0 \text{ is } \int_0^x \left( \frac{\sqrt{2}}{2} + \frac{5\sqrt{2}}{2}t - \frac{25\sqrt{2}}{4}t^2 \right) dt \\ = \frac{\sqrt{2}}{2}x + \frac{5\sqrt{2}}{4}x^2 - \frac{25\sqrt{2}}{12}x^3 \end{aligned}$$

4 :  $P(x)$

$\langle -1 \rangle$  each error or missing term

deduct only once for  $\sin\left(\frac{\pi}{4}\right)$   
evaluation error

deduct only once for  $\cos\left(\frac{\pi}{4}\right)$   
evaluation error

$\langle -1 \rangle$  max for all extra terms,  $+\dots$ ,  
misuse of equality

2 :  $\begin{cases} 1 : \text{magnitude} \\ 1 : \text{sign} \end{cases}$

1 : error bound in an appropriate  
inequality

2 : third-degree Taylor polynomial for  $G$   
about  $x = 0$

$\langle -1 \rangle$  each incorrect or missing term

$\langle -1 \rangle$  max for all extra terms,  $+\dots$ ,  
misuse of equality