



- D. A student combines 35.00 mL of $1.85 \times 10^{-3} \text{ M Mg(NO}_3)_2(aq)$ with 50.00 mL of $2.80 \times 10^{-4} \text{ M NaOH}(aq)$, as shown in the diagram. Calculate $[\text{Mg}^{2+}]$ after the two solutions are combined but before any reaction takes place. (Assume that volumes are additive.)



- E. The dissolution of magnesium hydroxide is represented by the following equation.



- Write the expression for the solubility product constant, K_{sp} .
 - After the two solutions are combined in beaker 3 as described in part D, but before any reaction takes place, $[\text{OH}^{-}] = 1.65 \times 10^{-4} \text{ M}$. Using your answer to part D, calculate the value of the reaction quotient, Q .
 - Using the reaction quotient, Q , predict whether a precipitate should form as the mixture in beaker 3 approaches equilibrium. Justify your answer.
- F. In a separate experiment, the student adds $\text{HNO}_3(aq)$ to decrease the pH of a saturated solution containing undissolved $\text{Mg(OH)}_2(s)$. Does the amount of undissolved $\text{Mg(OH)}_2(s)$ increase, decrease, or remain the same as the $\text{HNO}_3(aq)$ is added? Justify your answer.

5. Complete Lewis diagrams and some physical properties for compounds X and Y are given.

Compound	X	Y
Lewis diagram		
Molar mass	74.1 g/mol	90.2 g/mol
Boiling point	82°C	98°C

- A. Based on VSEPR theory, predict the geometry around the Si atom in compound Y.
- B. A student claims that compound Y has a higher boiling point than that of compound X because compound Y has stronger London dispersion forces. Do you agree or disagree? Justify your answer.
- C. An equimolar mixture of the two compounds is heated. When the mixture reaches 82°C, which compound will have the higher vapor pressure? Justify your answer.
- D. The mixture is heated to 198°C in a sealed, rigid 12.5 L container, at which point both substances are gases and the total pressure in the container is 2.30 atm. Calculate the number of moles of gas particles in the container.

Question 5: Short Answer**4 points**

A For the correct answer: **Point 01***Tetrahedral*

B For the correct answer and a valid justification: **Point 02**

Agree. Compound Y has a larger, more polarizable electron cloud because the Si atom has more occupied electron shells than the C atom, giving compound Y stronger London dispersion forces and a higher boiling point than compound X.

C For the correct answer and a valid justification. **Point 03**

Examples of acceptable responses may include the following:

- Compound X. Compound X has weaker intermolecular forces than compound Y, so molecules of X are more likely to be in the gas phase at 82°C and would therefore have a higher vapor pressure.*
- Compound X. At 82°C, compound X has reached its boiling point, but compound Y has not. Therefore, the proportion of X molecules in the vapor phase would be much greater than that of compound Y, giving compound X the higher vapor pressure.*

D For the correct calculated value: **Point 04**

$$PV = nRT$$

$$n = \frac{PV}{RT} = \frac{(2.30 \text{ atm})(12.5 \text{ L})}{(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{K}\cdot\text{mol}})(471 \text{ K})} = 0.744 \text{ mol}$$
