





2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



5. At high temperatures the compound C_4H_6 (1,3-butadiene) reacts according to the equation above. The rate of the reaction was studied at 625 K in a rigid reaction vessel. Two different trials, each with a different starting concentration, were carried out. The data were plotted in three different ways, as shown below.



2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

- (a) For trial 1, calculate the initial pressure, in atm, in the vessel at 625 K. Assume that initially all the gas present in the vessel is C_4H_6 .
- (b) Use the data plotted in the graphs to determine the order of the reaction with respect to C_4H_6 .
- (c) The initial rate of the reaction in trial 1 is $0.0010 \text{ mol}/(\text{L} \cdot \text{s})$. Calculate the rate constant, k , for the reaction at 625 K.

AP[®] CHEMISTRY
2016 SCORING GUIDELINES

Question 5 (continued)

- (a) For trial 1, calculate the initial pressure, in atm, in the vessel at 625 K. Assume that initially all the gas present in the vessel is C₄H₆.

<p>For trial 1, $\frac{n}{V} = 0.020 \text{ mol/L}$ (or assume the volume of the vessel is 1.0 L; the number of moles of C₄H₆ in the vessel would then be 0.020 mol).</p> <p>$PV = nRT$</p> <p>$P = \frac{nRT}{V} = \frac{(0.020 \text{ mol})(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(625 \text{ K})}{1.0 \text{ L}} = 1.0 \text{ atm}$</p>	<p>1 point is earned for a correct setup.</p> <p>1 point is earned for the correct answer.</p>
---	--

- (b) Use the data plotted in the graphs to determine the order of the reaction with respect to C₄H₆.

Second order (because the plot of $1/[\text{C}_4\text{H}_6]$ is a straight line).	1 point is earned for the correct order.
---	--

- (c) The initial rate of the reaction in trial 1 is 0.0010 mol/(L·s). Calculate the rate constant, k , for the reaction at 625 K.

<p>From the second-order rate law (differential form): $\text{rate} = k[\text{C}_4\text{H}_6]^2$</p> <p>$\Rightarrow k = \frac{\text{rate}}{([\text{C}_4\text{H}_6])^2} = \frac{0.0010 \text{ mol/(L} \cdot \text{s)}}{(0.020 \text{ mol/L})^2} = 2.5 \text{ L/(mol} \cdot \text{s)}$</p> <p>OR</p> <p>From the second-order rate law (integrated form):</p> <p>$\frac{1}{[\text{C}_4\text{H}_6]_t} = 2kt + \frac{1}{[\text{C}_4\text{H}_6]_0}$</p> <p>The coefficient of t is equal to $2k$ because of the reaction stoichiometry.</p> <p>The slope of the line in the plot of $\frac{1}{[\text{C}_4\text{H}_6]}$ versus time is $2k$.</p> <p>Thus slope = $5.0 \text{ L/(mol} \cdot \text{s)} = 2k$, therefore $k = 2.5 \text{ L/(mol} \cdot \text{s)}$.</p> <p><u>Note:</u> Students who choose the second method of determining k but omit the factor of 2, thereby getting an answer of $5.0 \text{ L/(mol} \cdot \text{s)}$, still earn the point.</p>	<p>1 point is earned for the correct value.</p>
---	---